



# Iron deficiency: new insights into diagnosis and treatment

Clara Camaschella<sup>1</sup>

<sup>1</sup>Vita Salute University, San Raffaele Scientific Institute, Milan, Italy

Iron deficiency and iron deficiency anemia are common conditions worldwide affecting especially children and young women. In developing countries, iron deficiency is caused by poor iron intake and/or parasitic infection, whereas vegetarian dietary choices, poor iron absorption, and chronic blood loss are common causes in high-income countries. Erythropoiesis stimulating agents can result in functional iron deficiency for erythropoiesis even when stores are iron-replete. Diagnosis of iron deficiency is straightforward, except when it occurs in the context of inflammatory disorders. Oral iron salts correct absolute iron deficiency in most patients, because low hepcidin levels facilitate iron absorption. Unfortunately frequent side effects limit oral iron efficacy. Intravenous iron is increasingly utilized, because currently available preparations allow rapid normalization of total body iron even with a single infusion and are effective also in functional iron deficiency and in iron deficiency associated with inflammatory disorders. The evidence is accumulating that these preparations are safe and effective. However, long-term safety issues of high doses of iron need to be further explored.

## Learning Objectives

- To diagnose absolute iron deficiency, iron deficiency anemia, and functional iron deficiency in the light of recent advances in iron metabolism
- To evaluate advantages and disadvantages of oral and intravenous iron therapy

Disorders of iron metabolism are common human pathologic conditions. Iron deficiency, defined as low body iron with or without anemia, is estimated to affect >2 billion people. Iron deficiency anemia is the most prevalent anemia worldwide roughly affecting 1 of 8 persons, with the highest prevalence in developing countries.<sup>1</sup> Still iron deficiency anemia often remains undiagnosed and untreated.

Iron is tightly regulated at both the cellular and systemic level. Systemic iron homeostasis is under the control of hepcidin and is dominated by the erythropoietic needs for hemoglobin synthesis.<sup>2</sup> Thus, individuals at increased risk of developing iron deficiency are young children and adolescents, because of expanding erythropoiesis during rapid growth, and postpartum women. Also regular blood donors may develop iron deficiency and anemia after multiple donations.<sup>3</sup> Insufficient iron intake, diets containing poorly bioavailable sources of iron, and parasitic infections account for most cases in low-income countries.<sup>1</sup> In high-income countries, iron deficiency anemia, usually with the exception of groups with increased iron needs, may point to pathologic conditions associated with poor iron absorption and/or chronic blood loss.

A vast literature primarily based on observational studies indicates that iron deficiency anemia may have a detrimental effect on cognitive development in children, on physical performance and work productivity in adults, and result in unfavorable pregnancy outcomes. In addition, the direct effect of iron deficiency on tissues other than erythropoiesis (as muscles, heart, and brain) even in the

absence of anemia is a matter of a debate in view of the potential benefit of correcting tissue iron deficiency.

The laboratory diagnosis of iron deficiency anemia is well established, except when it occurs in patients with chronic infections/inflammation. In these cases, the increased cytokines and hepcidin levels cause body iron maldistribution that confounds the diagnostic parameters. The diagnostic approaches and the differential diagnosis of iron deficiency anemia, as well as the different strategies to supplement iron, are reviewed in light of our improved knowledge of the regulation of iron homeostasis.

## Mechanism of adaptation to iron deficiency

Iron deficiency anemia usually develops slowly. As iron levels decline in the stores (iron deficiency) and in the circulation (iron restricted erythropoiesis) becoming insufficient for the full hemoglobinization of mature erythroblasts (iron deficiency anemia), the liver peptide hormone hepcidin is transcriptionally suppressed.<sup>4</sup> Indeed serum hepcidin levels are significantly lower in young women with a negative iron balance compared with males and postmenopausal women, and are even undetectable in serum of individuals with iron deficiency anemia.<sup>5,6</sup> The decrease of hepcidin enhances iron release into plasma through ferroportin from both enterocytes and macrophages in the attempt of maintaining normal transferrin.

Animal models have clarified the compensatory mechanisms used to supply iron to erythropoiesis. In the initial phases of iron deficiency erythropoietin is reduced by the IRP1-mediated repression of hypoxia inducible factor-2 (HIF-2 $\alpha$ ) translation, to balance erythropoiesis with the limited iron availability.<sup>7,8</sup> Decreased diferric transferrin, the ligand of both transferrin receptors (TFR1 and TFR2) not only impairs cell iron uptake by erythroid TFR1,<sup>2</sup> but also destabilizes the iron sensor TFR2 on both the hepatocyte<sup>2</sup> and erythroid cell surface.<sup>9</sup> Whereas this contributes to hepcidin suppression in hepatocytes, it disrupts the TFR2-erythropoietin receptor

Conflict-of-interest disclosure: The author declares no competing financial interests.

Off-label drug use: None disclosed.

**Table 1. Etiology of iron deficiency**

<b>Physiological causes</b>	Increased iron demand: infancy, adolescence, 2 <sup>nd</sup> and 3 <sup>rd</sup> trimester of pregnancy
<b>Insufficient intake</b>	Malnutrition, vegetarians, vegans, iron poor-foods
<b>Decreased absorption</b>	Gastrectomy, duodenal by pass, bariatric surgery, <i>H. pylori</i> infection, celiac sprue, atrophic gastritis, inflammatory bowel diseases, drugs (proton pump inhibitors, H <sub>2</sub> blockers) Genetic IRIDA
<b>Chronic blood loss</b>	Gastrointestinal tract: any benign or malignant lesion, hookworm, drugs (salicylates, corticosteroids, NSAID) Genitourinary system: heavy uterine bleeding, intravascular hemolysis (eg, PNH) Bleeding defects (eg, hereditary hemorrhagic telangiectasia) Regular blood donors
<b>Associated with other conditions</b>	Chronic kidney disease, inflammatory bowel diseases, heart failure, obesity

IRIDA indicates iron-refractory-iron-deficiency-anemia; NSAID, nonsteroidal anti-inflammatory drugs; and PNH, paroxysmal nocturnal hemoglobinuria. Adapted from Camaschella.<sup>14</sup>

interaction in erythroid cells with the resulting increased erythropoietin sensitivity.<sup>9</sup> This translates into decreased apoptosis and an increased number of erythroblasts, optimizing the use of the available iron to produce microcytic red cells.<sup>9</sup> This moderate erythropoiesis expansion increases the hepcidin inhibitor erythroferone.<sup>9,10</sup> At the same time, the transmembrane protease *TMPRSS6* is stimulated in iron deficiency in order to attenuate the BMP-SMAD signaling and to repress hepcidin synthesis.<sup>11</sup> These mechanisms cooperate to sustain iron concentrations ensuring effective erythropoiesis, while at the same time avoiding excessive restriction of iron to other cells. Degradation of the storage protein ferritin occurs through “ferritinophagy”, an autophagy process carried out in starvation through NCOA4, a newly identified cargo protein, which targets ferritin to lysosomes for degradation.<sup>12</sup> Whether this is important to recycle iron when cell iron is limited is an issue of great interest. Once iron deficiency has led to overt anemia, HIF-2 $\alpha$  increases erythropoietin production in the kidney and enhances the expression of the iron importer DMT1, the iron reductase DCYT-B, and the iron exporter ferroportin in duodenal enterocytes.<sup>13</sup>

### Traditional and emerging causes of iron deficiency

Groups of individuals at risk and traditional causes of iron deficiency and iron deficiency anemia are well known, summarized in Table 1 and will not be extensively discussed here. For thorough coverage the readers are referred to a recent review.<sup>14</sup>

Impaired iron absorption may result from surgical and medical conditions. Bariatric surgery, increasingly performed to control caloric intake or diabetes, is emerging as a potential cause of iron deficiency.<sup>15</sup> Post-operative iron deficiency is influenced by preoperative iron status, which is often low in obese patients, and is found more commonly in females. Medical disorders affecting the gastrointestinal tract (eg, rarely autoimmune atrophic gastritis, more frequently *Helicobacter pylori* gastritis and celiac sprue) impair iron absorption and lead to resistance to oral iron treatment.<sup>16,17</sup> In inflammatory bowel diseases iron deficiency may be due to both poor absorption and blood loss. A functional reduction of iron absorption secondary to high hepcidin and ferroportin degradation may occur in these disorders, as well as in any other chronic inflammatory conditions. Iron deficiency is increasingly reported in the inflammatory setting, a combination that is challenging for diagnosis and relevant for treatment. This problem is exemplified by anemia of chronic kidney disease, which is due to blood loss, especially at dialysis, enhanced hepcidin levels (because of increased production and reduced excretion) and treatment with erythropoiesis stimulating agents (ESAs).<sup>18</sup> These agents may induce an imbalance between increased iron requirement and

insufficient iron mobilization from stores, especially when hepcidin levels are elevated, leading to functional iron deficiency. Iron deficiency is increasingly reported in patients with heart failure,<sup>19</sup> and may be a component of anemia in cancer patients and in the elderly, when associated with inflammation, low erythropoietin, and often multiple medications. Mild iron deficiency anemia is present in severely obese subjects, because of low-grade inflammation, increased hepcidin levels, and reduced iron absorption.<sup>14</sup>

### Genetics and iron deficiency

Genetic forms of iron deficiency anemia are extremely rare. Iron-refractory-iron-deficiency anemia (IRIDA) is a recessive anemia caused by mutations in the transmembrane protease serine member 6 (*TMPRSS6*) gene, which encodes the liver protease that inhibits hepcidin transcription by cleaving the coreceptor of the BMP-SMAD pathway hemojuvelin.<sup>11</sup> High hepcidin levels in IRIDA patients limit iron release to plasma from absorption sites and stores, leading to a microcytic hypochromic anemia that is unresponsive to oral iron.<sup>20</sup> Such a disorder is usually suspected in children or young patients whose anemia is persistent and refractory to oral iron treatment, in the absence of gastrointestinal conditions that limit iron absorption.<sup>17</sup> IRIDA has raised interest in the identification of genetic susceptibility to iron deficiency. Genome Wide Association Studies, including a recent meta-analysis, indicate that *TMPRSS6* genetic variants associate with red cell and iron traits in Caucasian and Asian populations.<sup>21-23</sup> The most common associated *TMPRSS6* variant (rs855791, A736V) influences serum hepcidin levels.<sup>24</sup>

The genetic susceptibility to iron deficiency is of special interest for transfusion medicine because it has implications for blood donation. Blood donors, especially females, carriers of the allele (736A) of the *TMPRSS6* variant rs855791 associated with low hepcidin levels,<sup>24</sup> seem more resistant to the development of iron deficiency after blood donation as compared to blood donors with the high-hepcidin associated variant.<sup>25</sup> Male donor carriers of a common polymorphism (P582S) in *HIF-1 $\alpha$* , which influences protein stabilization and thus the response to hypoxia, were found to have higher hemoglobin and ferritin levels and were protected from iron deficiency following blood donations compared with donors homozygous for the wild-type allele.<sup>25</sup> These pilot studies indicate the need for further investigation with the aim of improving blood donor health, controlling iron deficiency, and personalizing blood donations.

**Table 2. Laboratory tests to evaluate the iron status**

	Iron deficiency	Functional iron deficiency	Iron deficiency anemia	IRIDA	Anemia of chronic disease	Iron deficiency and anemia of chronic disease	Normal values (adult subjects)
<b>Currently used tests</b>							
Serum iron	↓	N/↓	↓	↓	↓	↓	10-30 μMol/L
TSAT, %	≥16	N/↓	<16	<10	N/↓	N/↓	>16<45
Serum ferritin, μg/L	<30	N	<12	Variable	>100	<100	20-200 (F) 40-300 (M)
Hb g/dL	N	N	↓	↓	↓	↓	>12 (F) >13 (M)
MCV, fl	N	N	<80	↓↓	N/↓	↓	80-95
MCH, pg	N	N	<27	↓↓	N/↓	↓	27-34
<b>Other tests</b>							
sTFR	↑	↑	↑	↑	N/↑	Variable	†
sTFR/log ferritin	NA	NA	>2*	NA	<1*	>2*	
ZPP	N	↑	↑	↑	↑	↑	†
Serum hepcidin	↓	↓	↓↓	N/↑	↑	N/↑	†
CHR pg	<25	<29	↓	↓	↓	↓	31.2±1.6
BM iron staining	+	±	-	+	+++	+	±

N indicates normal; TSAT, transferrin saturation; F, females; M, males; Hb, hemoglobin; ZPP, Zn Protoporphyrin; CHR, reticulocyte Hb content; BM, bone marrow; MCV, mean corpuscular volume; MCH, mean corpuscular hemoglobin; and sTFR, soluble transferrin receptor.

\* According to Weiss and Goodnough.<sup>28</sup>

† Normal values are according to the method used.

Adapted from Camaschella.<sup>14</sup>

### Laboratory diagnosis of iron deficiency and iron deficiency anemia: iron parameters

Serum ferritin <30ng/mL is the most sensitive and specific test to identify isolated iron deficiency,<sup>26</sup> as it reflects low stores (Table 2). In the progression of the deficiency, because of low iron and increased transferrin synthesis, transferrin saturation (or the saturated iron binding capacity) drops <16%, di-ferric transferrin, the ligand of transferrin receptor, is reduced and iron supply to the bone marrow becomes insufficient. At this point serum ferritin is usually <12 ng/mL.<sup>4</sup> In parallel, soluble transferrin receptor is shed in the circulation by the proprotein convertase subtilisin/kexin 7 (PCSK7), which cleaves the receptor when unbound to its ligand.<sup>27</sup>

Both transferrin saturation and serum ferritin are consensually decreased in iron deficiency anemia. Transferrin saturation may also be decreased in the anemia of chronic inflammation, but ferritin levels are increased, reflecting macrophage iron sequestration. In addition, transferrin and soluble transferrin receptor are not increased in inflammation. Although the distinction of iron deficiency anemia from anemia of chronic disease is straightforward, there is no clear-cut test that diagnoses iron deficiency anemia in the setting of inflammation (Table 2). Low transferrin saturation is one criterion with ferritin levels empirically set at <100 ng/mL<sup>28</sup> or higher (<300 ng/mL) as in chronic kidney disease or in heart failure.<sup>14</sup> Other proposed tests, such as the levels of serum soluble transferrin receptor (increased in iron deficiency and normal/low in inflammation) or the ratio between soluble transferrin receptor and log ferritin levels,<sup>28</sup> are uncommonly used in clinical practice.

Perl's iron staining of bone marrow smears allows visualization of the absence of iron in both macrophages and erythroblasts in iron deficiency; the presence of iron in macrophages but not in erythroblasts occurs in anemia of chronic inflammation. Once the gold standard for the diagnosis of iron deficiency, Perl's staining is rarely assessed because it is qualitative and requires performance of an invasive procedure.<sup>4</sup> In general the combination of tests rather than a single test helps to ascertain the correct diagnosis (Table 2).

In IRIDA, a discrepancy is often observed with extremely low transferrin saturation levels and normal/borderline low serum ferritin levels. Iron parameters differentiate IRIDA from other microcytic anemias,<sup>29</sup> and a normal C-reactive protein level from anemia of inflammation. Measuring serum hepcidin might be useful in IRIDA, because levels are abnormally high (or normal but inappropriately high)<sup>30</sup> considering the iron deficiency. At present, hepcidin assays are used for research but are not widely available for diagnostic purposes.

### Laboratory tests: erythrocyte traits

Low mean corpuscular volume and mean corpuscular hemoglobin are typical of iron deficiency anemia. Red cell distribution width (RDW), a measure of variation of red cell volume is increased, especially in the initial phases of iron deficiency when newly produced small hypochromic red cells coexist with normal ones. The RDW is also increased after iron therapy that induces the production of better hemoglobinized cells in the context of the array of hypochromic microcytic ones. RDWs, as well as erythrocyte indexes, are not affected in anemia of chronic inflammation. The differential diagnosis of microcytic anemia has been recently reviewed.<sup>29</sup>

Decreased reticulocyte hemoglobin content, a parameter that measures the functional iron available for erythropoiesis over the previous 3-4 days, is an early sign of iron-restricted erythropoiesis, (eg, after treatment with ESA). On the other side its increase is a measure of an early (2-4 days after treatment start) response following intravenous iron therapy. The percentage of hypochromic red cells is less useful because it is not as early a sign as the reticulocyte hemoglobin content.<sup>26</sup>

Lack of iron affects the last step of heme synthesis leading to accumulation of protoporphyrin IX and incorporation of zinc instead of iron in the protoporphyrin ring. Increased red cell zinc protoporphyrins may be used as a screening test for iron deficiency. However, protoporphyrins may increase also in the rare sideroblastic anemias due to defects of enzymes in the heme synthetic pathway.

## Searching for the etiology of iron deficiency

Except when iron deficiency is clearly related to increased requirements, patients should be evaluated for the cause(s) of iron deficiency, usually gastrointestinal blood loss, or poor iron absorption. A gastrointestinal work-up should be undertaken to identify a source of blood loss and video capsule endoscopy may be useful in cases with negative conventional endoscopy to explore the small bowel as a source of occult bleeding. Aggressive work up is recommended in iron deficient men and postmenopausal women. Gynecologic evaluation is mandatory in females with heavy uterine bleeding.

Non-invasive tests such as the urea breath test or antibodies for *Helicobacter pylori* and anti-transglutaminase antibodies for celiac sprue are indicated after failure of oral iron treatment, even in young patients.<sup>17</sup> C-reactive protein and a search for inflammatory/neoplastic conditions are especially relevant in the elderly, because, besides dietary deficiency, multiple causes are possible including clonal hematopoiesis. In rare instances paroxysmal nocturnal hemoglobinuria, as well as other causes of chronic intravascular hemolysis, should be searched for in unexplained cases of iron deficiency.

Sequencing of the *TMPRSS6* gene is essential when suspecting IRIDA.<sup>30</sup>

## Therapeutic options

### Oral iron supplementation

Oral iron supplementation is the first line of treatment of iron deficiency anemia. Protocols for dosing and duration of treatment are well defined. Anemic adult patients are ideally treated with iron salts containing 100-200 mg elemental iron administered in divided doses fasting. Lower doses may be used with fewer side effects in individual patients. Children require liquid preparations (3-6 mg/Kg). The gold standard of oral iron treatment is ferrous sulfate; other effective compounds are iron fumarate, gluconate, carbonyl iron, and polysaccharide-iron complex. Slow release iron preparations should be avoided, because iron release occurs beyond the duodenum, which is the site of maximal absorption. However, oral iron is not easily tolerated. Dyspepsia, nausea, vomiting, abdominal pain, constipation, or diarrhea, likely due to the oxidative properties of iron on the gastrointestinal mucosa, occur frequently, especially when iron is taken fasting. A recent meta-analysis of randomized controlled trials has shown that gastrointestinal side effects are more common with oral iron compared with placebo or intravenous iron.<sup>31</sup> For this reason, few patients fully adhere to the prescribed dose for the entire duration of treatment that should last 3-6 months in order to replenish iron stores. Iron administration with meals or dose reduction may increase tolerance. Interference by diet and drugs of common use (eg, antacids, H<sub>2</sub> blockers, and proton pump inhibitors) may also reduce pharmacologic iron absorption.

Oral iron is effective in correcting anemia; however, results of the same treatment on the reversal of iron deficiency in other tissues are controversial. Few studies suggest that fatigue and quality-of-life improve in iron-deficient non-anemic females treated with oral iron. However, the studies are small, heterogeneous in design, and not always unbiased.<sup>32</sup> Iron supplementation in very young preschool children living in low-income areas may correct anemia, but a large meta-analysis of randomized trials failed to find evidence of improved cognitive performance or positive effect on growth and development.<sup>33</sup> In endemic malarial areas iron supplementation raises the concern of increasing parasitemia or

even clinical malaria.<sup>34</sup> Additional well-designed studies are needed to define the benefits/risks of iron supplementation in low-income countries.

Low-dose (37.5 mg) oral iron supplementation used to prevent iron deficiency in blood donors with normal hemoglobin levels, compared with no supplementation, showed a reduced time to recover the post-donation decrease in Hb concentration irrespective of basal ferritin levels.<sup>35</sup>

Anemic patients with malabsorption, *Helicobacter pylori* infection, atrophic gastritis, previous gastric or duodenal surgery, genetic IRIDA, as well as patients with elevated hepcidin levels driven by inflammation, do not respond or only partially respond to oral iron treatment. The use of oral iron is controversial in active inflammatory bowel disease. Eradication of *Helicobacter pylori* in infected patients or a gluten-free diet in asymptomatic celiac disease may normalize iron absorption.<sup>17</sup>

### Intravenous iron: indications and controversies

The only alternative to oral iron route is intravenous treatment. The intramuscular route has been abandoned because of the inconvenience of painful injection, dark discoloration of the skin, and development of sarcoma in the site of injection in treated animals. Intravenous iron is indicated when intestinal absorption is presumably poor, when rapid Hb increase is needed, as in severe anemia in the second to third trimester of pregnancy, or in chronic bleeding due to inherited defects, as in hereditary hemorrhagic telangiectasia (Table 3). With the intravenous route of administration, the exact dose of iron needed to normalize Hb levels and to replenish the stores can be calculated. Thus, it is not unexpected that hemoglobin response is better to intravenous than to oral iron, as documented in several studies.<sup>36</sup>

One drawback of intravenous iron has been the need for multiple infusions. However, some preparations on the market may replace the total dose needed to correct iron deficiency anemia (up to 1000 mg) in a single or a couple of infusions (Table 4).<sup>37</sup> This approach reduces the number of clinic visits and may be quite useful in selected patients.

Intravenous iron is more effective than oral iron when combined with ESA in chronic kidney disease and may decrease the needs of these agents in dialysis patients<sup>18</sup>; uncertainty remains on the use of ESAs in the anemia of cancer patients undergoing chemotherapy.<sup>38</sup>

In rare cases of severe anemia of various etiologies, intravenous iron was used in combination with ESA in lieu of blood transfusions in Jehovah's witnesses, who refuse allogeneic human blood. Few studies have explored the use of parenteral iron in the preoperative setting.<sup>39</sup> Although iron parameters in heart failure suggest iron maldistribution rather than iron deficiency, intravenous iron administration even in the absence of anemia has been shown to improve physical performances (6 minute walking test) and New York Heart Association functional class and quality-of-life, compared with placebo.<sup>40</sup> The hypothesis that intravenous iron may correct cardiomyocyte iron deficiency is intriguing and needs to be further explored.

Intravenous iron is contraindicated in the course of infections, in the first trimester of pregnancy, and in patients with a history of iron or of other drug allergy.<sup>41</sup>

**Table 3. Indications for parenteral iron therapy**

<b>Established indications</b>	
Failure of oral iron treatment	Iron intolerance Iron refractoriness: gastrectomy, duodenal by pass, <i>Helicobacter pylori</i> infection, celiac disease, atrophic gastritis, inflammatory bowel disease, genetic IRIDA
Need for a quick recovery	Severe anemia in the second and third trimester of pregnancy Chronic bleeding not manageable with oral iron, as in inherited bleeding disorders
Associated with ESA	Anemia of chronic kidney disease
Others	Substitution for blood transfusions when not accepted by patient for religious reasons
<b>Potential indications</b>	Anemia of chronic kidney disease (without treatment with ESA) Persistent anemia after the use of ESA in cancer patients under chemotherapy Anemia of chronic disease unresponsive to ESA alone
<b>Potential indications with insufficient supporting data</b>	Iron deficiency in chronic heart failure Transfusion sparing strategy in surgical patients*

Celiac disease and *H. pylori* infection should be considered if anemia remains refractory after specific (diet or triple therapy respectively) treatment.<sup>17</sup>

IRIDA indicates iron refractory iron deficiency anemia; and ESA, erythropoiesis stimulating agents.

\* Ongoing clinical trials.

Adapted from Camaschella.<sup>14</sup>

Immediate side effects of intravenous iron infusion may be nausea, vomiting, headache, flushing, myalgia, pruritus, arthralgia, and back and chest pain. Hypophosphatemia is usually transient and asymptomatic.

Hypersensitivity reactions, even anaphylactic shock, were reported with the use of high-molecular-weight iron dextran, whose use is now abandoned. Severe hypersensitivity reactions with the other preparations (Table 4) are exceptional. Simple recommendations to minimize the risk include slow administration with careful patient monitoring during and after treatment, an adequate clinical environment with trained staff and the avoidance of antihistamine premedication.<sup>41</sup> Taking into account the ability of iron to generate oxidative stress and to increase susceptibility to infections<sup>36</sup> the long-term side effects of large amounts of iron administered when iron stores are not depleted (eg, in inflammation) remain to be verified.

## Conclusion

Oral iron remains the mainstay of therapy for most patients with iron deficiency. However, the optimal mode of iron supplementation depends on several factors related to both the patient and the type of drug. Factors to be considered include the underlying pathologic condition, the degree of anemia, the need for a rapid Hb increase, tolerability or refractoriness to previous treatments, history of allergy, drug cost, and side effects. When planning to administer

a large iron dose in a single infusion the risk–benefit ratio should be evaluated, considering that the long-term effects of large iron doses remain uncertain with respect to iron overload, oxidative stress, and susceptibility to infections.

## Correspondence

Clara Camaschella, Vita Salute University, San Raffaele Scientific Institute, Via Olgettina 58, 20132 Milano, Italy; Phone: +39-0226437782; Fax: +39-0226434723; e-mail: [camaschella.clara@hsr.it](mailto:camaschella.clara@hsr.it)

## References

- Kassebaum NJ, Jasrasaria R, Naghavi M, et al. A systematic analysis of global anemia burden from 1990 to 2010. *Blood*. 2014;123(5):615–624.
- Hentze MW, Muckenthaler MU, Galy B, Camaschella C. Two to tango: regulation of Mammalian iron metabolism. *Cell*. 2010;142(1):24–38.
- Cable RG, Glynn SA, Kiss JE, et al. Iron deficiency in blood donors: analysis of enrollment data from the REDS-II Donor Iron Status Evaluation (RISE) study. *Transfusion*. 2011;51(3):511–522.
- Camaschella C, Hoffbrand AV, Hershko C. Iron Deficiency and Disorders of Haem Synthesis. In: Hoffbrand V, Higgs D, Keeling D, Mehta A, eds. *Post Graduate Haematology*. Ed 7. Oxford: Wiley-Blackwell; 2016.
- Traglia M, Girelli D, Biino G, et al. Association of HFE and TMPRSS6 genetic variants with iron and erythrocyte parameters is only in part dependent on serum hepcidin concentrations. *J Med Genet*. 2011;48(9):629–634.
- Galesloot TE, Vermeulen SH, Geurts-Moespot AJ, et al. Serum hepcidin: reference ranges and biochemical correlates in the general population. *Blood*. 2011;117(25):e218–e225.
- Anderson SA, Nizzi CP, Chang YI, et al. The IRP1-HIF-2alpha axis coordinates iron and oxygen sensing with erythropoiesis and iron absorption. *Cell Metab*. 2013;17(2):282–290.
- Ghosh MC, Zhang DL, Jeong SY, et al. Deletion of iron regulatory protein 1 causes polycythemia and pulmonary hypertension in mice through translational derepression of HIF2alpha. *Cell Metab*. 2013;17(2):271–281.
- Nai A, Lidonnici MR, Rausa M, et al. The second transferrin receptor regulates red blood cell production in mice. *Blood*. 2015;125(7):1170–1179.
- Kautz L, Jung G, Valore EV, Rivella S, Nemeth E, Ganz T. Identification of erythroferrone as an erythroid regulator of iron metabolism. *Nat Genet*. 2014;46(7):678–684.
- Silvestri L, Pagani A, Nai A, De Domenico I, Kaplan J, Camaschella C. The serine protease matrilysin-2 (TMPRSS6) inhibits hepcidin activation by cleaving membrane hemojuvelin. *Cell Metab*. 2008;8(6):502–511.

**Table 4. Iron preparations for intravenous use**

Formulation	Dose, mg (infusion, min)	Maximum dose, mg (infusion)
Ferric gluconate	125 (10-60)	250 (60 min)
Iron sucrose	100-400 (2-90)	300 (2 hr)
<b>Iron dextran (LMW)</b>	100 (2)	<b>1000 (1-4 hr)</b>
<b>Ferumoxytol*†</b>	510 (>1)	<b>510 (15-60 min)</b>
<b>Ferric carboxymaltose</b>	750-1000 (15-30)	<b>750-1000 (15-30 min)</b>
<b>Iron isomaltoside‡</b>	20 Kg (15)	<b>20 Kg (15 min)</b>

Formulations that can be administered as a total dose in a single infusion are indicated in bold.

LMW indicates low molecular weight.

\* Approved only in chronic kidney disease patients.

† FDA black box warning advises slow administration of the drug and observation of patients for 30 minutes post-infusion.

‡ Availability of the products may differ in different countries, eg, licensed only in Europe.

Adapted from Camaschella.<sup>14</sup>

12. Mancias JD, Wang X, Gygi SP, Harper JW, Kimmelman AC. Quantitative proteomics identifies NCOA4 as the cargo receptor mediating ferritinophagy. *Nature*. 2014;509(7498):105-109.
13. Mastrogriannaki M, Matak P, Peyssonnaud C. The gut in iron homeostasis: role of HIF-2 under normal and pathological conditions. *Blood*. 2013;122(6):885-892.
14. Camaschella C. Iron-deficiency anemia. *N Engl J Med*. 2015;372(19):1832-1843.
15. Khanbhai M, Dubb S, Patel K, Ahmed A, Richards T. The prevalence of iron deficiency anaemia in patients undergoing bariatric surgery. *Obes Res Clin Pract*. 2015;9(1):45-49.
16. Choung RS, Ditch IC, Nadeau AM, et al. Trends and racial/ethnic disparities in gluten-sensitive problems in the United States: findings from the National Health and Nutrition Examination Surveys from 1988 to 2012. *Am J Gastroenterol*. 2015;110(3):455-461.
17. Hershko C, Camaschella C. How I treat unexplained refractory iron deficiency anemia. *Blood*. 2014;123(3):326-333.
18. Macdougall IC, Bock AH, Carrera F, et al. FIND-CKD: a randomized trial of intravenous ferric carboxymaltose versus oral iron in patients with chronic kidney disease and iron deficiency anaemia. *Nephrol Dial Transplant*. 2014;29(11):2075-2084.
19. van Veldhuisen DJ, Anker SD, Ponikowski P, Macdougall IC. Anemia and iron deficiency in heart failure: mechanisms and therapeutic approaches. *Nat Rev Cardiol*. 2011;8(9):485-493.
20. Finberg KE, Heeney MM, Campagna DR, et al. Mutations in TMPRSS6 cause iron-refractory iron deficiency anemia (IRIDA). *Nat Genet*. 2008;40(5):569-571.
21. Benyamin B, Ferreira MA, Willemsen G, et al. Common variants in TMPRSS6 are associated with iron status and erythrocyte volume. *Nat Genet*. 2009;41(11):1173-1175.
22. Benyamin B, Esko T, Ried JS, et al. Novel loci affecting iron homeostasis and their effects in individuals at risk for hemochromatosis. *Nat Commun*. 2014;5:4926.
23. Gichohi-Wainaina WN, Towers GW, Swinkels DW, Zimmermann MB, Feskens EJ, Melse-Boonstra A. Inter-ethnic differences in genetic variants within the transmembrane protease, serine 6 (TMPRSS6) gene associated with iron status indicators: a systematic review with meta-analyses. *Genes Nutr*. 2015;10(1):442.
24. Nai A, Pagani A, Silvestri L, et al. TMPRSS6 rs855791 modulates hepcidin transcription in vitro and serum hepcidin levels in normal individuals. *Blood*. 2011;118(16):4459-4462.
25. Kiss JE. Laboratory and genetic assessment of iron deficiency in blood donors. *Clin Lab Med*. 2015;35(1):73-91.
26. Goodnough LT, Nemeth E, Ganz T. Detection, evaluation, and management of iron-restricted erythropoiesis. *Blood*. 2010;116(23):4754-4761.
27. Schwienbacher C, Serafin A, Zanon A, Pramstaller PP, Pichler I, Hicks AA. Involvement of proprotein convertase PCSK7 in the regulation of systemic iron homeostasis. *Hepatology*. 2013;58(5):1860-1861.
28. Weiss G, Goodnough LT. Anemia of chronic disease. *N Engl J Med*. 2005;352(10):1011-1023.
29. DeLoughery TG. Microcytic anemia. *N Engl J Med*. 2014;371(14):1324-1331.
30. De Falco L, Sanchez M, Silvestri L, et al. Iron refractory iron deficiency anemia. *Haematologica*. 2013;98(6):845-853.
31. Tolkien Z, Stecher L, Mander AP, Pereira DI, Powell JJ. Ferrous sulfate supplementation causes significant gastrointestinal side-effects in adults: a systematic review and meta-analysis. *PLoS One*. 2015;10(2):e0117383.
32. Pasricha SR, Low M, Thompson J, Farrell A, De-Regil LM. Iron supplementation benefits physical performance in women of reproductive age: a systematic review and meta-analysis. *J Nutr*. 2014;144(6):906-914.
33. Pasricha SR, Hayes E, Kalumba K, Biggs BA. Effect of daily iron supplementation on health in children aged 4-23 months: a systematic review and meta-analysis of randomised controlled trials. *Lancet Glob Health*. 2013;1(2):e77-e86.
34. Sazawal S, Black RE, Ramsan M, et al. Effects of routine prophylactic supplementation with iron and folic acid on admission to hospital and mortality in preschool children in a high malaria transmission setting: community-based, randomised, placebo-controlled trial. *Lancet*. 2006;367(9505):133-143.
35. Kiss JE, Brambilla D, Glynn SA, et al. Oral iron supplementation after blood donation: a randomized clinical trial. *JAMA*. 2015;313(6):575-583.
36. Litton E, Xiao J, Ho KM. Safety and efficacy of intravenous iron therapy in reducing requirement for allogeneic blood transfusion: systematic review and meta-analysis of randomised clinical trials. *BMJ*. 2013;347:f4822.
37. Auerbach M, Ballard H. Clinical use of intravenous iron: administration, efficacy, and safety. *Hematology Am Soc Hematol Educ Program*. 2010;2010:338-347.
38. Gurusamy KS, Nagendran M, Broadhurst JF, Anker SD, Richards T. Iron therapy in anaemic adults without chronic kidney disease. *Cochrane Database Syst Rev*. 2014;12:CD010640.
39. Richards T, Clevenger B, Keidan J, et al. PREVENTT: preoperative intravenous iron to treat anaemia in major surgery: study protocol for a randomised controlled trial. *Trials*. 2015;16:254.
40. Anker SD, Comin Colet J, Filippatos G, et al. Ferric carboxymaltose in patients with heart failure and iron deficiency. *N Engl J Med*. 2009;361(25):2436-2448.
41. Rampton D, Folkersen J, Fishbane S, et al. Hypersensitivity reactions to intravenous iron: guidance for risk minimization and management. *Haematologica*. 2014;99(11):1671-1676.