

Is There a Role for β -Adrenoceptor Agonists in the Management of Acute Lung Injury and the Acute Respiratory Distress Syndrome?

Danny F. McAuley and Michael A. Matthay

Departments of Medicine and Anesthesia, and the Cardiovascular Research Institute,, University of California at San Francisco, San Francisco, California, USA

Abstract

Despite improvements in general supportive care and ventilatory strategies designed to limit lung injury, no specific pharmacological therapy has yet proven to be efficacious in the management of acute lung injury (ALI) and the acute respiratory distress syndrome (ARDS). Based on experimental studies, as well as studies of the *ex-vivo* human lung, pulmonary edema fluid clearance from the alveolar space can be augmented by both inhaled and systemic β_2 -adrenoceptor agonists (β_2 -agonists). Additionally, in the presence of lung injury, β_2 -agonists may reduce lung vascular permeability. Treatment with β_2 -agonists may also increase the secretion of surfactant and have anti-inflammatory effects. In view of these potentially beneficial effects, β_2 -agonist therapy should be evaluated for the treatment of lung injury in humans, particularly because they are already in wide clinical use and do not seem to have serious adverse effects in critically ill patients.

Acute lung injury (ALI) and acute respiratory distress syndrome (ARDS) are characterized by a constellation of clinical, radiological and physiological abnormalities, resulting in acute respiratory failure due to alveolar epithelial and lung endothelial injury (figure 1).

Two well-designed epidemiological studies estimating the incidence of ALI and ARDS have been published recently. The Australia and New Zealand Intensive Care Society Clinical Trials Group have estimated the incidence of ALI and ARDS as 34 and 28 cases per 100 000 per year, respectively.^[1] The second study reported the incidence of ALI/ARDS in hospitals that participated in the ARDS Network low tidal volume trial^[2] and extrapolated these results to estimate incidence rates for the US. The average incidence of ALI/ARDS in the ARDS network registry was 2.2 cases per intensive care unit (ICU) bed per year.^[2] The incidence of ALI in the US has been estimated as 22–64 cases per 100 000 per year.^[3] Although improvements in general supportive care have contributed to a decrease in mortality over the last 10

years,^[4,5] mortality associated with ARDS remains high, between 30% and 65% depending on the population that has been studied.^[2,4,6]

A recent review provided a comprehensive overview of the potential role of pharmacotherapy in the management of ARDS.^[7] In their initial description, Ashbaugh et al.^[8] described using “a clinical trial of a variety of drugs, respirators and fluid regimen” with limited success. Despite active investigation over the past 30 years, little progress has been made in identifying an effective pharmacological agent for the treatment of this condition. However, there is increasing evidence that β -adrenoceptor agonists (β -agonists) may have a potential role in the management of ARDS. This review will discuss the pathophysiological basis for the mechanism of action of β_2 -agonists in ARDS and review the evidence supporting a potential role for their use in the management of this condition. The potential beneficial and adverse effects of β_2 -agonists in the treatment of patients with ALI/ARDS are presented in tables I and II.




Fig. 1. Clinical and physiological abnormalities leading to the development of acute lung injury (ALI)/acute respiratory distress syndrome (ARDS) following pulmonary epithelial and endothelial damage.

1. Lung Epithelial and Endothelial Function

1.1 Alveolar Epithelial Fluid Clearance

The normal alveolar epithelium acts as a tight barrier to prevent alveolar flooding and has an important role in the active transport of fluid from the alveolar to the interstitial space. Alveolar fluid clearance is dependent on active sodium and chloride transport. The alveolar type II cell appears to be responsible for most of the ion transport via the apical sodium and chloride conductive pathways and the basolateral sodium-potassium activated adenosine triphosphatase (Na^+ , K^+ -ATPase) activity, although the alveolar type I cell and distal airway epithelium may also contribute to sodium and chloride ion transport.^[9]

Early ALI/ARDS is characterized by diffuse alveolar epithelial and endothelial injury with increased pulmonary endothelial permeability and pulmonary edema. In the majority of patients with ALI/ARDS alveolar fluid clearance is impaired; maximal alveolar fluid clearance was associated with better clinical outcomes (figure 2).^[10] The ability of the alveolar epithelium to remove edema fluid is an important first step in recovery from ALI/ARDS and the

degree of alveolar epithelial injury is an important predictor of outcome.^[10-12] The normal mechanisms regulating alveolar fluid clearance appear to be intact in mild-to-moderate lung injury^[13,14] (figure 3) and in some instances are upregulated.^[15,16]

Several experimental studies in animals, as well as in the *ex-vivo* human lung, have demonstrated that β_2 -agonists accelerate the rate of alveolar fluid clearance.^[17-19] It has been hypothesized that the mechanism underlying increased alveolar fluid clearance induced by β_2 -agonists is due to an increase in intracellular cyclic adenosine monophosphate (cAMP), which results in increased sodium transport across alveolar type II cells by upregulation of the apical sodium and chloride pathways and Na^+ , K^+ -ATPase and probably the cystic fibrosis transmembrane conductance regulator (CFTR) activity.^[9] Suggested mechanisms for the upregulation of sodium transport proteins by cAMP include augmented sodium channel open probability, increases in Na^+ , K^+ -ATPase α -subunit phosphorylation, as well as increased trafficking of epithelial Na^+ channels (ENaC) to the apical membrane and Na^+ , K^+ -ATPases to the basolateral membrane.^[9] β_2 -Agonists have been shown to increase both the expression^[20] and activity^[21] of sodium transport channels in the type II alveolar cells, as well as CFTR.

Table I. Potential beneficial effects of β₂-adrenoceptor agonists in the treatment of acute lung injury/acute respiratory distress syndrome

Target	Action
Lung epithelial and endothelial function	Increases alveolar fluid clearance; reduces pulmonary endothelial permeability; epithelial cytoprotection
Neutrophil function	Decreases neutrophil adhesion via inhibition of neutrophil Mac-1 cell surface expression and reduced epithelial intercellular adhesion molecule-1 expression; decreases neutrophil chemotaxis; decreases release of neutrophil mediators via neutrophil membrane stabilization; down-regulation of neutrophil priming and decreased release of neutrophil oxidative products; induces neutrophil apoptosis
Cytokine production	Inhibits nuclear factor-kappa B activation and translocation; decreases pro-inflammatory cytokine production; increases interleukin-10 production
Coagulation	Decreases activation of coagulation system; increases fibrinolysis
Surfactant	Increases surfactant secretion
Respiratory mechanics	Reduces airflow resistance and peak airway pressure; reduces plateau pressure with improvement in respiratory system compliance

cAMP stimulation has also been shown to upregulate fluid transport in the presence of septic or hypovolemic shock,^[16,22] hydrostatic edema,^[23-25] and experimental hyperoxic lung injury^[14,26,27] (figure 3). β-Adrenergic stimulation restores alveolar fluid clearance in an animal model of ventilator-associated lung injury.^[28] In addition, β₂-agonists can overcome the depressant effects of hypoxia on alveolar fluid clearance.^[29,30] Also, salmeterol, a long-acting β₂-agonist, has been associated with a reduction in the incidence of high altitude pulmonary edema in susceptible patients.^[31] Patients susceptible to high altitude pulmonary edema have impaired basal transepithelial sodium and fluid transport, suggesting a genetic defect of sodium and fluid transport, which is likely to be further impaired by hypoxia. One possible mechanism for the beneficial effect of salmeterol may be upregulation of alveolar fluid clearance.^[31]

Denopamine, a β₁-agonist, causes a dose-dependent increase in alveolar fluid clearance in *ex vivo* rat and guinea pig lungs.^[32] Although high concentrations of denopamine increased intracellular cAMP levels, at lower concentrations, which caused an increase in alveolar fluid clearance, denopamine failed to increase cAMP levels, suggesting that the mechanism of action for improved alveolar fluid clearance is, at least in part, cAMP independent.^[32] However, β₂-adrenergic stimulation is more important in increasing alveolar epithelial sodium and fluid transport. Dopamine, at doses associated with only a β₁-agonist effect, whether given intra-alveolar or intravenously, had no effect on alveolar fluid clearance *in vivo* in rats.^[33] Additionally, the increase in alveolar fluid clearance induced by dobutamine was blocked by selective β₂-antagonists.^[33]

1.2 Pulmonary Endothelial Permeability

β-Adrenergic stimulation reduces pulmonary vascular endothelial permeability in animal models of ALI.^[34-37] It has also been reported that activation of β-adrenergic receptors can reduce vascular endothelial permeability in the systemic circulation.^[38] Isoproterenol (isoprenaline) has also been shown to attenuate the increase in endothelial permeability induced by high vascular pressure.^[39] One study demonstrated that β₂-agonists reduced lung edema in isolated rat lungs subjected to ischemia/reperfusion injury both by restoring pulmonary endothelial permeability and upregulating alveolar fluid transport.^[36] In this study, edema formation was maximal when epithelial fluid transport was inhibited. Isoproterenol reduced edema formation when epithelial fluid transport was inhibited, indicating improvement in endothelial permeability, but reduced edema formation further when epithelial fluid transport was not inhibited, indicating an additional effect on upregulation of alveolar fluid transport.^[36] In acid aspiration-induced lung injury, clinically relevant airspace concentrations of

Table II. Potential adverse effects of β₂-adrenoceptor agonists in the treatment of acute lung injury/acute respiratory distress syndrome

Tachycardia and arrhythmia
Hypokalemia
Hyperglycemia
Increased risk of nosocomial infection
Lack of effect due to extensive alveolar epithelium and endothelial injury
β-Adrenoceptor down-regulation with attenuation of effect
β-Adrenoceptor genetic polymorphisms associated with variable effect
Pulmonary edema in pregnancy associated with the use of β ₂ -agonists as tocolytics




Fig. 2. Plot of hospital mortality in two groups of patients with acute lung injury or acute respiratory distress syndrome; those with maximal alveolar fluid clearance (i.e. >14% per hour), and those with impaired alveolar fluid clearance (i.e. <14% per hour). The mean rate of alveolar fluid clearance in the maximal clearance group was $34 \pm 28\%$ per hour. These threshold levels were determined on the basis of extrapolations from studies in the *ex vivo* human lung. Alveolar fluid clearance was measured from the change in protein concentration between two pulmonary edema fluid samples during the first 4 hours after intubation. Columns represent the hospital mortality rate in each group (* $p < 0.02$ vs impaired alveolar fluid clearance) [reproduced from Ware and Matthay,^[10] with permission].

albuterol (salbutamol), administered following acid injury, reduce pulmonary edema both by increasing alveolar fluid clearance and decreasing endothelial permeability.^[40]

1.3 Hemodynamic Effects and Lung Edema Formation

In addition to causing vascular smooth muscle relaxation β_2 -agonists have positive chronotropic and inotropic effects.^[41] The resultant hemodynamic effect is an increase in cardiac output with an associated increase in systemic and pulmonary arterial pressure.^[42] Pulmonary edema formation increases as pulmonary arterial blood flow and pressure increases.^[43] It is possible that β_2 -agonists, by increasing pulmonary artery pressure or blood flow, an effect that occurs in animals in response to β_2 -agonists delivered to the airspaces,^[44] may increase edema formation. However, the net effect is an overall reduction in lung edema due to increased alveolar fluid resorption and a reduction in endothelial permeability.

1.4 Epithelial Cytoprotection and Repair

There is also some evidence that β_2 -agonists have a cytoprotective action. β_2 -agonists cause a reduction in ultrastructural epithelial damage caused by bacteria, bacterial toxins and inflammatory mediators such as elastase, probably by maintaining intracellular levels of cAMP.^[45,46] The process of cell migration and wound repair involves cAMP-dependent mechanisms. Isoproterenol increases migration of bovine epithelial cells and accelerates closure of mechanically and enzymatically induced injury of monolayers.^[47]

2. Neutrophil Function, Oxidative Stress and Inflammatory Mediators

The acute phase of ALI/ARDS is characterized by infiltration and activation of neutrophils and inflammatory cytokines. It is likely that neutrophils have a central role in the development of lung injury.^[48] Pulmonary neutrophil sequestration occurs early in the development of ALI/ARDS^[49] and neutrophil activation seems to be an important step in mediating tissue damage.^[50] There is considerable evidence of oxidative damage in ALI/ARDS,^[51,52] and activated neutrophils secrete a variety of oxidants and granular enzymes that probably contribute to the endothelial and epithelial injury.^[53,54] Furthermore, neutrophil-dependent oxidant injury to the alveolar epithelium can inhibit β -adrenergic dependent alveolar fluid clearance.^[55-57]

2.1 Neutrophil Function

2.1.1 Neutrophil Adhesion

β -Adrenergic induced elevation of intracellular cAMP inhibits neutrophil adhesion to bronchial epithelial cells^[58] and vascular endothelium^[59] through inhibition of neutrophil Mac-1 cell surface expression. In addition, β_2 -agonists reduce intercellular adhesion molecule (ICAM)-1 expression in stimulated bronchial epithelial cells.^[60] Furthermore, in animal models, β_2 -agonists have been shown to inhibit neutrophil adhesion to the vascular endothelium.^[61]




Fig. 3. The effects of terbutaline on alveolar fluid clearance in rats exposed to 100% oxygen (hyperoxic) compared with rats exposed to room air. Amiloride reduced the rate of active sodium transport by approximately 50% in both groups (data not shown). Alveolar fluid clearance increased to $42 \pm 7\%$ in rats exposed to room air and to $50 \pm 10\%$ in hyperoxic-injured rats after terbutaline stimulation (* $p < 0.05$ vs control rats, † $p < 0.05$ vs hyperoxia). These findings indicate that in a rat model of moderate hyperoxic lung injury alveolar epithelial fluid clearance mechanisms are intact and up-regulated by β -adrenoceptor agonists (reproduced from Garat et al.,^[14] with permission).

2.1.2 Neutrophil Accumulation

β_2 -Agonists decrease systemic blood neutrophilia in rats^[62] and pulmonary neutrophil sequestration *in vivo* in humans^[63] in response to platelet-activating factor. Although not all studies are consistent,^[64] the majority show that β_2 -agonists inhibit neutrophil chemotaxis.^[65-68] One study has demonstrated a dose-dependent effect, with therapeutic concentrations of β_2 -agonists increasing and suprathreshold concentrations inhibiting neutrophil chemotaxis.^[69] The mechanism by which β_2 -agonists inhibit chemotaxis may be due, only in part, to increased levels of cAMP.^[67]

2.1.3 Neutrophil Function and Mediator Release

The production of neutrophil oxidant products and oxidant-mediated cytotoxicity is reduced by β -adrenergic stimulation,^[70-72] although the dose of β_2 -agonists required to achieve this effect may be suprathreshold.^[71] In addition, there is evidence that β_2 -agonists may have a biphasic effect on release of oxidative products with lower concentrations increasing and suprathreshold concentrations inhibiting release of oxidative products.^[69,73] The mechanism by which β_2 -agonists reduce oxidative stress appears to be via both β -adrenoceptor stimulation and intrinsic reactive oxygen scavenging ability.^[71,74,75] There is also evidence that β -adrenergic stimulation may have a differential effect on intra- and extracellular oxidant production, with one study demonstrating that adrenaline (epinephrine) reduces the generation of extracellular oxygen metabolites likely to mediate tissue injury while maintaining intracellular oxidant production for phagocytosis.^[76] β -Adrenergic stimulation also causes down-regulation of neutrophil priming.^[77,78]

2.1.4 Apoptosis

When neutrophils fail to undergo apoptosis and die by necrosis, they induce an inflammatory response that contributes to tissue injury. There is some preliminary evidence that β_2 -agonists induce apoptosis in neutrophils. Apoptosis, as measured by nuclear morphology, was increased in isolated neutrophils incubated with salmeterol for 20 hours.^[79] The action of salmeterol was completely reversed by both a non-selective and a selective β_2 -antagonist. Similar effects were seen with a phosphodiesterase-4 inhibitor. These findings suggest the effect of salmeterol is mediated by β_2 -adrenergic receptor activation and elevation of cAMP.^[79]

2.2 Cytokines

A complex network of cytokines, pro-inflammatory and anti-inflammatory, are involved in the inflammatory response in ALI/ARDS. The balance between pro- and anti-inflammatory

cytokines is likely to be critical in the development and persistence of ALI/ARDS.^[80,81] Levels of bronchoalveolar lavage (BAL) fluid interleukin (IL)-8, are increased in patients at risk of ALI/ARDS who subsequently develop ALI/ARDS. Furthermore, pulmonary macrophages from patients who developed ARDS contained higher levels of IL-8 than those who did not, suggestive of the fact that pulmonary macrophages were, in part, the cellular source of IL-8. Levels of IL-8 in blood did not predict development of ALI/ARDS.^[82,83] Tumor necrosis factor (TNF)- α and IL-1 β activity have also been found to be increased in pulmonary edema fluid in the early phase of ALI/ARDS.^[84] High initial plasma or BAL fluid levels of TNF α , IL-1 β , IL-2, IL-6 and IL-8 are associated with a poor outcome. Furthermore, persistent elevation of these cytokines was associated with increased mortality.^[85,86] IL-8 is important in the recruitment of neutrophils to the injured lung.^[80,83,87] Treatment with anti-IL-8 monoclonal antibody in experimental animal models of ALI/ARDS decreases the magnitude of ALI.^[88,89]

β -agonists modulate the release of a range of cytokines. β -agonists decrease lipopolysaccharide (LPS)-stimulated release of TNF α in bovine alveolar macrophages and HL-60 cells differentiated to macrophages,^[90] rat myocardial cells,^[91] in human blood and promonocytic THP-1 cells,^[92-94] *in vivo* in mice^[95] and *in vivo* in healthy humans.^[94] Adrenaline and exogenous cAMP inhibit LPS stimulated IL-1 β production from mouse peritoneal macrophages.^[96] LPS-stimulated release of IL-6 from rat myocardial cells^[91] and human whole blood^[97] is also inhibited by β -agonists. In a murine model of endotoxin-induced lung injury, intravenous treatment with the β -agonists, dobutamine and dopexamine, reduced IL-6 and macrophage inflammatory protein-2 mRNA expression and caused a reduction in neutrophil infiltration in BAL.^[98] In one study, however, isoproterenol increased the release of IL-6 *in vivo* in mice.^[95]

Studies examining the effect of β_2 -agonists on IL-8 production suggest that they may have a cell specific effect with an increase in IL-8 production from bronchial epithelial cells^[99,100] and a decrease in IL-8 production from neutrophils^[101] and monocytes.^[93] LPS-induced production of IL-10, an anti-inflammatory cytokine, is increased from human whole blood^[94] and *in vivo* in mice.^[95] These effects are predominately mediated through a β -adrenoceptor stimulated increase in cAMP,^[91-93,96,97,99] although increased IL-10 production is mediated through a combined effect on α - and β -adrenergic receptors.^[94] This effect may be mediated, at least in part, through inhibition of nuclear factor- κ B activation and translocation.^[93]

3. Coagulation

According to some pathological studies, microvascular thrombosis is a characteristic feature of clinical lung injury.^[102] ALI/ARDS is characterized by local abnormalities of fibrin turnover with an increased procoagulant response and depressed fibrinolytic activity resulting in abnormal fibrin. In the lungs, tissue factor can also be produced by the endothelial cells on stimulation with IL-6 and other inflammatory factors, creating a hypercoagulable environment.^[103-105] The finding that an early increase in pulmonary deadspace, which may reflect, in part, pulmonary microvascular thrombosis, is associated with higher mortality in ALI/ARDS, indicates this procoagulant state may be an important etiological factor in the development of ALI/ARDS.^[106] β -Adrenergic stimulation can inhibit the procoagulant effects of endotoxin. Adrenaline has been shown to reduce activation of the coagulation system as measured by decreased circulating thrombin-anti-thrombin complexes and enhances fibrinolysis as measured by increased plasma levels of tissue-type plasminogen activator (TPA).^[107] Isoproterenol has also been shown to stimulate increased secretion of TPA in dogs and healthy humans, an effect which can be blocked by a non-selective β -antagonist.^[108,109] These data suggest β -agonists may have beneficial effects on the procoagulant state potentially attenuating coagulation-dependent lung injury.

4. Surfactant

Alveolar type II epithelial cells produce surfactant, which has an important role in reducing the surface tension at the alveolar air-liquid interface, stabilizing alveoli and terminal airways at low lung volumes to allow efficient gas exchange. In addition to the mechanical properties of surfactant, the surfactant associated proteins have other important functions, which include antibacterial host defense.^[110] There is increasing evidence that dysfunction of the alveolar epithelial type II cells may be an important mechanism of injury in ALI/ARDS, as reflected by a worse outcome in patients with a reduced level of surfactant protein-D (SP-D) in the alveolar edema fluid within 24 hours of the onset of ALI.^[111] One prior study^[112] found that lower levels of SP-D in fluid of patients with ALI/ARDS was associated with a higher mortality rate. The lower levels of surfactant proteins in the distal airspaces of the lung early in the course of ALI suggest that the alveolar type II epithelial cells are not producing or releasing surfactant proteins normally. Alternatively, it is possible there is an increase in lung epithelial permeability and accelerated loss of SP-D from the airspaces in the more injured patients. Surfactant proteins may also be nitrated and inactivated in patients with ALI/ARDS.^[113] Inter-

estingly, β_2 -agonists improve surfactant release *in vitro* from animal type II pneumocytes^[114-116] and the human fetal lung.^[117]

5. Respiratory Mechanics

Reduced lung compliance is characteristic of ALI/ARDS. However, increased airflow resistance has also been documented in animal models and in patients with ALI/ARDS.^[118] Furthermore, there is evidence of expiratory flow limitation.^[119] There is data to suggest that β_2 -agonists have a beneficial role in improving respiratory mechanics. Not surprisingly, β_2 -agonists reduce airflow resistance and peak airway pressure.^[120-122] In addition, in some but not all studies,^[120] β_2 -agonists reduce plateau pressure.^[121,122] The reduction in plateau airway pressure suggests an improvement in respiratory system compliance, possibly because of a reduction in alveolar edema and/or increased surfactant secretion.

6. Dosage and Delivery

Experimental studies demonstrate that β_2 -agonists stimulate alveolar fluid clearance in animal models and healthy *ex-vivo* human lungs when drug concentrations of 10^{-5} – 10^{-6} mol/L are achieved in alveolar fluid.^[19,44] Plasma and edema fluid concentrations of adrenaline and noradrenaline, measured in mechanically ventilated patients with pulmonary oedema due to ALI/ARDS are significantly below this therapeutic level for up regulating alveolar fluid clearance.^[123] This suggests it might be possible to pharmacologically produce further increases in the rate of edema clearance by administration of exogenous β -agonists. In a retrospective observational study in intubated, ventilated patients with pulmonary edema albuterol levels were measured in the pulmonary edema fluid from mechanically ventilated patients with pulmonary edema. After a total aerosolized dose of 3.5 ± 2.6 mg albuterol in the previous 6 hours patients with ALI (9 patients, 11 samples) had a median albuterol level of 1240 ng/mL, which is equivalent to $>10^{-6}$ mol/L.^[124] Therefore, administration of salbutamol either by inhalation or intravenously, at these doses, is likely to achieve physiologically efficacious levels to upregulate fluid clearance.

7. Differences Among β_2 -Adrenoceptor Agonists (β_2 -Agonists)

Salmeterol is more potent than albuterol in upregulating alveolar fluid clearance *in vivo* in rats but the maximal rate of alveolar fluid clearance was similar with both agents.^[40] In the *ex vivo* human lung, salmeterol was more potent than terbutaline in enhancing alveolar fluid clearance.^[19] The effect of β_2 -agonists in

the modulation of neutrophil function appears variable. Salmeterol appears to be inhibiting neutrophil recruitment and degranulation more effectively.^[68,125-127] In contrast, although albuterol can inhibit neutrophil recruitment, in comparison to other β_2 -agonists, it does not inhibit neutrophil oxidant generation.^[72,127] Another study showed adrenaline was approximately 100-fold more potent in inhibiting formation of oxygen radicals compared with dobutamine or dopamine.^[75] Because adrenaline has alpha effects, α -adrenergic stimulation may contribute to effects on neutrophil function. α -Adrenergic stimulation inhibits neutrophil accumulation and activation in the lungs attenuating the severity of ALI.^[128] On the basis of these findings, salmeterol theoretically might be more effective than other β_2 -agonists in the treatment of ALI/ARDS, although the studies are limited.

8. Potential Problems in the Use of β -Agonists to Treat Acute Lung Injury (ALI)/Acute Respiratory Distress Syndrome (ARDS)

It is possible that an extensively injured, denuded alveolar epithelium might make it difficult for any pharmacologic intervention aimed at improving epithelial function to be effective because of the extent of epithelial injury. However, this appears less likely since the normal mechanisms regulating alveolar fluid clearance are maintained in at least moderate lung injury.^[13] In addition, some experimental models indicate that neutrophil-dependent oxidant injury to the alveolar epithelium is more resistant to β -adrenergic-dependent upregulation of alveolar fluid clearance.^[55-57] However, because β_2 -agonists may inhibit neutrophil function, they may still improve clinical outcome by reducing neutrophil mediated tissue damage. Prolonged treatment with β_2 -agonists may lead to down-regulation of β -adrenoreceptors and subsequently attenuate the effect on stimulating the rate of alveolar fluid clearance.^[129-131] Although β -adrenoreceptor down-regulation has been a concern in patients with asthma treated long-term with β_2 -agonists, it is less likely that this would be a significant issue with short-term usage (7 days) of β_2 -agonists in patients with ALI/ARDS. In a recent mouse study in which high-dose albuterol was administered over 6 days, there was no reduction in the rate of alveolar fluid clearance.^[132]

β_2 -Agonists may also be associated with potential adverse effects. Tachycardia and arrhythmias could develop. However, in a randomized, controlled trial of intravenous albuterol there was no increase in incidence of significant hemodynamic instability.^[133] The risk of cardiovascular adverse effects is likely to be reduced if the drug is given as a metered dose inhaler or nebulizer

where the systemic concentration will be significantly lower.^[124] Hypokalemia and hyperglycemia, are other potential adverse effects which would need to be monitored and treated, particularly given recent evidence demonstrating the importance of strict glycemic control in improving outcome in critically ill patients.^[134] Finally, given the inhibitory effects of β_2 -agonists on neutrophil function, it is possible that the incidence of nosocomial infection may be increased. However, despite these potential concerns, experience with β_2 -agonists in the critically ill indicates that these drugs are usually well tolerated and suggests they are not likely to be associated with serious adverse effects if used in the treatment of ALI/ARDS.

In patients with asthma, it is well recognized that there is some inter-individual variation in response to β_2 -agonists^[135] that is, at least in part, due to genetic variation in the β_2 -adrenergic receptor and its related modulatory protein.^[136,137] Genetic polymorphisms are associated with bronchodilator response and tachyphylaxis to β_2 -agonists in patients with asthma, and it is possible that a similar association may occur with responses to albuterol therapy in patients with ALI.^[138,139]

Pulmonary edema occurs rarely in pregnancy and has been associated with the use of β_2 -agonists as tocolytics. However, the underlying mechanism is not clearly understood. It is difficult to resolve the experimental evidence that β_2 -agonists accelerate alveolar fluid clearance with the association between β_2 -agonists and maternal pulmonary edema.^[140] However, in one large series of 86 pregnant patients the development of pulmonary edema was not associated with β_2 -agonists.^[141]

9. Conclusions

In summary, therapy with β_2 -agonists is an attractive potential strategy in the treatment of ALI/ARDS for several reasons. They may (i) improve alveolar fluid clearance and hasten the resolution of alveolar edema; (ii) reduce pulmonary endothelial permeability; (iii) modulate neutrophil function and the inflammatory response; and (iv) perhaps increase surfactant secretion. Hypothetically, β_2 -agonists may reduce the severity of lung injury, improve gas exchange and respiratory mechanics and reduce the need for mechanical ventilation, and improve survival. Assuming that β_2 -agonists modulate mechanisms which are more important in the development and early exudative phase of ARDS we speculate that these agents are more likely to be effective if they are commenced early after the onset of ALI/ARDS. Since these drugs do not seem to modulate the later fibroproliferative phase of ALI/ARDS, it is proposed treatment should be for 7 days, since

treatment periods longer than this are unlikely to be associated with additional benefit.

The concept of treating ALI/ARDS patients with β_2 -agonists merits further study. A phase II study examining the role of intravenous albuterol in ALI/ARDS has recently been completed in the UK. In this prospective, double-blind, placebo-controlled study, 40 patients were randomized within 48 hours of onset of ALI/ARDS to intravenous albuterol or placebo for 7 days. Extravascular lung water, a measure of pulmonary edema, was reduced with a trend towards reduced mortality in the albuterol-treated group. Furthermore, the treatment was well tolerated. This study provides the first proof of principle that in humans with ALI/ARDS, sustained treatment with an intravenous β -agonists may be beneficial.^[133] Also the NIH sponsored ARDS network in the US is currently considering undertaking a phase III prospective, randomized, double-blind, placebo-controlled study of nebulized albuterol. In this study patients will be enrolled within 48 hours of onset of ALI/ARDS to nebulized albuterol or placebo for 7 days with mortality at 60 days as the primary outcome measure.

Acknowledgments

Danny F. McAuley is funded by a Research Fellowship from the Peel Medical Research Trust.

Drs Matthay and McAuley have no conflicts of interest directly relevant to the content of this review. No sources of funding were used to assist in the preparation of this manuscript.

References

- Bersten AD, Edibam C, Hunt T, et al. Incidence and mortality of acute lung injury and the acute respiratory distress syndrome in three Australian States. *Am J Respir Crit Care Med* 2002; 165 (4): 443-8
- The Acute Respiratory Distress Syndrome Network. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000; 342: 1301-8
- Goss CH, Brower RG, Hudson LD, et al. Incidence of acute lung injury in the United States. *Crit Care Med* 2003; 31 (6): 1607-11
- Milberg JA, Davis DR, Steinberg KP, et al. Improved survival of patients with acute respiratory distress syndrome (ARDS): 1983-1993. *JAMA* 1995; 273 (4): 306-9
- Baudouin S. Improved survival in ARDS: chance, technology or experience? *Thorax* 1998; 53 (4): 237-8
- Monchi M, Bellenfant F, Cariou A, et al. Early predictive factors of survival in the acute respiratory distress syndrome: a multivariate analysis. *Am J Respir Crit Care Med* 1998; 158 (4): 1076-81
- dos Santos CC, Chant C, Slutsky AS. Pharmacotherapy of acute respiratory distress syndrome. *Expert Opin Pharmacother* 2002; 3 (7): 875-88
- Ashbaugh DG, Bigelow DB, Petty TL, et al. Acute respiratory distress in adults. *Lancet* 1967; II (7511): 319-23
- Matthay MA, Folkesson HG, Clerici C. Lung epithelial fluid transport and the resolution of pulmonary edema. *Physiol Rev* 2002; 82 (3): 569-600
- Ware LB, Matthay MA. Alveolar fluid clearance is impaired in the majority of patients with acute lung injury and the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001; 163 (6): 1376-83
- Pittet JF, Mackersie RC, Martin TR, et al. Biological markers of acute lung injury: prognostic and pathogenetic significance. *Am J Respir Crit Care Med* 1997; 155 (4): 1187-205
- Matthay MA, Wiener-Kronish JP. Intact epithelial barrier function is critical for the resolution of alveolar edema in humans. *Am Rev Respir Dis* 1990; 142 (6 Pt 1): 1250-7
- Matthay MA, Folkesson HG, Verkman AS. Salt and water transport across alveolar and distal airway epithelia in the adult lung. *Am J Physiol Lung Cell Mol Physiol* 1996; 270: L487-503
- Garat C, Meignan M, Matthay MA, et al. Alveolar epithelial fluid clearance mechanisms are intact after moderate hyperoxic lung injury in rats. *Chest* 1997; 111 (5): 1381-8
- Pittet JF, Wiener-Kronish JP, Serikov V, et al. Resistance of the alveolar epithelium to injury from septic shock in sheep. *Am J Respir Crit Care Med* 1995; 151 (4): 1093-100
- Pittet JF, Wiener-Kronish JP, McElroy MC, et al. Stimulation of lung epithelial liquid clearance by endogenous release of catecholamines in septic shock in anesthetized rats. *J Clin Invest* 1994; 94 (2): 663-71
- Sakuma T, Okaniwa G, Nakada T, et al. Alveolar fluid clearance in the resected human lung. *Am J Respir Crit Care Med* 1994; 150 (2): 305-10
- Sakuma T, Suzuki S, Usuda K, et al. Preservation of alveolar epithelial fluid transport mechanisms in rewarmed human lung after severe hypothermia. *J Appl Physiol* 1996; 80 (5): 1681-6
- Sakuma T, Folkesson HG, Suzuki S, et al. Beta-adrenergic agonist stimulated alveolar fluid clearance in ex vivo human and rat lungs. *Am J Respir Crit Care Med* 1997; 155 (2): 506-12
- Minakata Y, Suzuki S, Grygorczyk C, et al. Impact of beta-adrenergic agonist on Na⁺ channel and Na⁺-K⁺-ATPase expression in alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 1998; 275 (2 Pt 1): L414-22
- Suzuki S, Zuege D, Berthiaume Y. Sodium-independent modulation of Na⁺(+)-K⁺(+)-ATPase activity by beta-adrenergic agonist in alveolar type II cells. *Am J Physiol Lung Cell Mol Physiol* 1995; 268 (6 Pt 1): L983-90
- Pittet JF, Brenner TJ, Modelska K, et al. Alveolar liquid clearance is increased by endogenous catecholamines in hemorrhagic shock in rats. *J Appl Physiol* 1996; 81 (2): 830-7
- Frank JA, Wang Y, Osorio O, et al. Beta-adrenergic agonist therapy accelerates the resolution of hydrostatic pulmonary edema in sheep and rats. *J Appl Physiol* 2000; 89 (4): 1255-65
- Campbell AR, Folkesson HG, Berthiaume Y, et al. Alveolar epithelial fluid clearance persists in the presence of moderate left atrial hypertension in sheep. *J Appl Physiol* 1999; 86 (1): 139-51
- Lane SM, Maender KC, Awender NE, et al. Adrenal adrenaline increases alveolar liquid clearance in a canine model of neurogenic pulmonary edema. *Am J Respir Crit Care Med* 1998; 158 (3): 760-8
- Lasnier JM, Wangenstein OD, Schmitz LS, et al. Terbutaline stimulates alveolar fluid resorption in hyperoxic lung injury. *J Appl Physiol* 1996; 81 (4): 1723-9
- Saldias FJ, Comellas A, Ridge KM, et al. Isoproterenol improves ability of lung to clear edema in rats exposed to hyperoxia. *J Appl Physiol* 1999; 87 (1): 30-5
- Saldias FJ, Lecuona E, Comellas AP, et al. Beta-adrenergic stimulation restores rat lung ability to clear edema in ventilator-associated lung injury. *Am J Respir Crit Care Med* 2000; 162 (1): 282-7
- Vivona M, Matthay MA, Freidlander G, et al. Hypoxia reduces alveolar epithelial sodium and fluid transport in rats: reversal by beta-2 adrenergic agonist treatment. *Am J Respir Cell Mol Biol* 2001; 25: 554-61
- Planes C, Blot-Chabaud M, Matthay MA, et al. Hypoxia and beta 2-agonists regulate cell surface expression of the epithelial sodium channel in native alveolar epithelial cells. *J Biol Chem* 2002; 277 (49): 47318-24
- Sartori C, Allemann Y, Duplain H, et al. Salmeterol for the prevention of high-altitude pulmonary edema. *N Engl J Med* 2002; 346 (21): 1631-6
- Sakuma T, Tsuchihara C, Ishigaki M, et al. Denopamine, a beta (1)-adrenergic agonist, increases alveolar fluid clearance in ex vivo rat and guinea pig lungs. *J Appl Physiol* 2001; 90 (1): 10-6

33. Tibayan FA, Chesnutt AN, Folkesson HG, et al. Dobutamine increases alveolar liquid clearance in ventilated rats by beta-2 receptor stimulation. *Am J Respir Crit Care Med* 1997; 156 (2 Pt 1): 438-44
34. Minnear FL, DeMichele MA, Moon DG, et al. Isoproterenol reduces thrombin-induced pulmonary endothelial permeability in vitro. *Am J Physiol Lung Cell Mol Physiol* 1989; 257 (5 Pt 2): H1613-23
35. Minnear FL, DeMichele MA, Leonhardt S, et al. Isoproterenol antagonizes endothelial permeability induced by thrombin and thrombin receptor peptide. *J Appl Physiol* 1993; 75 (3): 1171-9
36. Khimenko PL, Barnard JW, Moore TM, et al. Vascular permeability and epithelial transport effects on lung edema formation in ischemia and reperfusion. *J Appl Physiol* 1994; 77 (3): 1116-21
37. Adkins WK, Barnard JW, May S, et al. Compounds that increase cAMP prevent ischemia-reperfusion pulmonary capillary injury. *J Appl Physiol* 1992; 72 (2): 492-7
38. Dobbins DE, Buehn MJ, Dabney JM. Bradykinin-mediated edema formation is blocked by levorotatory but not dextrorotatory terbutaline. *Microcirc Endothelium Lymphatics* 1988; 4 (5): 377-97
39. Parker JC, Ivey CL. Isoproterenol attenuates high vascular pressure-induced permeability increases in isolated rat lungs. *J Appl Physiol* 1997; 83 (6): 1962-7
40. McAuley DR, Frank JA, Fang X, et al. Clinically relevant concentrations of beta2-adrenergic agonists stimulate maximal cyclic adenosine monophosphate-dependent airspace fluid clearance and decrease pulmonary edema in experimental acid-induced injury. *Crit Care Med* 2004; 32 (7): 1470-6
41. Lands AM, Arnold A, McAuliff JP, et al. Differentiation of receptor systems activated by sympathomimetic amines. *Nature* 1967; 214 (88): 597-8
42. Crane J, Burgess C, Beasley R. Cardiovascular and hypokalaemic effects of inhaled salbutamol, fenoterol, and isoprenaline. *Thorax* 1989; 44 (2): 136-40
43. Landolt CC, Matthay MA, Albertine KH, et al. Overperfusion, hypoxia, and increased pressure cause only hydrostatic pulmonary edema in anesthetized sheep. *Circ Res* 1983; 52 (3): 335-41
44. Berthiaume Y, Staub NC, Matthay MA. Beta-adrenergic agonists increase lung liquid clearance in anesthetized sheep. *J Clin Invest* 1987; 79 (2): 335-43
45. Dowling RB, Rayner CF, Rutman A, et al. Effect of salmeterol on *Pseudomonas aeruginosa* infection of respiratory mucosa. *Am J Respir Crit Care Med* 1997; 155 (1): 327-36
46. Dowling RB, Johnson M, Cole PJ, et al. Effect of salmeterol on *Haemophilus influenzae* infection of respiratory mucosa in vitro. *Eur Respir J* 1998; 11 (1): 86-90
47. Spurzem JR, Gupta J, Veys T, et al. Activation of protein kinase A accelerates bovine bronchial epithelial cell migration. *Am J Physiol Lung Cell Mol Physiol* 2002; 282 (5): L1108-16
48. Ware LB, Matthay MA. The acute respiratory distress syndrome. *N Engl J Med* 2000; 342 (18): 1334-49
49. Steinberg KP, Milberg JA, Martin TR, et al. Evolution of bronchoalveolar cell populations in the adult respiratory distress syndrome. *Am J Respir Crit Care Med* 1994; 150 (1): 113-22
50. Matthay MA. Conference summary: acute lung injury. *Chest* 1999; 116 (1 Suppl.): 119S-26S
51. Matthay MA, Geiser T, Matalon S, et al. Oxidant-mediated lung injury in the acute respiratory distress syndrome. *Crit Care Med* 1999; 27 (9): 2028-30
52. Quinlan GJ, Lamb NJ, Tilley R, et al. Plasma hypoxanthine levels in ARDS: implications for oxidative stress, morbidity, and mortality. *Am J Respir Crit Care Med* 1997; 155 (2): 479-84
53. Fujishima S, Aikawa N. Neutrophil-mediated tissue injury and its modulation. *Intensive Care Med* 1995; 21 (3): 277-85
54. Carden D, Xiao F, Moak C, et al. Neutrophil elastase promotes lung microvascular injury and proteolysis of endothelial cadherins. *Am J Physiol* 1998; 275 (2 Pt 2): H385-92
55. Pittet JF, Lu LN, Morris DG, et al. Reactive nitrogen species inhibit alveolar epithelial fluid transport after hemorrhagic shock in rats. *J Immunol* 2001; 166 (10): 6301-10
56. Modelska K, Matthay MA, Brown LA, et al. Inhibition of beta-adrenergic-dependent alveolar epithelial clearance by oxidant mechanisms after hemorrhagic shock. *Am J Physiol Lung Cell Mol Physiol* 1999; 276 (5 Pt 1): L844-57
57. Laffon M, Lu LN, Modelska K, et al. Alpha-adrenergic blockade restores normal fluid transport capacity of alveolar epithelium after hemorrhagic shock. *Am J Physiol Lung Cell Mol Physiol* 1999; 277 (4 Pt 1): L760-8
58. Bloemen PG, van den Tweel MC, Henricks PA, et al. Increased cAMP levels in stimulated neutrophils inhibit their adhesion to human bronchial epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 1997; 272 (4 Pt 1): L580-7
59. Derian CK, Santulli RJ, Rao PE, et al. Inhibition of chemotactic peptide-induced neutrophil adhesion to vascular endothelium by cAMP modulators. *J Immunol* 1995; 154 (1): 308-17
60. Oddera S, Silvestri M, Lantero S, et al. Downregulation of the expression of intercellular adhesion molecule (ICAM)-1 on bronchial epithelial cells by fenoterol, a beta2-adrenoceptor agonist. *J Asthma* 1998; 35 (5): 401-8
61. Bowden JJ, Sulakvelidze I, McDonald DM. Inhibition of neutrophil and eosinophil adhesion to venules of rat trachea by beta 2-adrenergic agonist formoterol. *J Appl Physiol* 1994; 77 (1): 397-405
62. Altenburg SP, Bozza PT, Martins MA, et al. Adrenergic modulation of the blood neutrophilia induced by platelet activating factor in rats. *Eur J Pharmacol* 1994; 256 (1): 45-9
63. Masclans JR, Barbera JA, MacNee W, et al. Salbutamol reduces pulmonary neutrophil sequestration of platelet-activating factor in humans. *Am J Respir Crit Care Med* 1996; 154 (2 Pt 1): 529-32
64. Tse RL, Phelps P, Urban D. Polymorphonuclear leukocyte motility in vitro: VI. Effect of purine and pyrimidine analogues: possible role of cyclic AMP. *J Lab Clin Med* 1972; 80 (2): 264-74
65. Rivkin I, Rosenblatt J, Becker EL. The role of cyclic AMP in the chemotactic responsiveness and spontaneous motility of rabbit peritoneal neutrophils: the inhibition of neutrophil movement and the elevation of cyclic AMP levels by catecholamines, prostaglandins, theophylline and cholera toxin. *J Immunol* 1975; 115 (4): 1126-34
66. Hill HR, Estensen RD, Quie PG, et al. Modulation of human neutrophil chemotactic responses by cyclic 3',5'-guanosine monophosphate and cyclic 3',5'-adenosine monophosphate. *Metabolism* 1975; 24 (3): 447-56
67. Harvath L, Robbins JD, Russell AA, et al. cAMP and human neutrophil chemotaxis: elevation of cAMP differentially affects chemotactic responsiveness. *J Immunol* 1991; 146 (1): 224-32
68. Silvestri M, Oddera S, Lantero S, et al. Beta 2-agonist-induced inhibition of neutrophil chemotaxis is not associated with modification of LFA-1 and Mac-1 expression or with impairment of polymorphonuclear leukocyte antibacterial activity. *Respir Med* 1999; 93 (6): 416-23
69. Llewellyn-Jones CG, Stockley RA. The effects of beta 2-agonists and methylxanthines on neutrophil function in vitro. *Eur Respir J* 1994; 7 (8): 1460-6
70. Braga PC, Mancini L, Guffanti EE, et al. Effects of nedocromil sodium on the oxidative burst of polymorphonuclear leukocytes: comparison with salbutamol. *Drugs Exp Clin Res* 1997; 23 (1): 33-8
71. Gillissen A, Wickenburg D, van Zwoll D, et al. Beta-2-agonists have antioxidant function in vitro: II. The effect of beta-2-agonists on oxidant-mediated cytotoxicity and on superoxide anion generated by human polymorphonuclear leukocytes. *Respiration* 1997; 64 (1): 23-8
72. Ottonello L, Morone P, Dapino P, et al. Inhibitory effect of salmeterol on the respiratory burst of adherent human neutrophils. *Clin Exp Immunol* 1996; 106 (1): 97-102
73. Calhoun WJ, Stevens CA, Lambert SB. Modulation of superoxide production of alveolar macrophages and peripheral blood mononuclear cells by beta-agonists and theophylline. *J Lab Clin Med* 1991; 117 (6): 514-22
74. Mirza ZN, Kato M, Kimura H, et al. Fenoterol inhibits superoxide anion generation by human polymorphonuclear leukocytes via beta-adrenoceptor-dependent and -independent mechanisms. *Ann Allergy Asthma Immunol* 2002; 88 (5): 494-500
75. Weiss M, Schneider EM, Tarnow J, et al. Is inhibition of oxygen radical production of neutrophils by sympathomimetics mediated via beta-2 adrenoceptors? *J Pharmacol Exp Ther* 1996; 278 (3): 1105-13

76. Kopprasch S, Gatzweiler A, Graessler J, et al. Beta-adrenergic modulation of FMLP- and zymosan-induced intracellular and extracellular oxidant production by polymorphonuclear leukocytes. *Mol Cell Biochem* 1997; 168 (1-2): 133-9
77. Barnett CCJ, Moore EE, Partrick DA, et al. Beta-adrenergic stimulation down-regulates neutrophil priming for superoxide generation, but not elastase release. *J Surg Res* 1997; 70 (2): 166-70
78. Opdahl H, Benestad HB, Nicolaysen G. Effect of beta-adrenergic agents on human neutrophil granulocyte activation with N-formyl-methionyl-leucyl-phenylalanine and phorbol myristate acetate. *Pharmacol Toxicol* 1993; 72 (4-5): 221-8
79. Lee E, Smith J, Robertson T, et al. Salmeterol and inhibitors of phosphodiesterase 4 (PDE4) induce apoptosis in neutrophils from asthmatics: beta-adrenergic receptor-mediated salmeterol activity and additive effects with PDE4 inhibitors [abstract]. *Am J Respir Crit Care Med* 1999; 159: A329
80. Martin TR. Lung cytokines and ARDS: Roger S. Mitchell lecture. *Chest* 1999; 116 (1 Suppl.): 2S-8S
81. Park WY, Goodman RB, Steinberg KP, et al. Cytokine balance in the lungs of patients with acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2001; 164 (10 Pt 1): 1896-903
82. Donnelly SC, Strieter RM, Kunkel SL, et al. Interleukin-8 and development of adult respiratory distress syndrome in at-risk patient groups. *Lancet* 1993; 341 (8846): 643-7
83. Donnelly SC, Strieter RM, Kunkel SL, et al. Chemotactic cytokines in the established adult respiratory distress syndrome and at-risk patients. *Chest* 1994; 105 (3 Suppl.): 98S-9S
84. Pugin J, Verghese G, Widmer MC, et al. The alveolar space is the site of intense inflammatory and profibrotic reactions in the early phase of acute respiratory distress syndrome. *Crit Care Med* 1999; 27 (2): 304-12
85. Meduri GU, Kohler G, Headley S, et al. Inflammatory cytokines in the BAL of patients with ARDS: persistent elevation over time predicts poor outcome. *Chest* 1995; 108 (5): 1303-14
86. Meduri GU, Headley S, Kohler G, et al. Persistent elevation of inflammatory cytokines predicts a poor outcome in ARDS: plasma IL-1 beta and IL-6 levels are consistent and efficient predictors of outcome over time. *Chest* 1995; 107 (4): 1062-73
87. Goodman RB, Strieter RM, Martin DP, et al. Inflammatory cytokines in patients with persistence of the acute respiratory distress syndrome. *Am J Respir Crit Care Med* 1996; 154 (3 Pt 1): 602-11
88. Folkesson HG, Matthay MA, Hebert CA, et al. Acid aspiration-induced lung injury in rabbits is mediated by interleukin-8-dependent mechanisms. *J Clin Invest* 1995; 96 (1): 107-16
89. Modelska K, Pittet JF, Folkesson HG, et al. Acid-induced lung injury: protective effect of anti-interleukin-8 pretreatment on alveolar epithelial barrier function in rabbits. *Am J Respir Crit Care Med* 1999; 160 (5 Pt 1): 1450-6
90. Gu Y, Seidel A. Influence of salbutamol and isoproterenol on the production of TNF and reactive oxygen species by bovine alveolar macrophages and calcitriol differentiated HL-60 cells. *Immunopharmacol Immunotoxicol* 1996; 18 (1): 115-28
91. Newman WH, Castresana MR, Webb JG, et al. Isoproterenol inhibits transcription of cardiac cytokine genes induced by reactive oxygen intermediates. *Anesthesiology* 2002; 96 (4): 947-54
92. Severn A, Rapson NT, Hunter CA, et al. Regulation of tumor necrosis factor production by adrenaline and beta-adrenergic agonists. *J Immunol* 1992; 148 (11): 3441-5
93. Farmer P, Pugin J. Beta-adrenergic agonists exert their "anti-inflammatory" effects in monocytic cells through the IkappaB/NF-kappaB pathway. *Am J Physiol Lung Cell Mol Physiol* 2000; 279 (4): L675-82
94. Van Der PT, Coyle SM, Barbosa K, et al. Adrenaline inhibits tumor necrosis factor-alpha and potentiates interleukin 10 production during human endotoxemia. *J Clin Invest* 1996; 97 (3): 713-9
95. Szabo C, Hasko G, Zingarelli B, et al. Isoproterenol regulates tumour necrosis factor, interleukin-10, interleukin-6 and nitric oxide production and protects against the development of vascular hyporeactivity in endotoxaemia. *Immunology* 1997; 90 (1): 95-100
96. Koff WC, Fann AV, Dunegan MA, et al. Catecholamine-induced suppression of interleukin-1 production. *Lymphokine Res* 1986; 5 (4): 239-47
97. Van Der PT, Jansen J, Ender E, et al. Noradrenaline inhibits lipopolysaccharide-induced tumor necrosis factor and interleukin 6 production in human whole blood. *Infect Immun* 1994; 62 (5): 2046-50
98. Dhingra VK, Uusaro A, Holmes CL, et al. Attenuation of lung inflammation by adrenergic agonists in murine acute lung injury. *Anesthesiology* 2001; 95 (4): 947-53
99. Linden A. Increased interleukin-8 release by beta-adrenoceptor activation in human transformed bronchial epithelial cells. *Br J Pharmacol* 1996; 119 (2): 402-6
100. Korn SH, Jerre A, Brattsand R. Effects of formoterol and budesonide on GM-CSF and IL-8 secretion by triggered human bronchial epithelial cells. *Eur Respir J* 2001; 17 (6): 1070-7
101. Au BT, Teixeira MM, Collins PD, et al. Effect of PDE4 inhibitors on zymosan-induced IL-8 release from human neutrophils: synergism with prostanoids and salbutamol. *Br J Pharmacol* 1998; 123 (6): 1260-6
102. Tomashefski Jr JF, Davies P, Boggis C, et al. The pulmonary vascular lesions of the adult respiratory distress syndrome. *Am J Pathol* 1983; 112 (1): 112-26
103. Idell S. Endothelium and disordered fibrin turnover in the injured lung: newly recognized pathways. *Crit Care Med* 2002; 30 (5 Suppl.): S274-80
104. Vervloet MG, Thijs LG, Hack CE. Derangements of coagulation and fibrinolysis in critically ill patients with sepsis and septic shock. *Semin Thromb Hemost* 1998; 24 (1): 33-44
105. Abraham E. Coagulation abnormalities in acute lung injury and sepsis. *Am J Respir Cell Mol Biol* 2000; 22 (4): 401-4
106. Nuckton TJ, Alonso JA, Kallet RH, et al. Pulmonary dead-space fraction as a risk factor for death in the acute respiratory distress syndrome. *N Engl J Med* 2002; 346 (17): 1281-6
107. Van Der PT, Levi M, Dentener M, et al. Adrenaline exerts anticoagulant effects during human endotoxemia. *J Exp Med* 1997; 185 (6): 1143-8
108. Probst A, Lill H, Strein K. Beta-adrenergic receptor mediated release of tissue plasminogen activator in anaesthetized dogs. *Thromb Res* 1988; 50 (1): 9-17
109. Chandler WL, Levy WC, Stratton JR. The circulatory regulation of TPA and UPA secretion, clearance, and inhibition during exercise and during the infusion of isoproterenol and phenylephrine. *Circulation* 1995; 92 (10): 2984-94
110. Hawgood S, Clements JA. Pulmonary surfactant and its apoproteins. *J Clin Invest* 1990; 86 (1): 1-6
111. Cheng IW, Ware LB, Greene KE, et al. Prognostic value of surfactant proteins A and D in patients with acute lung injury. *Crit Care Med* 2003; 31 (1): 20-7
112. Greene KE, Wright JR, Steinberg KP, et al. Serial changes in surfactant-associated proteins in lung and serum before and after onset of ARDS. *Am J Respir Crit Care Med* 1999; 160 (6): 1843-50
113. Zhu S, Ware LB, Geiser T, et al. Increased levels of nitrate and surfactant protein a nitration in the pulmonary edema fluid of patients with acute lung injury. *Am J Respir Crit Care Med* 2001; 163 (1): 166-72
114. Griesse M, Gobran LI, Rooney SA. Signal-transduction mechanisms of ATP-stimulated phosphatidylcholine secretion in rat type II pneumocytes: interactions between ATP and other surfactant secretagogues. *Biochim Biophys Acta* 1993; 1167 (1): 85-93
115. Bernhard W, Muller B, Von Wichert P. Beta-Adrenergic priming of rats in vivo modulates the effect of beta-agonist in vitro on surfactant phospholipid metabolism of isolated lungs. *Eur J Clin Invest* 1994; 24 (6): 393-9
116. Gobran LI, Rooney SA. Regulation of SP-B and SP-C secretion in rat type II cells in primary culture. *Am J Physiol Lung Cell Mol Physiol* 2001; 281 (6): L1413-9
117. Odom MJ, Snyder JM, Mendelson CR. Adenosine 3',5'-monophosphate analogs and beta-adrenergic agonists induce the synthesis of the major surfactant apoprotein in human fetal lung in vitro. *Endocrinology* 1987; 121 (3): 1155-63
118. Wright PE, Bernard GR. The role of airflow resistance in patients with the adult respiratory distress syndrome. *Am Rev Respir Dis* 1989; 139 (5): 1169-74
119. Koutsoukou A, Bekos B, Sotiropoulou C, et al. Effects of positive end-expiratory pressure on gas exchange and expiratory flow limitation in adult respiratory distress syndrome. *Crit Care Med* 2002; 30 (9): 1941-9

120. Pesenti A, Pelosi P, Rossi N, et al. Respiratory mechanics and bronchodilator responsiveness in patients with the adult respiratory distress syndrome. *Crit Care Med* 1993; 21 (1): 78-83
121. Morina P, Herrera M, Venegas J, et al. Effects of nebulized salbutamol on respiratory mechanics in adult respiratory distress syndrome. *Intensive Care Med* 1997; 23 (1): 58-64
122. Wright PE, Carmichael LC, Bernard GR. Effect of bronchodilators on lung mechanics in the acute respiratory distress syndrome (ARDS). *Chest* 1994; 106 (5): 1517-23
123. Sakuma T, Gu X, Maeda S, et al. Adrenaline dependent stimulation of alveolar fluid clearance in the human lung is CFTR dependent [abstract]. *Proc Am Thorac Soc* 2005; 2: A299
124. Atabai K, Ware LB, Snider ME, et al. Aerosolized beta (2)-adrenergic agonists achieve therapeutic levels in the pulmonary edema fluid of ventilated patients with acute respiratory failure. *Intensive Care Med* 2002; 28 (6): 705-11
125. Whelan CJ, Johnson M, Vardey CJ. Comparison of the anti-inflammatory properties of formoterol, salbutamol and salmeterol in guinea-pig skin and lung. *Br J Pharmacol* 1993; 110 (2): 613-8
126. Ramage L, Blair AL, Cree IA, et al. Effect of salmeterol on polymorphonuclear leukocyte (PMNL) chemiluminescence in vitro. *J Biolumin Chemilumin* 1993; 8 (5): 247-52
127. Anderson R, Feldman C, Theron AJ, et al. Anti-inflammatory, membrane-stabilizing interactions of salmeterol with human neutrophils in vitro. *Br J Pharmacol* 1996; 117 (7): 1387-94
128. Arcaroli J, Yang KY, Yum HK, et al. Effects of catecholamines on kinase activation in lung neutrophils after hemorrhage or endotoxemia. *J Leukoc Biol* 2002; 72 (3): 571-9
129. Morgan EE, Hodnichak CM, Stader SM, et al. Prolonged isoproterenol infusion impairs the ability of beta (2)-agonists to increase alveolar liquid clearance. *Am J Physiol Lung Cell Mol Physiol* 2002; 282 (4): L666-74
130. Fabisiak JP, Vesell ES, Rannels DE. Interactions of beta adrenergic antagonists with isolated rat alveolar type II pneumocytes: I. Analysis, characterization and regulation of specific beta adrenergic receptors. *J Pharmacol Exp Ther* 1987; 241 (2): 722-7
131. Geiser T, Jarreau PH, Atabai K, et al. Interleukin-1beta augments in vitro alveolar epithelial repair. *Am J Physiol Lung Cell Mol Physiol* 2000; 279 (6): L1184-90
132. Sartori C, Fang X, McGraw DW, et al. Selected contribution: long-term effects of beta (2)-adrenergic receptor stimulation on alveolar fluid clearance in mice. *J Appl Physiol* 2002; 93 (5): 1875-80
133. Perkins GD, McAuley DF, Gao F, et al. The beta agonist lung injury trial (BALTI) [abstract]. *Proc Am Thorac Soc* 2005; 2: A294
134. Van den BG, Wouters P, Weekers F, et al. Intensive insulin therapy in the critically ill patients. *N Engl J Med* 2001; 345 (19): 1359-67
135. Drazen JM, Israel E, Boushey HA, et al. Comparison of regularly scheduled with as-needed use of albuterol in mild asthma. *Asthma Clinical Research Network. N Engl J Med* 1996; 335 (12): 841-7
136. McGraw DW, Forbes SL, Kramer LA, et al. Polymorphisms of the 5' leader cistron of the human beta2-adrenergic receptor regulate receptor expression. *J Clin Invest* 1998; 102 (11): 1927-32
137. Parola AL, Kobilka BK. The peptide product of a 5' leader cistron in the beta 2 adrenergic receptor mRNA inhibits receptor synthesis. *J Biol Chem* 1994; 269 (6): 4497-505
138. Ohe M, Munakata M, Hizawa N, et al. Beta 2 adrenergic receptor gene restriction fragment length polymorphism and bronchial asthma. *Thorax* 1995; 50 (4): 353-9
139. Tan S, Hall IP, Dewar J, et al. Association between beta 2-adrenoceptor polymorphism and susceptibility to bronchodilator desensitisation in moderately severe stable asthmatics. *Lancet* 1997; 350 (9083): 995-9
140. Lamont RF. The pathophysiology of pulmonary oedema with the use of beta-agonists. *Br Obstet Gynaecol* 2000; 107 (4): 439-44
141. DiFederico EM, Burlingame JM, Kilpatrick SJ, et al. Pulmonary edema in obstetric patients is rapidly resolved except in the presence of infection or of nitroglycerin tocolysis after open fetal surgery. *Am J Obstet Gynecol* 1998; 179 (4): 925-33

Correspondence and offprints: Dr *Michael A. Matthay*, Cardiovascular Research Institute, University of California at San Francisco, 505 Parnassus Ave, M-917 Box 0624, San Francisco, CA 94143-0624, USA.
E-mail: mmat@itsa.ucsf.edu