

# Isolation and Characterization of an Iron-Containing Superoxide Dismutase From Water Lily, *Nuphar luteum*<sup>1</sup>

Received for publication June 29, 1981 and in revised form August 18, 1981

MARVIN L. SALIN AND SUSAN M. BRIDGES

Department of Biochemistry, Mississippi State University, Mississippi State, Mississippi 39762

## ABSTRACT

A cyanide-insensitive superoxide dismutase (EC 1.15.1.1) was purified from leaves of the water lily *Nuphar luteum* (L.) Sibth. and Smith Subsp. *macrophyllum* (Small) Beal. The enzyme had a molecular weight of 46,000 and was composed of two equally sized subunits. Metal analysis showed the protein to contain about 1 gram atom of iron per dimer. The iron-containing superoxide dismutase was sensitive to H<sub>2</sub>O<sub>2</sub> as well as to azide. Antibody to the protein did not cross-react with iron superoxide dismutase isolated from the eucaryote *Brassica* or with algal extracts.

The enzyme SOD<sup>2</sup> serves a protective role in all aerobic organisms by detoxifying the superoxide free radical in a dismutative reaction (10, 23):  $O_2^- + O_2^- + 2 H^+ \rightarrow H_2O_2 + O_2$ . Three metalloproteins have been found to exhibit SOD activity. The Cu-Zn enzyme is found in all eucaryotes, land plants, and fungi. It consists of two subunits and contains two atoms of Cu and two of Zn per dimer. The protein can be distinguished by its sensitivity to cyanide and H<sub>2</sub>O<sub>2</sub> (10). A Mn-containing SOD has been found in bacteria, as well as in the mitochondrial matrix of plants and animals. A metal content of one to four atoms of Mn per molecule has been reported (10, 22). The Mn isozyme is resistant to cyanide and to H<sub>2</sub>O<sub>2</sub> (10). The Fe-containing SOD, found predominantly in procaroytes, has a metal content of one to two atoms Fe per molecule and is cyanide-insensitive, but, unlike the Mn enzyme, it is inhibited by H<sub>2</sub>O<sub>2</sub> (10, 22).

SODs have been examined in photosynthetic organisms (2, 3, 19). The Cu-Zn isozyme is absent in procaroytic and most eucaryotic algae. However, the Mn and/or Fe enzymes are present (1, 2, 3). Land plants contain large amounts of the Cu-Zn isozyme plus an additional cyanide-resistant SOD, which was assumed to be the Mn protein (3, 19). We have recently reported on a survey of plant families for isozymes of SOD (6). Of 43 families of plants examined, three families—Ginkgoaceae, Cruciferae, and Nymphaeaceae—were found to contain the Fe enzyme.

In this paper, we report the results of our studies on the purification and characterization of Fe SOD from the water lily *Nuphar luteum*, a member of the Nymphaeaceae.

## MATERIALS AND METHODS

Cyt *c* (type III), xanthine oxidase (grade I), and nitroblue tetrazolium (type III) were obtained from Sigma Chemical Com-

<sup>1</sup> Supported by funds made available by the Mississippi Agricultural and Forestry Experiment Station. Mississippi Agricultural and Forestry Experiment Station Publication No. 4883.

<sup>2</sup> Abbreviations: SOD, superoxide dismutase; Fe SOD, iron-containing superoxide dismutase; Cu-Zn SOD, copper-zinc-containing superoxide dismutase.

pany. Iron SOD was purified from *Brassica campestris*, as previously reported (27), and Cu/Zn SOD was purified from wheat germ (5). *Nuphar luteum* (L.) Sibth. and Smith Subsp. *macrophyllum* (Small) Beal was collected and identified by Dr. Sidney McDaniel, Mississippi State University. Other vascular plants were grown locally. *Chlorella* sp. (UM 10 isolate), *Scenedesmus* sp. (UTEX 1588), and *Nostoc muscorum* (UTEX B1037) were generously supplied by Dr. Bailey Ward, University of Mississippi. Other algae were obtained from the R. C. Starr Collection, University of Texas.

SOD activity was assayed by the Cyt *c* reduction method (23), using a Perkin-Elmer Model 552 spectrophotometer. One unit of activity was defined as the amount of enzyme required to inhibit Cyt *c* reduction by 50%. Crude extracts were dialyzed extensively against 50 mM K-phosphate (pH 7.8) to deplete the extract of low mol wt reductants before assay of activity.

Analytical gel electrophoresis was performed on 7.5% acrylamide gels (8), with SOD activity localized using the photochemical method (4). Protein concentrations of extracts were determined by the method of Lowry *et al.* (17), while the concentration of purified protein was determined according to Murphy and Kies (25).

Mol wt was determined by gel exclusion on a Sephadex G-150 column. Subunit mol wt was determined by SDS gel electrophoresis, according to the method of Laemmli (16). Mol wt standards used for both mol wt and subunit mol wt determinations were: ribonuclease A, 13,700; chymotrypsinogen A, 25,000; ovalbumin, 43,000; and BSA, 67,000. Iron, copper, zinc, and manganese were assayed by atomic absorption spectrometry, using a Perkin-Elmer Model 305B atomic absorption photometer.

Purified enzyme was prepared for amino-acid analysis by extensive dialysis against H<sub>2</sub>O, followed by hydrolysis *in vacuo* in 6 M HCl at 110°C for 24 h, and then analyzed by a Beckman Model 120C amino-acid analyzer. Cysteine was determined as S-sulfo-cysteine after hydrolysis with boiling HCl, followed by reduction with DTT and sodium tetrathionate (13). Tryptophan was measured spectrophotometrically in 6 M guanidine hydrochloride, according to the method of Edelhoch (9).

The isoelectric point of the purified protein was determined on a Desaga-Brinkman double chamber thin layer electrofocusing apparatus. The enzyme was applied to a thin glass plate covered with ampholytes on a matrix of Sephadex G-75. After electrophoresis at 900 v for 3 h, a paper impression of the plate was made and stained for protein with Coomassie blue. The corresponding pH values were determined directly off the glass plate by insertion of a glass electrode (Broadley-James, Santa Ana, CA) into the gel matrix.

Antisera to purified *Nuphar* Fe SOD and wheat germ Cu-Zn SOD were produced by immunization of New Zealand white rabbits with a 2-ml emulsion consisting of 1 ml purified enzyme in 50 mM K-phosphate emulsified with 1 ml of Freund's Complete Adjuvant. One-half of the emulsion was injected into the footpads

Table I. Purification of Fe SOD from *N. luteum*

| Purification Step  | Volume | Protein Concentration | Total Protein | Total Activity | Specific Activity | Yield | Purification |
|--|--------|-----------------------|---------------|----------------|-------------------|-------|--------------|
|  | ml     | mg/ml                 | mg            | unit           | unit/mg           | %     | -fold        |
| Crude extract after centrifugation                                 | 615    | 3.2 <sup>a</sup>      | 1,968         | 51,718         | 26                | 100   | 1            |
| 35 to 65% (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> dialyzed | 112    | 2.0 <sup>a</sup>      | 224           | 46,368         | 207               | 89.7  | 7.9          |
| DEAE-cellulose   | 166    | 0.72 <sup>a</sup>     | 119           | 34,362         | 289               | 66.4  | 11.0         |
| CM-cellulose   | 211    | 0.49 <sup>a</sup>     | 103           | 32,705         | 318               | 63.2  | 12.2         |
| G-75   | 11.5   | 1.80 <sup>a</sup>     | 20.7          | 12,777         | 617               | 24.7  | 23.7         |
| G-75   | 3.5    | 1.04 <sup>b</sup>     | 3.6           | 7,248          | 2,013             | 14.0  | 77.4         |

<sup>a</sup> Lowry *et al.* (17).

<sup>b</sup> Murphy and Kies (25).




FIG. 1. Polyacrylamide gels of purified Fe SOD from *N. luteum*. A, Stained for protein with Coomassie blue (28  $\mu$ g); B, stained for SOD activity using nitroblue tetrazolium (50 ng).

of the rabbit and the remainder at multiple intramuscular sites. Subsequent injections at weekly intervals were entirely intramuscular with *Nuphar* Fe SOD emulsified in Freund's Incomplete Adjuvant. After a satisfactory titer was reached (3–4 weeks), the rabbits were sacrificed by cardiac puncture. The immunoglobulin was partially purified from the serum by precipitation with 33% (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, three times followed by dialysis into a buffer containing 0.025 M sodium tetraborate, 0.1 M boric acid, and 0.06 M NaCl, adjusted to pH 8.4.

Cross-reactivity of the antibodies with various purified proteins and cell-free extracts was tested by the Ouchterlony technique (26) of double immunodiffusion followed by staining with Coomassie Blue.

## RESULTS

**Purification of Enzyme.** The Fe SOD from *N. luteum* was purified using a modification of the procedure reported for *Brassica* Fe SOD (27). About 250 g of leaves were homogenized in a Waring Blender in 500 ml of grinding media containing 50 mM K-phosphate (pH 6.8), 0.3% Triton X-100, and 1 mM DTT. Cell debris was removed by filtration through eight layers of cheesecloth, followed by centrifugation at 10,000g for 10 min. The supernatant was brought to 30% saturation by addition of solid (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and stirred for 1 h at room temperature. The resulting precipitate was removed by centrifugation at 10,000g for 10 min, and the supernatant was brought to 65% saturation by additional solid (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>. After stirring for 1 h at room temperature, the precip-




FIG. 2. Absorption spectra of Fe SOD from *N. luteum*. Protein concentrations were 0.289 mg/ml (A) and 2.89 mg/ml (B).

itate was collected by centrifugation at 10,000g for 10 min, resuspended in 10 mM phosphate buffer (pH 6.8), and dialyzed for 24 h against 10 mM phosphate buffer (pH 6.8), with frequent changes of dialysis buffer.

The dialysate was clarified by centrifugation and then applied to a 37- × 2-cm DEAE column (Whatman DE-52) at 4°C equilibrated with 10 mM phosphate buffer (pH 6.8). The enzyme was eluted with a linear gradient of K-phosphate buffer (pH 6.8) (10–100 mM), and fractions with enzymic activity were pooled and dialyzed against 10 mM K-phosphate buffer (pH 5.8). The protein was then passed through a carboxymethyl cellulose (CM-32, Whatman) column (21.5 × 1.5 cm) equilibrated with 10 mM K-phosphate buffer (pH 5.8). SOD activity was not absorbed onto the column and was washed through with 50 ml of 10 mM K-phosphate buffer (pH 5.8). The eluate was concentrated to a volume of 2 ml by ultrafiltration and further purified by gel filtration on a Sephadex G-75 column (77 × 1.5 cm). Fractions containing the greatest amount of SOD activity were pooled and rechromatographed on a Sephadex G-75 column (77 × 1.5 cm). Table I summarizes the results of the purification. It can be estimated that 250 g of *Nuphar* contained approximately 25 mg of CN<sup>-</sup>-insensitive SOD. A yield of about 15% was recorded, and the enzyme had a specific activity of about 2,000. This value is lower than that for Cu/Zn SODs but is in agreement with values reported (15, 27, 30) in the literature for other Fe SODs. Unlike Fe SOD isolated and purified from *Brassica* (27), the *Nuphar*

protein, upon purification, was relatively stable at 4°C.

Figure 1 is a photograph of polyacrylamide gels electrophoresed and stained for activity and protein. Several bands of isozymic activity, corresponding to the protein are visible.

**Mol Wt.** The mol wt of the purified protein was determined by chromatography on a Sephadex G-150 column. By comparison with markers of known mol wt, the data obtained yielded a value of 46,200. Subunit size, determined by SDS-gel electrophoresis in the presence of  $\beta$ -mercaptoethanol (16), revealed a single band corresponding to a value of 23,000. It would appear, therefore, that the protein is composed of two equally sized subunits.

**Absorption Spectra.** Figure 2 shows the visible and UV absorption spectra of the purified protein. The enzyme exhibited a peak at 278 nm, typical of most proteins. Shoulders at 290 and 260 nm were also observed. A weak, broad absorption spectrum was observed in the range of 450 nm to 300 nm. Both UV and visible absorption spectra were similar to those of Fe SODs reported previously in the literature (14, 18, 24, 27, 30). The  $E_{1\text{cm}}^{1\%}$  at 278 nm was 18.3, corresponding to a molar extinction coefficient of  $8.4 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$ .

**Metal Analysis.** The enzyme was dialyzed extensively against 5 mM K-phosphate (pH 7.8), containing 0.5 mM EDTA, followed by buffer lacking EDTA and then subjected to atomic absorption analysis. A value of  $0.75 \pm 0.2$  atoms of iron per dimer was obtained. Less than 0.3 atoms of zinc per dimer was found, while manganese was observed to be present in a concentration of less than 0.07 atoms per dimer. Fe SODs with contaminating amounts of zinc have been reported previously (15).

**Isoelectric Point and Amino Acid Analysis.** An analysis of protein stains after isoelectric focusing showed one spot at a pH value of 4.8. This value is in the range of values reported for other Fe SODs (18, 19, 27).

Table II shows the amino acid composition of Fe SOD purified from *N. luteum*. For comparison, the amino acid compositions of Fe SOD isolated from *Plectonema boryanum* (24), *B. campestris* (27), *Escherichia coli* (30), *Spirulina platensis* (18), and *Euglena gracilis* (14) are included. Based upon comparative amino acid

content, there do not appear to be any major differences among the amino acid content of Fe SOD isolated from *Nuphar* and those iron enzymes reported in the literature. However, *Nuphar* appears to contain more serine, glycine, and methionine than do the other enzymes listed and less threonine, alanine, and valine on average.

As a further test of analogy between enzymes, a simple statistical method of comparing the sum of the squares of the difference between amino acids has been devised (20). The value obtained,  $\Delta Q$ , reveals relative sequence homology between proteins. Table III gives the  $\Delta Q$  values computed for 18 amino acids of eight different Fe SODs. The lower the number, the smaller the difference in amino acid content and, hence, the greater the relatedness. Based upon the values for  $\Delta Q$ , *Nuphar* Fe SOD is more closely related to that of the land plant *Brassica* than it is to any other iron enzyme. The *Brassica* enzyme is related to *Plectonema* and *Escherichia*. Neither of the two proteins appear to be related to the two *Euglena* Fe SOD isozymes. Therefore, the amino acid sequences of Fe SOD from *Nuphar* and *Brassica* appear to be related to each other but not to any of the Fe SOD isozymes from *Euglena* (14).

**Inhibitors of *Nuphar* Fe SOD.** Table IV shows the inhibitory effects of several compounds on *Nuphar* Fe SOD. SOD was incubated for 15 min with a particular compound in 50 mM K-phosphate (pH 7.8) and 0.1 mM EDTA. An aliquot of 10  $\mu\text{l}$  was then taken to determine SOD activity. Controls were routinely performed to ensure that the inhibitory compound added along with the enzyme did not interfere with the SOD assay. Like *Euglena* Fe SOD, as reported by Kanematsu and Asada (14), *Nuphar* Fe SOD was inactivated completely by incubation with 6 M guanidine-HCl. The addition of KCN, as expected, did not yield any inhibition. Hydrogen peroxide, at a concentration of 2 mM, led to about a 60% inhibition after 15 min of incubation, in agreement with previous studies (7). The addition of 10 mM azide caused about a 40% inhibition in *Nuphar* Fe SOD. This is in agreement with the findings for *Euglena* Fe SOD which was observed to be inhibited only 50% by azide (14).

Table II. Comparative Amino Acid Composition of Iron-Containing Superoxide Dismutases

| Amino Acid              | Residues <sup>a</sup> |   |                              |                               |                            |                     |                             |
|-------------------------|-----------------------|---|------------------------------|-------------------------------|----------------------------|---------------------|-----------------------------|
|                         | <i>N. luteum</i>      | <i>B. campestris</i><br>(27) <sup>d</sup> | <i>E. gracilis</i> I<br>(14) | <i>E. gracilis</i> II<br>(14) | <i>P. boryanum</i><br>(24) | <i>E. coli</i> (30) | <i>S. platensis</i><br>(18) |
|                         | per mole subunit      |   |                              |                               |                            |                     |                             |
| Lysine                  | 12                    | 11  | 4                            | 4                             | 12                         | 10                  | 10                          |
| Histidine               | 5                     | 4   | 7                            | 6                             | 5                          | 6                   | 6                           |
| Arginine                | 6                     | 5   | 10                           | 9                             | 2                          | 4                   | 0                           |
| Aspartic acid           | 23                    | 18  | 19                           | 20                            | 23                         | 22                  | 26                          |
| Threonine               | 6                     | 9   | 11                           | 11                            | 10                         | 13                  | 10                          |
| Serine                  | 15                    | 11  | 5                            | 5                             | 9                          | 10                  | 12                          |
| Glutamic acid           | 19                    | 16  | 22                           | 22                            | 16                         | 16                  | 17                          |
| Proline                 | 9                     | 10  | 13                           | 13                            | 10                         | 9                   | 8                           |
| Glycine                 | 22                    | 16  | 20                           | 20                            | 15                         | 16                  | 13                          |
| Alanine                 | 20                    | 17  | 26                           | 27                            | 24                         | 26                  | 25                          |
| Cysteine <sup>b</sup>   | 2                     | 2   | 2                            | 2                             | 1                          | 1                   | 1                           |
| Valine                  | 5                     | 11  | 11                           | 11                            | 11                         | 11                  | 9                           |
| Methionine              | 5                     | 1   | 3                            | 4                             | 2                          | 0                   | 3                           |
| Isoleucine              | 4                     | 6   | 6                            | 6                             | 3                          | 8                   | 7                           |
| Leucine                 | 16                    | 14  | 16                           | 17                            | 17                         | 15                  | 16                          |
| Tyrosine                | 6                     | 6   | 7                            | 7                             | 5                          | 6                   | 5                           |
| Phenylalanine           | 9                     | 7   | 10                           | 11                            | 13                         | 10                  | 14                          |
| Tryptophan <sup>c</sup> | 7                     | 4   | 8                            | 7                             | 6                          | 3                   | 3                           |

<sup>a</sup> Values given to the nearest integer.

<sup>b</sup> Determined as S-sulfocysteine after reduction with dithiothreitol and oxidation with sodium tetrathionate (13).

<sup>c</sup> Determined spectrophotometrically in 6 M guanidine HCl (9).

<sup>d</sup> Numbers in parentheses refer to references in "Literature Cited."

Table III. Comparison of  $\Delta Q$  Values of Fe SOD

Values for  $\Delta Q$  were calculated from the amino acid compositions of Fe SODs reported in this paper and in previous literature. The following equation was used (20):

$$\Delta Q = \sum_j (x_{ij} - X_j)^2$$

where the subscripts  $i$  and  $j$  indicate the enzymes compared, and  $X_j$  is the amino acid content, in mole percent content of a given amino acid of type  $j$ . In calculating  $\Delta Q$  values, 18 amino acids were used. Numbers in parenthesis refer to literature from which data was taken.

|                                   | (a) | (b) | (c) | (d) | (e) | (f) |
|-----------------------------------|-----|-----|-----|-----|-----|-----|
| (a) <i>Nuphar</i>                 |     |     |     |     |     |     |
| (b) <i>Brassica</i> (27)          | 39  |     |     |     |     |     |
| (c) <i>Plectonema</i> (24)        | 59  | 36  |     |     |     |     |
| (d) <i>Escherichia</i> (30)       | 74  | 30  | 22  |     |     |     |
| (e) <i>Spirulina</i> (18)         | 74  | 60  | 20  | 26  |     |     |
| (f) <i>Euglena</i> Fe SOD I (14)  | 88  | 59  | 71  | 58  | 105 |     |
| (g) <i>Euglena</i> Fe SOD II (14) | 85  | 59  | 61  | 51  | 91  | 2   |

Table IV. Effect of Various Compounds on *Nuphar* Fe SOD Activity

SOD was incubated at 25°C for 15 min in a buffer containing 50 mM K-phosphate (pH 7.8), 0.1 mM EDTA, and the compound indicated.

| Compound                             | Inhibition |
|--------------------------------------|------------|
|                                      | %          |
| NaN <sub>3</sub> , 10 mM             | 41         |
| Guanidine HCl, 6 M                   | 100        |
| H <sub>2</sub> O <sub>2</sub> , 2 mM | 64         |
| KCN, 1 mM                            | 0          |




FIG. 3. Ouchterlony double immunodiffusion of *Nuphar* SOD antibody and antigens. Center well, Partially purified immunoglobulin to *Nuphar* iron SOD (~320  $\mu$ g of protein). Other wells contain: A, Purified Fe SOD, from *Nuphar* (3.5  $\mu$ g); B, purified Fe SOD from *Brassica*, (4.6  $\mu$ g); C, crude extract from *Nuphar* leaves, (4 units SOD activity); and D, crude extract from *E. gracilis* (6 units SOD activity).

**Cross-Reactivities of Antigens Prepared to SODs.** Antibody to *Nuphar* Fe SOD was prepared from the serum of rabbits given injections of purified protein. Figure 3 shows the results of a typical Ouchterlony immunodiffusion experiment, in which antisera to *Nuphar* Fe SOD was placed in the center well, and purified *Nuphar* Fe SOD, as well as crude extracts from *Nuphar*, *Brassica*, and *Euglena*, were placed in the side wells. As can be seen, a precipitin line appeared between antisera and purified *Nuphar* Fe SOD, as well as between antisera and *Nuphar* crude extracts. *Brassica* and *Euglena* crude extracts did not cross-react with

*Nuphar* antibody.

Table V is a tabulation of some purified SODs and crude extracts which were examined for cross-reactivity to partially purified immunoglobulin to *Nuphar* Fe SOD. Only purified Fe SOD or crude extract from *Nuphar* cross-reacted with *Nuphar* Fe SOD antibody. None of the other plant or algal crude extracts showed any reaction. We conclude, therefore, that *Nuphar* Fe SOD does not contain antigenic sites in common with other Fe SODs. Table V also shows the results of cross-reactivity studies in which antibody prepared to wheat germ Cu-Zn SOD was tested for cross-reactivity against the purified iron enzyme, as well as against crude extracts of plant and algae. As expected, *Nuphar* Fe SOD did not cross-react with antisera to wheat germ Cu/Zn SOD. The wheat germ Cu-Zn antibody did show cross-reactivity against crude extracts from *Brassica*, thereby indicating a common antigenic determinant of Cu-Zn SODs from mustard and wheat germ. Asada *et al.* (1, 3) have previously shown that Cu-Zn SODs from land plants, including mosses and ferns, have antigens in common with spinach enzyme but not with Cu-Zn SOD isolated from bovine erythrocytes. Our finding of common determinants for *Brassica* and wheat germ agree with these earlier findings.

## DISCUSSION

Water lily Fe SOD, reported on in this paper, is similar in mol wt, subunit size, absorption spectra, and metal analysis to Fe SODs isolated from other sources (14, 18, 24, 27, 30). However, antibody prepared to water lily Fe SOD does not cross-react with purified Fe SODs or crude extracts from procaryotic or eucaryotic organisms known to contain Fe SOD. This implies an immunological uniqueness to the iron enzyme from water lily. Insight into sequence homology can be gained by analysis of least squares differences between amino acids of the iron isozyme reported in this paper and those SODs reported on in the literature (Table III). Fe SOD from water lily appears more closely related to Fe SOD from the eucaryotic land plant *B. campestris* than to many

Table V. Antigenic Cross-Reactivity of *Nuphar* Fe SOD Antibody and Wheat Germ Cu-Zn SOD Antibody to Plant Proteins and Cell-Free Extracts

Ouchterlony double immunodiffusion assay was used to test cross-reactivity. Immunoglobulin (~320  $\mu$ g) was placed in the center well and purified proteins or cell-free extracts in the outer wells. Formation of a precipitin line between a center well and side well indicates antigenic cross-reactivity (+).

| Species                     | Antibody             |                      |
|-----------------------------|----------------------|----------------------|
|                             | <i>Nuphar</i> Fe SOD | Wheat germ Cu-Zn SOD |
| <b>Purified Proteins</b>    |                      |                      |
| <i>Nuphar</i> Fe SOD        | +                    | -                    |
| <i>Brassica</i> Fe SOD      | -                    | -                    |
| Wheat germ Cu-Zn SOD        | -                    | +                    |
| <b>Cell-Free Extracts</b>   |                      |                      |
| <i>N. luteum</i>            | +                    | -                    |
| <i>Nympha odorata</i>       | -                    | -                    |
| <i>B. campestris</i>        | -                    | +                    |
| <i>Gingko biloba</i>        | -                    | -                    |
| Wheat germ                  | -                    | +                    |
| <i>E. coli</i>              | -                    | -                    |
| <i>E. gracilis</i>          | -                    | -                    |
| <i>Navicula pelliculosa</i> | -                    | -                    |
| <i>P. boryanum</i>          | -                    | -                    |
| <i>Scenedesmus</i> sp.      | -                    | -                    |
| <i>Nostoc muscorum</i>      | -                    | -                    |
| <i>Chlorella</i> sp.        | -                    | -                    |

of the blue-green and green algal Fe SODs, including those from *Euglena*. Clearly, work is needed to elucidate amino-acid sequences of the higher land plant Fe SODs.

We have previously reported that Fe SOD is a chloroplast stromal enzyme in *Brassica* leaves (28), yet absent in *Brassica* mitochondria (29). We intend to investigate further the subcellular localization of Fe SOD in water lily by using fluorescently labeled antibody.

Water lily, *N. luteum*, is a unique plant in terms of SODs. The plant is apparently devoid of Cu-Zn SOD but contains the iron isozyme (6). Several questions are raised by this anomaly. First, why is there an apparent lack of the Cu-Zn SOD? A possible explanation is that, in a relatively anaerobic pond bottom in which this organism is rooted, most copper would be found in the form of insoluble copper sulfide. As a result, copper might not be as readily absorbed and utilized as it is in land plants. A metal analysis of *Nuphar*, however, shows that its total copper content is not significantly different from that of land plants (M. L. Salin and S. M. Bridges, unpublished). Second, assuming an absence of Cu-Zn SOD in *Nuphar* and a necessity for dismutase activity (10), why has this plant expressed an iron enzyme as opposed to enhancing production of an already present manganese-containing SOD? Manganese-containing SODs are found in procaryotic and eucaryotic organisms (2, 10, 22) and have been shown to be dramatically responsive to environmental changes (11, 12). In contrast, Fe SOD was not responsive to enhancement of synthesis by environmental pressures (11, 12). A possible explanation for the occurrence of Fe SOD might be accounted for by an independent occurrence of gene transfer from a bacteria or algae to *Nuphar* or its predecessor. A case of gene transfer between a host fish and its symbiotic bacteria has recently been reported (21). Finally, we might wonder whether the genes for the Cu-Zn SOD are present in *Nuphar*, yet not expressed. Alternately, we might ask if genes for Fe SOD in other eucaryotes may be present, yet not expressed. Answers to many of these questions might be found by searching for potential genes coding for Fe SOD and Cu-Zn SOD in organisms where they are not expressed.

**Acknowledgments**—The authors would like to thank Dr. Sidney McDaniel, Department of Biological Sciences, Mississippi State University, for collecting much of the plant material. They would also like to acknowledge the assistance of Michael Aide for metal analysis by atomic absorption, William E. Poe for performing amino acid analysis, and Joseph K. Wong for assistance in obtaining antibody to Fe SOD.

#### LITERATURE CITED

- ASADA K, S KANEMATSU, K UCHIDA 1977 Superoxide dismutases in photosynthetic organisms: Absence of the cuprozinc enzyme in eukaryotic algae. *Arch Biochem Biophys* 179: 243-256
- ASADA K, S KANEMATSU 1978 Distribution of cuprozinc, manganic, and ferric superoxide dismutases in plants and fungi: an evolutionary aspect. In H Matsubara, T Yamanaka, eds, *Evolution of Protein Molecules*. Japan Scientific Societies Press, Tokyo, pp 361-372
- ASADA K, S KANEMATSU, S OKADA, T HAYAKAWA 1980 Phylogenetic distribution of three types of superoxide dismutase in organisms and cell organelles. *In JV Bannister, HAO Hill, eds, Chemical and Biochemical Aspects of Superoxide and Superoxide Dismutase*. Elsevier, New York, pp 136-153
- BEAUCHAMP CO, I FRIDOVICH 1971 Superoxide dismutase: improved assays and an assay applicable to acrylamide gel. *Anal Biochem* 44: 276-287
- BEAUCHAMP CO, I FRIDOVICH 1973 Isozymes of superoxide dismutase from wheat germ. *Biochim Biophys Acta* 317: 50-64
- BRIDGES SM, ML SALIN 1981 Distribution of iron-containing superoxide dismutase in vascular plants. *Plant Physiol* 68: 275-278
- BRITTON L, DP MALINOWSKI, I FRIDOVICH 1978 Superoxide dismutase and oxygen metabolism in *Streptococcus faecalis* and comparisons with other organisms. *J Bacteriol* 134: 229-236
- DAVIS BJ 1964 Disc gel electrophoresis. *Ann NY Acad Sci* 121: 404-427
- EDELHOCH H 1967 Spectroscopic determination of tryptophan and tyrosine in proteins. *Biochemistry* 6: 1948-1954
- FRIDOVICH I 1975 Superoxide dismutases. *Annu Rev Biochem* 44: 147-159
- HASSAN HM, I FRIDOVICH 1977 Enzymatic defenses against the toxicity of oxygen and of streptonigrin in *Escherichia Coli*. *J Bacteriol* 129: 1574-1583
- HASSAN HM, I FRIDOVICH 1977 Regulation of the synthesis of superoxide dismutase in *Escherichia coli*: Induction by methyl viologen. *J Biol Chem* 252: 7667-7672
- INGLIS AS, TY LIU 1970 The stability of cysteine and cystine during acid hydrolysis of proteins and peptides. *J Biol Chem* 245: 112-116
- KANEMATSU S, K ASADA 1979 Ferric and manganic superoxide dismutases in *Euglena gracilis*. *Arch Biochem Biophys* 195: 535-545
- KIRBY TW, JR LANCASTER JR, I FRIDOVICH 1981 Isolation and characterization of the iron-containing superoxide dismutase of *Methanobacterium bryantii*. *Arch Biochem Biophys* 210: 140-148
- LAEMMLI UK 1970 Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (Lond)* 227: 680-685
- LOWRY OH, NJ ROSEBROUGH, AL FARR, RJ RANDALL 1951 Protein measurement with the Folin phenol reagent. *J Biol Chem* 193: 265-275
- LUMSDEN J, R CAMMACK, DO HALL 1976 Purification and physicochemical properties of superoxide dismutase from two photosynthetic microorganisms. *Biochim Biophys Acta* 438: 380-392
- LUMSDEN J, L HENRY, DO HALL 1977 Superoxide dismutase in photosynthetic organisms. In AM Michelson, JM McCord, I Fridovich, eds, *Superoxide and Superoxide Dismutases*. Academic Press, New York, pp 437-450
- MARCHALONIS JJ, JK WELTMAN 1971 Relatedness among proteins: A new method of estimation and its application to immunoglobulins. *Comp Biochem Physiol* 38: 609-625
- MARTIN JP, I FRIDOVICH 1981 Evidence for a natural gene transfer from the ponyfish to its bioluminescent bacterial symbiont, *Photobacter leiognathi*: the close relationship between bacteriocuprein and the copper-zinc superoxide dismutase of teleost fishes. *J Biol Chem* 256: 6080-6089
- MCCORD JM 1979 Superoxide dismutases: occurrence, structure, function and evolution. In M Rattazzi, J Scandalios, GS Whitt, eds, *Current Topics in Biological and Medical Research*, Vol 3, Liss, New York, pp 1-21
- MCCORD JM, I FRIDOVICH 1969 Superoxide dismutase: an enzymatic function for erythropoietin. *J Biol Chem* 244: 6049-6055
- MISRA HP, BB KEELE JR 1975 The purification and properties of superoxide dismutase from a blue-green alga. *Biochim Biophys Acta* 379: 418-425
- MURPHY JB, MW KIES 1960 Note on spectrophotometric determination of proteins in dilute solutions. *Biochim Biophys Acta* 45: 382-384
- OUCHTERLONY O 1968 *Handbook of Immunodiffusion and Immunoelectrophoresis*. Ann Arbor Science Publishers, Ann Arbor, MI
- SALIN ML, SM BRIDGES 1980 Isolation and characterization of an iron-containing superoxide dismutase from a eucaryote, *Brassica campestris*. *Arch Biochem Biophys* 201: 369-374
- SALIN ML, SM BRIDGES 1980 Localization of superoxide dismutase in chloroplasts from *Brassica campestris*. *Z Pflanzenphysiol* 99: 37-45
- SALIN ML, SM BRIDGES 1981 Absence of the iron-containing superoxide dismutase in mitochondria from *Brassica campestris*. *Biochem J* 195: 229-233
- YOST FJ, I FRIDOVICH 1973 An iron-containing superoxide dismutase from *Escherichia coli*. *J Biol Chem* 248: 4905-4908