

Isolation and Characterization of *Trichoderma* spp. for Antagonistic Activity Against Root Rot and Foliar Pathogens

Krishna Kumar · N. Amaresan · S. Bhagat ·
K. Madhuri · R. C. Srivastava

Received: 22 June 2010 / Accepted: 23 November 2010 / Published online: 13 August 2011
© Association of Microbiologists of India 2011

Abstract *Trichoderma*, soil-borne filamentous fungi, are capable of parasitising several plant pathogenic fungi. Twelve isolates of *Trichoderma* spp. isolated from different locations of South Andaman were characterized for their cultural, morphological and antagonistic activity against soil borne and foliar borne pathogens. The sequencing of these isolates showed seven different species. The isolates revealed differential reaction patterns against the test pathogens viz., *Sclerotium rolfsii*, *Colletotrichum gloeosporioides* and *C. capsici*. However, the isolates, TND1, TWN1, TWC1, TGD1 and TSD1 were most effective in percentage inhibition of mycelial growth of test pathogens. Significant chitinase and β -1,3-glucanase activities of all *Trichoderma* isolates has been recorded in growth medium. *T. viride* was found with highest chitinase whereas *T. harzianum* was recorded with highest β -1,3-glucanase activities.

Keywords *Trichoderma* · Spice · Biocontrol · Andaman and Nicobar Islands

K. Kumar (✉) · N. Amaresan · S. Bhagat · K. Madhuri ·
R. C. Srivastava
Division of Field Crops, Central Agricultural Research Institute,
Port Blair 744 105, Andaman and Nicobar Islands, India
e-mail: krishnakingsh2000@yahoo.co.in

N. Amaresan
e-mail: amarcool82@yahoo.co.in

S. Bhagat
e-mail: sombhagat73@rediffmail.com

K. Madhuri
e-mail: madhurikbiotech@gmail.com

R. C. Srivastava
e-mail: ramesh_cari@yahoo.co.in

Introduction

The anamorphic fungal genus *Trichoderma* (Hypocreales, Ascomycota) is a cosmopolitan soil-borne fungi frequently found on decaying wood [1, 2] of which some are economically important producers of industrial enzymes (*Trichoderma reesei*) [3], antibiotics [4] and have been used as biocontrol (i.e., *T. harzianum*, *T. atroviride* and *T. asperellum*) agents against plant pathogens [5]. *Trichoderma* spp. are among the most frequently isolated soil fungi and present in plant root systems [6]. However, there is still considerable interest in finding more efficient mycoparasitic fungi especially within *Trichoderma* spp., which differ considerably with respect to their biocontrol effectiveness. It is important to isolate *Trichoderma* spp. having potentially higher antagonistic efficiency by the selection of isolates with high potential to secrete extra cellular lytic enzymes chitinase and β -1,3-glucanase. The lytic enzymes break down cell wall polysaccharides into short oligomers and by this way facilitate the hyperparasite to penetrate into the cytoplasm of the target fungi [7]. The aim of this study was screening of *Trichoderma* spp. for their antagonistic ability by dual cultures as well as their capability of producing lytic enzymes against the test pathogens.

Materials and Methods

Soil Samples and Isolation

Soil samples were collected from different ecological habitat of spice crops of district, South Andaman, Andaman and Nicobar Islands (India) for the isolation of *Trichoderma* spp. (Table 1). Samples were brought to

laboratory and stored at 4°C until used. Five-fold serial dilutions of each soil samples were prepared in sterilized distilled water and 0.5 ml diluted sample was poured on the surface of *Trichoderma* Specific Medium (TSM) [8]. Plates were incubated at $28 \pm 2^\circ\text{C}$ for 96 h. Morphologically different colonies appearing on the plates were purified in the Potato Dextrose Agar (PDA) (HiMedia, India). The purified isolates were preserved at 4°C and used during the course of study.

Phenotype Characters of the *Trichoderma* Isolates

The morphological and cultural characteristics of 12 isolates of *Trichoderma* were studied in four different media viz., OMA, CMD, PDA and TSM following the protocol of Samules et al. [9]. Mycelial discs (6 mm) of young growing culture of respective isolates of *Trichoderma* was inoculated in the periphery of the Petri plates containing above said media and incubated at $28 \pm 2^\circ\text{C}$ for one week. Colony radius was measured at 24, 48 and 72 h. each growth rate experiment was repeated three times in triplicate and the results were averaged for each isolate. Additional characters include presence of pigments, green conidia, odor and colony appearance were also noted.

Morphological observations were recorded from cultures grown on PDA plates. The following characters were measured; Phialide width at the widest pint, phialide length, conidium length and width and presence of chlamyospores. Each character was measured from water after initial wetting in 3% KOH for each isolates.

Fungal Growth Conditions and DNA Extraction

Cultures were maintained on PDA at 25°C was grown in potato dextrose broth for 3 days. Mycelial mat was collected on filter paper, washed with distilled water for 2–3 times, frozen and were used for DNA extraction. Genomic DNA was extracted using method of Raeder and Broda [10]. DNA resuspended in 50 µl of TE buffer and quantified by use of ethidium bromide fluorescence.

PCR Amplification and Sequencing of Amplification Products

Primers ITS1 (5'-TCC GTA GGT GAA CCT GCG G-3') and ITS4 (5'-TCC TCC GCT TAT TGA TAT GC-3') described by White et al. [11] were used to amplify a fragment of rDNA including ITS1 and ITS2 and the 5.8S rDNA gene. PCR amplifications were performed in a total volume of 50 µl by mixing 20 ng of the template DNA with 2.5 mM concentrations of each deoxynucleotide triphosphate, 1 µM concentrations of each primer and 3 U of Taq DNA polymerase in 10× Taq buffer A (GenEi). These reactions were subjected to initial denaturations of 1 min at 95°C, followed by 35 cycles of 1 min at 95°C, 30 s at 55°C and 1.5 min at 72°C, with a final extension of 10 min at 72°C using GeneAmp® PCR system 9700 (Applied Biosystems). The PCR products were resolved using a 1% agarose gel. Sequencing of purified PCR product was undertaken by SAP services, GeneI, Bangalore using ITS1 (forward primer).

Table 1 *Trichoderma* isolates studied, their identification and host

S. No.	Isolate designation ^a	Crop name ^b	Place of collection	Species identified	Accession number
1	TSD1	Cinnamon	Sippighat	<i>Trichoderma inhamatum</i>	GQ426033
2	TWN1	Nutmeg	Wandoor	<i>T. harzianum</i>	GQ426034
3	TGN1	Nutmeg	Guptapara	<i>T. harzianum</i>	GQ426035
4	TWD1	Cinnamon	Wandoor	<i>T. asperellum</i>	GQ426036
5	TND1	Cinnamon	New Mangulton	<i>T. erinaceum</i>	GQ426037
6	TJP1	Pepper	Jirkatang	<i>T. harzianum</i>	GQ426038
7	TWC2	Clove	Wandoor	<i>T. harzianum</i>	GQ426039
8	TMC1	Clove	Manjery	<i>T. longibrachiatum</i>	GQ426040
9	TWP1	Pepper	Wandoor	<i>Trichoderma</i> sp.	GQ426041
10	TCC1	Cinnamon	Calicut	<i>T. ovalisporum</i>	GQ426042
11	TGD1	Cinnamon	Guptapara	<i>T. viride</i>	GQ426043
12	TBC1	Clove	Brichgunj	<i>T. brevicompactum</i>	GQ426044

^a Coding system of isolates *T* *Trichoderma*, *S* place of collection and *D* crop name

^b Rhizosphere soil of various spice crops

Phylogenetic Analysis

ITS gene sequences of the isolates were compared with ITS sequences available by the BLAST search in the NCBI, GenBank database (<http://www.ncbi.nlm.nih.gov>). Multiple sequence alignment was performed using Clustal X (1.1). The method of Jukes and Cantor [12] was used to calculate evolutionary distances. Phylogenetic dendrogram was constructed by the neighbour-joining method and tree topologies were evaluated by performing bootstrap analysis of 100 data sets using MEGA 3.1 (Molecular Evolutionary Genetic Analysis). The sequences were submitted to GenBank under the following accession numbers GQ426033–GQ426044 (12 sequences).

Antagonistic Activity of *Trichoderma* Isolates

The dual culture technique described by Morton and Stroube [13] was used to test the antagonistic ability of *Trichoderma* spp. against soil borne pathogen *S. rolfsii* and foliar borne pathogens viz., *C. capsici* and *C. gloeosporioides*. The pathogens and *Trichoderma* were grown on PDA for a week at room temperature ($28 \pm 2^\circ\text{C}$). Small blocks of the target fungi (*C. capsici* and *C. gloeosporioides*) cut from the periphery were transferred to the Petri dish previously poured with PDA. After 2 days of *Colletotrichum* spp. growth, the *Trichoderma* spp. was transferred aseptically in the same plate of opposite end and were incubated at room temperature with alternate light and darkness for 7 days and observed periodically. For *S. rolfsii* both pathogen and *Trichoderma* were inoculated at the same time. The experiment was replicated thrice and percent growth inhibition was calculated by the formula of $I = (C - T)/C \times 100$, where C is mycelial growth in control plate, T is mycelial growth in test organisms inoculated plate and I is inhibition of mycelial growth.

Assay for Extra Cellular Enzyme Activity

Twelve isolates of *Trichoderma* were separately inoculated into 100 ml broth media with four combinations viz., Czapek Dox Broth alone, CDB + mycelial powder of *S. rolfsii* (10 g), CDB + mycelial powder of *C. gloeosporioides* (10 g) and CDB + mycelial powder of *C. capsici* (10 g) in Erlenmeyer flasks and incubated at $28 \pm 1^\circ\text{C}$ for 7 days with intermittent shaking at 125 rpm twice a day. The culture filtrate each isolate was harvested, filtered through the Whatman Filter Paper 42, centrifuged and assayed for chitinase and β -1,3-glucanase enzyme activities immediately.

β -1,3-Glucanase Activity

For assay of β -1,3-glucanase enzyme, 0.5 ml laminarin, 1.0 ml of 0.05 M citrate buffer (pH 4.8) and 0.5 ml culture filtrate was mixed and incubated at 40°C for 60 min. An equal volume of dinitrosalicylic acid reagent was added to the reaction mixture and warmed in boiling water for 15 min. The absorbance of reaction mixture was measured at 575 nm in a spectrophotometer and compared with standard graph drawn by following the same procedure but using different concentrations of glucose instead of culture filtrate. The quantity of reducing sugar was calculated from the glucose standards used in the assay and activity of β -1,3-glucanase was expressed in nkat/ml. One nkat corresponds to the release of 1 nmol glucose equivalent per second.

Chitinase Activity

A mixture of 0.5 ml culture filtrate, 0.5 ml suspension of colloidal chitin and 1.0 ml of McIlvaine's buffer (pH 4.0) was mixed and incubated at 37°C for 2 h in a water bath with constant shaking. The reaction was stopped by boiling 3 min in heated water bath. 3 ml potassium ferricyanide reagent was added and warmed in boiling water for 15 min. The amount of N-acetyl glucosamine (NAG) released was estimated following the methods of Reissing et al. [14]. The absorbance of reaction mixture was measured in a spectrophotometer at 420 nm. The amount of reducing sugar released was calculated from standard curves for NAG and chitinase enzyme activity was expressed in pkat (pmol/s) per millilitre.

Statistical Analysis

Statistical analysis were performed with the Agres and Agdata using completely randomized analyses of variances (ANOVA) was used to compare the biocontrol efficacy of *Trichoderma* isolates and means separated by Fisher's protected least significant difference (LSD). The significance of effects of *Trichoderma* on growth characteristics was determined by the magnitude of the F value ($P = 0.05$).

Results and Discussion

Morphological Characterization

Based on the observation of the conidia, phialides, colony texture, chlamyospore, conidiophore morphology the isolates were grouped into section Pachybasium B (clade Lixii/catoptron), *Trichoderma* (Rufa), *Trichoderma*

(Pachybasium A), Longibrachiatum and Pachybasium B (Lutea). Section Pachybasium consists of the isolates TSD1, TWN1, TGN1, TJPI and TWC1 where the conidiophores were much branched, which arise in distinct and continues ring like zones. The main branches mostly in groups of 2–3 and stand at right angle to the bearer and their length increased with the distance from the tip of the main branch which gave a conical or pyramidal appearance. In *Trichoderma* section (Rufa) (TWD1, TGD1 and TCC1) the hyphae, branched, smooth walled and colourless. The conidiophores were observed as less complicated and formed aerial hyphae. They produced smaller branches and ultimately a conifer-like branching system is formed. In Pachybasium B (Lutea) (TBC1) the conidiophores are smooth walled, verticillately branched in the Pachybasium type pattern. *Trichoderma* (section of Pachybasium A) consists of *T. asperellum* seen the conidiophores have terminating 2 or more phialides and primary branching arising at nearly 90 degrees to the main axis. In section Longibrachiatum (*H. orientalis*), the conidiophores were complicated and progressively longer, often paired, secondary branches. Phialides arising directly from secondary branches, typically not in whorls. Using the key of Rifai [15] in the earlier most of the strains isolated in these regions were identified as *T. harzianum*, *T. hamatum* and *T. viride* [16]. In fact, apart from *T. harzianum*, which, nevertheless accounted for the majority of isolates, a very interesting finding of this study was the coexistence of several species of *Trichoderma* (e.g. *T. harzianum*, *T. erinaceum*, *T. asperellum*, *T. ovalisporum*, *T. viride* and *T. brevicompactum*) in the Bay Island ecosystem.

Molecular Identification of *Trichoderma* Species

Sequence analysis of twelve isolates was done to confirm species identity, which initially has been done based solely on morphological parameters. Comparison of oligonucleotide fragments of rDNA sequences, which included the 5.8S gene and the flanking ITS1 and ITS2 regions, with reference sequences from public databases, showed that they were very similar.

Phylogenetic Analysis

The phylogenetic tree obtained by sequence analysis of ITS1 and ITS4 of 12 of our isolate sequences and the sequence of 22 *Trichoderma* spp. obtained from NCBI, GenBank is represented in Fig. 1. The ITS sequence was chosen for this analysis because it has been showed to be more informative with various sections of the genus *Trichoderma* [17–19]. Bootstrap analysis with 100 bootstrap replications demonstrated three major branches. On the basis of the bootstrap values, 34 *Trichoderma* spp. could be


divided into 5 groups. Group 1, the *T. harzianum* complex includes 6 of our isolates. This group is supported by bootstrap value of 92% contains 3 subgroups supported by bootstrap values more than 90%. Group II (*Lutea*) includes 3 strains representing *T. brevicompactum*. This cluster is supported by a bootstrap value of 83%. Group IV (*Rufa*) includes 3 different isolates includes *T. viride*, *T. ovalisporum* and *T. erinaceum*, this cluster is supported by more than 60% of bootstrap values. In the case of group III cluster which, consists of section *Longibrachiatum* (*H. orientalis*) and *Pachybasium* A (*T. asperellum*). In this analysis no grouping among the isolates was observed, and did not show any bootstrap supports suggests instability of the clade.

Antagonistic Activity and Enzyme Production of *Trichoderma* Isolates

Results from the dual culture test showed that all isolates of *Trichoderma* inhibited mycelial growth of *S. rolfsii* more than 50%. Two isolates TGD1 and TWN1 (*T. viride* and *T. harzianum*) showed statistically significant inhibition of mycelial growth (76.3%) over control against *S. rolfsii* (Table 2). The isolates TND1, TBC1, TCC1, TGN1 and TWC2 (*T. erinaceum*, *T. brevicompactum*, *T. ovalisporum*, *H. lixii* and *T. harzianum*) were next best isolates in inhibiting the growth of pathogen i.e., 73.3, 70.8, 72.5, 70.8 and 72.1 per cent over the control, respectively. Isolates TSD1 (*T. inhamatum*), TJPI (*H. lixii*) and TND1 (*T. erinaceum*) showed statistically significant inhibition of mycelial growth of *C. gloeosporioides* in dual culture (Table 2). The growth inhibitions of these isolates were 45.6, 43.3 and 42.2% respectively over the control. Similarly in the case of *C. capsici*, isolates TWC2 (*T. harzianum*) and TSD1 (*T. inhamatum*) showed significant inhibition of mycelial growth.

The production of chitinase and β -1,3-glucanase are summarized in Tables 3 and 4. All isolates showed significantly higher chitinase and β -1,3-glucanase activities with supplement of different carbon sources as substrates in the basal media. The highest chitinase enzyme activity was recorded with *T. viride* (42.0 U) in CDB + mycelial powder of *S. rolfsii*, followed by *T. harzianum* (40.3 U), *T. erinaceum* (37.1 U), *H. lixii* (36.4 U), *T. brevicompactum* (35.1 U), *T. ovalisporum* (34.2 U) and *Trichoderma* sp. (33.2 U). Chitinase activity of test isolates of *Trichoderma* was relatively less when the basal medium was added with mycelial powder of *C. gloeosporioides* and *C. capsici* as compare to *S. rolfsii* whereas highest chitinase activity was recorded with *T. harzianum*. The partial substitution of sucrose by same carbon sources in the basal medium has profound effect as the β -1,3-glucanase activity as compared to basal media (CDB). Highest β -1,3-glucanase enzyme activity was recorded with *T. harzianum* (22.6,

Fig. 1 Phylogram based on the ITS region of the genomic rRNA gene of 12 isolates and 22 representative strains of *Trichoderma* with isolate number indicated in parenthesis. The phylogram was generated after DNA distance based and neighbor-joining analysis of the data. Bootstrap 50% majority-rule consensus tree (1,000 replications), indicated above the nodes, was generated using MEGA 3.1



31.5, 28.6 and 30.2 U) in CDB, CDB + *S. rolsii*, CDB + *C. gloeosporioides* and CDB + *C. capsici* followed by *T. viride*, *T. erinaceum* and *T. inhamatum*. Among all carbon sources as amendment of media, mycelial powder of *S. rolsii* induced higher β -1,3-glucanase and chitinase activity than *C. gloeosporioides* and *C. capsici*.

The application of *Trichoderma* as biocontrol agent has been focused mainly in controlling root and soil-borne diseases while there are few reports on their application in controlling foliar diseases [20]. This stems from the fact that *Trichoderma* sp. are native (residents) in the soil but transients in the phylloplane [21]. The

antagonism by *Trichoderma* spp. against many soil-borne plant pathogens has been established [6, 22, 23]. Strong antagonism by *Trichoderma* spp. against a range of soil-borne plant pathogens has been reported [6, 23]. It is well known that *Trichoderma* spp. have the potential to produce cell wall-degrading enzymes by using the materials that are present in the growth medium [6]. Production of hydrolytic enzymes such as β -1,3-glucanase, chitinase, cellulase and proteinase increased significantly when *Trichoderma* spp. were grown in media supplemented with either autoclaved mycelium or purified host fungal cell walls [7, 24]. Ulhoa and Peberdy [25] found that products of chitin degradation also

Table 2 Antagonistic potential of *Trichoderma* isolates against *S. rolfsii*, *C. gloeosporioides* and *C. capsici*

S. No.	Isolate name	Mycelial growth inhibition* (%)		
		<i>S. rolfsii</i>	<i>C. gloeosporioides</i>	<i>C. capsici</i>
1	<i>T. inhamatum</i>	69.6 ^{bc}	45.6 ^a	48.9 ^{ab}
2	<i>T. longibrachiatum</i>	67.1 ^{cd}	35.6 ^{cd}	42.2 ^{de}
3	<i>T. brevicompactum</i>	70.8 ^{ab}	38.9 ^{bc}	36.7 ^e
4	<i>T. ovalisporum</i>	72.5 ^{abc}	27.8 ^d	43.3 ^{cd}
5	<i>T. viride</i>	76.3 ^a	30.0 ^{cd}	46.7 ^{bc}
6	<i>T. erinaceum</i>	73.3 ^{ab}	42.2 ^{ab}	42.2 ^{cd}
7	<i>T. harzianum</i>	70.8 ^{ab}	35.6 ^{cd}	46.7 ^{bc}
8	<i>T. asperellum</i>	69.6 ^{bc}	40.0 ^{bc}	38.9 ^{de}
9	<i>Trichoderma</i> sp.	62.1 ^d	28.9 ^d	42.2 ^{de}
10	<i>T. harzianum</i>	76.3 ^a	34.4 ^{cd}	45.6 ^{bc}
11	<i>T. harzianum</i>	72.1 ^{ab}	27.8 ^d	50.0 ^a
12	<i>T. harzianum</i>	56.3 ^e	43.3 ^{ab}	46.7 ^{bc}
	Control (mm)	80	30	30
	CD (0.05)	0.4387	0.3025	0.1946
	SEd	0.2126	0.1466	0.0943

Data were analyzed using one-way analysis of variance (values with different lower case letters are significantly different; $P \leq 0.05\%$)

* Values are mean of three replications

Table 3 Chitinase enzyme activity of *Trichoderma* isolates from Andaman and Nicobar Islands (pkat/ml)

Isolate	CDB	CDB + SR	CDB + CG	CDB + CC
<i>T. inhamatum</i>	22.3	32.5	28.4	30.1
<i>T. longibrachiatum</i>	19.2	30.2	25.3	28.1
<i>T. brevicompactum</i>	23.9	35.1	26.4	23.5
<i>T. ovalisporum</i>	22.7	34.2	22.3	19.6
<i>T. viride</i>	28.3	42.0	24.5	27.8
<i>T. erinaceum</i>	23.5	37.1	28.0	28.6
<i>T. asperellum</i>	21.0	22.0	28.0	26.5
<i>T. harzianum</i>	25.1	36.4	26.4	24.8
<i>Trichoderma</i> sp.	20.6	33.2	19.6	21.3
<i>T. harzianum</i>	32.2	40.3	30.2	32.6
<i>T. harzianum</i>	30.5	38.6	29.4	31.2
<i>T. harzianum</i>	20.0	29.6	22.7	20.5
SEd	0.403	0.321	0.369	0.426
CD (0.05)	0.833	0.878	0.763	0.878

Means of four replications; CDB Czapek Dox Broth, S substrate, SR *S. rolfsii*, CG *C. gloeosporioides*, CC *C. capsici*

regulate the chitinase synthesis in *T. harzianum*. Kumar and Gupta [2] reported that cell wall of *M. phaseolina* and *S. rolfsii* is known to have glucan and chitin that should have resulted in the induction of glucanase and chitinase in mycelial mat amended media. High β -1,3-glucanase and chitinase activities were detected in dual culture when *T. harzianum* parasitized *R. solani* and *S. rolfsii* compared with low levels of substrates or in absence of pathogen [26]. In present investigation, it was found that chitinase and β -1,3-glucanase enzyme activities was increased with the substitution of specific carbon source at 1% concentration. These findings is in

accordance with Ulhoa and Peberdy [25] where they suggested that chitinase activity was substrate's concentration dependent above 0.5% (w/v) chitin there was no further synthesis of chitinase in the growth medium by *T. harzianum* was increased up to 1% concentration, whereas β -1,3-glucanase enzyme production was increase up to 1% concentration of laminarin but decreased at higher concentrations. This may be due to the fact that at higher concentration of sugar activity of this enzyme was inhibited. The degree of mycelial inhibition of target pathogen by *T. erinaceum* is notable example as bio-control against pathogens tested.

Table 4 β -1,3-Glucanase enzyme activity of *Trichoderma* isolates from Andaman and Nicobar Islands (nkat/ml)

Isolate	CDB	CDB + SR	CDB + CG	CDB + CC
<i>T. inhamatum</i>	15.2	22.3	24.9	22.5
<i>T. longibrachiatum</i>	17.4	23.5	19.4	21.0
<i>T. brevicompactum</i>	14.8	15.0	17.5	18.6
<i>T. ovalisporum</i>	9.5	18.1	21.2	19.6
<i>T. viride</i>	20.5	24.2	25.5	28.3
<i>T. erinaceum</i>	11.3	29.8	22.3	24.0
<i>T. asperellum</i>	11.5	12.8	16.5	20.0
<i>T. harzianum</i>	13.4	19.3	24.3	25.0
<i>Trichoderma</i> sp.	11.2	16.4	17.6	15.3
<i>T. harzianum</i>	22.6	31.5	28.6	30.2
<i>T. harzianum</i>	19.4	28.1	26.8	27.7
<i>T. harzianum</i>	9.3	17.5	18.7	21.0
SEd	0.372	0.320	0.408	0.406
CD (0.05)	0.768	0.654	0.843	0.837

Means of four replications;
CDB Czapek Dox Broth, SR *S. rolfsii*, CG *C. gloeosporioides*, CC *C. capsici*

Acknowledgments This research was supported by grants from the project “Application of Microorganisms in Agriculture and Allied Sectors” of the NBAIM-ICAR, UP, Mau, India. We thank the Head Division of Field Crops, CARI for constant support and encouragements during the course of study.

References

- Samuels GJ (1996) *Trichoderma*: a review of biology and systematics of the genus. Mycol Res 100:923–935
- Kumar A, Gupta JP (1999) Variation in enzyme activity of tub-econazole tolerant biotypes of *Trichoderma viride*. Ind Phytopathol 52:263–266
- Kubicek CP, Penttila ME (1998) Regulation of production of plant polysaccharide degrading enzymes by *Trichoderma*. In: Harman GE, Kubicek CP (eds) *Trichoderma and Gliocladium enzymes biological control and commercial applications*, vol 2. Taylor and Francis Ltd, London, pp 49–71
- Sivasithamparam K, Ghisalberti EL (1998) Secondary metabolism in *Trichoderma* and *Gliocladium*. In: Kubicek CP, Harman GE (eds) *Trichoderma and Gliocladium basic biology taxonomy and genetics*, vol 1. Taylor and Francis Ltd, London, pp 139–191
- Harman GE (2000) Myth and dogmas of biocontrol; changes in perceptions derived from research on *Trichoderma harzianum* T-22. Plant Dis 84:377–393
- Harman GE, Howell CR, Viterbo A, Chet I, Lorito M (2004) *Trichoderma* species-opportunistic, avirulent plant symbionts. Nat Rev Microbiol 2:43–56
- Cruz L, Pinter-Toro JA, Benitez T, Llobell A (1995) Purification and characterization of an endo- β -1,3-glucanase from *Trichoderma harzianum* that is related to its mycoparasitism. J Bacteriol 177:1864–1877
- Elad Y, Chet I, Henis Y (1982) Degradation of plant pathogenic fungi by *Trichoderma harzianum*. Can J Microbiol 28:719–725
- Samuels GJ, Dodd SL, Gams W, Castlebury LA, Petrini O (2002) *Trichoderma* species associated with the green mold epidemic of commercially grown *Agaricus bisporus*. Mycologia 94:146–170
- Raeder U, Broda P (1985) Rapid preparation of DNA from filamentous fungi. Lett Appl Microbiol 1:17–20
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal DNA for phylogenetics. In: Innes MA, Gelfand DH, Sninsky JJ, White TJ (eds) PCR protocols: a guide to methods and applications. Academic, San Diego, CA, pp 315–322
- Jukes TH, Cantor CR (1969) Evolution of protein molecules. In: Munro HN (ed) Mammalian protein 242 metabolism. Academic, New York, pp 21–132
- Morton DT, Stroube NH (1955) Antagonistic and stimulatory effect of microorganism upon *Sclerotium rolfsii*. Phytopathology 45:419–420
- Reissing JL, Strominger JL, Leloir LF (1955) A modified colorimetric method for estimation of N-acetylamine sugars. J Biol Chem 217:959–966
- Rifai MA (1969) A revision of the genus *Trichoderma*. Mycol Pap 116:1–56
- Bhagat S (2008) Biocontrol potential of *Trichoderma* spp. from Andaman and Nicobar Islands. PhD thesis, BCKV, West Bengal, India
- Kuhls K, Lieckfeldt E, Samuels GJ, Kovacs W, Meyer W, Petrini O, Gams W, Borner T, Kubicek CP (1996) Molecular evidence that the asexual industrial fungus *Trichoderma reesei* is a clonal derivative of the ascomycete *Hypocrea jecorina*. Proc Natl Acad Sci USA 93:7755–7760
- Kuhls K, Lieckfeldt E, Samuels GJ, Meyer WC, Kubicek P, Borner T (1997) Revision of *Trichoderma* sec. *Longibrachiatum* including related teleomorphs based on analysis of ribosomal DNA internal transcribed spacer regions. Mycologia 89:442–460
- Ospina-Giraldo MD, Royse DJ, Thon MR, Chen X, Romaine CP (1998) Phylogenetic relationships of *Trichoderma harzianum* causing mushroom green mold in Europe and North America to other species of *Trichoderma* from world-wide sources. Mycologia 90:76–81
- Blakeman JP, Fokemma NJ (1982) Potential for biological control of plant disease on the phylloplane. Ann Rev Phytopathol 20:167–192
- Bankole SA (1990) Micro flora associated with cowpea phylloplane and their role in disease development and control. MSc thesis, University of Ibadan, Ibadan
- Bhagat S, Pan S (2007) Mass multiplication of *Trichoderma harzianum* on agricultural byproducts and their evaluation against seedling blight of mungbean and collar rot of groundnut. Ind J Agric Sci 77:583–588
- Bhagat S, Pan S (2008) Variability in production of extra cellular hydrolytic enzymes by *Trichoderma* spp. and induction of disease resistance in gram (*Cicer arietinum*). J Biol Conserv 22:57–66

24. Carasolio CA, Gutierrez A, Jimenez B, Van Monkngu M, Herrera Estrella A (1994) Primary structures and expression pattern of the 33 kDa chitinase gene from the mycoparasitic fungus *Trichoderma harzianum*. Proc Natl Acad Sci USA 91:10903–10907
25. Ulhoa CJ, Peberdy JF (1991) Regulation of chitinase synthesis in *Trichoderma harzianum*. J Gen Microbiol 137:2163–2169
26. Elad Y, Chet I (1983) Improved selective media for isolation of *Trichoderma* spp. and *Fusarium* spp. Phytoparasitica 11:55–58