



Isolation and Identification of Plant Growth Promoting Rhizobacteria from Cucumber Rhizosphere and Their Effect on Plant Growth Promotion and Disease Suppression

Shaikhul Islam¹, Abdul M. Akanda¹, Ananya Prova¹, Md. T. Islam² and Md. M. Hossain^{3*}

¹ Department of Plant Pathology, EXIM Bank Agricultural University, Chapainawabganj, Bangladesh, ² Department of Biotechnology, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh, ³ Department of Plant Pathology, Bangabandhu Sheikh Mujibur Rahman Agricultural University, Gazipur, Bangladesh

OPEN ACCESS

Edited by:

Anton Hartmann,
Helmholtz Zentrum München –
German Research Center
for Environmental Health, Germany

Reviewed by:

Abu Hena Mostafa Kamal,
University of Texas at Arlington, USA
Christian Staehelin,
Sun Yat-sen University, China

*Correspondence:

Md. M. Hossain
hossainmm@bsmrau.edu.bd

Specialty section:

This article was submitted to
Plant Biotic Interactions,
a section of the journal
Frontiers in Microbiology

Received: 16 August 2015

Accepted: 16 November 2015

Published: 02 February 2016

Citation:

Islam S, Akanda AM, Prova A,
Islam MT and Hossain MM (2016)
Isolation and Identification of Plant
Growth Promoting Rhizobacteria from
Cucumber Rhizosphere and Their
Effect on Plant Growth Promotion
and Disease Suppression.
Front. Microbiol. 6:1360.
doi: 10.3389/fmicb.2015.01360

Plant growth promoting rhizobacteria (PGPR) are the rhizosphere bacteria that may be utilized to augment plant growth and suppress plant diseases. The objectives of this study were to identify and characterize PGPR indigenous to cucumber rhizosphere in Bangladesh, and to evaluate their ability to suppress *Phytophthora* crown rot in cucumber. A total of 66 isolates were isolated, out of which 10 (PPB1, PPB2, PPB3, PPB4, PPB5, PPB8, PPB9, PPB10, PPB11, and PPB12) were selected based on their *in vitro* plant growth promoting attributes and antagonism of phytopathogens. Phylogenetic analysis of 16S rRNA sequences identified these isolates as new strains of *Pseudomonas stutzeri*, *Bacillus subtilis*, *Stenotrophomonas maltophilia*, and *Bacillus amyloliquefaciens*. The selected isolates produced high levels (26.78–51.28 $\mu\text{g mL}^{-1}$) of indole-3-acetic acid, while significant acetylene reduction activities (1.79–4.9 $\mu\text{mole C}_2\text{H}_4 \text{ mg}^{-1} \text{ protein h}^{-1}$) were observed in eight isolates. Cucumber plants grown from seeds that were treated with these PGPR strains displayed significantly higher levels of germination, seedling vigour, growth, and N content in root and shoot tissue compared to non-treated control plants. All selected isolates were able to successfully colonize the cucumber roots. Moreover, treating cucumber seeds with these isolates significantly suppressed *Phytophthora* crown rot caused by *Phytophthora capsici*, and characteristic morphological alterations in *P. capsici* hyphae that grew toward PGPR colonies were observed. Since these PGPR inoculants exhibited multiple traits beneficial to the host plants, they may be applied in the development of new, safe, and effective seed treatments as an alternative to chemical fungicides.

Keywords: PGPR, plant growth promotion, IAA production, biological nitrogen fixation, antagonism, *Phytophthora capsici*, disease suppression

INTRODUCTION

The cucumber (*Cucumis sativus*) is one of the most widely grown vegetable crops in the world, and is particularly prevalent on the Indian sub-continent. This crop is prone to massive attacks by *Phytophthora capsici* that causes crown rot and blight (Kim et al., 2008; Maleki et al., 2011). *P. capsici* infects susceptible hosts throughout the growing season at any growth stage, and causes

yield losses as high as 100% (Lee et al., 2001). This pathogen has a wide host range with more than 50 plant species including Cucurbitaceae, Leguminosae, and Solanaceae (Hausbeck and Lamour, 2004). Although fungicides can control the disease, their use is detrimental to the surrounding environment and to the viability and survival of beneficial rhizosphere microbes (Carson et al., 1962; Hussain et al., 2009; Heckel, 2012). Furthermore, the growing cost of pesticides and the consumer demand for pesticide-free food have led to a search for substitutes for these products. Thus, there has been a need to find effective alternatives to costly and environmentally degrading synthetic pesticides.

Rhizobacteria that benefit plants by stimulating growth and suppressing disease are referred to as plant growth promoting rhizobacteria (PGPR; Kloepper et al., 1980). PGPR have been tested as biocontrol agents for suppression of plant diseases (Gerhardson, 2002), and also as inducers of disease resistance in plants (Cattelan et al., 1999; Bargabus et al., 2002; Bais et al., 2004). In particular, strains of *Pseudomonas*, *Stenotrophomonas*, and *Bacillus* have been successfully used in attempts to control plant pathogens and increase plant growth (Bais et al., 2004; Idris et al., 2007; Liu et al., 2007; Messiha et al., 2007; Chen et al., 2009; El-Sayed et al., 2014). The widely recognized mechanisms of plant growth promotion by PGPR are production of phytohormones, diazotrophic fixation of nitrogen, and solubilization of phosphate. Mechanisms of biocontrol action include competition with phytopathogens for an ecological niche or substrate, as well as production of inhibitory compounds and hydrolytic enzymes that are often active against a broad spectrum of phytopathogens (Zhang and Yuen, 2000; Manjula et al., 2004; Haas and Défago, 2005; Stein, 2005; Detry et al., 2006; Konsoula and Liakopoulou-Kyriakides, 2006; Cazorla et al., 2007).

Many PGPR have been shown to reduce *Phytophthora* crown rot occurrence on various plants. Ahmed et al. (2003) demonstrated *in vitro* suppression of *P. capsici* by bacterial isolates from the aerial part and rhizosphere of sweet pepper. An endophytic bacterium isolated from black pepper stem and roots, *B. megaterium* IISRBP17 suppressed *P. capsici* on black pepper in greenhouse assays (Aravind et al., 2009). Zhang et al. (2010) demonstrated that PGPR strains used separately or in combinations had the potential to suppress *Phytophthora* blight on squash in the greenhouse. Shirzad et al. (2012) reported that some fluorescent pseudomonads isolated from different fields of East and West Azarbaijan and Ardebil provinces of Iran exhibited strong antifungal activity against *P. drechsleri* and controlled crown and root rot of cucumber caused by the pathogen. However, little is known about PGPRs with the potential to suppress crown rot caused by *P. capsici* on cucumber. Furthermore, the plant-growth-promoting and biocontrol efficacy of PGPR often depend upon the rhizosphere competence of the microbial inoculants (Lugtenberg and Kamilova, 2009). Rhizosphere competence refers to the survival and colonization potential of PGPR (Bulgarelli et al., 2013), and is thought to be highest for each PGPR strain when associated with its preferred host plant. This to some extent explains why some PGPR strains exhibiting

promise as biocontrol agents *in vitro* have variable biocontrol efficacy in the rhizosphere of a given crop under a given set of conditions. The identification and characterization of PGPR populations indigenous to cucumber rhizospheres is therefore critical to discovery of strains that can be utilized to improve growth and *Phytophthora* crown rot suppression in cucumber. The objectives of the present study were to isolate bacterial strains from the cucumber rhizosphere, to characterize these isolates on the basis of morphological and physiological attributes as well as by 16S rRNA sequence analysis, and to assess the plant growth promoting effects of these isolates *in vivo* and their ability to suppress *Phytophthora* crown rot in cucumber plants. To our knowledge, this is the first report of PGPR reducing *P. capsici* infection on cucumber.

MATERIALS AND METHODS

The Study Site

The experimental site was located at the Field Laboratory of the Department of Plant Pathology, Bangabandhu Sheikh Mujibur Rahman Agricultural University (BSMRAU), Gazipur, Bangladesh. The location of the site is at 24.09° N latitude and 90.25° E longitude with an elevation of 2–8 m. The study area is within the Madhupur Tract agro-ecological zone (AEZ 28). The soil used for pot experiments belongs to the Salna series and has been classified as “swallow red brown terrace soil” in the Bangladesh soil classification system, which falls under the order Inceptisol (Brammer, 1978). This soil is characterized by clay within 50 cm of the surface and is slightly acidic in nature. The pH value, cation exchange capacity (CEC) and electrical conductivity (EC) of bulk soil samples collected from the study site were 6.38, 6.78 meq 100 g⁻¹ soil and 0.6 dS m⁻¹, respectively. This soil contained 1.08% organic carbon (OC), 1.87% organic matter (OM), 0.10% nitrogen (N), 9 ppm phosphorus (P) and 0.20 meq 100 g⁻¹ soil exchangeable potassium (K).

Plant Material, Bacterial Isolation, and Pathogenic Organism

Cucumber (*Cucumis sativus* L.) variety Baromashi (Lal Teer Seed Company, Dhaka, Bangladesh) root samples were collected from the study site along with rhizosphere soil. For isolation of bacteria, 2–5 g of fresh roots were washed under running tap water and surface sterilized in 5% NaOCl for 1 min. After washing three times with sterilized distilled water (SDW), the root samples were ground with a sterilized mortar and pestle. Serial dilutions were prepared from the ground roots, and 100 μl aliquots from each dilution of 1 × 10⁻⁶, 1 × 10⁻⁷, and 1 × 10⁻⁸ CFU mL⁻¹ were spread on potato dextrose agar (PDA) plates and incubated for 2 days at 28 ± 2°C. Morphologically distinct bacterial colonies were selected for further purifications. The purified isolates were preserved temporarily in 20% glycerol solution at -20°C. The pathogen *P. capsici* was provided by Prof. W. Yuanchao, Nanjing Agricultural University, China.

Morphological and Biochemical Characterization of Bacterial Isolates

Colony morphology, size, shape, color, and growth pattern were recorded after 24 h of growth on PDA plates at $28 \pm 2^\circ\text{C}$ as described by Somasegaran and Hoben (1994). Cell size was observed by light microscopy. The Gram reaction was performed as described by Vincent and Humphrey (1970). A series of biochemical tests were conducted to characterize the isolated bacteria using the criteria of Bergey's Manual of Systematic Bacteriology (Bergey et al., 1994). For the KOH solubility test, bacteria were aseptically removed from Petri plates with an inoculating wire loop, mixed with 3% KOH solution on a clean slide for 1 min and observed for formation of a thread-like mass. The motility of each isolate was tested in sulfide indole motility (SIM) medium. Using a needle, strains were introduced into test tubes containing SIM, and were incubated at room temperature until the growth was evident (Kirsop and Doyle, 1991). Turbidity away from the line of inoculation was a positive indicator of motility. Catalase and oxidase tests were performed as described in Hayward (1960) and Rajat et al. (2012), respectively. To determine whether the rhizobacterial isolates are better suited to aerobic or anaerobic environments, the citrate test was conducted according to Simmons (1926) using Simmons citrate agar medium. All experiments were done following complete randomized design (CRD) with three replications for each isolate and repeated once.

Molecular Characterization of Bacterial Isolates

Culture DNA was obtained using the lysozyme-SDS-phenol/chloroform method (Maniatis et al., 1982). DNA was extracted with phenol-chloroform-isoamyl alcohol (25:24:1) and precipitated with isopropanol. The extracted DNA was treated with DNase-free RNase (Sigma Chemical Co., St. Louis, MO, USA) at a final concentration of 0.2 mg/ml at 37°C for 15 min, followed by a second phenol-chloroform-isoamyl alcohol extraction and isopropanol precipitation. Finally, the DNA pellet was re-suspended in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0), stored at -20°C , and used as template DNA in PCR to amplify the 16S rRNA for phylogenetic analysis.

16S rRNA gene amplification was performed by using the bacterial-specific primers, 27F (5'-AGAGTTTGATCCTGG CTCAG-3') and 1492R (5'-GGTTACCTTGTTACGACTT-3') (Reysenbach et al., 1992). PCR amplifications were performed with $1 \times$ Ex Taq buffer (Takara Bio Inc, Japan), 0.8 mM dNTP, 0.02 units μl^{-1} Ex Taq polymerase, 0.4 mg ml^{-1} BSA, and 1.0 μM of each primer. Three independent PCR amplifications were performed at an annealing temperature of 55°C (40 s), an initial denaturation temperature of 94°C (5 min), 30 amplification cycles with denaturation at 94°C (60 s), annealing (30 s), and extension at 72°C (60 s), followed by a final extension at 72°C (10 min). The PCR product was purified using Wizard[®] PCR Preps DNA Purification System (Promega, Madison, WI, USA). Purified double-stranded PCR fragments were directly sequenced with Big Dye Terminator Cycle sequencing kits (Applied Biosystems, Forster City, CA,

USA) using the manufacturer's instructions. Sequences for each region were edited using Chromas Lite 2.0¹. The 16S rRNA sequence of the isolate has been deposited in the GenBank database. The BLAST search program² was used to search for nucleotide sequence homology for the 16S region for bacteria. Highly homologous sequences were aligned, and neighbor-joining trees were generated using ClustalX version 2.0.11 and MEGA version 6.06. Bootstrap replication (1000 replications) was used as a statistical support for the nodes in the phylogenetic trees.

Bioassays for Plant Growth Promoting Traits

Biological Nitrogen Fixation

Nitrogenase activity of isolates was determined via the acetylene reduction assay/ethylene production assay as described in Hardy et al. (1968). Pure bacterial colonies were inoculated to an airtight 30 ml vial containing 10 ml nitrogen-free basal semi-solid medium, and were grown for 48 h at $28 \pm 2^\circ\text{C}$. Following pellicle formation, the bottles were injected with 10% (v/v) acetylene gas and incubated at $28 \pm 2^\circ\text{C}$ for 24 h. Ethylene production was measured using a G-300 Gas Chromatograph (Model HP 6890, USA) fitted with a Flame Ionization Detector and a Porapak-N column. Hydrogen and oxygen were used as a carrier gas, with a flow rate of 4 kg/cm^2 , and the column temperature was maintained at 165°C . The ethylene concentration calibration curve was plotted for each trial, and the viable cell numbers (cfu) of the isolate were determined. The rate of N_2 fixation was expressed as the quantity of ethylene accumulated ($\mu\text{mol C}_2\text{H}_4 \text{ mg}^{-1}$ protein h^{-1}) based on the standard curve and peak-area percentage.

Indole-3-Acetic Acid Production

For detection and quantification of indole-3-acetic acid (IAA) production by bacterial isolates, isolated colonies were inoculated into Jensen's broth (Sucrose 20 g, K_2HPO_4 1 g L^{-1} , $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ 0.5 g L^{-1} , NaCl 0.5 g L^{-1} , FeSO_4 0.1 g L^{-1} , NaMoO_4 0.005 g L^{-1} , CaCO_3 2 g L^{-1}) (Bric et al., 1991) containing 2 mg mL^{-1} L-tryptophan. The culture was incubated at $28 \pm 2^\circ\text{C}$ with continuous shaking at 125 rpm for 48 h (Rahman et al., 2010). Approximately 2 mL of culture solution was centrifuged at 15000 rpm for 1 min, and a 1 mL aliquot of the supernatant was mixed with 2 mL of Salkowski's reagent and incubated 20 min in darkness at room temperature (150 ml concentrated H_2SO_4 , 250 ml distilled water, 7.5 ml 0.5 M $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$) as described by Gordon and Weber (1951). IAA production was observed as the development of a pink-red color, and the absorbance was measured at 530 nm using a spectrophotometer. The concentration of IAA was determined using a standard curve prepared from pure IAA solutions (0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, and 65 $\mu\text{g ml}^{-1}$).

¹<http://www.technelysium.com.au/chromas.html>

²<http://blast.ncbi.nlm.nih.gov/Blast.cgi>

Preparation of Bacterial Inocula for Cucumber Seed Treatment

Bacterial strains were cultured in 250 ml conical flasks containing 200 ml yeast peptone broth on an orbital shaker at 120 rpm for 72 h at $28 \pm 2^\circ\text{C}$. Bacterial cells were collected via centrifugation at 15000 rpm for 1 min at 4°C , and each pellet was washed twice with SDW. The bacterial pellets were suspended in 0.6 ml SDW, vortexed and used for seed treatment. Approximately 30–31 cucumber seeds were surface sterilized in 5% NaOCl for 1 min and washed three times in SDW. Dry seeds were immersed in each bacterial suspension, and the preparation was stirred frequently for 5 min. The treated seeds were spread on a petri dish and air dried overnight at room temperature. The number of bacterial cells per seed, determined via serial dilutions, was approximately 10^8 CFU/seed.

Effect of Bacterial Seed Treatment on Germination and Vigour Index in Cucumber

In order to determine the effect of the isolates on germination and seedling vigour, 100 seeds inoculated with each isolate were incubated in ten 9-cm petri dishes on two layers of moistened filter paper. As a control treatment, seeds treated with water instead of bacterial suspensions were also established. In order to maintain sufficient moisture for germination, 5 ml distilled water was added to the petri dishes every other day, and seeds were incubated at $28 \pm 2^\circ\text{C}$ in a light incubator. Germination was considered to have occurred when the radicles were half of the seed length. The germination percentage was recorded every 24 h for 7 days. Root and shoot length were measured after the seventh day. The experiment was planned as a completely randomized design with 10 replications for each isolate.

$$\text{Germination rate (\%)} = \left(\frac{\text{number of seeds germinated}}{\text{total number of seeds}} \right) \times 100$$

$$\text{Vigour index} = \% \text{ germination} \times \text{total plant length}$$

Effect of Bacterial Seed Treatment on Growth and Nitrogen Content in Cucumber Plants

In order to test the ability of isolates to promote growth in cucumber plants, surface-sterilized cucumber seeds were inoculated with each isolate as described above. The soil from the study site described above was used as potting medium. After autoclaving twice at 24 h intervals at 121°C and 15 psi for 20 min, 180 g of the sterilized soil was placed in each sterilized plastic pot (9.5 cm \times 7.0 cm size). One pre-germinated cucumber seed was sown in each pot, and plants were grown 3 weeks in a net house with watering on alternate days. After harvest, the fresh weight, dry weight, and root and shoot lengths of the plants were measured. The shoots and roots were separated and dried in an

oven at $68 \pm 2^\circ\text{C}$ for 48 h, then ground for determination of tissue-N concentrations (Kjeldahl, 1883).

Root Colonization

Root colonization by bacterial isolates was determined according to the protocol of Hossain et al. (2008). Roots were harvested from plants at 7, 14, and 21 days of growth. Root systems were thoroughly washed with running tap water to remove adhering soil particles, then were rinsed three times with SDW and blotted to dryness. Roots were divided into top, middle, and bottom regions, and were weighed and homogenized in SDW. Serial dilutions were prepared on PDA plates, and the number of colony-forming units (cfu) per gram root was determined after 24 h of incubation at $28 \pm 2^\circ\text{C}$.

Pathogenicity of *P. capsici* in Cucumber

For preparation of zoospore inoculum, *P. capsici* was cultured on PDA plates at $18 \pm 2^\circ\text{C}$ for 7 days. Five-mm blocks were then cut from the culture plates and placed in petri dishes containing SDW. The petridishes were incubated in darkness at room temperature for 72 h, followed by a 1-h cold treatment at 4°C . Zoospore production was confirmed via light microscopy. In order to test the pathogenicity of *P. capsici* zoospores, cucumber seedlings were planted in pots containing 0, 500, 1000, or 1500 μl zoospores/pot. As 100% mortality was found in case 1000 μl zoospores/pot, two concentrations (500 and 1000 μl) of zoospores suspension per pot were fixed.

In Vitro Screening for Antagonism

To test antagonism of *P. capsici* by each isolate, the pathogen and bacteria were inoculated 3 cm apart on the same agar plate. Fungal growth on each plate was observed, and the zone of inhibition, if present, was determined as described in Riungu et al. (2008):

$$\% \text{ Inhibition of mycelial growth} = \frac{X-Y}{X} \times 100$$

Where,

X = Mycelial growth of pathogen in absence of antagonists

Y = Mycelial growth of pathogen in presence of antagonists

Morphologies of hyphae in the vicinity of bacterial colonies were observed under a light microscope (Meiji Techno: ML2600), and images were recorded with a digital camera (Model: Canon Digital IXUS 900 Ti). Each experiment was carried out following CRD with three replications for each isolate and repeated once.

Testing the Effect of Rhizobacterial Seed Treatment on Phytophthora Crown Rot of Cucumber

Cucumber seeds inoculated with each isolate were sown and grown for 7 days in sterilized plastic pots as described above. Seven-day-old seedlings were inoculated with 500 or 1000 μl zoospore suspension/pot as described in Deora et al.

(2005). Inoculated plants were kept inside humid chambers for 48 h. Each experiment included 12 plants per treatment with three replications. Surviving plants were counted 7 days after inoculation. Percent disease incidences (PDI) were calculated using the following formula:

$$\text{PDI} = \frac{\text{Number of infected plants}}{\text{Total number of inoculated plants}} \times 100\%$$

Percent protection by PGPR was calculated using following formula:

$$\% \text{ Protection} = \left[\frac{A-B}{A} \right] \times 100\%$$

Where,

A = PDI in non-inoculated control plants

B = PDI in PGPR-treated plants.

Statistical Analysis

Statistical analyses were performed using SPSS (Version 17) and Microsoft Office Excel 2007. A completely randomized design was used for all experiments, with 3–12 replications for each treatment. The data presented are from representative experiments that were repeated at least twice with similar results. Treatments were compared via ANOVA using the least significant differences test (LSD) at 5% ($P \leq 0.05$) probability level.

RESULTS

Strain Isolation and Biochemical and Molecular Characterization

We obtained a total of 66 rhizobacterial strains from the interior of cucumber roots. Ten isolates – PPB1, PPB2, PPB3, PPB4, PPB5, PPB8, PPB9, PPB10, PPB11, and PPB12 – were selected based on their ability to produce IAA, fix N_2 , and show *in vitro* antagonism against various pathogens in a preliminary screening. All isolates were rods producing fast-growing, round to irregular colonies with raised elevations and smooth surfaces. Reddish pigmentation was produced by PPB5, while no pigmentation was produced by other isolates (Supplementary Table S1). All 10 isolates were motile and reacted positively to the Gram staining, citrate, catalase and oxidase tests, but reacted negatively to the KOH solubility test (Table 1).

Phylogenetic trees constructed from 16S rRNA sequences showed that the selected isolates were mainly members of genus *Bacillus*, *Pseudomonas*, and *Stenotrophomonas* (Supplementary Figure S1). The sequences of the isolates PPB2, PPB5, PPB8, PPB9, and PPB11 showed 99% similarity with *Bacillus subtilis* and were submitted to GenBank under accession numbers KJ690255, KM008605, KM008606, KM092525 and KM092527, respectively (Table 1). Isolate PPB1 had 99% homology with *Pseudomonas stutzeri* and was submitted to GenBank under accession number KJ959616. Isolate PPB3 was identified as *Stenotrophomonas maltophilia* with GenBank accession number KJ959617. Isolates PPB4, PPB10, and PPB12 showed 99% sequence homology with

B. amyloliquefaciens and were submitted to GenBank under accession number KM008604, KM092526 and KM092528, respectively (Table 1).

Characterization for Plant Growth Promoting Traits

The plant growth promoting characteristics viz., IAA production and ARA were examined with the ten selected PGPR isolates. The results of the assays are presented in Table 2. In the presence of tryptophan, the isolated bacteria produced IAA in concentrations between 26.78 $\mu\text{g mL}^{-1}$ and 51.28 $\mu\text{g mL}^{-1}$. The highest and lowest amounts of IAA were produced by strain PPB5 and PPB3, respectively (Table 2). Nitrogenase activity, as determined by ARA, was not detected in PPB1 and PPB12 under the conditions tested. However, the ARA values ranged from 1.79 to 4.9 $\mu\text{mole C}_2\text{H}_4/\text{mg protein/h}$ for remaining isolates. PPB2 showed the highest activity, while the lowest was recorded for PPB11 (Table 1). The other isolates also reduced acetylene in significant amounts. Collectively, these results suggest that the isolates possess a number of traits associated with plant growth promotion.

Germination and Vigour Index Improvement in Cucumber

The effect of rhizobacterial treatment upon seed germination and vigour index of cucumber varied with different isolates. All treatments had a significant effect on the germination rate and vigour index compared to the control. The PGPR treatments increased the germination rate of cucumber seeds by 8.07–15.32% compared with the control, while the vigour index was increased by 98.62–148.05% (Figure 1). In both germination rate and vigour index, the maximum increase was obtained with the PPB9 treatment. These results suggest that rhizobacterial treatment could improve the germination and vigour of cucumber seeds.

Plant Growth Promotion in Cucumber

All isolates significantly increased the growth of cucumber compared to non-inoculated controls. Treatment with isolate PPB12 produced the maximum shoot and root lengths of 18.23 and 20.47 cm, corresponding to increases of 66.02 and 65.63% above control treatments (Figure 1). However, the maximum shoot and root weight enhancement was observed in PPB8-treated plants. Treatment with isolate PPB8 produced shoot fresh and dry weights of 5.29 and 0.60 g plant^{-1} , which were 79.32 and 100.00% higher than those of control plants. Similarly, treatment with isolate PPB8 produced root fresh and dry weights of 3.03 and 0.32 g plant^{-1} , corresponding to increase of 91.77 and 128.57% above control treatments.

N Concentration in Cucumber Plants

The N content in plant roots and shoots significantly increased due to inoculation treatments with rhizobacterial isolates (Figure 2). The shoot and root N content showed similar trends in response to different treatments; hence, the N content is reported as the total combined shoot and root N. The total N content in PGPR-treated plants ranged from 3.66 mg g^{-1} to

TABLE 1 | Biochemical and molecular analysis of the endophytic bacterial isolates.

Strains	Biochemical analysis						Molecular analysis	
	KOH Test	Gram reaction	Citrate Test	Catalase Test	Oxidase Test	IAA ($\mu\text{g/ml}$)	ARA ($\mu\text{mole C}_2\text{H}_4$ mg protein/h)	Identification based on 16S rRNA gene sequencing
PPB1	–	+ve	+	+	+	39.67 \pm 0.12*	0.00 \pm 0.00*	<i>Pseudomonas stutzeri</i>
PPB2	–	+ve	+	+	+	41.43 \pm 0.71	4.90 \pm 0.23	<i>Bacillus subtilis</i>
PPB3	–	+ve	+	+	+	26.78 \pm 0.69	4.55 \pm 0.38	<i>Stenotrophomonas maltophilia</i>
PPB4	–	+ve	+	+	+	50.18 \pm 0.23	2.41 \pm 0.12	<i>B. amyloliquefaciens</i>
PPB5	–	+ve	+	+	+	51.28 \pm 0.41	2.90 \pm 0.17	<i>B. subtilis</i> subsp. <i>subtilis</i>
PPB8	–	+ve	+	+	+	44.41 \pm 0.22	4.79 \pm 0.01	<i>B. subtilis</i>
PPB9	–	+ve	+	+	+	41.75 \pm 0.93	3.95 \pm 0.02	<i>B. subtilis</i> subsp. <i>spizizenii</i>
PPB10	–	+ve	+	+	+	38.43 \pm 0.82	1.83 \pm 0.01	<i>B. amyloliquefaciens</i>
PPB11	–	+ve	+	+	+	40.30 \pm 0.23	1.79 \pm 0.01	<i>B. subtilis</i> subsp. <i>subtilis</i>
PPB12	–	+ve	+	+	+	29.25 \pm 0.97	0.00 \pm 0.00	<i>B. amyloliquefaciens</i>

'–' corresponds to negative response; '+ve' and '+' correspond to positive responses. *Values are the Mean \pm SE. The experiment was repeated twice with three replicates for each isolate.

TABLE 2 | Comparative performance of PGPR in mycelia growth inhibition of *P. capsici* and Phytophthora crown rot disease suppression in cucumber plants.

Treatments	Pathogen suppression ^a (% <i>P. capsici</i> mycelial growth inhibition)	Disease suppression ^b (% Protection ^c)	
		500 μl Zoospores/pot ^d	1000 μl Zoospores/pot
PPB1	67.16 \pm 0.68e*	58.33 \pm 0.66c	33.33 \pm 2.18b
PPB2	70.01 \pm 0.85g	69.45 \pm 0.80e	38.84 \pm 1.70c
PPB3	69.08 \pm 0.91f	50.00 \pm 1.69b	50.00 \pm 1.54e
PPB4	62.07 \pm 0.11c	70.33 \pm 2.37f	45.68 \pm 2.37d
PPB5	65.94 \pm 0.53d	50.00 \pm 0.57b	33.33 \pm 1.69b
PPB8	82.05 \pm 0.55j	83.33 \pm 0.56h	77.78 \pm 2.25g
PPB 9	90.08 \pm 0.46k	66.67 \pm 1.87d	66.67 \pm 2.93f
PPB10	73.08 \pm 0.83h	73.67 \pm 1.53g	66.67 \pm 0.52f
PPB11	58.32 \pm 0.12b	88.83 \pm 1.67i	86.08 \pm 2.23h
PPB12	80.53 \pm 0.69i	50.00 \pm 1.15b	33.33 \pm 0.43b
Control	0.00 \pm 0.00a	0.00 \pm 0.00a	0.00 \pm 0.00a

*Values are the means \pm SE ($n = 12$). Data within the same column followed by different letters are significantly different.

^aPathogen suppression was measured as percent inhibition of radial growth of *P. capsici* by antagonistic PGPR in dual plate assay.

^bDisease suppression was expressed as percent protection due to treatment with PGPR relative to control (non-inoculated).

^cProtection (%) = $[(A - B)/A] \times 100$ in which, A = PDI in non-inoculated control plants and B = PDI in PGPR-treated plants.

^dSeven-day-old seedlings were inoculated with 500 or 1000 μl zoospore suspension/pot.

The data presented are from representative experiments that were repeated twice with similar results.

8.25 mg g⁻¹ N compared with 2.57 mg g⁻¹ N for non-inoculated control plants, a 42–221% increase in PGPR-treated plants over control plants (Figure 2). The highest N content was recorded in plants grown under PPB2 followed by PPB8, PPB3, PPB9 and other treatments.



Root Colonization

The ability to colonize the root system is essential for rhizobacteria to be effective plant growth promoters. The root colonization assays showed that all the tested isolates successfully colonized the roots of cucumber plants as tested after only 7 days of seedling growth. The inoculated populations were even higher on 21-day-old roots. Nevertheless, the root population densities varied widely among the isolates (Figure 3). The largest root populations were observed for strain PPB2, followed by PPB5 and

PPB9 (Figure 3). These results demonstrate specific interactions between cucumber plants and the rhizobacterial isolates.

In vitro Antagonism of *Phytophthora capsici*



All rhizobacterial isolates exhibited significant antagonistic activity against *P. capsici* on PDA. The largest inhibition zone was observed with PPB9 (90.08%) followed by PPB8 (82.05%) (Table 2). Distinct morphological alterations in *P. capsici* hyphae were also detected during dual cultures with the rhizobacterial isolates. Hyphal features observed in the vicinity of bacterial colonies included irregular and excessive branching, abnormal swelling of hyphal diameters, unusually long and pointed hyphal tips, and vacuolization leading to hyphal lysis (Figure 4).



Suppression of Phytophthora Crown Rot in Cucumber

All the selected PGPR strains showed consistent suppression of *Phytophthora* crown rot in the greenhouse experiments.

Compared with the control, the average disease protection at 500 μl zoospore suspension ranged from 50 to 88.83% after treatment with rhizobacterial isolates, while protection at 1000 μl zoospore suspension ranged from



33.33 to 86.08% (Table 2). At both inoculum rates, isolate PPB11 showed the highest disease reduction, and the lowest disease reduction was obtained with PPB5.

DISCUSSIONS

PGPR colonizing the surface or inner part of roots play important beneficial roles that directly or indirectly influence

plant growth and development (Glick et al., 1999; Gerhardt et al., 2009). In this study, 10 PGPR classified as *Pseudomonas stutzeri* (PPB1), *B. subtilis* (PPB2, 5, 8, 9, and 11), *S. maltophilia* (PPB3), and *B. amyloliquefaciens* (PPB4, 10, and 12) were isolated from the rhizosphere of cucumber plants. All the isolated PGPR were gram positive and motile rods, and tested positive for the ability to utilize citrate as a carbon source. Flagellar motility and citrate utilization are both thought to play a significant role in competitive root colonization and maintenance of bacteria in roots (Turnbull et al., 2001; Weisskopf et al., 2011). These strains also tested positive for oxidase and catalase activity. Standard microbiology references suggest that *S. maltophilia* is an oxidase-negative bacterium (Ryan et al., 2009). Recent data, however, suggest that some *S. maltophilia* are oxidase-positive (Carmody et al., 2011), and this was also the case for isolate PPB3 in this study. Our catalase test results corroborate prior studies showing that *B. subtilis*, *Pseudomonas stutzeri*, and *B. amyloliquefaciens* are catalase-positive (Merchant and Packer, 1999; Kamboh et al., 2009). *Bacillus* and *Pseudomonas* are the most frequently reported genera of PGPR (Laguette et al., 1994; Hallmann and Berg, 2006; Zahid et al., 2015). Similarly, most isolates in this study belong to genus *Bacillus*.

Treatment of cucumber seeds with the selected isolates significantly improved seedling emergence and growth. Several different mechanisms have been suggested for similar observations using other PGPR strains: PGPR might indirectly enhance seed germination and vigour index by reducing the incidence of seed mycoflora, which can be detrimental to plant growth (Begum et al., 2003). Duarah et al. (2011) found that amylase activity during germination was increased in rice and legume after inoculation with PGPR. The amylase hydrolyzes the starch into metabolizable sugars, which provide the energy for growth of roots and shoots in germinating seedlings (Beck and Ziegler, 1989; Akazawa and Nishimura, 2011). One of the most commonly reported mechanisms is the production of phytohormones such as IAA (Patten and Glick, 2002). All the selected isolates in this study produced IAA. Similar studies have shown that IAA production is very common among PGPR (Yasmin et al., 2004; Ng et al., 2012; Zahid et al., 2015). In fact, many isolates in this study produced higher IAA than previously reported strains (Yasmin et al., 2004; Banerjee et al., 2010; Ng et al., 2012). This is an important mechanism of plant growth promotion because IAA promotes root development and uptake of nutrients (Carrillo et al., 2002). It has long been proposed that phytohormones act to coordinate demand and acquisition of nitrogen (Kiba et al., 2011). Therefore, enhanced N-content found in inoculated plants could be due to increased N-uptake by the roots caused by hormonal effects on root morphology and activity. Nitrogen fixation may also play a role in plant growth promotion. All the selected isolates in this study except PPB1 and PPB12 showed acetylene reduction activity, which is a widely accepted surrogate for nitrogenase activity and N₂-fixing potential (Andrade et al., 1997). However, defensible proof of N₂-fixation needs the application of ¹⁵N as tracer of soil N or as ¹⁵N₂-gas and the

demonstration of significantly changed N-isotope-labeling in the plant biomass. Transfer of N between diazotrophic N-fixing rhizobacteria and the roots of several crops has been demonstrated (Islam et al., 2009; Abbasi et al., 2011; Tajini et al., 2012; Verma et al., 2013). It is interesting to note that in this study all isolates, including the two that demonstrated no acetylene reduction activity, enhanced the N content of cucumber. This suggests that while N₂ fixation may be an important mechanism of plant growth promotion, there may also be alternate mechanisms, like hormonal interactions and nutrient uptake or pathogen suppression, which might be more pronounced than the contribution of nitrogen fixation.

Results from our study indicate that PGPR strains applied as a seed treatment significantly reduced disease severity of Phytophthora crown rot on cucumber plants. The fungal antagonists *Pseudomonas stutzeri*, *B. subtilis*, *B. amyloliquefaciens*, and *S. maltophilia* were have been shown to be effective biocontrol agents in prior studies (Dunne et al., 2000; Zhang and Yuen, 2000; Dal Bello et al., 2002; Berg et al., 2005; Islam and Hossain, 2013; Erlacher et al., 2014). Competitive root tip colonization by PGPR strains might play an important role in the efficient control of soil-borne diseases. The crucial colonization level that must be reached has been estimated at 10⁵–10⁶ cfu g⁻¹ of root in the case of *Pseudomonas* sp., which protect plants from *Gaeumannomyces tritici* or *Pythium* sp. (Haas and Défago, 2005). Most of our selected strains were efficient colonizers of roots, since CFU counts for tested strains were more than 10⁷ cfu g⁻¹ root. However, the former study examined the root colonization by introduced bacteria under natural field soil conditions, while our study did under axenic conditions. In view of that, comparison between root colonization data obtained under these two conditions may not be accurate. Biological control of *P. capsici* can also result from antibiosis by the bacteria (Nakayama et al., 1999; Kawulka et al., 2004; Chung et al., 2008; Lim and Kim, 2010; Mousivand et al., 2012; Islam and Hossain, 2013). All the selected isolates exhibited moderate to high antagonistic activity against *P. capsici* *in vitro*, and caused clear morphological distortions such as abnormal branching, curling, swelling and lysis of the hyphae at the interaction zone in dual cultures. Excessive branching and curling accompanied by marked ultrastructural alterations including invagination of the hyphal membrane, disintegration or necrosis of hyphal cell walls, accumulation of excessive lipid bodies, enlarged and electron-dense vacuoles, and degradation of cytoplasm were potentially due to bacterial production of antibiotics and lytic enzymes (Deora et al., 2005; Islam and von Tiedemann, 2011). These antibiotics and lytic enzymes cause membrane damage and are particularly inhibitory to zoospores of Oomycete (de Souza et al., 2003; Beneduzi et al., 2012). Induced systemic resistance is most likely another mechanism by which bacteria suppress *P. capsici* (Zhang et al., 2010).

In the present study, we have isolated 10 new strains of PGPR from indigenous cucumber plants. These strains possessed

several plant growth promoting traits as well as antifungal activity, and were found to be efficient in controlling *Phytophthora* crown rot of cucumber seedlings. *In vitro* and *in vivo* evidence suggest that the selected isolates benefit cucumber plants via multiple modes of action including antibiosis against phytopathogens, competitive colonization, and plant growth promotion. This reveals the potential of these strains for biofertilizer applications and commercial use as biocontrol agents in the field. However, from the estimation of a PGPR-potential to a biofertilizer application, it requires a long way of greenhouse experiments with pot filled with different type of soils and finally, field experiments to find out the optimum formulations for the inoculums. Thus, the inoculants can perform close to its optimum potential. Future studies concerning commercialization and field applications of integrated stable bio-formulations as effective biocontrol strategies are in progress.

AUTHOR CONTRIBUTION

SI was involved in the planning and execution of the research work; collection, analysis and interpretation of the data; manuscript writing etc. following the suggestions and directions of the Major Professor. AMA served as the Member of the Dissertation Committee of SI and was involved in the planning of the work and editing of the manuscript. AP was actively involved in the original research work, data collection, analysis as well

as manuscript preparation. TI supplied the *Phytophthora capsici* inocula and oversaw the sequence work of the bacterial isolates and related description in the manuscript. MMH served as the Major Professor of SI and was involved in the research design and planning; analysis and interpretation of data; drafting as well as critical revision of the work for intellectual content.

All authors approve the final version of the manuscript for publication and agrees to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

ACKNOWLEDGMENTS

The authors would like to acknowledge the financial assistance from University Grant Commissions through Research Management Committee of Bangabandhu Sheikh Mujibur Rahman Agricultural University, Bangladesh.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2015.01360>

FIGURE S1 | Phylogenetic tree of 16S rRNA gene sequences showing the relationships among the isolates isolated from cucumber rhizosphere. The data of type strains of related species were from GenBank database.

REFERENCES

- Abbasi, M. K., Sharif, S., Kazmi, M., Sultan, T., and Aslam, M. (2011). Isolation of plant growth promoting rhizobacteria from wheat rhizosphere and their effect on improving growth, yield and nutrient uptake of plants. *Plant Biosyst.* 145, 159–168. doi: 10.1080/11263504.2010.542318
- Ahmed, A., Ezziyyani, M., Egea-Gilbert, C., and Candela, M. E. (2003). Selecting bacterial strains for use in the biocontrol of diseases caused by *Phytophthora capsici* and *Alternaria alternata* in sweet pepper plants. *Biol. Plant.* 47, 569–574.
- Akazawa, T., and Nishimura, H. (2011). Topographic aspects of biosynthesis, extracellular secretion, and intracellular storage of proteins in plant cells. *Annu. Rev. Plant Physiol.* 36, 441–472. doi: 10.1146/annurev.pp.36.060185.002301
- Andrade, G., Esteban, E., Velasco, L., Lorite, M. J., and Bedmar, E. J. (1997). Isolation and identification of N₂-fixing microorganisms from the rhizosphere of *Capparis spinosa* (L.). *Plant Soil* 197, 19–23. doi: 10.1023/A:100421190964
- Aravind, R., Kumar, A., Eapen, S. J., and Ramana, K. V. (2009). Endophytic bacterial flora in root and stem tissues of black pepper (*Piper nigrum* L.) genotype: isolation, identification and evaluation against *Phytophthora capsici*. *Let. Appl. Microbiol.* 48, 58–64. doi: 10.1111/j.1472-765X.2008.02486.x
- Bais, H. P., Fall, R., and Vivanco, J. M. (2004). Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production. *Plant Physiol.* 134, 307–319. doi: 10.1104/pp.103.028712
- Banerjee, S., Palit, R., Sengupta, C., and Standing, D. (2010). Stress induced phosphate solubilization by 'Arthrobacter' sp. and 'Bacillus' sp. isolated from tomato rhizosphere. *Aust. J. Crop. Sci.* 4, 378–383.
- Bargabus, R. L., Zidack, N. K., Sherwood, J. E., and Jacobsen, B. J. (2002). Characterization of systemic resistance in sugar beet elicited by a non-pathogenic, phyllosphere-colonizing *Bacillus mycoides*, biological control agents. *Physiol. Mol. Plant Pathol.* 61, 289–298. doi: 10.1006/pmpp.2003.0443
- Beck, E., and Ziegler, P. (1989). Biosynthesis and degradation of starch in higher plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 40, 95–117. doi: 10.1146/annurev.pp.40.060189.000523
- Begum, M., Rai, V. R., and Lokesh, S. (2003). Effect of plant growth promoting rhizobacteria on seed borne fungal pathogens in okra. *Indian Phytopathol.* 56, 156–158.
- Beneduzi, A., Ambrosini, A., and Passaglia, L. M. (2012). Plant growth-promoting rhizobacteria (PGPR): their potential as antagonists and biocontrol agents. *Genet. Mol. Biol.* 35, 1044–1051. doi: 10.1590/S1415-47572012000600020
- Berg, G., Krechel, A., Ditz, M., Sikora, R. A., Ulrich, A., and Hallmann, J. (2005). Endophytic and ectophytic potato-associated bacterial communities differ in structure and antagonistic function against plant pathogenic fungi. *FEMS Microbiol. Ecol.* 51, 215–229. doi: 10.1016/j.femsec.2004.08.006
- Bergey, D. H., Holt, J. G., and Noel, R. K. (1994). *Bergey's Manual of Systematic Bacteriology*, Vol. 1, 9th Edn (Baltimore, MD: Williams & Wilkins), 1935–2045.
- Brammer, H. (1978). *Rice Soils of Bangladesh*. In: *Soils and Rice*. Los Banos: International Rice Research Institute Publication, 35–55.
- Bric, J. M., Bostock, R. M., and Silverstone, S. E. (1991). Rapid in situ assay for indole acetic acid production by bacteria immobilized on a nitrocellulose membrane. *Appl. Environ. Microbiol.* 57, 535–538.
- Bulgarelli, D., Schlaeppi, K., Spaepen, S., Ver Loren van Themaat, E., and Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. *Annu. Rev. Plant Biol.* 64, 807–838. doi: 10.1146/annurev-arplant-050312-120106
- Carmody, L. A., Spilker, T., and Li Puma, J. J. (2011). Reassessment of *Stenotrophomonas maltophilia* phenotype. *J. Clin. Microbiol.* 49, 1101–1103. doi: 10.1128/JCM.02204-10

- Carrillo, A. E., Li, C. Y., and Bashan, Y. (2002). Increased acidification in the rhizosphere of cactus seedlings induced by *Azospirillum brasilense*. *Naturwissenschaften* 89, 428–432. doi: 10.1007/s00114-002-0347-6
- Carson, R., Darling, L., and Darling, L. (1962). *Silent Spring*. Cambridge, MA: Riverside Press.
- Cattelan, A. J., Hartel, P. G., and Fuhrmann, J. J. (1999). Screening for plant growth-promoting rhizobacteria to promote early soybean growth. *Soil Sci. Soc. Am. J.* 63, 1670–1680. doi: 10.2136/sssaj1999.6361670x
- Cazorla, F. M., Romero, D., Pérez-García, A., Lugtenberg, B. J., Vicente, A., and Bloemberg, G. (2007). Isolation and characterization of antagonistic *Bacillus subtilis* strains from the avocado rhizosphere displaying biocontrol activity. *J. Appl. Microbiol.* 103, 1950–1959. doi: 10.1111/j.1365-2672.2007.03433.x
- Chen, X. H., Koumoutsis, A., Scholz, R., Schneider, K., Vater, J., Süßmuth, R., et al. (2009). Genome analysis of *Bacillus amyloliquefaciens* FZB42 reveals its potential for biocontrol of plant pathogens. *J. Biotechnol.* 140, 27–37. doi: 10.1016/j.jbiotec.2008.10.011
- Chung, S., Kong, H., Buyer, J. S., Lakshman, D. K., Lydon, J., Kim, S. D., et al. (2008). Isolation and partial characterization of *Bacillus subtilis* ME488 for suppression of soilborne pathogens of cucumber and pepper. *Appl. Microbiol. Biotechnol.* 80, 115–123. doi: 10.1007/s00253-008-1520-4
- Dal Bello, G. M., Mónaco, C. I., and Simón, M. R. (2002). Biological control of seedling blight of wheat caused by *Fusarium graminearum* with beneficial rhizosphere microorganisms. *World J. Microbiol. Biotechnol.* 18, 627–636. doi: 10.1023/A:1016898020810
- Deora, A., Hashidoko, Y., Islam, M. T., and Tahara, S. (2005). Antagonistic rhizosphere bacteria induce diverse morphological alterations in *Peronosporomycete hyphae* during in vitro interaction. *Eur. J. Plant Pathol.* 112, 311–322. doi: 10.1007/s10658-005-4753-4
- de Souza, J. T., Arnould, C., Deulvot, C., Lemanceau, P., Gianinazzi-Pearson, V., and Raaijmakers, J. M. (2003). Effect of 2,4-diacetylphloroglucinol on *Pythium*: cellular responses and variation in sensitivity among propagules and species. *Phytopathology* 93, 966–975. doi: 10.1094/PHYTO.2003.93.8.966
- Detry, J., Rosenbaum, T., Lütz, S., Hahn, D., Jaeger, K. E., Müller, M., et al. (2006). Biocatalytic production of enantiopure cyclohexane-trans-1, 2-diol using extracellular lipases from *Bacillus subtilis*. *Appl. Microbiol. Biotechnol.* 72, 1107–1116. doi: 10.1007/s00253-006-0391-9
- Duarah, I., Deka, M., Saikia, N., and Deka, B. H. P. (2011). Phosphate solubilizers enhance NPK fertilizer use efficiency in rice and legume cultivation. *Biotech* 1, 227–238.
- Dunne, C., Loccoz, Y. M., de Bruijn, F. J., and O’Gara, F. (2000). Overproduction of an inducible extracellular serine protease improves biological control of *Pythium ultimum* by *Stenotrophomonas maltophilia* strain W81. *Microbiology* 146, 2069–2078. doi: 10.1099/00221287-146-8-2069
- El-Sayed, S. W., Akhkh, A., El-Naggar, M. Y., and Elbadry, M. (2014). In vitro antagonistic activity, plant growth promoting traits and phylogenetic affiliation of rhizobacteria associated with wild plants grown in arid soil. *Front. Microbiol.* 5:651. doi: 10.3389/fmicb.2014.00651
- Erlacher, A., Cardinale, M., Grosch, R., Grube, M., and Berg, G. (2014). The impact of the pathogen *Rhizoctonia solani* and its beneficial counterpart *Bacillus amyloliquefaciens* on the indigenous lettuce microbiome. *Front. Microbiol.* 5:175. doi: 10.3389/fmicb.2014.00175
- Gerhardson, B. (2002). Biological substitutes for pesticides. *Trends Biotechnol.* 20, 338–343. doi: 10.1016/S0167-7799(02)02021-8
- Gerhardt, K. E., Huang, X.-D., Glick, B. R., and Greenberg, B. M. (2009). Phytoremediation and rhizoremediation of organic soil contaminants: potential and challenges. *Plant Sci.* 176, 20–30. doi: 10.1016/j.plantsci.2008.09.014
- Glick, B. R., Patten, C. L., Holguin, G., and Penrose, D. M. (1999). *Biochemical and Genetic Mechanisms Used by Plant Growth Promoting Bacteria*. London: Imperial College Press.
- Gordon, S. A., and Weber, R. P. (1951). Colorimetric estimation of indole acetic acid. *Plant Physiol.* 26, 192–195. doi: 10.1104/pp.26.1.192
- Haas, D., and Défago, G. (2005). Biological control of soil-borne pathogens by fluorescent *Pseudomonads*. *Nat. Rev. Microbiol.* 3, 307–319. doi: 10.1038/nrmicro1129
- Hallmann, J., and Berg, G. (2006). “Spectrum and population dynamics of bacterial root endophytes,” in *Microbial Root Endophytes*, eds B. Schulz, C. Boyle, and T. Sieber (Heidelberg: Springer), 15–31.
- Hardy, R. W. F., Holstern, R. D., Jackson, E. K., and Burns, R. C. (1968). The acetylene-ethylene assay for N₂ fixation: laboratory and field evaluation. *Plant Physiol.* 43, 1185–1207. doi: 10.1104/pp.43.8.1185
- Hausbeck, M. K., and Lamour, K. H. (2004). *Phytophthora capsici* on vegetable crops: research progress and management challenges. *Plant Dis.* 88, 1292–1303. doi: 10.1094/PDIS.2004.88.12.1292
- Hayward, A. C. (1960). A method for characterizing *Pseudomonas solanacearum*. *Nature* 186:405. doi: 10.1038/186405a0
- Heckel, D. G. (2012). Insecticide resistance after silent spring. *Science* 337, 1612–1614. doi: 10.1126/science.1226994
- Hossain, M. M., Sultana, F., Kubota, M., and Hyakumachi, M. (2008). Differential inducible defense mechanisms against bacterial speck pathogen in *Arabidopsis thaliana* by plant-growth-promoting-fungus *Penicillium* sp. GP16-2 and its cell free filtrate. *Plant Soil* 304, 227–239. doi: 10.1007/s11104-008-9542-3
- Hussain, S., Siddique, T., Saleem, M., Arshad, M., and Khalid, A. (2009). “Impact of pesticides on soil microbial diversity, enzymes, and biochemical reactions,” in *Advances in Agronomy*, Vol. 102, eds D. L. Sparks and S. H. du Pont (Amsterdam: Elsevier), 159–200.
- Idris, E. E. S., Iglesias, D. J., Talon, M., and Borriss, R. (2007). Tryptophan-dependent production of indole-3-acetic acid (IAA) affects level of plant growth promotion by *Bacillus amyloliquefaciens* FZB42. *Mol. Plant Microbe Interact.* 20, 619–626. doi: 10.1094/MPMI-20-6-0619
- Islam, M. R., Madhaiyan, M., Deka Boruah, H. P., Yim, W., Lee, G., Saravanan, V. S., et al. (2009). Characterization of plant growth-promoting traits of free-living diazotrophic bacteria and their inoculation effects on growth and nitrogen uptake of crop plants. *J. Microbiol. Biotechnol.* 19, 1213–1222. doi: 10.4014/jmb.0903.3028
- Islam, M. T., and Hossain, M. M. (2013). “Biological control of peronosporomycete phytopathogen by bacterial antagonist,” in *Bacteria in Agrobiology: Disease Management*, ed D. K. Maheshwari (Heidelberg: Springer), 167–218.
- Islam, M. T., and von Tiedemann, A. (2011). 2, 4-Diacetylphloroglucinol suppresses zoosporogenesis and impairs motility of *Peronosporomycete zoospores*. *World J. Microbiol. Biotechnol.* 27, 2071–2079. doi: 10.1007/s11274-011-0669-7
- Kamboh, A. A., Rajput, N., Rajput, I. R., Khaskheli, M., and Khaskheli, G. B. (2009). Biochemical properties of bacterial contaminants isolated from livestock vaccines. *Pak. J. Nutr.* 8, 578–581. doi: 10.3923/pjn.2009.578.581
- Kawulka, K. E., Sprules, T., Diaper, C. M., Whittal, R. M., McKay, R. T., Mercier, P., et al. (2004). Structure of Subtilisin A, a cyclic antimicrobial peptide from *Bacillus subtilis* with unusual sulfur to α -carbon cross-links: formation and reduction of α -Thio- α -Amino Acid derivatives. *Biochemistry* 43, 3385–3395. doi: 10.1021/bi0359527
- Kiba, T., Kudo, T., Kojima, M., and Sakakibar, H. (2011). Hormonal control of nitrogen acquisition: roles of auxin, abscisic acid, and cytokinin. *J. Exp. Bot.* 62, 1399–1409. doi: 10.1093/jxb/erq410
- Kim, Y. C., Jung, H., Kim, K. Y., and Park, S. K. (2008). An effective biocontrol bioformulation against *Phytophthora* blight of pepper using growth mixtures of combined chitinolytic bacteria under different field conditions. *Eur. J. Plant Pathol.* 120, 373–382. doi: 10.1007/s10658-007-9227-4
- Kirsop, B. E., and Doyle, A. (1991). *Maintenance of Microorganisms and Cultured Cells: A Manual of Laboratory Methods*, 2nd Edn. London: Academic Press.
- Kjeldahl, J. (1883). A new method for the estimation of nitrogen in organic compounds. *Z. Anal. Chem.* 22:366. doi: 10.1007/BF01338151
- Kloepper, J. W., Leong, J., Teintze, M., and Schroth, M. N. (1980). Enhanced plant growth by siderophores produced by plant growth promoting rhizobacteria. *Nature* 268, 885–886. doi: 10.1038/286885a0
- Konsoula, Z., and Liakopoulou-Kyriakides, M. (2006). Thermostable α -amylase production by *Bacillus subtilis* entrapped in calcium alginate gel capsules. *Enzyme Microb. Technol.* 39, 690–696. doi: 10.1016/j.enzmictec.2005.12.002
- Laguerre, G., Attard, M. R., Revoy, F., and Amarger, N. (1994). Rapid identification of Rhizobia by restriction fragment length polymorphism analysis of PCR amplified 16S rRNA genes. *Appl. Environ. Microbiol.* 60, 56–63.

- Lee, B. K., Kim, B. S., Chang, S. W., and Hwang, B. K. (2001). Aggressiveness to pumpkin cultivars of isolates of *Phytophthora capsici* from pumpkin and pepper. *Plant Dis.* 85, 497–500. doi: 10.1094/PDIS.2001.85.5.497
- Lim, J. H., and Kim, S. D. (2010). Biocontrol of *Phytophthora* blight of red pepper caused by *Phytophthora capsici* using *Bacillus subtilis* AH18 and *B. licheniformis* K11 formulations. *J. Korean Soc. Appl. Biol. Chem.* 53, 766–773. doi: 10.3839/jksabc.2010.116
- Liu, C. H., Chen, X., Liu, T. T., Lian, B., Gu, Y., Caer, V., et al. (2007). Study of the antifungal activity of *Acinetobacter baumannii* LCH001 in vitro and identification of its antifungal components. *Appl. Microbiol. Biotechnol.* 76, 459–466. doi: 10.1007/s00253-007-1010-0
- Lugtenberg, B., and Kamilova, F. (2009). Plant-growth-promoting rhizobacteria. *Annu. Rev. Microbiol.* 63, 541–556. doi: 10.1146/annurev.micro.62.081307.162918
- Maleki, M., Mokhtarnejad, L., and Mostafaeae, S. (2011). Screening of rhizobacteria for biological control of cucumber root and crown rot caused by *Phytophthora drechsleri*. *Plant Pathol. J.* 27, 78–84. doi: 10.5423/PPJ.2011.27.1.078
- Maniatis, T., Fritsch, E. F., and Sambrook, J. (1982). *Molecular Cloning: A Laboratory Manual*. New York, NY: Cold Spring Harbor Laboratory.
- Manjula, K., Kishore, G. K., and Podile, A. R. (2004). Whole cells of *Bacillus subtilis* AF1 proved more effective than cell-free and chitinase-based formulations in biological control of citrus fruit rot and groundnut rust. *Can. J. Microbiol.* 50, 737–744. doi: 10.1139/W04-058
- Merchant, I. A., and Packer, R. A. (1999). *Veterinary Bacteriology and Virology*, 8th Edn. New Delhi: CBS Publishers and Distributors.
- Messiha, N. A. S., van Diepeningen, A. D., Farag, N. S., Abdallah, S. A., Janse, J. D., van Bruggen, A. H. C., et al. (2007). *Stenotrophomonas maltophilia*: a new potential biocontrol agent of *Ralstonia solanacearum*, causal agent of potato brown rot. *Eur. J. Plant Pathol.* 118, 211–225. doi: 10.1007/s10658-007-9136-6
- Mousivand, M., Jouzani, G. S., Monazah, M., and Kowsari, M. (2012). Characterization and antagonistic potential of some native biofilm-forming and surfactant-producing *Bacillus subtilis* strains against six pathotypes of *Rhizoctonia solani*. *J. Plant Pathol.* 94, 171–180. doi: 10.4454/jpp.v94i1.017
- Nakayama, T., Homma, Y., Hashidoko, Y., Mizutani, J., and Tahara, S. (1999). Possible role of xanthobaccins produced by *Stenotrophomonas* sp. strain SB-K88 in suppression of sugar beet damping-off disease. *Appl. Environ. Microbiol.* 65, 4334–4339.
- Ng, L. C., Sariah, M., Sariam, O., Radziah, O., and Abi, M. A. Z. (2012). Rice seed bacterization for promoting germination and seedling growth under aerobic cultivation system. *Aust. J. Crop. Sci.* 6, 170–175.
- Patten, C. L., and Glick, B. R. (2002). Regulation of indole acetic acid production in *Pseudomonas putida* GR12-2 by tryptophan and the stationary-phase sigma factor RpoS. *Can. J. Microbiol.* 48, 635–642. doi: 10.1139/w02-053
- Rahman, A., Sitepu, I. R., Tang, S. Y., and Hashidoko, Y. (2010). Salkowski's reagent test as a primary screening index for functionalities of rhizobacteria isolated from wild dipterocarp saplings growing naturally on medium-strongly acidic tropical peat soil. *Biosci. Biotechnol. Biochem.* 74, 2202–2208. doi: 10.1271/bbb.100360
- Rajat, R. M., Ninama, G. L., Mistry, K., Parmar, R., Patel, K., and Vegad, M. M. (2012). Antibiotic resistance pattern in *Pseudomonas aeruginosa* species isolated at a tertiary care hospital. *Ahmadabad. Natl. J. Med. Res.* 2, 156–159.
- Reysenbach, A. L., Giver, L. J., Wickham, G. S., and Pace, N. R. (1992). Differential amplification of rRNA genes by polymerase chain reaction. *Appl. Environ. Microbiol.* 58, 3417–3418.
- Riungu, G. M., Muthorni, J. W., Narla, R. D., Wagacha, J. M., and Gathumbi, J. K. (2008). Management of Fusarium head blight of wheat and deoxynivalenol accumulation using antagonistic microorganisms. *Plant Pathol. J.* 7, 13–19. doi: 10.3923/ppj.2008.13.19
- Ryan, R. P., Monchy, S., Cardinale, M., Taghavi, S., Crossman, L., Avison, M. B., et al. (2009). The versatility and adaptation of bacteria from the genus *Stenotrophomonas*. *Nat. Rev.* 7, 514–525. doi: 10.1038/nrmicro2163
- Shirzad, A., Fallahzadeh-Mamaghani, V., and Pazhouhandeh, M. (2012). Antagonistic potential of fluorescent pseudomonads and control of crown and root rot of cucumber caused by *Phytophthora drechsleri*. *Plant Pathol. J.* 28, 1–9. doi: 10.5423/PPJ.OA.05.2011.0100
- Simmons, J. S. (1926). A culture medium for differentiating organisms of typhoid-colon aerogenes groups and for isolation of certain fungi. *J. Infect. Dis.* 39:209. doi: 10.1093/infdis/39.3.209
- Somasegaran, P., and Hoben, H. J. (1994). *Handbook for Rhizobia: Methods in Legume-Rhizobium Technology*. New York, NY: Springer-Verlag.
- Stein, T. (2005). *Bacillus subtilis* antibiotics: structures, syntheses and specific functions. *Mol. Microbiol.* 56, 845–857. doi: 10.1111/j.1365-2958.2005.04587.x
- Tajini, F., MustaphaTrabelsi, M., and Drevon, J. J. (2012). Combined inoculation with *Glomus intraradices* and *Rhizobium tropici* CIAT899 increases phosphorus use efficiency for symbiotic nitrogen fixation in common bean (*Phaseolus vulgaris* L.). *Saudi J. Biol. Sci.* 19, 157–163. doi: 10.1016/j.sjbs.2011
- Turnbull, G. A., Morgan, J. A. W., Whipps, J. M., and Saunders, J. R. (2001). The role of bacterial motility in the survival and spread of *Pseudomonas fluorescens* in soil and in the attachment and colonisation of wheat roots. *FEMS Microbiol. Ecol.* 36, 21–31. doi: 10.1111/j.1574-6941.2001.tb00822.x
- Verma, J. P., Tiwari, K. N., and Kumar, A. (2013). Effect of indigenous *Mesorhizobium* spp. and plant growth promoting rhizobacteria on yields and nutrients uptake of chickpea (*Cicer arietinum* L.) under sustainable agriculture. *Ecol. Eng.* 51, 282–286. doi: 10.1016/j.ecoleng.2012.12.022
- Vincent, J. M., and Humphrey, B. (1970). Taxonomically significant group antigens in *Rhizobium*. *J. Gen. Microbiol.* 63, 379–382. doi: 10.1099/00221287-63-3-379
- Weisskopf, L., Heller, S., and Eberl, L. (2011). Burkholderia species are major inhabitants of white lupin cluster roots. *Appl. Environ. Microbiol.* 77, 7715–7720. doi: 10.1128/AEM.05845-11
- Yasmin, S., Bakar, M. A. R., Malik, K. A., and Hafeez, F. Y. (2004). Isolation, characterization and beneficial effects of rice associated plant growth promoting bacteria from Zanzibar soils. *J. Basic Microbiol.* 44, 241–252. doi: 10.1002/jobm.200310344
- Zahid, M., Abbasi, M. K., Hameed, S., and Rahim, N. (2015). Isolation and identification of indigenous plant growth promoting rhizobacteria from Himalayan region of Kashmir and their effect on improving growth and nutrient contents of maize (*Zea mays* L.). *Front. Microbiol.* 6:207. doi: 10.3389/fmicb.2015.00207
- Zhang, S., White, T. L., Martinez, M. C., McInroy, J. A., Kloepper, J. W., and Klassen, W. (2010). Evaluation of plant growth-promoting rhizobacteria for control of *Phytophthora* blight on squash under greenhouse conditions. *Biol. Control* 53, 129–135. doi: 10.1016/j.biocontrol.2009.10.015
- Zhang, Z., and Yuen, G. Y. (2000). The role of chitinase production by *Stenotrophomonas maltophilia* strain C3 in biological control of *Bipolaris sorokiniana*. *Phytopathology* 90, 384–389. doi: 10.1094/PHYTO.2000.90.4.384

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Islam, Akanda, Prova, Islam and Hossain. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.