

Protocol

Isolation of Mitochondria from Tissue Culture Cells

David A. Clayton¹ and Gerald S. Shadel^{2,3,4}

¹Janelia Farm Research Campus, Howard Hughes Medical Institute, Ashburn, Virginia 20147-2408; ²Department of Pathology, Yale University School of Medicine, New Haven, Connecticut 06520-8023; ³Department of Genetics, Yale University School of Medicine, New Haven, Connecticut 06520-8023

The number of mitochondria per cell varies substantially from cell line to cell line. For example, human HeLa cells contain at least twice as many mitochondria as smaller mouse L cells. This protocol starts with a washed cell pellet of 1–2 mL derived from $\sim 10^9$ cells grown in culture. The cells are swollen in a hypotonic buffer and ruptured with a Dounce or Potter-Elvehjem homogenizer using a tight-fitting pestle, and mitochondria are isolated by differential centrifugation.

MATERIALS

It is essential that you consult the appropriate Material Safety Data Sheets and your institution's Environmental Health and Safety Office for proper handling of equipment and hazardous materials used in this protocol.

RECIPES: Please see the end of this protocol for recipes indicated by <R>. Additional recipes can be found online at <http://cshprotocols.cshlp.org/site/recipes>.

Reagents

Cell pellet derived from $1\text{--}5 \times 10^9$ tissue culture cells

MS homogenization buffer (1 \times and 2.5 \times) <R>

MS homogenization buffer is an iso-osmotic buffer used to maintain the tonicity of the organelles and prevent agglutination.

RSB hypo buffer <R>

RSB is a hypotonic buffer used for swelling tissue culture cells.

Equipment

Centrifuge tubes

Dounce homogenizer (15 mL) with a tight-fitting B pestle or Potter-Elvehjem homogenizer (5 mL) with a Teflon pestle (see Steps 1 and 3)

Phase contrast microscope

⁴Correspondence: gerald.shadel@yale.edu

METHOD

The solutions, tubes, and homogenizer should be prechilled on ice. All centrifugation steps are at 4°C.

1. Resuspend the cell pellet in 11 mL of ice-cold RSB hypo buffer and transfer the cells to a 15-mL Dounce homogenizer.

Alternatively, as described by Frezza et al. (2007), resuspend the cell pellet in 9 mL of ice-cold RSB hypo buffer and transfer 3 mL of the cells at a time to a 5-mL Potter-Elvehjem homogenizer with a Teflon pestle.

2. Allow the cells to swell for 5–10 min. Check the progress of the swelling using a phase-contrast microscope.
3. Break open the swollen cells with several strokes of the B pestle. For each stroke, press the pestle straight down the tube, maintaining a firm, steady pressure.

If a Potter-Elvehjem homogenizer is used in Step 1, then break open the cells with the Teflon pestle rotating at ~1600 rpm.

4. Check the degree of homogenization with a phase-contrast microscope.

Naked nuclei (smooth spheres with obvious nucleoli inside), smaller organelles (dark, granular objects), and a small number of unbroken cells (large spheres with a granular appearance) should be present if cell lysis was successful. Eight to nine naked nuclei for every whole cell is a very good result. Trying for anything better usually results in increasing the number of damaged nuclei, which increases the number of mitochondria trapped in the nuclear pellet during the first centrifugation.

See Troubleshooting.

5. Immediately add 8 mL of 2.5× MS homogenization buffer to give a final concentration of 1× MS homogenization buffer. Cover the top of the homogenizer with Parafilm and mix by inverting a couple of times. (Save a portion of the homogenate if marker enzyme assays are to be performed later.)
6. Transfer the homogenate to a centrifuge tube for differential centrifugation. Rinse the homogenizer with a small amount of 1× MS homogenization buffer and add it to the homogenate. Bring the volume to 30 mL with 1× homogenization MS buffer.
7. Centrifuge the homogenate at 1300g for 5 min to remove nuclei, unbroken cells, and large membrane fragments.
8. Pour the supernatant into a clean centrifuge tube.

The top of the pellet will be loose, so be careful not to collect it with the supernatant.

9. Repeat Steps 6 and 7 two more times.
10. Transfer the supernatant to a clean centrifuge tube and pellet the mitochondria at 7,000g–17,000g for 15 min.
11. Discard the supernatant and wipe out the inside of the tube with a Kimwipe.
12. Wash the mitochondria by resuspending the pellet in 1× MS buffer and repeating the 7,000g–17,000g sedimentation.

*This wash is not necessary if a density gradient will be performed (see Protocol: **Purification of Mitochondria by Sucrose Step Density Gradient Centrifugation** [Clayton and Shadel 2014]).*

13. Discard the supernatant and resuspend the pellet in a buffer suitable for subsequent work.

The mitochondria can be stored at –80°C for at least 1 yr for some purposes (e.g., protein isolation).

TROUBLESHOOTING

Problem (Step 4): Too many or not enough cells have lysed.

Solution: Homogenization works best if the cells are resuspended in at least 5–10× the volume of the cell pellet and if the cell suspension occupies at least half the volume of the homogenizer. Homogenization should be performed as quickly as possible because it is performed in a hypotonic buffer. The

Dounce homogenizer disrupts swollen tissue culture cells by pressure change. As the pestle is pressed down, pressure around the cell increases. When the cell slips past the end of the pestle, the sudden decrease in pressure causes the cell to rupture. If the pestle is very tight fitting, there may be some mechanical breakage as well. If an excessive number of strokes are needed for good cell breakage, a tighter-fitting homogenizer is needed.

DISCUSSION

This basic protocol can be modified to suit special purposes. For example, if the mitochondria are being purified to isolate mitochondrial DNA, contamination with nuclei, not the small organelles, is a problem and the following modifications could be made: Harvest the cells in stationary growth phase when the fewest cells will be actively dividing, substitute CaCl_2 for MgCl_2 in the RSB hypo buffer to stabilize the nuclear membrane, omit washing the mitochondrial pellet, omit any density gradient purification, resuspend and lyse the mitochondrial pellet from Step 10, and purify the mitochondrial DNA from any remaining nuclear DNA (Hudson et al. 1968).

RECIPES

MS Homogenization Buffer (1×)

210 mM mannitol
70 mM sucrose
5 mM Tris-HCl (pH 7.5)
1 mM EDTA (pH 7.5)

The buffer should be ice cold before use.

MS Homogenization Buffer (2.5×)

525 mM mannitol
175 mM sucrose
12.5 mM Tris-HCl (pH 7.5)
2.5 mM EDTA (pH 7.5)

The buffer should be ice cold before use.

RSB Hypo Buffer

10 mM NaCl
1.5 mM MgCl_2
10 mM Tris-HCl (pH 7.5)

The buffer should be ice cold before use.

REFERENCES

- Clayton DA, Shadel GS. 2014. Purification of mitochondria by sucrose step density gradient centrifugation. *Cold Spring Harb Protoc* doi: 10.1101/pdb.prot080028.
- Frezza C, Cipolat S, Scorrano L. 2007. Organelle isolation: Functional mitochondria from mouse liver, muscle and cultured fibroblasts. *Nature Protocols* 2: 287–295.
- Hudson B, Clayton DA, Vinograd J. 1968. Complex mitochondrial DNA. *Cold Spring Harbor Symp Quant Biol* 33: 435–442.



Cold Spring Harbor Protocols

Isolation of Mitochondria from Tissue Culture Cells

David A. Clayton and Gerald S. Shadel

Cold Spring Harb Protoc; doi: 10.1101/pdb.prot080002

Email Alerting Service

Receive free email alerts when new articles cite this article - [click here](#).

Subject Categories

Browse articles on similar topics from *Cold Spring Harbor Protocols*.

[Cell Biology, general](#) (1379 articles)
[Subcellular Fractionation](#) (87 articles)

To subscribe to *Cold Spring Harbor Protocols* go to:
<http://cshprotocols.cshlp.org/subscriptions>
