

ISOMETRIC DILATIONS FOR INFINITE SEQUENCES OF NONCOMMUTING OPERATORS

GELU POPESCU

ABSTRACT. This paper develops a dilation theory for $\{T_n\}_{n=1}^\infty$ an infinite sequence of noncommuting operators on a Hilbert space, when the matrix $[T_1, T_2, \dots]$ is a contraction. A Wold decomposition for an infinite sequence of isometries with orthogonal final spaces and a minimal isometric dilation for $\{T_n\}_{n=1}^\infty$ are obtained. Some theorems on the geometric structure of the space of the minimal isometric dilation and some consequences are given. This results are used to extend the Sz.-Nagy-Foiaş lifting theorem to this noncommutative setting.

This paper is a continuation of [5] and develops a dilation theory for an infinite sequence $\{T_n\}_{n=1}^\infty$ of noncommuting operators on a Hilbert space \mathcal{H} when $\sum_{n=1}^\infty T_n T_n^* \leq I_{\mathcal{H}}$ ($I_{\mathcal{H}}$ is the identity on \mathcal{H}).

Many of the results and techniques in dilation theory for one operator [8] and also for two operators [3, 4] are extended to this setting.

First we extend Wold decomposition [8, 4] to the case of an infinite sequence $\{V_n\}_{n=1}^\infty$ of isometries with orthogonal final spaces.

In §2 we obtain a minimal isometric dilation for $\{T_n\}_{n=1}^\infty$ by extending the Schaffer construction in [6, 4]. Using these results we give some theorems on the geometric structure of the space of the minimal isometric dilation. Finally, we give some sufficient conditions on a sequence $\{T_n\}_{n=1}^\infty$ to be simultaneously quasi-similar to a sequence $\{R_n\}_{n=1}^\infty$ of isometries on a Hilbert space \mathcal{H} with $\sum_{n=1}^\infty R_n R_n^* = I_{\mathcal{H}}$.

In §3 we use the above-mentioned theorems to obtain the Sz.-Nagy-Foiaş lifting theorem [7, 8, 1, 4] in our setting.

In a subsequent paper we will use the results of this paper for studying the “characteristic function” associated to a sequence $\{T_n\}_{n=1}^\infty$ with $\sum_{n=1}^\infty T_n T_n^* \leq I_{\mathcal{H}}$.

1

Throughout this paper Λ stands for the set $\{1, 2, \dots, k\}$ ($k \in \mathbf{N}$) or the set $\mathbf{N} = \{1, 2, \dots\}$.

Received by the editors September 1, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 47A20; Secondary 47A45.

Key words and phrases. Isometric dilation, Wold decomposition, lifting theorem.

©1989 American Mathematical Society
0002-9947/89 \$1.00 + \$.25 per page

For every $n \in \mathbb{N}$ let $F(n, \Lambda)$ be the set of all functions from the set $\{1, 2, \dots, n\}$ to Λ and

$$\mathcal{F} = \bigcup_{n=0}^{\infty} F(n, \Lambda), \quad \text{where } F(0, \Lambda) = \{0\}.$$

Let \mathcal{H} be a Hilbert space and $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ be a sequence of isometries on \mathcal{H} . For any $f \in F(n, \Lambda)$ we denote by V_f the product $V_{f(1)}V_{f(2)} \cdots V_{f(n)}$ and $V_0 = I_{\mathcal{H}}$.

A subspace $\mathcal{L} \subset \mathcal{H}$ will be called *wandering* for the sequence \mathcal{V} if for any distinct functions $f, g \in \mathcal{F}$ we have

$$V_f \mathcal{L} \perp V_g \mathcal{L} \quad (\perp \text{ means orthogonal}).$$

In this case we can form the orthogonal sum

$$M_{\mathcal{V}}(\mathcal{L}) = \bigoplus_{f \in \mathcal{F}} V_f \mathcal{L}.$$

A sequence $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ of isometries on \mathcal{H} is called a Λ -orthogonal shift if there exists in \mathcal{H} a subspace \mathcal{L} , which is wandering for \mathcal{V} and such that $\mathcal{H} = M_{\mathcal{V}}(\mathcal{L})$.

This subspace \mathcal{L} is uniquely determined by \mathcal{V} : indeed we have $\mathcal{L} = \mathcal{H} \ominus (\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H})$. The dimension of \mathcal{L} is called the multiplicity of the Λ -orthogonal shift. One can show, by an argument similar to the classical unilateral shift, that a Λ -orthogonal shift is determined up to unitary equivalence by its multiplicity. It is easy to see that for $\Lambda = \{1\}$ we find again the classical unilateral shift.

Let us make some simple remarks whose proofs will be omitted.

Remark 1.1. If $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ is a Λ -orthogonal shift on \mathcal{H} , with the wandering subspace \mathcal{L} , then for any $n \in \mathbb{N}$, $\lambda \in \Lambda$ and $f \in F(n, \Lambda)$ we have

(a)

$$V_\lambda^* V_f = \begin{cases} V_{f(2)} V_{f(3)} \cdots V_{f(n)} & \text{if } f(1) = \lambda, \\ 0 & \text{if } f(1) \neq \lambda, \end{cases}$$

and $V_\lambda^* \ell = 0$ ($\ell \in \mathcal{L}$).

(b) $\sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* + P_{\mathcal{L}} = I_{\mathcal{H}}$, where $P_{\mathcal{L}}$ stands for the orthogonal projection from \mathcal{H} into \mathcal{L} .

Remark 1.2. If $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ is a Λ -orthogonal shift on \mathcal{H} then

- (a) $\lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* h\|^2 = 0$, for any $h \in \mathcal{H}$.
- (b) $V_\lambda^{*k} \rightarrow 0$ (strongly) as $k \rightarrow \infty$, for any $\lambda \in \Lambda$.
- (c) There exists no nonzero reducing subspace $\mathcal{H}_0 \subset \mathcal{H}$ for each V_λ ($\lambda \in \Lambda$) such that $(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda V_\lambda^*)|_{\mathcal{H}_0} = 0$.

Let us consider a model Λ -orthogonal shift.

Form the Hilbert space

$$l^2(\mathcal{F}, \mathcal{H}) = \left\{ (h_f)_{f \in \mathcal{F}}; \sum_{f \in \mathcal{F}} \|h_f\|^2 < \infty, h_f \in \mathcal{H} \right\}.$$

We embed \mathcal{H} in $l^2(\mathcal{F}, \mathcal{H})$ as a subspace, by identifying the element $h \in \mathcal{H}$ with the element $(h_f)_{f \in \mathcal{F}}$, where $h_0 = h$ and $h_f = 0$ for any $f \in \mathcal{F}$, $f \neq 0$.

For each $\lambda \in \Lambda$ we define the operator S_λ on $l^2(\mathcal{F}, \mathcal{H})$ by $S_\lambda((h_f)_{f \in \mathcal{F}}) = (h'_g)_{g \in \mathcal{F}}$, where $h'_0 = 0$ and for $g \in F(n, \Lambda)$ ($n \geq 1$)

$$h'_g = \begin{cases} h_0 & \text{if } g \in F(1, \Lambda) \text{ and } g(1) = \lambda, \\ h_f & \text{if } g \in F(n, \Lambda) \text{ (} n \geq 2 \text{), } f \in F(n-1, \Lambda) \text{ and } g(1) = \lambda, \\ & g(2) = f(1), g(3) = f(2), \dots, g(n) = f(n-1), \\ 0 & \text{otherwise.} \end{cases}$$

It is easy to see that $\{S_\lambda\}_{\lambda \in \Lambda}$ is the Λ -orthogonal shift, acting on $l^2(\mathcal{F}, \mathcal{H})$, with the wandering subspace \mathcal{H} .

This model plays an important role in this paper. The following theorem is our version of Wold decomposition for a sequence of isometries.

Theorem 1.3. *Let $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ be a sequence of isometries on a Hilbert space \mathcal{H} such that $\sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* \leq I_{\mathcal{H}}$.*

Then \mathcal{H} decomposes into an orthogonal sum $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$ such that \mathcal{H}_0 and \mathcal{H}_1 reduce each operator V_λ ($\lambda \in \Lambda$) and we have $(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda V_\lambda^)|_{\mathcal{H}_1} = 0$ and $\{V_\lambda|_{\mathcal{H}_0}\}_{\lambda \in \Lambda}$ is a Λ -orthogonal shift acting on \mathcal{H}_0 .*

This decomposition is uniquely determined, indeed we have

$$\mathcal{H}_1 = \bigcap_{n=0}^{\infty} \left(\bigoplus_{f \in F(n, \Lambda)} V_f \mathcal{H} \right);$$

$\mathcal{H}_0 = M_{\mathcal{F}}(\mathcal{L})$, where $\mathcal{L} = \mathcal{H} \ominus (\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H})$.

Proof. It is easy to see that the subspace $\mathcal{L} = \mathcal{H} \ominus (\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H})$ is wandering for \mathcal{V} .

Now let $\mathcal{H}_0 = M_{\mathcal{F}}(\mathcal{L})$ and $\mathcal{H}_1' = \mathcal{H} \ominus \mathcal{H}_0$. Observe that $k \in \mathcal{H}_1'$ if and only if $k \perp \bigoplus_{f \in \mathcal{F}_n} V_f \mathcal{L}$ for every $n \in \mathbb{N}$, where \mathcal{F}_n stands for $\bigcup_{k=0}^n F(k, \Lambda)$.

We have

$$\begin{aligned} \mathcal{L} \oplus \left(\bigoplus_{f \in F(1, \Lambda)} V_f \mathcal{L} \right) \oplus \cdots \oplus \left(\bigoplus_{g \in F(n, \Lambda)} V_g \mathcal{L} \right) &= \left[\mathcal{H} \ominus \left(\bigoplus_{f \in F(1, \Lambda)} V_f \mathcal{H} \right) \right] \\ &\oplus \left[\left(\bigoplus_{f \in F(1, \Lambda)} V_f \mathcal{H} \right) \ominus \left(\bigoplus_{f \in F(2, \Lambda)} V_f \mathcal{H} \right) \right] \\ &\oplus \cdots \oplus \left[\left(\bigoplus_{f \in F(n, \Lambda)} V_f \mathcal{H} \right) \ominus \left(\bigoplus_{f \in F(n+1, \Lambda)} V_f \mathcal{H} \right) \right] \\ &= \mathcal{H} \ominus \left(\bigoplus_{f \in F(n+1, \Lambda)} V_f \mathcal{H} \right). \end{aligned}$$

Thus $k \in \mathcal{H}'_1$ if and only if $k \in \bigoplus_{f \in F(n+1, \Lambda)} V_f \mathcal{H}$ for every $n \in \mathbf{N}$. Since

$$\bigoplus_{f \in F(n, \Lambda)} V_f \mathcal{H} \supset \bigoplus_{f \in F(n+1, \Lambda)} V_f \mathcal{H} \quad (n \in \mathbf{N})$$

it follows that

$$\mathcal{H}'_1 = \bigcap_{n=0}^{\infty} \left(\bigoplus_{f \in F(n, \Lambda)} V_f \mathcal{H} \right) = \mathcal{H}_1.$$

Let us notice that

$$\begin{aligned} V_\lambda \mathcal{H}_1 &\subset \bigcap_{n=0}^{\infty} \left(\bigoplus_{f \in F(n, \Lambda)} V_\lambda V_f \mathcal{H} \right) \subset \bigcap_{n=0}^{\infty} \left(\bigoplus_{g \in F(n+1, \Lambda)} V_g \mathcal{H} \right) = \mathcal{H}_1, \\ V_\lambda^* \mathcal{H}_1 &\subset \bigcap_{n=1}^{\infty} \left(V_\lambda^* \left(\bigoplus_{\substack{g \in F(n, \Lambda) \\ g(1)=\lambda}} V_g \mathcal{H} \right) \right) = \bigcap_{n=1}^{\infty} \left(\bigoplus_{f \in F(n-1, \Lambda)} V_f \mathcal{H} \right) = \mathcal{H}_1. \end{aligned}$$

Therefore \mathcal{H}_1 reduces each V_λ ($\lambda \in \Lambda$). Hence \mathcal{H}_0 also reduces each V_λ ($\lambda \in \Lambda$).

Since $\mathcal{H}_1 \subset \bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H}$ it follows that $(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda V_\lambda^*)|_{\mathcal{H}_1} = 0$. The fact that $\{V_\lambda|_{\mathcal{H}_0}\}_{\lambda \in \Lambda}$ is a Λ -orthogonal shift is obvious. The uniqueness of the decomposition follows by an argument similar to the classical Wold decomposition [8, Chapter I, Theorem 1.1]. The proof is completed.

Remark 1.4. The subspaces $\mathcal{H}_0, \mathcal{H}_1$ from Wold decomposition can be described as follows:

$$\begin{aligned} \mathcal{H}_0 &= \left\{ k \in \mathcal{H} : \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* k\|^2 = 0 \right\}, \\ \mathcal{H}_1 &= \left\{ k \in \mathcal{H} : \sum_{f \in F(n, \Lambda)} \|V_f^* k\|^2 = \|k\|^2 \text{ for every } n \in \mathbf{N} \right\}. \end{aligned}$$

We call the sequence $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ in Theorem 1.3 *pure* if $\mathcal{K}_1 = 0$, that is, if \mathcal{V} is a Λ -orthogonal shift on \mathcal{H} .

2

Let $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ a sequence of contractions on a Hilbert space \mathcal{H} such that $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \leq I_{\mathcal{H}}$.

We say that a sequence $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ of isometries on a Hilbert space $\mathcal{K} \supset \mathcal{H}$ is a minimal isometric dilation of \mathcal{T} if the following conditions hold:

- (a) $\sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* \leq I_{\mathcal{K}}$.
- (b) \mathcal{H} is invariant for each V_λ^* ($\lambda \in \Lambda$) and $V_\lambda^*|_{\mathcal{H}} = T_\lambda^*$ ($\lambda \in \Lambda$).
- (c) $\mathcal{K} = \bigvee_{f \in \mathcal{F}} V_f \mathcal{H}$.

Let D_* on \mathcal{H} and D on $\bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda$ (\mathcal{H}_λ is a copy of \mathcal{H}) be the positive operators uniquely defined by $D_* = (I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} T_\lambda T_\lambda^*)^{1/2}$ and $D = D_T$, where T stands for the matrix $[T_1, T_2, \dots]$ and $D_T = (I - T^*T)^{1/2}$.

Let us denote $\mathcal{D}_* = \overline{D_* \mathcal{H}}$ and $\mathcal{D} = \overline{D(\bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda)}$.

Theorem 2.1. *For every sequence $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ of noncommuting operators on a Hilbert space \mathcal{H} such that $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \leq I_{\mathcal{H}}$, there exists a minimal isometric dilation $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ on a Hilbert space $\mathcal{K} \supset \mathcal{H}$, which is uniquely determined up to an isomorphism.*

Proof. Let us consider the Hilbert space $\mathcal{K} = \mathcal{H} \oplus l^2(\mathcal{F}, \mathcal{D})$. We embed \mathcal{H} and \mathcal{D} into \mathcal{K} in a natural way. For each $\lambda \in \Lambda$ we define the isometry $V_\lambda: \mathcal{K} \rightarrow \mathcal{K}$ by the relation

$$(2.1) \quad V_\lambda(h \oplus (d_f)_{f \in \mathcal{F}}) = T_\lambda h \oplus (D(\underbrace{0, \dots, 0}_{\lambda-1 \text{ times}}, h, 0, \dots) + S_\lambda(d_f)_{f \in \mathcal{F}})$$

where $\{S_\lambda\}_{\lambda \in \Lambda}$ is Λ -orthogonal shift on $l^2(\mathcal{F}, \mathcal{D})$ (see §1).

Obviously, for any $\lambda, \mu \in \Lambda, \lambda \neq \mu$ we have $\text{range } S_\lambda \perp \text{range } S_\mu$ and

$$(T_\mu^* T_\lambda h, h') = -(D^2(\underbrace{0, \dots, 0}_{\lambda-1 \text{ times}}, h, 0, \dots), (\underbrace{0, \dots, 0}_{\mu-1 \text{ times}}, h', 0, \dots)).$$

Hence, taking into account (2.1), it follows that

$$\text{range } V_\lambda \perp \text{range } V_\mu \quad (\lambda, \mu \in \Lambda, \lambda \neq \mu)$$

therefore $\sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* \leq I_{\mathcal{K}}$.

It is easy to show that \mathcal{H} is invariant for each V_λ^* ($\lambda \in \Lambda$) and $V_\lambda^*|_{\mathcal{H}} = T_\lambda^*$ ($\lambda \in \Lambda$).

Finally, we verify that $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ is the minimal isometric dilation of \mathcal{T} .

Let $\mathcal{H}_1 = \mathcal{H} \vee (\bigvee_{f \in \mathcal{F}(1, \Lambda)} V_f \mathcal{H})$ and

$$\mathcal{H}_n = \mathcal{H}_{n-1} \vee \left(\bigvee_{f \in \mathcal{F}(1, \Lambda)} V_f \mathcal{H}_{n-1} \right) \quad \text{if } n \geq 2.$$

It is easy to see that $\mathcal{H}_1 = \mathcal{H} \oplus \mathcal{D}$ and

$$\mathcal{H}_n = \mathcal{H} \oplus \mathcal{D} \oplus \left(\bigoplus_{f \in F(1, \Lambda)} S_f \mathcal{D} \right) \oplus \cdots \oplus \left(\bigoplus_{f \in F(n-1, \Lambda)} S_f \mathcal{D} \right) \quad \text{if } n \geq 2.$$

Clearly $\mathcal{H}_n \subset \mathcal{H}_{n+1}$ and we have

$$\bigvee_1^\infty \mathcal{H}_n = \mathcal{H} \oplus M_{\mathcal{F}}(\mathcal{D}) = \mathcal{H} \oplus l^2(\mathcal{F}, \mathcal{D}) = \mathcal{K}.$$

Therefore $\mathcal{K} = \bigvee_{f \in \mathcal{F}} V_f \mathcal{H}$.

Following Theorem 4.1 in [8, Chapter I] it is easy to show that the minimal isometric dilation \mathcal{V} of \mathcal{T} is unique up to a unitary operator. To be more precise, let $\mathcal{V}' = \{V'_\lambda\}_{\lambda \in \Lambda}$ be another minimal isometric dilation of \mathcal{T} , on a Hilbert space $\mathcal{H}' \supset \mathcal{H}$. Then there exists a unitary operator $U: \mathcal{H} \rightarrow \mathcal{H}'$ such that $V'_\lambda U = UV_\lambda$ ($\lambda \in \Lambda$) and $Uh = h$ for every $h \in \mathcal{H}$.

This completes the proof.

Remark 2.2. For each $\lambda \in \Lambda$, $\bar{V}_\lambda^{*n} \rightarrow 0$ (strongly) as $n \rightarrow \infty$ if and only if $T_\lambda^{*n} \rightarrow 0$ (strongly) as $n \rightarrow \infty$.

From this remark and Theorem 2.1 one can easily deduce Proposition 1.1 in [5].

The following is a generalization of [2] or Theorem 1.2 in [8, Chapter II].

Proposition 2.3. Let $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ be the minimal isometric dilation of $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$. Then \mathcal{V} is pure if and only if

$$(2.2) \quad \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2 = 0$$

for any $h \in \mathcal{H}$.

Proof. Assume that \mathcal{V} is pure. Then, by Theorem 1.3 it follows that \mathcal{V} is a Λ -orthogonal shift on the space $\mathcal{K} \supset \mathcal{H}$ of the minimal isometric dilation of \mathcal{T} .

Taking into account Remark 1.2 and the fact that for each $f \in \mathcal{F}$, $V_f^*|_{\mathcal{H}} = T_f^*$, we have

$$\lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2 = \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* h\|^2 = 0 \quad \text{for any } h \in \mathcal{H}.$$

Conversely, assume that (2.2) holds. We claim that

$$(2.3) \quad \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* k\|^2 = 0 \quad \text{for any } k \in \mathcal{K} = \bigvee_{f \in \mathcal{F}} V_f \mathcal{H}.$$

By (2.2) we have

$$\lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* h\|^2 = 0 \quad (h \in \mathcal{H}).$$

For each $k \in \bigvee_{f \in \mathcal{F}; f \neq 0} V_f \mathcal{H}$ and any $\varepsilon > 0$, there exists

$$k_\varepsilon = \sum_{g \in \mathcal{F}; g \neq 0} V_g h_g \quad (h_g \in \mathcal{H})$$

such that $\|k - k_\varepsilon\| < \varepsilon$. (Here \sum' stands for a finite sum.)

Since the isometries V_λ ($\lambda \in \Lambda$) have orthogonal final spaces, it follows that

$$\lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* k\|^2 = \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* (k - k_\varepsilon)\|^2 \leq \|k - k_\varepsilon\|^2 < \varepsilon^2,$$

for any $\varepsilon > 0$. Thus, (2.3) holds and by Remark 1.4 we have that \mathcal{V} is pure. This completes the proof.

Corollary 2.4. *If $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \leq rI_{\mathcal{H}}$, $r < 1$, then the minimal isometric dilation of $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ is pure.*

Now let us establish when the minimal isometric dilation $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ cannot contain a Λ -orthogonal shift. The notations being the same as above we have

Proposition 2.5. $\sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* = I_{\mathcal{H}}$ if and only if $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* = I_{\mathcal{H}}$.

Proof. (\Rightarrow) Since $V_\lambda^*|_{\mathcal{H}} = T_\lambda^*$ ($\lambda \in \Lambda$) it follows that $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* h = h$ ($h \in \mathcal{H}$).

(\Leftarrow) If $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* = I_{\mathcal{H}}$ then $\sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2 = \|h\|^2$ for any $n \in \mathbb{N}$ and $h \in \mathcal{H}$. Taking into account Theorem 1.3 let us assume that there exists $k \in \mathcal{H} \ominus (\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H})$, $k \neq 0$. Using Remark 1.4 it follows that

$$(2.4) \quad \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|V_f^* k\|^2 = 0.$$

On the other hand, since

$$\mathcal{H} = \mathcal{H} \vee \left(\bigvee_{\substack{f \in \mathcal{F} \\ f \neq 0}} V_f \mathcal{H} \right) \quad \text{and} \quad \bigvee_{\substack{f \in \mathcal{F} \\ f \neq 0}} V_f \mathcal{H} \subset \bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H}$$

it follows that $k \in \mathcal{H}$ and by (2.4) that $\lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|T_f^* k\|^2 = 0$, contradiction. Thus we have $\sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* = I_{\mathcal{H}}$ and the proof is complete.

Dropping out the minimality condition in the definition of the isometric dilation of a sequence $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$, we can prove the following.

Proposition 2.6. *For any sequence $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ of operators on a Hilbert space \mathcal{H} such that $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \leq I_{\mathcal{H}}$ there exists an isometric dilation $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ on a Hilbert space $\mathcal{K} \supset \mathcal{H}$ such that $\sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* = I_{\mathcal{K}}$.*

Proof. Taking into account Theorems 2.1 and 1.3, we show, without loss of generality, that the Λ -orthogonal shift $\mathcal{S} = \{S_\lambda\}_{\lambda \in \Lambda}$ on $\mathcal{H}_0 = l^2(\mathcal{F}, \mathcal{E})$ (\mathcal{E} is

a Hilbert space) can be extended to a sequence $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ of isometries on a Hilbert space $\mathcal{H}_0 \supset \mathcal{H}_0$ such that

$$(2.5) \quad \sum_{\lambda \in \Lambda} V_\lambda V_\lambda^* = I_{\mathcal{H}_0} \quad \text{and} \quad V_\lambda|_{\mathcal{H}_0} = S_\lambda \quad (\lambda \in \Lambda).$$

Consider the Hilbert space

$$\mathcal{H} = [l^2(\mathcal{F}, \mathcal{E}) \ominus \mathcal{E}] \oplus l^2(\mathcal{F}, \mathcal{E}).$$

We embed $l^2(\mathcal{F}, \mathcal{E})$ into \mathcal{H} by identifying the element $\{e_f\}_{f \in \mathcal{F}} \in l^2(\mathcal{F}, \mathcal{E})$ with the element $0 \oplus \{e_f\}_{f \in \mathcal{F}} \in \mathcal{H}$.

Let us define the isometries V_λ ($\lambda \in \Lambda$) on \mathcal{H} . For $\lambda \geq 2$ we set $V_\lambda = S_\lambda|_{l^2(\mathcal{F}, \mathcal{E}) \ominus \mathcal{E}} \oplus S_\lambda$.

Consider the countable set

$$\mathcal{F}' = \{f \in \mathcal{F} \setminus F(1, \Lambda) : f(1) = 1\} \cup F(1, \Lambda) \cup \{0\}$$

and a one-to-one map $\gamma : \mathcal{F} \setminus \{0\} \rightarrow \mathcal{F}'$.

For $\{e_f^*\}_{f \in \mathcal{F} \setminus \{0\}} \oplus \{e_f\}_{f \in \mathcal{F}} \in \mathcal{H}$ the isometry V_1 is defined as follows

$$\begin{aligned} V_1(0 \oplus \{e_f\}_{f \in \mathcal{F}}) &= 0 \oplus S_1(\{e_f\}_{f \in \mathcal{F}}), \\ V_1(\{e_f^*\}_{f \in \mathcal{F} \setminus \{0\}} \oplus 0) &= \{e_g'^*\}_{g \in \mathcal{F} \setminus \{0\}} \oplus \{e_g'\}_{g \in \mathcal{F}}, \end{aligned}$$

where

$$e_g'^* = \begin{cases} e_f^* & \text{if } g = \gamma(f), \\ 0 & \text{otherwise} \end{cases}$$

and

$$e'_0 = e_f^* \quad \text{if } \gamma(f) = 0, \quad e'_g = 0 \quad \text{if } g \in \mathcal{F} \setminus \{0\}.$$

Now it is easy to see that the relations (2.5) hold.

Following the classification of contractions from [8] we give, in what follows, a classification of the sequences of contractions.

Let $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ on a Hilbert space \mathcal{H} such that $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \leq I_{\mathcal{H}}$.

Consider the following subspace of \mathcal{H} :

$$(2.6) \quad \mathcal{H}_0 = \left\{ h \in \mathcal{H} : \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2 = 0 \right\},$$

$$(2.7) \quad \mathcal{H}_1 = \left\{ h \in \mathcal{H} : \sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2 = \|h\|^2 \text{ for any } n \in \mathbf{N} \right\}.$$

Remark 2.7. The subspaces \mathcal{H}_0 and \mathcal{H}_1 are orthogonal and invariant for each operator T_λ^* ($\lambda \in \Lambda$).

Proof. Taking into account Theorem 2.1, 1.3 and Remark 1.4 the proof is immediately.

Thus, the Hilbert space \mathcal{H} decomposes into an orthogonal sum $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2$.

For each $k \in \{0, 1, 2\}$ we shall denote by $C^{(k)}$ (respectively $C_{(k)}$) the set of all sequences $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ on \mathcal{H} for which we have $\mathcal{H}_k = \{0\}$ (respectively $\mathcal{H} = \mathcal{H}_k$).

Let us mention that \mathcal{H}_1 is the largest subspace in \mathcal{H} on which the matrix

$$\begin{bmatrix} T_1^* \\ T_2^* \\ \vdots \end{bmatrix}$$

acts isometrically.

Consequently, a sequence $\mathcal{T} \in C^{(1)}$ will be also called completely noncoisometric (c.n.c).

In the particular case when $\mathcal{T} = \{T\}$ ($\|T\| \leq 1$) we have that $\mathcal{T} \in C^{(1)}$ if and only if T^* is completely nonisometric, that is, if there is no nonzero invariant subspace for T^* on which T^* is an isometry.

We continue this section with the study of the geometric structure of the space of the minimal isometric dilation.

For this, let $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ be a sequence of operators on a Hilbert space \mathcal{H} such that $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \leq I_{\mathcal{H}}$ and $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ be the minimal isometric dilation of \mathcal{T} on the Hilbert space $\mathcal{K} = \mathcal{H} \oplus l^2(\mathcal{F}, \mathcal{D})$ (see Theorem 2.1).

Considering the subspaces of \mathcal{K}

$$\mathcal{L} = \bigvee_{\lambda \in \Lambda} (V_\lambda - T_\lambda)\mathcal{H} \quad \text{and} \quad \mathcal{L}_* = \overline{\left(I_{\mathcal{K}} - \sum_{\lambda \in \Lambda} V_\lambda T_\lambda^* \right) \mathcal{H}}$$

we can generalize some of the results from [8, Chapter II, §§1,2] concerning the geometric structure of the space of the minimal isometric dilation.

Theorem 2.8. (i) *The subspaces \mathcal{L} and \mathcal{L}_* are wandering subspaces for \mathcal{V} and*

$$\dim \mathcal{L} = \dim \mathcal{D}; \quad \dim \mathcal{L}_* = \dim \mathcal{D}_*.$$

(ii) *The space \mathcal{K} can be decomposed as follows:*

$$\mathcal{K} = \mathcal{R} \oplus M_{\mathcal{F}}(\mathcal{L}_*) = \mathcal{H} \oplus M_{\mathcal{F}}(\mathcal{L}),$$

and the subspace \mathcal{R} reduces each operator V_λ ($\lambda \in \Lambda$).

(iii) $\mathcal{L} \cap \mathcal{L}_* = 0$.

(iv) *The subspace \mathcal{R} reduces to $\{0\}$ if and only if $\mathcal{T} \in C_{(0)}$.*

Proof. The Wold decomposition (see Theorem 1.3) for the minimal isometric dilation \mathcal{V} on the space $\mathcal{K} = \mathcal{H} \oplus l^2(\mathcal{F}, \mathcal{D})$ gives $\mathcal{K} = \mathcal{R} \oplus M_{\mathcal{F}}(\mathcal{L}'_*)$, where $\mathcal{R} = \bigcap_{n=0}^\infty [\bigoplus_{f \in F(n, \Lambda)} V_f \mathcal{H}]$ reduces each operator V_λ ($\lambda \in \Lambda$) and $\mathcal{L}'_* = \mathcal{H} \ominus (\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H})$ is a wandering subspace for \mathcal{V} .

It is easy to see that $\mathcal{L}'_* = \mathcal{L}_*$ and that the operator $\Phi_* : \mathcal{L}_* \rightarrow \mathcal{D}_*$ defined by

$$\Phi_* \left(I_{\mathcal{K}} - \sum_{\lambda \in \Lambda} V_\lambda T_\lambda^* \right) h = D_* h \quad (h \in \mathcal{H})$$

is unitary. Hence it follows that $\dim \mathcal{L}_* = \dim \mathcal{D}_*$. Equation (2.1) yields

$$\sum_{\lambda \in \Lambda} (V_\lambda - T_\lambda)h_\lambda = 0 \oplus D((h_\lambda)_{\lambda \in \Lambda}) \quad \text{for } (h_\lambda)_{\lambda \in \Lambda} \in \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda$$

(\mathcal{H}_λ is a copy of \mathcal{H}).

By this relation we deduce that there exists a unitary operator $\Phi: \mathcal{L} \rightarrow \mathcal{D}$ defined by equation

$$\Phi \left(\sum_{\lambda \in \Lambda} (V_\lambda - T_\lambda)h_\lambda \right) = D((h_\lambda)_{\lambda \in \Lambda})$$

and hence that $\dim \mathcal{L} = \dim \mathcal{D}$.

The fact that \mathcal{L} is a wandering subspace for \mathcal{V} and that $\mathcal{H} \perp M_{\mathcal{F}}(\mathcal{L})$ follows from the form of the isometries V_λ ($\lambda \in \Lambda$) defined by (2.1).

Taking into account the minimality of \mathcal{H} it follows that $\mathcal{H} = \mathcal{H} \oplus M_{\mathcal{F}}(\mathcal{L})$.

Let us now show that $\mathcal{L} \cap \mathcal{L}_* = 0$. First we need to prove that

$$(2.8) \quad \mathcal{L}_* \oplus \left(\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H} \right) = \mathcal{H} \oplus \mathcal{L}.$$

This follows from the fact that, for an element $u \in \mathcal{H}$, the possibility of a representation of the form

$$u = \left(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda T_\lambda^* \right) h_0 + \sum_{\lambda \in \Lambda} V_\lambda h_\lambda, \quad h_0 \in \mathcal{H}, (h_\lambda)_{\lambda \in \Lambda} \in \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda,$$

is equivalent to the possibility of a representation of the form

$$u = h^{(0)} + \sum_{\lambda \in \Lambda} (V_\lambda - T_\lambda)h^{(\lambda)}, \quad h^{(0)} \in \mathcal{H}, (h^{(\lambda)})_{\lambda \in \Lambda} \in \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda.$$

Indeed, we have only to set

$$h_0 = h^{(0)} - \sum_{\lambda \in \Lambda} T_\lambda h^{(\lambda)}, \quad h_\lambda = T_\lambda^* h^{(0)} + h^{(\lambda)}$$

and, conversely,

$$h^{(0)} = \sum_{\lambda \in \Lambda} T_\lambda h_\lambda + \left(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \right) h_0, \quad h^{(\lambda)} = h_\lambda - T_\lambda^* h_0.$$

Thus (2.8) holds. On the other hand, since

$$\mathcal{L} \subset \left(\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H} \right) \vee \mathcal{H} \quad \text{and} \quad \bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H} \subset \mathcal{L} \oplus \mathcal{H}$$

we have that $\mathcal{H} \vee \left(\bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H} \right) = \mathcal{H} \oplus \mathcal{L}$. This relation and (2.8) show that $\mathcal{L} \cap \mathcal{L}_* = \{0\}$.

The statement (iv) is contained in Proposition 2.3. The proof is complete.

Proposition 2.9. For every sequence $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ of operators on \mathcal{H} and for its minimal isometric dilation $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ on \mathcal{K} , we have

$$(2.9) \quad M_{\mathcal{G}}(\mathcal{L}) \vee M_{\mathcal{G}}(\mathcal{L}_*) = \mathcal{H} \ominus \mathcal{H}_1$$

where \mathcal{H}_1 is given by (2.7).

In particular, if \mathcal{T} is c.n.c., then

$$(2.10) \quad M_{\mathcal{G}}(\mathcal{L}) \vee M_{\mathcal{G}}(\mathcal{L}_*) = \mathcal{H}.$$

Proof. Taking into account Theorem 2.8 and that $\mathcal{H}_1 \subset \mathcal{H}$ it follows that $\mathcal{H}_1 \perp M_{\mathcal{G}}(\mathcal{L}) \vee M_{\mathcal{G}}(\mathcal{L}_*)$.

Now let $k \in \mathcal{H}$ be such that $k \perp M_{\mathcal{G}}(\mathcal{L})$ and $k \perp M_{\mathcal{G}}(\mathcal{L}_*)$.

From the same theorem it follows that $k \in \mathcal{H}$ and $k \perp V_f \mathcal{L}_*$ for every $f \in \mathcal{F}$. Hence we have

$$0 = \left(k, V_f \left(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda T_\lambda^* \right) h \right) = (T_f^* k, h) - \sum_{\lambda \in \Lambda} (T_\lambda^* T_f^* k, T_\lambda^* h)$$

for every $h \in \mathcal{H}$.

Choosing $h = T_f^* k$ ($f \in \mathcal{F}$) we obtain

$$\|T_f k\|^2 = \sum_{\lambda \in \Lambda} \|T_\lambda^* T_f^* k\|^2$$

for any $f \in \mathcal{F}$.

Hence we deduce

$$\sum_{g \in F(n, \Lambda)} \|T_g^* k\|^2 = \|k\|^2$$

for any $n \in \mathbb{N}$. We conclude that $k \in \mathcal{H}_1$. Conversely, for every $k \in \mathcal{H}_1$ it is easy to see that $k \perp M_{\mathcal{G}}(\mathcal{L}) \vee M_{\mathcal{G}}(\mathcal{L}_*)$. The relation (2.10) follows because for \mathcal{T} c.n.c. we have $\mathcal{H}_1 = \{0\}$.

The last aim of this section is to generalize some of the results from [8, Chapter II, §3]. Throughout $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ is the minimal isometric dilation of $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$. The space of the minimal isometric dilation is

$$(2.11) \quad \mathcal{K} = \mathcal{R} \oplus M_{\mathcal{G}}(\mathcal{L}_*) = \mathcal{H} \oplus l^2(\mathcal{F}, \mathcal{D}).$$

Proposition 2.10. For every $h \in \mathcal{H}$ we have

$$(2.12) \quad P_{\mathcal{H}} h = \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} V_f T_f^* h$$

and consequently

$$(2.13) \quad \|P_{\mathcal{H}} h\|^2 = \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2$$

where $P_{\mathcal{H}}$ denotes the orthogonal projection of \mathcal{K} into \mathcal{H} .

Proof. An easy computation shows that

$$\begin{aligned} & \left\| \sum_{f \in F(n+1, \Lambda)} V_f T_f^* h - \sum_{f \in F(n, \Lambda)} V_f T_f^* h \right\|^2 \\ &= \sum_{f \in F(n+1, \Lambda)} \|T_f^* h\|^2 - \sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2 \leq 0 \end{aligned}$$

for every $n \in \mathbf{N}$. This implies the convergence of $\{\sum_{f \in F(n, \Lambda)} V_f T_f^* h\}_{n=1}^\infty$ the sequence in \mathcal{H} . Setting

$$k = \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} V_f T_f^* h,$$

let us show that $k = P_{\mathcal{R}} h$, i.e. $k \perp M_{\mathcal{F}}(\mathcal{L}_*)$ and $h - k \in M_{\mathcal{F}}(\mathcal{L}_*)$.

Since for every $g \in \mathcal{F}$ there exists $n_0 \in \mathbf{N}$ such that

$$\sum_{f \in F(n, \Lambda)} V_f T_f^* h \perp V_g \mathcal{L}_*$$

for any $n \geq n_0$, it follows that $k \perp M_{\mathcal{F}}(\mathcal{L}_*)$.

On the other hand we have

$$\begin{aligned} h - \sum_{f \in F(n, \Lambda)} V_f T_f^* h &= \left(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda T_\lambda^* \right) h + \sum_{f \in F(1, \Lambda)} V_f \left(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda T_\lambda^* \right) T_f^* h \\ &+ \dots + \sum_{f \in F(n-1, \Lambda)} V_g \left(I_{\mathcal{H}} - \sum_{\lambda \in \Lambda} V_\lambda T_\lambda^* \right) T_g^* h \in M_{\mathcal{F}}(\mathcal{L}_*) \end{aligned}$$

and therefore

$$h - k = \lim_{n \rightarrow \infty} \left(h - \sum_{f \in F(n, \Lambda)} V_f T_f^* h \right) \in M_{\mathcal{F}}(\mathcal{L}_*).$$

This ends the proof.

Proposition 2.11. Let $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ be a sequence of operators on \mathcal{H} such that the matrix $[T_1, T_2, \dots]$ is an injection. Then $\overline{P_{\mathcal{R}} \mathcal{H}} = \mathcal{R}$.

Proof. Let us suppose that there exists $k \in \mathcal{R}$, $k \neq 0$ such that $k \perp P_{\mathcal{R}} \mathcal{H}$, or equivalently, such that $k \perp M_{\mathcal{F}}(\mathcal{L}_*)$ and $k \perp \mathcal{H}$.

By Theorem 2.8 we have $\mathcal{H} = \mathcal{H} \oplus M_{\mathcal{F}}(\mathcal{L})$. It follows that $k \in M_{\mathcal{F}}(\mathcal{L})$ and hence $k = \sum_{f \in \mathcal{F}} V_f l_f$ where $l_f \in \mathcal{L}$ ($f \in \mathcal{F}$) and $\sum_{f \in \mathcal{F}} \|l_f\|^2 < \infty$. Since $k \neq 0$ there exists $f_0 \in \mathcal{F}$, such that $V_{f_0} l_{f_0} \neq 0$ and

$$V_{f_0}^* k = l_{f_0} + \sum_{\substack{g \in \mathcal{F} \\ g \neq f_0}} V_g l'_g \quad (l'_g \in \mathcal{L}).$$

One can easily show that for every $g \in \mathcal{F}$, $g \neq 0$, $V_g \mathcal{L} \perp \mathcal{L}_*$. Since $V_{f_0}^* k \perp \mathcal{L}_*$ it follows that $l_{f_0} \perp \mathcal{L}_*$. By the relation (2.8) we deduce that $l_{f_0} \in \bigoplus_{\lambda \in \Lambda} V_\lambda \mathcal{H}$.

Therefore, there exists a nonzero $\bigoplus_{\lambda \in \Lambda} h_\lambda \in \bigoplus_{\lambda \in \Lambda} \mathcal{H}_\lambda$ such that $l_{f_0} = \sum_{\lambda \in \Lambda} V_\lambda h_\lambda$. Since $\mathcal{L} \perp \mathcal{H}$, it follows that $\sum_{\lambda \in \Lambda} T_\lambda h_\lambda = 0$ which is a contradiction with the hypothesis.

Thus $\overline{P_{\mathcal{R}} \mathcal{H}} = \mathcal{R}$ and the proof is complete.

For each $\lambda \in \Lambda$ let us denote by R_λ the operator $V_\lambda|_{\mathcal{R}}$. Taking into account the Wold decomposition (Theorem 1.3) we have $\sum_{\lambda \in \Lambda} R_\lambda R_\lambda^* = I_{\mathcal{R}}$.

The following theorem is a generalization of Proposition 3.5 in [8, Chapter II].

Proposition 2.12. *Let $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ a sequence of operators on \mathcal{H} such that $\mathcal{T} \in C^{(0)}$ and the matrix $[T_1, T_2, \dots]$ is an injective contraction.*

Then \mathcal{T} is quasi-similar to $\{R_\lambda\}_{\lambda \in \Lambda}$, i.e., there exists a quasi-affinity Y from \mathcal{R} to \mathcal{H} such that $T_\lambda Y = Y R_\lambda$ for every $\lambda \in \Lambda$.

Proof. According to Proposition 2.10 we have

$$\begin{aligned} V_\lambda^* P_{\mathcal{R}} h &= \lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} V_\lambda^* V_f T_f^* h \\ &= \lim_{n \rightarrow \infty} \sum_{g \in F(n-1, \Lambda)} V_g T_g^* T_\lambda^* h = P_{\mathcal{R}} T_\lambda^* h \end{aligned}$$

for all $h \in \mathcal{H}$ and each $\lambda \in \Lambda$.

Setting $X = P_{\mathcal{R}}|_{\mathcal{H}}$ it follows that $R_\lambda^* X = X T_\lambda^*$ for every $\lambda \in \Lambda$. Let us show that X is a quasi-affinity.

Since $\mathcal{T} \in C^{(0)}$ we have that

$$\lim_{n \rightarrow \infty} \sum_{f \in F(n, \Lambda)} \|T_f^* h\|^2 = 0 \quad \text{for every nonzero } h \in \mathcal{H}.$$

By Proposition 2.10 we deduce that $P_{\mathcal{R}} h \neq 0$ for every nonzero $h \in \mathcal{H}$, i.e., X is an injection.

On the other hand, Proposition 2.11 shows that $\overline{X\mathcal{H}} = \mathcal{R}$.

If we take $Y = X^*$, this finishes the proof.

3

In this section we extend the Sz.-Nagy-Foiaş lifting theorem [7, 8, 1, 4] to our setting.

Let $\mathcal{T} = \{T_\lambda\}_{\lambda \in \Lambda}$ be a sequence of operators on \mathcal{H} with $\sum_{\lambda \in \Lambda} T_\lambda T_\lambda^* \leq I_{\mathcal{H}}$ and $\mathcal{V} = \{V_\lambda\}_{\lambda \in \Lambda}$ be the minimal isometric dilation of the Hilbert space $\mathcal{H} = \mathcal{H} \oplus l^2(\mathcal{T}, \mathcal{D})$ (see Theorem 2.1).

Consider the following subspaces of \mathcal{H}

$$\mathcal{H}_1 = \mathcal{H} \vee \left(\bigvee_{f \in F(1, \Lambda)} V_f \mathcal{H} \right)$$

and

$$\mathcal{H}_n = \mathcal{H}_{n-1} \vee \left(\bigvee_{f \in F(1, \Lambda)} V_f \mathcal{H}_{n-1} \right) \quad \text{for } n \geq 2.$$

Note that $\mathcal{H}_n \subset \mathcal{H}_{n+1}$ and that all the space \mathcal{H}_n ($n \geq 1$) are invariant for each operator V_λ^* ($\lambda \in \Lambda$).

As in [7, 8, 1, 4] the n -stepped dilation of \mathcal{T} is the sequence $\mathcal{T}_n = \{(T_\lambda)_n\}_{\lambda \in \Lambda}$ of operators defined by $(T_\lambda)_n^* = V_\lambda^*|_{\mathcal{H}_n}$ ($n \geq 1, \lambda \in \Lambda$).

One can easily show that \mathcal{V} is the minimal isometric dilation on \mathcal{T}_n and that \mathcal{T}_{n+1} is the one-step dilation of \mathcal{T}_n .

Let us observe that $\mathcal{H}_1 = \mathcal{H} \oplus \mathcal{D}$ and

$$\mathcal{H}_n = \mathcal{H} \oplus \mathcal{D} \oplus \left(\bigoplus_{f \in F(1, \Lambda)} S_f \mathcal{D} \right) \oplus \cdots \oplus \left(\bigoplus_{f \in F(n-1, \Lambda)} S_f \mathcal{D} \right) \quad (n \geq 2)$$

where $\mathcal{S} = \{S_\lambda\}_{\lambda \in \Lambda}$ is the Λ -orthogonal shift acting on $l^2(\mathcal{F}, \mathcal{D})$.

Now Lemma 2 and Theorem 3 in [4] can be easily extended to our setting. Thus, we omit the proofs in what follows.

Lemma 3.1. *Let P_n be the orthogonal projection from \mathcal{H} into \mathcal{H}_n .*

Then $\bigvee_{n \geq 1} \mathcal{H}_n = \mathcal{H}$ and for each $\lambda \in \Lambda$ we have

$$(T_\lambda)_n^* P_n \rightarrow V_\lambda^* \quad (\text{strongly}) \text{ as } n \rightarrow \infty.$$

Let $\mathcal{T}' = \{T'_\lambda\}_{\lambda \in \Lambda}$ be another sequence of operators on a Hilbert space \mathcal{H}' with $\sum_{\lambda \in \Lambda} T'_\lambda T'^*_\lambda \leq I_{\mathcal{H}'}$, and $\mathcal{V}' = \{V'_\lambda\}_{\lambda \in \Lambda}$ be the minimal isometric dilation of \mathcal{T}' acting on the Hilbert space $\mathcal{H}' = \mathcal{H}' \oplus l^2(\mathcal{F}, \mathcal{D}')$.

Theorem 3.2. *Let $A: \mathcal{H} \rightarrow \mathcal{H}'$ be a contraction such that for each $\lambda \in \Lambda$ $T'_\lambda A = AT_\lambda$. Then there exists a contraction $B: \mathcal{H} \rightarrow \mathcal{H}'$ such that for each $\lambda \in \Lambda$ $V'_\lambda B = BV_\lambda$ and $B^*|_{\mathcal{H}'} = A^*$.*

REFERENCES

1. R. G. Douglas, P. S. Muhly and C. M. Pearcy, *Lifting commuting operators*, Michigan Math. J. **15** (1968), 385–395.
2. C. Foiaş, *A remark on the universal model for contractions of G. C. Rota*, Comm. Acad. R. P. Romane **13** (1963), 349–352.
3. A. E. Frazho, *Models for noncommuting operators*, J. Funct. Anal. **48** (1982).
4. —, *Complements to models for noncommuting operators*, J. Funct. Anal. **59** (1984), 445–461.
5. G. Popescu, *Models for infinite sequences of noncommuting operators*, INCREST preprint, no. 23/1986.
6. J. J. Schaffer, *On unitary dilations of contractions*, Proc. Amer. Math. Soc. **6** (1955), 322.
7. B. Sz.-Nagy and C. Foiaş, *Dilation des commutants*, C. R. Acad. Sci. Paris Sér. A **266** (1968), 201–212.
8. —, *Harmonic analysis on operators on Hilbert space*, North-Holland, Amsterdam, 1970.

DEPARTMENT OF MATHEMATICS, INCREST, BD. PACII 220, 79622 BUCHAREST, ROMANIA