



Isotopic ($\delta^2\text{H}$) Analysis of Stored Lipids in Migratory and Overwintering Monarch Butterflies (*Danaus plexippus*): Evidence for Southern Critical Late-Stage Nectaring Sites?

Keith A. Hobson^{1,2*}, Oscar R. García-Rubio³, Rogelio Carrera-Treviño⁴, Libesha Anparasan¹, Kevin J. Kardynal², Jeremy N. McNeil¹, Eligio García-Serrano⁵ and Blanca X. Mora Alvarez¹

¹ Department of Biology, University of Western Ontario, London, ON, Canada, ² Environment and Climate Change Canada, Saskatoon, SK, Canada, ³ Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, Santiago de Querétaro, Mexico, ⁴ Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Nuevo León, Nuevo León, Mexico, ⁵ Fondo Para la Conservación de la Mariposa Monarca, Zitácuaro, Mexico

OPEN ACCESS

Edited by:

Jennifer M. Gleason,
University of Kansas, United States

Reviewed by:

Chip Taylor,
University of Kansas, United States
Victoria Pocius,
Pennsylvania State University (PSU),
United States

*Correspondence:

Keith A. Hobson
khobson6@uwo.ca

Specialty section:

This article was submitted to
Behavioral and Evolutionary Ecology,
a section of the journal
Frontiers in Ecology and Evolution

Received: 15 June 2020

Accepted: 14 September 2020

Published: 09 October 2020

Citation:

Hobson KA, García-Rubio OR, Carrera-Treviño R, Anparasan L, Kardynal KJ, McNeil JN, García-Serrano E and Mora Alvarez BX (2020) Isotopic ($\delta^2\text{H}$) Analysis of Stored Lipids in Migratory and Overwintering Monarch Butterflies (*Danaus plexippus*): Evidence for Southern Critical Late-Stage Nectaring Sites? *Front. Ecol. Evol.* 8:572140. doi: 10.3389/fevo.2020.572140

Monarch butterflies (*Danaus plexippus*) fuel their migration and overwinter energy needs through accumulated fat stores derived from plant nectars. Determining origins of these fuels is crucial to effective conservation programs. We used stable-hydrogen ($\delta^2\text{H}$) and carbon ($\delta^{13}\text{C}$) isotope measurements in stored lipids of monarchs raised under laboratory conditions as a proof of principle for the isotopic spatial sourcing of stored lipids. We then applied this approach to wild specimens collected from 2015 to 2018 to infer spatial information on nectaring by fall migrants through northeast Mexico and at the Mexican overwinter sites. Migrating monarchs derived from wide geographic natal origins but lipid $\delta^2\text{H}$ values from migratory cohorts were not related to natal origin. Instead, migrants exploited isotopically similar nectar sources. Distributions of lipid $\delta^2\text{H}$ values in overwintering monarchs were broader and more negative by $\sim 40\%$ suggesting more transport of lipids from higher latitudes or additional nectaring while migrating at higher elevations though northeastern to central Mexico. Our work establishes a new isotopic technique for tracking origins of stored lipids in monarchs and other migratory animals and emphasizes the importance of nectar availability in the southern portion of the range, and especially the nectar corridor through central Mexico.

Keywords: lipids, overwintering, stable isotopes, deuterium, carbon-13, nectar corridors, migratory fueling

INTRODUCTION

Migrant species have evolved complex physiological and behavioral mechanisms that facilitate long-distance movement, including the ability to efficiently convert dietary macromolecules to energy for immediate use or stored as lipid, an energy dense fuel (McWilliams et al., 2004). Understanding the nutritional ecology of migrant animals, especially those using stopover sites

to obtain resources to fuel the next leg of their journey is crucial for the development of effective strategies for the management of migrant animal populations (Moore et al., 1993; Lennox et al., 2016).

The eastern North American population of the monarch butterfly (*Danaus plexippus*) is an iconic example of insect long-distance migration where millions of adults migrate from their natal sites in the United States and Canada to the overwintering sites in the high-altitude oyamel fir (*Abies religiosa*) forests of central Mexico. This journey is fueled by nectar (Brown and Chippendale, 1974; Brower et al., 2006) obtained at sites along the migratory pathway (Beall, 1948; Gibo and McCurdy, 1993; Brower et al., 2006). Lipid stores generally increase as the butterflies approach the wintering grounds (Brower et al., 2006). Because they rarely feed on flowers during the 4–5 month overwintering period, they must sustain themselves on lipid stores acquired prior to arrival at the winter roost sites, which provide a thermal environment conducive to lipid conservation (Alonso-Mejia et al., 1997; Williams and Brower, 2015). It was proposed that resources acquired primarily in Texas provides the bulk of the lipids required for both overwintering and subsequent reproduction in spring (Brower, 1985). However, in 2011–2012, fall monarchs in Texas had lower than average lipid reserves, due to drought conditions, yet those at the overwintering sites apparently had normal levels (reviewed by Brower et al., 2015). This difference in lipid levels would suggest that, at least in some years, monarchs actively nectar along the floral corridor in Mexico leading to and perhaps including their overwintering sites. To date, the real importance of nectaring sites along the migratory journey and especially in Mexico remains anecdotal despite their likely importance (Saunders et al., 2019).

Understanding where monarchs obtain resources to fuel both migratory flight and overwintering stores would provide new insights into the migratory strategy of this species and subsequently to overall conservation efforts (Brower et al., 2006). More and more attention is being directed to conditions experienced by migrating monarchs during the fall migration with respect to overall population demographics and conservation (Saunders et al., 2019; Taylor et al., 2019). If there is an intrinsic marker where lipids were synthesized, it could be used, retrospectively, to infer locations of key fueling sites. One approach to investigate this possibility is the use of naturally occurring stable isotopes of several elements (Hobson and Wassenaar, 2019) as stable isotope values in foodwebs can vary predictably across large spatial gradients due to a variety of biogeochemical processes and that these spatial patterns are reflected in consumers. The first studies to use stable-hydrogen isotope measurements ($\delta^2\text{H}$) to infer natal origin of any migrant species were carried out using wing chitin, a tissue that is metabolically inert following synthesis, performed on monarchs at their roost sites in central Mexico (Wassenaar and Hobson, 1998; Hobson et al., 1999). Since then, continental patterns, or isoscapes, of both precipitation $\delta^2\text{H}$ and milkweed $\delta^{13}\text{C}$ have been used to study patterns of monarch movements in North America (Miller et al., 2012; Flockhart et al., 2013, 2017; Altizer et al., 2015).

We speculate that despite the numerous metabolic pathways involved in lipid synthesis, the $\delta^{13}\text{C}$ and $\delta^2\text{H}$ values of stored neutral lipids could provide valuable provenance information of monarchs as lipids are composed primarily of carbon and hydrogen derived from plant nectars. This composition is particularly true of palmitic and oleic acids which are the most abundant stored lipids in monarchs (Cenedella, 1971; Brower et al., 2006). While a North American plant leaf water $\delta^2\text{H}$ isoscape has been developed that shows a close concordance with patterns of amount-weighted precipitation $\delta^2\text{H}$ (West et al., 2008, 2010), to our knowledge, no plant $\delta^2\text{H}$ carbohydrate isoscapes exist. Foodwebs driven by plant primary production show well-established continental gradients in $\delta^2\text{H}$ and so we expect plant nectars and the lipids synthesized from them by monarchs to follow the continent wide $\delta^2\text{H}$ precipitation isoscape as demonstrated for chitins and keratins of numerous migratory taxa (Hobson and Wassenaar, 2019).

Our approach was to first demonstrate that stable isotope measurements could be used to trace nutritional origins of stored lipids and then to apply this proof of concept to wild migrating and overwintering monarchs in Mexico. We measured $\delta^2\text{H}$ and $\delta^{13}\text{C}$ in (i) lipids from individual monarchs raised in the laboratory on nectar sources comprised of isotopically known carbohydrate and water values, and (ii) wing and lipid $\delta^2\text{H}$ of monarchs collected at different sites along the migratory pathway in northeastern to central Mexico and at the Mexican overwinter sites. We postulate that if the isotopic distribution of lipid $\delta^2\text{H}$ values were independent of natal origin (as derived from wing $\delta^2\text{H}$) but were tightly clustered, this would suggest adults fed at geographically restricted sites that could potentially be identified. We discuss the relevance of our findings to the ultimate goal of tracing geographic origins of lipids in migratory monarchs and other insects and how our approach can have significant implications for conservation.

MATERIALS AND METHODS

Laboratory Study

Eggs were obtained from adult monarchs collected in July 2019 on the Environmental Science Western Field Station (WFS) (43.07°N, 81.34°W) and reared at 25°C, 65% RH and 16L:8D. The larvae were reared in individual containers and provided pieces of washed fresh, field collected milkweed (*Asclepias syriaca*) leaves each day. The newly emerged adults were sexed, caged separately to deter mating, and ten individuals (five male and five female) were assigned to one of four isotopically distinct dietary treatments, an approach allowing us to directly trace isotopically contributions from dietary macromolecules to ultimate lipid stores. The isotopic values of the treatment nectars were: (1) high $\delta^{13}\text{C}$ (C4 cane sugar: mean \pm SD, $-12.4 \pm 0.03\%$, $n = 5$) with high $\delta^2\text{H}$ (spiked) water ($182 \pm 6.1\%$, $n = 3$), (2) high $\delta^{13}\text{C}$ with low $\delta^2\text{H}$ (unspiked tap) water ($-55 \pm 2.0\%$, $n = 3$), (3) low $\delta^{13}\text{C}$ (C3 beet sugar: $-26.4 \pm 0.1\%$, $n = 5$) with low $\delta^2\text{H}$ water and (4) low $\delta^{13}\text{C}$ with high $\delta^2\text{H}$ water ($n = 5$). Deuterium spiked water was prepared by adding a quantity of 99.9 atom % heavy water (deuterium oxide) (Sigma Aldrich). The

nectar, at a 15% concentration (with cane or beet sugar as the nutrient sugar source), was prepared following the methodology developed by Orley Taylor (personal communication) that avoids fermentation. Adults were fed daily *ad libitum* for 5 days, after which they were sacrificed, the fat body extracted using 2:1 (v:v) mixture of chloroform:methanol. The lipids were recovered following evaporation in a fume hood for > 24 h (23°C, 101.3 kPa), then frozen (−20°C) until stable isotope analyses were conducted.

Field Study

Monarchs were sampled between 2015 and 2018 (Mexican permits SGPA/DGVS/002159/18, SGPA/DGVS/002299/18, SGPA/DGVS/09159/17, SGPA/DGVS/10763/16, SGPA/DGVS/06299/15), at several sites in northeastern to central Mexico along the fall migration route and at overwintering colonies (Figure 1 and Supplementary Table S1). In 2015–2016 overwintering monarchs were collected at Sierra Chincua, El Rosario, and Piedra Herrada colonies. Evidence for isotopic change or discrimination through the period of lipid use when fasting at colonies would complicate interpretation of lipid

isotope values based on timing. So, to test this, in 2017–2018 monarch lipids were sampled monthly from December to February at the Cerro Pelon colony. All monarch samples were placed in individual paper envelopes, labeled and then frozen (−20°C) and shipped to Canada for stable isotope analysis.

Stable Isotope Analyses

Samples of 0.35 ± 0.02 mg of wing were weighed into pressed silver 3.5×5 mm capsules, and analyzed using a Eurovector Uniprep autosampler (Milan, Italy) carousel attached to a Eurovector 3000 Elemental Analyzer, coupled to a Thermo Delta V Plus isotope ratio mass spectrometer (Bremen, Germany) in continuous flow mode with helium carrier gas. After the samples were loaded, the Uniprep autosampler (heated to 60°C) was vacuum evacuated and subsequently flushed with dry helium twice to remove adsorbed atmospheric moisture from the crushed silver capsules. Two USGS keratin standards, EC-01 (formerly CBS: Caribou Hoof Standard) and EC-02 (KHS: Kudu Horn Standard of Environment Canada) were included every ten samples. Samples were combusted at 1350°C using glassy carbon. Values of $\delta^2\text{H}$ of non-exchangeable hydrogen




FIGURE 1 | General study area in Northeastern Mexico. Note that the range shown in northeast Mexico includes potential non-migrants.

were derived using the comparative equilibration approach of Wassenaar and Hobson (2003) and calibrated to Vienna Standard Mean Ocean Water (VSMOW) using EC-01 ($\pm 1.9\%$ 1 SD, $n = 18$, accepted $\delta^2\text{H} = -197.0\%$) and EC-02 ($\pm 1.6\%$, $n = 17$, accepted $\delta^2\text{H} = -54.1\%$). We used these former calibration standard values because our original assignment algorithms were based on these values (Hobson et al., 1999). This approach results in identical assignments if new calibration standards are used with correspondingly modified assignment algorithms (see below; Soto et al., 2017). Overall (within-run) measurement error for EC-01 and EC-02 $\delta^2\text{H}$ was $\sim 2\%$.

Lipid samples were analyzed in an identical fashion to wing samples but were calibrated against the USGS water standards sealed in silver tubes VSMOW2 (0‰) and USGS 46 (-235.8%) because no international standards for animal lipids are available and we were interested only in relative lipid $\delta^2\text{H}$ values (note that lipids have no exchangeable H). We compared calibration results with several other laboratory standards including the keratins CBS (-157%) and KHS (-35.3%) and found consistent offsets. Overall measurement error for these standards relative to the VSMOW scale and based on within-run replicate measurements was $\pm 3\%$. For the laboratory experiment, water $\delta^2\text{H}$ analyses were performed using an Off Axis Integrated Cavity Output Spectroscopy (OA-ICOS) using a Los Gatos Research DLT-100 laser spectrometer (Mountain view California). We used two calibrated reference waters (INV1 $\delta^2\text{H} = -217.7\%$ and ROD3 $\delta^2\text{H} = -3.9\%$, respectively) to normalize raw delta values to the VSMOW scale. Precisions as determined by replicate analyses of samples and reference waters were ± 1 and 0.1% respectively.

Concurrent $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses were performed using a Costech Elemental Analyzer coupled to a Thermo Delta Plus XL isotope ratio mass spectrometer (Thermo Instruments, Bremen, Germany) operated in continuous flow mode with helium carrier gas, but only $\delta^{13}\text{C}$ values are reported here. Two standards, USGS-40 and USGS-41, were included for every ten samples, and two internal laboratory standards, powdered keratin (MP Biomedicals Inc., Cat No. 90211, Lot No.9966H) and IAEA-CH-6 were included to monitor instrument drift and provide a check on accuracy over the course of each analytical session. Values of $\delta^{13}\text{C}$ were calibrated to VPDB using USGS-40 ($\pm 0.1\%$ 1 SD, $n = 8$, accepted $\delta^{13}\text{C} = -26.4\%$) and USGS-41 ($\pm 0.1\%$, $n = 8$, accepted $\delta^{13}\text{C} = +37.6\%$). Measurement error was $\pm 0.1\%$ for $\delta^{13}\text{C}$.

Assignment to Origins

We depicted the origins of wild monarchs caught in Mexico with a likelihood-based assignment method (Hobson et al., 2009b; Wunder, 2010), using the wing chitin $\delta^2\text{H}$ isoscape ($\delta^2\text{H}_w$) and an amount-weighted precipitation-to-wing calibration algorithm derived by Hobson et al. (1999). This was used to convert amount-weighted mean annual precipitation $\delta^2\text{H}$ ($\delta^2\text{H}_p$) isoscapes (Terzer et al., 2013; IAEA/WMO, 2015) into $\delta^2\text{H}_w$ isoscapes. We used the 9.3% residual SD error from this regression in our assignments. We then created a spatial layer representing the known geographic range for the eastern breeding population of the monarch butterfly and used

this as a mask (i.e., clip) to limit our assignment to origin analyses (Figure 2).

We estimated the likelihood that a cell (pixel) within the $\delta^2\text{H}_w$ isoscape represented a potential origin for a sample by using a normal probability density function to estimate the likelihood function based on the observed $\delta^2\text{H}_w$, and thus depicted the likely origins of each monarch by assigning individuals to the $\delta^2\text{H}_w$ isoscape one at a time. We arbitrarily chose a 2:1 odds ratio (see Van Wilgenburg et al., 2012) to include only those pixels (coded 1) with at least a 67% probability of origin vs. all others (coded 0). This resulted in a map of binary values for each assigned individual representing presence (1) or absence (0) for each cell in the isoscape. We then summed the results of individual assignments by stacking the surfaces. This approach thus provides assignment maps that show the number of individuals in a sample assigned to any given pixel.

Statistical Analyses

When data were normally distributed (Shapiro Wilk's test), we compared isotopic data among groups using Analysis of Variance (ANOVA) and Tukey's *post-hoc* comparisons with statistical significance set at $p = 0.05$. In cases where groups were not significantly different, they were pooled for further analysis. Simple regression was used to establish isotopic response variables in monarch tissues with isotope values of dietary treatments. The above analyses were completed in SPSS Version 24 (IBM Corp, 2016).


Geographic assignments to origin were made using functions within the R statistical computing environment (R Core Team, 2019) using scripts employing the "raster" v. 3.0-12 (Hijmans, 2020) and "maptools" v. 0.9-9 (Bivand and Lewin-Koh, 2019) packages. Thus, the final assignment surface depicted the number of individuals co-assigned at each pixel based on the odds criteria.

RESULTS

Laboratory Experiments

No effect of sex was found for either $\delta^{13}\text{C}$ [$F_{(1, 37)} = 0.10$, $p = 0.75$] or $\delta^2\text{H}$ [$F_{(1, 37)} = 0.035$, $p = 0.966$] so the individual isotope data sets were pooled. The $\delta^{13}\text{C}$ values of lipids were normally distributed (Shapiro Wilk, $p > 0.05$) and were significantly affected by treatment [$F_{(3, 37)} = 20.46$, $p < 0.001$], with Tukey *post-hoc* analyses revealing differences based on the expected contrasts of C3 and C4 sugars (Supplementary Table S2). By regressing lipid $\delta^{13}\text{C}$ values against sugar $\delta^{13}\text{C}$ values for the unspiked water treatment (since tap water corresponded best to environmental waters experienced by monarchs), we obtained the relationship $\delta^{13}\text{C}_{\text{lipid}} = 0.59 * \delta^{13}\text{C}_{\text{sugar}} - 15.08$, indicating that nectar contributed $\sim 59\%$ of total carbon in the lipid.

Similarly, $\delta^2\text{H}$ values of lipids were normally distributed (Shapiro Wilk, $p > 0.05$) and there was a significant effect of treatment [$F_{(3, 37)} = 8.96$, $p = 0.00017$], with Tukey *post-hoc* analyses revealing differences based on the expected contrasts of spiked and unspiked nectar waters (Supplementary Table S2). The contribution of carbohydrate-based H to stored lipids was



evaluated by examining the response in lipid $\delta^2\text{H}$ with a change from beet to cane sugar under a constant unspiked water nectar source, which provided the relationship $\delta^2\text{H}_{\text{lipid}} = 0.27 * \delta^2\text{H}_{\text{sugar}} - 279.6$ indicating that sugar contributed $\sim 27\%$ of the hydrogen in adult monarch lipids. A similar analysis examining the response in lipid $\delta^2\text{H}$ with a change from tap water to spiked water under a constant C3 nectar sugar resulted in the relationship $\delta^2\text{H}_{\text{lipid}} = 0.19 * \delta^2\text{H}_{\text{water}} - 272.6$ revealing that nectar water contributed about 19% of hydrogen in the fat body lipids. As determined in controlled studies involving metabolically inactive chitins (Hobson et al., 1999), our captive studies provided strong evidence for a close isotopic coupling between nectar sugars and water to synthesized lipids in monarchs. This provided the proof of concept required to interpret isotopic values in lipids of wild migrating monarchs.

Migration and Overwinter Samples

Wing $\delta^2\text{H}$ values of monarchs captured during migration in northeast through central Mexico indicated that individuals potentially originated from many geographic regions of the

breeding grounds. Both collection site and/or year did result in some differences in mean $\delta^2\text{H}_w$ values during migration [Supplementary Table S3; $F_{(5, 442)} = 3.60$, $p = 0.003$], with Nuevo Leon, 2017 being significantly different from both Nuevo Leon and Queretaro in 2016 (Tukey's *post hoc*, $p < 0.05$). However, $\delta^2\text{H}_w$ lipid values did not differ across samples taken during migration [Figure 3 and Supplementary Table S3, $F_{(5, 442)} = 1.74$, $p = 0.13$]. There was no relationship between wing and lipid $\delta^2\text{H}$ values (Figure 3; $r^2 < 0.006$, $p > 0.71$) indicating that body lipid during migration was independent of natal origin and that the overall mean lipid $\delta^2\text{H}$ values were remarkably constant regardless of wing $\delta^2\text{H}$ values. For the purpose of illustration, we depicted origins of migrants for a subset of three sampling periods/locations in northeast to central Mexico (Figure 3): The first group was Queretaro and Nuevo Leon (2016) ($n = 130$; $\delta^2\text{H}_w = -110.4 \pm 12.4\text{‰}$; $\delta^2\text{H}_l = -324.3 \pm 14.8\text{‰}$; Figures 3A,B) and Queretaro and Guanajuato (2017) ($n = 99$; $\delta^2\text{H}_w = -107.8 \pm 13.7\text{‰}$; $\delta^2\text{H}_l = -329.5 \pm 7.8\text{‰}$; Figures 3C,D). The third group, Coahuila and Nuevo Leon (2017), was found to be bimodal for wing $\delta^2\text{H}$ values which presents problems for




FIGURE 3 | Depictions of potential natal locations and the relationship between lipid $\delta^2\text{H}$ and wing $\delta^2\text{H}$ for monarchs sampled during migration for **(A,B)** Queretaro and Nuevo Leon (2016) ($n = 130$), **(C,D)** Queretaro and Guanajuato (2017) ($n = 99$), **(E,F)** Coahuila and Nuevo Leon (2017) with wing $\delta^2\text{H} < -103\text{‰}$ ($n = 118$) and **(G,H)** Coahuila and Nuevo Leon (2017) with wing $\delta^2\text{H} > -103\text{‰}$ ($n = 98$). The legend represents the number of individuals at each pixel with wing $\delta^2\text{H}$ values consistent with those predicted for that location based on a 2:1 odds ratio. In all cases, results reveal that monarchs represented diverse breeding origins but there was, in general, no relationship between natal origin and accumulated lipid stores.

isotopic assignments (Wunder, 2010) and so, based on inspection of the distribution of δ^2H_w values (**Supplementary Figure S1**) we split this into one group with δ^2H_w values $< -103\text{‰}$ ($n = 118$; $\delta^2H_w = -116.8 \pm 8.6\text{‰}$, $\delta^2H_l = -327.4 \pm 20.1\text{‰}$; **Figures 3E,F**) and the other with δ^2H_w values $> -103\text{‰}$ ($n = 98$; $\delta^2H_w = -91.9 \pm 8.0\text{‰}$, $\delta^2H_l = -324.4 \pm 16.1\text{‰}$; **Figures 3G,H**).


A similar comparison for overwintering monarchs in 2015–2016 and 2017–2018 found that wing δ^2H values differed among colonies [$F_{(3, 87)} = 9.57$, $p < 0.001$] with two homogenous isotopic groupings (**Supplementary Table S3**). Lipid δ^2H values for the same sample also differed among overwintering collections [$F_{(4,289)} = 12.69$, $p < 0.001$] and formed two homogenous isotopic groups with 2015–2016 groups El Rosario, Piedra Herrada, Sierra Chincua and the 2017–2018 Cerro Pelon group (overall mean $-376.8 \pm 29.6\text{‰}$) showing more depleted values compared with Cerro Pelon in 2015–2016 ($-331.5 \pm 21.6\text{‰}$). For illustrative purposes, we depicted origins of monarchs sampled in overwinter sites in the single season 2015–2016 (**Figure 4**): Cerro Pelon ($n = 25$; $\delta^2H_w = -114.6 \pm 12.2\text{‰}$, $\delta^2H_l = -340.7 \pm 29.1\text{‰}$; **Figures 4A,B**) and the combined Sierra Chincua, El Rosario and Piedra Herrada sites ($n = 63$; $\delta^2H_w = -121.6 \pm 39.7\text{‰}$, $\delta^2H_l = -372.0 \pm 35.7\text{‰}$; **Figures 4C,D**). Overall, across all

years and sites, overwintering monarchs ($n = 294$) had lower lipid δ^2H than migrating individuals [$n = 448$; migration: $-326.4 \pm 15.6\text{‰}$; overwinter: $-357.6 \pm 32.4\text{‰}$; $F_{(1, 740)} = 300.7$, $p < 0.001$; **Figure 5**].

In 2018 at the Cerro Pelon overwinter site, there was no effect of sampling date or sex on lipid δ^2H values [$F_{(3, 84)} = 1.65$, $p = 0.16$; **Supplementary Figure S2** and **Supplementary Table S4**]. Lipids were not quantitatively sampled for these specimens but we found no significant relationships ($r^2 < 0.006$ in both cases) between lipid δ^2H values and either dry mass or wing length.

DISCUSSION

Our isotopic approach to investigating where and how monarch butterflies nectar to fuel the last leg of their fall migration and accumulate the necessary lipids for overwintering are encouraging and suggest a fundamentally new approach to evaluating critical nectaring strategies in this species and other animals. Our captive experiments clearly showed that stored lipids track the isotopic values of carbohydrate and water components of nectar diets. Despite a broad catchment origin of






FIGURE 5 | Difference in lipid $\delta^2\text{H}$ values (‰) between migrating and overwintering monarch butterflies based on all lipid samples analyzed across sites and years (Supplementary Table S1). Not all specimens were run for wing $\delta^2\text{H}$ analyses and so sample sizes may differ from those depicted in Figures 3, 4.

monarchs sampled across years along their Mexican migration route, monarch lipid $\delta^2\text{H}$ values were not related to origin but were clustered indicating more local nectaring *en route* and consistent with more southern nectar sources. Lipid $\delta^2\text{H}$ values of overwintering monarchs were more variable, and more negative $\delta^2\text{H}$ values compared to active migrants suggests local nectaring at higher altitudes in some individuals or the selective loss of monarchs with higher lipid $\delta^2\text{H}$ values detected in migration. Taken together, our results support the need to maintain nectaring corridors in the south including through northeast Mexico and in the vicinity of the overwintering sites. We encourage further isotopic research to ultimately derive means of probabilistically assigning monarch lipids to geographic origins via the establishment of species-specific “liposcapes” (Hobson and Wassenaar, 2019).

Our experimental analysis of captive monarchs allowed to accumulate lipids post emergence on manipulated nectar revealed that body lipids isotopically tracked dietary nectar and water sources. We estimated that, over the 5-day post-emergence period, about 59% of the carbon and 27% of the hydrogen in body lipids was derived from the sugar component of nectar and that the water in nectar contributed to about 19% of the hydrogen. The remaining components of body lipids were presumably derived from lipids formed at the larval stage. So, while we clearly have demonstrated that stored lipids can be quantitatively linked to dietary nectars, a prerequisite for applying stable

isotope techniques to ultimately track sources of nectars for wild migrating monarchs, the larval effect in our study needs to be considered. However, for monarchs migrating in the fall and at overwintering sites, we can presume that little if any larval lipid was present. In fact, it is well known that monarchs are able to rapidly synthesize lipids when conditions permit (Beall, 1948; Brown and Chippendale, 1974).

Although nectaring occurs at wintering colonies, its prevalence depends on several factors and is generally considered to be minimal and the lipid component of body mass ranges from about 73% at arrival to 37% at departure on spring migration (Brower, 1985, reviewed by Calvert and Lawton, 1993). We investigated if there was evidence for isotopic discrimination occurring through the overwintering period at the colony of Cerro Pelon to determine if neutral lipids were differentially mobilized during the winter thereby resulting in isotopic changes in the remaining lipid stores. We found no evidence for lipid isotopic change January–March and this finding supports the potential use of stored bulk lipids in individuals at overwinter colonies as a means of inferring past nectaring history.

Why might overwintering monarchs have more depleted lipid $\delta^2\text{H}$ values compared to migrants in northeast Mexico? Overwintering monarchs do require water, the degree to which depends on ambient temperature and levels of dehydration (Calvert and Lawton, 1993) and colonies are often located near a water source. This situation presents a potential hydrogen



FIGURE 6 | A migrating monarch butterfly through Nuevo Leon, Mexico. Active nectaring on flowers by migrating monarchs throughout the region is commonly observed. Photo by Omar Franco.

source to overwintering monarchs but lipids, consisting primarily of C-H bonds will not exchange with H from body water (Wassenaar, 2019). Rather, we suspect that some overwintering individuals had used nectar to form lipids at higher elevations, possibly close to the overwinter sites rather than at higher latitudes in the United States or Canada. Indeed, we expect higher $\delta^2\text{H}$ values at lower elevations and that the altitude effect (Clark and Fritz, 1997) would result in significantly lower precipitation $\delta^2\text{H}$ values at the ~ 3000 m elevations of colonies along the Transvolcanic Mountain belt, in general, and the Sierra Madre Oriental region, in particular (Wassenaar et al., 2009, see also Hobson et al., 2009a). Although our migration and wintering lipid samples were collected across years, with the exception of Cerro Pelon sampled in 2015–2016, our results indicate that monarch lipids were more variable in their $\delta^2\text{H}$ values at the overwinter colonies and also generally had lower lipid $\delta^2\text{H}$ values (by $\sim 37\text{‰}$) compared to those sampled during migration in northeastern Mexico. Further research is required to evaluate isotopic difference in stored lipids of monarchs during migration in northeast Mexico and those ultimately used by monarchs at overwintering colonies in order to evaluate the importance of late-stage nectaring just prior to and during the arrival phase at overwinter colonies. The prediction of more negative $\delta^2\text{H}$ values of lipids at high-elevation overwintering sites compared to lower elevation

migration sites from Texas and northeastern Mexico provides a potentially valuable approach to tracing the importance of flowers near the colonies.

There is much scope for future research into ways in which the stable isotope approach can be used to refine our understanding of origins of lipid synthesis in migrating monarchs and other Lepidoptera. Our captive studies using both monarchs and armyworms were necessarily of short duration (~ 6 days post eclosion) and so it was not possible to entirely remove the larval influence on adult lipid stores. Ideally, these experiments would be repeated on individuals that were held on manipulated diets for long periods and then induced to go into diapause and rapidly accumulate fat stores thereby mimicking conditions experienced during migration. Regressions from such experiments would better link $\delta^2\text{H}$ values of stored lipids to nectar sources integrated during migration and overwintering sites. However, because monarchs are nectar generalists (Tooker et al., 2002) and we expect plant nectar to vary considerably among and within plant species especially in water and sugar content (Abrahamczyk et al., 2017), an ultimate means of creating lipid-specific isoscapes (“liposcapes”) will be to either sample lipids of wild monarchs from known (i.e., constrained) nectaring sites on the breeding grounds prior to migration or captive monarchs allowed to nectar at outdoor pens exposed to natural precipitation (see Hobson et al., 1999). Such sampling would ideally be conducted across a

large geographic gradient in precipitation $\delta^2\text{H}$. That approach has proven to be extremely successful in assigning several species of migrant birds to molt origin (Hobson and Wassenaar, 2019). Of course, stored bulk lipids can be derived from single geographic sites or represent an integration of many sites spread over large spatial gradient in environmental $\delta^2\text{H}$ values. As such, in tissues like the fat body of insects, the isotope approach will always represent a probabilistic average compared to metabolically inert tissues like wing chitins. Clearly, evaluating lipid turnover rates in migrating monarchs would allow insights into expected bulk lipid isotope values. Here again, captive studies using monarchs in a “migratory phase” could expose individuals to a diet-switch (i.e., from a depleted to an enriched ^2H nectar source) that would then allow the rate of lipid synthesis in the fat body to be quantified. Further manipulations could also expose individuals to different levels of flight exercise. Finally, we fully recognize that the path ahead should involve compound-specific (fatty acid) vs. bulk lipid isotopic analyses that promise to more accurately reflect metabolic pathways leading to lipid synthesis (Whiteman et al., 2019). Since stored lipids in monarchs are primarily palmitic and oleic acids (Brower et al., 2006), the compound-specific approach focused on these fatty acids should be rewarding.

We stress that our study was conducted largely through a period of drought in southern Texas and northeast Mexico and we fully expect the dynamics of lipid synthesis and storage in migrating monarchs to change depending on fall climate conditions. Such profound influences of climate on the migratory and overwinter success of monarchs (e.g., Saunders et al., 2019) means that the wing and lipid isotopic sampling of migrant and overwintering monarchs should be conducted annually or at least every few years to elucidate nectar origins. Overwhelmingly Our work provides support for maintaining nectar corridors in the southern portion of the range and especially in Mexico along the migration route¹ and potentially in the immediate vicinity of the overwintering sites (Figure 6).

¹<https://monarchwatch.org/blog/2019/03/15/monarch-population-status-38/>

REFERENCES

- Abrahamczyk, S., Kessler, M., Hanley, D., Karger, D. N., Muller, M. P., Knauer, A. C., et al. (2017). Pollinator adaptation and the evolution of floral nectar sugar composition. *J. Evol. Biol.* 30, 112–127. doi: 10.1111/jeb.12991
- Alonso-Mejia, A., Rendon-Salina, E., Montesinos-Patino, E., and Brower, L. P. (1997). Use of lipid reserves by monarch butterflies overwintering in Mexico: implications for conservation. *Ecol. Appl.* 7, 934–947.
- Altizer, S. M., Hobson, K. A., Davis, A. K., de Roode, J. C., and Wassenaar, L. I. (2015). Do healthy monarchs migrate farther? Tracking natal origins of parasitized vs. uninfected monarch butterflies overwintering in Mexico. *PLoS One* 10:e0141371. doi: 10.1371/journal.pone.0141371
- Beall, G. (1948). The fat content of a butterfly *Danaus plexippus* Linn. As affected by migration. *Ecology* 29, 80–94. doi: 10.2307/1930346
- Bivand, R. S., and Lewin-Koh, N. (2019). *Maptools: Tools for Reading and Handling Spatial Objects, Version 0.8-39*. Available online at: <http://CRAN.R-project.org/package=maptools> (accessed September 20, 2020).
- Brower, L. P. (1985). “New perspectives on the migration biology of the monarch butterfly, *Danaus plexippus* L.” in *Migration: Mechanisms and Adaptive Significance*, ed. M. A. Rankin (Ustin, TX: University of Texas), 748–785.
- Brower, L. P., Fink, L. S., Kiphart, R. J., Pocius, V., Zubietta, R. R., and Ramirez, M. I. (2015). “Effect of the 2010–2011 drought on the lipid content of monarchs

DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, to any qualified researcher.

AUTHOR CONTRIBUTIONS

KH conceived the study. KH and JM wrote the first draft. BM, OG-R, RC-T, and EG-S collected the monarch specimens. LA conducted the captive study. KK assisted with assignments and figures. All authors edited final manuscript.

FUNDING

This work was supported by the University of Western Ontario and an NSERC Discovery grant (2017-04430) to KH.

ACKNOWLEDGMENTS

Stable isotope analyses were performed by the KH at the LSIS-AFAR stable isotope facility. We thank Dr. Isabel Ramirez for assisting with sample collections that formed part of her contract with the Commission for Environmental Cooperation. We also thank Orley (Chip) Taylor and reviewer for their constructive comments on an earlier version of the manuscript.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fevo.2020.572140/full#supplementary-material>

- migrating through Texas to overwintering sites in Mexico,” in *Monarchs in a Changing World: Biology and Conservation of an Iconic Butterfly*, eds K. S. Oberhauser, K. R. Nail, and S. Altizer (Ithaca: Cornell University Press), 117–129.
- Brower, L. P., Fink, L. S., and Walford, P. (2006). Fueling the fall migration of the monarch butterfly. *Integrat. Comp. Biol.* 46, 1123–1142. doi: 10.1093/icb/icl029
- Brown, J. J., and Chippendale, G. M. (1974). Migration of the monarch butterfly *Danaus plexippus*: energy sources. *J. Insect Physiol.* 20, 1117–1130. doi: 10.1016/0022-1910(74)90218-2
- Calvert, W. H., and Lawton, R. O. (1993). “Comparative phenology of variation in size, weight, and water content of eastern North American monarch butterflies at five overwintering sites in Mexico,” in *Biology and Conservation of the Monarch Butterfly*, eds S. B. Malcolm and M. P. Zalucki (Los Angeles: Natural History Museum of Los Angeles), 299–307.
- Cenedella, R. J. (1971). The lipids of the female monarch butterfly, *Danaus plexippus*, during fall migration. *Insect Biochem.* 1, 244–247. doi: 10.1016/0020-1790(71)90077-1
- Clark, I. D., and Fritz, P. (1997). *Environmental Isotopes in Hydrogeology*. London: CRC Press.
- Flockhart, D. T., Brower, L. P., Ramirez, M. I., Hobson, K. A., Wassenaar, L. I., Altizer, S., et al. (2017). Regional climate on the breeding grounds predicts

- variation in the natal origin of monarch butterflies overwintering in Mexico over 38 years. *Glob. Chang. Biol.* 23, 2565–2576. doi: 10.1111/gcb.13589
- Flockhart, T., Wassenaar, L. I., Martin, T., Hobson, K. A., Wunder, M., and Norris, D. R. (2013). Tracking multi-generational colonization of the breeding grounds by monarch butterflies in eastern North America. *Proc. Roy. Soc. Lond. B* 280, 1–8. doi: 10.1155/2009/705780
- Gibo, D. L., and McCurdy, J. A. (1993). Lipid accumulation by migrating monarch butterflies (*Danaus plexippus* L.). *Can. J. Zool.* 71, 76–82. doi: 10.1139/z93-012
- Hijmans, R. J. (2020). *raster: Geographic Data Analysis and Modeling, Version 2.5-8*. Available online at: <http://CRAN.R-project.org/package=raster> (accessed September 20, 2020).
- Hobson, K. A., Van Wilgenburg, S. L., Larson, K., and Wassenaar, L. I. (2009a). A feather hydrogen isotope for Mexico. *J. Geochem. Explor.* 102, 63–70. doi: 10.1016/j.gexplo.2009.02.002
- Hobson, K. A., and Wassenaar, L. I. (eds) (2019). *Tracking Animal Migration with Stable Isotopes*, 2nd Edn, London: Academic Press.
- Hobson, K. A., Wassenaar, L. I., and Taylor, O. R. (1999). Stable isotopes (δD and $\delta^{13}C$) are geographic indicators of natal origins of monarch butterflies in eastern North America. *Oecologia* 120, 397–404. doi: 10.1007/s004420050872
- Hobson, K. A., Wunder, M. B., Van Wilgenburg, S. L., Clark, R. G., and Wassenaar, L. I. (2009b). A Method for investigating population declines of migratory birds using stable isotopes: origins of harvested lesser scaup in North America. *PLoS One* 4:e7915. doi: 10.1371/journal.pone.0007915
- IAEA/WMO (2015). *Global Network of Isotopes in Precipitation. The GNIP Database*. Available online at: <http://www.iaea.org/water> (accessed May 17, 2020).
- IBM Corp (2016). *Released 2016. IBM SPSS Statistics for Windows, Version 24.0*. Armonk, NY: IBM Corp.
- Lennox, R. J., Chapman, J. M., Souliere, C. M., Tudorache, C., Wikelski, M., Metcalfe, J. D., et al. (2016). Conservation physiology of animal migration. *Conserv. Physiol.* 4:cov072. doi: 10.1093/conphys/cov072
- McWilliams, S. R., Guglielmo, C. G., Pierce, B., and Klaassen, M. (2004). Flying, fasting and feeding in birds during migration: a nutritional and physiological ecology perspective. *J. Avian Biol.* 35, 377–393. doi: 10.1111/j.0908-8857.2004.03378.x
- Miller, N. G., Wassenaar, L. I., Hobson, K. A., and Norris, D. R. (2012). Migratory connectivity of the monarch butterfly (*Danaus plexippus*): patterns of spring re-colonization in eastern North America. *PLoS One* 7:e0031891. doi: 10.1371/journal.pone.0031891
- Moore, F. R., Gauthreaux, S. A., Kerlinger, P., and Simons, T. R. (1993). "Habitat requirements during migration: Important link in conservation," in *Ecology and Management of Neotropical Migratory Birds*, eds T. E. Martin and D. M. Finch (Fort Collins, CO: USDA Forest Service), 121–144.
- R Core Team (2019). *R: A Language and Environment for Statistical Computing, Version 3.3.2*. Vienna: The R Foundation for Statistical Computing.
- Saunders, S. P., Ries, L., Neupane, N., Ramírez, M. I., García-Serrano, E. U., Rendón-Salinas, E., et al. (2019). Multiscale seasonal factors drive the size of winter monarch colonies. *Proc. Nat. Acad. Sci. U.S.A.* 116, 8609–8614. doi: 10.1073/pnas.1805114116
- Soto, D. X., Koehler, G., Wassenaar, L. I., and Hobson, K. A. (2017). Determination of stable hydrogen isotopic compositions of complex organic materials: contrasting the role of exchangeable hydrogen and residual moisture. *Rapid Comm. Mass Spectrom.* 31, 1193–1203.
- Taylor, O. R., Lovett, J. P., Gibo, D. L., Weiser, E. L., Thogmartin, W. E., Semmens, D. J., et al. (2019). Is the timing, pace and success of the monarch migration associated with sun angle? *Front. Ecol. Evol.* 7:442. doi: 10.3389/fevo.2019.00442
- Terzer, S., Wassenaar, L. I., Araguás-Araguás, L. J., and Aggarwal, P. K. (2013). Global isoscapes for $\delta^{18}O$ and δ^2H in precipitation: improved prediction using regionalized climatic regression models. *Hydrol. Earth Syst. Sci.* 17, 4713–4728. doi: 10.5194/hess-17-4713-2013
- Tooker, J. F., Reigel, P. F., and Hanks, L. M. (2002). Nectar sources of dayflying lepidoptera of Central Illinois. *Ann. Entomol. Soc. Am.* 95, 84–96. doi: 10.1603/0013-8746(2002)095[0084:nsodf]2.0.co;2
- Van Wilgenburg, S. L., Hobson, K. A., Brewster, K. R., and Welker, J. M. (2012). Addressing uncertainty in assessing dispersal in threatened migratory species using stable hydrogen isotope analysis of feathers. *Endang. Spec. Res.* 16, 17–29. doi: 10.3354/esr00383
- Wassenaar, L. I. (2019). "Introduction to conducting stable isotope measurements for animal migration studies," in *Tracking Animal Migration with Stable Isotopes*, 2nd Edn, eds K. A. Hobson and L. I. Wassenaar (London: Academic Press), 25–51. doi: 10.1016/b978-0-12-814723-8.00002-7
- Wassenaar, L. I., and Hobson, K. A. (1998). Natal origins of migratory Monarch Butterflies at wintering colonies in Mexico: New isotopic evidence. *Proc. Nat. Acad. Sci. U.S.A.* 95, 15436–15439. doi: 10.1073/pnas.95.26.15436
- Wassenaar, L. I., and Hobson, K. A. (2003). Comparative equilibration and online technique for determination of non-exchangeable hydrogen of keratins for use in animal migration studies. *Isotopes Environ. Health Stud.* 39, 1–7.
- Wassenaar, L. I., Van Wilgenburg, S. L., Larson, K., and Hobson, K. A. (2009). A Groundwater Isoscape (δD , $\delta^{18}O$) for Mexico. *J. Geochem. Explor.* 102, 123–136. doi: 10.1016/j.gexplo.2009.01.001
- West, J. B., Kreuzer, H. W., and Ehleringer, J. R. (2010). "Approaches to plant hydrogen and oxygen isoscapes generation," in *Isoscapes*, eds J. West, G. Bowen, T. Dawson, and K. Tu (Dordrecht: Springer).
- West, J. B., Sobek, A., and Ehleringer, J. R. (2008). A simplified GIS approach to modeling global leaf water isoscapes. *PLoS One* 3:e2447. doi: 10.1371/journal.pone.0002447
- Whiteman, J. P., Elliott Smith, E. A., Besser, A. C., and Newsome, S. D. (2019). A guide to using compound-specific stable isotope analysis to study the fates of molecules in organisms and ecosystems. *Diversity* 11, 1–18.
- Williams, E. H., and Brower, L. P. (2015). "Microclimate protection of overwintering monarchs provided by Mexico's high-elevation oyamel fir forests," in *Monarchs in a Changing World: Biology and Conservation of an Iconic Butterfly*, eds K. S. Oberhauser, K. R. Nail, and S. Altizer (Ithaca: Cornell University Press), 109–116.
- Wunder, M. (2010). "Using isoscapes to model probability surfaces for determining geographic origins," in *Isoscapes*, eds J. West, G. Bowen, T. Dawson, and K. Tu (Dordrecht: Springer).

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Hobson, García-Rubio, Carrera-Treviño, Anparasan, Kardynal, McNeil, García-Serrano and Mora Alvarez. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.