

Iteration-complexity of block-decomposition algorithms and the alternating direction method of multipliers *

Renato D. C. Monteiro[†] Benar F. Svaiter[‡]

August 1, 2010

(Revised: August 22, 2011, August 8, 2012 and October 30, 2012)

Abstract

In this paper, we consider the monotone inclusion problem consisting of the sum of a continuous monotone map and a point-to-set maximal monotone operator with a separable two-block structure and introduce a framework of block-decomposition prox-type algorithms for solving it which allows for each one of the single-block proximal subproblems to be solved in an approximate sense. Moreover, by showing that any method in this framework is also a special instance of the hybrid proximal extragradient (HPE) method introduced by Solodov and Svaiter, we derive corresponding convergence rate results. We also describe some instances of the framework based on specific and inexpensive schemes for solving the single-block proximal subproblems. Finally, we consider some applications of our methodology to establish for the first time: i) the iteration-complexity of an algorithm for finding a zero of the sum of two arbitrary maximal monotone operators and, as a consequence, the ergodic iteration-complexity of the Douglas-Rachford splitting method, and; ii) the ergodic iteration-complexity of the classical alternating direction method of multipliers for a class of linearly constrained convex programming problems with proper closed convex objective functions.

2000 Mathematics Subject Classification: Primary, 90C60, 49M27, 90C25; Secondary, 47H05, 47N10, 64K05, 65K10.

Key words: decomposition; complexity; monotone operator; inclusion problem; proximal; extragradient

1 Introduction

A broad class of optimization, saddle point, equilibrium and variational inequality (VI) problems can be posed as the *monotone inclusion problem*, namely: finding x such that $0 \in T(x)$, where T is a maximal monotone point-to-set operator. The proximal point method, proposed by Rockafellar [23], is a classical iterative scheme for solving the monotone inclusion problem which generates a sequence $\{z_k\}$ according to

$$\|z_k - (\lambda_k T + I)^{-1}(z_{k-1})\| \leq e_k, \quad \sum_{k=1}^{\infty} e_k < \infty.$$

This method has been used as a generic framework for the design and analysis of several implementable algorithms. Observe that $\{e_k\}$ is a (summable) sequence of errors bounds.

*The title of the first version of this work was “Iteration-complexity of block-decomposition algorithms and the minimization augmented Lagrangian method”.

[†]School of Industrial and Systems Engineering, Georgia Institute of Technology, Atlanta, GA, 30332-0205. (email: monteiro@isye.gatech.edu). The work of this author was partially supported by NSF Grants CCF-0808863 and CMMI-0900094 and ONR Grants ONR N00014-08-1-0033 and N00014-11-1-0062.

[‡]IMPA, Estrada Dona Castorina 110, 22460-320 Rio de Janeiro, Brazil (email: benar@impa.br). The work of this author was partially supported by CNPq grants 480101, 474944, 303583, 302962, FAPERJ grant E-26/102.821, E-26/102.940 and PRONEX-Optimization

New inexact versions of the proximal point method which uses instead relative error criteria were proposed by Solodov and Svaiter [25, 26, 27, 28]. In this work we use one of these variants namely the hybrid proximal-extragradient (HPE) method [25] to develop and analyze block decomposition algorithms, and we now briefly discuss this method. The *exact* proximal point iteration from z with stepsize $\lambda > 0$ is given by $z_+ = (\lambda T + I)^{-1}(z)$, which is equivalent to

$$v \in T(z_+), \quad \lambda v + z_+ - z = 0. \quad (1)$$

In each step of the HPE, the above *proximal system* is solved inexactly with $(z, \lambda) = (z_{k-1}, \lambda_k)$ to obtain $z_k = z_+$ as follows. For a given constant $\sigma \in [0, 1]$, a triple $(\tilde{z}, \tilde{v}, \varepsilon) = (\tilde{z}_k, \tilde{v}_k, \varepsilon_k)$ is found such that

$$\tilde{v} \in T^\varepsilon(\tilde{z}), \quad \|\lambda \tilde{v} + \tilde{z} - z\|^2 + 2\lambda \varepsilon \leq \sigma^2 \|\tilde{z} - z\|^2, \quad (2)$$

where T^ε denotes the ε -enlargement [1] of T . (It has the property that $T^\varepsilon(z) \supset T(z)$ for each z .) Note that this construction relaxes both the inclusion and the equation in (1). Finally, instead of choosing \tilde{z} as the next iterate z_+ , the HPE method computes the next iterate z_+ by means of the following *extragradient* step:

$$z_+ = z - \lambda \tilde{v}.$$

Iteration complexity results for the HPE method were established in [15] and these results depend on the distance of the initial iterate to the solution set instead of the diameter of the feasible set.

By viewing Korpelevich's method as well as Tseng's modified forward-backward splitting (MF-BS) method [30] as special cases of the HPE method, the authors have established in [15, 16] the pointwise and ergodic iteration-complexities of these methods applied to either: monotone variational inequalities problems, the monotone inclusion problems for the sum of a Lipschitz continuous monotone map with a maximal monotone operator whose resolvent is assumed to be easily computable, convex-concave saddle point problems, or a large class of linearly constrained convex programming problems, including for example cone programming and problems whose objective functions converge to infinity as the boundaries of their domain are approached. In the context of variational inequality problems, we should mention that prior to [15, 16] Nemirovski [17] has established the ergodic iteration-complexity of Korpelevich's method under the assumption that the feasible set of the problem is bounded, and Nesterov [18] has established the ergodic iteration-complexity of a new dual extrapolation algorithm whose termination depends on the guess of a ball centered at the initial iterate.

In this paper, we continue along the same line of investigation as in our previous papers [15] and [16], which is to use the HPE method as a general framework to derive iteration-complexity results for specific algorithms for solving various types of structured monotone inclusion problems. More specifically, we consider the monotone inclusion problem consisting of the sum of a continuous monotone map and a point-to-set maximal monotone operator with a separable two-block structure, namely:

$$0 \in T(x, y) := \left\{ \begin{pmatrix} F_x(x, y) + a \\ F_y(x, y) + b \end{pmatrix} : a \in A(x), b \in B(y) \right\}. \quad (3)$$

We introduce a general block-decomposition HPE (BD-HPE) framework in the context of this inclusion problem, which allows for each one of the single-block proximal subproblems to be solved in an approximate sense. More specifically, given a pair $((x, y), \lambda) = ((x_{k-1}, y_{k-1}), \lambda_k)$, an instance of the BD-HPE framework computes an approximate solution $((\tilde{x}, \tilde{y}), (\tilde{v}_x, \tilde{v}_y), \varepsilon)$ of (1) (in the sense of (2)) with T given by (3) by first computing an approximate solution $(\tilde{x}, \tilde{v}_x, \varepsilon_x)$ of (1) with $T = F_x(\cdot, y) + A(\cdot)$, then computing an approximate solution $(\tilde{y}, \tilde{v}_y, \varepsilon_y)$ of (1) with $T = F_y(\tilde{x}, \cdot) + B(\cdot)$, and finally setting $\varepsilon = \varepsilon_x + \varepsilon_y$. Moreover, by showing that any method in this framework is also a special instance of the HPE method, we derive convergence rate results for the BD-HPE framework based on the ones developed in [15] for the HPE method.

Subsequently, we describe some ways of implementing the BD-HPE framework based on specific and inexpensive schemes for solving the single-block proximal subproblems. We also consider some applications of our methodology introduced here to establish for the first time: i) the iteration-complexity of an algorithm for finding a zero of the sum of two arbitrary maximal monotone operators and, as a consequence, the ergodic

iteration-complexity of the Douglas-Rachford splitting method, and; ii) the ergodic iteration-complexity of the classical alternating direction method of multipliers (ADMM) for a class of linearly constrained convex programming problems with proper closed convex objective functions.

The ADMM was first introduced in [10, 11]. Recently, there has been some growing interest in the ADMM for solving large scale linear cone programming (see for example [5, 4, 21, 12, 14]). However, to the best of our knowledge, no iteration-complexity analysis for the ADMM have been established so far. Development and analysis of splitting and block-decomposition (BD) methods is by now a well-developed area, although algorithms which allow a relative error tolerance in the solution of the proximal subproblems have been studied in just a few papers. In particular, Ouorou [20] discusses an ε -proximal decomposition using the ε -subdifferential and a relative error criterion on ε . Projection splitting methods for the sum of arbitrary maximal monotone operators using a particular case of the HPE error tolerance for solving the proximal subproblems were presented in [7, 8]. The use of the HPE method for studying BD methods was first presented in [24]. We observe however that none of these works deal with the derivation of iteration-complexity bounds. More recently, Chambolle and Pock [6] have developed and established iteration-complexity bounds for a BD method, which solves the proximal subproblems exactly, in the context of saddle-point problems with a bilinear coupling.

This paper is organized as follows. Section 2 contains two subsections. Subsection 2.1 reviews some basic definitions and facts on convex functions and the definition and some basic properties of the ε -enlargement of a point-to-set maximal monotone operator. Subsection 2.2 reviews the HPE method and the global convergence rate results obtained for it in [15]. Section 3 introduces the BD-HPE framework for solving a special type of monotone inclusion problem mentioned above and shows that any instance of the framework can be viewed as a special case of the HPE method. As a consequence, global convergence rate results for the BD-HPE framework are also obtained in this section using the general theory outlined in Subsection 2.2. Section 4 describes specific schemes for solving the single-block proximal subproblems based on a small number (one or two) of resolvent evaluations. Section 5 describes some instances of the BD-HPE framework which, are not only interesting in their own right, but also illustrate the use of the different schemes for solving the single-block proximal subproblems. It contains three subsections as follows. Subsection 5.1 discusses a specific instance of the BD-HPE framework where both single-block proximal subproblems are solved exactly. Subsection 5.2 gives another instance of the BD-HPE framework in which both single-block proximal subproblems are approximately solved by means of a Tseng's type scheme. Subsection 5.3 studies a BD method for a large class of linearly constrained convex optimization problems, which includes cone programs and problems whose objective functions converge to infinity as the relative boundaries of their domain are approached. Section 6 considers the monotone inclusion problem consisting of the sum of two maximal monotone operators and show how it can be transformed to an equivalent monotone inclusion problem with two-block structure of the aforementioned type, which can then be solved by any instance of the BD-HPE framework. Section 7 considers the ADMM for solving a class of linearly constrained convex programming problems with proper closed convex objective functions and shows that it can be interpreted as a specific instance of the BD-HPE framework applied to a two-block monotone inclusion problem.

1.1 Notation

We denote the set of real numbers by \mathbb{R} and nonnegative numbers by \mathbb{R}_+ . For a real symmetric matrix E , we denote its largest eigenvalue by $\theta_{\max}(E)$. The domain of definition of a one-to-one function F is denoted by $\text{Dom } F$. The effective domain of a function $f : \mathbb{R}^n \rightarrow [-\infty, \infty]$ is defined as $\text{dom } f := \{x \in \mathbb{R}^n : f(x) < \infty\}$.

2 Technical background

This section contains two subsections. In the first one, we review some basic definitions and facts about convex functions and ε -enlargement of monotone multi-valued maps. This subsection also reviews the weak transportation formula for the ε -subdifferentials of closed convex functions and the ε -enlargements of maximal

monotone operators. The second subsection reviews the HPE method and the global convergence rate results obtained for it in [15].

2.1 The ε -subdifferential and ε -enlargement of monotone operators

Let \mathcal{Z} denote a finite dimensional inner product space with inner product and associated norm denoted by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$. A point-to-set operator $T : \mathcal{Z} \rightrightarrows \mathcal{Z}$ is a relation $T \subset \mathcal{Z} \times \mathcal{Z}$ and

$$T(z) = \{v \in \mathcal{Z} \mid (z, v) \in T\}.$$

Alternatively, one can consider T as a multi-valued function of \mathcal{Z} into the family $\wp(\mathcal{Z}) = 2^{(\mathcal{Z})}$ of subsets of \mathcal{Z} . Regardless of the approach, it is usual to identify T with its graph defined as

$$Gr(T) = \{(z, v) \in \mathcal{Z} \times \mathcal{Z} \mid v \in T(z)\}.$$

The domain of T , denoted by $\text{Dom } T$, is defined as

$$\text{Dom } T := \{z \in \mathcal{Z} : T(z) \neq \emptyset\}.$$

An operator $T : \mathcal{Z} \rightrightarrows \mathcal{Z}$ is *affine* if its graph is an affine manifold. Clearly, if T is affine, then the following implication holds:

$$\left. \begin{array}{l} \alpha_i \geq 0, \quad i = 1, \dots, k \\ \alpha_1 + \dots + \alpha_k = 1 \\ v_i \in T(z_i), \quad 1, \dots, k \end{array} \right\} \implies \sum_{i=1}^k \alpha_i v_i \in T \left(\sum_{i=1}^k \alpha_i z_i \right). \quad (4)$$

Moreover, $T : \mathcal{Z} \rightrightarrows \mathcal{Z}$ is *monotone* if

$$\langle v - \tilde{v}, z - \tilde{z} \rangle \geq 0, \quad \forall (z, v), (\tilde{z}, \tilde{v}) \in Gr(T),$$

and T is *maximal monotone* if it is monotone and maximal in the family of monotone operators with respect to the partial order of inclusion, i.e., $S : \mathcal{Z} \rightrightarrows \mathcal{Z}$ monotone and $Gr(S) \supset Gr(T)$ implies that $S = T$.

In [1], Burachik, Iusem and Svaiter introduced the ε -enlargement of maximal monotone operators. In [15] this concept was extended to a generic point-to-set operator in \mathcal{Z} as follows. Given $T : \mathcal{Z} \rightrightarrows \mathcal{Z}$ and a scalar ε , define $T^\varepsilon : \mathcal{Z} \rightrightarrows \mathcal{Z}$ as

$$T^\varepsilon(z) = \{v \in \mathcal{Z} \mid \langle z - \tilde{z}, v - \tilde{v} \rangle \geq -\varepsilon, \quad \forall \tilde{z} \in \mathcal{Z}, \forall \tilde{v} \in T(\tilde{z}), \quad \forall z \in \mathcal{Z}. \quad (5)$$

We now state a few useful properties of the operator T^ε that will be needed in our presentation.

Proposition 2.1. *Let $T, T' : \mathcal{Z} \rightrightarrows \mathcal{Z}$. Then,*

- a) *if $\varepsilon_1 \leq \varepsilon_2$, then $T^{\varepsilon_1}(z) \subset T^{\varepsilon_2}(z)$ for every $z \in \mathcal{Z}$;*
- b) *$T^\varepsilon(z) + (T')^{\varepsilon'}(z) \subset (T + T')^{\varepsilon + \varepsilon'}(z)$ for every $z \in \mathcal{Z}$ and $\varepsilon, \varepsilon' \in \mathbb{R}$;*
- c) *T is monotone if, and only if, $T \subset T^0$;*
- d) *T is maximal monotone if, and only if, $T = T^0$;*

Observe that items a) and d) of the above proposition imply that, if $T : \mathcal{Z} \rightrightarrows \mathcal{Z}$ is maximal monotone, then

$$T(z) \subset T^\varepsilon(z) \quad \forall z \in \mathcal{Z}, \quad \varepsilon \geq 0,$$

so that $T^\varepsilon(z)$ is indeed an enlargement of $T(z)$.

Note that, due to the definition of T^ε , the verification of the inclusion $v \in T^\varepsilon(z)$ requires checking an infinite number of inequalities. This verification is feasible only for specially-structured instances of operators T . However, it is possible to compute points in the graph of T^ε using the following *weak transportation formula* [2]. This formula will be used in the complexity analysis of the ergodic mean.

Theorem 2.2 ([2, Theorem 2.3]). *Suppose that $T : \mathcal{Z} \rightrightarrows \mathcal{Z}$ is maximal monotone. Let $z_i, v_i \in \mathcal{Z}$ and $\varepsilon_i, \alpha_i \in \mathbb{R}_+$, for $i = 1, \dots, k$, be such that*

$$v_i \in T^{\varepsilon_i}(z_i), \quad i = 1, \dots, k, \quad \sum_{i=1}^k \alpha_i = 1,$$

and define

$$z^a := \sum_{i=1}^k \alpha_i z_i, \quad v^a := \sum_{i=1}^k \alpha_i v_i, \\ \varepsilon^a := \sum_{i=1}^k \alpha_i [\varepsilon_i + \langle z_i - z^a, v_i - v^a \rangle] = \sum_{i=1}^k \alpha_i [\varepsilon_i + \langle z_i - z^a, v_i \rangle].$$

Then, the following statements hold:

- a) $\varepsilon^a \geq 0$ and $v^a \in T^{\varepsilon^a}(z^a)$;
- b) if, in addition, $T = \partial f$ for some proper lower semi-continuous convex function f and $v_i \in \partial_{\varepsilon_i} f(z_i)$ for $i = 1, \dots, k$, then $v^a \in \partial_{\varepsilon^a} f(z^a)$.

Finally, we refer the reader to [3, 29] for further discussion on the ε -enlargement of a maximal monotone operator.

For a scalar $\varepsilon \geq 0$, the ε -subdifferential of a function $f : \mathcal{Z} \rightarrow [-\infty, +\infty]$ is the operator $\partial_\varepsilon f : \mathcal{Z} \rightrightarrows \mathcal{Z}$ defined as

$$\partial_\varepsilon f(z) = \{v \mid f(\tilde{z}) \geq f(z) + \langle \tilde{z} - z, v \rangle - \varepsilon, \forall \tilde{z} \in \mathcal{Z}\}, \quad \forall z \in \mathcal{Z}. \quad (6)$$

When $\varepsilon = 0$, the operator $\partial_\varepsilon f$ is simply denoted by ∂f and is referred to as the subdifferential of f . The operator ∂f is trivially monotone if f is proper. If f is a proper lower semi-continuous convex function, then ∂f is maximal monotone [22].

The conjugate f^* of f is the function $f^* : \mathcal{Z} \rightarrow [-\infty, \infty]$ defined as

$$f^*(v) = \sup_{z \in \mathcal{Z}} \langle v, z \rangle - f(z), \quad \forall v \in \mathcal{Z}.$$

The following result lists some useful properties about the ε -subdifferential of a proper convex function.

Proposition 2.3. *Let $f : \mathcal{Z} \rightarrow (-\infty, \infty]$ be a proper convex function. Then,*

- a) $\partial_\varepsilon f(z) \subset (\partial f)^\varepsilon(z)$ for any $\varepsilon \geq 0$ and $z \in \mathcal{Z}$;
- b) $\partial_\varepsilon f(z) = \{v \mid f(z) + f^*(v) \leq \langle z, v \rangle + \varepsilon\}$ for any $\varepsilon \geq 0$ and $z \in \mathcal{Z}$;
- c) if $v \in \partial f(z)$ and $f(\tilde{z}) < \infty$, then $v \in \partial_\varepsilon f(\tilde{z})$, where $\varepsilon := f(\tilde{z}) - [f(z) + \langle \tilde{z} - z, v \rangle]$.

For the following definitions, assume that $Z \subset \mathcal{Z}$ is a nonempty closed convex set. The *indicator function* of Z is the function $\delta_Z : \mathcal{Z} \rightarrow [0, \infty]$ defined as

$$\delta_Z(z) = \begin{cases} 0, & z \in Z, \\ \infty, & \text{otherwise,} \end{cases}$$

and the *normal cone operator* of Z is the point-to-set map $N_Z : \mathcal{Z} \rightrightarrows \mathcal{Z}$ given by

$$N_Z(z) = \begin{cases} \emptyset, & z \notin Z, \\ \{v \in \mathcal{Z}, \mid \langle \tilde{z} - z, v \rangle \leq 0, \forall \tilde{z} \in Z\}, & z \in Z. \end{cases} \quad (7)$$

Clearly, the normal cone operator N_Z of Z can be expressed in terms of δ_Z as $N_Z = \partial \delta_Z$. The *orthogonal projection* $P_Z : \mathcal{Z} \rightarrow \mathcal{Z}$ onto Z is defined as

$$P_Z(z) = \operatorname{argmin}_{z' \in Z} \|z' - z\| \quad \forall z \in \mathcal{Z}.$$

It is well-known that P_Z is the resolvent of the normal cone operator, that is, $P_Z = (\lambda N_Z + I)^{-1}$ for every $\lambda > 0$.

2.2 The hybrid proximal extragradient method

This subsection reviews the HPE method and corresponding global convergence rate results obtained in [15].

Let $T : \mathcal{Z} \rightrightarrows \mathcal{Z}$ be maximal monotone operator. The monotone inclusion problem for T consists of finding $z \in \mathcal{Z}$ such that

$$0 \in T(z).$$

We also assume throughout this section that this problem has a solution, that is, $T^{-1}(0) \neq \emptyset$.

We next review the hybrid proximal extragradient method introduced in [25] for solving the above problem and state the iteration-complexity results obtained for it in [15].

Hybrid Proximal Extragradient Method:

- 0) Let $z_0 \in \mathcal{Z}$ and $0 \leq \sigma \leq 1$ be given and set $k = 1$;
- 1) choose $\lambda_k > 0$ and find $\tilde{z}_k, \tilde{v}_k \in \mathcal{Z}$, $\sigma_k \in [0, \sigma]$ and $\varepsilon_k \geq 0$ such that

$$\tilde{v}_k \in T^{\varepsilon_k}(\tilde{z}_k), \quad \|\lambda_k \tilde{v}_k + \tilde{z}_k - z_{k-1}\|^2 + 2\lambda_k \varepsilon_k \leq \sigma_k^2 \|\tilde{z}_k - z_{k-1}\|^2; \quad (8)$$

- 2) define $z_k = z_{k-1} - \lambda_k \tilde{v}_k$, set $k \leftarrow k + 1$, and go to step 1.

end

We now make several remarks about the HPE method. First, the HPE method does not specify how to choose λ_k and how to find \tilde{z}_k, \tilde{v}_k and ε_k as in (8). The particular choice of λ_k and the algorithm used to compute \tilde{z}_k, \tilde{v}_k and ε_k will depend on the particular implementation of the method and the properties of the operator T . Second, if $\tilde{z} := (\lambda_k T + I)^{-1} z_{k-1}$ is the *exact* proximal point iterate, or equivalently

$$\tilde{v} \in T(\tilde{z}), \quad (9)$$

$$\lambda_k \tilde{v} + \tilde{z} - z_{k-1} = 0, \quad (10)$$

for some $\tilde{v} \in \mathcal{Z}$, then $(\tilde{z}_k, \tilde{v}_k) = (\tilde{z}, \tilde{v})$ and $\varepsilon_k = 0$ satisfies (8). Therefore, the error criterion (8) relaxes the inclusion (9) to $\tilde{v} \in T^\varepsilon(\tilde{z})$ and relaxes equation (10) by allowing a small error relative to $\|\tilde{z}_k - z_{k-1}\|$.

We now state a few results about the convergence behaviour of the HPE method. The proof of the following result can be found in Lemma 4.2 of [15]. It provides a computable estimate of how much the square of the distance of an arbitrary solution to an iterate of the HPE method decreases from one iteration to the next one.

Proposition 2.4. *For any $z^* \in T^{-1}(0)$, the sequence $\{\|z^* - z_k\|\}$ is non-increasing and*

$$\|z^* - z_0\|^2 \geq \|z^* - z_k\|^2 + \sum_{i=1}^k \left[\|\tilde{z}_i - z_{i-1}\|^2 - (\|\lambda_i \tilde{v}_i + \tilde{z}_i - z_{i-1}\|^2 + 2\lambda_i \varepsilon_i) \right] \quad (11)$$

$$\geq \|z^* - z_k\|^2 + (1 - \sigma^2) \sum_{i=1}^k \|\tilde{z}_i - z_{i-1}\|^2. \quad (12)$$

The proof of the following result which establishes the convergence rate of the residual $(\tilde{v}_k, \varepsilon_k)$ of z_k can be found in Theorem 4.4 of [15].

Theorem 2.5. *Assume that $\sigma < 1$ and let d_0 be the distance of z_0 to $T^{-1}(0)$. Then, for every $k \in \mathbb{N}$, $\tilde{v}_k \in T^{\varepsilon_k}(\tilde{z}_k)$ and there exists an index $i \leq k$ such that*

$$\|\tilde{v}_i\| \leq d_0 \sqrt{\frac{1 + \sigma}{1 - \sigma} \left(\frac{1}{\sum_{j=1}^k \lambda_j^2} \right)}, \quad \varepsilon_i \leq \frac{\sigma^2 d_0^2 \lambda_i}{2(1 - \sigma^2) \sum_{j=1}^k \lambda_j^2}. \quad (13)$$

Theorem 2.5 estimate the quality of the best among the iterates $\tilde{z}_1, \dots, \tilde{z}_k$. We will refer to these estimates as the *pointwise* complexity bounds for the HPE method.

We will now describe alternative estimates for the HPE method which we refer to as the *ergodic* complexity bounds. The next result describes the convergence properties of an ergodic sequence associated with $\{\tilde{z}_k\}$.

Theorem 2.6. *For every $k \in \mathbb{N}$, define*

$$\tilde{z}_k^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i \tilde{z}_i, \quad \tilde{v}_k^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i \tilde{v}_i, \quad \varepsilon_k^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i (\varepsilon_i + \langle \tilde{z}_i - \tilde{z}_k^a, \tilde{v}_i \rangle), \quad (14)$$

where $\Lambda_k := \sum_{i=1}^k \lambda_i$. Then, for every $k \in \mathbb{N}$,

$$\tilde{v}_k^a = \frac{1}{\Lambda_k} (z_0 - z_k) \in T^{\varepsilon_k^a}(\tilde{z}_k^a), \quad \|\tilde{v}_k^a\| \leq \frac{2d_0}{\Lambda_k}, \quad (15)$$

and

$$0 \leq \varepsilon_k^a \leq \frac{1}{2\Lambda_k} [2\langle \tilde{z}_k^a - z_0, z_k - z_0 \rangle - \|z_k - z_0\|^2] \leq \frac{2d_0^2}{\Lambda_k} (1 + \rho_k), \quad (16)$$

where d_0 is the distance of z_0 to $T^{-1}(0)$, and

$$\rho_k := \frac{1}{d_0} \|\tilde{z}_k^a - z_k^a\| \leq \frac{1}{d_0} \max_{i=1, \dots, k} \|\tilde{z}_i - z_i\|, \quad \text{where } z_k^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i z_i. \quad (17)$$

Moreover, the sequence $\{\rho_k\}$ is bounded under either one of the following situations:

a) if $\sigma < 1$ then

$$\rho_k \leq \frac{\sigma \sqrt{\tau_k}}{\sqrt{(1 - \sigma^2)}}, \quad \text{where } \tau_k = \max_{i=1, \dots, k} \frac{\lambda_i}{\Lambda_k} \leq 1; \quad (18)$$

b) $\text{Dom } T$ is bounded, in which case

$$\rho_k \leq \frac{D}{d_0} + 1,$$

where $D := \sup\{\|y - y'\| : y, y' \in \text{Dom } T\}$ is the diameter of $\text{Dom } T$.

Proof. The bounds (15) and (16) and statement a) follow immediately from Proposition 4.6 and the proof of Theorem 4.7 of [15]. Let z^* be the closest point to z_0 lying in $T^{-1}(0)$. Relation (17), the triangle inequality for norms, Proposition 2.4 and the definition of D imply that

$$\rho_k \leq \frac{1}{d_0} \max_{i=1, \dots, k} (\|\tilde{z}_i - z^*\| + \|z^* - z_i\|) \leq \frac{1}{d_0} (D + \|z^* - z_0\|) = \frac{D}{d_0} + 1. \quad \square$$

Note that the rate of convergence (16) in Theorem 2.6 is only useful if we know how to bound ρ_k , which is the case when $\sigma < 1$ or $\text{Dom } T$ is bounded. When $\sigma = 1$ and $\text{Dom } T$ is unbounded, we do not know how to bound ρ_k in the general setting of the HPE method. However, we will show that ρ_k can still be bounded in special cases of the HPE method when $\sigma = 1$. Our interest in this extreme case is due to the fact that the ADMM (see Section 7) can be viewed as a special implementation of the HPE method with $\sigma = 1$.

3 The BD-HPE framework

In this section, we introduce the BD-HPE framework for solving a special type of monotone inclusion problem consisting of the sum of a continuous monotone map and a point-to-set maximal monotone operator with a separable block-structure. Recall from Section 1 that the acronym BD-HPE stands for ‘‘block decomposition

hybrid proximal extragradient". As suggested by its name and formally proved in this section, the BD-HPE framework is a special case of the HPE method. Using this fact and the results of Subsection 2.2, global convergence rate bounds are then established for the BD-HPE framework.

Throughout this paper, we let \mathcal{X} and \mathcal{Y} denote finite dimensional inner product spaces with associated inner products both denoted by $\langle \cdot, \cdot \rangle$ and associated norms both denoted by $\| \cdot \|$. We endow the product space $\mathcal{X} \times \mathcal{Y}$ with the canonical inner product defined as

$$\langle (x, y), (x', y') \rangle = \langle x, x' \rangle + \langle y, y' \rangle, \quad \forall (x, y), (x', y') \in \mathcal{X} \times \mathcal{Y}.$$

The associated norm, also denoted by $\| \cdot \|$ for shortness, is then given

$$\|(x, y)\| = \sqrt{\|x\|^2 + \|y\|^2}, \quad \forall (x, y) \in \mathcal{X} \times \mathcal{Y}.$$

Our problem of interest in this section is the monotone inclusion problem of finding (x, y) such that

$$(0, 0) \in [F + (A \otimes B)](x, y), \tag{19}$$

or equivalently,

$$0 \in F_x(x, y) + A(x), \quad 0 \in F_y(x, y) + B(y), \tag{20}$$

where $F(x, y) = (F_x(x, y), F_y(x, y)) \in \mathcal{X} \times \mathcal{Y}$ and the following conditions are assumed:

- A.1** $A : \mathcal{X} \rightrightarrows \mathcal{X}$ and $B : \mathcal{Y} \rightrightarrows \mathcal{Y}$ are maximal monotone operators;
- A.2** $F : \text{Dom } F \subset \mathcal{X} \times \mathcal{Y} \rightarrow \mathcal{X} \times \mathcal{Y}$ is a continuous map such that $\text{Dom } F \supset \text{cl}(\text{Dom } A) \times \mathcal{Y}$;
- A.3** F is monotone on $\text{Dom } A \times \text{Dom } B$;
- A.4** there exists $L_{xy} > 0$ such that

$$\|F_x(x, y') - F_x(x, y)\| \leq L_{xy} \|y' - y\|, \quad \forall x \in \text{Dom } A, \quad \forall y, y' \in \mathcal{Y}. \tag{21}$$

We now make a few remarks about the above assumptions. First, it can be easily seen that A.1 implies that the operator $A \otimes B : \mathcal{X} \times \mathcal{Y} \rightrightarrows \mathcal{X} \times \mathcal{Y}$ defined as

$$(A \otimes B)(x, y) = A(x) \times B(y), \quad \forall (x, y) \in \mathcal{X} \times \mathcal{Y},$$

is maximal monotone. Moreover, in view of the proof of Proposition A.1 of [16], it follows that $F + (A \otimes B)$ is maximal monotone. Second, without loss of generality, we have assumed in A.2 that F is defined in $\text{cl}(\text{dom } A) \times \mathcal{Y}$ instead of a set of the form $\text{cl}(\text{dom } A) \times \Omega$ for some closed convex set $\Omega \supset \text{dom } B$ (e.g., $\Omega = \text{cl}(\text{dom } B)$). Indeed, if F were defined on the latter set only, then it would be possible to extend it to the whole set $\text{cl}(\text{dom } A) \times \mathcal{Y}$ by considering the extension $(x, y) \in \mathcal{X} \times \mathcal{Y} \rightarrow F(x, P_\Omega(y))$, which can be easily seen to satisfy A.2-A.4. Note that evaluation of this extension requires computation of a projection onto Ω . Third, assumption A.4 is needed in order to estimate how much an iterate found by the block decomposition scheme below violates the proximal point equation for (19).

The exact proximal point iteration for this problem is: given $(x_{k-1}, y_{k-1}) \in \mathcal{X} \times \mathcal{Y}$, let (x_k, y_k) be the solution of the proximal inclusion subproblem:

$$0 \in \lambda[F_x(x, y) + A(x)] + x - x_{k-1}, \tag{22}$$

$$0 \in \lambda[F_y(x, y) + B(y)] + y - y_{k-1}. \tag{23}$$

In this section, we are interested in BD methods for solving (19), where the k -th iteration consists of finding an approximate solution \tilde{x}_k of the subproblem

$$0 \in \lambda[F_x(x, y_{k-1}) + A(x)] + x - x_{k-1}, \tag{24}$$

then computing an approximate solution \tilde{y}_k of

$$0 \in \lambda[F_y(\tilde{x}_k, y) + B(y)] + y - y_{k-1}, \quad (25)$$

and finally using the pair $(\tilde{x}_k, \tilde{y}_k)$ to obtain the next iterate (x_k, y_k) . Note that if (24) and (25) are solved exactly, then the pair $(\tilde{x}_k, \tilde{y}_k)$ will satisfy the proximal point equation (22)-(23) with residual $(r_x, r_y) := (F_x(\tilde{x}_k, \tilde{y}_k) - F_x(\tilde{x}_k, y_{k-1}), 0)$, that is, the inclusion in (22)-(23) with its left hand side replaced by (r_x, r_y) . Moreover, Assumption A.4 provides a way to control this residual. Note also that A.2 ensures that the method outlined above is well-defined. Indeed, we can show that $\tilde{x}_k \in \text{cl}(\text{dom } A)$ but can not guarantee that $y_{k-1} \in \text{cl}(\text{dom } B)$, which explains the need to assume that $\text{Dom } F \supset \text{cl}(\text{dom } A) \times \mathcal{Y}$.

To formalize the method outlined in the previous paragraph, we now state the BD-HPE framework.

Block-decomposition HPE framework:

0) Let $(x_0, y_0) \in \mathcal{X} \times \mathcal{Y}$, $\sigma, \sigma_x \in [0, 1]$, and $\tilde{\sigma}_x, \sigma_y \in [0, 1]$ be given and set $k = 1$;

1) choose $\lambda_k > 0$ such that

$$\sigma_k := \left\{ \theta_{\max} \left(\begin{bmatrix} \sigma_x^2 & \lambda_k \tilde{\sigma}_x L_{xy} \\ \lambda_k \tilde{\sigma}_x L_{xy} & \sigma_y^2 + \lambda_k^2 L_{xy}^2 \end{bmatrix} \right) \right\}^{1/2} \leq \sigma; \quad (26)$$

2) compute $\tilde{x}_k, \tilde{a}_k \in \mathcal{X}$ and $\varepsilon_k^x \geq 0$ such that

$$\tilde{a}_k \in A^{\varepsilon_k^x}(\tilde{x}_k), \quad \|\lambda_k[F_x(\tilde{x}_k, y_{k-1}) + \tilde{a}_k] + \tilde{x}_k - x_{k-1}\|^2 + 2\lambda_k \varepsilon_k^x \leq \sigma_x^2 \|\tilde{x}_k - x_{k-1}\|^2; \quad (27)$$

$$\|\lambda_k[F_x(\tilde{x}_k, y_{k-1}) + \tilde{a}_k] + \tilde{x}_k - x_{k-1}\|^2 \leq \tilde{\sigma}_x^2 \|\tilde{x}_k - x_{k-1}\|^2; \quad (28)$$

3) compute $\tilde{y}_k, \tilde{b}_k \in \mathcal{Y}$ and $\varepsilon_k^y \geq 0$ such that

$$\tilde{b}_k \in B^{\varepsilon_k^y}(\tilde{y}_k), \quad \|\lambda_k[F_y(\tilde{x}_k, \tilde{y}_k) + \tilde{b}_k] + \tilde{y}_k - y_{k-1}\|^2 + 2\lambda_k \varepsilon_k^y \leq \sigma_y^2 \|\tilde{y}_k - y_{k-1}\|^2; \quad (29)$$

4) set

$$(x_k, y_k) = (x_{k-1}, y_{k-1}) - \lambda_k[F(\tilde{x}_k, \tilde{y}_k) + (\tilde{a}_k, \tilde{b}_k)], \quad (30)$$

$k \leftarrow k + 1$, and go to step 1.

end

We now make a few remarks about the BD-HPE framework. First, instead of using constants $\sigma_x, \tilde{\sigma}_x$, and σ_y in (27), (28), and (29), respectively, we could use variable factors $\sigma_{k,x} \leq \sigma_x, \tilde{\sigma}_{k,x} \leq \tilde{\sigma}_x$ and $\sigma_{k,y} \leq \sigma_y$, respectively, just like in the HPE method. However, for sake of simplicity, we will deal only with the case where these factors are constant. Second, even though we have assumed in step 0) that $\sigma_y < 1$, we observe that this condition is implied by (26) and the fact that $\sigma \leq 1$. Third, (27) implies that (28) holds for any $\tilde{\sigma}_x \in [\sigma_x, 1)$. Hence, when $\sigma_x < 1$, we can simply choose $\tilde{\sigma}_x$ to be equal to σ_x as long as (26) holds with $\tilde{\sigma}_x = \sigma_x$. Fourth, if $\sigma_x < 1$, then the assumption that $\sigma_y < 1$ implies that there always exists $\lambda_k > 0$ satisfying (26). Fifth, if $\sigma_x = 1$, then the assumption that $\sigma \leq 1$ implies that there exists $\lambda_k > 0$ satisfying (26) if, and only if, $\tilde{\sigma}_x = 0$, in which case we must have $\sigma = 1$. Sixth, there are relevant instances of the BD-HPE framework in which (27) and (28) hold with $\sigma_x = 1$ and $\tilde{\sigma}_x = 0$, and (26) holds with $\sigma = 1$, or equivalently, $\sigma_y^2 + \lambda_k^2 L_{xy}^2 \leq 1$. Finally, assumption (26) does not necessarily imply that $\sigma_x < 1$ (see the latter remark).

The following result shows that, under inequality (26), any instance of the BD-HPE framework is also an instance of the HPE method of Section 2.2 applied to the monotone inclusion (19).

Proposition 3.1. *Consider the sequences $\{(x_k, y_k)\}$, $\{(\tilde{x}_k, \tilde{y}_k)\}$, $\{(\tilde{a}_k, \tilde{b}_k)\}$, $\{\lambda_k\}$ and $\{(\varepsilon_k^x, \varepsilon_k^y)\}$ generated by*

the BD-HPE framework. Then, for every $k \in \mathbb{N}$,

$$F(\tilde{x}_k, \tilde{y}_k) + (\tilde{a}_k, \tilde{b}_k) \in [F + (A \otimes B)^{\varepsilon_k^x + \varepsilon_k^y}](\tilde{x}_k, \tilde{y}_k) \subset (F + A \otimes B)^{\varepsilon_k^x + \varepsilon_k^y}(\tilde{x}_k, \tilde{y}_k) \quad (31)$$

and

$$\left\| \lambda_k [F(\tilde{x}_k, \tilde{y}_k) + (\tilde{a}_k, \tilde{b}_k)] + (\tilde{x}_k, \tilde{y}_k) - (x_{k-1}, y_{k-1}) \right\|^2 + 2\lambda_k(\varepsilon_k^x + \varepsilon_k^y) \leq \sigma_k^2 \|(\tilde{x}_k, \tilde{y}_k) - (x_{k-1}, y_{k-1})\|^2.$$

As a consequence, any instance of the BD-HPE framework is a special case of the HPE method for the inclusion problem (19) with $\tilde{v}_k = F(\tilde{x}_k, \tilde{y}_k) + (\tilde{a}_k, \tilde{b}_k)$ and $\varepsilon_k = \varepsilon_k^x + \varepsilon_k^y$ for every $k \in \mathbb{N}$.

Proof. Using the inclusions in (27) and (29), definition (5) and the definition of ε_k , we have for every $(a, b) \in (A \otimes B)(x, y)$ that

$$\langle (\tilde{x}_k, \tilde{y}_k) - (x, y), (\tilde{a}_k, \tilde{b}_k) - (a, b) \rangle = \langle \tilde{x}_k - x, \tilde{a}_k - a \rangle + \langle \tilde{y}_k - y, \tilde{b}_k - b \rangle \geq -(\varepsilon_k^x + \varepsilon_k^y),$$

which shows that $(\tilde{a}_k, \tilde{b}_k) \in (A \otimes B)^{\varepsilon_k^x + \varepsilon_k^y}(\tilde{x}_k, \tilde{y}_k)$, and hence that (31) holds, in view of statements b) and c) of Proposition 2.1. Let

$$r_k^x := \lambda_k(F_x(\tilde{x}_k, y_{k-1}) + \tilde{a}_k) + \tilde{x}_k - x_{k-1}, \quad r_k^y := \lambda_k(F_y(\tilde{x}_k, \tilde{y}_k) + \tilde{b}_k) + \tilde{y}_k - y_{k-1} \quad (32)$$

Then,

$$\lambda_k[F(\tilde{x}_k, \tilde{y}_k) + (\tilde{a}_k, \tilde{b}_k)] + (\tilde{x}_k, \tilde{y}_k) - (x_{k-1}, y_{k-1}) = (r_k^x + \lambda_k(F_x(\tilde{x}_k, \tilde{y}_k) - F_x(\tilde{x}_k, y_{k-1})), r_k^y),$$

which, together with (21), (26) and (32), and the inequalities in (27), (28) and (29), imply

$$\begin{aligned} & \left\| \lambda_k [F(\tilde{x}_k, \tilde{y}_k) + (\tilde{a}_k, \tilde{b}_k)] + (\tilde{x}_k, \tilde{y}_k) - (x_{k-1}, y_{k-1}) \right\|^2 + 2\lambda_k(\varepsilon_k^x + \varepsilon_k^y) \\ & \leq \|r_k^x + \lambda_k(F_x(\tilde{x}_k, \tilde{y}_k) - F_x(\tilde{x}_k, y_{k-1}))\|^2 + \|r_k^y\|^2 + 2\lambda_k(\varepsilon_k^x + \varepsilon_k^y) \\ & \leq (\|r_k^x\| + \lambda_k\|F_x(\tilde{x}_k, \tilde{y}_k) - F_x(\tilde{x}_k, y_{k-1})\|)^2 + \|r_k^y\|^2 + 2\lambda_k(\varepsilon_k^x + \varepsilon_k^y) \\ & \leq (\|r_k^x\| + \lambda_k L_{xy} \|\tilde{y}_k - y_{k-1}\|)^2 + \|r_k^y\|^2 + 2\lambda_k(\varepsilon_k^x + \varepsilon_k^y) \\ & \leq \lambda_k^2 L_{xy}^2 \|\tilde{y}_k - y_{k-1}\|^2 + 2\lambda_k L_{xy} \|r_k^x\| \|\tilde{y}_k - y_{k-1}\| + (\|r_k^x\|^2 + 2\lambda_k \varepsilon_k^x) + (\|r_k^y\|^2 + 2\lambda_k \varepsilon_k^y) \\ & \leq \lambda_k^2 L_{xy}^2 \|\tilde{y}_k - y_{k-1}\|^2 + 2\lambda_k \tilde{\sigma}_x L_{xy} \|\tilde{x}_k - x_{k-1}\| \|\tilde{y}_k - y_{k-1}\| + \sigma_x^2 \|\tilde{x}_k - x_{k-1}\|^2 + \sigma_y^2 \|\tilde{y}_k - y_{k-1}\|^2 \\ & \leq \sigma_k^2 (\|\tilde{x}_k - x_{k-1}\|^2 + \|\tilde{y}_k - y_{k-1}\|^2) = \sigma_k^2 \|(\tilde{x}_k, \tilde{y}_k) - (x_{k-1}, y_{k-1})\|^2. \quad \square \end{aligned}$$

We now state two iteration-complexity results for the BD-HPE framework which are direct consequences of Proposition 3.1 and Theorems 2.5 and 2.6. The first (pointwise) one is about the behaviour of the sequence $\{(\tilde{x}_k, \tilde{y}_k)\}$ and the second (ergodic) one is in regards to an ergodic sequence associated with $\{(\tilde{x}_k, \tilde{y}_k)\}$.

Theorem 3.2. *Assume that $\sigma < 1$ and consider the sequences $\{(\tilde{x}_k, \tilde{y}_k)\}$, $\{(\tilde{a}_k, \tilde{b}_k)\}$, $\{\lambda_k\}$ and $\{(\varepsilon_k^x, \varepsilon_k^y)\}$ generated by the BD-HPE framework and let d_0 denote the distance of the initial point $(x_0, y_0) \in \mathcal{X} \times \mathcal{Y}$ to the solution set of (19). Then, for every $k \in \mathbb{N}$,*

$$(\tilde{a}_k, \tilde{b}_k) \in A^{\varepsilon_k^x}(\tilde{x}_k) \times B^{\varepsilon_k^y}(\tilde{y}_k),$$

and there exists $i \leq k$ such that

$$\left\| F(\tilde{x}_i, \tilde{y}_i) + (\tilde{a}_i, \tilde{b}_i) \right\| \leq d_0 \sqrt{\frac{1 + \sigma}{1 - \sigma} \left(\frac{1}{\sum_{j=1}^k \lambda_j^2} \right)}, \quad \varepsilon_i^x + \varepsilon_i^y \leq \frac{\sigma^2 d_0^2 \lambda_i}{2(1 - \sigma^2) \sum_{j=1}^k \lambda_j^2}.$$

Proof. This result follows immediately from Proposition 3.1 and Theorem 2.5. \square

Theorem 3.3. Consider the sequences $\{(\tilde{x}_k, \tilde{y}_k)\}$, $\{(\tilde{a}_k, \tilde{b}_k)\}$ and $\{(\varepsilon_k^x, \varepsilon_k^y)\}$ generated by the BD-HPE framework and define for every $k \in \mathbb{N}$:

$$(\tilde{x}_k^a, \tilde{y}_k^a) = \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i (\tilde{x}_i, \tilde{y}_i), \quad (\tilde{a}_k^a, \tilde{b}_k^a) = \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i (\tilde{a}_i, \tilde{b}_i), \quad \tilde{F}_k^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i F(\tilde{x}_i, \tilde{y}_i), \quad (33)$$

and

$$\varepsilon_{k,F}^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i \langle (\tilde{x}_i, \tilde{y}_i) - (\tilde{x}_k^a, \tilde{y}_k^a), F(\tilde{x}_i, \tilde{y}_i) \rangle \geq 0, \quad (34)$$

$$\varepsilon_{k,A}^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i (\varepsilon_i^x + \langle \tilde{x}_i - \tilde{x}_k^a, \tilde{a}_i \rangle) \geq 0, \quad (35)$$

$$\varepsilon_{k,B}^a := \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i \left(\varepsilon_i^y + \langle \tilde{y}_i - \tilde{y}_k^a, \tilde{b}_i \rangle \right) \geq 0, \quad (36)$$

where $\Lambda_k := \sum_{i=1}^k \lambda_i$. Let d_0 denote the distance of the initial point $(x_0, y_0) \in \mathcal{X} \times \mathcal{Y}$ to the solution set of (19). Then, for every $k \in \mathbb{N}$,

$$\tilde{F}_k^a \in F^{\varepsilon_{k,F}^a}(\tilde{x}_k^a, \tilde{y}_k^a), \quad (\tilde{a}_k^a, \tilde{b}_k^a) \in A^{\varepsilon_{k,A}^a}(\tilde{x}_k^a) \times B^{\varepsilon_{k,B}^a}(\tilde{y}_k^a), \quad (37)$$

and

$$\left\| \tilde{F}_k^a + (\tilde{a}_k^a, \tilde{b}_k^a) \right\| \leq \frac{2d_0}{\Lambda_k}, \quad \varepsilon_{k,F}^a + \varepsilon_{k,A}^a + \varepsilon_{k,B}^a \leq \frac{2d_0^2}{\Lambda_k} (1 + \eta_k), \quad (38)$$

where

$$\eta_k := \frac{2}{1 - \sigma_{xy}} \left(1 + \frac{1}{(1 - \sigma_y)^2} \right)^{1/2} \sqrt{\tilde{\sigma}_x^2 + \sigma_y^2 + \lambda_k^2 L_{xy}^2} \leq \frac{2\sqrt{2}\sigma}{1 - \sigma_{xy}} \left(1 + \frac{1}{(1 - \sigma_y)^2} \right)^{1/2} \quad (39)$$

and $\sigma_{xy} := \max\{\tilde{\sigma}_x, \sigma_y\}$. Moreover, if F is affine, then $\tilde{F}_k^a = F(\tilde{x}_k^a, \tilde{y}_k^a)$. Also, if A (resp., B) is affine and $\varepsilon_k^x = 0$ (resp., $\varepsilon_k^y = 0$) for every $k \in \mathbb{N}$, then $\tilde{a}_k^a \in A(\tilde{x}_k^a)$ (resp., $\tilde{b}_k^a \in B(\tilde{y}_k^a)$).

Proof. First, note that (37) follows immediately from the definitions in (33), (34), (35) and (36), the inclusions in (27) and (29), and Theorem 2.2. In view of Proposition 3.1, any instance of the BD-HPE framework is a special case of the HPE method with $\tilde{v}_k = F(\tilde{x}_k, \tilde{y}_k) + (\tilde{a}_k, \tilde{b}_k)$ and $\varepsilon_k = \varepsilon_k^x + \varepsilon_k^y$ for every $k \in \mathbb{N}$. Since, in this case, the quantities \tilde{z}_k^a , \tilde{v}_k^a and ε_k^a defined in Theorem 2.6 are equal to $(\tilde{x}_k^a, \tilde{y}_k^a)$, $F_k^a + (\tilde{a}_k^a, \tilde{b}_k^a)$ and $\varepsilon_{x,F}^a + \varepsilon_{x,F}^a + \varepsilon_{x,F}^a$, respectively, it follows from the conclusions of this theorem that the first inequality in (38) holds and

$$\varepsilon_{x,F}^a + \varepsilon_{x,F}^a + \varepsilon_{x,F}^a \leq \frac{2d_0^2}{\Lambda_k} (1 + \rho_k), \quad (40)$$

where

$$\rho_k := \frac{1}{d_0} \max_{i=1, \dots, k} \|(\tilde{x}_i, \tilde{y}_i) - (x_i, y_i)\|. \quad (41)$$

Noting the definition of η_k in (39), we now claim that $\rho_k \leq \eta_k$, which, together with (40), clearly implies the second inequality in (38). Indeed, let (x^*, y^*) be a solution of (19) such that $\|(x_0, y_0) - (x^*, y^*)\| = d_0$. Due to Proposition 2.4, we know that

$$\begin{aligned} \|(x_k, y_k) - (x_{k-1}, y_{k-1})\| &\leq \|(x_k, y_k) - (x^*, y^*)\| + \|(x_{k-1}, y_{k-1}) - (x^*, y^*)\| \\ &\leq \|(x_0, y_0) - (x^*, y^*)\| + \|(x_0, y_0) - (x^*, y^*)\| = 2d_0. \end{aligned} \quad (42)$$

It follows from (29) and (30) that

$$\|\tilde{y}_k - y_{k-1}\| \leq \|\tilde{y}_k - y_k\| + \|y_k - y_{k-1}\| \leq \sigma_y \|\tilde{y}_k - y_{k-1}\| + \|y_k - y_{k-1}\|,$$

and hence that

$$\|\tilde{y}_k - y_{k-1}\| \leq \frac{\|y_k - y_{k-1}\|}{1 - \sigma_y}, \quad \|\tilde{y}_k - y_k\| \leq \sigma_y \|\tilde{y}_k - y_{k-1}\| \leq \frac{\sigma_y \|y_k - y_{k-1}\|}{1 - \sigma_y}. \quad (43)$$

Also, it follows from (21), (28) and (30) that

$$\begin{aligned} \|\tilde{x}_k - x_k\| - \lambda_k L_{xy} \|\tilde{y}_k - y_{k-1}\| &\leq \|\tilde{x}_k - x_k + \lambda_k [F_x(\tilde{x}_k, y_{k-1}) - F_x(\tilde{x}_k, \tilde{y}_k)]\| \\ &\leq \tilde{\sigma}_x \|\tilde{x}_k - x_{k-1}\| \leq \tilde{\sigma}_x (\|\tilde{x}_k - x_k\| + \|x_k - x_{k-1}\|), \end{aligned}$$

and hence that

$$\|\tilde{x}_k - x_k\| \leq \frac{\tilde{\sigma}_x \|x_k - x_{k-1}\| + \lambda_k L_{xy} \|\tilde{y}_k - y_{k-1}\|}{1 - \tilde{\sigma}_x}. \quad (44)$$

Adding the second inequality in (43) to inequality (44) and using (42), the first inequality in (43) and the definition of $\sigma_{x,y}$ in the statement of the theorem, we conclude that

$$\begin{aligned} \|(\tilde{x}_k, \tilde{y}_k) - (x_k, y_k)\| &\leq \|\tilde{x}_k - x_k\| + \|\tilde{y}_k - y_k\| \\ &\leq \frac{1}{1 - \sigma_{xy}} (\tilde{\sigma}_x \|x_k - x_{k-1}\| + \sigma_y \|y_k - y_{k-1}\| + \lambda_k L_{xy} \|\tilde{y}_k - y_{k-1}\|) \\ &\leq \frac{1}{1 - \sigma_{xy}} \sqrt{\tilde{\sigma}_x^2 + \sigma_y^2 + \lambda_k^2 L_{xy}^2} \sqrt{\|x_k - x_{k-1}\|^2 + \|y_k - y_{k-1}\|^2 + \|\tilde{y}_k - y_{k-1}\|^2} \\ &\leq \frac{1}{1 - \sigma_{xy}} \sqrt{\tilde{\sigma}_x^2 + \sigma_y^2 + \lambda_k^2 L_{xy}^2} \sqrt{4d_0^2 + (1 - \sigma_y)^{-2} \|y_k - y_{k-1}\|^2} \\ &\leq \frac{2d_0}{1 - \sigma_{xy}} \left(1 + \frac{1}{(1 - \sigma_y)^2}\right)^{1/2} \sqrt{\tilde{\sigma}_x^2 + \sigma_y^2 + \lambda_k^2 L_{xy}^2}. \end{aligned}$$

The last estimate together with (39) and (41) clearly imply our claim that $\rho_k \leq \eta_k$. Finally, note that the second inequality in (39) follows from (26) and the assumption that $\tilde{\sigma}_x \leq \sigma_x$, and that the last assertion of the theorem follows from implication (4) about an affine map T . \square

We observe that, if we had assumed in Theorem 3.3 that $\sigma < 1$ (or $\text{Dom } A \times \text{Dom } B$ is bounded), then its proof would be much simpler since in this case we could have used the last part of Theorem 2.6 to establish boundedness of $\{\rho_k\}$. However, as observed earlier, our interest in the case where $\sigma = 1$ is due to the fact that the ADMM (see Section 7) can be viewed as an instance of the HPE method with $\sigma = 1$. To handle the case $\sigma = 1$, the proof of Theorem 3.3 uses inequality (16), which depends on the quantity ρ_k . As shown in this proof, all that is required for showing boundedness of $\{\rho_k\}$ is the assumption that $\sigma \leq 1$ and $\max\{\tilde{\sigma}_x, \sigma_y\} < 1$.

Theorems 3.2 (resp., Theorem 3.3) requires condition (26) to hold for some $\sigma < 1$ (resp., $\sigma \leq 1$), which in turn implies that $\sigma_y < 1$. We have seen that any these assumptions implies that the BD-HPE framework is a special case of the HPE method (see Proposition 3.1). We conjecture whether iteration-complexity bounds can be established for the BD-HPE framework, or some subclass of it, under some weaker assumption, i.e., one which either allows condition (26) to hold for some $\sigma > 1$, or σ_y to be equal to 1.

4 Approximate solutions of the proximal subproblems

In this section, we describe some specific procedures for finding approximate solutions $(\tilde{x}_k, \tilde{a}_k, \varepsilon_k^x)$ and $(\tilde{y}_k, \tilde{b}_k, \varepsilon_k^y)$ of (24) and (25) according to steps 2 and 3, respectively, of the BD-HPE framework. We should emphasize that such solutions can be found by other procedures which are not discussed below.

The problem of finding approximate solutions as above can be cast in the following general form. Throughout this section, we assume that

B.1) $C : \mathcal{X} \rightrightarrows \mathcal{X}$ is a maximal monotone operator;

B.2) $G : \text{Dom } G \subset \mathcal{X} \rightarrow \mathcal{X}$ is a continuous map which is monotone on $\text{cl}(\text{Dom } C) \subset \text{Dom } G$.

Given $x \in \mathcal{X}$ and $\lambda > 0$ together with tolerances $\sigma, \tilde{\sigma} \geq 0$, our goal is to describe specific procedures for computing a triple $(\tilde{x}, \tilde{c}, \varepsilon) \in \mathcal{X} \times \mathcal{X} \times \mathbb{R}_+$ such that

$$\tilde{c} \in C^\varepsilon(\tilde{x}), \quad \|\lambda(G(\tilde{x}) + \tilde{c}) + \tilde{x} - x\|^2 + 2\lambda\varepsilon \leq \sigma^2 \|\tilde{x} - x\|^2, \quad (45)$$

$$\|\lambda(G(\tilde{x}) + \tilde{c}) + \tilde{x} - x\| \leq \tilde{\sigma} \|\tilde{x} - x\|. \quad (46)$$

We note that conditions B.1 and B.2 imply that $G + C$ is a maximal monotone operator (see the proof of Proposition A.1 of [16]). This implies that, for any $\lambda > 0$, the resolvent of $G + C$, namely the map $[I + \lambda(G + C)]^{-1}$ is a single-valued map defined over the whole \mathcal{X} . The following simple result shows that when the resolvent of $G + C$ is computable, (45) and (46) can be solved exactly.

Proposition 4.1. *For any $x \in \mathcal{X}$ and $\lambda > 0$, the triple $(\tilde{x}, \tilde{c}, \varepsilon)$ defined as*

$$\tilde{x} := [\lambda(G + C) + I]^{-1}(x), \quad \tilde{c} := \frac{1}{\lambda}(x - \tilde{x}) - G(\tilde{x}), \quad \varepsilon = 0 \quad (47)$$

satisfies (45) and (46) for any $\sigma \geq 0$ and $\tilde{\sigma} \geq 0$.

Proof. Using the three identities in (47), we easily see that $\tilde{c} \in C(\tilde{x})$ and

$$\|\lambda(G(\tilde{x}) + \tilde{c}) + \tilde{x} - x\|^2 + 2\lambda\varepsilon = 0 \leq \sigma^2 \|\tilde{x} - x\|^2. \quad \square$$

Now we deal with the case in which C is the sum of a differentiable convex function with Lipschitz continuous gradient and a maximal monotone operator T for which the resolvent of $G + T$ is easy to compute. Note that this case describes a meaningful situation in which it is possible to compute a triple $(\tilde{x}, \tilde{c}, \varepsilon)$ for which the smallest provably σ satisfying (45) is positive while the smallest $\tilde{\sigma}$ satisfying (46) is zero.

Proposition 4.2. *Assume that $C = \partial f + T$, where $T : \mathcal{X} \rightrightarrows \mathcal{X}$ is maximal monotone and $f : \mathcal{X} \rightarrow (-\infty, \infty]$ is a proper closed convex function such that f is differentiable on $\text{cl}(\text{dom } T) \subset \text{int}(\text{dom } f)$, ∇f is L -Lipschitz continuous on $\text{cl}(\text{Dom } T)$. Then, for any $x \in \text{Dom } T$ and $\lambda > 0$, the triple $(\tilde{x}, \tilde{c}, \varepsilon)$ defined as*

$$\tilde{x} = [I + \lambda(G + T)]^{-1}(x - \lambda \nabla f(x)), \quad \tilde{c} = \frac{1}{\lambda}(x - \tilde{x}) - G(\tilde{x}), \quad \varepsilon = \frac{L}{2} \|\tilde{x} - x\|^2 \quad (48)$$

satisfies (45) and (46) for any $\sigma \geq \sqrt{\lambda L}$ and $\tilde{\sigma} \geq 0$. Moreover, $\tilde{c} \in (\partial_\varepsilon f + T)(\tilde{x})$.

Proof. First observe that the last two identities in (48) imply that

$$\|\lambda(G(\tilde{x}) + \tilde{c}) + \tilde{x} - x\|^2 + 2\lambda\varepsilon = 2\lambda\varepsilon \leq 2\lambda \left(\frac{L}{2} \|\tilde{x} - x\|^2 \right) = \lambda L \|\tilde{x} - x\|^2,$$

and hence that $(\tilde{x}, \tilde{c}, \varepsilon)$ satisfies (45) and (46) for any $\sigma \geq \sqrt{\lambda L}$ and $\tilde{\sigma} \geq 0$. It remains to show that $\tilde{c} \in C^\varepsilon(\tilde{x})$. Using the definition of \tilde{x} , we have

$$\frac{1}{\lambda}(x - \tilde{x}) - \nabla f(x) \in (G + T)(\tilde{x}),$$

and hence, $\tilde{c} \in \nabla f(x) + T(\tilde{x})$, due to the definition of \tilde{c} . We now claim that $\nabla f(x) \in \partial_\varepsilon f(\tilde{x})$, from which we conclude that

$$\tilde{c} \in (\partial_\varepsilon f + T)(\tilde{x}) \subset [(\partial f)^\varepsilon + T](\tilde{x}) \subset (\partial f + T)^\varepsilon(\tilde{x}) = C^\varepsilon(\tilde{x}),$$

where the second and third inclusions follow from Proposition 2.3(a) and Proposition 2.1, and the equality follows from the definition of C . To prove the claim, note that Proposition 2.3(c) with $v = \nabla f(x)$ implies that $\nabla f(x) \in \partial_{\varepsilon'} f(\tilde{x})$, where

$$\varepsilon' := f(\tilde{x}) - f(x) - \langle \nabla f(x), \tilde{x} - x \rangle \leq \frac{L}{2} \|\tilde{x} - x\|^2 =: \varepsilon,$$

where the inequality is due to the fact that ∇f is L -Lipschitz continuous on $\text{cl}(\text{Dom } T) \supset \text{Dom } T \ni \tilde{x}, x$, and $\text{cl}(\text{Dom } T)$ is convex. \square

In contrast to Proposition 4.2, the next result shows how one can obtain an approximate solution $(\tilde{x}, \tilde{c}, \varepsilon)$ of (45) for which $\varepsilon = 0$. A special case of it (in which $\Omega = \mathbb{R}^n$) forms the basis of Tseng's modified forward-backward splitting algorithm (see [30]).

Proposition 4.3. *Assume that $G : \mathcal{X} \rightarrow \mathcal{X}$ is L -Lipschitz continuous on a closed convex set Ω such that $\text{Dom } C \subset \Omega \subset \text{Dom } G$. Then, for any $x \in \mathcal{X}$ and $\lambda > 0$, the triple $(\tilde{x}, \tilde{c}, \varepsilon)$ defined as*

$$\tilde{x} = (I + \lambda C)^{-1}(x - \lambda G(P_\Omega(x))), \quad \tilde{c} = \frac{1}{\lambda}(x - \tilde{x}) - G(P_\Omega(x)), \quad \varepsilon = 0 \quad (49)$$

satisfies (45) and (46) for any $\sigma, \tilde{\sigma} \geq \lambda L$.

Proof. First note that the first two identities in (49) imply that $\tilde{x} \in \text{Dom } C \subset \Omega$ and $\tilde{c} \in C(\tilde{x})$, and hence that $\tilde{c} \in C^\varepsilon(\tilde{x})$ in view of the definition of ε and Proposition 2.1(c). Also, relation (49), the inclusion $\tilde{x} \in \Omega$ and the assumption that G is L -Lipschitz continuous on Ω imply that

$$\begin{aligned} \|\lambda(G(\tilde{x}) + \tilde{c}) + \tilde{x} - x\|^2 + 2\lambda\varepsilon &= \|\lambda(G(\tilde{x}) + \tilde{c}) + \tilde{x} - x\|^2 = \lambda^2\|G(\tilde{x}) - G(P_\Omega(x))\|^2 \\ &= \lambda^2\|G(P_\Omega(\tilde{x})) - G(P_\Omega(x))\|^2 \leq \lambda^2 L^2\|P_\Omega(\tilde{x}) - P_\Omega(x)\|^2 \leq (\lambda L)^2\|\tilde{x} - x\|^2, \end{aligned}$$

where the last inequality follows from the fact that P_Ω is a nonexpansive map. We have thus shown that $(\tilde{x}, \tilde{c}, \varepsilon)$ satisfies (45) and (46) for any $\sigma, \tilde{\sigma} \geq \lambda L$. \square

Note that the above formula for \tilde{x} is in terms of the resolvent $(I + \lambda C)^{-1}$ of C , which must be easily computable so that \tilde{x} can be obtained. Observe also that for the case where $C = N_X$ for some closed convex set $X \subset \mathcal{X}$, we have $(I + \lambda N_X)^{-1} = P_X$ and the above expression for \tilde{x} reduces to $\tilde{x} = P_X(x - \lambda G(P_\Omega(x)))$.

The next result describes a way of computing an approximate solution $(\tilde{x}, \tilde{c}, \varepsilon)$ of (45) which forms the basis of Korpelevich's method (see [13]) for solving monotone variational inequalities and its generalized version (see for example [19, 16]) for solving monotone hemi-variational inequalities. In contrast to Proposition 4.3, it assumes that C is a subdifferential and it needs to evaluate two resolvents of G in order to compute $(\tilde{x}, \tilde{c}, \varepsilon)$.

Proposition 4.4. *Assume that $C = \partial g$, where $g : \mathcal{X} \rightarrow (-\infty, \infty]$ is a closed proper convex function and G is L -Lipschitz continuous on $\text{dom } g$. Then, for any $x \in \text{dom } g$ and $\lambda > 0$, the triple $(\tilde{x}, \tilde{c}, \varepsilon)$ defined as*

$$\tilde{x} = (I + \lambda \partial g)^{-1}(x - \lambda G(x)), \quad \tilde{c} = \frac{1}{\lambda}[x - x^+] - G(\tilde{x}), \quad \varepsilon := g(\tilde{x}) - [g(x^+) + \langle \tilde{x} - x^+, \tilde{c} \rangle], \quad (50)$$

where

$$x^+ := (I + \partial g)^{-1}(x - \lambda G(\tilde{x})), \quad (51)$$

satisfies (45) and (46) for any $\sigma, \tilde{\sigma} \geq \lambda L$.

Proof. First observe that ε is well-defined since $\tilde{x}, x^+ \in \text{dom } g$, in view of their definition in (50) and (51), respectively. We first prove that $\tilde{c} \in C^\varepsilon(\tilde{x})$. Indeed, the definition of \tilde{c} and x^+ in (50) and (51), respectively, imply that $\tilde{c} \in \partial g(x^+)$. Hence, it follows from Proposition 2.3(c) and the definition of ε in (50) that $\tilde{c} \in \partial_\varepsilon g(\tilde{x})$, and hence that $\tilde{c} \in (\partial g)^\varepsilon(\tilde{x}) = C^\varepsilon(\tilde{x})$, in view of Proposition 2.3(a).

To show that the inequality in (45) holds for any $\sigma \geq \lambda L$, define

$$p = \frac{1}{\lambda}[x - \lambda G(x) - \tilde{x}]. \quad (52)$$

Using the definition of \tilde{x} and p in (50) and (52), respectively, we conclude that $p \in \partial g(\tilde{x})$. This fact and the last identity in (50) then imply that

$$\varepsilon = g(\tilde{x}) - g(x^+) - \langle \tilde{c}, \tilde{x} - x^+ \rangle = -[g(x^+) - g(\tilde{x}) - \langle p, x^+ - \tilde{x} \rangle] + \langle p - \tilde{c}, \tilde{x} - x^+ \rangle \leq \langle p - \tilde{c}, \tilde{x} - x^+ \rangle.$$

This, together with the second identity in (50), then imply that

$$\begin{aligned} \|\lambda(G(\tilde{x}) + \tilde{c}) + \tilde{x} - x\|^2 + 2\lambda\varepsilon &= \|\tilde{x} - x^+\|^2 + 2\lambda\varepsilon \leq \|\tilde{x} - x^+\|^2 + 2\lambda\langle p - \tilde{c}, \tilde{x} - x^+ \rangle \\ &= \|\lambda(p - \tilde{c}) + \tilde{x} - x^+\|^2 - \lambda^2\|p - \tilde{c}\|^2 \leq \|\lambda(p - \tilde{c}) + \tilde{x} - x^+\|^2 \\ &= \|\lambda(G(x) - G(\tilde{x}))\|^2 \leq (\lambda L\|x - \tilde{x}\|)^2, \end{aligned}$$

where the last equality follows from (52) and the second identity in (50), and the last inequality is due to the assumption that G is L -Lipschitz continuous on $\text{dom } g \supset \{x, \tilde{x}\}$. We have thus shown that (45) and (46) hold for any $\sigma, \tilde{\sigma} \geq \lambda L$. \square

5 Specific examples of BD-HPE methods

The goal of this section is to illustrate how the different procedures discussed in Section 4 for constructing triples $(\tilde{x}_k, \tilde{a}_k, \varepsilon_k^x)$ and $(\tilde{y}_k, \tilde{b}_k, \varepsilon_k^y)$ satisfying (27), (28), and (29), respectively, can be used to obtain specific instances of the BD-HPE framework presented in Section 3. This section is divided into three subsections. In the first one, we discuss a specific instance of the the BD-HPE framework in which (24) and (25) are both solved exactly (see Proposition 4.1). In the second subsection, we give another instance of BD-HPE framework in which these two proximal subproblems are approximately solved by means of Tseng's scheme presented in Proposition 4.3. In the third subsection we study a BD method for a large class of linearly constrained convex optimization problems, which includes cone programs whose objective functions converge to infinity as the relative boundaries of their domain are approached.

5.1 Exact BD-HPE method

In this subsection, we consider a special case of the general BD-HPE framework where the subproblems (24) and (25) are solved exactly and specialize the iteration-complexity bounds of Theorems 3.2 and 3.3 to the current setting.

In this subsection, we assume that we know how to solve the proximal subproblems (24) and (25) exactly. More precisely, we consider the following special case of the BD-HPE framework.

Exact BD-HPE method:

- 0) Let $(x_0, \tilde{y}_0) \in \mathcal{X} \times \mathcal{Y}$ and $\sigma \in (0, 1]$ be given and set $\lambda = \sigma/L_{xy}$ and $k = 1$;
- 1) compute $(\tilde{x}_k, \tilde{y}_k) \in \mathcal{X} \times \mathcal{Y}$ as

$$\tilde{x}_k = [I + \lambda(F_x(\cdot, \tilde{y}_{k-1}) + A)]^{-1}(x_{k-1}), \quad \tilde{y}_k = [I + \lambda(F_y(\tilde{x}_k, \cdot) + B)]^{-1}(\tilde{y}_{k-1}). \quad (53)$$

- 2) set $x_k = \tilde{x}_k - \lambda[F_x(\tilde{x}_k, \tilde{y}_k) - F_x(\tilde{x}_k, \tilde{y}_{k-1})]$ and $k \leftarrow k + 1$, and go to step 1;

end

The following result shows that the above algorithm is indeed a special case of the BD-HPE framework in which subproblems (24) and (25) are solved exactly (see Proposition 4.1).

Lemma 5.1. *Consider the sequences $\{(x_k, y_k)\}$ and $\{(\tilde{x}_k, \tilde{y}_k)\}$ generated by the exact BD-HPE method, and for each $k \in \mathbb{N}$, define $\varepsilon_k^x = \varepsilon_k^y = 0$, $\lambda_k = \lambda$,*

$$\tilde{a}_k = \frac{1}{\lambda}(x_{k-1} - \tilde{x}_k) - F_x(\tilde{x}_k, \tilde{y}_{k-1}), \quad \tilde{b}_k = \frac{1}{\lambda}(\tilde{y}_{k-1} - \tilde{y}_k) - F_y(\tilde{x}_k, \tilde{y}_k), \quad y_{k-1} = \tilde{y}_{k-1}. \quad (54)$$

Then, for every $k \in \mathbb{N}$, (26), (27), (28), (29) and (30) hold with $\sigma_x = \tilde{\sigma}_x = \sigma_y = 0$. As a consequence, the exact BD-HPE method is a special instance of the BD-HPE framework in which $\sigma_x = \tilde{\sigma}_x = \sigma_y = 0$.

Proof. The definition of $\{\lambda_k\}$ clearly implies that (26) holds with $\sigma_x = \tilde{\sigma}_x = \sigma_y = 0$. Using the definition of ε_k^x , ε_k^y and λ_k , and relations (53) and (54), and applying Proposition 4.1 twice to the pairs $(G, C) = (F(\cdot, \tilde{y}_{k-1}), A)$ and $(G, C) = (F(\tilde{x}_k, \cdot), B)$, we conclude that (27), (28) and (29) hold with $\sigma_x = \tilde{\sigma}_x = \sigma_y = 0$. Moreover, (30) follows from (54) and the update rule of x_k in step 2 of the exact BD-HPE method. \square

The following result, which is an immediate consequence of the previous result and Theorem 3.3, establishes the iteration-complexity of the exact BD-HPE method.

Theorem 5.2. *Consider the sequences $\{x_k\}$ and $\{(\tilde{x}_k, \tilde{y}_k)\}$ generated by the exact BD-HPE method, and define the sequence $\{(\tilde{a}_k, \tilde{b}_k)\}$ according to (54). Moreover, for each $k \in \mathbb{N}$, define:*

$$(\tilde{x}_k^a, \tilde{y}_k^a) = \frac{1}{k} \sum_{i=1}^k (\tilde{x}_i, \tilde{y}_i), \quad (\tilde{a}_k^a, \tilde{b}_k^a) = \frac{1}{k} \sum_{i=1}^k (\tilde{a}_i, \tilde{b}_i), \quad \tilde{F}_k^a := \frac{1}{k} \sum_{i=1}^k F(\tilde{x}_i, \tilde{y}_i),$$

and

$$\varepsilon_{k,F}^a := \frac{1}{k} \sum_{i=1}^k \langle (\tilde{x}_i, \tilde{y}_i) - (\tilde{x}_k^a, \tilde{y}_k^a), F(\tilde{x}_i, \tilde{y}_i) \rangle \geq 0, \\ \varepsilon_{k,A}^a := \frac{1}{k} \sum_{i=1}^k \langle \tilde{x}_i - \tilde{x}_k^a, \tilde{a}_i \rangle \geq 0, \quad \varepsilon_{k,B}^a := \frac{1}{k} \sum_{i=1}^k \langle \tilde{y}_i - \tilde{y}_k^a, \tilde{b}_i \rangle \geq 0.$$

Let d_0 denote the distance of the initial point $(x_0, \tilde{y}_0) \in \mathcal{X} \times \mathcal{Y}$ to the solution set of (19). Then, for every $k \in \mathbb{N}$, the following statements hold:

a) $(\tilde{a}_k, \tilde{b}_k) \in A(\tilde{x}_k) \times B(\tilde{y}_k)$, and if $\sigma < 1$, there exists $i \leq k$ such that

$$\|F(\tilde{x}_i, \tilde{y}_i) + (\tilde{a}_i, \tilde{b}_i)\| \leq \frac{L_{xy}d_0}{\sigma\sqrt{k}} \sqrt{\frac{1+\sigma}{1-\sigma}};$$

b) we have

$$\tilde{F}_k^a \in F^{\varepsilon_{k,F}^a}(\tilde{x}_k^a, \tilde{y}_k^a), \quad (\tilde{a}_k^a, \tilde{b}_k^a) \in A^{\varepsilon_{k,A}^a}(\tilde{x}_k^a) \times B^{\varepsilon_{k,B}^a}(\tilde{y}_k^a),$$

and

$$\|\tilde{F}_k^a + (\tilde{a}_k^a, \tilde{b}_k^a)\| \leq \frac{2L_{xy}d_0}{k\sigma}, \quad \varepsilon_{k,F}^a + \varepsilon_{k,A}^a + \varepsilon_{k,B}^a \leq \frac{2L_{xy}d_0^2}{k\sigma} (1 + 2\sqrt{2}\sigma).$$

Also, if F is affine, then $\tilde{F}_k^a = F(\tilde{x}_k^a, \tilde{y}_k^a)$. In addition, if A (resp., B) is affine, then $\tilde{a}_k^a \in A(\tilde{x}_k^a)$ (resp., $\tilde{b}_k^a \in B(\tilde{y}_k^a)$).

Proof. This result follows immediately from Lemma 5.1 and Theorems 3.2 and 3.3 by specializing the latter two results to the case where $\sigma_x = \tilde{\sigma}_x = \sigma_y = 0$, $\lambda_k = \lambda := \sigma/L_{xy}$ and $\varepsilon_k^x = \varepsilon_k^y = 0$ for every $k \in \mathbb{N}$. \square

5.2 An inexact BD-HPE method based on Tseng's procedure

In this subsection, we describe an inexact BD-HPE method based on Tseng's procedure described in Proposition 4.3.

We start by describing the general assumptions of this subsection. In addition to conditions A.1) to A.4) of Subsection 3, we also impose the following condition:

A.5 there exist scalars $L_{xx}, L_{yy} \geq 0$ and a closed convex set Ω_x such that $\text{Dom } A \subset \Omega_x$, $\Omega_x \times \mathcal{Y} \subset \text{Dom } F$, and:

- $F_x(\cdot, y)$ is L_{xx} -Lipschitz continuous on Ω_x for every $y \in \mathcal{Y}$;
- $F_y(x, \cdot)$ is L_{yy} -Lipschitz continuous on \mathcal{Y} for every $x \in \Omega_x$.

Tseng's based inexact BD-HPE method:

0) Let $(x_0, y_0) \in \mathcal{X} \times \mathcal{Y}$, $\sigma \in (0, 1]$ and $\lambda \in (0, \sigma/\tilde{L}]$ be given, where

$$\tilde{L} := \left\{ \theta_{\max} \left(\begin{bmatrix} L_{xx}^2 & L_{xx}L_{xy} \\ L_{xx}L_{xy} & L_{yy}^2 + L_{xy}^2 \end{bmatrix} \right) \right\}^{1/2}, \quad (55)$$

and set $k = 1$;

1) set $x'_{k-1} := P_{\Omega_x}(x_{k-1})$ and compute $(\tilde{x}_k, \tilde{y}_k) \in \mathcal{X} \times \mathcal{Y}$ as

$$\tilde{x}_k = [I + \lambda A]^{-1}(x_{k-1} - \lambda F_x(x'_{k-1}, y_{k-1})), \quad \tilde{y}_k = [I + \lambda B]^{-1}(y_{k-1} - \lambda F_y(\tilde{x}_k, y_{k-1})); \quad (56)$$

2) compute (x_k, y_k) as

$$x_k := \tilde{x}_k - \lambda[F_x(\tilde{x}_k, \tilde{y}_k) - F_x(x'_{k-1}, y_{k-1})], \quad y_k := \tilde{y}_k - \lambda[F_y(\tilde{x}_k, \tilde{y}_k) - F_y(\tilde{x}_k, y_{k-1})], \quad (57)$$

set $k \leftarrow k + 1$, and go to step 1.

end

It is easy to see that (55) and the assumption that $L_{xy} > 0$ imply that

$$\xi := \frac{1}{\tilde{L}} \max\{L_{xx}, L_{yy}\} < 1, \quad (58)$$

Proposition 5.3. *Tseng's based inexact BD-HPE method is a special instance of the BD-HPE framework, where $\sigma_x = \tilde{\sigma}_x = \lambda L_{xx}$ and $\sigma_y = \lambda L_{yy}$, and for every $k \in \mathbb{N}$,*

$$\lambda_k = \lambda, \quad \varepsilon_k^x = \varepsilon_k^y = 0,$$

and

$$\tilde{a}_k = \frac{1}{\lambda}(x_{k-1} - \tilde{x}_k) - F_x(x'_{k-1}, y_{k-1}), \quad \tilde{b}_k = \frac{1}{\lambda}(y_{k-1} - \tilde{y}_k) - F_y(\tilde{x}_k, y_{k-1}). \quad (59)$$

Proof. Applying Proposition 4.3 to the quintuple $(G, C, \Omega, x, \sigma) = (F_x(\cdot, y_{k-1}), A, \Omega_x, x_{k-1}, \lambda L_{xx})$, and also to the quintuple $(G, C, \Omega, x, \sigma) = (F_y(\tilde{x}_k, \cdot), B, \mathcal{Y}, y_{k-1}, \lambda L_{yy})$, and noting (57) and the definition of \tilde{a}_k and \tilde{b}_k , we conclude that (30) holds and that $(\tilde{x}_k, \tilde{a}_k)$ and $(\tilde{y}_k, \tilde{b}_k)$ satisfy (27), (28), and (29), respectively, with $\sigma_x = \tilde{\sigma}_x = \lambda L_{xx}$, $\sigma_y = \lambda L_{yy}$ and $\varepsilon_k^x = \varepsilon_k^y = 0$. It remains to show that $\lambda_k = \lambda$ satisfies (26) and that $\max\{\tilde{\sigma}_x, \sigma_y\} < 1$. Indeed, using the fact that $\tilde{\sigma}_x = \lambda L_{xx}$ and $\sigma_y = \lambda L_{yy}$, the definition of \tilde{L} in (55), the assumption that $\lambda \leq \sigma/\tilde{L}$ and $\sigma \leq 1$, and (58), we easily see that (26) holds and that

$$\max\{\tilde{\sigma}_x, \sigma_y\} = \lambda \max\{L_{xx}, L_{yy}\} \leq \frac{\sigma}{\tilde{L}} \max\{L_{xx}, L_{yy}\} = \sigma \xi \leq \xi < 1. \quad \square$$

The following convergence rate result now follows as an immediate consequence of Proposition 5.3 and Theorems 3.2 and 3.3.

Theorem 5.4. *Consider the sequences $\{(x_k, y_k)\}$, $\{(\tilde{x}_k, \tilde{y}_k)\}$ generated by Tseng's based inexact BD-HPE method with $\lambda = \sigma/\tilde{L}$, where \tilde{L} is given by (55). Define the sequence $\{(\tilde{a}_k, \tilde{b}_k)\}$ according to (59) and the sequences $\{(\tilde{x}_k^a, \tilde{y}_k^a)\}$, $\{(\tilde{a}_k^a, \tilde{b}_k^a)\}$, $\{\tilde{F}_k^a\}$, $\{\varepsilon_{k,F}^a\}$, $\{\varepsilon_{k,A}^a\}$ and $\{\varepsilon_{k,B}^a\}$ as in Theorem 5.2. Let d_0 denote the distance of the initial point $(x_0, y_0) \in \mathcal{X} \times \mathcal{Y}$ to the solution set of (19). Then, for every $k \in \mathbb{N}$, the following statements hold:*

a) $(\tilde{a}_k, \tilde{b}_k) \in A(\tilde{x}_k) \times B(\tilde{y}_k)$, and if $\sigma < 1$, there exists $i \leq k$ such that

$$\|F(\tilde{x}_i, \tilde{y}_i) + (\tilde{a}_i, \tilde{b}_i)\| \leq \frac{\tilde{L}d_0}{\sigma\sqrt{k}} \sqrt{\frac{1+\sigma}{1-\sigma}};$$

b) we have

$$\tilde{F}_k^a \in F^{\varepsilon_k^a, F}(\tilde{x}_k^a, \tilde{y}_k^a), \quad (\tilde{a}_k^a, \tilde{b}_k^a) \in A^{\varepsilon_k^a, A}(\tilde{x}_k^a) \times B^{\varepsilon_k^a, B}(\tilde{y}_k^a),$$

and

$$\left\| \tilde{F}_k^a + (\tilde{a}_k^a, \tilde{b}_k^a) \right\| \leq \frac{2\tilde{L}d_0}{k\sigma}, \quad \varepsilon_k^a \leq \frac{2\tilde{L}d_0^2}{k\sigma}(1 + \eta),$$

where

$$\eta := \frac{2\sqrt{2}\sigma}{1 - \xi\sigma} \left(1 + \frac{1}{(1 - \xi\sigma)^2} \right)^{1/2}$$

and ξ is defined in (58).

Also, if F is affine, then $\tilde{F}_k^a = F(\tilde{x}_k^a, \tilde{y}_k^a)$. In addition, if A (resp., B) is affine, then $\tilde{a}_k^a \in A(\tilde{x}_k^a)$ (resp., $\tilde{b}_k^a \in B(\tilde{y}_k^a)$).

Proof. This result follows immediately from Proposition 5.3 and Theorems 3.2 and 3.3 by specializing the latter two results to the case where $\sigma_x = \tilde{\sigma}_x = \lambda L_{xx}$, $\sigma_y = \lambda L_{yy}$, $\lambda_k = \lambda := \sigma/\tilde{L}$, and $\varepsilon_k^x = \varepsilon_k^y = 0$ for every $k \in \mathbb{N}$, and using the fact that $\max\{\tilde{\sigma}_x, \sigma_y\} \leq \sigma\xi$ (see the proof of Proposition 5.3). \square

We observe that it is possible to transform the bounds in terms of \tilde{L} in the above result to bounds in terms of the quantities L_{xx} , L_{yy} and L_{xy} by using the estimate

$$\tilde{L} \leq \sqrt{L_{xx}^2 + L_{xy}^2 + L_{yy}^2} \leq \sqrt{2}\tilde{L},$$

which follows immediately from the definition of \tilde{L} in (55).

5.3 An inexact BD method for convex optimization

In this subsection, we are interested in developing a specific instance of the BD-HPE framework for solving a class of linearly constrained convex optimization.

In this subsection, we consider the following optimization problem:

$$\min\{f(y) + h(y) : Cy = d\}, \tag{60}$$

where the following assumptions are made:

- O.1)** $C : \mathcal{Y} \rightarrow \mathcal{X}$ is a nonzero linear map and $d \in \mathcal{X}$;
- O.2)** $f, h : \mathcal{Y} \rightarrow (-\infty, \infty]$ are proper closed convex functions;
- O.3)** $\text{dom}(h) \subset \text{dom}(f)$ and there exist a point $\hat{y} \in \text{ri}(\text{dom } h) \cap \text{ri}(\text{dom } f)$ such that $C\hat{y} = d$;
- O.4)** the solution set of (60) is non-empty;
- O.5)** f is differentiable on $\text{cl}(\text{dom } h)$ and ∇f is L -Lipschitz continuous on $\text{cl}(\text{dom } h)$.

We now make some observations. First, under the above assumptions, y^* is an optimal solution if, and only if, it satisfies the condition

$$0 \in \partial f(y) + \partial h(y) + N_{\mathcal{M}}(y), \tag{61}$$

where $\mathcal{M} := \{y \in \mathcal{Y} : Cy = d\}$. Second, the above assumptions also guarantee that $\partial f + \partial h + N_{\mathcal{M}}$ is maximal monotone.

Clearly, y^* is an optimal solution if, and only if, there exists $x^* \in \mathcal{X}$ such that the pair $(y, x) = (y^*, x^*)$ satisfies

$$Cy - d = 0, \quad \nabla f(y) + \partial h(y) - C^*x \ni 0, \tag{62}$$

or equivalently, to the inclusion problem (19) with x and y swapped, where

$$F(x, y) := \begin{pmatrix} Cy - d \\ -C^*x \end{pmatrix}, \quad \forall (x, y) \in \mathcal{X} \times \mathcal{Y}, \quad A(\cdot) = 0, \quad B(\cdot) = \partial f(\cdot) + \partial h(\cdot).$$

We now state the algorithm that we are interested in studying in this subsection.

An inexact BD method for (60):

0) Let $(x_0, \tilde{y}_0) \in \mathcal{X} \times \mathcal{Y}$ and $0 < \sigma \leq 1$ be given, let $\lambda > 0$ be such that

$$\lambda L + \lambda^2 \|C\|^2 = \sigma^2. \quad (63)$$

and set $k = 1$;

1) compute

$$\tilde{x}_k = x_{k-1} - \lambda(C\tilde{y}_{k-1} - d), \quad \tilde{y}_k = (I + \lambda\partial h)^{-1}[\tilde{y}_{k-1} - \lambda(\nabla f(\tilde{y}_{k-1}) - C^*\tilde{x}_k)], \quad (64)$$

2) set $x_k = \tilde{x}_k + \lambda C(\tilde{y}_{k-1} - \tilde{y}_k)$ and $k \leftarrow k + 1$, and go to step 1.

end

Define

$$\bar{\xi} := (\lambda L)^{1/2} < 1, \quad (65)$$

where the inequality is due to (63) and the assumption that $C \neq 0$ (see O.1).

Proposition 5.5. *The above inexact BD method for (60) is a special instance of the BD-HPE framework, where $\sigma_x = \bar{\sigma}_x = 0$, $\sigma_y = (\lambda L)^{1/2}$ and, for every $k \in \mathbb{N}$,*

$$\lambda_k = \lambda, \quad \varepsilon_k^x = 0, \quad \varepsilon_k^y = \frac{L}{2} \|\tilde{y}_k - \tilde{y}_{k-1}\|^2, \quad (66)$$

and

$$\tilde{a}_k = 0, \quad \tilde{b}_k = \frac{1}{\lambda}(\tilde{y}_{k-1} - \tilde{y}_k) + C^*\tilde{x}_k \in [\partial_{\varepsilon_k^y} f + \partial h](\tilde{y}_k), \quad y_{k-1} = \tilde{y}_{k-1}, \quad (67)$$

Proof. Applying Proposition 4.1 with $G \equiv C\tilde{y}_{k-1} - d$, $C \equiv 0$ and $x = x_{k-1}$, and noting the definition of \tilde{a}_k , we conclude that $(\tilde{x}_k, \tilde{a}_k)$ satisfies (27) and (28) with $\sigma_x = 0$ and $\bar{\sigma}_x = 0$, respectively. Applying Proposition 4.2 with $T = \partial h$, $G \equiv -C^*\tilde{x}_k$ and $x = \tilde{y}_{k-1}$ and noting the definition of \tilde{b}_k , we conclude that $(\tilde{y}_k, \tilde{b}_k, \varepsilon_k^y)$ satisfies (29) with $\sigma_y = (\lambda L)^{1/2}$ and that the inclusion in (67) holds. Moreover, (67) and the update rule for x_k in step 2 of the algorithm imply that (30) holds. Also, by (65), we have

$$\max\{\tilde{\sigma}_x, \sigma_y\} = (\lambda L)^{1/2} = \bar{\xi} < 1. \quad (68)$$

In addition, using (63) and the fact that $\sigma_x = \bar{\sigma}_x = 0$, $\sigma_y^2 = \lambda L$ and $L_{xy} = \|C\|$, we easily see that that λ satisfies (26). \square

We are now ready to state the convergence rate result for the inexact BD method for (60).

Theorem 5.6. *Consider the sequences $\{x_k\}$ and $\{(\tilde{x}_k, \tilde{y}_k)\}$ generated by the inexact BD method for (60), and the sequences $\{\tilde{b}_k\}$ and $\{\varepsilon_k^y\}$ defined as in (67) and (66). Moreover, for each $k \in \mathbb{N}$, define*

$$(\tilde{x}_k^a, \tilde{y}_k^a) = \frac{1}{k} \sum_{i=1}^k (\tilde{x}_i, \tilde{y}_i), \quad \tilde{b}_k^a = \frac{1}{k} \sum_{i=1}^k \tilde{b}_i,$$

and

$$\varepsilon_{k,B}^a := \frac{1}{k} \sum_{i=1}^k \left(\varepsilon_k^y + \langle \tilde{y}_i - \tilde{y}_k^a, \tilde{b}_i \rangle \right) \geq 0.$$

Let d_0 denote the distance of the initial point $(x_0, \tilde{y}_0) \in \mathcal{X} \times \mathcal{Y}$ to the solution set of (62). Then, for every $k \in \mathbb{N}$, the following statements hold:

a) $\tilde{b}_k \in (\partial_{\varepsilon_k^y} f + \partial h)(\tilde{y}_k)$, and if $\sigma < 1$, there exists $i \leq k$ such that

$$\left\| (C\tilde{y}_i - d, -C^* \tilde{x}_i + \tilde{b}_i) \right\| \leq \frac{d_0}{\lambda \sqrt{k}} \sqrt{\frac{1+\sigma}{1-\sigma}}, \quad \varepsilon_i^y \leq \frac{\sigma^2 d_0^2}{(1-\sigma^2)\lambda k};$$

b) we have

$$\tilde{b}_k^a \in \partial_{\varepsilon_{k,B}^a} (f + h)(\tilde{y}_k^a), \tag{69}$$

and

$$\left\| (C\tilde{y}_k^a - d, -C^* \tilde{x}_k^a + \tilde{b}_k^a) \right\| \leq \frac{2d_0}{k\lambda}, \quad \varepsilon_{k,B}^a \leq \frac{2d_0^2}{k\lambda} (1 + \bar{\eta}),$$

where

$$\bar{\eta} := \frac{2\sqrt{2}\sigma}{1-\bar{\xi}} \left(1 + \frac{1}{(1-\bar{\xi})^2} \right)^{1/2}$$

and $\bar{\xi}$ is defined in (65).

Proof. This result follows immediately from Proposition 5.5 and Theorems 3.2 and 3.3 by specializing the latter two results to the case where $\lambda_k = \lambda := \sigma\lambda$, $\sigma_x = \tilde{\sigma}_x = 0$, $\sigma_y = (\lambda L)^{1/2}$, and ε_k^x and ε_k^y are given by (37), and using the fact that, by (68), $\max\{\tilde{\sigma}_x, \sigma_y\} = \bar{\xi}$. Observe also that (69) follows Theorem 2.2(b) and the fact that $\tilde{b}_k \in \partial_{\varepsilon_k^y} (f + h)(\tilde{y}_k)$, in view of statement a). \square

Note that it is possible to replace λ in the above estimates using its explicit formula:

$$\frac{1}{\lambda} = \frac{L + \sqrt{L^2 + 4\sigma^2\|C\|^2}}{2\sigma^2} \leq \frac{L}{\sigma^2} + \frac{\|C\|}{\sigma}.$$

We can interpret the above theorem in terms of the optimality condition (62) as follows. Defining $\tilde{v}_i = \tilde{b}_i - C^* \tilde{x}_i$ and $\varepsilon_i = \varepsilon_i^y$, it follows from its statement a) that

$$\tilde{v}_i \in \partial_{\varepsilon_i} (f + h)(\tilde{y}_i) - C^* \tilde{x}_i, \quad \|C\tilde{y}_i - d\| = \mathcal{O}(1/k^{1/2}), \quad \|\tilde{v}_i\| = \mathcal{O}(1/k^{1/2}), \quad \varepsilon_i = \mathcal{O}(1/k),$$

while, defining $\tilde{v}_k^a = \tilde{b}_k^a - C^* \tilde{x}_k^a$ and $\varepsilon_k^a = \varepsilon_{k,B}^a$, its statement b) implies that

$$\tilde{v}_k^a \in \partial_{\varepsilon_k^a} (f + h)(\tilde{y}_k^a) - C^* \tilde{x}_k^a, \quad \|C\tilde{y}_k^a - d\| = \mathcal{O}(1/k), \quad \|\tilde{v}_k^a\| = \mathcal{O}(1/k), \quad \varepsilon_k^a = \mathcal{O}(1/k).$$

Finally, we have shown in this section that the inexact BD method for solving (60) presented in this subsection is a special case of the BD-HPE framework in which the first equation (resp., second inclusion) in (62) is identified with the first (resp., second) inclusion in (20). Clearly, it is also possible to derive a variant of the BD method presented in this subsection which is also a special case of the BD-HPE framework but with the second inclusion (resp., first equation) in (62) identified with the first (resp., second) inclusion in (20). Note that for the latter variant, if the procedure of Proposition 4.2 is used for approximately solving the second inclusion in (62), then we have $\sigma_x = (\lambda L)^{1/2}$, $\tilde{\sigma}_x = 0$ and $\sigma_y = 0$, which allows us to choose a larger λ than the one in (63), namely $\lambda > 0$ such that $\max\{\lambda L, \lambda^2\|C\|^2\} = \sigma^2$. Note also that the latter variant provides a meaningful instance of the BD-HPE framework where $\tilde{\sigma}_x < \sigma_x$.

6 Sum of two maximal monotone operators

In this section, we consider the monotone inclusion problem consisting of the sum of two maximal monotone operators and show how it can be transformed to a problem of form (19), which can then be solved by any instance of the BD-HPE framework.

Consider the problem

$$0 \in (A + B)(x) \quad (70)$$

where $A, B : \mathcal{X} \rightrightarrows \mathcal{X}$ are maximal monotone. We assume that the resolvents of A and B are easily computable. Note that (70) is equivalent to the existence of $b \in \mathcal{X}$ such that

$$-b \in A(x), \quad b \in B(x),$$

or equivalently,

$$0 \in b + A(x), \quad 0 \in -x + B^{-1}(b).$$

Hence, the inclusion problem (70) is equivalent to the monotone inclusion problem

$$0 \in [F + (A \otimes B^{-1})](x, b), \quad (71)$$

where

$$F(x, b) = (b, -x). \quad (72)$$

Hence, we can apply any instance of the BD-HPE framework, and in particular the exact BD-HPE method of Subsection 5.1, to the inclusion problem (71) in order to compute an approximate solution of (70). In the following two subsections, we will discuss in detail these approaches, stating a general BD-HPE framework for (71) in Subsection 6.1, and an exact BD-HPE method for (71) in Subsection 6.2.

6.1 Block-decomposition HPE framework for (70)

We start by stating a general BD-HPE framework for solving (70).

Inexact BD-HPE framework for (70):

- 0) Let $(x_0, b_0) \in \mathcal{X} \times \mathcal{X}$, $\tilde{\sigma}_x, \sigma_y \in [0, 1)$ and $\sigma, \sigma_x \in [0, 1]$ be given and set $k = 1$;
- 1) choose $\lambda_k > 0$ such that (26) holds with $L_{xy} = 1$;
- 2) compute $\tilde{x}_k, \tilde{a}_k \in \mathcal{X}$ and $\varepsilon_k^x \geq 0$ such that

$$\tilde{a}_k \in A^{\varepsilon_k^x}(\tilde{x}_k), \quad \|\lambda_k[b_{k-1} + \tilde{a}_k] + \tilde{x}_k - x_{k-1}\|^2 + 2\lambda_k\varepsilon_k^x \leq \sigma_x^2 \|\tilde{x}_k - x_{k-1}\|^2; \quad (73)$$

$$\|\lambda_k[b_{k-1} + \tilde{a}_k] + \tilde{x}_k - x_{k-1}\| \leq \tilde{\sigma}_x \|\tilde{x}_k - x_{k-1}\|; \quad (74)$$

- 3) compute $\tilde{y}_k, \tilde{b}_k \in \mathcal{X}$ and $\varepsilon_k^y \geq 0$ such that

$$\tilde{y}_k \in (B^{-1})^{\varepsilon_k^y}(\tilde{b}_k), \quad \|\lambda_k[-\tilde{x}_k + \tilde{y}_k] + \tilde{b}_k - b_{k-1}\|^2 + 2\lambda_k\varepsilon_k^y \leq \sigma_y^2 \|\tilde{b}_k - b_{k-1}\|^2; \quad (75)$$

- 3) set

$$(x_k, b_k) = (x_{k-1}, b_{k-1}) - \lambda_k[(\tilde{b}_k, -\tilde{x}_k) + (\tilde{a}_k, \tilde{y}_k)] = (x_{k-1}, b_{k-1}) - \lambda_k(\tilde{b}_k + \tilde{a}_k, \tilde{y}_k - \tilde{x}_k), \quad (76)$$

$k \leftarrow k + 1$ and go to step 1.

end

Clearly, the above framework is nothing else but the BD-HPE framework of Section 3 for the monotone inclusion problem (71). Note that it is stated in terms of B^{-1} . It is possible to state a version of it which replaces condition (75) based on B^{-1} by a sufficient condition based on B as described by the following result.

Proposition 6.1. *If $\tilde{x}_k, \tilde{y}_k, b_{k-1}, \tilde{b}_k \in \mathcal{X}$, $\lambda_k > 0$ and $\varepsilon_k^y \geq 0$ satisfy*

$$\tilde{b}_k \in (B)^{\varepsilon_k^y}(\tilde{y}_k), \quad \|\lambda_k^{-1}[\tilde{b}_k - b_{k-1}] + \tilde{y}_k - \tilde{x}_k\|^2 + 2\lambda_k^{-1}\varepsilon_k^y \leq \left(\frac{\sigma_y}{1 + \sigma_y}\right)^2 \|\tilde{y}_k - \tilde{x}_k\|^2, \quad (77)$$

then (75) holds.

Proof. Since $(B^{-1})^\varepsilon = (B^\varepsilon)^{-1}$, the inclusion on (77) implies that $\tilde{y}_k \in (B^{-1})^{\varepsilon_k^y}$. Using the inequality in (77) and the triangle inequality for norms, we have

$$\begin{aligned} \|\tilde{y}_k - \tilde{x}_k\| &\leq \|\lambda_k^{-1}[\tilde{b}_k - b_{k-1}]\| + \|\lambda_k^{-1}[\tilde{b}_k - b_{k-1}] + \tilde{y}_k - \tilde{x}_k\| \\ &\leq \|\lambda_k^{-1}[\tilde{b}_k - b_{k-1}]\| + \frac{\sigma_y}{1 + \sigma_y} \|\tilde{y}_k - \tilde{x}_k\| \end{aligned}$$

Hence, $\|\tilde{y}_k - \tilde{x}_k\| \leq (1 + \sigma_y)\|\lambda_k^{-1}[\tilde{b}_k - b_{k-1}]\|$, which combined with the inequality in (77) yields

$$\|\lambda_k^{-1}[\tilde{b}_k - b_{k-1}] + \tilde{y}_k - \tilde{x}_k\|^2 + 2\lambda_k^{-1}\varepsilon_k^y \leq \sigma_y^2 \|\lambda_k^{-1}[\tilde{b}_k - b_{k-1}]\|^2.$$

To end the proof, multiply the above inequality by λ_k^2 . □

Note that the pair $(\tilde{y}, \tilde{b}) = (\tilde{y}_k, \tilde{b}_k)$ in the above result is an approximate solution of the proximal point equation $\lambda_k^{-1}[\tilde{b} - b_{k-1}] + (\tilde{y} - \tilde{x}_k) = 0$ and $\tilde{y} \in B(\tilde{b})$ in the sense described in the paragraph after the statement of the HPE method in Section 2.2.

The specialization of Theorems 3.2 and 3.3 for the above method are as follows.

Theorem 6.2. *Consider the sequences $\{\lambda_k\}$, $\{(\varepsilon_k^x, \varepsilon_k^y)\}$, $\{(\tilde{x}_k, \tilde{b}_k)\}$ and $\{(\tilde{a}_k, \tilde{y}_k)\}$ generated by the BD-HPE method for (70). Moreover, for every $k \in \mathbb{N}$, define:*

$$\begin{aligned} (\tilde{x}_k^a, \tilde{y}_k^a, \tilde{a}_k^a, \tilde{b}_k^a) &= \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i (\tilde{x}_i, \tilde{y}_i, \tilde{a}_i, \tilde{b}_i), \\ \varepsilon_{k,A}^a &= \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i (\varepsilon_i^x + \langle \tilde{x}_i - \tilde{x}_k^a, \tilde{a}_i \rangle), \quad \varepsilon_{k,B}^a = \frac{1}{\Lambda_k} \sum_{i=1}^k \lambda_i (\varepsilon_i^y + \langle \tilde{b}_i - \tilde{b}_k^a, \tilde{y}_i \rangle). \end{aligned}$$

Let d_0 denote the distance of the initial point $(x_0, b_0) \in \mathcal{X} \times \mathcal{X}$ to the solution set of (71), i.e.:

$$d_0 := \min \left\{ (\|x - x_0\|^2 + \|b - b_0\|^2)^{1/2} : -b \in A(x), b \in B(x) \right\}.$$

Then, for every $k \in \mathbb{N}$, the following statements hold:

a) $(\tilde{a}_k, \tilde{b}_k) \in A^{\varepsilon_k^x}(\tilde{x}_k) \times B^{\varepsilon_k^y}(\tilde{y}_k)$, and, if $\sigma < 1$, then there exists $i \leq k$ such that

$$\left\| (\tilde{b}_i + \tilde{a}_i, -\tilde{x}_i + \tilde{y}_i) \right\| \leq d_0 \sqrt{\frac{1 + \sigma}{1 - \sigma} \left(\frac{1}{\sum_{j=1}^k \lambda_j^2} \right)}, \quad \varepsilon_i^x + \varepsilon_i^y \leq \frac{\sigma^2 d_0^2 \lambda_i}{2(1 - \sigma^2) \sum_{j=1}^k \lambda_j^2};$$

b) $(\tilde{a}_k^a, \tilde{b}_k^a) \in A^{\varepsilon_{k,A}^a}(\tilde{x}_k^a) \times B^{\varepsilon_{k,B}^a}(\tilde{y}_k^a)$ and

$$\left\| (\tilde{b}_k^a + \tilde{a}_k^a, -\tilde{x}_k^a + \tilde{y}_k^a) \right\| \leq \frac{2d_0}{\Lambda_k}, \quad \varepsilon_{k,A}^a + \varepsilon_{k,B}^a \leq \frac{2d_0^2}{\Lambda_k} (1 + \eta_k),$$

where η_k is defined in (39) with $L_{xy} = 1$.

Proof. This result follows as an immediate consequence of Theorems 3.2 and 3.3 applied to (71)-(72) and noting that in this case F is affine and $L_{xy} = 1$. □

6.2 Exact BD-HPE method for (70)

In this subsection, we state an exact BD-HPE method for (70) and corresponding convergence rate results.

Exact BD-HPE method for (70):

0) Let $x_0, \tilde{b}_0 \in \mathcal{X}$ and $\lambda \in (0, 1]$ be given, and set $k = 1$;

1) compute $\tilde{x}_k, \tilde{b}_k \in \mathcal{X}$ as

$$\tilde{x}_k = (I + \lambda A)^{-1}(x_{k-1} - \lambda \tilde{b}_{k-1}), \quad \tilde{b}_k = (I + \lambda B^{-1})^{-1}(\tilde{b}_{k-1} + \lambda \tilde{x}_k);$$

2) set $x_k = \tilde{x}_k - \lambda(\tilde{b}_k - \tilde{b}_{k-1})$ and go to step 1.

end

The method above is nothing else but the exact BD-HPE method of Subsection 5.1 applied to (71)-(72) with variable b replacing variable y and vice-versa.

The following well-known result describes how the resolvent of B^{-1} used in step 1) can be computed using the resolvent of B .

Lemma 6.3. *Let $b, u \in \mathcal{X}$ be given. Then,*

$$b = (I + \lambda B^{-1})^{-1}(u) \Leftrightarrow b = u - \lambda(I + \lambda^{-1}B)^{-1}(\lambda^{-1}u).$$

In view of the above result, \tilde{b}_k may be computed as

$$\tilde{b}_k = \tilde{b}_{k-1} + \lambda \tilde{x}_k - \lambda(I + \lambda^{-1}B)^{-1}(\lambda^{-1}\tilde{b}_{k-1} + \tilde{x}_k).$$

The following iteration-complexity bounds for solving the inclusion problem (70) can now be stated as an immediate consequence of Theorem 5.2.

Theorem 6.4. *Consider the sequences $\{x_k\}$ and $\{(\tilde{x}_k, \tilde{b}_k)\}$ generated by the BD-HPE method for (70) and define the sequence $\{(\tilde{a}_k, \tilde{y}_k)\}$ as*

$$\tilde{a}_k = \frac{1}{\lambda}(x_{k-1} - \tilde{x}_k) - \tilde{b}_{k-1}, \quad \tilde{y}_k = \frac{1}{\lambda}(\tilde{b}_{k-1} - \tilde{b}_k) + \tilde{x}_k. \quad (78)$$

Moreover, for every $k \in \mathbb{N}$, define:

$$\begin{aligned} (\tilde{x}_k^a, \tilde{y}_k^a, \tilde{a}_k^a, \tilde{b}_k^a) &= \frac{1}{k} \sum_{i=1}^k (\tilde{x}_i, \tilde{y}_i, \tilde{a}_i, \tilde{b}_i), \\ \varepsilon_{k,A}^a &= \frac{1}{k} \sum_{i=1}^k \langle \tilde{x}_i - \tilde{x}_k^a, \tilde{a}_i \rangle, \quad \varepsilon_{k,B}^a = \frac{1}{k} \sum_{i=1}^k \langle \tilde{b}_i - \tilde{b}_k^a, \tilde{y}_i \rangle. \end{aligned}$$

Let d_0 denote the distance of the initial point $(x_0, \tilde{b}_0) \in \mathcal{X} \times \mathcal{X}$ to the solution set of (71), i.e.:

$$d_0 := \min \left\{ \left(\|x - x_0\|^2 + \|b - \tilde{b}_0\|^2 \right)^{1/2} : -b \in A(x), b \in B(x) \right\}.$$

Then, for every $k \in \mathbb{N}$, the following statements hold:

a) $(\tilde{a}_k, \tilde{b}_k) \in A(\tilde{x}_k) \times B(\tilde{y}_k)$, and if $\lambda < 1$, there exists $i \leq k$ such that

$$\left\| (\tilde{b}_i + \tilde{a}_i, -\tilde{x}_i + \tilde{y}_i) \right\| \leq \frac{d_0}{\lambda \sqrt{k}} \sqrt{\frac{1+\lambda}{1-\lambda}};$$

b) $(\tilde{a}_k^a, \tilde{b}_k^a) \in A^{\varepsilon_{k,A}^a}(\tilde{x}_k^a) \times B^{\varepsilon_{k,B}^a}(\tilde{y}_k^a)$ and

$$\left\| \left(\tilde{b}_k^a + \tilde{a}_k^a, -\tilde{x}_k^a + \tilde{y}_k^a \right) \right\| \leq \frac{2d_0}{k\lambda}, \quad \varepsilon_{k,A}^a + \varepsilon_{k,B}^a \leq \frac{2d_0^2}{k\lambda} \left(1 + 2\sqrt{2}\lambda \right).$$

Proof. This result follows as an immediate consequence of Theorem 5.2 applied to (71)-(72) and noting that in this case F is affine and $L_{xy} = 1$, and hence $\lambda = \sigma$. \square

We end this section by discussing the special case of the exact BD-HPE method for (70) in which $\lambda = 1$. It can be easily shown that this algorithm is equivalent to the Douglas-Rachford splitting method (see for example [9]). Hence, Theorem 6.4(b) for $\lambda = 1$ gives an ergodic complexity estimation of the Douglas-Rachford method. However, as far as we know, the exact BD-HPE method for (70) with $\lambda < 1$ is new.

7 Convergence of the alternating direction method of multipliers

In this section, we consider the ADMM for solving a large class of linearly constrained convex programming problems with proper closed convex objective functions and show that it can be interpreted as a specific instance of the BD-HPE framework applied to a two-block monotone inclusion problem.

We assume in this section that \mathcal{X} , \mathcal{Y} and \mathcal{S} are inner product spaces whose inner products and associated norms are denoted by $\langle \cdot, \cdot \rangle$ and $\| \cdot \|$. Consider the problem

$$\min\{f(y) + g(s) : Cy + Ds = c\} \tag{79}$$

where $c \in \mathcal{X}$, $C : \mathcal{Y} \rightarrow \mathcal{X}$ and $D : \mathcal{S} \rightarrow \mathcal{X}$ are linear operators, and $f : \mathcal{Y} \rightarrow (-\infty, \infty]$ and $g : \mathcal{S} \rightarrow (-\infty, \infty]$ are proper closed convex functions. Throughout this section, we also assume that the resolvent of ∂f and ∂g can be computed exactly.

The Lagrangian function $L : (\mathcal{Y} \times \mathcal{S}) \times \mathcal{X} \rightarrow (-\infty, \infty]$ for problem (79) is defined as

$$\mathcal{L}(y, s; x) = f(y) + g(s) - \langle x, Cy + Ds - c \rangle. \tag{80}$$

We make the following assumptions throughout this section:

C.1) there exists a saddle point of \mathcal{L} , i.e., a point $(y^*, s^*; x^*)$ such that $\mathcal{L}(y^*, s^*; x^*)$ is finite and

$$\min_{(y,s) \in \mathcal{Y} \times \mathcal{S}} \mathcal{L}(y, s; x^*) = \mathcal{L}(y^*, s^*; x^*) = \max_{x \in \mathcal{X}} \mathcal{L}(y^*, s^*; x); \tag{81}$$

C.2) $\text{ri}(\text{dom } g^*) \cap \text{range } D^* \neq \emptyset$;

C.3) C is injective.

For a scalar $\rho \geq 0$, the ρ -augmented Lagrangian function $\mathcal{L}_\rho : (\mathcal{Y} \times \mathcal{S}) \times \mathcal{X} \rightarrow (-\infty, \infty]$ associated with (79) is defined as

$$\mathcal{L}_\rho(y, s; x) := f(y) + g(s) + \langle x, c - Cy - Ds \rangle + \frac{\rho}{2} \|Cy + Ds - c\|^2.$$

We next state the alternating direction method of multipliers applied to problem (79).

Alternating direction method of multipliers:

0) Let $\rho > 0$ and $(x_0, \tilde{y}_0) \in \mathcal{X} \times \mathcal{Y}$ be given and set $k = 1$;

1) compute $\tilde{s}_k \in \mathcal{S}$ as

$$\tilde{s}_k = \operatorname{argmin}_s \{ \mathcal{L}_\rho(\tilde{y}_{k-1}, s; x_{k-1}) \} = \operatorname{argmin}_s \left\{ g(s) - \langle D^* x_{k-1}, s \rangle + \frac{\rho}{2} \| C\tilde{y}_{k-1} + Ds - c \|^2 \right\}. \quad (82)$$

and $\tilde{y}_k \in \mathcal{Y}$ as

$$\tilde{y}_k = \operatorname{argmin}_y \{ \mathcal{L}_\rho(y, \tilde{s}_k; x_{k-1}) \} = \operatorname{argmin}_y \left\{ f(y) - \langle C^* x_{k-1}, y \rangle + \frac{\rho}{2} \| Cy + D\tilde{s}_k - c \|^2 \right\}. \quad (83)$$

2) set $x_k = x_{k-1} - \rho(C\tilde{y}_k + D\tilde{s}_k - c)$ and $k \leftarrow k + 1$, and go to step 1;

end

Our goal in the remaining part of this section is to show that the ADMM is a special case of the exact BD-HPE method of Subsection 5.1 for a specific monotone inclusion problem of the form (19). As a by-product, we also derive convergence rate results for the ADMM.

We start by giving a preliminary technical result about the ADMM.

Proposition 7.1. *Let $(x_{k-1}, \tilde{y}_{k-1}) \in \mathcal{X} \times \mathcal{Y}$ be given. Then, the following statements hold:*

a) $\tilde{s}_k \in \mathcal{S}$ is an optimal solution of (82) if, and only if, the point

$$\tilde{x}_k := x_{k-1} - \rho(C\tilde{y}_{k-1} + D\tilde{s}_k - c), \quad (84)$$

satisfies

$$\tilde{s}_k \in \partial g^*(D^* \tilde{x}_k); \quad (85)$$

b) if $(\tilde{s}_k, \tilde{y}_k, x_k)$ are computed according to the k -iteration of the ADMM, then

$$0 \in \partial(g^* \circ D^*)(\tilde{x}_k) + C\tilde{y}_{k-1} - c + \frac{1}{\rho}(\tilde{x}_k - x_{k-1}), \quad (86)$$

$$0 \in \partial f(\tilde{y}_k) - C^* \tilde{x}_k + \rho C^* C(\tilde{y}_k - \tilde{y}_{k-1}), \quad (87)$$

$$x_k = \tilde{x}_k - \rho C(\tilde{y}_k - \tilde{y}_{k-1}). \quad (88)$$

Proof. By (84) and the optimality conditions of (82), we have that \tilde{s}_k is an optimal solution of (82) if, and only if,

$$D^* \tilde{x}_k = D^*[x_{k-1} - \rho(C\tilde{y}_{k-1} + D\tilde{s}_k - c)] \in \partial g(\tilde{s}_k),$$

which in turn is equivalent to (85). On the other hand, (85) implies that $D\tilde{s}_k \in D\partial g^*(D^* \tilde{x}_k) \subset \partial(g^* \circ D^*)(\tilde{x}_k)$. Combining the latter inclusion with (84), we obtain (86). Moreover, (83) implies that

$$0 \in \partial f(\tilde{y}_k) - C^* x_{k-1} + \rho C^*(C\tilde{y}_k + D\tilde{s}_k - c).$$

Combining the above equation with (84), we obtain (87). Finally, (88) follows immediately (84) and the update rule for x_k in step 3 of the ADMM. \square

Proposition 7.2. *Given $(x_{k-1}, \tilde{y}_{k-1}) \in \mathcal{X} \times \mathcal{Y}$, define*

$$\hat{x}_k := (\rho \partial(g^* \circ D^*) + I)^{-1}[x_{k-1} + \rho(c - C\tilde{y}_{k-1})], \quad \hat{w}_k := \frac{1}{\rho}(x_{k-1} - \hat{x}_k) + c - C\tilde{y}_{k-1}.$$

Then, the following statements hold:

- a) $\tilde{s}_k \in \mathcal{S}$ is an optimal solution of (82) if, and only if, $\tilde{s}_k \in D^{-1}(\hat{w}_k) \cap \partial g^*(D^*\hat{x}_k)$;
b) if condition C.2) holds, then $D^{-1}(\hat{w}_k) \cap \partial g^*(D^*\hat{x}_k) \neq \emptyset$, and hence the set of optimal solutions of (82) is nonempty;
c) if condition C.3) holds, then the set of optimal solutions of (83) is nonempty.

Proof. First, observe that $(\tilde{x}, \tilde{w}) = (\hat{x}_k, \hat{w}_k)$ is the unique solution of

$$0 = \rho(C\tilde{y}_{k-1} - c + \tilde{w}) + \tilde{x} - x_{k-1}, \quad \tilde{w} \in (g^* \circ D^*)(\tilde{x}). \quad (89)$$

a) Assume first that \tilde{s}_k is an optimal solution of (82). By Proposition 7.1, we conclude that \tilde{x}_k given by (84) satisfies (85). Then,

$$D\tilde{s}_k \in D\partial g^*(D^*\tilde{x}_k) \subset \partial(g^* \circ D^*)(\tilde{x}_k),$$

which, together with (84), imply that $(\tilde{x}, \tilde{w}) = (\tilde{x}_k, D\tilde{s}_k)$ satisfies (89). Hence, in view of the observation made at the beginning of this proof, we conclude that $\tilde{x}_k = \hat{x}_k$ and $D\tilde{s}_k = \hat{w}_k$. These identities and inclusion (85) then imply that $\tilde{s}_k \in D^{-1}(\hat{w}_k) \cap \partial g^*(D^*\hat{x}_k)$.

Conversely, assume that $\tilde{s}_k \in D^{-1}(\hat{w}_k) \cap \partial g^*(D^*\hat{x}_k)$. Then, $\hat{w}_k = D\tilde{s}_k$, and hence $(\hat{x}_k, D\tilde{s}_k) = (\hat{x}_k, \hat{w}_k)$ satisfies (89). In particular,

$$\hat{x}_k = x_{k-1} - \rho(C\tilde{y}_{k-1} - c + D\tilde{s}_k) = \tilde{x}_k,$$

where the latter equality is due to (84). Since $\tilde{x}_k = \hat{x}_k$ and, by assumption, $\tilde{s}_k \in \partial g^*(D^*\hat{x}_k)$, we conclude that (85) holds, and hence that \tilde{s}_k is an optimal solution of (82).

b) Using the fact that (\hat{x}_k, \hat{w}_k) satisfies (89), we conclude that

$$\hat{w}_k \in \partial(g^* \circ D^*)(\hat{x}_k) = D(\partial g^*(D^*\hat{x}_k)).$$

where the latter equality is due to Assumption C.2). The latter inclusion clearly implies that $D^{-1}(\hat{w}_k) \cap \partial g^*(D^*\hat{x}_k) \neq \emptyset$.

c) This statement follows from the fact that, under Assumption C.3, the objective function of (83) is strongly convex. \square

We will now derive the aforementioned monotone inclusion problem of the form (19). Let $X \times Y := \text{dom } f \times (D^*)^{-1}(\text{dom } g^*)$ and $\Psi : X \times Y \rightarrow \mathbb{R}$ be defined as

$$\begin{aligned} \Psi(y, x) &= \min_s \mathcal{L}(y, s; x) = f(y) + \langle x, c - Cy \rangle + \left(\min_s g(s) - \langle D^*x, s \rangle \right) \\ &= f(y) + \langle x, c - Cy \rangle - g^*(D^*x). \end{aligned}$$

It is easy to see that the pair (y^*, x^*) as in Assumption C.1) satisfies

$$\max_{x \in X} \min_{(y, s) \in Y \times \mathcal{S}} \mathcal{L}(y, s; x) = \max_{x \in X} \min_{y \in Y} \Psi(y, x) = \Psi(y^*, x^*) = \min_{y \in Y} \max_{x \in X} \Psi(y, x) = \min_{(y, s) \in Y \times \mathcal{S}} \max_{x \in X} \mathcal{L}(y, s; x) \in \mathbb{R}.$$

The latter condition is in turn equivalent to (y^*, x^*) being a solution of the inclusion problem:

$$0 \in \partial f(y) - C^*x, \quad 0 \in \partial(g^* \circ D^*)(x) + Cy - c. \quad (90)$$

Under the assumption that C^*C is nonsingular, the latter inclusion problem is clearly equivalent to

$$0 \in (\rho C^*C)^{-1}(\partial f(y) - C^*x), \quad 0 \in \rho[\partial(g^* \circ D^*)(x) + Cy - c]. \quad (91)$$

and hence to inclusion problem (19) with

$$F(x, y) = \left[\begin{array}{c} \rho(Cy - c) \\ -(\rho C^*C)^{-1}C^*x \end{array} \right], \quad A(x) = \rho\partial(g^* \circ D^*)(x), \quad B(y) = (\rho C^*C)^{-1}\partial f(y). \quad (92)$$

If \mathcal{U} is an inner product space with inner product also denoted by $\langle \cdot, \cdot \rangle$, then a symmetric positive definite operator $M : \mathcal{U} \rightarrow \mathcal{U}$ defines another inner product, denoted by $\langle \cdot, \cdot \rangle_M$, as follows:

$$\langle u, u' \rangle_M = \langle u, Mu' \rangle, \quad \forall u, u' \in \mathcal{U}.$$

We will denote the norm associated with the above inner product by $\|\cdot\|_M$. Moreover, when $M = \tau I$ for some $\tau > 0$, where I denotes the identity operator, we denote the norm $\|\cdot\|_M$ simply by $\|\cdot\|_\tau$.

Proposition 7.3. *Assume that C^*C is nonsingular, $\text{ri}(\text{dom } g^*) \cap \text{Range}(D^*)$ is non-empty and consider the inner products $\langle \cdot, \cdot \rangle_{\rho^{-1}}$ in \mathcal{X} and $\langle \cdot, \cdot \rangle_{\tilde{C}_\rho}$ in \mathcal{Y} , where $\tilde{C}_\rho := \rho C^*C$. Then, the following statements hold:*

- a) *the map F defined in (92) is monotone with respect to the inner product $\langle \cdot, \cdot \rangle_{\rho^{-1}} + \langle \cdot, \cdot \rangle_{\tilde{C}_\rho}$ in $\mathcal{X} \times \mathcal{Y}$ and the operators A and B defined in (92) are maximal monotone with respect to $\langle \cdot, \cdot \rangle_{\rho^{-1}}$ and $\langle \cdot, \cdot \rangle_{\tilde{C}_\rho}$, respectively;*
- b) *F satisfies (21) with $L_{xy} = 1$;*
- c) *the sequence $\{(x_k, \tilde{y}_k)\}$ generated by the ADMM together with the sequence $\{\tilde{x}_k\}$ defined in (84) correspond to the same sequence $\{(x_k, \tilde{x}_k, \tilde{y}_k)\}$ generated by the exact BD-HPE method applied to the inclusion problem (19) with (F, A, B) given by (92) and with $\sigma = 1$ (or equivalently, $\lambda = 1$).*

Proof. Monotonicity of F in the rescaled space $\mathcal{X} \times \mathcal{Y}$ holds trivially. Maximal monotonicity of A (resp., B) in \mathcal{X} (resp., \mathcal{Y}) endowed with the norm $\|\cdot\|_{\rho^{-1}}$ (resp., $\|\cdot\|_{\tilde{C}_\rho}$) follows from the fact that this operator is the subdifferential of $g^* \circ D^*$ (resp., f) in the rescaled space. For b), observe that

$$\|F_x(x, y) - F_x(x, y')\|_{\rho^{-1}}^2 = \|\rho C(y - y')\|_{\rho^{-1}}^2 = \rho \|C(y - y')\|^2 = \langle y - y', (\rho C^*C)(y - y') \rangle = \|y - y'\|_{\tilde{C}_\rho}^2.$$

Statement c) follows immediately from Proposition 7.1 by noting that relations (86)-(88) reduce to the recursive formulas for obtaining $(x_k, \tilde{x}_k, \tilde{y}_k)$ in the the exact BD-HPE method applied to the inclusion problem (19) with (F, A, B) given by (92) and with $\lambda = 1$. \square

As a consequence of the previous proposition, we obtain the following convergence rate result.

Theorem 7.4. *Consider the sequence $\{(x_k, \tilde{y}_k)\}$ generated by the ADMM and the sequence $\{\tilde{x}_k\}$ defined according to (84). Consider also the sequence $\{(\tilde{a}_k, \tilde{b}_k)\}$ defined as*

$$\tilde{a}_k = x_{k-1} - \tilde{x}_k - \rho(C\tilde{y}_{k-1} - c), \quad \tilde{b}_k = \tilde{y}_{k-1} - \tilde{y}_k + \tilde{C}_\rho^{-1}C^*\tilde{x}_k, \quad (93)$$

where $\tilde{C}_\rho := \rho C^*C$. Moreover, define the sequences $\{(\tilde{x}_k^a, \tilde{y}_k^a)\}$, $\{(\tilde{a}_k^a, \tilde{b}_k^a)\}$ and $\{(\varepsilon_{k,x}^a, \varepsilon_{k,y}^a)\}$ as

$$(\tilde{x}_k^a, \tilde{y}_k^a) = \frac{1}{k} \sum_{i=1}^k (\tilde{x}_i, \tilde{y}_i), \quad (\tilde{a}_k^a, \tilde{b}_k^a) = \frac{1}{k} \sum_{i=1}^k (\tilde{a}_i, \tilde{b}_i), \quad (94)$$

and

$$\varepsilon_{k,x}^a = \frac{1}{k} \sum_{i=1}^k \left\langle \tilde{x}_i - \tilde{x}_k^a, \frac{1}{\rho} \tilde{a}_i \right\rangle, \quad \varepsilon_{k,y}^a = \frac{1}{k} \sum_{i=1}^k \left\langle \tilde{y}_i - \tilde{y}_k^a, \tilde{C}_\rho \tilde{b}_i \right\rangle. \quad (95)$$

Then, for every $k \in \mathbb{N}$,

$$\frac{1}{\rho} \tilde{a}_k^a \in \partial_{\varepsilon_{k,x}^a} (g^* \circ D^*)(\tilde{x}_k^a), \quad \tilde{C}_\rho \tilde{b}_k^a \in \partial_{\varepsilon_{k,y}^a} f(\tilde{y}_k^a), \quad (96)$$

and

$$\left(\rho \left\| C\tilde{y}_k^a - c + \frac{1}{\rho} \tilde{a}_k^a \right\|^2 + \frac{1}{\rho} \left\| -C^* \tilde{x}_k^a + \tilde{C}_\rho \tilde{b}_k^a \right\|_{(C^*C)^{-1}}^2 \right)^{1/2} \leq \frac{2d_0}{k}, \quad \varepsilon_{k,x}^a + \varepsilon_{k,y}^a \leq \frac{2(1 + 2\sqrt{2})d_0^2}{k}, \quad (97)$$

where d_0 is the distance of the initial point $(x_0, \tilde{y}_0) \in \mathcal{X} \times \mathcal{Y}$ to the solution set of (90) with respect to the inner product $\langle \cdot, \cdot \rangle_{\rho^{-1}} + \langle \cdot, \cdot \rangle_{\tilde{C}_\rho}$ in $\mathcal{X} \times \mathcal{Y}$, namely d_0 is the infimum of

$$\sqrt{\frac{1}{\rho} \|x_0 - x^*\|^2 + \rho \|\tilde{y}_0 - y^*\|_{\tilde{C}^*C}^2}$$

over the set of all solutions (x^*, y^*) of (90).

Proof. By Proposition 7.3(c), we know that $\{(x_k, \tilde{x}_k, \tilde{y}_k)\}$ is the sequence generated by applying the exact BD-HPE method to (91) with $\sigma = 1$ and $L_{xy} = 1$, and hence $\lambda = 1$. Hence, it follows from Lemma 5.1 and Theorem 5.2, relations (92), (93), and the definition of F and the fact that F is affine, that the last inequality in (97) holds,

$$\tilde{a}_k \in \rho \partial(g^* \circ B^*)(\tilde{x}_k), \quad \tilde{b}_k \in \tilde{C}_\rho^{-1} \partial f(\tilde{y}_k), \quad (98)$$

and

$$\left(\|\rho(C\tilde{y}_k^a - c) + \tilde{a}_k\|_{\rho^{-1}}^2 + \|\tilde{b}_k^a - \tilde{C}_\rho^{-1}C^*\tilde{x}_k^a\|_{\tilde{C}_\rho}^2 \right)^{1/2} \leq \frac{2d_0}{k}.$$

Now, using the definition of \tilde{C}_ρ and the norm induced by this operator, we easily see that the latter inequality is equivalent to the first inequality in (97). Moreover, (96) follows from (95) and (98) and Theorem 2.2(b). \square

We now translate the above result stated more in the context of the inclusion problem (90) and the exact BD-HPE method to the context of the original optimization problem (79) and the ADMM, respectively.

Theorem 7.5. *Consider the sequence $\{(x_k, \tilde{y}_k, \tilde{s}_k)\}$ generated by the ADMM and the sequences $\{\tilde{x}_k\}$ and $\{\tilde{b}_k\}$ defined according to (84) and (93). Moreover, consider the sequences $\{(\tilde{x}_k^a, \tilde{y}_k^a)\}$, $\{\tilde{b}_k^a\}$, $\{(\varepsilon_{k,x}^a, \varepsilon_{k,y}^a)\}$ defined in (94) and (95), and define for every $k \in \mathbb{N}$:*

$$\tilde{s}_k^a := \frac{1}{k} \sum_{i=1}^k \tilde{s}_i, \quad r_k^x := C\tilde{y}_k^a + D\tilde{s}_k^a - c, \quad r_k^y := \rho C^*C\tilde{b}_k^a - C^*\tilde{x}_k^a. \quad (99)$$

Then,

$$0 \in \partial g_{\varepsilon_{k,x}^a}(\tilde{s}_k^a) - D^*\tilde{x}_k^a, \quad r_k^y \in \partial_{\varepsilon_{k,y}^a} f(\tilde{y}_k^a) - C^*\tilde{x}_k^a, \quad (100)$$

and

$$\left(\rho \|r_k^x\|^2 + \frac{1}{\rho} \|r_k^y\|_{(C^*C)^{-1}}^2 \right)^{1/2} \leq \frac{2d_0}{k}, \quad \varepsilon_{k,x}^a + \varepsilon_{k,y}^a \leq \frac{2(1+2\sqrt{2})d_0^2}{k}, \quad (101)$$

where d_0 is defined as in Theorem 7.4.

Proof. First note that (84) and the definition of \tilde{a}_k in (93) imply that

$$\tilde{a}_k = \rho D\tilde{s}_k, \quad (102)$$

which together with (95) imply

$$\varepsilon_{k,x}^a = \frac{1}{k} \sum_{i=1}^k \left\langle \tilde{x}_i - \tilde{x}_k^a, \frac{1}{\rho} \tilde{a}_i \right\rangle = \frac{1}{k} \sum_{i=1}^k \langle \tilde{x}_i - \tilde{x}_k^a, D\tilde{s}_i \rangle = \frac{1}{k} \sum_{i=1}^k \langle D^*\tilde{x}_i - D^*\tilde{x}_k^a, \tilde{s}_i \rangle.$$

This identity, (85) and Theorem 2.2(b) then imply that

$$\tilde{s}_k^a \in \partial_{\varepsilon_{k,x}^a} g^*(D^*\tilde{x}_k^a),$$

from which the first inclusion in (100) follows. The second inclusion in (100) follows from the definition of \tilde{C}_ρ in Theorem 7.4, the second inclusion in (96) and the definition of r_k^y in (99). In addition, the estimates in (101) follow from (97), the definition of \tilde{C}_ρ , (99), (102), and the definition of \tilde{a}_k^a in (94). \square

We should emphasize that the analysis of this section requires both subproblems (82) and (83) to be solved exactly. We conjectured whether this assumption can be relaxed so as to allow the subproblems (or one of them) to be solved approximately.

8 Concluding remarks

In this paper, we have presented a general framework, namely the BD-HPE framework, of BD methods and obtained broad convergence rate results for it. As a consequence, we have derived for the first time convergence rate results for the classical ADMM by showing that it can be viewed as a special instance of the BD-HPE framework. We have also proposed new BD algorithms and derived their respective convergence rate results. These include a new splitting method for finding a zero of the sum of two maximal monotone operators and a new BD method based on Tseng's modified forward-backward splitting procedure. The analysis of the latter uses of an important feature of the BD-HPE framework, i.e., that it allows the one-block subproblems to be solved only approximately.

We also note that Nemirovski [17] and Nesterov [18] have previously established $\mathcal{O}(1/k)$ ergodic iteration-complexity bounds similar to the ones derived in this paper for specific algorithms to solve VIs and saddle-point problems. Hence, the various ergodic iteration-complexity bounds obtained in this paper extend their complexity bounds to a broader class of algorithms and problems other than VIs. Moreover, Monteiro and Svaiter [15, 16] have previously established pointwise complexity bounds for hemi-VIs and saddle point problems similar to the ones derived in this paper.

Finally, we make some remarks about a recent work of Chambolle and Pock [6] in light of the development in this paper. They have studied the monotone inclusion problem

$$0 \in K^*y + \partial g(x), \quad 0 \in -Kx + \partial f^*(y),$$

where K is a linear map and f, g are proper closed convex functions, and analyzed the convergence rate of an algorithm based on the exact evaluation of the resolvents of ∂g and ∂f^* (or ∂f). Their analysis, in contrast to ours, is heavily based on the fact that the above monotone inclusion problem is the optimality condition associated with the saddle point problem

$$\min_x \max_y \langle Kx, y \rangle + g(x) - f^*(y).$$

It can be shown, by means of a rescaling procedure, that their method and assumptions coincide with the exact BD-HPE method for the above inclusion problem (see Section 5.1) with the assumption that $\sigma < 1$. It should be noted however that, in contrast to our analysis, theirs does not deal with the extreme case of $\sigma = 1$ which, as mentioned earlier, is crucial to the analysis of the ADMM.

References

- [1] Regina S. Burachik, Alfredo N. Iusem, and B. F. Svaiter. Enlargement of monotone operators with applications to variational inequalities. *Set-Valued Anal.*, 5(2):159–180, 1997.
- [2] Regina S. Burachik, Claudia A. Sagastizábal, and B. F. Svaiter. ϵ -enlargements of maximal monotone operators: theory and applications. In *Reformulation: nonsmooth, piecewise smooth, semismooth and smoothing methods (Lausanne, 1997)*, volume 22 of *Appl. Optim.*, pages 25–43. Kluwer Acad. Publ., Dordrecht, 1999.
- [3] Regina Sandra Burachik and B. F. Svaiter. Maximal monotone operators, convex functions and a special family of enlargements. *Set-Valued Anal.*, 10(4):297–316, 2002.
- [4] Samuel Burer. Optimizing a polyhedral-semidefinite relaxation of completely positive programs. *Math. Program. Comput.*, 2(1):1–19, 2010.
- [5] Samuel Burer and Dieter Vandembussche. Solving lift-and-project relaxations of binary integer programs. *SIAM J. Optim.*, 16(3):726–750, 2006.
- [6] Antonin Chambolle and Thomas Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *Journal of Mathematical Imaging and Vision*, 40:120–145, 2011.

- [7] Jonathan Eckstein and B. F. Svaiter. A family of projective splitting methods for the sum of two maximal monotone operators. *Math. Program.*, 111(1-2, Ser. B):173–199, 2008. Published as IMPA Tech. Rep. A 238/2003 in 2003.
- [8] Jonathan Eckstein and B. F. Svaiter. General projective splitting methods for sums of maximal monotone operators. *SIAM J. Control Optim.*, 48(2):787–811, 2009.
- [9] F. Facchinei and J.-S. Pang. *Finite-dimensional variational inequalities and complementarity problems, Volume II*. Springer-Verlag, New York, 2003.
- [10] Daniel Gabay and Bertrand Mercier. A dual algorithm for the solution of nonlinear variational problems via finite element approximation. *Comput. Math. Appl.*, 2:17–40, 1976.
- [11] R. Glowinski and A. Marroco. Sur l’approximation, par éléments finis d’ordre un, et la résolution, par pénalisation-dualité, d’une classe de problèmes de Dirichlet non linéaires. 1975.
- [12] Florian Jarre and Franz Rendl. An augmented primal-dual method for linear conic programs. *SIAM J. Optim.*, 19(2):808–823, 2008.
- [13] G. M. Korpelevič. An extragradient method for finding saddle points and for other problems. *Èkonom. i Mat. Metody*, 12(4):747–756, 1976.
- [14] Jérôme Malick, Janez Povh, Franz Rendl, and Angelika Wiegele. Regularization methods for semidefinite programming. *SIAM J. Optim.*, 20(1):336–356, 2009.
- [15] R. D. C. Monteiro and B. F. Svaiter. On the complexity of the hybrid proximal projection method for the iterates and the ergodic mean. *SIAM Journal on Optimization*, 20:2755–2787, 2010.
- [16] R. D. C. Monteiro and B. F. Svaiter. Complexity of variants of Tseng’s modified F-B splitting and Korpelevich’s methods for hemi-variational inequalities with applications to saddle-point and convex optimization problems. *SIAM Journal on Optimization*, 21:1688–1720, 2012.
- [17] A. Nemirovski. Prox-method with rate of convergence $O(1/t)$ for variational inequalities with Lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 15:229–251, 2005.
- [18] Yurii Nesterov. Dual extrapolation and its applications to solving variational inequalities and related problems. *Math. Program.*, 109(2-3, Ser. B):319–344, 2007.
- [19] M. A. Noor. An extraresolvent method for monotone mixed variational inequalities. *Math. Comput. Modelling*, 29(3):95–100, 1999.
- [20] Adam Ouorou. Epsilon-proximal decomposition method. *Math. Program.*, 99(1, Ser. A):89–108, 2004.
- [21] J. Povh, F. Rendl, and A. Wiegele. A boundary point method to solve semidefinite programs. *Computing*, 78(3):277–286, 2006.
- [22] R. T. Rockafellar. On the maximal monotonicity of subdifferential mappings. *Pacific J. Math.*, 33:209–216, 1970.
- [23] R. Tyrrell Rockafellar. Monotone operators and the proximal point algorithm. *SIAM J. Control Optimization*, 14(5):877–898, 1976.
- [24] M. V. Solodov. A class of decomposition methods for convex optimization and monotone variational inclusions via the hybrid inexact proximal point framework. *Optim. Methods Softw.*, 19(5):557–575, 2004.
- [25] M. V. Solodov and B. F. Svaiter. A hybrid approximate extragradient-proximal point algorithm using the enlargement of a maximal monotone operator. *Set-Valued Anal.*, 7(4):323–345, 1999.

- [26] M. V. Solodov and B. F. Svaiter. A hybrid projection-proximal point algorithm. *J. Convex Anal.*, 6(1):59–70, 1999.
- [27] M. V. Solodov and B. F. Svaiter. An inexact hybrid generalized proximal point algorithm and some new results on the theory of Bregman functions. *Math. Oper. Res.*, 25(2):214–230, 2000.
- [28] M. V. Solodov and B. F. Svaiter. A unified framework for some inexact proximal point algorithms. *Numer. Funct. Anal. Optim.*, 22(7-8):1013–1035, 2001.
- [29] B. F. Svaiter. A family of enlargements of maximal monotone operators. *Set-Valued Anal.*, 8(4):311–328, 2000.
- [30] Paul Tseng. A modified forward-backward splitting method for maximal monotone mappings. *SIAM J. Control Optim.*, 38(2):431–446 (electronic), 2000.