SCIENTIFIC REPORTS

Received: 4 August 2017 Accepted: 8 November 2017 Published online: 17 November 2017

OPEN Ketamine's antidepressant effect is mediated by energy metabolism and antioxidant defense system

Katja Weckmann^{1,6}, Michael J. Deery², Julie A. Howard², Renata Feret², John M. Asara³, Frederik Dethloff¹, Michaela D. Filiou⁴, Jamie Iannace¹, Christiana Labermaier¹, Giuseppina Maccarrone¹, Christian Webhofer¹, Larysa Teplytska¹, Kathryn Lilley², Marianne B. Müller⁵ & Christoph W. Turck¹

Fewer than 50% of all patients with major depressive disorder (MDD) treated with currently available antidepressants (ADs) show full remission. Moreover, about one third of the patients suffering from MDD does not respond to conventional ADs and develop treatment-resistant depression (TRD). Ketamine, a non-competitive, voltage-dependent N-Methyl-D-aspartate receptor (NMDAR) antagonist, has been shown to have a rapid antidepressant effect, especially in patients suffering from TRD. Hippocampi of ketamine-treated mice were analysed by metabolome and proteome profiling to delineate ketamine treatment-affected molecular pathways and biosignatures. Our data implicate mitochondrial energy metabolism and the antioxidant defense system as downstream effectors of the ketamine response. Specifically, ketamine tended to downregulate the adenosine triphosphate (ATP)/adenosine diphosphate (ADP) metabolite ratio which strongly correlated with forced swim test (FST) floating time. Furthermore, we found increased levels of enzymes that are part of the 'oxidative phosphorylation' (OXPHOS) pathway. Our study also suggests that ketamine causes less protein damage by rapidly decreasing reactive oxygen species (ROS) production and lend further support to the hypothesis that mitochondria have a critical role for mediating antidepressant action including the rapid ketamine response.

Mood disorders including major depressive disorder (MDD) are a leading cause of disability worldwide^{1,2}. The prevalence for mood disorders has been estimated at 21 million people across 28 European countries with an annual cost of 106 billion Euro^{3,4}. Globally, approximately 350 million people suffer from MDD, with women more likely to be affected than men^{5,6}. The currently available antidepressants (ADs) require chronic long-term treatment of several weeks before a therapeutic effect is achieved^{7,8}. Less than 50% of all patients treated with current ADs show full remission⁹. Moreover, about one third of the patients suffering from MDD do not respond to conventional ADs and suffer from treatment-resistant depression (TRD)10. Fifteen percent of severely depressed individuals commit suicide, the second leading cause of death for ages 15- to 29-years⁶.

In 2000 Berman et al. demonstrated for the first time a rapid antidepressant effect of a low dose of ketamine within 2 h and 4 h of treatment that lasts for up to 10 days¹¹. Ketamine is a non-competitive, voltage-dependent N-Methyl-D-aspartate receptor (NMDAR) antagonist¹²⁻¹⁷. The antidepressant effect of ketamine is also observed in patients suffering from TRD^{18,19}. Moreover, ketamine diminishes suicidal ideation²⁰. However, psychomimetic side effects of ketamine have prevented its use as a first-line drug²¹⁻²⁵.

¹Max Planck Institute of Psychiatry, Department of Translational Research in Psychiatry, Munich, Germany. ²Cambridge Centre for Proteomics, Cambridge System Biology Centre, University of Cambridge, Cambridge, UK. ³Division of Signal Transduction, Beth Israel Deaconess Medical Center and Department of Medicine, Harvard Medical School, Boston, USA. ⁴Max Planck Institute of Psychiatry, Department of Stress Neurobiology and Neurogenetics, Munich, Germany. ⁵Experimental Psychiatry, Department of Psychiatry and Psychotherapy & Focus Program Translational Neuroscience, Johannes Gutenberg University Medical Center, Mainz, Germany. ⁶Present address: Institute of Pathobiochemistry, Johannes Gutenberg University, Medical School, Mainz, Germany. Correspondence and requests for materials should be addressed to M.B.M. (email: Marianne.Mueller@unimedizinmainz.de) or C.W.T. (email: turck@psych.mpg.de)

A low dose of ketamine was initially reported to act through the activation of the mammalian target of rapamycin complex 1 (mTORC1) causing an increase of synaptic protein levels and an elevated number of spines and neuronal activity in the medial prefrontal cortex of rats²⁶. Furthermore, increased levels of postsynaptic proteins required for formation, maturation, and function of new spines were observed 2 h after ketamine treatment and lasted up to 72 h²⁶. More recent studies of ketamine's antidepressant effects implicate a ketamine metabolite as the actual active compound. Results from studies in mice treated with the ketamine metabolite (2 R, 6 R)-hydroxynorketamine (HNK) suggest that the antidepressant-like effect of the drug is independent of NMDAR binding and implicate the α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPAR) and its signaling pathway to be critical for the activity²⁷.

In previous studies we have shown an important role of mitochondrial energy metabolism for ketamine's antidepressant mode of action. A time-dependent metabolomics profiling study implicated 'glycolysis' and the 'citrate cycle' pathways. In the present paper we report on an extension of our analyses on ketamine's fast acting antidepressant activity that further support a critical role of mitochondrial pathways impacting cellular oxidative stress defense mechanisms²⁸.

Results

Our previous study on ketamine's mode of action suggested an involvement of mitochondrial energy metabolism, a finding that is also supported by its link to glutamate neurotransmission. Glutamate is metabolized from α -ketoglutarate, a 'citrate cycle' metabolite, and can be recycled through this pathway. Interestingly, glutamate neurotransmission also stimulates glucose utilization and 'glycolysis', the upstream pathway of the OXPHOS pathway. Therefore, we investigated whether ketamine's fast antidepressant-like effects are reflected by metabolite ratios that are part of the 'citrate cycle' and the connected OXPHOS pathway, including ATP/ADP, NADH/NAD and GTP/GDP ratios. In addition, we correlated these metabolites with antidepressant-like behavior using the FST floating time as a readout. Interestingly, the ATP/ADP metabolite ratio tended to be decreased 2 h and 24 h after a single injection of a low dose of ketamine. In addition, ATP and ADP levels significantly and strongly correlated with the FST floating time at 24 h (Fig. 1A). ATP is produced from ADP by OXPHOS complex V and NADH serves as an electron donor for OXPHOS complex I^{28,29}. The NADH/NAD ratio neither showed a significant difference nor a significant correlation with the FST floating time (Fig. 1B). GTP is synthesized from GDP as a by-product of succinate-CoA to succinate conversion in the 'citrate cycle'. Succinate is an electron donor for the OXPHOS complex II subunit A that we have previously shown to be elevated 2 h after ketamine treatment²⁸. We found that the GTP/GDP metabolite ratio is indeed significantly elevated 2h upon ketamine treatment and tended to be downregulated at the 24h time point. In addition, GTP levels significantly correlated with FST floating time at both time points (Fig. 1C).

Based on the above results we next wanted to investigate whether AMP-activated protein Kinase (AMPK) is affected by ketamine treatment. AMPK is the major energy status sensor that promotes catabolic pathways that generate ATP and inhibits anabolic pathways that consume ATP. A low ATP/ADP ratio activates AMPK resulting in its phosphorylation, leads to decreased activities of pathways that consume energy (anabolic pathways) and increased activities of pathways that produce energy (catabolic pathways)³⁰⁻³⁶. Previous data from a study in rats suggest that ketamine's antidepressant-like effects are mediated by an increased anabolic rate, thereby inducing cell growth and differentiation²⁶. The result of an elevated anabolism is a higher energy demand. This is supported by our results of an increased pAMPK/AMPK ratio and decreased ATP/ADP ratio and decreased ATP levels at the 24 h time point²⁸. While total AMPK protein levels are not significantly changed, the phosphorylated form of the protein, pAMPK, shows a statistically significant increase at the 24 h time point (Fig. 2, Supplementary Figure 1A and B).

We next investigated whether the observed energy status alterations upon ketamine treatment are associated with changes in the expression of proteins related to cellular energy metabolism. Time-dependent quantitative proteomics profiling analyses were performed 2 h and 24 h after a single injection of a low dose of ketamine. Five mice per time point and treatment group were chosen and MF and CF from ketamine- and vehicle-treated mice were isolated from the hippocampus and quantified using ¹⁵N metabolically labelled reference proteins by LC-MS/MS analyses.

After proteomics data processing and missing value imputation by k-nearest neighbor (KNN), a total of 889 hippocampal MF and 1173 CF proteins were processed by statistical analyses for the 2h time point. For the 24h time point 929 MF and 1091 CF proteins were used for further statistical analyses (Supplemental Table S1). The protein profiles allowed a separation of ketamine- and vehicle-treated mice using multivariate PLS-DA for both time points (Fig. 3). The quality criteria of PLS-DA models were assessed by accuracy, R^2 , and Q^2 values. They indicate good models for the 2h time point for MF and CF proteins of ketamine- compared to vehicle-treated animals. The PLS-DA models were only weak for the 24h time point, which might result in false-positives (Supplemental Table S2). To increase the robustness of the statistical analyses we combined the VIP-score of the PLS-DA with SAM analyses in order to obtain a reliable list of MF and CF proteins that significantly contribute to ketamine's antidepressant effect. This resulted in 61 MF and 95 CF proteins that were statistically significant altered at the 2h time point when comparing ketamine- to vehicle-treated mice (Supplemental Table S3). Consistent with previous analyses by us we found proteins that are part of major energy metabolism pathways with altered levels after ketamine treatment²⁸. The most significant alterations were detected already 2h after a single injection of ketamine, in line with the rapid antidepressant effect of ketamine. Subsequent pathway enrichment analyses highlighted altered OXPHOS, ⁵purine metabolism, 'fructose and mannose metabolism,' proteasome', 'protein processing in endoplasmic reticulum' and 'ribosome' (Table 1). We found that already at the 2 h time point a low dose of ketamine resulted in increased levels of several OXPHOS proteins including complex I NADH dehydrogenase [ubiquinone] flavoprotein 2 (Ndufv2), cytochrome c1, heme protein (Cyc1), and complex I assembly factor (Tmem126B) and decreased levels of NADH dehydrogenase [ubiquinone] complex I and



Figure 1. (A) Time-dependent hippocampal analysis of adenosine triphosphate (ATP)/adenosine diphosphate (ADP) metabolite ratio and correlation analyses of ATP (2h, $p \ge 0.05$ and 24h, $p \le 0.05$) and ADP (2h, $p \ge 0.05$ and 24h, $p \le 0.05$) with the forced swim test (FST) floating time. (B) Time-dependent hippocampal analysis of nicotinamide adenine dinucleotide (reduced) (NADH)/nicotinamide adenine dinucleotide (oxidized) (NAD) metabolite ratio. (C) Time-dependent hippocampal analysis of guanosine triphosphate (GTP)/guanosine diphosphate (GDP) ratio and correlation analyses of GTP (2h and 24h, $p \le 0.05$) and GDP (2h and 24h, $p \ge 0.05$) after a single injection of a low dose of ketamine with FST floating time. The results from ketamine-treated animals are normalized to the ones obtained from vehicle treatment (mean value = 1). N = 5 mice per group and time point. *p ≤ 0.05 . P-values were determined by Student's t-test and significance analysis of microarrays and other omic-datasets (SAM). Error bars represent s.e.m. The correlation coefficient r was calculated by Pearson. The linear regression line is only shown for significant ($p \le 0.05$) correlation coefficients.

assembly factor 7 (Ndufaf7) (Supplemental Table S3 and Fig. 4A). Furthermore, Ndufv2, Cyc1, Ndufaf7 and Tmem126B protein levels significantly correlated with floating time in the FST (Fig. 4B). These results are consistent with our previous metabolomics profiling study that had also indicated mitochondrial energy metabolism, specifically OXPHOS, to be affected by ketamine treatment²⁸.

Increased OXPHOS activity can result in higher ROS levels. The phenomenon referred to as oxidative stress causes an imbalance of ROS and the cellular defense system made up of antioxidant molecules and enzymes like peroxiredoxins (Prdxs) that contribute to the antioxidant capacity of the cell^{37–41}. We therefore next analysed ket-amine's effects on oxidative stress. Cytoplasmic Prdx1 and mitochondrial Prdx3 protein levels were significantly lower at the 2 h and 72 h time points, respectively (Fig. 5A, Supplementary Figure 1C). In addition, we found decreased total antioxidant capacity (TAC) 72 h after ketamine injection (Fig. 5B, supplementary Figure 1D). ROS radicals promoting oxidative stress result in cellular damage by reacting with deoxyribonucleic acid (DNA), ribonucleic acid (RNA), lipids, and proteins. Protein modifications by ROS become apparent through the introduction of carbonyl groups into protein side chains. Since ROS are difficult to measure due to their highly reactive nature we assessed their presence indirectly by the analysis of carbonylated (damaged) proteins³⁷. Our results indicate that ketamine results in statistically significant reduction of protein carbonylation 72 h after ketamine treatment (Fig. 5C).



Figure 2. Time-dependent Western blot analysis of AMP-activated protein kinase (AMPK) and phosphorylated AMPK (pAMPK) levels and pAMPK/AMPK ratios reflecting enzyme activation. The results from ketamine-treated animals are normalised to the ones obtained from vehicle treatment (mean value = 1). N = 5 mice per group and time point. *p \leq 0.10, *p \leq 0.05. P-values were determined by Student's t-test. Error bars represent s.e.m.

Discussion

Ketamine treatment reduced the ATP/ADP metabolite ratio at 2 h and 24 h and both ATP and ADP levels significantly correlated with FST floating time at 24 h suggesting an energy deficit. This is consistent with our previous results where we found a significant reduction of ATP levels at $24 h^{28}$. Also in line with this is our observation of an increased AMPK activation following ketamine treatment. In the current study we observed several metabolite level and metabolite ratio changes that are part of the 'citrate cycle' and 'glycolysis' as early as 2 h after ketamine administration, reflecting the fast antidepressant effect of the drug. The 'citrate cycle' is connected to the OXPHOS pathway which produces energy in the form of ATP. 'Glycolysis' is a major catabolic cytoplasmic pathway where glucose is degraded to pyruvate, which is further metabolized by pyruvate dehydrogenase complex to acetyl-CoA that enters the 'citrate cycle'. The latter is a mitochondrial matrix pathway that is coupled to the OXPHOS pathway²⁹. We observed metabolite ratio changes that are either due to protein level or protein activity alterations of OXPHOS complex II enzymatic subunit A that processes these metabolites^{28,42}. Our previous time-dependent hippocampal metabolomics profiling analyses indicated statistically significant biomolecular changes that are part of several pathways including the 'citrate cycle' and 'glycolysis'28. Mitochondrial abnormalities including alterations in energy metabolism have been implicated in the pathobiology of affective disorders⁴³⁻⁴⁵. Taken together, these findings provide further evidence for an important role of mitochondrial energy metabolism in ketamine's mode of action to generate ATP which compensates for the increased energy demand at activated synapses. We propose that ketamine treatment causes a cellular energy deficit by first promoting anabolic processes (that consume ATP) through the activation of mTORC1. This is counterbalanced in a second phase by the activation of AMPK that promotes catabolic pathways (to generate ATP).

Ketamine affects glutamate neurotransmission and is tightly linked to mitochondrial energy metabolism. Upon release, glutamate is taken up by glial cells and subsequently metabolized to glutamine. In addition, glutamate is metabolized from α -ketoglutarate, a 'citrate cycle' metabolite, and can also be recycled through this pathway. Interestingly, glutamate neurotransmission also stimulates glucose utilization and 'glycolysis', the upstream pathway of the OXPHOS pathway⁴⁶. Magnetic resonance spectroscopy analysis of rat medial prefrontal cortex with ¹³C-labeled glucose demonstrated ¹³C-enrichment of glutamate, glutamine and GABA upon ketamine treatment, reflecting neurotransmitter release coupled to glycolytic and 'citrate cycle' activity. In addition, other studies have shown elevated medial prefrontal cortex metabolism and an involvement of mitochondria in MDD pathobiology^{43,47-49}. Mitochondrial respiration in peripheral blood mononuclear cells from depressed patients is lower compared to cells from control subjects and correlates with depressive symptom severity⁵⁰. Furthermore, mitochondrial disorders are often comorbid with psychiatric disorders including MDD and patients frequently suffer from TRD. These patients have a primary impairment of mitochondrial functioning through either nuclear or mitochondrial DNA mutations. Mitochondrial impairment mostly affects high energy consuming organs like muscles and brain. The mitochondrial disorder is often recognized only later in life of MDD patients⁵¹⁻⁵³. Previous studies have connected the antidepressant response with drugs elevating ATP levels. Spectroscopy data have shown decreased NPT levels (mainly ATP) in the brain of depressed patients. Another magnetic resonance spectroscopy study could demonstrate lower NPT levels (mainly ATP) in fluoxetine responders compared with non-responders. Previous data from a study in rats have suggested that the antidepressant-like effect observed after a single injection of ketamine is mediated by an increased anabolic rate-mediating cell growth and differentiation, processes that are highly energy consuming^{54,55}. The high ketamine treatment efficacy for TRD patients might be due to the drug's beneficial effects on mitochondrial energy metabolism. We submit that mitochondria



▲ Ketamine-treated

+ vehicle-treated

Figure 3. Multivariate partial least squares-discriminant analysis (PLS-DA) of (**A**) membrane-associated (MF2h) and (**B**) cytoplasmic (CF2h) proteins at the 2h time point and (**C**) membrane-associated (MF24h) and (**D**) cytoplasmic (CF24h) proteins 24h after a single injection of a low dose of ketamine (3 mgkg^{-1}) or vehicle using all quantified proteins. N = 5 mice per group and time point.

.....

Pathway	Number of proteins	p-value	FDR
Protein processing in endoplasmic reticulum	6	0,001	0,084
Purine metabolism	6	0,001	0,084
Ribosome	5	0,002	0,084
Fructose and mannose metabolism	3	0,002	0,084
Oxidative phosphorylation (OXPHOS)	5	0,002	0,084
Proteasome	3	0,003	0,100

Table 1. Pathway enrichment analysis of significantly altered hippocampal proteins (PLS-DA, variable influence of projection (VIP)-score 1.0, significance analysis of microarrays and other omic-datasets (SAM), false-discovery rate (FDR) 0.10, and SAM q 0.10) 2 h after a single injection of a low dose of ketamine (3 mg kg⁻¹). N = 5 mice per group and time point.

.....

play a critical role in the development of MDD as well as AD treatment response, at least in a subset of patients with a defined symptomatology.

In order to better understand the biomolecular pathways and cellular mechanisms underlying ketamine's mode of action, a time-dependent proteomics profiling analysis was also performed. We identified several



Figure 4. (A) Hippocampal OXPHOS protein levels 2 h after a single injection of a low dose of ketamine (3 mg kg^{-1}) . Complex I NADH dehydrogenase [ubiquinone] flavoprotein 2 (Ndufv2), cytochrome c1, heme protein (Cyc1), and complex I assembly factor (Tmem126B) levels were determined by quantitative proteomics profiling. (B) Correlation analyses of Ndufv2, Cyc1, Ndufaf7, and Tmem126B protein levels with forced swim test (FST) floating time 2 h after a single injection of ketamine ($p \le 0.05$). The results from ketamine-treated animals are normalised to the ones obtained from vehicle treatment (mean value = 1). N = 5 mice per group and time point, ***p ≤ 0.001 . P-values were determined by Student's t-test and significance analysis of microarrays and other omic-datasets (SAM). Error bars represent s.e.m. The correlation coefficient r was calculated by Pearson. The linear regression line is only shown for significant ($p \le 0.05$) correlation coefficients.

pathways that were affected and associated with mitochondrial energy metabolism. The PLS-DA showed a time-dependent separation of ketamine- and vehicle-treated mice that was evident by significant protein level alterations and overrepresented pathways. Of the pathways that we found to be affected by ketamine treatment we focused our attention on OXPHOS, in line with results previously obtained by us that had implicated mitochondrial energy metabolism to play a major role in the drug's mode of action. Ketamine affects the OXPHOS pathway as is evident by several complex I protein levels (Ndufv2, Cyc1, Ndufaf7 and Tmem126B) that correlate with FST floating time in a treatment time-dependent manner.

Taken together, we propose a mechanism by which ketamine stimulates glutamatergic neurotransmission and neuronal activity, ultimately resulting in a long-term potentiation (LTP)-like process. This activation results in the activation of mTORC1. The mTORC1 mediated anabolic processes include protein synthesis and mitochondrial biogenesis within 30 minutes and 1 h upon ketamine treatment. Pretreatment with rapamycin, an mTOR inhibitor, prevents ketamine's antidepressant-like effect²⁶. Furthermore, other studies indicated that mTOR inhibition decreases mitochondrial respiration, whereas mTOR hyperactivation elevates the expression of oxidative metabolism genes and mitochondrial DNA copy number⁵⁶⁻⁵⁸. Hence, ketamine may not only increase protein translation through mTORC1, but also mitochondrial biogenesis. LTP-like processes require energy in the form of ATP followed by AMPK activation, which we found in the present study, in order to promote catabolic processes to generate ATP. Ketamine's antidepressant-like effects are known to last for one to two weeks. This long-term effect can be explained by the observed LTP-like process. Late phase LTP with spine formation and strengthening of synaptic connections are known to last from days to weeks. Increased OXPHOS activity can result in elevated ROS levels, a phenomenon referred to as oxidative stress, which causes protein, DNA, RNA and lipid modifications and damage. We found that protein carbonylation decreases over time until it reaches significance at the 72 h time point. This can be explained by either decreased ROS levels or alternatively by an activation of a cellular defense system that captures ROS. The cellular defense system includes antioxidant molecules and enzymes such as Prdxs that combat ROS. In addition, a protein quality control system that degrades damaged proteins counteracts oxidative stress^{38–41,59,60}. We found that total antioxidant capacity (TAC) and cytoplasmic Prdx1 levels were lower at the 2 h and 72 h time points and mitochondrial Prdx3 levels lower at the 72 h time point in ketamine- vs. vehicle-treated mice. The reduced protein carbonylation we observed in ketamine- vs. vehicle-treated mice could be indicative of increased protein quality control processes in the cell. This is supported by our proteomic data that indicate an upregulation of the 'proteasome' pathway at the 2h time point. Taken together, our combined proteomic results suggest that ketamine administration in mice leads to increased degradation of damaged proteins and homeostatic redox adjustment.

The close connection between the glutamatergic system and mitochondrial energy metabolism make a compelling case for the here observed facilitative effects of energy metabolism and the antioxidant defense system following ketamine administration. Further experiments including blocking pathway enzymes and receptors



Figure 5. Hippocampal analyses of the antioxidant defense system after a single injection of a low dose of ketamine (3 mg kg^{-1}) . (A) Time-dependent Western blot analysis of peroxiredoxin 1 (Prdx1) and peroxiredoxin 3 (Prdx3). N = 5 mice per group and time point. (B) Time-dependent total antioxidant capacity (TAC) analysis. N = 5 mice per group and time point. (C) OxyBlot analysis of the time-dependent protein damage by protein carbonylation in the hippocampus after ketamine treatment. N = 4 mice per group and time point. The results from ketamine-treated animals are normalised to the ones obtained from vehicle treatment (mean value = 1). *p ≤ 0.05 , **p ≤ 0.01 . P-values were determined by Student's t-test. Error bars represent s.e.m.

including OXPHOS complex I and AMPK that we found to be involved as well as different ketamine doses are required to confirm these findings. We only assessed ketamine's behavioral effect using FST floating time and did not carry out any additional behavioral analyses like the learned helplessness, chronic mild stress and novelty

suppressed feeding tests that have been previously reported by others. However, using several consecutive behavioral stress assays will in all likelihood affect the animals' metabolome and proteome and therefore skew the data.

Methods

Animals and ketamine treatment. The animals and ketamine treatment were housed and carried out as previously described²⁸. The experiments were performed in accordance with European Communities Council Directive 86/609/EEC. The protocols were approved by the committee for the Care and Use of Laboratory Animals of the Government of Upper Bavaria, Germany.

Forced swim test. The forced swim test and floating time analysis were carried out as previously described²⁸.

Isolation of cytoplasmic protein fraction (CF) and membrane-associated protein fraction **(MF).** CF and MF were prepared by repeated tissue homogenization and extraction of non-MF proteins and solubilization of MF proteins with sodium dodecyl sulfate (SDS). Tissues were homogenized for 30 s in 2 M NaCl, 10 mM Hepes/NaOH, pH 7.4, containing 1 mM ethylenediaminetetraacetic acid (EDTA), phosphatase inhibitor cocktail 2 and 3 (Sigma-Aldrich, Munich, Germany), protease inhibitor cocktail tablets 'cOmplete' (Roche Diagnostics, Mannheim, Germany), incubated for 10 min and homogenized for 30 s and further with a ultrasonicator for 3×10 s on ice. Homogenates were centrifuged at 16100 g at 4 °C for 20 min with the supernatant containing CF proteins. Pellets were rehomogenized in 0.1 M Na₂CO₃ and 1 mM EDTA containing 1 mM EDTA, phosphatase inhibitor cocktail 2 and 3 (Sigma-Aldrich, Munich, Germany), protease inhibitor cocktail tablets 'cOmplete' (Roche Diagnostics, Mannheim, Germany), pH 11.3, mixed at 4 °C for 30 min and collected by centrifugation (16100 g at 4 °C for 20 min). Subsequently, pellets were extracted with 5 M urea, 100 mM NaCl, 10 mM Hepes, pH 7.4 and 1 mM EDTA containing 1 mM EDTA, phosphatase inhibitor cocktail 2 and 3 (Sigma-Aldrich, Munich, Germany), protease inhibitor cocktail tablets 'cOmplete' (Roche Diagnostics, Mannheim, Germany) and then washed twice with 0.1 M Tris-HCl, containing 1 mM EDTA, phosphatase inhibitor cocktail 2 and 3 (Sigma-Aldrich, Munich, Germany), protease inhibitor cocktail tablets 'cOmplete' (Roche Diagnostics, Mannheim, Germany) pH 7.6. Pellets were solubilized in 2% SDS, 50 mM dithiothreitol (DTT) and 0.1 M Tris-HCl, containing 1 mM EDTA, phosphatase inhibitor cocktail 2 and 3 (Sigma-Aldrich, Munich, Germany), protease inhibitor cocktail tablets 'cOmplete' (Roche Diagnostics, Mannheim, Germany) pH 7.6, at 90 °C for 1 min and stored at -20 °C until further analysis.

Western blot. Hippocampal MF and CF proteins from 8-week-old male C57BL/6 mice treated with ketamine for 2h, 14h, 24h and 72h were fractionated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and Western blot was performed based on standard protocols. After electrophoresis, proteins were transferred to PVDF membranes (Immobilon-P, Millipore, Billerica, USA). Primary antibodies were against adenosinmonophosphate-activated protein kinase (AMPK, Abcam, Cambridge, UK), phosphorylated AMPK (pAMPK, Cell Signaling, Merck, Darmstadt, Germany), peroxiredoxin 1 (Prdx1, Abcam, Cambridge, UK), peroxiredoxin 3 (Prdx3, Abcam, Cambridge, UK). Anti-rabbit, anti-mouse and anti-goat ECL horseradish peroxidase-linked secondary antibodies (GE Healthcare Life Sciences, Little Chalfont, Buckinghamshire, UK) were used. The densitometric analyses were performed with the Image Lab software (Bio-Rad, Munich, Germany).

Protein oxidation analysis. The OxyBlotTM Kit (Merck Millipore, Darmstadt, Germany) was used according to the manufacturer's instructions.

Total antioxidant capacity (TAC). The TAC Assay Kit (BioVision, Milpitas, USA) was used according to the manufacturer's instructions. Absorbance was measured at 570 nm using a plate reader (iMarkTm Microplate Absorbance Reader, Bio-Rad, Munich, Germany).

Protein sample preparation for quantitative proteomics analyses. Hippocampal CF and MF proteins were mixed 1:1 with CF and MF ¹⁵N-labeled protein standards, respectively. The latter were derived from C57BL/6 mice that had been fed a ¹⁵N-labelled diet (Silantes, Munich, Germany) for 12 weeks. $50 \mu g$ of the ¹⁴N/¹⁵N protein mixture were separated by SDS-PAGE. Separated proteins were stained with Coomassie Brilliant Blue for 20 min and destained overnight. Each gel lane was cut into 16 2.5 mm slices per biological replicate and each slice further cut into smaller pieces.

The gel pieces were then covered with $100 \,\mu$ l of $25 \,\text{mM} \,\text{NH}_4\text{HCO}_3/50\%$ acetonitrile (ACN) for complete destaining and mixed for $10 \,\text{min}$ at room temperature. The supernatant was discarded and this step repeated twice. Proteins were reduced with $75 \,\text{mL} \,1x \,\text{DTT}/25 \,\text{mM} \,\text{NH}_4\text{HCO}_3$ and incubated at $56\,^\circ\text{C}$ for $30 \,\text{min}$ in the dark. The supernatant was discarded and for alkylation $100 \,\mu$ l iodoacetamide (IAM) was added to the gel pieces and mixed for $30 \,\text{min}$ at room temperature. The supernatant was discarded and the gel pieces washed twice with $100 \,\mu$ l $25 \,\text{mM} \,\text{Na}_4\text{HCO}_3/50\%$ ACN and incubated for $10 \,\text{min}$ at room temperature. The supernatant was discarded and gel pieces dried for $20 \,\text{min}$ at room temperature. Proteins were digested with $50 \,\mu$ l trypsin solution ($5 \,\text{mg}/\mu$ l trypsin/ $25 \,\text{mM} \,\text{NH}_4\text{HCO}_3$) overnight at $37\,^\circ\text{C}$. Peptides were extracted from the gel pieces by incubation in $50 \,\mu$ l 2% formic acid (FA)/50% ACN for $20 \,\text{min}$ at $37\,^\circ\text{C}$ followed by $5 \,\text{min}$ sonication. This step was repeated twice with $50 \,\mu$ L of $1\% \,\text{FA}/50\% \,\text{ACN}$. The supernatants were then combined and dried (SpeedVac Plus, SC 210 A, Savant, Hyannis, MA, USA). The pellet was stored at $-20\,^\circ\text{C}$.

Liquid chromatography tandem mass spectrometry (LC-MS/MS). Hippocampal MF and CF proteins were identified and quantified with a Dionex Ultimate 3000 RSLC nanoUPLC (Thermo Fisher Scientific, Waltham, USA) online coupled to a QExactiveTM OrbitrapTM mass spectrometer (Thermo Fisher Scientific, Waltham, USA). Separation of peptides was performed by reversed-phase chromatography at a flow rate of 300 nl/ min (Thermo Scientific PepMap C18, 2 µm particle size, 100 A pore size, $75 \mu m \times 50 \text{ cm}$). Peptides were loaded onto a pre-column (Thermo Scientific PepMap 100 C18. 5 µm particle size, 100 A pore size, $300 \mu m \times 5 \text{ mm}$) in 0.1% FA for 3 min at a flow rate of 10μ l/min. Chromatography was performed using the following solvents: solvent A water, 0.1% FA, solvent B 80% ACN, 20% water, 0.1% FA. A linear gradient of 2–40% B for 30 min was used. The LC eluant was sprayed into the mass spectrometer by means of an easy-spray source (Thermo Fisher Scientific, Waltham, USA). All *m/z* values of eluting ions were measured in the Orbitrap mass analyzer, set at a resolution of 70000. Data dependent scans (Top 20) were employed to automatically isolate and generate fragment ions by high energy collisional dissociation in the quadrupole mass analyser and measurement of the resulting fragment ions was performed in the Orbitrap analyzer, set at a resolution of 17500. Peptide ions with charge states of 2+ to 4+ and above were selected for fragmentation.

Orbitrap raw files were converted to mzXML files using MSConvert software. The in-house software package iSPY, which was adapted from an earlier version of a peptide quantitation program known as iTracker, was used to identify and quantify peptides^{61,62}. The software was used to convert mzXML to mgf files that were then imported into Mascot and searched against the SwissProt Mouse database (November 2013) and a decoy database. The search was then performed using TAXONOMY = Mus. The databases were searched using the following settings: variable modifications of carbamidomethyl, oxidation), 20 ppm peptide tolerance, 0.1 Da MS/MS tolerance, 2 missed cleavages and peptide charge states of +2, +3, or +4. In iSPY, Mascot dat output files were run through Percolator for improved identification. Non-unique peptides were discarded. Only peptides with a protein type 1 error of less than 0.01 were kept in the final dataset. The heavy and light peak intensities for each peptide were calculated in iSPY using retention time and sequence information from the MS1 spectra and Mascot search, respectively. Briefly, the intensities for a pre-specified number of isotopomeric peaks were calculated by scanning through a retention time window spanning a set distance on either side of the maximum intensity value. The ¹⁴N and ¹⁵N peptide isotopic peaks from the MS1 dataset were used to compare the theoretical mass difference between the heavy and light peptides, and the typical isotopic distribution patterns. Only quantifiable peptides for which both heavy and a light peak intensities were identified in five replicates were included in the dataset.

Isolation of polar metabolites from mouse tissue and metabolomics analyses. Samples were obtained, measured and analysed as previously described^{28,63}.

Statistics and data analyses. *Identification of significant metabolite and protein level alterations.* Metabolite intensities as well as protein ratios were median-normalised and auto-scaled for statistical analysis. Significant protein level changes 2 h and 24 h after ketamine treatment were identified by multivariate partial least squares-discriminant analyses (PLS-DA) and high-dimensional feature selection significance analysis of microarrays and other omic-datasets (SAM) using MetaboAnalyst^{64,65}. The quality of the PLS-DA models were assessed for R², Q² and accuracy values with variable influence of projection (VIP)-score ≥ 1.0 and for SAM with q ≤ 0.10 and false discovery rate (FDR) $\leq 0.10^{66}$. We improved robustness of our data analyses and increased confidence in significantly altered metabolites and proteins as well as in the subsequent analyses of overrepresented KEGG pathways by only considering the overlap between the two different statistical methods.

Identification of significantly enriched pathways. Pathway analyses were performed using MetaboAnalyst⁶⁴ applying a hypergeometric algorithm for overrepresentation analysis and relative-betweeness centrality for pathway topology analysis. Pathways were considered affected if they were significantly enriched for all significantly altered metabolites ($P_{Holm-corrected} \leq 0.05$). Proteomic pathway enrichment was assessed by String with an FDR $\leq 0.10^{67,68}$.

Calculation of metabolite pair ratios. The metabolite pair ratios were calculated as described²⁸.

All data generated or analysed during this study are included in this published article (and its Supplementary Information files).

References

- 1. Olesen, J. et al. Consensus document on European brain research. J Neurol Neurosurg Psychiatry 77(Suppl 1), i1-49 (2006).
- Murray, C. J. & Lopez, A. D. Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. Lancet 349, 1498–1504, https://doi.org/10.1016/S0140-6736(96)07492-2 (1997).
- 3. Agid, Y. et al. How can drug discovery for psychiatric disorders be improved? Nat Rev Drug Discov 6, 189–201, https://doi. org/10.1038/nrd2217 (2007).
- Kessler, R. C. et al. The global burden of mental disorders: an update from the WHO World Mental Health (WMH) surveys. Epidemiol Psichiatr Soc 18, 23–33 (2009).
- 5. Kessler, R. C. Epidemiology of women and depression. J Affect Disord 74, 5-13 (2003).
- 6. Organization, W. H. Depression. (2016).
- 7. Trivedi, M. H. et al. Evaluation of outcomes with citalopram for depression using measurement-based care in STAR*D: implications for clinical practice. Am J Psychiatry 163, 28–40, https://doi.org/10.1176/appi.ajp.163.1.28 (2006).
- Sonnenberg, C. M., Deeg, D. J., Comijs, H. C., van Tilburg, W. & Beekman, A. T. Trends in antidepressant use in the older population: results from the LASA-study over a period of 10 years. J Affect Disord 111, 299–305, https://doi.org/10.1016/j. jad.2008.03.009 (2008).
- 9. Racagni, G. & Popoli, M. Cellular and molecular mechanisms in the long-term action of antidepressants. *Dialogues Clin Neurosci* **10**, 385–400 (2008).
- Kessler, R. C. *et al.* The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). JAMA 289, 3095–3105, https://doi.org/10.1001/jama.289.23.3095 (2003).
- 11. Berman, R. M. et al. Antidepressant effects of ketamine in depressed patients. Biol Psychiatry 47, 351-354 (2000).
- 12. Hirota, K. & Lambert, D. G. Ketamine: its mechanism(s) of action and unusual clinical uses. Br J Anaesth 77, 441-444 (1996).

- 13. Krystal, J. H. et al. NMDA agonists and antagonists as probes of glutamatergic dysfunction and pharmacotherapies in neuropsychiatric disorders. Harv Rev Psychiatry 7, 125–143 (1999).
- Krystal, J. H. et al. Glutamate and GABA systems as targets for novel antidepressant and mood-stabilizing treatments. Mol Psychiatry 7(Suppl 1), S71–80, https://doi.org/10.1038/sj.mp.4001021 (2002).
- Krystal, J. H., Sanacora, G. & Duman, R. S. Rapid-acting glutamatergic antidepressants: the path to ketamine and beyond. *Biol Psychiatry* 73, 1133–1141, https://doi.org/10.1016/j.biopsych.2013.03.026 (2013).
- Kashiwagi, K. et al. Channel blockers acting at N-methyl-D-aspartate receptors: differential effects of mutations in the vestibule and ion channel pore. Mol Pharmacol 61, 533–545 (2002).
- Kotermanski, S. E. & Johnson, J. W. Mg2+ imparts NMDA receptor subtype selectivity to the Alzheimer's drug memantine. J Neurosci 29, 2774–2779, https://doi.org/10.1523/JNEUROSCI.3703-08.2009 (2009).
- aan het Rot, M. et al. Safety and efficacy of repeated-dose intravenous ketamine for treatment-resistant depression. Biol Psychiatry 67, 139–145, https://doi.org/10.1016/j.biopsych.2009.08.038 (2010).
- Diazgranados, N. et al. A randomized add-on trial of an N-methyl-D-aspartate antagonist in treatment-resistant bipolar depression. Arch Gen Psychiatry 67, 793–802, https://doi.org/10.1001/archgenpsychiatry.2010.90 (2010).
- Larkin, G. L. & Beautrais, A. L. A preliminary naturalistic study of low-dose ketamine for depression and suicide ideation in the emergency department. Int J Neuropsychopharmacol 14, 1127–1131, https://doi.org/10.1017/S1461145711000629 (2011).
- 21. Chatterjee, M., Ganguly, S., Srivastava, M. & Palit, G. Effect of 'chronic' versus 'acute' ketamine administration and its 'withdrawal' effect on behavioural alterations in mice: implications for experimental psychosis. *Behav Brain Res* 216, 247–254, https://doi. org/10.1016/j.bbr.2010.08.001 (2011).
- Chatterjee, M., Verma, R., Ganguly, S. & Palit, G. Neurochemical and molecular characterization of ketamine-induced experimental psychosis model in mice. *Neuropharmacology* 63, 1161–1171, https://doi.org/10.1016/j.neuropharm.2012.05.041 (2012).
- Swerdlow, N. R., Taaid, N., Oostwegel, J. L., Randolph, E. & Geyer, M. A. Towards a cross-species pharmacology of sensorimotor gating: effects of amantadine, bromocriptine, pergolide and ropinirole on prepulse inhibition of acoustic startle in rats. *Behav Pharmacol* 9, 389–396 (1998).
- Cilia, J., Hatcher, P., Reavill, C. & Jones, D. N. (+/-) Ketamine-induced prepulse inhibition deficits of an acoustic startle response in rats are not reversed by antipsychotics. *J Psychopharmacol* 21, 302–311, https://doi.org/10.1177/0269881107077718 (2007).
- Javitt, D. C. & Zukin, S. R. Recent advances in the phencyclidine model of schizophrenia. *Am J Psychiatry* 148, 1301–1308 (1991).
 Li, N. *et al.* mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science* 329, 000 (2000).
- 959–964, https://doi.org/10.1126/science.1190287 (2010).
 27. Zanos, P. *et al.* NMDAR inhibition-independent antidepressant actions of ketamine metabolites. *Nature* 533, 481–486, https://doi.org/10.1038/nature17998 (2016).
- Weckmann, K., Labermaier, C., Asara, J. M., Müller, M. B. & Turck, C. W. Time-dependent metabolomic profiling of Ketamine drug action reveals hippocampal pathway alterations and biomarker candidates. *Transl Psychiatry* 4, e481, https://doi.org/10.1038/ tp.2014.119 (2014).
- 29. Jeremy M. Berg, J. L. T. and Lubert Stryer. Biochemistry. 5th edition. (2002).
- Amodeo, G. A., Rudolph, M. J. & Tong, L. Crystal structure of the heterotrimer core of Saccharomyces cerevisiae AMPK homologue SNF1. Nature 449, 492–495, https://doi.org/10.1038/nature06127 (2007).
- Corton, J. M., Gillespie, J. G., Hawley, S. A. & Hardie, D. G. 5-aminoimidazole-4-carboxamide ribonucleoside. A specific method for activating AMP-activated protein kinase in intact cells? *Eur J Biochem* 229, 558–565 (1995).
- Townley, R. & Shapiro, L. Crystal structures of the adenylate sensor from fission yeast AMP-activated protein kinase. *Science* 315, 1726–1729, https://doi.org/10.1126/science.1137503 (2007).
- Xiao, B. et al. Structural basis for AMP binding to mammalian AMP-activated protein kinase. Nature 449, 496–500, https://doi. org/10.1038/nature06161 (2007).
- 34. Xiao, B. *et al.* Structure of mammalian AMPK and its regulation by ADP. *Nature* **472**, 230–233, https://doi.org/10.1038/nature09932 (2011).
- Xiao, B. et al. Structural basis of AMPK regulation by small molecule activators. Nat Commun 4, 3017, https://doi.org/10.1038/ ncomms4017 (2013).
- Barnes, K. et al. Activation of GLUT1 by metabolic and osmotic stress: potential involvement of AMP-activated protein kinase (AMPK). J Cell Sci 115, 2433–2442 (2002).
- Lushchak, V. I. Free Radicals, Reactive Oxygen Species, Oxidative Stresses And Their Classifications. Ukr Biochem J 87, 11–18 (2015).
- Ruszkiewicz, J. & Albrecht, J. Changes in the mitochondrial antioxidant systems in neurodegenerative diseases and acute brain disorders. *Neurochem Int* 88, 66–72, https://doi.org/10.1016/j.neuint.2014.12.012 (2015).
- Begara-Morales, J. C. et al. Antioxidant Systems are Regulated by Nitric Oxide-Mediated Post-translational Modifications (NO-PTMs). Front Plant Sci 7, 152, https://doi.org/10.3389/fpls.2016.00152 (2016).
- 40. Wang, X. & Hai, C. Novel insights into redox system and the mechanism of redox regulation. *Mol Biol Rep*, https://doi.org/10.1007/s11033-016-4022-y (2016).
- 41. Rhee, S. G. Overview on Peroxiredoxin. Mol Cells 39, 1-5, https://doi.org/10.14348/molcells.2016.2368 (2016).
- Petersen, A. K. *et al.* On the hypothesis-free testing of metabolite ratios in genome-wide and metabolome-wide association studies. BMC Bioinformatics 13, 120, https://doi.org/10.1186/1471-2105-13-120 (2012).
- Webhofer, C. *et al.* Proteomic and metabolomic profiling reveals time-dependent changes in hippocampal metabolism upon paroxetine treatment and biomarker candidates. *J Psychiatr Res* 47, 289–298, https://doi.org/10.1016/j.jpsychires.2012.11.003 (2013).
- Filiou, M. D. et al. Proteomics and metabolomics analysis of a trait anxiety mouse model reveals divergent mitochondrial pathways. Biol Psychiatry 70, 1074–1082, https://doi.org/10.1016/j.biopsych.2011.06.009 (2011).
- Scaini, G. *et al.* Evaluation of Krebs cycle enzymes in the brain of rats after chronic administration of antidepressants. *Brain Res Bull* 82, 224–227, https://doi.org/10.1016/j.brainresbull.2010.03.006 (2010).
- 46. Magistretti, P. J., Pellerin, L., Rothman, D. L. & Shulman, R. G. Energy on demand. Science 283, 496-497 (1999).
- Webhofer, C. *et al.* Metabolite profiling of antidepressant drug action reveals novel drug targets beyond monoamine elevation. *Transl Psychiatry* 1, e58, https://doi.org/10.1038/tp.2011.56 (2011).
- Glancy, B. & Balaban, R. S. Role of mitochondrial Ca2+ in the regulation of cellular energetics. *Biochemistry* 51, 2959–2973, https:// doi.org/10.1021/bi2018909 (2012).
- Chowdhury, G. M. *et al.* Transiently increased glutamate cycling in rat PFC is associated with rapid onset of antidepressant-like effects. *Mol Psychiatry* 22, 120–126, https://doi.org/10.1038/mp.2016.34 (2017).
- Karabatsiakis, A. *et al.* Mitochondrial respiration in peripheral blood mononuclear cells correlates with depressive subsymptoms and severity of major depression. *Transl Psychiatry* 4, e397, https://doi.org/10.1038/tp.2014.44 (2014).
- Anglin, R. E., Rosebush, P. I. & Mazurek, M. F. Treating psychiatric illness in patients with mitochondrial disorders. *Psychosomatics* 51, 179; author reply 179–180, https://doi.org/10.1176/appi.psy.51.2.179 (2010).
- Anglin, R. E., Garside, S. L., Tarnopolsky, M. A., Mazurek, M. F. & Rosebush, P. I. The psychiatric manifestations of mitochondrial disorders: a case and review of the literature. J Clin Psychiatry 73, 506–512, https://doi.org/10.4088/JCP.11r07237 (2012).

- Anglin, R. E., Tarnopolsky, M. A., Mazurek, M. F. & Rosebush, P. I. The psychiatric presentation of mitochondrial disorders in adults. J Neuropsychiatry Clin Neurosci 24, 394–409, https://doi.org/10.1176/appi.neuropsych.11110345 (2012).
- Moore, C. M., Christensen, J. D., Lafer, B., Fava, M. & Renshaw, P. F. Lower levels of nucleoside triphosphate in the basal ganglia of depressed subjects: a phosphorous-31 magnetic resonance spectroscopy study. Am J Psychiatry 154, 116–118 (1997).
- Volz, H. P. et al. 31P magnetic resonance spectroscopy in the frontal lobe of major depressed patients. Eur Arch Psychiatry Clin Neurosci 248, 289–295 (1998).
- Cunningham, J. T. et al. mTOR controls mitochondrial oxidative function through a YY1-PGC-1alpha transcriptional complex. Nature 450, 736–740, https://doi.org/10.1038/nature06322 (2007).
- Morita, M. et al. mTORC1 controls mitochondrial activity and biogenesis through 4E-BP-dependent translational regulation. Cell Metab 18, 698–711, https://doi.org/10.1016/j.cmet.2013.10.001 (2013).
- Ramanathan, A. & Schreiber, S. L. Direct control of mitochondrial function by mTOR. Proc Natl Acad Sci USA 106, 22229–22232, https://doi.org/10.1073/pnas.0912074106 (2009).
- Coux, O., Tanaka, K. & Goldberg, A. L. Structure and functions of the 20S and 26S proteasomes. Annu Rev Biochem 65, 801–847, https://doi.org/10.1146/annurev.bi.65.070196.004101 (1996).
- 60. Kastle, M. & Grune, T. Protein oxidative modification in the aging organism and the role of the ubiquitin proteasomal system. *Curr Pharm Des* **17**, 4007–4022 (2011).
- 61. Gutteridge, A. *et al.* Nutrient control of eukaryote cell growth: a systems biology study in yeast. *BMC Biol* **8**, 68, https://doi.org/10.1186/1741-7007-8-68 (2010).
- Marondedze, C., Groen, A. J., Thomas, L., Lilley, K. S. & Gehring, C. A Quantitative Phosphoproteome Analysis of cGMP-Dependent Cellular Responses in Arabidopsis thaliana. *Mol Plant* 9, 621–623, https://doi.org/10.1016/j.molp.2015.11.007 (2016).
- 63. Yuan, M., Breitkopf, S. B., Yang, X. & Asara, J. M. A positive/negative ion-switching, targeted mass spectrometry-based metabolomics platform for bodily fluids, cells, and fresh and fixed tissue. *Nat Protoc* 7, 872-881, https://doi.org/10.1038/nprot.2012.024 (2012).
- Xia, J., Psychogios, N., Young, N. & Wishart, D. S. MetaboAnalyst: a web server for metabolomic data analysis and interpretation. Nucleic Acids Res 37, W652–660, https://doi.org/10.1093/nar/gkp356 (2009).
- Tusher, V. G., Tibshirani, R. & Chu, G. Significance analysis of microarrays applied to the ionizing radiation response. Proc Natl Acad Sci USA 98, 5116–5121, https://doi.org/10.1073/pnas.091062498 (2001).
- 66. Svante, W., Michael, S. & Lennart, E. In Chemometrics and Intelligent Laboratory Systems 58, 109-130 (2001).
- Hosack, D. A., Dennis, G., Sherman, B. T., Lane, H. C. & Lempicki, R. A. Identifying biological themes within lists of genes with EASE. *Genome Biol* 4, R70, https://doi.org/10.1186/gb-2003-4-10-r70 (2003).
- Szklarczyk, D. et al. The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. Nucleic Acids Res 45, D362–D368, https://doi.org/10.1093/nar/gkw937 (2017).

Acknowledgements

KW was supported by the International Max Planck Research School for Molecular and Cellular Life Sciences. The technical support by Maria Holzapfel is gratefully acknowledged.

Author Contributions

K.W., M.B.M. and C.W.T. designed research; K.W., M.D., J.H., R.F. performed research; F.D., C.W., M.F., J.I., C.L., G.M., L.T., J.M.A. and K.L. contributed analytic tools; K.W. analysed data; K.W. and C.W.T. wrote the paper.

Additional Information

Supplementary information accompanies this paper at https://doi.org/10.1038/s41598-017-16183-x.

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2017