



Key Concepts for Estimating the Burden of Surgical Conditions and the Unmet Need for Surgical Care

Citation

Bickler, Stephen, Doruk Ozgediz, Richard Gosselin, Thomas Weiser, David Spiegel, Renee Hsia, Peter Dunbar, Kelly McQueen, and Dean Jamison. 2009. Key concepts for estimating the burden of surgical conditions and the unmet need for surgical care. *World Journal of Surgery* 34(3): 374-380.

Published Version

doi:10.1007/s00268-009-0261-6

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:4742706>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Key Concepts for Estimating the Burden of Surgical Conditions and the Unmet Need for Surgical Care

Stephen Bickler · Doruk Ozgediz · Richard Gosselin ·
Thomas Weiser · David Spiegel · Renee Hsia ·
Peter Dunbar · Kelly McQueen · Dean Jamison

Published online: 28 October 2009

© The Author(s) 2009. This article is published with open access at Springerlink.com

Abstract

Background Surgical care is emerging as a crucial issue in global public health. Methodology is needed to assess the impact of surgical care from a public health perspective.

Methods A consensus opinion of a group of surgeons, anesthesiologists, and public health experts was established regarding the methodology for estimating the burden of surgical conditions and the unmet need for surgical care.

Results For purposes of analysis, we define *surgical conditions* as any disease state requiring the expertise of a surgically trained provider. Abnormalities resulting from a surgical condition or its treatment are termed *surgical sequelae*. *Surgical care* is defined as any measure that reduces the rates of physical disability or premature death associated with a surgical condition. To measure the burden

of surgical conditions and unmet need for surgical care we propose using cumulative disability-adjusted life-year (DALY) curves generated from age-specific population-based data. This conceptual framework is based on the premise that surgically associated disability and death is determined by the incidence of surgical conditions and the quantity and quality of surgical care. The *burden of surgical conditions* is defined as the total disability and premature deaths that would occur in a population should there be no surgical care; the *unmet need for surgical care* is defined as the potentially treatable disability and premature deaths due to surgical conditions. Burden of surgical conditions should be expressed as DALYs and unmet need as potential DALYs avertable.

S. Bickler (✉)

Division of Pediatric Surgery, University of California
at San Diego, San Diego, USA
e-mail: sbickler@ucsd.edu

D. Ozgediz

Division of Pediatric Surgery, University of Toronto,
555 University Avenue, Toronto, ON, Canada M5G 1X8

R. Gosselin

School of Public Health, University of California at Berkeley,
50 University Hall #7360, Berkeley, CA 94720-7360, USA

T. Weiser

Department of Health Policy and Management, Harvard
University, Kresge 4th Floor, 677 Huntington Avenue,
Boston, MA 02115, USA

D. Spiegel

University of Pennsylvania Children's Hospital of Philadelphia,
34th Street and Civic Center Boulevard, Philadelphia,
PA 19104, USA

R. Hsia

University of California, Box 1377, 1001 Potrero Avenue 1E21,
San Francisco, CA 94143-1377, USA

P. Dunbar

Department of Anesthesia, University of Washington, PO Box
35640BB-1469, Health Sciences, 1959 NE Pacific Street,
Seattle, WA 98195-6540, USA

K. McQueen

Harvard Humanitarian Initiative, Harvard University,
14 Story Street, 2nd Floor, Cambridge, MA 02138, USA

D. Jamison

Department of Global Health, University of Washington,
Seattle, WA 98195, USA

Conclusions Methodology is described for estimating the burden of surgical conditions and unmet need for surgical care. Using this approach it will be feasible to estimate the global burden of surgical conditions and help clarify where surgery fits among other global health priorities. These methods need to be validated using population-based data.

Introduction

Over the past several years there has been a growing interest in the role of surgery in global public health [1–4]. Inspired by the work of economist Dean Jamison [5] and the chapter “Surgery” in the *Disease Control Priorities in Developing Countries*, 2nd edition [6] on the cost-effectiveness of surgical care in the developing world, a Global Burden of Surgical Disease Working Group (GBSDWG) was formed in 2008 [7]. This group of surgeons, anesthesiologists, emergency physicians, and public health experts met for the first time in Seattle on April 16–17, 2008 to begin a dialogue that explores the role of surgery in public health, priority areas of research, and how best to advocate for support of global surgical programs.

The overwhelming consensus of attendees at this meeting pointed to a critical lack of data concerning the true portion of the global burden of diseases that could be alleviated by surgical expertise. Current methodology was inadequate for measuring the impact of surgical care. During these discussions, several key problems were identified that have impeded accurate measurement of the global burden of surgical conditions: First, basic definitions related to surgical conditions have not been clarified. Second, surgery is a procedure-oriented specialty; and previous global burden of diseases estimates have not been approached from an intervention perspective. Third, the large number of surgical diagnoses and treatments and wide range in patient ages complicate the analysis.

In this article, we propose definitions for assessing surgery from a public health perspective. We also describe a conceptual framework within which we can estimate the burden of surgical conditions and the unmet need for surgical care.

Methods

At the 2008 GBSDWG meeting, a committee was designated to work on surgical definitions. The committee included five surgeons, two anesthesiologists, an emergency room physician, and a health economist. All members of the committee had global health experience, and six members had advanced degrees in public health. Over the subsequent year, the committee worked together to develop a consensus

opinion on definitions and methodology for assessing surgical care from a public health perspective.

Results

Table 1 summarizes our proposed definitions for analyzing surgical care from a public health perspective. The definitions were crafted with the recognition that not all surgical conditions require a procedure; nonphysicians or general doctors, rather than surgical specialists, often provide surgical care in low-income countries; and multiple surgical sequelae can result from a single surgical condition.

Two parameters are described for assessing the public health impact of surgical conditions and surgical care. The severity of a surgical condition is represented as a surgical disability weight (DW). As used in the Disease Control Priorities Project [8], surgical disability weight is a health state valuation expressed on a ratio scale between 0 (full health) and 1 (states equivalent to death).

To measure the impact of surgical care, we propose a parameter called the “value of surgical care” (VSC), which is the relative ability to prevent or reverse a surgical disability. In a manner similar to disability weight, the value of surgical care is expressed on a ratio scale (between 0 and 1) according to how effectively the intervention reverses the disability. Surgical care is curative when the value of surgical care matches the disability weight ($VSC = DW$) or partially restorative when the value of surgical care is less than the disability weight ($VSC < DW$). In situations where surgical care results in a disability greater than the initial condition, the complication is calculated as a separate surgical condition with its own disability weight.

The method we describe assigns a DALY¹ value to each surgical condition and a DALY averted value to each surgical intervention. Using this approach, it is possible to measure quantitatively the burden of surgical conditions and the impact of surgical care.


Figure 1 presents a conceptual framework for estimating the burden of surgical conditions and the unmet need for surgical care. The framework is based on the premise that surgically related DALYs within a population are determined by the incidence of surgical conditions and the quantity and quality of surgical care. For analysis purposes, a cumulative incidence function of surgical DALYs is proposed. Cumulative surgical DALYs can be calculated from age-specific data (Appendix 1).

As shown in Fig. 1, average cumulative DALY curves can vary depending on the level of surgical care. In the best

¹ Disability adjusted life year. The DALY combines time lost due to premature mortality (years of life lost, or YLL) and time lived with the disability (years of healthy life lost due to disability, or YLD). Thus, $DALY = YLL + YLD$.

Table 1 Terminology for analyzing surgical care from a public health perspective

Surgical condition	disease state requiring the expertise of a surgically trained provider
Surgical sequelae	abnormalities that result from a surgical condition or its treatment.
Surgical disability	physical deficit associated with a surgical sequela
Surgical care	any intervention directed at reducing the disability or premature death associated with a surgical condition
Surgical procedure	the suturing, incision, excision, or manipulation of tissue; or other invasive procedure that usually, but not always, requires local, regional, or general anesthesia
Disability weight	measure of the relative valuations of a health state on an interval scale
Value of surgical care	measure of the relative ability to prevent or reverse a surgical disability

**Fig. 1** Conceptual framework for estimating the burden of surgical conditions and the need for surgical care. Surgical disability and premature death in a population is a time-dependent function that relates to the incidence of surgical conditions and the quantity and quality of surgical care. The potential, actual, and unpreventable curves represent the age-specific cumulative disability-adjusted life-years (DALYs) with different levels of surgical care. See Table 2 for the definitions describing the need for surgical care and the burden of surgical conditions

case scenario (when surgical care is ideal), the only surgical disability and premature death in a population is that which is unpreventable. In the worst case scenario (when surgical

care is nonexistent), surgical disability and death reaches the maximum possible. Because surgical care is never completely absent, nor ever ideal, the actual surgical DALY curve lies somewhere between these two curves.

Table 2 summarizes terminology for describing the need for surgical care within a population. The *met need* refers to the surgical care that has been provided, and *unmet need* refers to potentially treatable disability and death due surgical conditions. The *unmeetable need* is the disability and premature death that is unavoidable, even with the best surgical care. The *burden of surgical conditions* is defined as the total disability and premature death that would occur in a population should there be no surgical care. The burden of surgical conditions is expressed as DALYs, met need for surgical care as DALYs averted, and unmet need as potential DALYs avertable. Appendix 2 shows an example of how these parameters can be calculated using population-based data.

Discussion

There is an ongoing debate about the relative importance of surgical care in global public health. Definitions of the burden of surgical conditions and the impact of surgical care are critical to clarifying the relative priority of surgery within global health. To date, neither the global burden of surgical conditions nor the effect of surgical treatment has been quantified with existing measures [9]. As major disparities in surgical care exist between high and low-income countries, an estimate of regionally specific disability-adjusted life-years that can be averted by surgical interventions is also needed [6].

In this article, we outline a conceptual framework for estimating the burden of surgical conditions and unmet need for surgical care. Our definition for surgical conditions is broader than the definition used in the Disease Control Priorities Project, where surgical conditions were defined as “any treatment that includes suture, incision, excision, manipulation, or other invasive procedure that

Table 2 Proposed terminology for describing and measuring burden of surgical conditions and need for surgical care

Term	Definition	Unit of measure
Burden of surgical conditions	The disability and premature death that would exist in a population without any surgical care	DALYs
Met need for surgical care	The disability and premature death in a population that has been prevented or corrected with surgical care	DALYs averted
Unmet need for surgical care	The disability and premature death in a population that is preventable or correctable with surgical care	DALYs potentially avertable
Unmeetable need for surgical care	The disability and premature death in a population that is unpreventable or uncorrectable with even the best surgical care	DALYs unavertable

DALYs disability-adjusted life-years

usually, but not always, requires local, regional, or general anesthesia” [6]. The main consideration in broadening this definition is the fact that surgical conditions do not always require a surgical procedure. Examples include the care of most head injuries and nonoperative management of blunt abdominal injuries (e.g., splenic injury in a child). A clear advantage of the broader definition is that it more accurately reflects the surgical workload, which may be critical when limited resources are being allocated. While the ratio of nonoperative to operative surgical care is likely to vary by specialty, limited data suggest that it may be substantial. In a community-based study of pediatric surgical conditions in West Africa, only 46% of children presenting with a surgical condition required a surgical procedure [10]. Research is needed to determine what this ratio might be in other areas of surgery.

To be consistent with the latest Global Burden of Disease (GBD) study, we use the term *surgical sequelae* to describe any abnormality that results from a surgical condition or its treatment [11]. As used in the current GBD study, sequelae refer to the combination of health states that result from particular causes. This terminology arose from confusion in past GBD studies as to why something was a cause versus sequela versus risk factor. We do not use the term “surgical cause” in any of our definitions, as the definition of a surgical condition is based on a clinical problem rather than an established etiology. The latest round of the GBD study, funded by the Bill and Melinda Gates Foundation, is scheduled to be completed by November 2010. The final study is expected to produce specific DALY, YLL, and YLD estimates for more than 220 diseases/injuries and 40-plus risk factors by age range and sex for 21 regions of the world.

We recognize that there may be confusion about what constitutes surgical conditions and sequelae and that the availability of surgical providers may alter rates of surgical consultation. However, it should be feasible to define surgical conditions and sequelae using the International Classification of Disease (ICD) system. The ICD is a coding system of diseases and signs, symptoms, abnormal findings, complaints, social circumstances, and external causes of injury or diseases as well as therapeutic interventions, as classified by the World Health Organization (WHO) [12]. As of 2008, the ICD-10 coding system includes 68,064 diagnoses and 86,917 procedure codes [13]. The increased detail in the ICD-10, compared to the ICD-9, should make it easier to reach a consensus on which codes represent surgical conditions.

Given the large number of diagnoses and procedures in ICD-10, it may be unrealistic to gather information (or estimate the burden) on the full spectrum of surgical conditions and/or interventions. A more practical approach might be to focus on a subset of these conditions or

procedures. For example, the short list of “essential” surgical procedures (outlined by the WHO) could be expanded from facilities-based data gathered in low-income countries as well as published material. Procedures such as cesarean section, abscess drainage, laparotomy, and fracture care are likely to account for a significant percentage of interventions in most settings.

Recognizing that in many parts of the world surgical procedures are not done by fully trained surgeons, we use the term “surgically trained provider” rather than surgeon. In many other low-income countries, general practitioners may perform surgical procedures. Twenty-five Sub-Saharan countries utilize nonphysician clinicians, and almost half of them perform minor surgical procedures [14]. In Uganda, a study of five general hospitals reported that more than 5000 surgical procedures were performed annually by general practitioners [15]. With only nine orthopedic surgeons for than 20 million people, Malawi has trained orthopedic clinical officers to deliver most of the musculoskeletal services for the country [16]. In addition, 90% of the cesarean sections at the district hospital level in Malawi are carried out by surgically trained clinical officers [17]. In Mozambique, *técnicos de cirurgia* have performed major surgery in district hospitals since 1989 [18]. They perform 92% of emergency obstetric care and 65% of major general surgery at the district hospital level. Other examples of surgically trained providers include nonphysician anesthesia providers [19]. A surgically trained provider implies a level of expertise in making a diagnosis, formulating a treatment plan (including the decision whether an operation is necessary), performing a procedure, and recognizing and treating any complications.

Our method for estimating the burden of surgical conditions and the unmet need for surgical care is based on the concept of “cumulative incidence.” Cumulative incidence refers to the number of new cases that occur in a population over a period of time and is expressed in terms of the number of people at risk in the population at the beginning of the study. A key feature of cumulative incidence analysis is that an average age-specific cumulative risk curve can be plotted using population-based data. This is done under the assumption that age-specific incidence rates remain constant in the future. “Lifetime risk” of disease is a variation of the cumulative incidence concept and has been used by clinicians, researchers, and policymakers to assess the burden of a wide variety of diseases [20–26].

We use disability weights and values for surgical care to calculate cumulative surgical DALYs. This approach allows a DALY value to be assigned to each surgical condition and surgical intervention. In the case of rendered surgical care, the units of measure are expressed as DALYs averted. This approach is fundamentally the same as that used by McCord and Chowdhury [27] and Debas et al. [6], with the exception that their calculations were based on the

percentage of averted risk. We believe that the “value of surgical care” concept allows a much more detailed analysis of surgical services. One concern is that it might be difficult to assign disability weights and values of surgical care to the large number of surgical conditions and interventions. Developing an approach that utilizes an average value for disability weight and value of surgical care could circumvent this problem.

The terminology we use to describe met and unmet need for surgical care is based on definitions used in obstetric need studies [28]. In these studies, obstetric need was estimated using census data, number of females at reproductive ages, and birth rates. Unfortunately, these methods are not as easily applied to the much broader spectrum of surgical conditions. Nevertheless, using terminology that is consistent with the obstetric literature should lessen any confusion.

An important feature of our proposed methodology is that the results will be comparable to other burden of disease studies. “DALYs” have been selected as the measure of burden of surgical conditions and “DALYs averted” as a measure of the impact of surgical care. This should be helpful in determining where surgical care fits among other global health priorities.

We recognize that the DALY continues to evolve in response to criticism and that this summary measure does not take into account contextual variables. Whereas the impairment associated with a particular disease process may be the same throughout the world, the disability associated with each condition will necessarily be determined by social and cultural variables in each environment. Also, disability weights have not yet been calculated for many of the surgical diseases. We suggest that research be undertaken to revise calculations for disability weights associated with surgical conditions, aimed at achieving a consensus opinion from health professionals (and possibly lay people) in both high- and low-income environments.

Finally, even with a strategy to measure the burden of surgical conditions and the unmet need for surgical care at the global level, there are still major obstacles to obtaining the needed estimates. Foremost is a profound lack of data on the incidence and prevalence of surgical conditions. This is especially the case in low- and middle-income countries, where hospital data are of limited value because of access problems and comprehensive surgical databases do not exist. Ideally, prospective community-based surveys will be undertaken in low- and middle-income countries to acquire the needed data.

Conclusions

We have described a strategy for measuring the impact of surgical care from a public health perspective. Using this

methodology, it should be feasible to estimate the burden of surgical conditions and unmet need for surgery in a wide variety of populations.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Appendix 1

Let the age groups be indexed by i , and the interval length and the rate of age-specific disability-adjusted life-years (DALYs) of each age group be represented by n and m , respectively. In the absence of competing deaths, the cumulative surgical DALYs from birth to the end of the i th age interval is given by the following equation:

$$\text{Cumulative surgical DALYs} = \sum_{j \leq i} (n_j \times m_j).$$

Appendix 2

The burden of surgical conditions and unmet need for surgical care can be calculated from population-based data using basic epidemiologic data. Consider the bogus data set of surgical patients shown in Table 3. Included are data from ten patients from a population of 100 treated for surgical conditions during a 1-year period. The patient’s age, sex, diagnoses, and treatment were arbitrarily selected to illustrate the utility of our method over a wide spectrum of patient ages and clinical conditions.

The age-specific values for years of life lost (YLL) are from Table 1.1 (Global Burden of Disease, or GBD, study) in ref. 29. Unless otherwise specified, disability weights for the surgical conditions are from Annex Table 3 (GBD study) in ref. 29. Disability weights of 1.0 were assigned to the patients with anorectal atresia, typhoid bowel perforation, strangulated inguinal hernia, and severe head injury under the assumption that these conditions would be fatal without surgical care.

The categories of the resulting DALYs (averted, potentially avertable, unavertable) were assigned by the surgical care/outcome. If patients’ conditions were corrected by surgery, they were classified as “averted” and as “potentially avertable” if surgical care could have corrected the problem. The disability and death associated with severe head and spinal cord injury was classified as unavertable, as no care could have changed the outcome.

Table 4 summarizes data from the bogus data set. Data are categorized by the age groups used in the GBD study. We arbitrarily selected a population of 100 for our analysis and divided the population into different age groups using

Table 3 Bogus data set of patients used to illustrate calculations of burden of surgical conditions and unmet need for surgical care ($n = 10$ fictitious surgical patients)

Patient characteristics				Calculation of potential DALYs			Category of DALYs ^d		
Age (years)	Sex	Condition	Surgical care/outcome	YLL ^a	DW ^b	Potential DALYs ^c	Averted	Potentially avertable	Unavertable
0	Female	Cleft lip	Not repaired; survived	33.12	0.098	3.2	–	3.2	–
0	Male	Anorectal atresia	Not repaired; died	33.01	1	33.0	–	33	–
10	Female	20% Burn, short term	Treated; survived	37.62	0.186	7.0	7	–	–
15	Male	Typhoid bowel perforation	Not repaired; died	36.8	1	36.8	–	36.8	–
20	Male	Strangulated inguinal hernia	Repaired; survived	35.02	1	35.0	35	–	–
25	Male	Severe head injury	Treated; died	32.53	1	32.5	–	–	32.5
25	Male	Spinal cord injury	Treated; paraplegic	32.53	0.725	23.6	–	–	23.6
30	Female	Rectovaginal fistula	Repaired; survived	29.92	0.43	12.9	12.9	–	–
45	Male	Gastric perforation	Repaired; survived	20.17	1	20.2	20.2	–	–
60	Female	Strangulated inguinal hernia	Treated; survived	12.22	1	12.2	12.2	–	–
Total						216.4	87.3	73.0	56.1

YLL years of life lost, DW disability weight

^a Values are from Table 1.1 in ref. 29

^b See text for how these values were assigned

^c Potential surgical DALYS = YLL × DW

^d Category of DALYs assigned by surgical care/outcome

Table 4 Calculating the burden of surgical conditions and unmet need for surgical care using population-based data

Parameter	Age groups (years)					Total	
	0–4	5–14	15–24	25–59	60+		
Raw data by age group ^a							
No. of cases	2.0	1.0	2.0	4.0	1.0	10.0	
Population	19.0	28.0	19.0	30.0	4.0	100.0	
DALYs averted	0.0	7.0	35.0	33.1	12.2	87.3	
DALYs potentially avertable	36.2	0.0	36.8	0.0	0.0	73.0	
DALYs unavertable	0.0	0.0	0.0	56.1	0.0	56.1	
Total potential DALYs	36.2	7.0	71.8	89.2	12.2	216.4	
Age-specific incidence (DALYs /10,000 population/year) ^b							
DALYs averted	0.0	70.0	350.0	94.6	58.1 ^c	572.7	Met need ^d
DALYs potentially avertable	724.0	0.0	368.0	0.0	0.0	1092.0	Unmet need ^d
DALYs unavertable	0.0	0.0	0.0	160.3	0.0	160.3	Unmeetable need ^d
Total potential DALYs	724.0	70.0	718.0	254.9	58.1	1825.0	Burden of surgical conditions ^e

^a Data are from the bogus data set in Table 3

^b Incidence (DALYs/10,000 population/year) = [(DALYs per age group) × (10,000)]/[(population size) × (years in age group)]

^c Calculation based on an expected life expectancy of 81 years

^d Met, unmet, and unmeetable need are the sums of the age-specific DALYs for each category

^e Burden of surgical conditions is the sum of the potential DALYs for each age group

age distribution data for a population in Sub-Saharan Africa (Annex Table 11, GBD study [29]).

Age-specific incidence rates for averted, potentially avertable, unavertable, and total DALYs are shown in the

lower half of Table 4. Incidence rates are expressed as DALYs/10,000 population/year. As described in Appendix A, age-specific incidence data can be used to calculate cumulative DALY curves. Note that the total sum total of

the age group DALYs equals the cumulative surgical DALYs. Moreover, the values for cumulative DALYs (averted, potentially avertable, unmettable, total) represent the met, unmet, and unmettable need and the burden of surgical conditions.

References

1. PLoS Medicine Editors (2008) A crucial role for surgery in reaching the UN Millennium. *Development Goals PLoS Med* 5:1165–1167
2. Farmer P, Kim J (2008) Surgery and global health: a view from beyond the OR. *World J Surg* 32:533–536
3. Weiser TG, Regenbogen SE, Thompson KD et al (2008) An estimation of the global volume of surgery: a modelling strategy based on available data. *Lancet* 372:139–144
4. Ozgediz D, Jamison D, Cherian M et al (2008) The burden of surgical conditions and access to surgical care in low- and middle-income countries. *Bull World Health Organ* 86:646–647
5. Jamison D (ed) (2006) *Disease control priorities in developing countries*, 2nd edn. Oxford University Press, New York
6. Debas H, Gosselin R, McCord C (2006) Surgery. In: Jamison D et al (eds) *Disease control priorities in developing countries*. 2nd edition. Oxford University Press, New York
7. Global Burden of Surgical Disease Working Group (2009) <http://www.gsd2008.org/>. Accessed 26 Aug 2009
8. Disease Control Priorities Project (2009) <http://www.dcp2.org/pubs/GBD>. Accessed 26 Aug 2009
9. Bickler S, Spiegel D (2008) Global surgery—defining a research agenda. *Lancet* 372:90–92
10. Bickler SW, Telfer ML, Sanno-Duanda B (2008) Need for paediatric surgery care in an urban area of The Gambia. *Trop Doct* 33:91–94
11. Debas H, Jamison D, Lozano R et al (2009) Assessing the burden of surgical disease based on GBD 2005. BoSD Working Group Meeting, Chicago
12. WHO international classification of diseases (ICD). WHO, Geneva
13. American Health Information Management Association (AHIMA) (2009) <http://www.ahima.org/icd10/faq.asp>. Accessed 26 Aug 2009
14. Mullan F, Frehywot S (2007) Non-physician clinicians in 47 Sub-Saharan African countries. *Lancet* 370(9605):2158–2163
15. Ozgediz D, Galukande M, Mabweijano J et al (2008) The neglect of the global surgical workforce: experience and evidence from Uganda. *World J Surg* 32:1208–1215
16. Mkandawire N, Ngulube C, Lavy C (2008) Orthopaedic clinical officer program in Malawi: a model for providing orthopaedic care. *Clin Orthop* 466:2385–2391
17. Chilopora G, Pereira C, Kamwendo F et al (2007) Postoperative outcome of caesarean sections and other major emergency obstetric surgery by clinical officers and medical officers in Malawi. *Hum Resource Health* 14(5):17
18. Pereira C, Cumbi A, Malalane R et al (2007) Meeting the need for emergency obstetric care in Mozambique: work performance and histories of medical doctors and assistant medical officers trained for surgery. *BJOG* 114:1530–1533
19. Hodges SC, Mijumbi C, Okello M et al (2007) Anaesthesia services in developing countries: defining the problems. *Anaesthesia* 62:4–11
20. Lloyd-Jones DM, Larson MG, Beiser A et al (1999) Lifetime risk of developing coronary heart disease. *Lancet* 9(353):89–92
21. Garfinkel L (1995) Probability of developing or dying of cancer United States, 1991. *Stat Bull Metropolitan Insurance Co* 76:31–37
22. Schouten LJ, Straatman H, Kiemeny LA et al (1994) Cancer incidence: life table risk versus cumulative risk. *J Epidemiol Community Health* 48:596–600
23. Cummings SR, Black DM, Rubin SM (1989) Lifetime risks of hip, Colles', or vertebral fracture and coronary heart disease among white postmenopausal women. *Arch Intern Med* 149:2445–2448
24. Seshadri S, Wolf PA, Beiser A et al (1997) Lifetime risk of dementia and Alzheimer's disease: the impact of mortality on risk estimates in the Framingham Study. *Neurology* 49:1498–1504
25. Feuer EJ, Wung L, Boring CC et al (1993) The lifetime risk of developing breast cancer. *J Natl Cancer Inst* 85:892–897
26. Hopper JL, Southey MC, Dite GS et al (1999) Population-based estimate of the average age-specific cumulative risk of breast cancer for a defined set of protein-truncating mutations in BRCA-1 and BRCA-2. *Cancer Epidemiol Biomarkers Prev* 8:741–747
27. McCord D, Chowdhury Q (2003) A cost-effective small hospital in Bangladesh: what it can mean for emergency obstetric care. *Int J Gynaecol Obstet* 81:83–92
28. Tackling unmet need for major obstetric interventions: Part 1 concepts, general principles and international network. UON network—unmet need for major obstetric interventions. <http://www.uonn.org/uonn/pdf/Guide1.pdf>. Accessed 26 Aug 2009
29. Murray CJL, Lopez AD (1996) *The global burden of disease: a comprehensive assessment of mortality and disability from diseases, Injuries, and risk factors in 1990 and projected to 2020*. World Health Organization, Geneva