

Knowledge and Confidence Level Among Emergency Healthcare Workers in Airway Management and Resuscitation of Suspected COVID-19 Patients: A Cross Sectional Study in Malaysia

Azlan Helmy [Abd Samat](#), ¹*Dr Em Med*, Mohd Hisham [Isa](#), ¹*Dr Em Med*, Dazlin Masdiana [Sabardin](#), ¹*MEMed*, Shamsuriani [Md Jamal](#), ¹*MEMed*, Mohd Johar [Jaafar](#), ¹*MSurg (Orth and Trauma)*, Faizal Amri [Hamzah](#), ¹*MMed (EM)*, Alias [Mahmud](#), ¹*PhD*, Amirudin [Sanip](#), ¹*MD*, Syed Mohammad [Syed Ali](#), ¹*MD*, Ismail [Mohd Saiboon](#), ¹*MSurg (Orth and Trauma)*

Abstract

Introduction: This study aims to evaluate the knowledge and confidence of emergency healthcare workers (EHCW) in facing the COVID-19 pandemic. **Materials and Methods:** A cross-sectional online study using a validated questionnaire was distributed to doctors (MD), assistant medical officers (AMO), and staff nurses (SN) at an urban tertiary Emergency Department. It comprised of 40 knowledge and 10 confidence-level questions related to resuscitation and airway management steps. **Results:** A total of 135 from 167 eligible EHCW were enrolled. 68.9% (n = 93) had high knowledge while 53.3% (n = 72) possessed high confidence level. Overall knowledge mean score was 32.96/40 (SD = 3.63) between MD (33.88±3.09), AMO (32.28±4.03), and SN (32.00±3.60), $P = 0.025$. EHCWs with a length of service (LOS) between 4–10 years had the highest knowledge compared to those with LOS <4-year (33.71±3.39 versus 31.21±3.19 $P = 0.002$). Airway-related knowledge was significantly different between the designations and LOS ($P = 0.002$ and $P = 0.003$, respectively). Overall, EHCW confidence level against LOS showed significant difference [F (2, 132) = 5.46, $P = 0.005$] with longer LOS showing better confidence. MD showed the highest confidence compared to AMO and SN (3.67±0.69, 3.53±0.68, 3.26±0.64) $P = 0.049$. The majority EHCW were confident in performing high-quality chest-compression, and handling of Personal Protective Equipment but less than half were confident in resuscitating, leading the resuscitation, managing the airway or being successful in first intubation attempt. **Conclusions:** EHCW possessed good knowledge in airway and resuscitation of COVID-19 patients, but differed between designations and LOS. A longer LOS was associated with better confidence, but there were some aspects in airway management and resuscitation that needed improvement.

Ann Acad Med Singap. 2020;49:643–51

Key words: Airway; Confidence; COVID-19; Knowledge; Resuscitation (CPR).

Introduction

The World Health Organization (WHO) has declared the severe acute respiratory syndrome coronavirus 2 disease (also known as COVID-19) a pandemic on 11 March 2020. Since then, COVID-19 has spread to more than 200 countries globally¹ and the mortality rate ranges from 1 to 20% depending on the country.^{2, 3} Many countries were caught off guard when this disease struck. In fact, from the day of the fifth confirmed death, the country with the highest COVID-19 mortality had shifted from Wuhan, China to Europe, then to the United States of America and subsequently, to South

America.⁴ Furthermore, at the time this article was written in early August, 2020, there were more than 18 million confirmed cases with more than 680 000 deaths reported worldwide from January 2020.^{1,4}

In the early stages of the disease, infected patients show symptoms that are indistinguishable from upper respiratory tract infection (URTI). It then progresses to more severe and critical conditions such as severe breathlessness, high grade fever, haemodynamic instability or circulatory collapse.^{5–7} Like any other collapsed patient, suspected COVID-19 patients require cardiopulmonary resuscitation (CPR) with airway management, but

¹ Department of Emergency Medicine, Faculty of Medicine, Universiti Kebangsaan Malaysia, Malaysia

Address for Correspondence: Prof Dr Ismail Mohd Saiboon, Department of Emergency Medicine, Faculty of Medicine, University Kebangsaan Malaysia, Jalan Yaacob Latif, Bandar Tun Razak, Cheras, 56000 Kuala Lumpur, Malaysia

Email: ismail@ppukm.ukm.edu.my

with special precautionary measures, which need to be understood and taken by the resuscitation team.^{8,9} Emergency health care workers (EHCW) need special training to approach suspected COVID-19 cases because of the high risks of transmission from patients and spread among themselves.¹⁰ Within a short period of time, EHCW ought to learn new knowledge, adapt to new policies and guidelines, and overcome their fears and anxieties while maintaining professionalism in delivering emergency care services.¹¹ A cross sectional study in Pakistan among HCW revealed positive findings on knowledge and attitude towards COVID-19.¹² However, this study did not focus on EHCW as it involved doctors, nurses and pharmacists in general. In addition, confidence in airway management and resuscitation was not explicitly explored.¹² Meanwhile, a study in Wuhan, China involving the frontline HCWs including doctors, nurses and paramedics showed good correlation between knowledge, confidence level and attitudes in handling COVID-19 cases.¹³

In Malaysia, the threat of COVID-19 became apparent when Singapore reported its first imported COVID-19 case from Wuhan, China on 23 January 2020, followed by identification of close contacts of the case in Johor, Malaysia.^{6,14} Less than 2 days after the first case was reported in Singapore, Malaysia reported its first COVID-19-positive case on 25 January 2020; an imported case from Wuhan, China.¹⁵ The situation became worse in Malaysia in March after the International Health Regulations (IHR) Malaysia received information on one positive case in Brunei. The patient had attended a mass religious gathering in Seri Petaling Mosque, Selangor, Malaysia, from 27 February to 1 March 2020, which was attended by more than 10,000 participants from numerous countries, with the majority coming from Malaysia.¹⁶ At the time this article was written, the country has recorded more than 9,000 confirmed cases, with a mortality rate of around 1.4% (125 deaths).¹⁷ All hospitals' HCW were alerted to the potential influx of high-risk patients throughout the country.

We set out to determine the readiness of EHCW in terms of knowledge and confidence level in dealing with this deadly disease. This is crucial as only with good level of preparedness and sufficient knowledge will the EHCW be able to face this situation confidently and with a high degree of safety.⁹ Furthermore, specialised training should be introduced to train all in mitigating this problem.⁹

Methodology

This is a cross-sectional study to investigate the knowledge and confidence level of EHCW working in

the Emergency Department (ED) of a tertiary teaching hospital located in Kuala Lumpur. The EHCW were categorised into 3 designations: medical doctors (MD), assistant medical officers (AMO) and staff nurses (SN). It was conducted from 1 April 2020 to 31 May 2020. During this period, the government enforced strict lockdown measures aimed at containing the spread of the disease. The inclusion criterion was all EHCW who were active for duty during the study period. We excluded those who declined to participate in the study.

This study collected data using an online self-administered questionnaire that can be obtained through the link <https://forms.gle/jVi46LkZfAQuuEb78> (Figure 1). The questionnaire was designed on a Google form and its link was shared with all the emergency department healthcare workers via a dedicated WhatsApp group (Whatsapp Messenger Version 2.20.193.9. Whatsapp Inc. Boston, MA). The link was also shared directly with individual EHCW who were in the contact lists of the investigators or via e-mail communication.

A survey instrument was designed based on guidelines, reports, course materials on resuscitation and respiratory disease including COVID-19.^{8,18–20} The questionnaire was divided into two parts. Part A was related to the demography of the respondents, comprising age, gender, EHCW designation categories and length of service (LOS) which was defined as the number of years of working as a healthcare provider.

Part B consisted of 40 questions that were related to knowledge, and 10 questions related to confidence level, respectively. Among the knowledge-related questions, 23 were about airway management and the other 17 were about resuscitation measures. These questions required the respondents to select the answer in each statement to be either true or false.

The 10 questions related to confidence level consisted of 4 questions on airway-related confidence (ARC) (Q2, Q3, Q7, Q9), 4 questions on resuscitation-related confidence (RRC) (Q1, Q4, Q5, Q10), and the remaining 2 questions were on personal protective equipment (PPE) (Q6, Q8).

The respondents indicated their answers based on a 5-point Likert scale; strongly agree = 5; agree = 4; undetermined = 3; disagree = 2; and strongly disagree = 1. The questionnaire was validated by 3 local emergency physicians and a pilot study was conducted to obtain the reliability of this questionnaire. Cronbach alpha (α) score for the questionnaire was 0.748.

The sample size calculated by Krejcie and Morgan formula was 118, assuming a response rate of 50%,

Fig. 1. Questionnaire

1. I have read the information in the Patient Information Sheet including information regarding the risk in this study and understand that I may freely choose to withdraw from this study at any time without reason and without having to give an explanation and understand that my anonymity will be ensured in the write-up. I voluntarily agree to be part of this research study.
 Yes No

Part A: Demographic Details

2. Designation
 Specialist Medical Officer House officer Assistant Medical Officer Nurse

3. Gender
 Male Female

4. Age
 20-30 31-40 41-50 51-60

5. How long you have been in service?
 Less than 1 year 1-3 years 4-6 years 7-10 years More than 10 years

6. Do you have co-morbid?
 Yes No

7. If you have co-morbid, please specify:

Part B: Knowledge
During the resuscitation of suspected Covid-19 patient, the virus can spread to you via:

8. Bag valve mask ventilation
 True False

9. Blood taking
 True False

10. Chest compression
 True False

11. Endotracheal intubation
 True False

COVID-19 virus can be spread via:

12. Inhalation through the patient airway
 True False

13. Mucosa through patient vomitus
 True False

14. 3. Needle prick through patient blood
 True False

15. Usage of non-invasive ventilation (NIV)
 True False

A patient was brought into the PUI Centre unresponsive, with no breathing and no pulse. What are the immediate steps that need to be done?

16. Administer 1V. Adrenaline 1mg stat
 True False

17. Administer oxygen via non-rebreather mask 15L/min
 True False

18. Attach ECG electrodes

37. Minimum of 4 people are required to attend to intubation of SARI patient in the emergency department
 True False

38. Pre-oxygenation with 15L/min high flow mask oxygen is recommended
 True False

39. Suctioning of the airway prior to intubation increases risk of aerosolization
 True False

Regarding intubation of suspected COVID-19 patient:

40. Wearing a face shield does not improve safety unless the intubator also wears N95 mask
 True False

41. During intubation of a suspected COVID-19 case, an adequate dose of muscle relaxant must be given in order to achieve total paralysis
 True False

42. If there is a failure of intubation with the conventional approach, the intubating personnel should resort to cricothyroidotomy immediately
 True False

43. Risk of aerosolization increases during a change of tubing of ventilator
 True False

Regarding intensive care management of a suspected Covid-19 patient:

44. End-tidal CO2 capnography is NOT recommended in determining the correct placement of ETT
 True False

45. Intubated SARI patient can be managed in normal Non-Covid General ICU
 True False

46. Suctioning of the patient's oral pharyngeal with Yankauer suction is permissible
 True False

47. The ventilator tubing should be sealed at all connection points
 True False

Part C: Confidence
 On a scale of 1-5, with 1 being strongly disagree, 2 is disagree, 3 is indeterminate, 4 is agree and 5 being strongly agree. How would you rate your confidence level on resuscitation of PUI/COVID positive patient during the COVID-19 pandemic?

48. You are comfortable resuscitating a PUI patient
 1 2 3 4 5

49. You are well trained to manage the airway
 1 2 3 4 5

50. You are confident in managing the airway
 1 2 3 4 5

51. You are comfortable in doing High-Quality CPR on the patient
 1 2 3 4 5

True False

19. Check your team PPE for any breach, then start chest compression
 True False

Before you start performing one-man compression-only CPR, you MUST:

20. Cover the victim's nose and mouth with a 3-ply surgical mask
 True False

21. Cover the victim's mouth and nose with non-re-breather face mask
 True False

22. Give two rescue breaths via 2-man technique with a good seal
 True False

23. Perform pre-cordial chest thump
 True False

During resuscitation of a suspected PUI or patient in Zone A, to ensure a high-quality CPR:

24. The chest compressor is not allowed to change with other team members even though he is not doing it effectively, to avoid contamination
 True False

25. The compressor should change with the person who administers the drug for every 2 minutes
 True False

26. The compressor should stop compressing the chest during the attempt of endotracheal intubation.
 True False

27. The runner can be called in to take over chest compression if the compressor is not doing it well.
 True False

Regarding etiquette of PPE during resuscitation:

28. Application of Tyvek suite is a level 2 PPE
 True False

29. Cross-checking of PPE among team members is mandatory to ensure PPE not breach
 True False

30. If PPE is breached during resuscitation, one must immediately step out from the resuscitation area
 True False

31. The chest compressor who is wearing Tyvek PPE should switch role with the airway person who is also wearing a Tyvek PPE after intubation was complete
 True False

During resuscitation of a suspected COVID-19 patient:

32. Application of non-rebreather mask without connection to oxygen can protect against aerosolization
 True False

33. Chest compression does not generate aerosolization
 True False

34. Defibrillation is considered to generate aerosolization
 True False

35. 4. Usage of low flow nasal cannula 3L/min can minimize generation of aerosol
 True False

Regarding airway management in suspected COVID-19 patient:

36. Intubation of SARI (Severe Acute Respiratory Infection) requires application of Tyvek
 True False

52. You are comfortable to lead the resuscitation team
 1 2 3 4 5

53. You are comfortable checking your PPE for any breach
 1 2 3 4 5

54. You are comfortable in voicing your concern to your senior if the need arises
 1 2 3 4 5

55. You are confident of achieving success on your 1st intubation attempt
 1 2 3 4 5

56. You know what to do if you notice your PPE is breached during resuscitation
 1 2 3 4 5

57. You know what to do if the patient suddenly vomits on you during intubation attempt
 1 2 3 4 5

95% confidence interval (CI), Z of 1.96 and margin of error of 5%. A further 10% was added to counteract any errors in completing the questionnaire, resulting in a final sample size of 130.

The results were analysed descriptively for the number of respondents, percentage, and the mean for overall and each category of designation and LOS using SPSS version 26 (IBM Corp, Armonk, NY). We analysed the mean knowledge score and confidence level for each category of EHCW based on the different designation and LOS using one-way ANOVA and Post Hoc Tukey HSD. Further analyses were done on the confidence level of the three-separate cluster of airway, resuscitation and PPE. Chi squared test and Pearson's rank correlation were used to compare differences in knowledge and confidence level of EHCW by demographic characteristics.

This study complied with, and received the approval from, the Medical Research Ethics Committee (MREC), Universiti Kebangsaan Malaysia (approval no. FF-2020-185). In addition, the confidentiality of all the respondents was guaranteed, and all respondents agreed to participate in the study voluntarily by signing the consent form before answering the questionnaire.

Results

Out of the 167 total EHCW eligible for the study, 135 responded to the questionnaire and were included in the final analysis. These respondents consisted of: MD 45.9% ($n = 62$), AMO 37.8% ($n = 51$) and SN 16.3% ($n = 22$). Of these, 51.9% ($n = 70$) were males, and 68.2% were aged between 31 and 40 years. In terms of working experience, 83.0% ($n = 112$) had less than 10 years of experience. Only 11.9% ($n = 16$) of them had comorbid illnesses. Table 1 shows details of the demography.

Knowledge of EHCW on Managing COVID-19

Overall, this study revealed that 68.9% ($n = 93$) of EHCW had good knowledge, and 53.3% ($n = 72$) had high confidence level in resuscitation and airway management of suspected COVID-19 patients (Table 1). Table 2 displays the mean scores of knowledge among different designations and LOS. Out of total score of 40, the mean score for overall knowledge was 32.96 ($SD = 3.63$). One-way ANOVA showed significant difference of knowledge scores between the different designations and LOS with $P = 0.025$ and 0.004 , respectively. Post hoc analyses revealed a significant difference between MD ($M = 33.88$, $SD = 3.09$) and AMO ($M = 32.28$, $SD = 4.03$) with $P = 0.049$. In terms

of LOS, the overall knowledge level for 4–10 years ($M = 33.71$, $SD = 3.39$) was significantly higher than LOS <4 years ($M = 31.21$, $SD = 3.19$), with $P = 0.002$. However, there was no statistically significant difference between LOS duration 4–10 years and >10 years of service.

Subsequent analyses of the 23 questions on airway related knowledge (ARK) using one-way ANOVA revealed significant difference between designations with $P = 0.002$. Post hoc Tukey HSD test indicated significant difference between MD and SN ($M = 20.67$, $SD = 1.99$ versus $M = 18.68$, $SD = 2.70$) with $P = 0.002$. An analysis based on different LOS showed statistically significant difference ($F(2, 132) = 6.262$, $P = 0.003$). Post hoc comparison showed that the ARK for LOS <4 years ($M = 18.82$, $SD = 2.35$) was significantly lower than LOS 4–10 years ($M = 20.50$, $SD = 2.32$) with $P = 0.002$.

Analyses of the 17 questions on Resuscitation Related knowledge (RRK) using one-way ANOVA revealed no significant difference between designation and LOS with [$F(2, 132) = 1.59$, $P = 0.207$] and [$F(2, 132) = 2.017$, $P = 0.137$], respectively (Table 2).

Confidence Level with Designation and Length of Service

Table 3 displays the mean confidence scores among different designations and LOS. There was a significant difference in overall confidence level among the different designations, based on one-way ANOVA test, [$F(2, 132) = 3.081$, $P = 0.049$]. Post hoc tests on different designations revealed that MD ($M = 3.67$, $SD = 0.69$) confidence level was significantly different from SN ($M = 3.26$, $SD = 0.64$) with $P = 0.039$. However, the overall confidence level between MD and AMO showed no significant difference with $P = 0.516$.

Similarly, one-way ANOVA comparison of overall confidence level with LOS showed significant difference [$F(2, 132) = 5.46$, $P = 0.005$]. Post hoc test showed the confidence level of those with LOS <4 years ($M = 3.24$, $SD = 0.61$) was significantly different from those with LOS 4–10 years ($M = 3.60$, $SD = 0.71$) with $P = 0.029$. A comparison between LOS <4 years and LOS >10 years revealed significant mean differences ($M = 3.80$, $SD = 0.59$) with $P = 0.006$.

The mean score for ARC among the EHCW was 3.35 ($SD = 0.77$) with no statistically significant difference [$F(2, 132) = 1.264$, $P = 0.286$] based on ANOVA. A confidence level score of 3.55 differentiates between high and low confidence groups (Table 3). Our results revealed all designations had low confidence in managing the airway.

Table 1. Differences in Knowledge and Confidence Level of by Demographics (N = 135)

Particulars		Number (n)	%	Poor Knowledge n (%)	Moderate Knowledge, n (%)	Good knowledge n (%)	χ^2 (P)	Poor Confidence n (%)	High Confidence n (%)	χ^2 (P)
Overall		135	100	1(0.7)	41(30.4)	93(68.9)		63(46.7)	72(53.3)	
Designation	MD	62	45.9	0(0.0)	13(21.0)	49(79.0)	6.790	27(43.5)	35(56.5)	4.95
	AMO	51	37.8	1(2.0)	20(39.2)	30(58.8)	(0.147)	21(41.2)	30(58.8)	(0.084)
	SN	22	16.3	0(0.0)	8(36.4)	14(63.6)		15(68.2)	7(31.8)	
Sex	Male	65	48.1	1(1.5)	22(33.8)	42(64.6)	1.908	26(40)	39(60)	2.24
	Female	70	51.9	0(0.0)	19(27.1)	51(72.9)	(0.385)	37(52.9)	33(47.1)	(0.135)
Age group (Years)	20–30 y.o	37	27.4	1(2.7)	18(48.6)	18(48.6)	12.10	25(67.6)	12(32.4)	8.95
	31–40 y.o	90	68.2	0(0.0)	20(22.2)	70(77.8)	(0.017)	35(38.9)	55(61.1)	(0.011)
	41–50 y.o	8	4.4	0(0.0)	3(37.5)	5(62.5)		3(37.5)	5(62.5)	
Work experience (Years)	< 4y	33	24.4	0(0.0)	18(54.5)	15(45.5)	12.689	22(66.7)	11(33.3)	7.613
	4–10y	78	57.8	1(1.3)	17(21.8)	60(76.9)	(0.013)	33(42.3)	45(57.7)	(0.022)
	>10 y	24	17.8	0(0.0)	6(25.0)	18(75.0)		8(33.3)	16(66.7)	
Co-morbid	Present	16	11.9	0(0.0)	6(37.5)	10(62.5)	0.546	4(25)	12(75)	3.42
	Absent	119	88.1	1(0.8)	35(29.4)	83(69.7)	(0.761)	59(49.6)	60(50.4)	(0.064)

$P < 0.05$ was considered significant. Bold values show significant differences.

Total scores for knowledge ranged from 0–40. A score of <24 was set for poor knowledge, 24–31 for moderate, ≥ 32 for good knowledge on resuscitation and airway management of COVID-19 patients.

Total scores for confidence level ranged from 0–5. A mean score of ≥ 3.55 was set for good confidence level, <3.55 set for low confidence level in resuscitation and airway management of COVID-19 patients.

AMO = Assistant Medical Officers; MD = Medical Doctors; SN = Staff Nurses

ANOVA analysis of ARC against LOS showed statistically significant difference ($F(2, 132) = 9.143, P = 0.000$). Post hoc test showed the ARC for those with LOS <4 years ($M = 2.89, SD = 0.72$) was significantly different from those with LOS 4–10 years and >10 years ($M = 3.45, SD = 0.76$) and ($M = 3.64, SD = 0.61$), respectively, with $P = 0.001$. However, there was no statistically significant difference between the group of 4–10 years and those with > 10 years of service.

An analysis of RRC against designation and LOS with one way ANOVA showed significant difference [$F(2, 132) = 4.655, P = 0.011$]. Post hoc analysis showed significant difference between MD ($M = 3.84, SD = 0.63$) and SN ($M = 3.35, SD = 0.65$), with $P = 0.011$. However, this was not the case with LOS, $P = 0.055$ (Table 3).

There was a significant association between overall confidence level and knowledge scores with $P = 0.017$. In terms of confidence level, 72 respondents showed high, while 63 had low, confidence. Within the group with high confidence level, 79% ($n = 57$) also possessed good knowledge. However, among those in the low confidence group, only 57% ($n = 36$) of the respondents possessed good knowledge.

An analysis of individual attributes of confidence revealed a greater proportion of EHCW had high confidence level in PPE inspection [$n = 107 (79.3\%)$], taking action on breached-PPE [$n = 103 (76.3\%)$] and performing high quality chest compression [$n = 82 (60.1\%)$]. Notably less than half of the EHCW were confident in either resuscitating, leading the

Table 2. Knowledge of Airway and Resuscitation on COVID-19 Patients Among Different Professions (by designation) and Length of Service (LOS)

Knowledge			
Profession	Airway (/23) [%]	Resuscitation (/17) [%]	Overall Mean (/40) [%]
MD	20.67 (SD 1.99) [89.8]	13.21 (SD 1.79) [77.7]	33.88 (SD 3.09) [84.7]
SN	18.68 (SD 2.70) [81.2]	13.32 (SD 1.89) [78.3]	32.00 (SD 3.60) [80.0]
AMO	19.67 (SD 2.45) [85.5]	12.61 (SD 2.28) [74.1]	32.28 (SD 4.03) [80.7]
Total	19.96 (SD 2.39) [86.8]	13.00 (SD 2.01) [76.5]	32.96 (SD 3.63) [82.4]
ANOVA	P = 0.002	<i>P</i> = 0.207	P = 0.025
Length of service			Total
<4 y	18.82 (SD 2.35) [81.8]	12.39 (SD 1.68) [72.9]	31.21 (SD 3.19) [78.0]
4–10 y	20.50 (SD 2.32) [89.1]	13.21 (SD 1.97) [77.7]	33.71 (SD 3.39) [84.3]
>10 y	19.79 (SD 2.17) [86.1]	13.17 (SD 2.44) [77.5]	32.96 (SD 4.21) [82.4]
Total	19.96 (SD 2.39) [86.8]	13.00 (SD 2.01) [76.5]	32.96 (SD 3.62) [82.4]
ANOVA	P = 0.003	<i>P</i> = 0.137	P = 0.004

Level of significance *P* < 0.05

Table 3. Levels of Confidence on Airway and Resuscitation of COVID-19 Patients Among Different Designations and Lengths of Service.

Confidence			
Profession	Airway	Resuscitation	Av. Mean
MD	3.41 (SD 0.84)	3.84 (SD 0.63)	3.67 (SD 0.69)
SN	3.11 (SD 0.68)	3.35 (SD 0.65)	3.26 (SD 0.64)
AMO	3.37 (SD 0.78)	3.63 (SD 0.71)	3.53 (SD 0.68)
Total	3.35 (SD 0.77)	3.68 (SD 0.68)	3.55 (SD 0.69)
	<i>P</i> = 0.286	P = 0.011	P = 0.049
Length of service			
<4 y	2.89 (SD 0.72)	3.48 (SD 0.60)	3.24 (SD 0.61)
4–10 y	3.45 (SD 0.76)	3.67 (SD 0.72)	3.60 (SD 0.71)
>10 y	3.64 (SD 0.61)	3.91 (SD 0.61)	3.80 (SD 0.59)
Total	3.35 (SD 0.77)	3.68 (SD 0.68)	3.55 (SD 0.69)
	P = 0.000	<i>P</i> = 0.055	P = 0.005

Level of significance *P* < 0.05

resuscitation, managing the airway, or achieving first-attempt-intubation-success in management of COVID-19 patients (Table 4).

Discussion

This study, which was the first to be carried out among the EHCW in Malaysia during the initial period of the COVID-19 pandemic, revealed that the majority of our EHCW were knowledgeable but lacked confidence, especially in managing airway and handling resuscitation of suspected COVID-19 patients. Airway management poses a risk of aerosol generation with inadequate training, and may potentially jeopardise personnel safety and increase the spread of the disease.²¹ The overall knowledge of COVID-19 among EHCW was good. Our study had similar findings with those of Bhagavatula et al²², and Zhou et al. which showed that MD possess higher knowledge level than SN and paramedics, which directly influenced their attitudes and confidence levels.¹³ This may be related to the role of MD in resuscitation as team leader and the person who performs the intubation procedure. Nevertheless, these previous studies showed a significant majority (>90%) of the respondents had good knowledge, whereas our findings only showed 68.9%. This is possibly due to the scope of knowledge covered in our study, which focused on resuscitation and technicalities of airway management, as opposed to COVID-19 transmission in general. The findings are significant in the context of the current situation, as

EHCW knowledge is essential in reducing the risk of transmission during aerosol generating procedure (AGP) and safety during resuscitation in general.

As people working at the frontline, EHCW should be knowledgeable in the resuscitation of COVID-19 patients. Adequate knowledge is paramount as the disease is extremely dangerous and easily spreads through direct contact or respiratory secretions including during resuscitation.²³ Previous studies have shown that the level of knowledge of HCW is related to the length of service. A study by Mohamad et al. on junior doctors showed that 68.3% did not have sufficient knowledge to perform resuscitation, including CPR.²⁴ This is in line with our findings where junior EHCW of fewer than 4 years LOS had lower knowledge compared to their senior colleagues.

Our results also showed higher confidence level in airway management among the senior EHCW especially those with more than 10 years of working experience. This is probably due to their previous involvement in outbreaks such as SARS, H1N1, MERS-CoV and exposures to various level of difficulties of airway management. It is similar to a study done in Korea that found confidence level in managing the airway correlated with experience, and the rate of successful intubation was higher among senior physicians.²⁵ Meanwhile, a local survey among junior MD supports our findings, which showed inadequate advanced airway and resuscitation exposure leading to lack of confidence in

Table 4. Confidence of EHCW Across Different Attributes of COVID-19 Patient Management

No	Attributes	Confident (based on Likert scale 4 and 5 (n, %))	Not Confident (based on Likert scale of 1– 3) (n, %)	Mean scores
Q1	Comfortable resuscitating a PUI patient	50 (37.0%)	85 (63.0%)	3.24 (SD = 0.87)
Q2	Well trained to manage the airway	61 (45.2%)	74 (54.8%)	3.28 (SD = 0.97)
Q3	Confident in managing the airway	58 (43%)	77 (57.0%)	3.35 (SD = 0.87)
Q4	Comfortable in doing High Quality Chest Compression	82 (60.1%)	53 (39.3%)	3.66 (SD = 0.94)
Q5	Confident to lead the resuscitation team	47 (34.8%)	88 (65.2%)	3.11 (SD = 1.03)
Q6	Confident to check for breached PPE	107 (79.3%)	28 (20.7%)	4.14 (SD = 0.86)
Q7	Comfortable in voicing concern to seniors	91 (67.4%)	44 (32.6%)	3.87 (SD = 0.91)
Q8	Confident successful intubation on 1 st attempt	49 (36.3%)	86 (63.7%)	3.22 (SD = 1.03)
Q9	Confidence on action to take if PPE breached during resuscitation	103 (76.3%)	32 (23.7%)	4.07 (SD = 0.89)
Q10	Comfortable in handling the patient’s vomitus during intubation	74 (54.8%)	61 (45.2%)	3.54 (SD = 0.98)

patient management.²⁶ Nevertheless, our study revealed no significant difference in confidence in airway management between MD, SN and AMO. Our findings support the current recommendations for COVID-19 airway management to be led by the most experienced provider.⁸

Most EHCW were comfortable with the correct usage of PPE and its appropriate level. This finding was similarly discovered in a Singaporean study done on Avian influenza pandemic preparedness.²⁷ Houghton et al. reported that clear guidelines and training sessions were among the essential factors to increase adherence towards infection control and prevention.²⁸ Knowledge and perception of disease transmission also promote compliance to PPE guidelines.^{13,29,30} In our practice, self-inspection and buddy-system are used to check for proper PPE application and ensure it is working in order.

The 3 confidence attributes that received the lowest scores were resuscitation of COVID-19 patients, success at first intubation attempt, and leading the resuscitation team. The confidence in managing airway and resuscitation depends on the clinical experience such as the number of cases encountered and the total airway procedures performed by the EHCW.³¹ Nevertheless, with the uniqueness of managing a COVID-19 patient's airway, it was not surprising that the EHCW demonstrated lower confidence levels in these aspects.

In the present study, a significant number of respondents showed high knowledge scores in resuscitation and airway management of COVID-19 patients, but had low levels of confidence. On the other hand, adequate knowledge was not always translated into confidence in patient management. Adequate training and experience,^{32,33} perception of the disease,²² comorbidities, social stigma are possible confounding factors which may influence the level of confidence of the HCW. One of the contributing factors to the reduction in confidence among EHCW was the exponential rise in the number of COVID-19 cases and widespread dissemination of this news.

Limitations and Recommendations

This study has several limitations. It was a single-centre study with a relatively small sample size involving the EHCW working in the ED. The majority of the respondents were doctors which limited its generalisation for the emergency workforce, which consists predominantly of nurses. As this study was designed to focus on determining the knowledge and confidence level of EHCW, the competency level

through psychomotor assessment and the associations of these factors with patients' outcome were not explored. To the best of our knowledge, currently, there is no published local data on this subject, hence the inability to compare with other facilities with similar settings.

Future multicentre studies with equally distributed designations among EHCW respondents are recommended to address these limitations. The association between knowledge, confidence and competence level with patients' outcome should also be the main focus for subsequent studies. Therefore, based on the findings from this study, we recommend that specialised training on airway management and resuscitation for suspected COVID-19 patients be conducted regularly for EHCW to improve their confidence and, hence, their readiness to face this pandemic.

Conclusions

EHCW shows good knowledge of airway management and resuscitation of suspected COVID-19 patients differs between different designations and LOS. In addition, the confidence level correlates positively with LOS. Although generally, the confidence level of EHCW was good, airway management, leading resuscitation and success of first attempt intubation for suspected COVID-19 patients were identified as areas for improvement. Future studies should explore this idiosyncrasy.

Acknowledgements

The authors would like to thank Pn Saadah Ahmad from the Research Unit Department of Emergency Medicine, UKM for her invaluable statistical input in this study and also the Department of Emergency Medicine, Hospital Canselor Tuanku Muhriz, for allowing us to conduct this study.

REFERENCES

1. Worldometers. Worldometer. COVID-19 Coronavirus pandemic. : Worldometer; 2020 [cited 2020 4th August 2020]. Available from: <https://www.worldometers.info/coronavirus/>.
2. Baud D, Qi X, Nielsen-Saines K, Musso D, Pomar L, Favre G. Real estimates of mortality following COVID-19 infection. *Lancet Infect Dis.* 2020 Jul;20(7):773.
3. Mizumoto K, Chowell G. Estimating the risk of 2019 novel coronavirus death during the course of the outbreak in China, 2020. medRxiv. 2020.
4. Hannah Ritchie, Esteban Ortiz-Ospina, Diana Beltekian, Edouard Mathieu, Joe Hasell, Bobbie Macdonald, et al. Statistics and Research: Coronavirus (COVID-19) Deaths University of Oxford: University of Oxford; 2020 [cited 4 August 2020]. Available from: <https://ourworldindata.org/covid-deaths>.
5. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet.* 2020 Feb 15;395(10223):497–506.

6. Hsu LY, Chia PY, Lim JF. The Novel Coronavirus (SARS-CoV-2) Pandemic. *Ann Acad Med Singap.* 2020 Mar 16;49(3):105–7.
7. Goh KJ, Choong MC, Cheong EH, Kalimuddin S, Duu Wen S, Phua GC, et al. Rapid Progression to Acute Respiratory Distress Syndrome: Review of Current Understanding of Critical Illness from COVID-19 Infection. *Ann Acad Med Singap.* 2020 March 16;49(3):108–18.
8. Brewster DJ, Chrimes NC, Do TB, Fraser K, Groombridge CJ, Higgs A, et al. Consensus statement: Safe Airway Society principles of airway management and tracheal intubation specific to the COVID-19 adult patient group. *Med J Aust.* 2020 Jun; 212(10):472–481.
9. Mohd Hisham I, Azlan Helmy AS, Ismail MS. The Preparation, Delivery and Outcome of COVID-19 Pandemic Training Program among the Emergency Healthcare Frontliners (EHFs): The Malaysian Teaching Hospital Experience. *Medicine and Health* 2020; 15(1):250–65.
10. Wang J, Zhou M, Liu F. Exploring the reasons for healthcare workers infected with novel coronavirus disease 2019 (COVID-19) in China. *J Hosp Infect.* 2020 May;105(1):100–1.
11. Ho CS, Chee CY, Ho RC. Mental health strategies to combat the psychological impact of COVID-19 beyond paranoia and panic. *Ann Acad Med Singap.* 2020 Mar 16;49(1):155–160.
12. Saqlain M, Munir MM, Ur Rehman S, Gulzar A, Naz S, Ahmed Z, et al. Knowledge, attitude, practice and perceived barriers among healthcare professionals regarding COVID-19: A Cross-sectional survey from Pakistan. *J Hosp Infect* 2020 Jul;105(3):419–23.
13. Zhou M, Tang F, Wang Y, Nie H, Zhang L, You G, et al. Knowledge, attitude and practice regarding COVID-19 among health care workers in Henan, China. *J Hosp Infect.* 2020 Jun;105 (2):183–87.
14. Dr Noor Hisham Abdullah. Kenyataan Akhbar Ketua Pengarah Kesihatan Malaysia pada 24 Januari 2020- Tindakan KKM Bagi Pengesanan Kontak kepada Kes Pertama Positif Novel Coronavirus di negara Singapura pada 24 Januari 2020 Kuala Lumpur: Ministry of Health Malaysia; 2020.
15. Dr Noor Hisham Abdullah. Kenyataan Akhbar Ketua Pengarah Kesihatan Malaysia pada 25 Januari 2020- Pengesanan Kes Baharu yang disahkan dijangkiti 2019 Novel Coronavirus (2019-nCoV) di Malaysia Kuala Lumpur: Ministry of Health Malaysia; 25 Januari 2020.
16. Shah AUM, Safri SNA, Thevadas R, Noordin NK, Abd Rahman A, Sekawi Z, et al. COVID-19 Outbreak in Malaysia: Actions Taken by the Malaysian Government. *Int J Infect Dis.* 2020 Aug;97:108–116.
17. Ministry of health Malaysia. Latest Update on COVID-19 in Malaysia: Ministry of health Malaysia; 2020 [cited 2020 6 August 2020]. Available from: <http://covid-19.moh.gov.my/terkini>.
18. Academy of Emergency Sciences Manila. AHA COVID 19 PALS Algorithm 2020 [update 9 April 2020; cited 10 June 2020]. Available from: <https://www.facebook.com/aesciencesmanila/posts/covid-blue-cpr-in-the-time-of-covid-19-last-edited-april-10-2020-whats-new1-adde/839713119771071/>.
19. Edelson DP, Sasson C, Chan PS, Atkins DL, Aziz K, Becker LB, et al. Interim Guidance for Basic and Advanced Life Support in Adults, Children, and Neonates With Suspected or Confirmed COVID-19: From the Emergency Cardiovascular Care Committee and Get With the Guidelines®-Resuscitation Adult and Pediatric Task Forces of the American Heart Association in Collaboration with the American Academy of Pediatrics, American Association for Respiratory Care, American College of Emergency Physicians, The Society of Critical Care Anesthesiologists, and American Society of Anesthesiologists: Supporting Organizations: American Association of Critical Care Nurses and National EMS Physicians. *Circulation.* 23 June 2020;141(25):e933–e943.
20. Saiboon IM, Apoo FN, Jamal SM, Bakar AA, Yatim FM, Jaafar JM, et al. Improving the position of resuscitation team leader with simulation (IMPORTS); a pilot cross-sectional randomized intervention study. *Medicine (Baltimore).* 2019 Dec;98(49):e18201.
21. Raboud J, Shigayeva A, McGeer A, Bontovics E, Chapman M, Gravel D, et al. Risk factors for SARS transmission from patients requiring intubation: a multicentre investigation in Toronto, Canada. *PLoS One.* 2010 May 19;5(5).
22. Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Novel coronavirus (COVID-19) knowledge and perceptions: a survey on healthcare workers. *JMIR Public Health Surveill.* 2020 Apr 30;6(2):e19160.
23. Nolan J, Monsieurs K, Bossaert L, Böttiger B, Greif R, Lott C, et al. European Resuscitation Council COVID-19 Guidelines Executive Summary. *Resuscitation.* 2020 Aug;153:45–55.
24. Mohammed Z, Arafa A, Saleh Y, Dardir M, Taha A, Shaban H, et al. Knowledge of and attitudes towards cardiopulmonary resuscitation among junior doctors and medical students in Upper Egypt: cross-sectional study. *Int J Emerg Med.* 2020 Apr 22;13(1):19.
25. Kim C, Kang HG, Lim TH, Choi BY, Shin Y-j, Choi HJ. What factors affect the success rate of the first attempt at endotracheal intubation in emergency departments? *Emerg Med J.* 2013 Nov;30(11):888–92.
26. Chew K, Hashairi FM, Zarina ZI, Farid AS, Yazid MA, Hisamuddin N. A survey on the knowledge, attitude and confidence level of adult cardiopulmonary resuscitation among junior doctors in Hospital Universiti Sains Malaysia and Hospital Raja Perempuan Zainab II, Kota Bharu, Kelantan, Malaysia. *Med J Malaysia.* 2011 Mar;66(1):56–9.
27. Wong TY, Koh GC, Cheong SK, Lee HY, Fong YT, Sundram M, et al. Concerns, perceived impact and preparedness in an avian influenza pandemic--a comparative study between healthcare workers in primary and tertiary care. *Ann Acad Med Singap.* 2008 Feb;37(2):96.
28. Houghton C, Meskell P, Delaney H, Smalle M, Glenton C, Booth A, et al. Barriers and facilitators to healthcare workers' adherence with infection prevention and control (IPC) guidelines for respiratory infectious diseases: a rapid qualitative evidence synthesis. *Cochrane Database Syst Rev.* 2020 Apr 21;4(4):CD013582.
29. Barati M, Bashirian S, Jenabi E, Khazaei S, Karimi-Shahanjari N, Zareian S, et al. Factors Associated with Preventive Behaviours of COVID-19 among Hospital Staff in Iran in 2020: An Application of the Protection Motivation Theory. *J Hosp Infect.* 2020 Jul;105(3):430–433.
30. Shi Y, Wang J, Yang Y, Wang Z, Wang G, Hashimoto K, et al. Knowledge and attitudes of medical staff in Chinese psychiatric hospitals regarding COVID-19. *Brain Behav Immun Health.* 2020 Apr;4:100064.
31. Awad Z, Pothier D. Management of surgical airway emergencies by junior ENT staff: a telephone survey. *J Laryngol Otol.* 2007 Jan;121(1):57–60.
32. Mark ME, LoSavio P, Husain I, Papagiannopoulos P, Batra PS, Tajudeen BA. Effect of Implementing Simulation Education on Health Care Worker Comfort With Nasopharyngeal Swabbing for COVID-19. *Otolaryngol Head Neck Surg.* 2020 Aug; 163(2):271–274. doi: 10.1177/0194599820933168.
33. Mouli TC, Davuluri A, Vijaya S, Priyanka ADY, Mishra SK. Effectiveness of simulation based teaching of ventilatory management among non-anaesthesiology residents to manage COVID 19 pandemic-A Quasi experimental cross sectional pilot study. *Indian J Anaesth.* 2020 May;64(Suppl 2):S136–S140. doi: 10.4103/ija.IJA_452_20. Epub 23 May 2020.