

## Knowledge, Attitudes, and Practice Regarding Infection Control Measures Among Dental Students in Central India



### Medical science

**KEYWORDS :** Infection, dental.

**Nilesh Agrawal**

Assistant professor, Dept. of Community Medicine, Travancore medical college, Kollam

**\* Dr. Pranay Gandhi**

Assistant professor, Dept. of Community Medicine, IGGMC, Nagpur. \* corresponding Author

**Sunita Sharma**

Junior resident-3, Dept. of ENT, SRTR medical college, Ambajogai

### ABSTRACT

*A questionnaire study was conducted among 245 dental students from Nagpur city, Central India, to determine their level of knowledge, attitudes, and practice regarding infection control measures and if any correlation exists among the knowledge, attitudes, and practice scores. We found that 61.2 percent of the dental students had not been vaccinated with hepatitis B. Use of face mask, gloves, eyewear, and protective clothing as standard infection control measures was practiced only by two students. Mean knowledge, attitude, and practice scores were 3.75 (1.01), 3.40 (0.75), and 3.35 (1.04), respectively. Significant linear correlation was seen between attitude and practice scores ( $r=0.20$ ,  $p\leq 0.01$ ). The level of knowledge and practice of infection control measures was poor among dental students. We recommend rigorous training on infection control measures prior to graduation and mandatory hepatitis B immunization of students before exposure to clinical practice.*

### Introduction

The dental clinic is an environment where disease transmission occurs easily.<sup>1</sup> Prevention of cross infection in the dental clinic is therefore a crucial aspect of dental practice, and dental clinic workers must adopt certain basic routines while practicing. Dental health care professionals (DHCPs) are at risk of infections caused by various microorganisms such as Mycobacterium tuberculosis, hepatitis B and hepatitis C viruses, staphylococci, streptococci, herpes simplex virus types 1, human immunodeficiency virus (HIV), mumps, influenza, and rubella. Infections may be transmitted in the dental operator through several routes, including direct contact with blood, oral fluids, or other secretions; indirect contact with contaminated instruments, operator equipment, or environmental surfaces; or contact with airborne contaminants present in either droplet splatter or aerosols of oral and respiratory fluids.<sup>2,3</sup>

With this in mind, the aim of this work was to analyze the knowledge, attitudes, and practice regarding infection control measures among dental students in Nagpur city, Central India. The study also explored if any correlation exists among the knowledge, attitudes, and practice scores.

### Methodology:

A questionnaire study was conducted among dental students (third year, final year, interns) of a Dental college in Nagpur city, Central India. The sample was comprised of eighty-six third-year dental students, eighty-two final-year students, and seventy-seven interns. Interns are graduates from the same school, with postings equally distributed in various departments during the stipulated one year. Training in infection control is mainly provided in the first, second, and third years of dental school. The study population of 245 dental students voluntarily completed a questionnaire consisting of fifteen questions. The percentages of students in third year, fourth year, and internships who responded were 91.5 percent, 91.1 percent, and 89.5 percent, respectively.

The questionnaire was framed with the help of experts in the field. The questionnaire kept the study group in mind, and questions were linked to curriculum content of infection control. A self-administrated questionnaire consisting of fifteen close-ended items was used for data collection. The dental students were given the questionnaire in the classrooms and asked to fill it out without discussing it in fifteen minutes. The interns were given the same questionnaire in various departments. There were six questions to assess knowledge, four questions to assess attitude, and five questions to judge infection control practices of the re-

spondents.

Informed consent and university clearance were granted for the study. The questionnaire was pretested on a random sample of dental students to ensure practicability, validity, and interpretation of responses. The validity of the questionnaire was assessed using Cronbach's alpha internal consistency coefficient. Chi-square test was used to compare categorical variables. Analysis of variance (ANOVA) was used to compare mean of knowledge, attitudes, and practice scores. Kendall's test was used to compute the correlation among knowledge, attitudes, and practice scores. A  $p$  value of  $\leq 0.05$  was considered significant for all statistical analyses. Statistical analysis was performed using the Statistical Package for Social Sciences (SPSS) software for Windows version 13.

### Results:

Of the 245 dental students, sixty-three were males and 182 females. No significant differences were noted between males and females

#### 1. Distribution of dental students in study according to gender and class, by number and percentage of respondents in each group.

Gender	3 <sup>rd</sup> year	4 <sup>th</sup> year	Interns	total	P value
Male	26 (22.4%)	21 (17.2%)	16 (12.3%)	63 (25.7%)	
Female	60 (77.6%)	61 (82.8%)	61 (87.7%)	182 (74.3%)	
Total	86 (35.1%)	82 (33.5%)	77 (31.4%)	245 (100%)	0.38

Only 39.2 percent of the undergraduates used antiseptic solution to wash their hands before and after patient examination. The majority (92.2 percent) of the students considered isolation to be an important infection control measure. Surprisingly, 61.2 percent of the undergraduate dental students had not been vaccinated with hepatitis B vaccine. Use of face mask and gloves as an infection control measure was practiced by 69.8 percent (Table 2<sub>1</sub>). The majority (92.2 percent) of the students considered the importance of isolation in infection control, and 96.7 percent confirmed that ineffective sterilization during clinical practice can transmit infection from one patient to another.

## 2. Students' knowledge, attitudes, and practice regarding infection control measures, by number and percentage of total respondents to each item

Question	Response	Number	Percentage
Q1. Do you wash your hands before and after patient examination?	Yes	234	95.5%
	No	11	4.5%
Q2. With what do you wash your hands?	Plain soap	139	56.7%
	Detergent	10	4.1%
	Antiseptic solution	97	39.2%
Q3. Do you prefer oral mouth rinse before commencement of any treatment procedure?	Yes	138	55.4%
	No	107	43.6%
Q4. Do you think isolation is important in infection control?	Yes	226	92.2%
	No	19	7.8%
Q5. With which of the following vaccines have you been vaccinated ?	Hepatitis B	94	38.4%
	Tetanus	29	11.8%
	Tuberculosis	1	0.4%
	None	121	49.4%
Q6. Which of the following do you use to sterilize instruments in dental clinic?	Autoclave	231	94.3%
	Boiling	3	1.2%
	Washing	11	4.5%
Q7. Minimum time required for sterilization in autoclave?	5 min	3	1.2%
	10 min	28	11.4%
	15 min	214	87.4%
Q8. Temperature for sterilization in autoclave?	100° C	3	1.2%
	120° C	236	96.4%
	150° C	6	2.4%
Q9. Which of the following has the highest rate of transmission via saliva?	Hepatitis B	79	32.2%
	AIDS	26	10.6%
	Tuberculosis	123	50.2%
	Don't know	17	6.9%
Q10. What immediate action should be taken in case of direct blood contact with an HIV patient?	Anti-HIV immunoglobulins	86	35.1%
	Anti-HIV drugs	73	29.8%
	Blood tests to be carried out	74	30.2%
	Don't know	12	4.9%
Q11. Odds of HIV transmission after a single contaminated needlestick injury?	0.1%–0.4%	91	37.1%
	1%–4%	94	38.4%
	10%–40%	28	11.4%
	70%–90%	32	13.1%
Q12. As a clinician, what protective measures do you take to prevent yourself from injury?	Face mask and gloves	172	69.8%
	Eyewear	4	1.6%

Question	Response	Number	Percentage
	Protective clothing	68	27.8%
	All the above	2	0.8%
Q13. After use of gloves for a patient, what do you do with them?	Dispose of them	171	69.8%
	Reuse them after wash	4	1.6%
	Reuse them after sterilization	70	28.6%
Q14. Ineffective sterilization during clinical practice can transmit infection from one patient to another?	Yes	237	96.7%
	No	5	2.0%
	Don't know	3	1.3%
Q15. Apart from instrument sterilization, disinfection of dental chair, clinic, dental office is required?	Yes	224	91.4%
	No	13	5.3%
	Don't know	8	3.3%

Knowledge, attitudes, and practice scores were calculated separately. Each correct response added a score of 1. Mean knowledge, attitudes, and practice scores were 3.75 (1.01), 3.40 (0.75), and 3.35 (1.04) respectively. Significant differences between the groups were noted for means of knowledge ( $\leq 0.05$ ) and practice scores ( $\leq 0.01$ ).

## 3. Mean (SD) of knowledge, attitudes, and practice scores regarding infection control

Group	Knowledge Mean (SD)	Attitude Mean (SD)	Practice Mean (SD)
Maximum score: Knowledge: 6, Attitudes: 4, Practice: 5.			
* $\leq 0.05$ , ** $\leq 0.01$			
Interns	3.48 (0.91)	3.42 (0.67)	3.18 (1.03)
Final-year students	3.70 (0.98)	3.47 (0.83)	3.25 (1.10)
Third-year students	4.03 (1.06)	3.30 (0.75)	3.18 (0.92)
Total	3.75 (1.01)	3.40 (0.75)	3.35 (1.04)
p value	$\leq 0.05^*$	0.31	$\leq 0.01^{**}$

Significant linear correlation was seen between attitude and practice scores ( $r=0.20$ ,  $p\leq 0.01$ ). No differences were noted for knowledge-attitude and knowledge-practice scores.

## 4. Correlation among knowledge, attitudes, and practice scores

Variables	Correlation Coefficient	p value
Knowledge-Attitude	0.047	0.46
Attitude-Practice	0.204	$\leq 0.01$
Knowledge-Practice	-0.024	0.71

## Discussion:

The most surprising result of the study was that 61.2 percent of the undergraduate dental students had not been vaccinated with hepatitis B vaccine. The Dental Council of India has made hepatitis B vaccination mandatory for dental students prior to admission, although the school has not listed it as a requirement.

The level of knowledge about and compliance with infection control measures was poor among the students. Attributable reasons could be inadequate training for infection control measures, inadequate supply of personal protective equipment, and carelessness. Similar results were found in studies by Askarian and Assadian,<sup>28</sup> Henrique et al.,<sup>29</sup> Taiwo and Aderinokun,<sup>1</sup> and Ogden et al.<sup>30</sup> regarding dental students in Iran, Brazil, Nigeria, and UK.

Most students (94.3 percent) in our study used an autoclave to sterilize instruments. Henrique et al.<sup>29</sup> conducted a ten-year study to assess attitudes and behavior of dental students concerning infection control rules. In 1995, most students used an autoclave to sterilize instruments (83.8 percent), and this percentage increased in 2005 (95.9 percent). No student could describe the correct pressure, temperature, and sterilization time in either 1995 or 2005. However, in our study, 87.4 percent and 96.4 percent of the students answered correctly about temperature and sterilization time, respectively. The high percentage of correct answers to questions about sterilization procedure revealed a good knowledge.

The majority (92.2 percent) believed in the importance of isolation in infection control. Also, 91.4 percent of the students thought that disinfection of the dental chair, clinic, and dental office is required apart from instrument sterilization. With respect to attitudes towards adherence to infection control measures, the majority of the respondents believed infection control measures to be necessary. Viewing the responses indicates that

dental students in the study have a positive attitude towards infection control measures. Ramesh and Anuradha,<sup>31</sup> in a study done with Bangalore and Chennai dentists found that although attitudes towards treating patients with infectious disease were positive, more knowledge about infection control was needed.

Means of knowledge, attitudes, and practice scores were 3.75 (1.01), 3.40 (0.75) and 3.35 (1.04), respectively. Significant differences between the groups (third year, fourth year, and interns) were noted for knowledge and practice scores ( $p \leq 0.05$ ;  $p \leq 0.01$ ). This was a reflection of students' forgetting material over time. The finding suggests the importance and need of rigorous infection control training prior to graduation. The topic of infection control requires a proactive approach throughout the course.

Personnel are more likely to comply with an infection control program and exposure control plan if they understand its rationale. Clearly written policies, procedures, and guidelines can help ensure consistency, efficiency, and effective coordination of activities.

### Conclusion:

Our findings indicate a lack of understanding of the basics of infection control and the prevention of transmission of communicable infectious diseases not only in a large percentage of our medical students, but also in graduate students who took part in this study.

## REFERENCE

1. Taiwo JO, Aderinokun GA. Assessing cross infection prevention measures at the dental clinic, University College Hospital, Ibadan. *Afr J Med Sci* 2002; 31(3):213-7. | 2. Verrusio AC, Neidle EA, Nash KD, Silverman S, Horowitz AM, Wagner KS. The dentist and infectious diseases: a national survey of attitudes and behavior. *J Am Dent Assoc* 1989; 118:553-62. | 3. Girdler NM, Matthews RW, Scully C. Use and acceptability of rubber gloves for outpatient dental treatment. *J Dent* 1987;15:209-12. | 4. Mosley JW, Edwards VM, Casey G, Redecker AG, White E. Hepatitis B virus infection in dentists. *N Engl J Med* 1975; 293:729-34. | 5. Centers for Disease Control. Recommendations for prevention of HIV transmission in health care settings. *MMWR Morb Mortal Wkly Rep* 1987; 36(Suppl):35-185. | 6. Centers for Disease Control and Prevention. Guidelines for infection control in dental health care settings. 2003: recommendations and reports. Dec 19 2003/52 (RR17). Atlanta: Centers for Disease Control and Prevention, 2003. | 7. Rustage KJ, Rothwell PS, Brook IM. Evaluation of a dedicated dental procedure glove for clinical dentistry. *Br Dent J* 1987;103:193-5. | 8. Crawford TJ. State of the art practical infection control in dentistry. *J Am Dent Assoc* 1985;110:629-33. | 9. Burke FJT, Wilson NHF, Bogge HFJ. Glove wearing by dental surgery assistants. *Dent Update* 1993;20:385-7. | 10. Allen AL, Organ RJ. Occult blood accumulation under the finger nails: a mechanism for the spread of bloodborne infection. *J Am Dent Assoc* 1982;105:455-9. | 11. Machado-Carvalho HP, Martins TCPM, Ramos-Jorge ML, Magela-Machado D, Paiva SM, Pordeus IA. Management of occupational bloodborne exposure in a dental teaching environment. *J Dent Educ* 2007;71(12):1348-55. | 12. Kohn WG, Collins AS, Cleveland JL, Harte JA, Eklund KJ, Malvitz DM. Guidelines for infection control in dental health care settings. *MMWR Recommendations Rep* 2003; 52(17):1-61. | 13. Centers for Disease Control and Prevention. Updated U.S. Public Health Service guidelines for the management of occupational exposures to HBV, HCV, and HIV and recommendations for postexposure prophylaxis. *MMWR* 2001; 50:1-52. | 14. Cleveland JL, Barker L, Gooch BF, Beltrami EM, Cardo D. Use of HIV postexposure prophylaxis by dental health care personnel: an overview and updated recommendations. *J Am Dent Assoc* 2002; 133:1619-30. | 15. Hazelkorn HM. Do dentists have sufficient information about their patients to control infection? *J Dent Educ* 1990;54(2):149-52. | 16. Duffy RE, Cleveland JL, Hutin YJ, Cardo D. Evaluating infection control practices among dentists in Vâlcea, Romania, in 1998. *Infect Control Hosp Epidemiol* 2004;25:570-5. | 17. Elkarrim IA, Abdulla ZA, Yahia NA, Al Qudah A, Ibrahim YE. Basic infection control procedures in dental practice in Khartoum-Sudan. *Int Dent J* 2004;54:413-7. | 18. Di Giuseppe D, Marinelli P, Angelillo IF. A survey of knowledge, attitudes, and behavior of Italian dentists toward immunization. *Vaccine* 2007;25:1669-75. | 19. Mehtar S, Shisana O, Mosala T, Dunbar R. Infection control practices in public dental care services: findings from one South African province. *J Hosp Infect* 2007;66:65-70. | 20. Myers R, Larson E, Cheng B, Schwartz A, Da Silva K, Kunzel C. Hand hygiene among general practice dentists: a survey of knowledge, attitudes, and practices. *J Am Dent Assoc* 2008;139:948-57. | 21. Utomi IL. Attitudes of Nigerian dentists towards hepatitis B vaccination and use of barrier techniques. *West Afr J Med* 2005;24:223-6. | 22. Freire DN, Pordeus IA, Paixão HH. Observing the behavior of senior dental students in relation to infection control practices. *J Dent Educ* 2000;64(3):352-6. | 23. Galli MG, Tesauro M, Bianchi A, Consonni M. Evaluation of Milan University dental students' knowledge of health and hygiene risks related to clinical work. *Minerva Stomatol* 2006;55:391-400. | 24. Qudeimat MA, Farrah RY, Owais AI. Infection control knowledge and practices among dentists and dental nurses at a Jordanian University teaching center. *Am J Infect Control* 2006;34:218-22. | 25. Souza RA, Namen FM, Galan J, Vieira C, Sedano HO. Infection control procedures among senior dental students in Rio de Janeiro State, Brazil. *J Public Health Dent* 2006;66:282-4. | 26. Sofola OO, Folayan MO, Denloye OO, Okeighemen SA. Occupational exposure to bloodborne pathogens and management of exposure incidents in Nigerian dental schools. *J Dent Educ* 2007;71(8):832-7. | 27. Acosta-Gío AE, Borges-Yañes AS, Flores M, Herrera A, Jerónimo J, Martínez M, et al. Infection control attitudes and perceptions among dental students in Latin America: implications for dental education. *Int Dent J* 2008;58:187-93. | 28. Askarian M, Assadian O. Infection control practices among dental professionals in Shiraz Dentistry School, Iran. *Arch Iranian Med* 2009; 12(1):48-51. | 29. Henrique M, Claudia T, Braz F, Lúcia A, Martins S, Almeida I. Attitudes and behavior of dental students concerning infection control rules: a study with a 10-year interval. *Braz Dent J* 2009;20(3). | 30. Ogden GR, Bahrami M, Sivarajasangam V, Phillips G. Dental students' knowledge and compliance in cross-infection control procedures at a UK dental hospital. *Oral Dis* 1997;3(1):25-30. | 31. Ramesh N, Anuradha KP. A survey on infection control knowledge, attitude, and practice among Bangalore and Chennai dentists. *J Indian Dent Assoc* 2000;71:116-7. | 32. Association for Professionals in Infection Control and Epidemiology. APIC position paper: immunization. *Am J Infect Control* 1999;27:52-3. | 33. Leggat PA, Kedjarune U. Bacterial aerosols in the dental clinic: a review. *Int Dent J* 2001;51:39-44. |