

# RADIOCARBON EVIDENCE OF THE MIDDLE TO UPPER PALAEOOLITHIC TRANSITION IN SOUTHWESTERN EUROPE

## LA TRANSICIÓN DEL PALEOLÍTICO MEDIO AL SUPERIOR EN EL SUROESTE DE EUROPA EN BASE A LAS DATACIONES RADIOCARBÓNICAS

OLAF JÖRIS (\*)  
ESTEBAN ÁLVAREZ FERNÁNDEZ (\*\*)  
BERNHARD WENINGER (\*\*\*)

### ABSTRACT

In the present paper we systematically evaluate the radiometric database underlying the Middle to Upper Palaeolithic transition in southwestern Europe. The different models which attempt to explain the demographical processes underlying this transition rely to a large degree on radiocarbon chronology. We observe that: 1) with increasing age, dates on bone samples show large offsets against those on charcoal, often underestimating these for several thousand years BP and; 2) there is no proof for a persistence of Middle Palaeolithic industries into the time of the earliest Aurignacian in SW Europe. These data contradict the “*Ebro-Frontier*” model that distinguishes Late Middle Palaeolithic industries in the SW of the Iberian Peninsula from early Aurignacian ones in the NE. On the contrary, our data 3) imply a model of interregional shifts of populations contracting during severe cold and arid phases and expanding under warmer, interstadial conditions, raising ideas on a regional *in situ* development of the SW European Aurignacian out of Latest Middle Palaeolithic industries made by Neanderthals some 40.0 kyr cal BC.

### RESUMEN

*Se presenta un estudio sistemático sobre la información radiométrica disponible para la transición Paleolítica*

(\*) Forschungsbereich Altsteinzeit des Römisch-Germanischen Zentralmuseums, Schloß Monrepos. 56567 Neuwied. Germany. Correo electrónico: joeris.monrepos@rz-online.de

(\*\*) Área de Prehistoria. Dpto. de Prehistoria, Historia Antigua y Arqueología. Fac. de Geografía e Historia. University of Salamanca. 37002 Salamanca. Spain. Correo electrónico: estebanalfer@hotmail.com

(\*\*\*) <sup>14</sup>C-Labor. Institut für Ur- und Frühgeschichte. University of Cologne. Weyertal 125. 50923 Cologne. Germany. Correo electrónico: b.weninger@uni-koeln.de

Recibido:1-VIII-03; aceptado: 25-VIII-03

*Medio-Paleolítico Superior en el Suroeste de Europa. Los diferentes modelos para explicar el proceso demográfico que subyace en esta transición dependen en gran medida de la cronología radiocarbónica. Se observa que: 1) a mayor antigüedad las fechas sobre hueso muestran una mayor desvisación frente a las muestras sobre carbón, a menudo infravalorando estas varios miles de años BP y 2) que no hay pruebas de perduración de industrias de Paleolítico Medio durante las fases tempranas del Auriñaciense en el SW de Europa. Estos datos contradicen el modelo de “frontera del Ebro” que distingue industrias de Paleolítico Medio Tardío en el SW de la Península Ibérica de las industrias del Auriñaciense temprano en el NE. Por el contrario, 3) nuestros datos implican un modelo de cambios de población interregional que se contrae durante las fases áridas y de frío severo y que se expande durante las fases más cálidas de los interstadios, surgiendo la idea de un desarrollo regional del Auriñaciense del SW europeo a partir de las industrias del Paleolítico Medio Tardío realizadas por los Neanderthales hace 40 kyr cal BC.*

**Key words:** Middle to Upper Palaeolithic transition. Colonization model. “*Ebro-Frontier*”. Radiometric dates. Radiocarbon calibration. Palaeoclimate change. Population dynamics.

**Palabras clave:** Transición. Paleolítico Medio-Superior. Modelo de colonización. “*Frontera del Ebro*”. Datación. Calibración. Cambio paleoclimático. Dinámica de población.

### 1. INTRODUCTION

The disappearance of Neanderthals is one of the most controversial questions in the study of homi-

CULTURE	HOMINID
MIDDLE UPPER PALAEOOLITHIC (MUP)	anatomically modern humans (AMH)
Aurignacian (Au.)	Vogelherd: AMH
Châtelper. LATE MIDDLE PALAEOOLITHIC (LMP)	NEANDERTHALS

Fig. 1. Cultural sequence of Late Middle Palaeolithic to Middle Upper Palaeolithic technocomplexes in SW Europe compared with the hominid record.

mid evolution. Neanderthal fossil remains throughout Europe are exclusively known from pre-Aurignacian sites (Fig. 1), while early anatomically modern humans are exclusively tied to Upper Palaeolithic (UP) and younger technocomplexes (Churchill and Smith 2000; cf. Gambier 1997). Churchill and Smith evaluate the evidence for “*the makers of the Early Aurignacian in Europe*” and assert modern human presence in Europe “*almost certainly*” by ca. 32.0 kyr BP, and – based on Rieks 1931 discovery of a modern human skull (“Stetten 1”) from the base of the Aurignacian layer V of the southern German Vogelherd cave (Riek 1932; 1934) – they see a “*strong possibility*” that modern humans “*were there by ca. 36 ka BP*”. Although the new radiocarbon dates from the same stratum vary by several thousand years BP, they confirm the antiquity of the Aurignacian deposits at Vogelherd cave (Conard and Bolus 2003). But stratigraphic attribution of hominid fossils can rarely be established with certainty since skeletal remains may have been reworked in into older layers, a problem especially relevant for old excavations, which are often less well-documented. A good example is the recent dating of a perforated shell (*Littorina* sp.), found as a grave-good in a Cro-Magnon burial at the famous eponymous rock-shelter of the Gra-

vettian period, which had formerly been attributed to an Aurignacian horizon at this site (Henry-Gambier 2002; cf. Djindjan *et al.* 1999). Being aware of such difficulties Cabrera Valdés *et al.* (2000: 91) state that the “*only well-known human evidence for the time range 40-35 ka is either of Neanderthal or undiagnostic type [...], while the presence of modern human types is not certainly recognized in Europe until about 30 ka*”.

Nevertheless, the majority of models that today undertake efforts to explain the origin of anatomically modern humans is based on the “*Out of Africa*” hypothesis (Stringer and Andrews 1988; Stringer 2003), postulating colonization of Europe by early *Homo sapiens* with a simultaneous contraction of Neanderthal dispersal as expressed in the geographical spread of Middle Palaeolithic (MP) sites (Bocquet-Appel and Demars 2000). Recent studies in molecular genetics of fossil Neanderthal remains (Krings *et al.* 1997; 1999; Ovchinnikov *et al.* 2000) and new finds of early *Homo sapiens* in Ethiopia (White *et al.* 2003; cf. Clark *et al.* 2003) have provided substantial support for the “*Out of Africa*” hypothesis, whereby anatomically modern humans immigrated from Africa through the Near East into Europe replacing indigenous Neanderthal populations (Stringer 2003). This process – roughly placed between 40.0 and 30.0 kyr ago – is generally assumed to be unidirectional, with modern humans spreading rapidly through Central Europe, finally arriving on the Iberian Peninsula (Zilhão and d’Errico 1999).

## 2. THE “EBRO-FRONTIER” AND RECENT CONTROVERSIES

According to the European colonization scenario, anatomically modern humans reached the Iberian Peninsula last. Consequently, most authors today believe that modern humans – following the arrival of the Aurignacian north of the Ebro basin (as represented by the antiquity of early Aurignacian radiocarbon dates from the North of the Iberian Peninsula) – and Neanderthals in the southwestern part of the peninsula coexisted for many thousands of years: i.e. between 40.0 and 30.0 kyr ago (Fig. 2; Zilhão 2000a; 2000b; Zilhão and d’Errico 1999). In view of this “*Ebro-Frontier*”-model the Iberian Peninsula represents the major Neanderthal refugium (Vega Toscano 1993; vgl. Zilhão 1993) before their final replacement during the later Aurig-



Fig. 2. J. Zilhão's "Ebro-Frontier", separating earliest anatomically modern human populations in Europe (as likely to be represented by 'archaic Aurignacian', 'Aurignacian 0' and 'Aurignacian I' inventories) from latest Middle Palaeolithic ones in the southwest of the Iberian Peninsula (modified after: Zilhão 2000a; 2000b; with addition from the "Stage Three Project database" of last interpleniglacial radiometric dates for European archaeological sites: Davies 2000, online; cf. annotation 1).

(arc.) Au. – (archaic) Aurignacian; MP – Middle Palaeolithic; UP – Upper Palaeolithic; LMP – Latest Middle Palaeolithic.

nacian, following ca. 33,500  $^{14}\text{C}$  BP (Vega Toscano 1990; Zilhão 1993).

This paradigm of a quite recent colonization of Europe by anatomically modern humans has been challenged during recent years by an alternative model of an *in situ* development of the Aurignacian out of the preceding regional Latest Middle Palaeolithic (LMP), based on the results of new excavations at El Castillo cave (Cabrera Valdés *et al.* 2001).

Due to the nature of population advances and the sparse typological evidence (i.e. the dispersal of Aurignacian I type split-based points in the northern part of the Iberian Peninsula; cf. Zilhão 2000a; 2000b) the Ebro Frontier model relies to a great extent on methods of chronometric dating, each with its own implicit chronological notions. The alternative hypothesis of an Aurignacian *in situ* development in SW Europe is based on a series of

chronometric data, most of which are also radiocarbon measurements.

In the present paper we have aimed to test the "Ebro-Frontier"-model and to evaluate the possibility of an Aurignacian *in situ* development, using the "Stage Three Project database" of last interpleniglacial (Oxygene Isotope Stage 3 = OIS 3) radiometric dates for European archaeological sites (1), completed for the Iberian Peninsula as a part of assessments of the demographic processes underlying the transition from the LMP to the Earliest Upper Palaeolithic (EUP) in SW Europe.

### 3. THE MIDDLE TO UPPER PALAEOLITHIC TRANSITION AND AURIGNACIAN ORIGINS

Throughout the different regions of Europe, including its southwestern part, i.e. France and the Iberian Peninsula, regional archaeological records best displayed in numerous stratigraphies of cave sites and rock shelters, show exclusively Aurignacian industries post-dating LMP ones attributed to Neanderthals (Fig. 1). At several sites the latest MP is found in sediments formed under interstadial conditions of moderate to temperate climate attributed to the Hengelo-period (Carbonell *et al.* 2000), the most significant warm interval in the second half of OIS 3 (2).

In France and Northern Spain the Châtelperronian (3), which is restricted to these areas, is of

(1) Davies 2000 (online): ARCH-DBASE.XLS at <http://www.esc.ac.uk/oistage3/secure/arch-dbase.xls> (August 2000 with bibliography for the cited dates; cf. Tab. 1 and Tab. 2). Some further extended databases have been compiled by Bocquet-Appel and Demars (2000): <http://intarch.ac.uk/antiquity/additional/bocqtable1.html> and by d'Errico and Sánchez Goñi (2003 at QSR website: <http://www.elsevier.nl/locate/quascirev>), and Zilhão and d'Errico (1999).

(2) According to radiocarbon measurements available for Hengelo-interstadial deposits this warm interval most likely correlates with interstadial 12 of the Greenland ice cores (Greenland Interstadial 12 = GI 12; Jöris 2003). This temperate intervall possibly continues until GI 11 (cf. Fig. 7).

(3) Besides the rich evidence of Châtelperronian in France (Bosinski 1987; Demars 1996), in Northern Spain, level 10 from Cueva Morín, the small inventories from Labeko Koba IX and Ekain X, and those from El Pendo and A Valiña with their stratigraphical problems (Maíllo Fernández 2003) are the only inventories known from Cantabria. From Catalunya only few Châtelperronian points have been recorded embedded in Aurignacian inventories (cf. Canal i Roquet and Carbonell i Roura 1989: 337; l'Arbreda, Cova Pau, Reclau Viver). Besides the problems concerning El Pendo (Montes Barquín and Sanguino González 2001), stratigraphical disturbances may also account to explain the interstratifications of Châtelperronian with Aurignacian levels at Roc de Combe (Rigaud 2001) and Le Piage in France (d'Errico *et al.* 1998; Rigaud 2001).

ten sandwiched between LMP and Aurignacian layers (Carbonell *et al.* 2000; d'Errico *et al.* 1998). The few hominid fossil remains currently known may imply that the Châtelperronian should also to be linked to Neanderthals (Hublin *et al.* 1996; cf. Asmus 1964). Therefore, one may assign the Châtelperronian to the Middle Palaeolithic, as can be also argued from the evolution of lithic technology (Gouedo 1990; cf. Bodu 1990) and tool spectra (de Sonneville-Bordes 1972; Bosinski 1987; 1990), both showing links to LMP industries.

In summary, the MP/UP transition of SW Europe comprises a well-established sequence of LMP, Châtelperronian-, and EUP Aurignacian-type techno-complexes (Fig. 1).

### 3.1. Radiocarbon evidence

During recent years numerous chronometric dates have been assembled from a large number of sites dating to the LMP and EUP on the European continent (Bocquet-Appel and Demars 2000; Davies 2001). The dates have been compared and interpreted in context with archaeological findings (d'Errico and Sánchez Goñi 2003; Zilhão and d'Errico 1999) (4). Most of the dates, with radiocarbon-measurements forming the bulk of dates from the entire corpus of radiometric age-determinations, have been obtained from sites in SW Europe, altogether providing more than 700 radiocarbon measurements  $>17,500$   $^{14}\text{C}$  BP, backed by more than 200 *non*-radiocarbon dates (TL/OSL; U-series; ESR). Due to its comprehensiveness the combined date list is a valuable instrument for studies on the demographic processes underlying the models in question.

However, due to a number of limitations of the different chronometric dating methods employed, in particular radiocarbon, caution in the meaning and interpretation of dates has repeatedly been expressed (e.g. Djindjian 1999; Pettitt 1999; Pettitt and Pike 2001). It is also most important to acknowledge that many of the  $^{14}\text{C}$ -measurements on which the MP/UP transition is based range close to the technical limits of the dating equipment. With increasing age the dates become, as a rule, less reliable (one of the principles of the method underlying radiocarbon dating). This is reflected in increas-

ing standard deviations as well as in the higher number of infinite ('greater than') age determinations – both parameters that largely depend on the technical equipment of the laboratory.

In the past developing radiocarbon methods did indeed produce a few dozen age determinations in the range of 10.0-12.5 half-lives of radiocarbon. These were measured on large peat samples (*ca.* 100 g carbon) by a combination of thermal isotope enrichment with large  $^{14}\text{C}$ - $\beta$ -counting systems (cf. Grootes 1977). Today, however, the technical limits of most modern laboratories for routine  $^{14}\text{C}$ - (AMS)-measurements remain close to 9.5 half-lives i.e. *ca.* 55,000  $^{14}\text{C}$  BP (Fig. 3). While some radiocarbon laboratories promise reliability of their measurements up to at least 40,000  $^{14}\text{C}$  BP (Hedges and Pettitt 1999), others have much lower technical age limits around 5.0-7.5 half-lives of radiocarbon age (5), i.e. the time of transition from LMP to EUP. Such problems principally effect radiocarbon dating and, consequently, the interpretation of dating results.

The most comprehensive study so far on the beginning of the EUP and the earliest occurrences of the Aurignacian in Europe has been undertaken by Zilhão and d'Errico (1999) who, using a selected data set, (1) confirm the validity of the "Ebro-Frontier"-model and (2) state that no Aurignacian older than 36,500  $^{14}\text{C}$  BP (i.e. north of the Ebro basin) withstands their criteria of evaluation of sample taphonomy (6). This picture of earliest Aurignacian presence in Europe results from the omission of all radiocarbon dates older than 36,500  $^{14}\text{C}$  BP on charcoal samples that appear systematically older relative to bone.

(5) Despite corresponding claims documented in the proposed analytical dating errors, we recognise that some radiocarbon laboratories, notably in earlier years, cannot achieve reproducible  $^{14}\text{C}$ -measurements of such high age. A fair portion of the equipment earlier used in  $\beta$ -decay counting quite apparently had rather large and at any rate often widely varying counter backgrounds, with statistical variations not always according to Poisson statistics, so that the reliability of the archaeological radiocarbon data available today is not in all cases beyond reasonable credence. Quality and reliability of dates thus relate to the age of the laboratory and the year a specific sample had been dated, rather than systematic offsets between conventional and mass spectrometric methods of radiocarbon dating as stated by d'Errico and Sánchez Goñi (2003).

(6) It is noteworthy that any dates are worthless without interpretation of the circumstances that may have contributed to the date, but the taphonomy and context of samples for radiometric dating are difficult to evaluate years after excavation. Any such 'evaluation' finally remains a selection judged by the authors' personal criteria. Testing the available data for integrity of their statistical properties should, at least, be carried out.

(4) See annotation 1.




Fig. 3. Decay of radiocarbon with time relative to its initial activity in relation to the dating limits of radiocarbon laboratories. The Middle to Upper Palaeolithic transition (MP/UP) is close to the detection limits of most radiocarbon laboratories.

### 3.2. A material matter

In their analyses of data from the MP/UP transition Zilhão and d'Errico (1999) recognize that bone samples systematically turn out younger than charcoal ones, often diverging by several thousand  $^{14}\text{C}$  BP. This phenomenon is independently and best illustrated by the example of the recently published  $^{14}\text{C}$  data series from Kostienki 14 in Russia (Sinityn *et al.* 2002) (7). The main reason for Zilhão and d'Errico (1999) to trust the bone dates more than the ones produced from charcoal is that charcoal is often judged as 'mobile' material sensitive to stratigraphical disturbance, e.g. by processes of bioturbation. Based on the physical properties of the radiocarbon method of dating, there is no reason to rely more on the dating of bone rather than on charcoal samples, especially when the archaeological context is clearly identified, as is most likely to be the case with charcoal from hearths. For a variety of reasons, namely (1) the smaller initial content of datable carbon, (2) the well-known higher susceptibility of bone towards contamination with younger carbon, and (3) the disappearance of bone collagen due to decay (e.g. Schwarcz 2001), higher reliability should be given charcoal *versus* bone samples, especially when the samples are close to the detection limits of the radiocarbon method of dating.

Taken at face value, the  $^{14}\text{C}$  age-distributions of

the MP and UP of SW Europe more or less overlap in the entire period from around 38,000 to 19,000  $^{14}\text{C}$  BP (Fig. 4, left), creating a "Coexistence Effect" (Conard and Bolus 2003). This results from the exponentially decreasing rate of  $^{14}\text{C}$ -decay and the rapidly increasing susceptibility towards contamination in older samples, as shown by theoretical estimates of the surplus of modern carbon on dating (Grootes 1977). Due to the fact that the youngest LMP samples exclusively derive from samples of bone or burnt (cremated) bone, their young age-determinations are most likely due to effects of contamination and alterations of the physical properties of sample material during processes of combustion (cf. Schwarcz 2001; Gillespie 1997) and/or to difficulties in the techniques and methods of dating burnt bone (Lanting and Brindley 1999; cf. Gillespie 1997). It is often impossible to establish whether the material is contaminated or not, and apart from obvious anomalous samples, it is still difficult to judge until which age LMP dates may be considered reliable.

Having filtered for sample material type (i.e. bone *versus* charcoal), radiocarbon age-distributions for bone dates (Fig. 4, centre) appear similar to those that have already appeared in the entire data set, while a completely different picture of age-distributions is given with charcoal (Fig. 4, right). This is most clearly seen in MP charcoal dates that appear systematically older than those obtained on bone. Direct comparisons of both bone and charcoal samples, as undertaken exemplarily for the SW European MP on the one hand (Fig. 5, left) and the Aurignacian of the same region on the other (Fig. 5, centre; cf. Zilhão and d'Errico 1999), emphasize

(7) At Kostienki 14 a layer of volcanic ash most likely to be linked with the Campanian Ignimbrite eruption (Fedele *et al.* 2002) confirms the great antiquity of charcoal data from the underlying strata, while horse bone samples show remarkable offsets towards younger ages (Sinityn *et al.* 2002).




Fig. 4. Age-distributions [kyr  $^{14}\text{C}$  BP] of Middle Palaeolithic (to the left of each diagram) radiocarbon-dates of SW Europe compared with those available for the Upper Palaeolithic (to the right of each diagram) of the same region comprising all  $^{14}\text{C}$ -data (left), those sorted for bone (center) and samples on charcoal (right).

In the construction method applied to the resulting  $^{14}\text{C}$ -dispersion graphs, each individual radiocarbon date has been defined by its given median value and standard deviation. The corresponding individual Gaussian curves have been added, to give a curve of the summed  $^{14}\text{C}$  dating probability (Geyh 1969). Because each Gaussian curve is normalised with equal area, using this method each date/sample is given equal weight, independent of dating precision.

the complex difficulties with which radiocarbon-bone samples are generally endowed (Jöris *et al.* 2001; Schwarcz 2001). Demographic modelling of the processes underlying the MP/UP transition (Fig. 5, right) is thus strongly affected by the material dated.

### 3.3. Aurignacian origins

Against the background that different sample material may result in entirely different age-distributions, an 'alternative' radiocarbon chronology based solely on charcoal samples for the earliest occurrences of the Aurignacian in Europe (1) pushes back earliest appearance of the Aurignacian to *ca.* 38,300  $^{14}\text{C}$  BP and (2) produces two distinct geographical clusters of possible Aurignacian origin

(Fig. 6), i.e. the southeastern Central European region to the East and the Pyrenean and Cantabrian area to the SW (Jöris 2003). (3) The results do not, however, reflect a geographical pattern that one would assume in a model of anatomically modern humans colonizing Europe from the East to the West.

Since the Aurignacian lacks any readily apparent 'cultural predecessors' outside of Europe, it cannot be excluded that this culture may have evolved locally within these two regions in Europe (cf. Cabrera Valdés *et al.* 2001). The geographical dispersal of the oldest sites dated using  $^{14}\text{C}$ -charcoal samples could imply Aurignacian genesis out of LMP leaf-point industries in the East and out of the Châtelperronian in the SW. In these regions youngest reliable radiocarbon dates on charcoal for the LMP range between 38,800 and 35,900  $^{14}\text{C}$  BP.




Fig. 5. Age-distributions [kyr <sup>14</sup>C BP] of radiocarbon samples on bone (to the left of the left and centre diagrams) compared with those derived from charcoal samples (to the right of the left and centre diagrams) for Middle Palaeolithic and Aurignacian technocomplexes in comparison with age-distributions based on charcoal samples (right) from the Middle Palaeolithic (MP) and Aurignacian (Au.).

In the construction method applied to the resulting <sup>14</sup>C-dispersion graphs, each individual radiocarbon date has been defined by its given median value and standard deviation. The corresponding individual Gaussian curves have been added, to give a curve of the summed <sup>14</sup>C dating probability (Geyh 1969). Because each Gaussian curve is normalised with equal area, by this method each date/sample is given equal weight, independent of dating precision.




Fig. 6. Europe at the Middle to Upper Palaeolithic transition with Châtelperronian sites in the SW (Bosinski 1987; cf. Demars 1996), Uluzzian in Italy (Gioia 1990; Palma di Cesnola 1993), and leaf-point industries in south-central and southeastern Europe (Allsworth-Jones 1986; Bolus and Rück 2000) as well as 'last appearance data' (LAD's) for Middle Palaeolithic levels <40,000 and >35,000 <sup>14</sup>C BP (white underlain) and East European leaf-point industries (grey underlain) in comparison with first appearance data (FAD's) for the Aurignacian in Europe based on radiocarbon dates >35,000 <sup>14</sup>C BP (black underlain).

All radiocarbon FAD's and LAD's derive from charcoal samples and are given in kyr <sup>14</sup>C BP, in case of repeated measurements given as weighted means (cf. Jöris 2003; cf. Tab. 1-2).

Palaeogeography of Europe corresponds to the glacial maximum of the last glaciation at around 22.5 kyr cal BC with ice margins dotted and coastline lowered for some 120 m.

#### 4. THE EARLIEST AURIGNACIAN AND THE LATEST MIDDLE PALAEOLITHIC OF SOUTHWESTERN EUROPE

The question of whether or not the Aurignacian originated from the regional LMP or is the result of anatomically modern humans colonizing Europe can best be approached from the point of view of available radiometric data (Tab. 1-2). In terms of this the “*Ebro-Frontier*”-model between *ca.* 40,000 and 30,000  $^{14}\text{C}$  BP is testable, since it requires two basic lines of evidence, which are (1) proof for Aurignacian presence NE of the Ebro basin at times for which (2) contemporaneous LMP is manifested in




Fig. 7. Calibration of the radiocarbon scale [kyr  $^{14}\text{C}$  BP] between 45,000 and 25,000  $^{14}\text{C}$  BP (Jöris and Weninger 1999a; 1999b; 2000, updated; see annotation 8) in relation to the Greenland GISP2 ice core record ( $‰$  PDB  $\delta^{18}\text{O}$  after: Stuiver and Grootes 2000), in the time-window 44.0–28.0 kyr cal BC. GI – Greenland interstadial; GS – Greenland stadial.




Fig. 8. Calibrated radiocarbon dates on charcoal of Middle Palaeolithic and Aurignacian of SW Europe (data identical with those in Fig. 5, right) against the GISP2 palaeoclimate record from Fig. 7, showing the abrupt ending of the Latest Middle Palaeolithic (LMP) at the onset of the Aurignacian (Au.) in SW Europe. For regional differences the Aurignacian is distinguished into A, B, C1, C2, C3.

the region SW. Both lines can contribute to the colonization model in question, assuming that LMP lithic industries indeed represent Neanderthals, and that anatomically modern humans are the artisans of the Aurignacian.

While first appearance data (FAD's) of the Aurignacian in Europe – based on  $^{14}\text{C}$ -charcoal samples – indeed demonstrate an early Aurignacian presence NE of the Ebro basin (Fig. 6), the second line of evidence is especially problematic to establish. Problems arise –as described in the preceding chapter–, in determining up to which age young LMP radiocarbon dates can still be regarded as reliable and, more importantly since only a few LMP sites in the southern part of the Iberian Peninsula are published in detail, with the exception of Gibraltar.

To derive ‘realistic’ latest-appearance estimates (LAD's) for LMP sites of that region, it is necessary to compare the radiocarbon dates available with *non*-radiocarbon age-determinations. But direct data pairs of different dating methods, radiocarbon *versus non*-radiocarbon, are unfortunately rare. Furthermore, it is necessary to acknowledge that the radiocarbon time-scale requires calibration to cor-



rect for variations in past atmospheric  $^{14}\text{C}$ -levels (Fig. 7) (8) in order to achieve compatibility with the results from other *non*-radiocarbon methods of dating. When calibrated, the  $^{14}\text{C}$ -time interval covering the MP/UP transition proposed by different authors roughly corresponds to the period 40.6 – 30.2 kyr cal BC (Tab. 3), showing a remarkably clear distinction between LMP *versus* Aurignacian age-distributions on the calendric scale (Fig. 8).

#### 4.1. Earliest Aurignacian evidence...

The oldest Aurignacian sites in SW Europe dated by radiocarbon are situated in the North of the Iberian Peninsula (Fig. 2; Fig. 9). These sites – in sum – form the base of the first line of arguments for the “*Ebro-Frontier*”-model, stating that the Aurignacian starts around 36,500  $^{14}\text{C}$  BP in most of Europe (Zilhão and d’Errico 1999), whereas the Aurignacian in the southwestern part of the Iberian Peninsula does not date before 33,500  $^{14}\text{C}$  BP (Vega Toscano 1990; Zilhão 1993).

(8) We note, however, that all radiocarbon “age”-estimates are measured on the conventional  $^{14}\text{C}$ -scale, which is very precisely defined as a dimensionless logarithmic ratio (Mook 1983), so that an independent calibration of all  $^{14}\text{C}$ -ages is necessary if we wish to sensibly discuss the chronological implications of the  $^{14}\text{C}$ -data. Intensive progress in the construction of the calibration curves required for such age-transfer of (uncalibrated) radiocarbon dates into the calendric dimension has been made during the last few years (eg. Jöris and Weninger 1999a; 1999b; 2000; <http://www.calpal.de>: CALPAL-2003; cf. van der Plicht 2002), with the period in question being best recorded in the data sets from Suigetsu in Japan (atmospheric; Kitagawa and van der Plicht 2000) and PS2644 in the North Atlantic (marine; Voelker *et al.* 2000). Between 46.0 and 28.0 kyr  $^{14}\text{C}$  BP (45.0-30.0 kyr cal BC) the combined ‘synthetical’ calibration data set (CALPAL-2003) allows for rough calendric estimates of the  $^{14}\text{C}$ -time scale (Fig. 7), representing a pattern of highly fluctuating  $^{14}\text{C}$ -levels, with periods of extremely high production of radiocarbon (steep parts in the calibration curve), i.e. between 42.0 and 35.0 kyr  $^{14}\text{C}$  BP, and others with limited production rates, resulting in long  $^{14}\text{C}$ -age-plateaux, i.e. between 39.0 and 35.0 kyr cal BC, normally accompanied with extreme age-distortions (cf. Beck *et al.* 2001). It is this complex ‘pattern’ of the calibration curve that allows for improvement of dating precision for periods of rising atmospheric radiocarbon contents as well as the age-distortions which make it difficult to interpret radiocarbon dates that fall into the period of extended *plateaux*.

Due to the method of construction of the CALPAL-2003 record, i.e. the transferral of radiocarbon-dates from marine oxygen isotope records into the GISP2 Greenland ice core age model (Jöris and Weninger 1998; 1999a; 1999b; cf. Voelker *et al.* 1998; 2002),  $^{14}\text{C}$ -ages are reliably linked with palaeoclimate signatures (Fig. 7). This allows for high-precision age-transferral of  $^{14}\text{C}$ -measurements into a calendric age-model, combined with its positioning within the record of OIS 3 palaeoclimate change that is characterised by extremely rapid oscillations between cold and dry (glacial) and more temperate (interstadial) conditions (for the Iberian Peninsula cf. Carbonell *et al.* 2000; d’Errico and Sánchez Goni 2003).

Evidence for an Aurignacian presence in SW Europe dating prior to 36,500  $^{14}\text{C}$  BP is sparse, and a single radiocarbon age on bone dated to 40,000 ± 1400  $^{14}\text{C}$  BP (OxA-3727) obtained from the basal layer at Reclau Viver may be due to a reworked sample (Zilhão and d’Errico 1999). The upper part of the same level contains an Aurignacian I with split-based points (Canal i Roquet and Carbonell i Roura 1989). Similar problems seem to account for a radiocarbon measurement (GifA-97185: 37,200 ± 150  $^{14}\text{C}$  BP; bone) from layer G at Caminade-Est, where “*systematic refitting work carried out by J.-G. Bordes (1998) has demonstrated that around 30% of the archaeological material included in this layer comes from the underlying Mousterian deposits, to which the dated sample could conceivably be related*” (Zilhão and d’Errico 1999, 17). The overlying layer F at this site has produced an Aurignacian I with split-based points, and a radiocarbon date of 35,400 ± 1100  $^{14}\text{C}$  BP (GifA-97186) on bone (Rigaud 2001).

Whereas early Aurignacian sites are dispersed in SW France and Northern Spain (Fig. 2), oldest radiocarbon evidence for the Aurignacian (Tab. 1; Fig. 9) is exclusively restricted to the provinces of Cantabria and Catalunya in Northern Spain. Here the sites of Abric Romaní, L’Arbreda and El Castillo are most controversially discussed, due to their history of research, the antiquity of radiometric measurements with radiocarbon dates going back to > 36,500  $^{14}\text{C}$  BP, and typological arguments.

A series of radiocarbon measurements of charcoal samples from the new excavations of V. Cabrera Valdés immediately in front of the cave entrance at El Castillo gave results ranging between 41,100 ± 1700 (OxA-2477) and 37,100 ± 2200 (OxA-2473)  $^{14}\text{C}$  BP (Tab. 2) (Cabrera Valdés and Bischoff 1989; Cabrera Valdés *et al.* 1996, 2000; 2001) for a layer inside the cave labeled “*Aurignacian Delta*” by H. Obermaier (Cabrera Valdés 1984; cf. Cabrera Valdés *et al.* 1996). Zilhão and d’Errico (1999) critically request (1) the stratigraphic correlation of the layer outside the cave with the inner deposits, (2) the cultural attribution of the recent finds, and (3) the association of the samples dated with the lithic industry, which they think has more affinities with Mousterian or Châtelperronian than the Aurignacian. According to Zilhão and d’Errico (1999), the Aurignacian of El Castillo is present only inside the cave in the upper portion of Obermaier’s “*Aurignacian Delta*”, while the lower part of this layer contains more MP types, and likely

**AURIGNACIEN > 33.5 kyr 14C BP and/or > 36.5 kyr non-14C BP**

LEVEL	CULTURE	WD	MAT.	METH.	LAB-NO.	DATE	STD+	STD-	t-val.
<b>14C on CHARCOAL</b>									
<i>Gr. Tourmal (Bize, Aude, France)</i>									
G (sq. P31)	Au.	748	charcoal	14C	Ly-1898	>35800			
F (sq. LM32)	arc. Au.	746	charcoal	14C	Ly-1895	>29000			
C (sq. L31)	arc. Au.	747	charcoal	14C	Ly-1031	>34200			
<i>Esquicho-Grapaou (Sainte-Anastasia, Gard, France)</i>									
SCL1B	Au. 0	372	charcoal	14C	MC-2161	34540	2000	2000	
<i>Isturitz (Saint-Martin-d'Aberoué, Pyrénées Atlantiques, France)</i>									
V1 26	Au.	708	charcoal	14C	GifA-98233	34630	560	560	
U27, 4d	Au.	707	charcoal	14C	GifA-98232	36510	610	610	
	Au.	707-708	charcoal	14C	weighted mean	35490	413		
<i>La Vina (Manzaneda, Asturias, Spain)</i>									
XIII inf.	Au. I	92	charcoal	14C	Ly-6390	36500	750	750	
<b>14C on CHARCOAL vs. 14C on BONE</b>									
<i>L'Arbreda (Serinyà, Girona, Spain)</i>									
CE 103 [Level H]	Au.	151	bone	14C	OxA-3729	37340	1000	1000	
B1 [Level H] sq. E2	arc. Au.	147	charcoal	14C	AA-3779	37700	1000	1000	0,53
B1 [Level H] sq. E2	arc. Au.	148	charcoal	14C	AA-3780	37700	1000	1000	0,53
B1 [Level H] sq. E2	arc. Au.	150	charcoal	14C	AA-3782	38700	1200	1200	0,30
B1 [Level H] sq. E2	arc. Au.	149	charcoal	14C	AA-3781	39900	1300	1300	1,13
B1 [Level H] sq. E2	arc. Au.	147-150	charcoal	14C	weighted mean	38307	552		
BE 111 [Level H]	arc. Au.	152	bone	14C	OxA-3730	35480	820	820	
<b>14C on CHARCOAL vs. 14C on UNKNOWN MATERIAL</b>									
<i>Beneito (Muro, Alicante, Spain)</i>									
VIII	Au.	160	?	14C	Gif-7650	26040	890	890	
VIII	Au.	159	charcoal	14C	AA-1388	33900	1100	1100	
<b>14C on CHARCOAL vs. U-SERIES</b>									
<i>Romani (Capellades, Barcelona, Spain)</i>									
above lev. 2		142	charcoal	14C	USGS-2839	36600	1300	1300	
2	arc. Au.	135	charcoal	14C	AA-8037A*	35400	810	810	
2	arc. Au.	136	charcoal	14C	AA-8037B*	37900	1000	1000	
2	arc. Au.	135-136	charcoal	14C	weighted mean*	36390	629		0,35
2	arc. Au.	137	charcoal	14C	NZA-2311	36590	640	640	0,07
2	arc. Au.	141	charcoal	14C	AA-6608	36740	920	920	0,10
2	arc. Au.	140	charcoal	14C	AA-7395	37290	990	990	0,61
2	arc. Au.	all level 2	charcoal	14C	weighted mean	36644	373		
2	arc. Au.	144	travertine	U-series	?	43000	1000	1000	
below lev. 2		143	charcoal	14C	USGS-2840	35000	500	500	
					* = identical sample				
<b>14C on BONE</b>									
<i>Flageolet I (Bézenac, Dordogne, France)</i>									
XI	Au. I	321	bone	14C	Ly-2727	>31500			
XI	Au. I	323	bone	14C	GifA-95538	32040	850	850	0,90
XI	Au. I	320	bone	14C	OxA-598	33800	1800	1800	0,42
XI	Au. I	322	bone	14C	GifA-95559	34300	1100	1100	1,03
XI	Au. I	320+322-23	bone	14C	weighted mean	32997	630		
<i>Pataud (Les-Eyzies-de-Tayac, Dordogne, France)</i>									
14	Au. 0	258	burnt bone	14C	GrN-4610	33300	760	760	0,28
14	Au. 0	259	bone	14C	GrN-4720	33330	410	410	0,38
14	Au. 0	257	burnt bone	14C	GrN-4507	34250	675	675	0,97
14	Au. 0	257-259		14C	weighted mean	33529	318		

Tab 1.

<b>Mollet (Serinyà, Girona, Spain)</b>								
0.6-0.8m	Au.	158	bone	14C	OxA-3728	33780	730	730
<b>Combe Sauniere (Sarriac-sur-l'Isle, Dordogne, France)</b>								
VIII	Au.	724	bone	14C	OxA-6507	34000	850	850
<b>Caminade (La Canéda, Dordogne, France)</b>								
D21	Au. II	450	bone	14C	GifA-97187	34140	990	990
<b>Roc de Combe (Payrignac, Lot, France)</b>								
7c	Au. I	433	bone	14C	OxA-1263	34800	1200	1200
<b>Castanet (Sergeac, Dordogne, France)</b>								
inferieur	Au.	712	bone	14C	GifA-97313	35200	1100	1100
<b>Caminade (La Canéda, Dordogne, France)</b>								
F	Au. I	448	bone	14C	GifA-97186	35400	1100	1100
<b>Caminade (La Canéda, Dordogne, France)</b>								
G	Au. I?	449	bone	14C	GifA-97185	37200	1500	1500
<b>Reclau Viver (Serinyà, Girona, Spain)</b>								
T III	arc. Au.?	165	bone	14C	OxA-3727	40000	1400	1400

**14C on BONE vs. 14C on UNKNOWN MATERIAL**

<b>Tuto de Camalhot (Saint-Jean-des-Verges, Ariège, France)</b>								
	Au. I?	156	bone	14C	Gif-2941	24200	600	600
	Au. I		?	14C	?	34750	570	570
<b>Ferrassie (Savignac-de-Miremont, Dordogne, France)</b>								
K6	Au. I	344	?	14C	GrN-5751	33220	570	570
K6	Au. I	343	bones	14C	Gif-4279	>35000		

**14C on UNKNOWN MATERIAL**

<b>Arenillas (Cantabria, Spain)</b>								
II	Au.	87	?	14C	GrN-?	33870	1700	1700 0,20
II	Au.	88	?	14C	GrN-?	34660	1600	1600 0,19
II	Au.	87-88	?	14C	weighted mean	34289	1165	
<b>Pataud (Les-Eyzies-de-Tayac, Dordogne, France)</b>								
?	Au. I	265	?	14C	GrN-3230	34760	1000	1000

**TL / OSL**

<b>Gato Preto (Rio Maior, Santarém, Portugal)</b>								
	Au.	1	burnt silex	TL	BM-?	38100	3900	3900

Tab 1 (Cont.). Radiometric dates for the SW European Aurignacian. WD – No. in William Davies datelist (see annotation 1); MAT. – material dated; METH. – dating method; STD – standard deviation; *t-val.* – *t*-value. (arc.) Au. – (archaic) Aurignacian.

correlates with layer 18 of V. Cabrera Valdés' excavation. Due to the fact that Obermaier did not observe a sterile horizon between the upper and lower portions of the "Aurignacian Delta"-sediment, one may conclude that the Aurignacian immediately followed  $38,679 \pm 744$   $^{14}\text{C}$  BP, which is the weighted mean (WM) of five radiocarbon measurements on charcoal samples from *sub*-unit b of layer 18. The underlying *sub*-unit c has produced five additional radiocarbon dates – again all on charcoal – resulting in a WM of  $40,621 \pm 750$   $^{14}\text{C}$  BP paired with three ESR dates on bone (Rink *et al.* 1996; WM:  $38.4 \pm 2.6$  kyr BP).

However, a consistent group of radiocarbon age-determinations (Tab. 1), significantly older than  $36,500$   $^{14}\text{C}$  BP, has been produced for the Aurigna-

cian layers of L' Arbreda (Bischoff *et al.* 1989; Canal i Roquet and Carbonell i Roura 1989). Four charcoal samples taken in an artificial 5cm-horizon (5,50-5,55m below surface) in square E2 (BE 111) immediately sheltered by the travertine wall in the lower Aurignacian, have produced a WM of  $38,307 \pm 552$   $^{14}\text{C}$  BP (Bischoff *et al.* 1989). Furthermore a radiocarbon measurement of  $35,480 \pm 820$   $^{14}\text{C}$  BP (OxA-3730) on bone assigned to the same cultural unit is available. A few metres to one side (CE 103) a radiocarbon measurement, again on bone, ages the upper Aurignacian level to  $37,340 \pm 1000$   $^{14}\text{C}$  BP (OxA-3729), whereas one further measurement (Gif-6422) obtained earlier, is considered to be too young. Against the homogeneity of the available data we can hardly follow the vague assumptions of




Fig. 9. Radiocarbon dates (first appearance data: FAD's) for the SW European Aurignacian, sorted for material dated and given in kyr  $^{14}\text{C}$  BP, when required as weighted means.

Zilhão and d'Errico (1999) who claim possible stratigraphical disturbances in the various parts of the area excavated, and cannot assign the spatial origin of samples to areas that may be stratigraphically problematic. On the contrary: The samples from square E2 come from the very base of the Aurignacian deposits, immediately next to the main profile published by Canal i Roquet and Carbonell i Roura (1989).

Although the Aurignacian of L'Arbreda displays some similarities with that of "layer 2" at Abric Romaní (e.g. in terms of the presence of Dufour bladelets; Carbonell *et al.* 2000, 18), G. Laplace and N. Soler have emphasized the likelihood that the small assemblage from this layer may have resulted out of palimpsests of different occupations, one during the late Gravettian, another during the Aurignacian, based on the similarities of some (6) backed tools with Gravettian points (Canal i Roquet and Carbonell i Roura 1989; Carbonell *et al.* 1994). This 'Aurignacian' layer from the initial excavations conducted by A. Romaní between 1909 and 1929 ("layer 2") is preserved in remnant deposits – labeled "level A" – along the back wall of the central part of the rock-shelter as well as in the Coveta Nord, where it is stratified between two travertine horizons (Bischoff *et al.* 1994). "Level A" – al-

though only 2cm in thickness – produced "abundant faunal remains, dispersed charcoal and artifacts" (Bischoff *et al.* 1994: 544) and most likely represents a living floor sealed by the overlying travertine. Seven radiocarbon measurements on charcoals were obtained from the remnant "level A" deposits at three different places of the site all together. Five of these form a WM of  $36,644 \pm 373$   $^{14}\text{C}$  BP, with individual dates ranging from  $37,290 \pm 990$  (AA-7395) to  $36,390 \pm 629$  (WM of AA-8037A and AA-8037B)  $^{14}\text{C}$  BP. These dates are in strong contrast to two measurements from the radiocarbon laboratory in Waikato, New Zealand, which – due to possible contamination with younger carbon – are significantly younger (NZA-1817; NZA-1818). The same laboratory has also produced a date of  $36,590 \pm 640$   $^{14}\text{C}$  BP (NZA-2311, contained in the WM given above). All these dates are stratigraphically consistent with a further radiocarbon measurement of a charcoal sample that was embedded in the travertine (USGS-2839:  $36,600 \pm 1300$   $^{14}\text{C}$  BP; Bischoff *et al.* 1994). Furthermore, the great antiquity of these deposits is confirmed by U-series dating of this travertine (Bischoff *et al.* 1994; with corrected U/Th-ages of between 39.1 and 42.9 kyr BP given in Carbonell *et al.* 1994) (9). Concerning the difficulties with the archaeological assignment of "layer 2" Zilhão and d'Errico (1999), plead for the stratigraphical reliability of this sequence. Whereas it is highly likely that the radiocarbon measurements from "level A" material do indeed date the Aurignacian, stratigraphical mixing on top of the Aurignacian layer may also have played a role, close to the wall of the rock-shelter, and only a few metres away from the sampled area (as observed by A. Romaní).

Contra the interpretation of Zilhão and d'Errico (1999) the evidence from Castillo, L'Arbreda, and Romaní strongly indicates earliest Aurignacian presence in the North of the Iberian Peninsula before  $36,500$   $^{14}\text{C}$  BP (Fig. 10), confirmed by the consistency of both radiometric dating and stratigraphy. Insignificantly younger radiocarbon measurements – all on charcoal – derive from the lower part of layer XIII at La Viña (Ly-6390:  $36,500 \pm 750$   $^{14}\text{C}$  BP), from Isturitz (WM of two dates:  $35,490 \pm 413$   $^{14}\text{C}$  BP), and possibly at Tournal G (oldest date: Ly-1898:  $>35,800$   $^{14}\text{C}$  BP).

(9) Although less precise than radiocarbon dates, due to the high standard deviations, the results from U-series dating at Romaní approximately fit the ranges of the calibrated radiocarbon measurements (cf. CALPAL-2003: <http://www.calpal.de>).




Fig. 10. Aurignacian (Au.) and archaic Aurignacian (arc. Au.)  $>36,500$   $^{14}\text{C}$  BP in the northern part of the Iberian Peninsula and its relation to Latest Middle Palaeolithic (LMP) industries.

Towards the North (north of  $44.0^\circ$  N; Fig. 9) radiocarbon FAD's of the oldest regional Aurignacian turn out slightly younger (starting with  $35,400 \pm 1100$   $^{14}\text{C}$  BP [GifA-97186] in layer F of Caminade-Est; Tab. 1), and in the south of the Iberian Peninsula, the oldest Aurignacian is present in Beneito VIII (AA-1388:  $33,900 \pm 1100$   $^{14}\text{C}$  BP on charcoal). Further to the South and to the West radiocarbon dates for the Aurignacian are even younger than  $33,500$   $^{14}\text{C}$  BP (Vega Toscano 1990; Zilhão 1993), with the Portuguese site of Gato Preto, dated by TL to  $38.1 \pm 3.9$  kyr BP, representing the oldest Aurignacian in the SW of the Iberian Peninsula.

#### 4.2. ... and evidence for the latest Middle Palaeolithic

For SW Europe the “*Ebro-Frontier*”-model assumes LMP industries at times when Aurignacian people had already populated the North and Northeast of the Iberian Peninsula (Zilhão 1993; cf. Fig. 2). For the second line of arguments underlying the “*Ebro-Frontier*”-model it is thus important to verify the contemporaneity of LMP industries in the SW of the Iberian Peninsula with the earliest Aurignacian in the North and Northeast, i.e. evidence persisting significantly later than  $38,300$   $^{14}\text{C}$  BP. This is all the more difficult since Aurignacian layers in general overlay LMP ones and are subject to some reworking of deposits. Relying more on  $^{14}\text{C}$ -bone data for the MP/UP transition, Zilhão and d'Errico (1999: 10) note the “*apparent contradiction between stratigraphy and dating*” because the radiometric dates do imply a significant chronological overlap. They try to explain this contradiction

as “*simply an artifact of serious errors of method and interpretation in the use of radiometric results*”.

Due to the problems of possible sample contamination and the fact that – as a trend – radiocarbon charcoal data are shown to be more reliable than bone data, we will base our study solely on radiocarbon dates on charcoal as well as on the chronometric data available from the different *non*-radiocarbon methods of dating (e.g. TL/OSL, U-series; ESR, Tab. 2; Fig. 11), in order to trace reliable LMP LAD's in SW Europe

Regarding the charcoal-dated LMP (incl. Châtelperronian) of SW Europe only two sites – Fuentes de San Cristobal (Mousterian) and Morín (layer 10: Châtelperronian) – have produced finite radiocarbon age-determinations younger than  $38,300$   $^{14}\text{C}$  BP. At Morín the same sample (SI-951) has produced two contradictory radiocarbon measurements, one with an extremely high standard deviation of several thousand BP. These measurements along with a single sample (OxA-8591) for the LMP at Fuentes de San Cristobal with a high standard deviation ( $\pm 1900$ ), cannot be interpreted as reliable proof for a long persistence of the LMP in SW-Europe.

One sample from the basal LMP layer X at Beneito in the SE of the Iberian Peninsula has been dated to  $30,160 \pm 680$   $^{14}\text{C}$  BP (unknown lab.-no.), whereas a second sample (AA-1387) on charcoal, obtained from the same layer, dates to  $38,800 \pm 1900$   $^{14}\text{C}$  BP and is thus similar in age to the WM of the radiocarbon measurements available for the LMP layer 18b of El Castillo ( $38,679 \pm 744$ ). Sample AA-1388 from the overlaying Aurignacian level VIII at Beneito confirms the antiquity of the AA-1387 sample.

Other sites that have produced radiocarbon age-determinations younger than  $38,300$   $^{14}\text{C}$  BP have either been dated several decades ago (e.g. Ermitons, Quinçay, Tournal, Brugas) or the dates have been obtained from “*inadequate dating material*” (Zilhão 2000a: e.g. Columbeira) that is likely to be contaminated with modern carbon (Zilhão 1997: 35). Possible contamination is furthermore likely for some radiocarbon measurements from LMP layers that have been independently dated by *non*-radiocarbon methods. At Zafarraya (Hublin *et al.* 1995), for example, teeth of *Capra ibex* were dated by radiocarbon as well as by U-series methods, but due to a variety of reasons mostly related to the stratigraphical properties of this site, these samples are

## MIDDLE PALAEOLITHIC < 40.0 kyr 14C BP and/or < 43.5 kyr non-14C BP

LEVEL	CULTURE	WD	MAT.	METH.	LAB-NO.	DATE	STD+	STD-	t-val.
<b>14C on CHARCOAL</b>									
<b>St-Marcel (Bidon, Ardèche, France)</b>									
G2	MP	761	charcoal	14C	Ly-2901	>25000			
<b>Devil's Tower (Gibraltar)</b>									
3	MP	256	charcoal	14C	GrN-2488	>30000			
<b>Trozoul (Trébeurden, Bretagne, France)</b>									
hearth	MP?	505	charcoal	14C	Gif-1312	>35000			
<b>Los Moros I (Gabasa, Huesca, Spain)</b>									
a	MP	*	charcoal	14C	OxA-5671	>39000			
a+c	MP	*	charcoal	14C	OxA-5672	>45000			
* = Blasco Sánchez and Montes Ramírez 1997									
<b>Cueva Morin (Villanueva, Villaescusa, Cantabria, Spain)</b>									
10	Chat.	100	charcoal	14C	SI-951*	27777	577	577	
10	Chat.	99	charcoal	14C	SI-951-A*	35874	6777	6777	
* = identical sample									
<b>Fuentes de San Cristobal (Veracruz, Huesca, Spain)</b>									
	MP	*	charcoal	14C	OxA-8591	36000	1900	1900	
* = Rosell Ardèvol et al. 2000									
<b>Barbas III (Creysse, Dordogne, France)</b>									
4	MP	731	charcoal	14C	?	38300	500	500	0,37
4	MP	732	charcoal	14C	GifA-93050	43500	2200	2200	2,19
4	MP	731-732	charcoal	14C	weighted mean	38555	488		
<b>Castillo (Puente Viesgo, Cantabria, Spain)</b>									
18b-upper	MP	124	charcoal	14C	AA-2407	37700	1800	1800	0,50
18b-upper	MP	123	charcoal	14C	AA-2406	38500	1800	1800	0,09
18b	MP	127	charcoal	14C	OxA-2473	37100	2200	2200	0,68
18b	MP	126	charcoal	14C	OxA-2474	38500	1300	1300	0,12
18b-lower	MP	125	charcoal	14C	OxA-2475	40700	1600	1600	1,15
18b	MP	123-127	charcoal	14C	weighted mean	38679	744		
20b2	MP	128	charcoal	14C	GifA-89144	39300	1900	1900	0,48
20b2	MP	129	charcoal	14C	GifA-92506	43300	2900	2900	0,85
20b2	MP	128-129	charcoal	14C	weighted mean	40501	1589		
<b>L'Arbreda (Serinyà, Girona, Spain)</b>									
BE 116 (level I) sq. E2	MP	154	charcoal	14C	AA-3777	34100	750	750	
BE 116 (level I) sq. E2	MP	155	charcoal	14C	AA-3776	39400	1400	1400	0,92
BE 116 (level I) sq. E2	MP	156	charcoal	14C	AA-3778	41400	1600	1600	0,24
BE 116 (level I) sq. E2	MP	153	charcoal	14C	OxA-3731	44560	2400	2400	1,39
BE 116 (level I) sq. E2	MP	153+155-56	charcoal	14C	weighted mean	40961	965		
<b>Gorham's Cave (Gibraltar)</b>									
context 24*	transitional	239	charcoal	14C	OxA-7857	32280	420	420	
context 18**	transitional	240	charcoal	14C	OxA-7791	42200	1100	1100	
* = remnant combustion zone									
** = combustion zone									
<b>14C on CHARCOAL vs. 14C on BONE</b>									
<b>Ermítos (Sales de Llierca, Girona, Spain)</b>									
IV	MP	163	bone	14C	OxA-3725	33190	660	660	
IV	MP	164	charcoal	14C	CSIC-197/IAB-3	36430	1800	1800	
<b>L'Arbreda (Serinyà, Girona, Spain)</b>									
BE 116	MP	153	bone	14C	OxA-3731	44560	2400	2400	
BE 116	MP	154	charcoal	14C	AA-3777	34100	750	750	
BE 116	MP	155	charcoal	14C	AA-3776	39400	1400	1400	0,49
BE 116	MP	156	charcoal	14C	AA-3778	41400	1600	1600	0,59
BE 116	MP	155-156	charcoal	14C	weighted mean	40267	1054		
<b>14C on CHARCOAL vs. 14C on HUMUS</b>									
<b>Quincay (Quincay, Vienne, France)</b>									
black unit	Chat.	577	charcoal	14C	Ly-790	11910	200	200	
black unit	Chat.	578	humus	14C	Ly-791	20300	500	500	

Tab 2.

## 14C on CHARCOAL vs. 14C on UNKNOWN MATERIAL

**Beneito (Muro, Alicante, Spain)**

basal; X (D1)	MP	162	?	14C	?	30160	680	680	
basal; X (D1)	MP	161	charcoal	14C	AA-1387	38800	1900	1900	

## 14C on CHARCOAL vs. TL / OSL on SEDIMENT / BURNT SILEX

**Tournal (Bize, Aude, France)**

C (Sq. K29)	MP	749	charcoal	14C	Ly-1676	33600	1300	1300	
	MP	744	burnt silex?	TL?	Gif-?	33000	4000	4000	0,00
	MP	745	burnt silex?	TL?	Gif-?	33000	8000	8000	0,00
	MP	744-745	burnt silex?	TL?	weighted mean	33000	3578		

**Gorham's Cave (Gibraltar)**

context 22/22D*	MP	245	sediment	OSL	MacGor-3	35000	7000	7000	
context 22/22D*	MP	244	charcoal	14C	OxA-6075	45300	1700	1700	0,59
context 22/22D*	MP	246	charcoal	14C	OxA-7790	51700	3300	3300	1,39
context 22/22D*	MP	244+246	charcoal	14C	weighted mean	46642	1511		

\* = combustion zone (=G?)

**Brugas (Vallabrix, Gard, France)**

4	MP	679	charcoal	14C	Ly-2351	29000	860	860	
4	MP	678	charcoal	14C	Ly-2038	>32000			
4	MP	676	burnt silex	TL	Gif-?	58200	7500	7500	0,58
4	MP	677	burnt silex	TL	Gif-?	65900	9900	9900	0,27
4	MP	674	burnt silex	TL	?	63000	5800	5800	0,01
4	MP	675	burnt silex	TL	Gif-?	69000	10800	10800	0,55
4	MP	674-677	burnt silex	TL	weighted mean	63063	3843		

## 14C on CHARCOAL vs. ESR

**Castillo (Puente Viesgo, Cantabria, Spain)**

18c	MP	121	bone	ESR	?	36200	4100	4100	0,46
18c	MP	122	bone	ESR	?	39900	4600	4600	0,28
18c	MP	*	bone	ESR	?	40000	5000	5000	0,28
18c	MP	121-122+	bone	ESR	weighted mean	38427	2610		
18c	MP	120	charcoal	14C	GifA-89147	39500	2000	2000	0,36
18c	MP	119	charcoal	14C	OxA-2478	39800	1400	1400	0,29
18c	MP	116	charcoal	14C	AA-2405	40000	2100	2100	0,12
18c	MP	117	charcoal	14C	OxA-2476	40700	1500	1500	0,26
18c	MP	118	charcoal	14C	OxA-2477	41100	1700	1700	0,45
18c	MP	116-120	charcoal	14C	weighted mean	40621	750		

\* = Cabrera Valdés et al. 2001

## 14C on CHARCOAL / CARBONACEOUS EARTH vs. U-SERIES

**Columbeira (Bombarral, Portugal)**

16=7	MP	57	cc/ce*	14C	Gif-2703**	26400	750	750	
16=7	MP	58	tooth	U-series	SMU-235E1	35876	27299	35583	0,34
16=7	MP	59	tooth	U-series	SMU-238E1	54365	22240	27525	0,26
16=7	MP	58-59	tooth	U-series	weighted mean	46990	17240		
20=8	MP	54	cc/ce*	14C	Gif-2704**	28900	950	950	
20=8	MP	55	tooth	U-series	SMU-236E1***	60927	27405	35522	0,39
20=8	MP	56	tooth	U-series	SMU-236E1***	101487	38406	55919	0,61
20=8	MP	55-56	tooth	U-series	weighted mean	74620	22310		

\* = charcoal/carbonaceous earth

\*\* = inadequate dating material (Zilhao 2000)

\*\*\* = identical sample

## 14C on BONE / TEETH vs. U-SERIES

**Zafarraya (Alcaucin, Málaga, Spain)**

I (3-7)	MP	167	tooth	14C	Gif-9140-II*	29800	600	600	
I (3-7)	MP	168	tooth	U-series	Gif-9140-II*	25100	1300	1300	0,55
I (3-7)	MP	169	tooth	U-series	Gif-9140-II*	26900	2700	2700	0,30
I (3-7)	MP	170	tooth	U-series	Gif-9140-II*	31700	3600	3600	1,50
I (3-7)	MP	168-170	tooth	U-series	weighted mean	26038	1114		
I (8)	MP	172	tooth	U-series	Gif/LSM-9140-I*	31700	3600	3600	
I (8)	MP	171	tooth	14C	Gif/LSM-9140-I*	31800	550	550	
D	MP	173	tooth	U-series		33400	2000	2000	

\* = identical samples, respectively

Tab. 2 (Cont.).

## 14C on BONE vs. ESR

**Combe Sauniere (Sarriac-sur-l'Isle, Dordogne, France)**

X	Chat.	723	tooth	ESR	?	36400	2500	2500	
X	Chat.	722	bone	14C	OxA-6504	33000	900	900	2,30
X	Chat.	720	bone	14C	OxA-6503	35900	1100	1100	0,36
X	Chat.	721	bone	14C	OxA-6503*	38100	1000	1000	2,30
X	Chat.	720-722	bone	14C	weighted mean	35449	572		

\* = tripeptide

## 14C on SHELL vs. U-series

**Figueira Brava (Sesimbra, Portugal)**

2?	MP?	9	shell	14C	ICEN-387	30930	700	700	
2	MP	11	tooth	U-series	SMU-232E1*	30561	11759	10725	0,33
2	MP	12	tooth	U-series	SMU-232E2*	44806	15889	13959	0,50
2	MP	11-12	tooth	U-series	weighted mean	35600	9450		

\* = identical sample

## TL / OSL

**Conceicao (Alcochete, Portugal)**

C-topo	MP*	70	sediment	TL/OSL	QTSL-CNC-11	27200	2500	2500	
--------	-----	----	----------	--------	-------------	-------	------	------	--

\* = MP lies in layer D below C

**Roche a Pierrot (Saint-Cesaire, Charente-Maritime, France)**

10	MP	390	?	TL	?	40900	2500	2500	0,20
11	MP	391	?	TL	?	38200	3300	3300	0,55
12	MP	392	?	TL	?	42400	4800	4800	0,41
	MP	390-392	?	TL	weighted mean	40281	1840		

**Carigüela (Pinar, Granada, Spain)**

MP	189	silex	TL	TB-2	28000			
MP	187	silex	TL	TB-5	31000-35000			
MP	188	silex	TL	TB-1	32000			
MP	186	silex	TL	TB-12	39000			
MP	185	silex	TL	TB-3	46000			
MP	184	silex	TL	TB-9a	48000			

## TL vs. ESR

**Le Moustier (Le Moustier, Dordogne, France)**

K	MP/Chat.?	801	burnt silex	TL	Gif-?	42600	3700	3700	0,37
J	MP	800	burnt silex	TL	Gif-?	40300	2600	2600	0,23
I	MP	799	burnt silex	TL	Gif-?	40900	5000	5000	0,03
K-I	MP	799-801	burnt silex	ESR	weighted mean	41036	1957		
H[2a-7c]	MP	798	tooth	ESR: LU	?	41000	2600	2600	0,22
H[2a-7c]	MP	797	tooth	ESR: EU	?	39700	2400	2400	0,20
H[2a-7c]	MP	797-798	tooth	ESR	weighted mean	40298	1764		
H2-H9	MP	796	burnt silex	TL	Gif-?	42500	2000	2000	

**El Pendo (Escobedo de Camargo, Cantabria, Spain)**

26	MP	*		ESR	?	33700	1300	1300	
28	MP	*		ESR	?	14300	300	300	
32 (H)	MP	*		ESR	?	30500	300	300	
B	MP	**		ESR	?	<40000			
D	MP	**	burnt silex?	TL	?	83790	8291	8291	
F	MP	**	burnt silex?	TL	?	39626	3864	3864	

\* = Montes Barquin and Sanguino González 2001; \*\* = Montes Barquin 2000

## U-SERIES

**Gruta do Escoural (Évora, Montemor-o-Novo, Portugal)**

test 3a (90-100)	MP	33	tooth	U-series	SMU-248*	26400	5800	5500	1,33
test 3a (80-90)	MP	34	tooth	U-series	SMU-249*	39800	10000	9000	0,61
test 3a (60-70)	MP	35	tooth	U-series	SMU-250	48900	11000	10000	0,98
test 3a	MP	33-35	tooth	U-series	weighted mean	33070	4560		

\* = low U content (Zilhao 2000)

**Foz do Enxarrique (Ródão, Portugal)**

C	MP	6	tooth	U-series	SMU-225	32938	1055	1055	0,74
C	MP	7	tooth	U-series	SMU-226	34088	800	800	0,29
C	MP	8	tooth	U-series	SMU-224	34093	920	920	0,27
C	MP	6-8	tooth	U-series	weighted mean	33806	524		

Tab. 2 (Cont.).



<b>Almonda (Casais Martanes, Santarém, Portugal)</b>								
EVS cone	MP	26	tooth	U-series	SMU-231E1*	35000	2000	2000
					* = low 230Th/232Th ratio (Zilhao 2000)			
<b>Almonda (Casais Martanes, Santarém, Portugal)</b>								
EVS cone	MP	26	tooth	U-series	SMU-231E1*	35000	2000	2000
					* = low 230Th/232Th ratio (Zilhao 2000)			

**U-SERIES vs. ESR**

<b>Bajondillo (Torremolinos, Málaga, Spain)</b>									
basal level	MP	226	travertine	ESR: EU		25300	2530	2530	0,48
basal level	MP	227	travertine	ESR: LU		26500	3975	3975	0,04
basal level	MP	228	travertine	U-series		27300	1700	1700	0,30
basal level	MP	226-228		all	weighted mean	26658	1330		

Tab. 2 (Cont.). Radiometric dates for the SW European Middle Palaeolithic excluding radiocarbon measurements on bone samples. WD – No. in William Davies datelist (see annotation 1); MAT. – material dated; METH. – dating method; STD – standard deviation; *t-val.* – *t*-value. MP – Middle Palaeolithic; Chat. – Châtelperronian.

to be regarded “*highly questionable and/or irrelevant*” in contribution to the issue discussed (Pettitt and Pike 2001: 416). A ‘transitional industry’ from Gorham’s Cave may be due to stratigraphic palimpsests, since both radiocarbon measurements obtained on charcoal diverge by some 10,000 <sup>14</sup>C BP (OxA-7857 for context 24: 32,280 ± 420 <sup>14</sup>C BP vs. OxA-7791 for context 18: 42,000 ± 1100 <sup>14</sup>C BP). Stratigraphical inversion of layers most plausibly account for the El Pendo series (Montes Barquín 2000; Montes Barquín and Sanguino González 2001).

At Brugas and Columbeira *non*-radiocarbon age-determinations all dating older than 45.0 kyr ago have produced results significantly older than radiocarbon measurements. In contrast, two radiocarbon measurements on charcoal at Gorham’s Cave (context 22/22D) are much older than the OSL-dated sediment. At Figueira Brava a radiocarbon measurement on a *Patella* sp. shell has been dated to 30,930 ± 700 <sup>14</sup>C BP (ICEN-387). Although Zilhão (2000) attributes the date to the LMP level 2, its precise stratigraphic provenience remains unclear. Two U-series measurements of the




Fig. 11. Radiometric dates (‘last appearance data’: LAD’s) for the SW European Middle Palaeolithic, with radiocarbon measurements (left) sorted for material dated and given in kyr <sup>14</sup>C BP, when required as weighted means, and *non*-radiometric measurements (right) sorted for method of dating and given in kyr BP, when required as weighted means.

same sample gave strongly divergent results with extremely high standard deviations. Whereas the younger date was given as  $30.6 \pm 11.8$  kyr BP (SMU-232E1), the older one resulted in  $44.8 \pm 15.9$  kyr BP (SMU-232E2).

For the period under discussion in general, *non*-radiocarbon methods of dating have to be regarded as less reliable due to the often extremely high standard deviations, when compared with  $^{14}\text{C}$  (e.g. Schwarcz 2001). Even in cases where radiocarbon measurements appear in agreement with results obtained from the various *non*-radiocarbon dating techniques, this may be simply coincidental (10).

The LMP from Tournal C is dated to  $33,600 \pm 1300$   $^{14}\text{C}$  BP (Ly-1667) by radiocarbon and to  $33.0 \pm 3.6$  kyr BP by TL on burnt silex (WM of two dates), whereas a sample from the overlaying Aurignacian of level G is older than  $35,800$   $^{14}\text{C}$  BP (Ly-1898; Tavoso 1976; cf. Tab. 1). The Châtelperronian of Combe Sauniere has been dated by ESR ( $36.4 \pm 2.5$  kyr BP; Tab. 2) as well as by radiocarbon on two bone samples. Although the WM of the radiocarbon measurements ( $35,449 \pm 572$   $^{14}\text{C}$  BP) is largely in agreement with the date produced by ESR, the individual measurements are strongly divergent. Moreover, OxA-6503 gave a date of  $35,900 \pm 1100$   $^{14}\text{C}$  BP, whereas a tripeptide-measurement of the same sample resulted in  $38,100 \pm 1000$   $^{14}\text{C}$  BP, close to the date that we propose for the earliest Aurignacian in the North of the Iberian Peninsula (L'Arbreda; cf. Fig. 10).

*Non*-radiocarbon dates younger than 30.0 kyr BP have been obtained from Bajondillo (ESR and U-series) with a WM of  $26.7 \pm 1.3$  kyr BP and from Conceição (TL), for a level overlaying the LMP (QTSL-CNC-11:  $27.2 \pm 2.5$  kyr BP). TL-dates from Roche à Pierrot (St. Césaire) and Carigüela – although rather heterogeneous – indicate that LMP layers are older than 40.0 kyr BP. The same accounts for the TL and ESR-measurements obtained from the Le Moustier sequence.

A low  $^{230}\text{Th}/^{232}\text{Th}$ -ratio explains the relatively young age for the MP finds from the EVS-cone at Almonda (Zilhão 2000b), and low U-contents for the samples SMU-248 and SMU-249 at Gruta do Escoural (Zilhão 2000b) point to the higher reliability

of the SMU-250-sample, giving an age of  $48.9 \pm 11.0$  kyr BP.

The only consistent series of U/Th-measurements comes from Foz do Enxarrique C with an WM of  $33.8 \pm 0.5$  kyr BP. Unfortunately the relation between LMP lithic artefacts and the faunal material dated at this site remains unclear (cf. Zilhão 1997).

## 5. CONCLUSIONS AND DISCUSSION

Our analyses of the radiometric evidence for the MP/UP transition in SW Europe has augmented a range of criteria for quality control in data mining as well as in terms of advanced numeric data processing. The results emphasize sample material, problems of possible adhering contamination, and on measuring precision. Comparisons of bone and charcoal radiocarbon dates points to extreme discrepancies between these sample categories, with bone dates being systematically too young (Jöris *et al.* 2001; cf. Fig. 4-5). These findings do not refute the reliability of radiocarbon measurements of bone in general, but call for circumspection in evaluation of quality and reliability of such dates. Against this background, bone dates may contribute less than initially hoped to the understanding of the chronological issues surrounding the transition from the LMP to the EUP in Europe, and in particular we must remain cautious when interpreting  $^{14}\text{C}$ -bone data close to the detection limits of the radiocarbon method (Fig. 3).

### 5.1. On the age of the MP/UP transition in SW Europe

Evaluation of LMP radiometric dates from SW Europe  $<40,000$   $^{14}\text{C}$  BP and  $<43.5$  kyr *non*- $^{14}\text{C}$  BP (Tab. 2; Fig. 11) has shown that minimal reliability – if any – is given for dates younger than  $38,300$   $^{14}\text{C}$  BP. Based on radiocarbon dates obtained on charcoal, SW European LMP LAD's range between  $35,900$   $^{14}\text{C}$  BP (Morín, level 10: SI-951-A) and  $38,800$   $^{14}\text{C}$  BP (Beneito: AA-1387), with a WM of Beneito-, Castillo- (level 18b), Barbas-, Ermitons-, Fuentes de San Cristobal- and Morín-LAD's of  $38,391 \pm 381$   $^{14}\text{C}$  BP (Fig. 11, left). This SW European LMP LAD is statistically identical with the FAD of the archaic Aurignacian at L'Arbreda (WM:  $38,307 \pm 552$   $^{14}\text{C}$  BP) and is in overall agree-

(10) Strikingly, at some sites radiocarbon measurements and *non*-radiocarbon methods have produced identical ages – at least within the errors contained in the standard deviations (e.g. Combe Sauniere, Zafarraya). This is the more surprising, since the calibration records that are available today, imply that radiocarbon age-determinations underestimate the calendric scale for several thousand BP within the time-period under discussion.

14C-Age [BP]	Cal Age [cal BC] 68% (95%)	Cal Age [cal BP] 68% (95%)	Notes
40000 ± 800	40550 ± 430 (41410–39690)	42500 ± 430 (43360–41640)	(1)
38300 ± 766	39890 ± 290 (40470–39310)	41840 ± 290 (42420–41260)	(2)
36500 ± 730	38610 ± 940 (40490–36730)	40560 ± 940 (42440–38680)	(3)
35400 ± 708	37950 ± 1080 (40110–35790)	39900 ± 1080 (42060–37740)	(4)
33500 ± 670	36360 ± 1710 (39780–32940)	38310 ± 1710 (41730–34890)	(5)
30000 ± 600	31900 ± 700 (33300–30500)	33850 ± 700 (35250–32450)	(1)
28000 ± 560	30190 ± 690 (31570–28810)	32140 ± 690 (33520–30760)	(6)

- (1) commonly accepted limits of the MP/UP transition  
 (2) FAD for the Aurignacian in SW Europe, this work  
 (3) FAD for the Aurignacian in SW Europe after Zilhão and d'Errico (1999)  
 (4) FAD for the Aurignacian in France, this work  
 (5) FAD for the Aurignacian in the SW of the Iberian Peninsula after Vega Toscano (1990) and Zilhão (1993)  
 (6) average of youngest dates for the European MP after various authors  
 (e.g.: Bocquet-Appel and Demars 2000; Zilhão 1993; cf. Hublin et al. 1995)

Tab 3. Calibration of seven fictive radiocarbon measurements [BP] spanning around the Middle to Upper Palaeolithic (MP/UP) transition with typical errors in the range of 2% of total radiocarbon BP. Calibration was established using the CAL-PAL-2003 data set (see annotation 8). FAD – First appearance data.

ment with LMP LAD's available for other parts of Europe (Fig. 6), implying a sudden and simultaneous ending of the MP all over Europe.

To summarize, we note

- that the LMP discontinues over the whole of SW Europe at ca. 38,300 <sup>14</sup>C BP,
- at a time for which similar LMP LAD's and Aurignacian FAD's can be fixed in various parts of Europe (Fig. 6).

In terms of calibrated radiocarbon-‘years’ this transition falls into the shift from the interstadial conditions of GI 11 (GI = *Greenland interstadial*) to the stadial ones of GS 11 (GS = *Greenland stadial*) at around 39.9 kyr cal BC (Fig. 8; cf. Tab. 3) (11).

We can therefore confirm the high antiquity of the oldest Aurignacian in the North of the Iberian Peninsula, i.e. Cantabria and Catalunya, but - in contrast to the radiocarbon record – evidence from the *non*-radiocarbon methods of dating for the LMP (Tab. 2; Fig. 11, bottom) does not withstand criticism and the data available are too weak to prove LMP persistence in the southern half of the Iberian Peninsula until 33,500 <sup>14</sup>C BP or later, at least not until the Aurignacian of Gato Preto.

(11) Whereas our FAD's for the Aurignacian of SW Europe preceded those proposed by Zilhão and d'Errico (1999) by some 1800 <sup>14</sup>C BP, in calibrated terms FAD-differences would be in the range of only 1.3 kyr cal, still placing the transition into GS 11.

Although the available radiometric dates represent regionally different patterns of hominid presence, we cannot confirm the existence of an “*Ebro Frontier*”, that geographically distinguishes between Aurignacian industries to the NE and LMP ones in the SW against the background of these data. Rather our data implies that the southern part of the Iberian Peninsula was already void of MP hominids long before the appearance of the first Aurignacians.

## 5.2. Climate controlled population dynamics?

Based on radiocarbon evidence, the oldest Aurignacian sites of SW Europe cluster in the North of the Iberian Peninsula (Fig. 12: A; cf. Fig. 9), while the southern French Aurignacian does not date to before 35,400 <sup>14</sup>C BP and Aurignacian sites from the southern part of the Iberian Peninsula do not pre-date 33,500 <sup>14</sup>C BP (Vega Toscano 1990; Zilhão 1993). This pattern may best be explained by a northward spread (Fig. 12: B) of Aurignacian populations in a severe cold phase (GS 9 = Heinrich-event 4: H4; Fig. 8) shortly after GS 11. In a recent study F. d'Errico and M. F. Sánchez Goñi (2003) have characterized the highly arid desert-steppe-like H4-environments found over large areas of the Iberian Peninsula as inhospitable: conditions that




Fig. 12. Model of regional population shifts (arrows) around the transition from the Latest Middle Palaeolithic (LMP) to the Aurignacian in SW Europe (clockwise: LMP-A-B-C; cf. Fig. 8), covering the time span of *ca.* 42,000 – 28,000 <sup>14</sup>C BP over a period of strong climatic fluctuations during the last interpleniglacial (OIS 3; cf. Fig. 2, 7-8). GS – Greenland stadial.

may have triggered a northward shift of populations (Fig. 8, 12: B), followed by a further expansion into the NE as well as into the SW at the onset of GI 8 (Fig. 12: C).

### 5.3. Makers of the Aurignacian

Based on the chronometric dates that cover the European MP/UP transition there is neither radio-

metric nor stratigraphic indication for incoming populations that may have colonized Europe from East to West (12), nor is there reliable proof for a late persistence of LMP younger than 38,300 <sup>14</sup>C BP. Against these data, the geographical pattern of both LMP LAD's and Aurignacian FAD's (i.e. the MP/UP transition) can be best explained by a contraction of regional population dispersal into glacial refugia during GS 11. Such an interpretation would imply local *in situ* developments (13) of Aurignacian industries simultaneously in two different areas of Europe, separating the SW European Aurignacian without leaf-points from that of south-central and southeastern European sites which are characterized by the addition of a few of these artefact types. Consequently, this would imply that Neanderthals did indeed produce at least the earliest Aurignacian industries (cf. Churchill and Smith 2000).

Since the hominid fossil evidence is sparse and not entirely unambiguous during the Aurignacian, it would appear that unambiguous evidence of anatomically modern humans is not known prior to the European Middle UP.

## ACKNOWLEDGEMENTS

We thank V. Cabrera Valdés and A. F. Bernaldo de Quirós for permission to study the bone industry from the new excavations at El Castillo. We kindly acknowledge F. d'Errico, A. J. M. Maíllo Fernández, I. Castanedo Tapia, M. Street, St. Wenzel and J. Zilhão for fruitful discussions and contributions to this paper. Furthermore we thank F. d'Errico and W. Davies for supplying literature and data prior to publication. N. Barton, O. Bar-Yosef, C. Gamble, Chr. Stringer and – especially – E. Turner we thank for valuable comments on an earlier draft of this article.

## BIBLIOGRAPHY

- ALLSWORTH-JONES, Ph. 1986: *The Szeletian and the Transition from Middle to Upper Palaeolithic in Central Europe*. Clarendon Press, Oxford.
- (12) An infinite radiocarbon date (> 35,000 <sup>14</sup>C BP) obtained from a newly discovered human mandible from Romania (Gibbons 2003: 894) does not prove the presence of Aurignacian early modern humans in Europe. Since the new fossil displays both anatomically modern human and archaic features one has to await further results from palaeoanthropology as well as new datations.
- (13) Although such *in situ* development cannot be proven by the LMP industry of El Castillo 18 that Cabrera Valdés *et al.* (2001) attribute to an Aurignacian with archaic elements.
- ASMUS, G. 1964: "Kritische Bemerkungen und neue Gesichtspunkte zur jungpaläolithischen Bestattung von Combe Capelle, Périgord". *Eiszeitalter und Gegenwart* 15: 181-6.
- BECK, J. W.; RICHARDS, D. A.; EDWARDS, R. L.; SILVERMAN, B. W.; SMART, P. L.; DONAHUE, D. J.; HERERRA-OSTERHELD, S.; BURR, G. S.; CALSOYAS; L.; JULL, A. J. T. and BIDDULPH, D. 2001: "Extremely Large Variations of Atmospheric <sup>14</sup>C Concentration During the Last Glacial Period". *Science* 292: 2453-8.
- BISCHOFF, J. L., LUDWIG, K., GARCÍA, J. F., CARBONELL, E., VAQUERO, M., STAFFORD, TH. W. and JULL, A. J. T. 1994: "Dating of the Basal Aurignacian Sandwich at Abric Romani (Catalunya, Spain) by Radiocarbon and Uranium-Series". *Journal of Archaeological Science* 21 (4): 541-51.
- BISCHOFF, J. L., SOLER, N., MAROTO, J. and JULIA, R. 1989: "Abrupt Mousterian/Aurignacian Boundary at c. 40 ka bp: Accelerator <sup>14</sup>C Dates from l'Arbreda Cave (Catalunya, Spain)". *Journal of Archaeological Science* 16: 563-76.
- BLASCO SÁNCHEZ, M. F. and MONTES RAMÍREZ, L. 1997: "Los helénidos del yacimiento musteriense de Gabasa 1". *BOLSKAN* 14: 9-27.
- BODU, P. 1990: "L'application de la méthode des remontages à l'étude du matériel lithique des premiers niveaux châtelperroniens de la Grotte du Renne à Arcy-sur-Cure (Yonne)". In C. Farizy (dir.): *Paléolithique moyen récent et Paléolithique supérieur ancien en Europe. Ruptures et transitions: examen critique des documents archéologiques*. Actes du Colloque international de Nemours (9-11 Mai 1988). Mémoires du Musée de Préhistoire d'Ile de France 3. Nemours: 309-12.
- BOLUS, M. and RÜCK, O. 2000: "Eine Blattspitze aus Wittislingen, Lkr. Dillingen a. d. Donau (Bayern). Zur südwestlichen Verbreitungsgrenze spätmittelpaläolithischer Blattspitzeninventare". *Archäologisches Korrespondenzblatt* 30: 165-72.
- BOQUET-APPEL, J.-P. and DEMARS, P. Y. 2000: "Neanderthal Contraction and Modern Human Colonization of Europe". *Antiquity* 74: 544-52.
- BORDES, J. G. 1998: *L'Aurignacian 0 en Périgord: analyse des données. Un exemple d'application d'une méthode de quantification des remontages d'intérêt stratigraphique: Caminade Est, couche G*. Diplôme d'Etudes Approfondies. Université Bordeaux I (manuscrit).
- BOSINSKI, G. 1987: "Die große Zeit der Eiszeitjäger. Europa zwischen 40.000 und 10.000 v. Chr.". *Jahrbuch des Römisch-Germanischen Zentralmuseums Mainz* 34 (1): 3-139.
- 1990: *Homo Sapiens. L'histoire des chasseurs du Paléolithique supérieur en Europe* (40.000-10.000 av. J.-C.). Errance. Paris.
- CABRERA VALDÉS, V. 1984: *El yacimiento de la cueva*

- de "El Castillo". *Puente Viesgo, Santander*. Bibliotheca Praehistorica Hispana XXII. C.S.I.C. Madrid.
- CABRERA VALDÉS, V. and BISCHOFF, J. L. 1989: "Accelerator C-14 Dates for Early Upper Paleolithic (Basal Aurignacian) at El Castillo Cave (Spain)". *Journal of Archaeological Science* 16: 577-84.
- CABRERA VALDÉS, V., MAÍLLO FERNÁNDEZ, J. M., LLORET, M. and BERNALDO DE QUIRÓS, F. 2001: "La transition vers le Paléolithique supérieur dans la grotte du Castillo (Cantabrie, Espagne): la couche 18". *L'Anthropologie* 105 (4): 505-32.
- CABRERA VALDÉS, V., PIKE-TAY, A., LLORET, M. and BERNALDO DE QUIRÓS, F. 2000: "Continuity Patterns in the Middle-Upper Palaeolithic Transition on Cantabrian Spain". In C. B. Stringer, R. N. E. Barton and J. C. Finlayson (eds.): *Neanderthals on the Edge*. Papers from a Conference Marking the 150<sup>th</sup> Anniversary of the Forbes' Quarry Discovery. Oxbow Books. Gibraltar: 85-93.
- CABRERA VALDÉS, V., VALLADAS, H., BERNALDO DE QUIRÓS, F. and HOYOS, M. 1996: "La transition Paléolithique moyen-Paléolithique supérieur à El Castillo (Cantabrie): Nouvelles datations per le carbone-14". *Comptes-Rendus de l'Académie des Sciences de Paris* 322 (2): 1093-8.
- CANAL I ROQUET, J. and CARBONELL I ROURA, R. 1989: *Catalunya paleolítica*. Patronat Francesc Eiximenis. Girona.
- CARBONELL, E., GIRALT, S. and VAQUERO, M. 1994: "Abric Romaní (Capellades, Barcelone, Espagne): une importante séquence anthropisée au Pléistocène supérieur". *Bulletin de la Société Préhistorique Française* 91 (1): 47-55.
- CARBONELL, E., M. VAQUERO, J. MAROTO, J., J. M. RANDO and MALLOL, C. 2000: "A Geographic Perspective on the Middle to Upper Palaeolithic Transition in the Iberian Peninsula". In O. Bar-Yosef and D. Pilbeam (eds.): *The Geography of Neandertals and Modern Humans in Europe and the Greater Mediterranean*. Peabody Museum of Archaeology and Ethnology. Harvard University. Cambridge: 5-34.
- CHURCHILL, S. E. and SMITH, FR. H. 2000: "Markers of the Early Aurignacian of Europe". *Yearbook of Physical Anthropology* 43: 61-115.
- CLARK, J. D.; BEYENE, Y.; WOLDEGABRIEL, G.; HART, W. K.; RENNE, P. R.; GILBERT, H.; DEFLEUR, A.; SUWA, G.; KATOH, S.; LUDWIG, K.; BOISSERIE, J.-R.; ASFAW, B. and WHITE, T. D. 2003: "Stratigraphic, Chronological and Behavioural Contexts of Pleistocene *Homo sapiens* from Middle Awash, Ethiopia". *Science* 423: 747-52.
- CONARD, N. J. and BOLUS, M. 2003: "Radiocarbon Dating the Appearance of Modern Humans and Timing of Cultural Innovations in Europe: New Results and New Challenges". *Journal of Human Evolution* 44: 331-71.
- DAVIES, W. 2001: "A Very Model of a Modern Human Industry: New Perspectives on the Origins and Spread of the Aurignacian in Europe". *Proceedings of the Prehistoric Society* 67: 195-217.
- DEMARS, P.Y. 1996: "La place du Piage et de Roc-de-Combe (Lot) dans la transition du Paleolithique moyen au Paléolithique supérieur". *Bulletin. Préhistorique du Sud-Ouest, Nouvelles Études* 3: 11-35.
- d'ERRICO, F. and SÁNCHEZ GOÑI, M. F. 2003: "Neanderthal Extinction and the Millennial Scale Climatic Variability of OIS 3". *Quaternary Science Reviews* (e. p.).
- d'ERRICO, F.; ZILHÃO, J.; BAFFIER, D.; JULIEN, M. and PELEGRIN, J. 1998: "Neanderthal Acculturation in Western Europe? A Critical Review of the Evidence and its Interpretation". *Current Anthropology* 39. Supplement, S-1-S44.
- DJINDJIAN, F. 1999: "Datations <sup>14</sup>C du Paléolithique supérieur européen. Bilan et perspectives". In J. Evin, J., C. Oberlin, J. P. Daugas and J. F. Salles (eds.): *Actes du 3ème Congr. Int. "Archéologie et <sup>14</sup>C": Rev. d'Archéométrie Suppl. 1999 et Soc. Préhist. Franç. Mémoire* 26. Lyon: 171-9.
- DJINDJIAN, F.; KOZŁOWSKI, J. and OTTE, M. 1999: "Le paléolithique supérieur en Europe". Armand Colin. Paris 1999.
- FEDELE, F. G.; GIACCIO, B.; ISAIA, R. and ORSI, G. 2002: "Ecosystem Impact of the Campagnian Ignimbrite Eruption in Late Pleistocene Europe". *Quaternary Research* 57: 420-4.
- GAMBIER, D. 1997: "Modern Humans at the Beginning of the Upper Palaeolithic in France". In G. A. Clark and C. M. Willwermet (eds.): *Conceptual Issues in Modern Human Origins Research*. Aldine de Gruyter: 117-131.
- GEYH, M. R. 1969: "Versuch einer chronologischen Gliederung des marinen Holozän an der Nordseeküste mit Hilfe der statistischen Auswertung von <sup>14</sup>C-Daten". *Zeitschrift der Deutschen Geologischen Gesellschaft* 188: 351-60.
- GIBBONS, A. 2003: "Meeting: Physical Anthropology and Palaeoanthropology". *Science* 300: 893-4.
- GILLESPIE, R. 1997: "Burnt and Unburnt Carbon: Dating Charcoal and Burnt Bone from the Willandra Lakes, Australia". *Radiocarbon* 39 (3): 239-50.
- GIOIA, P. 1990: "An aspect of the transition between Middle and Upper Palaeolithic in Italy: the Uluzzian". In: C. Farizy (dir.): *Paléolithique moyen récent et Paléolithique supérieur ancien en Europe. Ruptures et transitions: examen critique des documents archéologiques*. Actes du Colloque international de Nemours (9-11 Mai 1988). Mémoires du Musée de Préhistoire d'Île de France 3. Nemours: 241-50.
- GOUEDO, J.-M. 1990: "Les technologies lithiques du Châtelperronien de la couche X de la grotte du Renne d'Arcy-sur-Cure (Yonne)". In C. Farizy (dir.): *Paléolithique moyen récent et Paléolithique supérieur ancien en Europe. Ruptures et transitions: examen critique des*

- documents archéologiques*. Actes du Colloque international de Nemours (9-11 Mai 1988). Mémoires du Musée de Préhistoire d'Ile de France 3. Nemours: 305-8.
- GROOTES, P. M. 1977: *Thermal Diffusion Isotopic Enrichment and Radiocarbon Dating Beyond 50.000 Years BP*. Groningen.
- HEDGES, R. and PETTITT, P. 1999: "On the Validity of Archaeological Radiocarbon Dates beyond 30,000 Years BP". In J. Evin, J. C. Oberlin, J. P. Daugas and J. F. Salles (eds.): *Actes du 3ème Congr. Int. "Archéologie et <sup>14</sup>C"*: *Rev. d'Archéométrie Suppl. 1999 et Soc. Préhist. Franc. Mémoire* 26. Lyon: 137-41.
- HENRY-GAMBIER, D. 2002: "Les fossiles de Cro-Magnon (Les Eyzies-de-Tayac, Dordogne): Nouvelles données sur leur position chronologique et leur attribution culturelle". *Paleo* 14: 201-4.
- HUBLIN, J.-J.; BARROSO RUIZ, C.; MEDINA LARA, P.; FONTUGNE, M. and REYSS, J.L. 1995: "The Mousterian Site of Zafarraya (Andalucia, Spain): Dating and Implications on the Paleolithic Peopling Processes of Western Europe". *Comptes-Rendus de l'Académie des Sciences de Paris* 321 (2 a): 931-7.
- HUBLIN, J. J.; SPOOR, F.; BRAUN, F.; ZONNEVELD, F. and CONDEMI, S. 1996: "A Late Neanderthal Associated with Upper Palaeolithic Artefacts". *Nature* 381: 224-6.
- JÖRIS, O. 2003: "Zur chronostratigraphischen Stellung der spätmittelpaläolithischen Keilmessergruppen. Der Versuch einer kulturgeographischen Abgrenzung einer mittelpaläolithischen Formengruppe und ihr europäischer Kontext". *Ber. Röm.-German. Komm.* 84 (e.p.).
- JÖRIS, O.; ÁLVAREZ FERNÁNDEZ, E. and WENINGER, B. 2001: "Radiocarbon Evidence of the Middle to Upper Palaeolithic Transition on the Iberian Peninsula". XIVe Congrès de l'UISPP, 2-8 September (Pre-Prints), Liège: 141.
- JÖRIS, O. and WENINGER, B. 1998: "Extension of the <sup>14</sup>C Calibration Curve to ca. 40,000 calBC by Synchronizing Greenland <sup>18</sup>O/<sup>16</sup>O Ice Core Records and North Atlantic Foraminifera Profiles: a Comparison with U/Th Coral Data". *Radiocarbon* 40: 495-504.
- 1999 a: "Possibilities of Calendric Conversion of Radiocarbon Data for the Glacial Periods". In J. Evin, J. C. Oberlin, J. P. Daugas and J. F. Salles (eds.): *Actes du 3ème Congr. Int. "Archéologie et <sup>14</sup>C"*. *Rev. d'Archéométrie Suppl. 1999 et Soc. Préhist. Franc. Mémoire* 26. Lyon: 87-92.
  - 1999 b: "Calendric Age-Conversion of Glacial Radiocarbon Data at the Transition from the Middle to Upper Palaeolithic in Europe". *Bull. Soc. Préhist. Luxembourgeoise* 18: 43-55.
  - 2000: "Approaching the Calendric Age Dimension at the Transition from the Middle to Upper Palaeolithic in Europe". In J. Orschied and G.-Chr. Weniger (eds.): *Neanderthals and Modern Humans - Discussing the Transition: Central and Eastern Europe from 50.000-30.000 BP*. Wissenschaftliche Schriften des Neanderthal Museums 2. Mettmann: 13-9.
- KITAGAWA, H. and VANDER PLICHT, J. 2000: "Atmospheric Radiocarbon Beyond 11.900 calBP from Lake Suigetsu Laminated Sediments". *Radiocarbon* 42: 369-80.
- KRINGS, M.; GEISERT, H.; SCHMITH, R.; KRAINITZKI, H. and PÄÄBO, Sv. 1999: "DNA Sequence of the Mitochondrial Hypervariable Region II from the Neanderthal Type Specimen". *Proc. Nat. Acad. Sci. U.S.A.* 96: 5581-5.
- KRINGS, M.; STONE, A.; SCHMITZ, R. W.; KRAINITZKI, H.; STONEKING, M. and PÄÄBO, Sv. 1997: "Neanderthal DNA Sequences and the Origin of Modern Humans". *Cell* 90: 19-30.
- LANTING, J. N. and A. L. BRINDLEY 1999: "Fechando hueso cremado: la base científica". *Trabajos de Prehistoria* 56 (2): 137-40.
- MAÍLLO FERNÁNDEZ, J. M. 2003: *La transición Paleolítico Medio-Superior en Cantabria: análisis tecnológico de la industria lítica de Cueva Morín*. Doctoral thesis UNED, Madrid.
- MONTES BARQUÍN, R. 2000: "Actuaciones arqueológicas en la cueva de El Pendo (Escobedo de Camargo)". En R. Ontañón Peredo (Coord.): *Actuaciones Arqueológicas en Cantabria, 1984-1999*. Gobierno de Cantabria, Consejería de Cultura. Santander: 253-8.
- MONTES BARQUÍN, R. and SANGUINO GONZÁLEZ, J. 2001: *La Cueva de El Pendo. Actuaciones Arqueológicas 1994-2000*. Consejería de Cultura, Turismo y Deporte del Gobierno de Cantabria. Santander.
- MOOK, W. G. 1983: "<sup>14</sup>C Calibration Curves Depending on Sample Time Width". *Proc. Symp. <sup>14</sup>C and Archaeology* 8: 571-625.
- OVCHINNIKOV, I. V.; GÖTHERSTRÖM, A.; ROMANOVA, G. P.; KHARITONOV, V. M.; LIDÉN, K. and GOODWIN, W. 2000: "Molecular Analysis of Neanderthal DNA from the Northern Caucasus". *Nature* 404: 490-3.
- PALMA DI CESNOLA, A. 1993: *Il Paleolitico superiore in Italia*. Firenze 1993.
- PETTITT, P. 1999: "Neanderthal Extinction: Radiocarbon Chronology, Problems, Prospects and an Interpretation of the Existing Data". In J. Evin, J. C. Oberlin, J. P. Daugas and J. F. Salles (eds.): *Actes du 3ème Congr. Int. Archéologie et <sup>14</sup>C"*. *Rev. d'Archéométrie Suppl. 1999 et Soc. Préhist. Franc. Mémoire* 26. Lyon: 165-70.
- PETTITT, P. B. and PIKE, A. W. G. 2001: "Blind in a Cloud of Data. Problems with the Chronology of Neanderthal Extinction and Anatomically Modern Human Expansion". *Antiquity* 75: 415-20.
- van der PLICHT, J. 2002: "Calibration of the <sup>14</sup>C Time Scale: Towards the Complete Dating Range". *Netherlands Journal of Geosciences / Geologie en Mijnbouw* 81: 85-96.
- RIEK, G. 1932: "Paläolithische Station mit Tierplastiken und

- menschlichen Skelettresten bei Stetten ob Lontal". *Germania* 16: 1-8.
- 1934: *Die Eiszeitjägerstation am Vogelherd im Lontal, I: Die Kulturen*. Akademische Buchhandlung Franz F. Heine. Tübingen.
- RIGAUD, J. Ph. 2001: "A propos de la contemporanéité du Castelperronien et de l'Aurignacien ancien dans le nord-est de l'Aquitaine: une révision des données et ses implications". In J. Zilhão, Th. Aubry and A. F. Carvalho (eds.): *Les premières hommes modernes de la Péninsule Ibérique*. Actes du Colloque de la Commission VIII de l'UISPP (Vila Nova de Foz Cõa, 22-24 Octobre 1998). Instituto Português de Arqueologia. Lisboa: 61-8.
- RINK, J.; SCHWARCZ, H.P.; LEE, H.K.; CABRERA VALDÉS, V.; BERNALDO DE QUIRÓS, F. and HOYOS, M. 1996: "ESR Dating of Tooth Enamel: Comparison with AMS C-14 at El Castillo Cave, Spain". *Journal of Archaeological Science* 23: 945-51.
- ROSELL ARDÈVOL, J.; HUGUET PÁMIÉS, R.; AÈME-NE, M.; ANGELUCCI, D. E.; CANALS SOLOMÓ, A.; PASTÓ MARÍN, I. and RODRÍGUEZ ÁLVAREZ, J. P. 2000: "El yacimiento de las Fuentes de San Cristóbal (Veracruz, Huesca): un nuevo enclave del Paleolítico medio en el Prepirineo". En N. Bicho, E. Carbonell i Roura, B. Hockett, A. Moure Romanillo, L. Raposo, M. Santonja Gómez and G. Vega Toscano (Coor.): *Actas del 3º Congreso de Arqueología Peninsular* (Vila Real, 21-27 de Setembro de 1999) Vol. 2: *Paleolítico da Península Ibérica*. ADECAP-UTAD. Porto: 235-49.
- SCHWARCZ, H. P. 2001: "Dating Bones and Teeth: The Beautiful and the Dangerous". In Ph. V. Tobias, M. A. Raath, J. Moggi-Cecchi and G. A. Doyle (eds.): *Humanity from African Naissance to Coming Millennium*. Colloquia in Human Biology and Palaeanthropology. Firenze University Press. Firenze: 249-56.
- SINITSYN, A. A.; HAESAERTS, P., DAMBLON, F.; VAN DER PLICH, J. and FORMAN, S. L. 2002: "New Absolute Dates in Radiocarbon Sequences of Kostenki 14 (Markina Gora)". *Oso 'nmosti Raebitija Berchiego Paleolita Vostoëioj Evropy. Kostenki v kontekste paleolita Evrazii. Trudy kostenkovskoj ekspedicii*. Institut Istorii Material'noj Kultury RAS, Serija: Issledovani-ja I. Sankt Petersburg: 250-5.
- SONNEVILLE-BORDES, D. 1972: "Environnement et culture de l'homme du Périgordien ancien dans le sud-ouest de la France: données récentes". In F. Bordes (ed.): *The Origin of Homo Sapiens*. Proceedings of the Paris Symposium, 2-5 September 1969. UNESCO. Paris: 141-6.
- STRINGER, Chr. 2003: "Human evolution: Out of Ethiopia". *Science* 423: 692-5.
- STRINGER, C. B. and ANDREWS, P. 1988: "Genetic and Fossil Evidence for the Origin of Modern Humans". *Science* 239: 1263-8.
- STUIVER, M. and GROOTES, P. M. 2000: "GISP2 Oxygen Isotope Ratios". *Quaternary Research* 53 : 277-84.
- TAVOSO, A. 1976: "La Grotte Tournal ou Grande Grotte de Bize". In H. de Lumley (dir.): *Provence et Languedoc Médierranéen sites Paléolithiques et Néolithiques*. Livret-Guide de l'Excursion C2, IX Congrès UISPP (Nice, 13-18 Septembre 1976). Nice: 232-9.
- VEGA TOSCANO, L. G. 1990: "La fin du Paléolithique moyen au Sud de l'Espagne. Ses implications dans le contexte de la Péninsule Ibérique". In C. Farizy (ed.): *Paléolithique moyen récent et Paléolithique supérieur ancien en Europe*. 3. Mémoires du Musée de Préhistoire d'Ile de France. Nemours: 169-76.
- 1993: "El tránsito del Paleolítico Medio al Paleolítico Superior en el sur de la Península Ibérica". In V. Cabrera Valdés (ed.): *El origen del Hombre Moderno en el suroeste de Europa*. UNED. Madrid: 147-70.
- VOELKER, A.; SARNTHEIN, M.; GROOTES, P. M.; ERLKENKEUSER, H.; LAJ, C.; MAZAUD, A.; NADEAU, M. J. and SCHLEICHER, M. 1998: "Correlation of Marine <sup>14</sup>C-ages from the Nordic Seas with the GISP2 Isotopic Record: Implications for Radiocarbon Calibration Beyond 25 ka BP". *Radiocarbon* 40: 517-34.
- VOELKER, A. H. L.; GROOTES, P. M.; NADEAU, M. J. and SARNTHEIN, M. 2000: "<sup>14</sup>C Levels in the Icelandic Sea from 25-53 ka and their Link to the Earth's Magnetic Field Intensity". *Radiocarbon* 42: 437-52.
- WHITE, T.; ASFAW, B.; DEGUSTA, D.; GILBERT, H.; RICHARDS, G. D.; SUWA, G. and HOWELL, F. C. 2003: "Pleistocene *Homo sapiens* from Middle Awash, Ethiopia". *Science* 423: 742-7.
- ZILHÃO, J. 1993: "Le passage du Paléolithique Moyen au Paléolithique Supérieur dans le Portugal". In V. Cabrera Valdés (ed.): *El origen del Hombre Moderno en el suroeste de Europa*. UNED. Madrid: 127-45.
- 1997: *O Paleolítico Superior da Estremadura Portuguesa*. Edições Colibri. Faculdade de Letras de Lisboa. Lisboa. I
- 2000 a: "Nature and Culture in Portugal from 30.000 to 20.000 BP". In W. Roebroeks, M. Mussi, Svoboda, J. and K. Fennema (eds.): *Hunters of the Golden Age. The Mid-Upper Palaeolithic of Eurasia 30.000–20.000 BP*. University of Leiden. Leiden: 337-54.
- 2000 b: "The Ebro Frontier: A Model for the Late Extinction of Iberian Neanderthals". In C. B. Stringer, R. N. E. Barton and J. C. Finlayson (eds.): *Neanderthals on the Edge*. Papers from a Conference Marking the 150<sup>th</sup> Anniversary of the Forbes' Quarry Discovery, Gibraltar. Oxbow Books. Oxford: 111-21.
- ZILHÃO, J. and d'ERRICO, F. 1999: "The Chronology and Taphonomy of the Earliest Aurignacian and Its Implications for the Understanding of Neandertal Extinction". *Journal of World Prehistory* 13: 1-68.
- ZUBROW, E. 1989: "The Demographic Modelling of Neanderthal Extinction". In P. Mellars and C. Stringer (eds.): *The Human Revolution*. Princeton University Press. New Jersey: 121-31.