



Vol. 12 (2007), Paper no. 41, pages 1131–1150.

Journal URL

<http://www.math.washington.edu/~ejpecp/>

Large Deviations for the largest eigenvalue of rank one deformations of Gaussian ensembles

Mylène Maïda
Université Paris-Sud
Laboratoire de mathématiques
Bâtiment 425
Faculté des Sciences
91405 Orsay Cedex, France.
mylene.maida@math.u-psud.fr

Abstract

We establish a large deviation principle for the largest eigenvalue of a rank one deformation of a matrix from the GUE or GOE. As a corollary, we get another proof of the phenomenon, well-known in learning theory and finance, that the largest eigenvalue separates from the bulk when the perturbation is large enough.

A large part of the paper is devoted to an auxiliary result on the continuity of spherical integrals in the case when one of the matrix is of rank one, as studied in (12).

Key words: Large deviations, spiked models, spectral radius, Itzykson-Zuber integrals.

AMS 2000 Subject Classification: Primary 60F10, 15A52.

Submitted to EJP on September 26, 2006, final version accepted July 20, 2007.

1 Introduction

We consider in this paper rank one deformations of matrices from Gaussian ensembles, that is matrices which can be written $W_N + A_N$, with W_N from the Gaussian Orthogonal (or Unitary) Ensemble and A_N rank one deterministic, real symmetric if W_N is from the GOE and Hermitian if W_N is from the GUE.

Since the fifties, the classical Gaussian ensembles (see Mehta (19)) have been extensively studied. Various results for the global regime were established (Wigner semicircle law (22), large deviations for the spectral measure (5)...); the statistics of the spacings between eigenvalues were investigated for example in (8; 7), as well as the behaviour of extremal eigenvalues (Tracy-Widom distribution (21)). In the meantime, people got interested in the universality of some of these results. In this context, it is natural to look at various deformations of these ensembles, for example the rank one deformations we are interested in.

This so-called “deformed Wigner ensemble” was studied in (16) and (6), where the authors focused mainly on the problem of the local spacings and in (20) and (11), where they studied the behaviour of the largest eigenvalue. In this framework, our goal in this paper will be to establish a large deviation principle for the largest eigenvalue of $X_N = W_N + A_N$, that we denote in the sequel by x_N^* . Note that our result can also be seen as a generalization of the result established in (4) for the largest eigenvalue of a matrix distributed according to the GOE. If we denote by θ the unique non zero eigenvalue of A_N , the joint law of the eigenvalues x_1, \dots, x_N of $X_N = W_N + A_N$ is given by

$$\mathbb{Q}_N^\theta(dx_1, \dots, dx_N) = \frac{1}{Z_N^{\beta, \theta}} \prod_{i < j} |x_i - x_j|^\beta I_N^\beta(\theta, X_N) e^{-\frac{N}{2} \sum_{i=1}^N x_i^2} dx_1 \dots dx_N, \quad (1)$$

where I_N^β is the spherical integral defined by

$$I_N^\beta(\theta, X_N) := \int e^{N \text{tr}(U X_N U^* A_N)} dm_N^\beta(U) = \int e^{N \theta (U X_N U^*)_{11}} dm_N^\beta(U),$$

with m_N^β the Haar probability measure on \mathcal{O}_N the orthogonal group of size N if $\beta = 1$, on the unitary group \mathcal{U}_N if $\beta = 2$ and $Z_N^{\beta, \theta}$ is a normalizing constant. The fact that the joint law of the eigenvalues of X_N and that $I_N^\beta(\theta, X_N)$ depend on A_N only through its non zero eigenvalue θ comes from the unitary invariance respectively of the law of W_N and of the Haar measure m_N^β .

Our main result is the following

Theorem 1.1. *For $\beta = 1$ or 2 , if $\theta \geq 0$, then under \mathbb{Q}_N^θ , the largest eigenvalue $x_N^* = \max\{x_1, \dots, x_N\}$ satisfies a large deviation principle in the scale N , with good rate function K_θ^β defined as follows:*

- If $\theta \leq \sqrt{\frac{\beta}{2}}$,

$$K_\theta^\beta(x) = \begin{cases} +\infty, & \text{if } x < \sqrt{2\beta} \\ \int_{\sqrt{2\beta}}^x \sqrt{z^2 - 2\beta} dz, & \text{if } \sqrt{2\beta} \leq x \leq \theta + \frac{\beta}{2\theta}, \\ M_\theta^\beta(x), & \text{if } x \geq \theta + \frac{\beta}{2\theta}, \end{cases}$$

$$\text{with } M_\theta^\beta(x) = \frac{1}{2} \int_{\sqrt{2\beta}}^x \sqrt{z^2 - 2\beta} dz - \theta x + \frac{1}{4}x^2 + \frac{\beta}{4} - \frac{\beta}{4} \log \frac{\beta}{2} + \frac{1}{2}\theta^2.$$

- If $\theta \geq \sqrt{\frac{\beta}{2}}$,

$$K_\theta^\beta(x) = \begin{cases} +\infty, & \text{if } x < \sqrt{2\beta} \\ L_\theta^\beta(x), & \text{if } x \geq \sqrt{2\beta}, \end{cases}$$

$$\text{with } L_\theta^\beta(x) = \frac{1}{2} \int_{\theta + \frac{\beta}{2\theta}}^x \sqrt{z^2 - 2\beta} dz - \theta \left(x - \left(\theta + \frac{\beta}{2\theta} \right) \right) + \frac{1}{4} \left(x - \left(\theta + \frac{\beta}{2\theta} \right) \right)^2.$$

One can see in particular that K_θ^β differs from the rate function for the deviations of the largest eigenvalue of the non-deformed model that was obtained in (4).

Note that in the case when $\theta < 0$, similar results would hold for the smallest eigenvalue of the deformed ensemble. We let the precise statement to the reader and assume in the sequel that $\theta > 0$.

Remark 1.2. *Let us mention that, although we did not investigate this point in full details, very similar results can be obtained with our techniques in the case of sample covariance matrices for the so-called “single spike model” that is matrices of the form XX^* , where X is a $p \times n$ matrix, whose column vectors are iid Gaussian (real or complex) with a covariance matrix $\text{diag}(a, 1, 1, \dots, 1)$, with a single spike $a > 1$ (see below for references).*

We have to mention an important corollary of Theorem 1.1 :

Corollary 1.3. *For $\beta = 1$ or 2 , under \mathbb{Q}_N^θ , x_N^* converges almost surely to the edge of the support of the semicircle law σ_β as long as $\theta \leq \theta_c := \sqrt{\frac{\beta}{2}}$ and separates from the support when $\theta > \theta_c$. In this case, it converges to $\theta + \frac{\beta}{2\theta}$.*

This allows us to give a new proof, via large deviations, to this known phenomena which is crucial for applications to finance and learning theory (cf. for example (15; 18)).

On the mathematical level, this kind of phase transition has been pointed out and proved by several authors in the case of non-white sample covariance matrices (cf. for example (3) for the complete analysis in the complex Gaussian case, (10), (9) for more general models, (17) for statistical applications to PCA).

The organisation of the paper is as follows : as we can see in (1) above, the expression of the joint law \mathbb{Q}_N^θ of the eigenvalues involves the spherical integrals I_N^β in the case when one of the matrices is of rank one. We got the asymptotics of this quantity in (12) but we will need a precise continuity result of these spherical integrals to which Section 2 is devoted. In Section 3, we prove Theorem 1.1. Finally, in a very short Section 4, we show how to derive Corollary 1.3 from this Large Deviation Principle.

2 Continuity of spherical integrals

The question we want to address in this section is the continuity, in a topology to be prescribed, of $I_N^\beta(\theta, B_N)$, in its second argument B_N . The matrix B_N is supposed to be symmetric if $\beta = 1$ and Hermitian if $\beta = 2$. Due to invariance property of the Haar measure, we can always assume that B_N is real diagonal.

We denote by $\lambda_1(B_N), \dots, \lambda_N(B_N)$ the eigenvalues of B_N in decreasing order and we let $\hat{\nu}_{B_N} := \frac{1}{N} \sum_{i=2}^N \delta_{\lambda_i(B_N)}$; d is the Dudley distance defined on probability measures by

$$d(\mu, \nu) = \sup \left\{ \left| \int f d\mu - \int f d\nu \right|; |f(x)| \vee \left| \frac{f(x) - f(y)}{x - y} \right| \leq 1, \forall x \neq y \right\}.$$

The following continuity property holds

Proposition 2.1. *For $\beta = 1$ or 2 , for any $\theta > 0$ and any $\kappa > 0$, there exists a function $g_\kappa : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ going to zero at zero such that, for any $\delta > 0$ and N large enough, if B_N and B'_N are two sequences of real diagonal matrices such that $d(\hat{\nu}_{B_N}, \hat{\nu}_{B'_N}) \leq N^{-\kappa}$ and $|\lambda_1(B_N) - \lambda_1(B'_N)| \leq \delta$, with $\sup \|B_N\|_\infty < \infty$ then*

$$\left| \frac{1}{N} \log I_N^\beta(\theta, B_N) - \frac{1}{N} \log I_N^\beta(\theta, B'_N) \right| \leq g_\kappa(\delta).$$

Remark 2.2. *According to Theorem 6 of (12), we know that, for some values of θ , the limit of $\frac{1}{N} \log I_N^\beta(\theta, B_N)$ as N goes to infinity depends not only on the limiting spectral measure of B_N but also on the limit of $\lambda_1(B_N)$. Therefore $\frac{1}{N} \log I_N^\beta(\theta, B_N)$ cannot be continuous in the spectral measure of B_N but we have also to localize $\lambda_1(B_N)$. That is precisely the content of Proposition 2.1 above. We also refer the reader to the remarks made in (12) on point (3) of Lemma 14 therein.*

A key step to show Proposition 2.1 is to get an equivalent as explicit as possible of $\frac{1}{N} \log I_N^\beta(\theta, B_N)$. This is given by

Lemma 2.3. *If B_N has spectral radius uniformly bounded in N , then for any $\delta > 0$, for N large enough,*

$$\left| \frac{1}{N} \log I_N^\beta(\theta, B_N) - \left(\theta v_N - \frac{\beta}{2N} \sum_{i=1}^N \log \left(1 + \frac{2\theta}{\beta} v_N - \frac{2\theta}{\beta} \lambda_i(B_N) \right) \right) \right| \leq \delta,$$

where v_N is the unique solution in $\left[\lambda_N(B_N) - \frac{\beta}{2\theta}, \lambda_1(B_N) - \frac{\beta}{2\theta} \right]^c$ of the equation

$$\frac{\beta}{2\theta} \frac{1}{N} \sum_{i=1}^N \frac{1}{v_N + \frac{\beta}{2\theta} - \lambda_i(B_N)} = 1.$$

This lemma can be regarded as a generalization to any value of θ of the second point of Lemma 14 in (12).

The remaining of this section is devoted to its proof. For the sake of simplicity, we prove in full details the case $\beta = 1$ and leave to the reader the changes to the other cases.

2.1 Some preliminary inequalities

Notation and remarks.

- We denote by $\lambda_1 \geq \dots \geq \lambda_N$ the eigenvalues of B_N in decreasing order.
- For $\xi \in]0, 1/2[$, we define

$$K_N(\xi) := \{j \in \{1, \dots, N\} / \lambda_j \in (\lambda_1 - N^{-\xi}, \lambda_1]\},$$

and we denote by j_0 the cardinality of $K_N(\xi)$.

- The eigenvalues $\lambda_1 \geq \dots \geq \lambda_N$ being fixed, we can see that the function $x \mapsto \frac{1}{N} \sum_{i=1}^N \frac{1}{x - \lambda_i}$ is strictly decreasing taking negative values on $(-\infty, \lambda_N)$ and strictly decreasing taking positive values on (λ_1, ∞) . Therefore, v_N as introduced in Lemma 2.3, is well defined and we can define similarly \tilde{v}_N as the unique solution in $\left[\lambda_N(B_N) - \frac{1}{2\theta}, \lambda_1(B_N) - \frac{1}{2\theta}\right]^c$ of the equation

$$\frac{1}{N} \frac{1}{2\theta} \sum_{i=j_0+1}^N \frac{1}{\tilde{v}_N + \frac{1}{2\theta} - \lambda_i} = 1.$$

Moreover, if $\theta > 0$, one can easily see that $v_N + \frac{1}{2\theta}$ and $\tilde{v}_N + \frac{1}{2\theta}$ both lie in (λ_1, ∞) .

- \mathbb{E} and \mathbb{V} denotes respectively the expectation and the variance under the standard Gaussian measure on \mathbb{R}^N .
- As $1 + 2\theta v_N - 2\theta \lambda_1 > 0$, we can define the probability measure on \mathbb{R}^N given by

$$P_N(dg_1, \dots, dg_N) = (2\pi)^{-\frac{N}{2}} \prod_{i=1}^N \left[\sqrt{1 + 2\theta v_N - 2\theta \lambda_i} e^{-\frac{1}{2}(1+2\theta v_N - 2\theta \lambda_i)g_i^2} dg_i \right].$$

We denote by \mathbb{E}_{P_N} and \mathbb{V}_{P_N} respectively the expectation and the variance under P_N .

- Similarly, we define the probability measure on \mathbb{R}^{N-j_0} given by

$$\tilde{P}_N(dg_{j_0+1}, \dots, dg_N) = (2\pi)^{-\frac{N-j_0}{2}} \prod_{i=j_0+1}^N \left[\sqrt{1 + 2\theta \tilde{v}_N - 2\theta \lambda_i} e^{-\frac{1}{2}(1+2\theta \tilde{v}_N - 2\theta \lambda_i)g_i^2} dg_i \right].$$

We denote by $\mathbb{E}_{\tilde{P}_N}$ and $\mathbb{V}_{\tilde{P}_N}$ respectively the expectation and the variance under \tilde{P}_N .

Before going to the proof of Lemma 2.3, we enumerate hereafter some inequalities on the quantities we have just introduced, that will be useful further.

Fact 2.4. *Let $\theta > 0$. We have the following inequalities :*

1. For $i \geq j_0 + 1$, $v_N + \frac{1}{2\theta} - \lambda_i \geq N^{-\xi}$ and $\tilde{v}_N + \frac{1}{2\theta} - \lambda_i \geq N^{-\xi}$.
2. $v_N + \frac{1}{2\theta} - \lambda_1 \geq \frac{j_0}{2\theta N} - N^{-\xi}$ and $\forall i$, $v_N + \frac{1}{2\theta} - \lambda_i \geq \frac{1}{2\theta N}$.
3. $\tilde{v}_N \leq v_N \leq \lambda_1 \leq \tilde{v}_N + \frac{1}{2\theta} \leq v_N + \frac{1}{2\theta}$.
4. For any $\delta > 0$, for any $\xi \in (0, 1/2)$, for any $\varepsilon > 0$, there exists N_0 such that for any $N \geq N_0$ such that $j_0 \leq \delta N^{1-\frac{\xi}{2}}$, $|v_N - \tilde{v}_N| \leq \varepsilon$.

Proof :

1. As mentioned in the remarks above, $v_N + \frac{1}{2\theta} \geq \lambda_1$. For $i \geq j_0 + 1$, $\lambda_i \leq \lambda_1 - N^{-\xi}$ so that $v_N + \frac{1}{2\theta} \geq \lambda_1 \geq \lambda_i + N^{-\xi}$. The same holds for \tilde{v}_N .
2. We have the following inequality

$$j_0 \frac{1}{v_N + \frac{1}{2\theta} - (\lambda_1 - N^{-\xi})} \leq \sum_{i=1}^{j_0} \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} \leq 2\theta N,$$

where the right inequality comes from the fact that $\sum_{i=1}^N \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} = 2\theta N$ and for all i , $v_N + \frac{1}{2\theta} - \lambda_i \geq 0$ and the left one is inherited from the definition of j_0 . Putting the leftmost and rightmost terms together, we get the first inequality announced in point (2) above. The second one is even simpler : we have that $\sum_{i=1}^N \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} = 2\theta N$ and each term is positive so that any of them is smaller than $2\theta N$.

3. Suppose that $v_N < \tilde{v}_N$, then for all $i \geq j_0 + 1$, $v_N + \frac{1}{2\theta} - \lambda_i < \tilde{v}_N + \frac{1}{2\theta} - \lambda_i$

$$2\theta N = \sum_{i=j_0+1}^N \frac{1}{\tilde{v}_N + \frac{1}{2\theta} - \lambda_i} < \sum_{i=j_0+1}^N \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i},$$

but the rightmost term is smaller or equal to $2\theta N$. Therefore, $\tilde{v}_N \leq v_N$.

The λ_i 's being in decreasing order, we get that $2\theta = \frac{1}{N} \sum_{i=1}^N \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} \leq \frac{1}{v_N + \frac{1}{2\theta} - \lambda_1}$,

this gives $v_N + \frac{1}{2\theta} \leq \lambda_1 + \frac{1}{2\theta}$, i.e. $v_N \leq \lambda_1$.

4. Let $\varepsilon > 0$, $\delta > 0$ and $\xi \in (0, 1/2)$ be fixed. From (3), we have that

$$0 \leq v_N - \tilde{v}_N \leq v_N + \frac{1}{2\theta} - \lambda_1$$

If $v_N + \frac{1}{2\theta} - \lambda_1 \leq \varepsilon$, then $|v_N - \tilde{v}_N| \leq \varepsilon$.

Otherwise, we can write, thanks to (3), that

$$\sum_{i=j_0+1}^N \frac{1}{\tilde{v}_N + \frac{1}{2\theta} - \lambda_i} - \sum_{i=j_0+1}^N \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} \geq (v_N - \tilde{v}_N) \sum_{i=j_0+1}^N \frac{1}{(v_N + \frac{1}{2\theta} - \lambda_i)^2}.$$

From Cauchy-Schwarz inequality, we get

$$\sum_{i=1}^{j_0} \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} \geq (v_N - \tilde{v}_N) \frac{1}{N - j_0} \left(\sum_{i=j_0+1}^N \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} \right)^2.$$

Therefore,

$$|v_N - \tilde{v}_N| \leq \frac{(N - j_0)y_N}{(2\theta N - y_N)^2} \quad \text{with} \quad y_N = \sum_{i=1}^{j_0} \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i}.$$

Now, we are in the case when $v_N + \frac{1}{2\theta} - \lambda_1 \geq \varepsilon$, so that $y_N \leq \frac{j_0}{\varepsilon}$. For N such that $j_0 \leq \delta N^{1-\frac{\xi}{2}}$,

$$|v_N - \tilde{v}_N| \leq \frac{(N - j_0)\delta N^{1-\frac{\xi}{2}}}{\varepsilon(2\theta N - \delta N^{1-\frac{\xi}{2}})^2} \leq \varepsilon,$$

where the last inequality holds for N large enough. \square

2.2 Proof of Lemma 2.3

We first prove **the upper bound**: the starting point will be the same as in (12). It is a well known fact that the first column vector of a random orthogonal matrix distributed according to the Haar measure on \mathcal{O}_N has the same law as a standard Gaussian vector in \mathbb{R}^N divided by its Euclidian norm. Therefore, we can write

$$I_N(\theta, B_N) = \mathbb{E} \left(\exp \left\{ N\theta \frac{\sum_{i=1}^N \lambda_i g_i^2}{\sum_{i=1}^N g_i^2} \right\} \right).$$

From concentration for the norm of a Gaussian vector (cf. (12) for details), we get that, for any κ such that $0 < \kappa < 1/2$,

$$1 \leq \frac{I_N(\theta, B_N)}{\mathbb{E} \left(\mathbf{1}_{\mathcal{A}_N(\kappa)} \exp \left\{ N\theta \frac{\sum_{i=1}^N \lambda_i g_i^2}{\sum_{i=1}^N g_i^2} \right\} \right)} \leq \delta(\kappa, N), \quad (2)$$

where $\mathcal{A}_N(\kappa) = \left\{ \left| \frac{\|g\|^2}{N} - 1 \right| \leq N^{-\kappa} \right\}$ and $\delta(\kappa, N)$ goes to one at infinity for any $0 < \kappa < 1/2$.

From there, we have

$$\begin{aligned} I_N(\theta, B_N) &\leq \delta(\kappa, N) e^{N\theta v_N + N^{1-\kappa}\theta(M+v_N)} \mathbb{E} \left[\mathbf{1}_{\mathcal{A}_N(\kappa)} \exp \left\{ \theta \sum_{i=1}^N \lambda_i g_i^2 - \theta v_N \sum_{i=1}^N g_i^2 \right\} \right] \\ &\leq \delta(\kappa, N) e^{N\theta v_N + N^{1-\kappa}\theta(M+v_N)} \prod_{i=1}^N \left[\sqrt{1 + 2\theta v_N - 2\theta \lambda_i} \right]^{-1}, \end{aligned}$$

where M is the uniform bound on the spectral radius of B_N and we use that $P_N(\mathcal{A}_N(\kappa)) \leq 1$. Therefore, for any $\delta > 0$, we get that for N large enough,

$$\frac{1}{N} \log I_N(\theta, B_N) \leq \theta v_N - \frac{1}{2} \sum_{i=1}^N \log(1 + 2\theta v_N - 2\theta \lambda_i) + \delta.$$

For the **proof of the lower bound**, we have to treat two distinct cases. In both cases, the starting point, inherited from (2), is the following:

$$I_N(\theta, B_N) \geq \delta(\kappa, N) e^{N\theta v_N - N^{1-\kappa}\theta(M+v_N)} \mathbb{E} \left[\mathbf{1}_{\mathcal{A}_N(\kappa)} \exp \left\{ \theta \sum_{i=1}^N \lambda_i g_i^2 - \theta v_N \sum_{i=1}^N g_i^2 \right\} \right],$$

but our strategy will be different according to whether there is a lot of eigenvalues at the vicinity of the largest eigenvalue λ_1 or not, that is according to the size of j_0 defined above.

For $\kappa \in]0, 1/2[$ fixed, we choose in the sequel ξ such that $0 < \xi \leq \frac{1}{2} - \kappa$.

• **First case :** N is such that $j_0 \geq \delta N^{1-\frac{\xi}{2}}$.

In this case, the situation is very similar to what happens with a small θ , we therefore follow the proof of (12). Indeed, we write

$$\mathbb{E} \left[\mathbf{1}_{\mathcal{A}_N(\kappa)} \exp \left\{ \theta \sum_{i=1}^N \lambda_i g_i^2 - \theta v_N \sum_{i=1}^N g_i^2 \right\} \right] = \prod_{i=1}^N \left[\sqrt{1 + 2\theta v_N - 2\theta \lambda_i} \right]^{-1} P_N(\mathcal{A}_N(\kappa)),$$

and show that, for N large enough, $P_N(\mathcal{A}_N(\kappa)) \geq 1/2$.

An easy computation gives that

$$\mathbb{V}_{P_N} \left[\frac{1}{N} \|g\|^2 \right] = \frac{2}{N^2} \sum_{i=1}^N \frac{1}{(1 + 2\theta v_N - 2\theta \lambda_i)^2}$$

and our goal is to show that this variance decreases fast enough.

From (2) in Fact 2.4, we have

$$v_N + \frac{1}{2\theta} - \lambda_i \geq v_N + \frac{1}{2\theta} - \lambda_1 \geq \frac{j_0}{2\theta N} - N^{-\xi} \geq \frac{\delta}{2\theta} N^{-\frac{\xi}{2}} - N^{-\xi} \geq \frac{\delta}{2\theta} N^{-\xi}.$$

This gives that

$$\mathbb{V}_{P_N} \left[\frac{1}{N} \|g\|^2 \right] \leq \frac{2}{N^2 \delta^2} N^{2\xi} N \leq \frac{2}{\delta^2} N^{2\xi-1}$$

Therefore, by Chebichev inequality,

$$P_N(\mathcal{A}_N(\kappa)^c) \leq \frac{2}{\delta^2} N^{2\xi+2\xi-1} \leq \frac{1}{2},$$

where the last inequality holds for N large enough with our choice of ξ .

• **Second case** : N is such that $j_0 \leq \delta N^{1-\frac{\xi}{2}}$.

The strategy will be a bit different : separating the eigenvalues of B_N that are in $K_N(\xi)$ and the others, we get

$$\begin{aligned} I_N(\theta, B_N) &\geq e^{N\theta\tilde{v}_N - N^{1-\kappa}\theta(M+\tilde{v}_N)} \mathbb{E} \left(\mathbf{1}_{\left| \frac{1}{N} \|\mathbf{g}\|^2 - 1 \right| \leq N^{-\kappa}} \exp \left\{ \theta \sum_{i=1}^N \lambda_i g_i^2 - \theta \tilde{v}_N \sum_{i=1}^N g_i^2 \right\} \right) \\ &\geq e^{N\theta\tilde{v}_N - N^{1-\kappa}\theta(M+\tilde{v}_N)} \mathbb{E} \left(\mathbf{1}_{\left| \frac{1}{N} \sum_{i=j_0+1}^N g_i^2 - 1 \right| \leq \frac{N-\kappa}{2}} \exp \left\{ \theta \sum_{i=j_0+1}^N \lambda_i g_i^2 - \theta \tilde{v}_N \sum_{i=j_0+1}^N g_i^2 \right\} \right) \\ &\quad \mathbb{E} \left(\mathbf{1}_{\left| \frac{1}{N} \sum_{i=1}^{j_0} g_i^2 \right| \leq \frac{N-\kappa}{2}} \exp \left\{ \theta \sum_{i=1}^{j_0} \lambda_i g_i^2 - \theta \tilde{v}_N \sum_{i=1}^{j_0} g_i^2 \right\} \right). \end{aligned}$$

The first term will be treated similarly to what we made in the first case. We can easily check that $\mathbb{E}_{\tilde{P}_N} \left(\frac{1}{N} \sum_{i=j_0+1}^N g_i^2 \right) = 1$ and $\text{Var}_{\tilde{P}_N} \left(\frac{1}{N} \sum_{i=j_0+1}^N g_i^2 \right) = \frac{2}{N^2} \sum_{i=j_0+1}^N \frac{1}{(1+2\theta\tilde{v}_N - 2\theta\lambda_i)^2}$.

From (1) in Fact 2.4, we get that $\mathbb{V}_{\tilde{P}_N} \left(\frac{1}{N} \sum_{i=j_0+1}^N g_i^2 \right) \leq \frac{1}{2\theta^2 N^2} N^{2\xi} N \leq \frac{1}{2\theta^2} N^{2\xi-1}$. Therefore

$$\tilde{P}_N \left(\left| \frac{1}{N} \sum_{i=j_0+1}^N g_i^2 - 1 \right| \geq \frac{N^{-\kappa}}{2} \right) \leq \frac{2}{\theta^2} N^{2\kappa+2\xi-1},$$

which goes to zero with our choice of ξ .

This gives that for N large enough,

$$\mathbb{E} \left(\mathbf{1}_{\left| \frac{1}{N} \sum_{i=j_0+1}^N g_i^2 - 1 \right| \leq \frac{N-\kappa}{2}} \exp \left\{ \theta \sum_{i=j_0+1}^N \lambda_i g_i^2 - \theta \tilde{v}_N \sum_{i=j_0+1}^N g_i^2 \right\} \right) \geq \frac{1}{2} \prod_{i=j_0+1}^N \frac{1}{\sqrt{1+2\theta\tilde{v}_N - 2\theta\lambda_i}}. \quad (3)$$

We now go to the last term. From (3) in Fact 2.4, we have that $\tilde{v}_N \leq \lambda_1$, so that for any $i \leq j_0$, $\lambda_i - \tilde{v}_N \geq -N^{-\xi}$. Therefore,

$$\exp \left\{ \theta \left[\sum_{i=1}^{j_0} (\lambda_i - \tilde{v}_N) g_i^2 \right] \right\} \geq \exp \left(-\theta N^{-\xi} j_0 \right) \geq \exp \left(-\theta \delta N^{1-\frac{3\xi}{2}} \right),$$

so that

$$\begin{aligned} \mathbb{E} \left(\mathbf{1}_{\left| \frac{1}{N} \sum_{i=1}^{j_0} g_i^2 \right| \leq \frac{N-\kappa}{2}} \exp \left\{ \theta \sum_{i=1}^{j_0} \lambda_i g_i^2 - \theta \tilde{v}_N \sum_{i=1}^{j_0} g_i^2 \right\} \right) &\geq \mathbb{P} \left(\left| \frac{1}{N} \sum_{i=1}^{j_0} g_i^2 \right| \leq \frac{N-\kappa}{2} \right) \cdot \exp \left(-\theta \delta N^{1-\frac{3\xi}{2}} \right) \\ &\geq \frac{1}{2} \cdot \exp \left(-\theta \delta N^{1-\frac{3\xi}{2}} \right), \end{aligned} \quad (4)$$

where the last inequality is again obtained through Chebichev inequality.

Putting together (3) and (4), we get that for N large enough

$$\theta\tilde{v}_N - \frac{1}{2N} \sum_{i=j_0+1}^N \log(1 + 2\theta\tilde{v}_N - 2\theta\lambda_i) - \frac{\delta}{4} \leq \frac{1}{N} \log I_N(\theta, B_N).$$

The last step is now to prove that, for N large enough,

$$\left| \left(\theta v_N - \frac{1}{2N} \sum_{i=1}^N \log(1 + 2\theta v_N - 2\theta\lambda_i) \right) - \left(\theta\tilde{v}_N - \frac{1}{2N} \sum_{i=j_0+1}^N \log(1 + 2\theta\tilde{v}_N - 2\theta\lambda_i) \right) \right| \leq \frac{\delta}{2}. \quad (5)$$

On one side, we have that

$$\begin{aligned} & \left| \left(\theta v_N - \frac{1}{2N} \sum_{i=j_0+1}^N \log(1 + 2\theta v_N - 2\theta\lambda_i) \right) - \left(\theta\tilde{v}_N - \frac{1}{2N} \sum_{i=j_0+1}^N \log(1 + 2\theta\tilde{v}_N - 2\theta\lambda_i) \right) \right| \\ & \leq |v_N - \tilde{v}_N| \left(\theta + \frac{1}{2N} \sum_{i=j_0+1}^N \frac{1}{\tilde{v}_N + \frac{1}{2\theta} - \lambda_i} \right) \leq \frac{\delta}{4}, \end{aligned}$$

where we use (4) in Fact 2.4, with N large enough and $\varepsilon = \frac{\delta}{2\theta}$.

From (1) in Fact 2.4, we have that $\frac{1}{N} \leq 1 + 2\theta v_N - 2\theta\lambda_i$ and from (2), $v_N \leq \lambda_1 \leq M$, where M is the uniform bound on the spectral radius of B_N , so that $1 + 2\theta v_N - 2\theta\lambda_i \leq 1 + 2\theta M$. Therefore, for N large enough,

$$\begin{aligned} \left| \frac{1}{2N} \sum_{i=1}^{j_0} \log(1 + 2\theta v_N - 2\theta\lambda_i) \right| & \leq \frac{1}{2N} j_0 \log N \\ & \leq \frac{1}{2N} \delta N^{1-\frac{\varepsilon}{2}} \log N \leq \frac{\delta}{4}. \end{aligned}$$

This concludes the proof of Lemma 2.3. \square

2.3 Proof of Proposition 2.1

Let $\kappa > 0$ be fixed and $(B_N)_{N \in \mathbb{N}}$ and $(B'_N)_{N \in \mathbb{N}}$ two sequences of matrices. We denote by $\lambda_1 \geq \dots \geq \lambda_N$ and $\lambda'_1 \geq \dots \geq \lambda'_N$ the eigenvalues of B_N and B'_N respectively, both in decreasing order.

We assume that $d(\hat{\nu}_{B_N}, \hat{\nu}_{B'_N}) \leq N^{-\kappa}$ and $|\lambda_1 - \lambda'_1| \leq \delta$ for N large enough and that there exists M such that $\sup_N \|B_N\|_\infty < M$ and $\sup_N \|B'_N\|_\infty < M$.

We introduce also the following notations: $H_{B_N}(z) = \frac{1}{N} \sum_{i=1}^N \frac{1}{z - \lambda_i}$ and $H_{B'_N}(z) = \frac{1}{N} \sum_{i=1}^N \frac{1}{z - \lambda'_i}$.

- **First case :** N and θ are such that $2\theta \in H_{B_N}((\lambda_1 + 2\delta, +\infty)) \cap H_{B'_N}((\lambda'_1 + 2\delta, +\infty))$.

In this frame work, continuity has been established in Lemma 14 of (12). It comes from Lemma 2.3 and the fact that, for any $\lambda \in \cup_{N_0 \geq 0} \cap_{N \geq N_0} (\text{supp } \hat{\nu}_{B_N} \cap \text{supp } \hat{\nu}_{B'_N})$, $z \mapsto (z - \lambda)^{-1}$ is continuous bounded (with a norm independent of δ) on $\cup_{N_0 \geq 0} \cap_{N \geq N_0} ((\lambda_1 + 2\delta, +\infty) \cap (\lambda'_1 + 2\delta, +\infty))$.

- **Second case :** N and θ are such that $2\theta \notin H_{B_N}((\lambda_1 + 2\delta, +\infty))$ and $2\theta \notin H_{B'_N}((\lambda'_1 + 2\delta, +\infty))$.

In this case, $2\theta \notin H_{B_N}((\lambda_1 + 2\delta, +\infty)) \Rightarrow v_N + \frac{1}{2\theta} \in (\lambda_1, \lambda_1 + 2\delta)$ and similarly $v'_N + \frac{1}{2\theta} \in (\lambda'_1, \lambda'_1 + 2\delta)$ so that

$$|v_N - v'_N| \leq 3\delta.$$

Thanks to Lemma 2.3, we know that it is enough to study

$$\Delta_N := \left| \frac{1}{N} \sum_{i=1}^N \log \left(v_N + \frac{1}{2\theta} - \lambda_i \right) - \frac{1}{N} \sum_{i=1}^N \log \left(v'_N + \frac{1}{2\theta} - \lambda'_i \right) \right|.$$

As $d(\hat{\nu}_{B_N}, \hat{\nu}_{B'_N}) \leq N^{-\kappa}$, we proceed as in the proof of Lemma 5.1 in (14) and define a permutation σ_N that allows to put in pairs all but $(N^{1-\kappa} \wedge N\delta)$ of the λ_i 's with a corresponding $\lambda'_{\sigma_N(i)}$ which lies at a distance less than δ from λ_i .

As in (14), we denote by \mathcal{J}_0 the set of indices i such that we have such a pairing. Then we have

$$\begin{aligned} \Delta_N &\leq \frac{1}{N} \sum_{i \in \mathcal{J}_0} \max \left(\frac{1}{v_N + \frac{1}{2\theta} - \lambda_i}, \frac{1}{v'_N + \frac{1}{2\theta} - \lambda'_{\sigma_N(i)}} \right) (|v_N - v'_N| + |\lambda_i - \lambda'_{\sigma_N(i)}|) \\ &\quad + \frac{1}{N} \sum_{i \in \mathcal{J}_0^c} \left| \log \left(v_N + \frac{1}{2\theta} - \lambda_i \right) - \log \left(v'_N + \frac{1}{2\theta} - \lambda'_i \right) \right| \\ &\leq \left(\frac{1}{N} \sum_{i=1}^N \frac{1}{v_N + \frac{1}{2\theta} - \lambda_i} + \frac{1}{N} \sum_{i=1}^N \frac{1}{v'_N + \frac{1}{2\theta} - \lambda'_{\sigma_N(i)}} \right) 4\delta \\ &\quad + \frac{1}{N} \sum_{i \in \mathcal{J}_0^c} \left| \log \left(v_N + \frac{1}{2\theta} - \lambda_i \right) - \log \left(v'_N + \frac{1}{2\theta} - \lambda'_i \right) \right| \\ &\leq 16\theta\delta + \frac{2}{N} [N^{1-\kappa} \wedge N\delta] \left[|\log(2N\theta)| \vee \log \left(2M + \frac{1}{2\theta} \right) \right], \end{aligned}$$

where we used once again that

$$\frac{1}{2N\theta} \leq v_N + \frac{1}{2\theta} - \lambda_i \leq 2M + \frac{1}{2\theta}$$

so that we get the required continuity in this second case.

- **Third case :** N and θ are such that $2\theta \in H_{B_N}((\lambda_1 + 2\delta, +\infty))$ and $2\theta \notin H_{B'_N}((\lambda'_1 + 2\delta, +\infty))$.

In this case, we proceed exactly as in the second case. The only point is that establishing that v_N cannot be far from v'_N will be a bit more involved. We address this point in detail.

On one side we have from Fact 2.4 that

$$\lambda'_1 \leq v'_N + \frac{1}{2\theta} \leq \lambda'_1 + 2\delta. \tag{6}$$

On the other side, as $|\lambda_1 - \lambda'_1| \leq \delta$, $\lambda'_1 + 2\delta$ is greater than λ_1 and the map $B_N \mapsto H_{B_N}$ is continuous outside the support of all the spectral measures so that

$$\left| H_{B'_N}(\lambda'_1 + 2\delta) - H_{B_N}(\lambda'_1 + 2\delta) \right| \leq C(\delta),$$

with the function C going to zero at zero.

Furthermore, $H_{B'_N}$ is decreasing on $(\lambda'_1, +\infty)$ so that $H_{B'_N}(\lambda'_1 + 2\delta) < 2\theta$, yielding

$$H_{B_N}(\lambda'_1 + 2\delta) \leq 2\theta + C(\delta),$$

and H_{B_N} being decreasing

$$H_{B_N}(\lambda_1 + 3\delta) \leq 2\theta + C(\delta) = H_{B_N}\left(v_N + \frac{1}{2\theta}\right) + C(\delta),$$

what implies

$$\lambda_1 \leq v_N + \frac{1}{2\theta} \leq \lambda_1 + 3\delta + K(\delta),$$

with the function K going to zero at zero. and, together with (6) this gives that

$$|v_N - v'_N| \leq 5\delta + K(\delta).$$

Now the same estimates as in the second case above lead to the same conclusion. This gives Proposition 2.1. \square

3 Large deviations for x_N^*

The goal of this section is to prove the large deviation principle for x_N^* , the largest eigenvalue of a matrix from the deformed Gaussian ensemble, announced in the introduction in Theorem 1.1.

A first step will be to prove the following

Proposition 3.1. *For $\beta = 1$ or 2 and $\theta > 0$, if we define*

$$\mathbb{P}_N^\theta(dx_1, \dots, dx_N) = \frac{1}{Z_N^\beta} \prod_{i < j} |x_i - x_j|^\beta I_N^\beta(\theta, X_N) e^{-\frac{N}{2} \sum_{i=1}^N x_i^2} dx_1 \dots dx_N, \tag{7}$$

with Z_N^β the normalizing constant in the case $\theta = 0$, and we let

$$F_\theta^\beta(x) := \begin{cases} +\infty, & \text{if } x < \sqrt{2\beta} \\ -\frac{\beta}{2} + \frac{\beta}{2} \log \frac{\beta}{2} - \Phi_\beta(x, \sigma_\beta) - I_{\sigma_\beta}^\beta(x, \theta), & \text{otherwise,} \end{cases}$$

where σ_β denotes the semicircle law whose density on \mathbb{R} is given by $\frac{1}{\beta\pi} \mathbf{1}_{[-\sqrt{2\beta}, \sqrt{2\beta}]} \sqrt{2\beta - t^2} dt$, for $\mu \in \mathcal{P}(\mathbb{R})$ and $x \in \mathbb{R}$,

$$\Phi_\beta(x, \mu) = \beta \int \log |x - y| d\mu(y) - \frac{1}{2}x^2,$$

and $I_\mu^\beta(x, \theta) = \lim_{N \rightarrow \infty} \frac{1}{N} \log I_N^\beta(\theta, B_N)$, where B_N has limiting spectral measure μ and limiting largest eigenvalue x , then we have the following large deviations bounds :

1. there exists a function $f_\theta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ going to infinity at infinity such that for all N

$$\mathbb{P}_N^\theta \left(\max_{i=1 \dots N} |x_i| \geq M \right) \leq e^{-N f_\theta(M)}.$$

2. For any x , for any M such that $|x| < M$,

$$\lim_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_N^\theta(x \leq x_N^* \leq x + \delta, \max_{1 \dots N} |x_i| \leq M) \leq -F_\theta^\beta(x)$$

3. For any x ,

$$\lim_{\delta \downarrow 0} \liminf_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_N^\theta(x \leq x_N^* \leq x + \delta) \geq -F_\theta^\beta(x)$$

Remark 3.2. The function $I_\mu^\beta(x, \theta) = \lim_{N \rightarrow \infty} \frac{1}{N} \log I_N^\beta(\theta, B_N)$ is well defined by virtue of Theorem 6 in (12) (an explicit expression for it will be given in Section 3.2 below).

3.1 Proof of Proposition 3.1

- We first prove the “exponential tightness” property (1).

It is more convenient to rewrite (7) as

$$\mathbb{P}_N^\theta(dx_1, \dots, dx_N) = \frac{e^{\frac{N}{2}\theta^2}}{Z_N^\beta} \prod_{i < j} |x_i - x_j|^\beta e^{-\frac{N}{2} \text{tr}(X_N - A_N)^2} dx_1 \dots dx_N.$$

Now, a well known inequality (see for example Lemma 2.3 in (2)) gives that

$$\text{tr}(X_N - A_N)^2 \geq \min_{\pi} \sum_{i=1}^N |x_k - a_{\pi(k)}|^2,$$

where the minimum is taken over all permutations π of $\{1, \dots, N\}$. But all a_k 's are zero, except one of them, let's say a_1 , which is equal to θ . As the law of the x_j 's is invariant by permutations,

we can assume that $\pi_*^{-1}(1) = 1$, where π_* is the permutation for which the minimum is reached. Therefore

$$\operatorname{tr}(X_N - A_N)^2 \geq (x_1 - \theta)^2 + \sum_{i=2}^N x_i^2.$$

We can now use the very same estimates as in Lemma 6.3 in (4) to get (1). More precisely, we can write

$$|(x - \theta) + \theta - x_j|^\beta e^{-\frac{x_j^2}{2}} \leq e^{\frac{(x-\theta)^2}{4}},$$

for x large enough, so that, for M large enough,

$$\mathbb{P}_N^\theta \left(\max_{i=1 \dots N} |x_i| \geq M \right) \leq N \mathbb{P}_N^\theta (|x_1| \geq M) \leq \frac{Z_{N-1}^\beta}{Z_N^\beta} e^{-\frac{1}{4}N(M-\theta)^2 + \frac{N}{2}\theta^2}.$$

From Selberg formula (cf for example proof of Proposition 3.1 in (5)), we can show that

$$\frac{1}{N} \log \frac{Z_{N-1}^\beta}{Z_N^\beta} \xrightarrow{N \rightarrow \infty} C. \text{ This concludes the proof of (1).}$$

- For all $x < \sqrt{2\beta}$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_N^\theta(x_N^* \leq x) = -\infty. \tag{8}$$

Indeed, we know from Theorem 1.1 in (5) that the spectral measure of W_N satisfies a large deviation principle in the scale N^2 with a good rate function whose unique minimizer is the semicircle law σ_β . We can check that adding a deterministic matrix of bounded rank (uniformly in N) does not affect the spectral measure in this scale so that the spectral measure of X_N satisfies the same large deviation principle.

Therefore, if we let $x < \sqrt{2\beta}$, $f \in \mathcal{C}_b(\mathbb{R})$ such that $f(y) = 0$ if $y \leq x$ but $\int f d\sigma_\beta > 0$ and if we consider the closed set $F := \{\mu / \int f d\mu = 0\}$, we have that

$$\mathbb{Q}_N^\theta(x_N^* \leq x) \leq \mathbb{Q}_N^\theta \left(\frac{1}{N} \sum_{i=1}^N f(x_i) = 0 \right) \leq \mathbb{Q}_N^\theta(\hat{\mu}_N \in F),$$

where $\hat{\mu}_N := \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$ is the spectral measure of X_N . As $\sigma_\beta \notin F$,

$$\limsup_{N \rightarrow \infty} \frac{1}{N^2} \log \mathbb{Q}_N^\theta(x_N^* \leq x) < 0.$$

Furthermore, as we saw above, $\frac{1}{N} \log \frac{Z_{N-1}^\beta}{Z_N^\beta} \xrightarrow{N \rightarrow \infty} C$, the same holds for \mathbb{P}_N^θ and we immediately deduce what gives immediately (8).

- Let now $x \geq \sqrt{2\beta}$ and $\delta > 0$.

Let $M > |x|$ and δ small enough so that $M \geq |x + \delta|$. One important remark is that, by invariance by permutation, we have,

$$\mathbb{P}_N^\theta(x \leq x_N^* \leq x + \delta, \max_{i=1 \dots N} |x_i| \leq M) \leq N \mathbb{P}_N^\theta(x \leq x_1 \leq x + \delta, x_1 \geq \max_{i=2 \dots N} x_i, \max_{i=1 \dots N} |x_i| \leq M).$$

We introduce now the following notations :

- $\hat{\pi}_N := \frac{1}{N-1} \sum_{i=2}^N \delta_{x_i}$,
- \mathbb{P}_N^{N-1} is the measure on \mathbb{R}^{N-1} such that, for each Borel set E , we have

$$\mathbb{P}_N^{N-1}(\lambda \in E) = \mathbb{P}_{N-1}^0 \left(\sqrt{1 - \frac{1}{N}} \lambda \in E \right).$$

With these notations, we have

$$\begin{aligned} B &:= \mathbb{P}_N^\theta(x \leq x_N^* \leq x + \delta, \max_{i=1 \dots N} |x_i| \leq M) \\ &\leq \int_x^{x+\delta} dx_1 \int_{[-M, M]^{N-1}} e^{(N-1)\Phi_\beta(x_1, \hat{\pi}_N)} \cdot C_N^\beta \cdot I_N^\beta(\theta, X_N) \cdot d\mathbb{P}_{N-1}^0(x_2, \dots, x_N), \end{aligned}$$

where $C_N^\beta := N \frac{Z_{N-1}^\beta}{Z_N^\beta} \cdot \left(1 - \frac{1}{N}\right)^{\beta \frac{N(N-1)}{4}}$.

Let $0 < \kappa < \frac{1}{4}$, we have

$$\begin{aligned} B &\leq C_N^\beta \cdot \int_x^{x+\delta} dx_1 \int_{\substack{\hat{\pi}_N \in B(\sigma_\beta, N^{-\kappa}), \\ \max_{i=2 \dots N} x_i \leq x_1}} e^{(N-1)\Phi_\beta(x_1, \hat{\pi}_N)} I_N^\beta(\theta, X_N) \cdot d\mathbb{P}_N^{N-1}(x_2, \dots, x_N) \\ &\quad + (2M)^N e^{NM\theta} C_N^\beta \mathbb{P}_N^{N-1}(\hat{\pi}_N \notin B(\sigma_\beta, N^{-\kappa})), \quad (9) \end{aligned}$$

where $B(\sigma_\beta, N^{-\kappa})$ is the ball of size $N^{-\kappa}$ centered at σ_β , for the Dudley distance (defined at the very beginning of Section 2).

We first show that the second term is exponentially negligible. We have

$$\mathbb{P}_N^{N-1}(\hat{\pi}_N \notin B(\sigma_\beta, N^{-\kappa})) \leq \mathbb{P}_N^{N-1}(\|F_{N-1} - F_\beta\|_\infty \geq N^{-\kappa}),$$

where F_{N-1} and F_β are respectively the (cumulative) distribution function of $\hat{\pi}_N$ and σ_β . We know from the result of Bai in (1) that

$$\|\mathbb{E}_N^{N-1} F_{N-1} - F_\beta\| = O(N^{-\frac{1}{4}}),$$

where \mathbb{E}_N^{N-1} is the expectation under \mathbb{P}_N^{N-1} , so that

$$\mathbb{P}_N^{N-1}(\|F_{N-1} - F_\beta\|_\infty \geq N^{-\kappa}) \leq \mathbb{P}_N^{N-1} \left(\|F_{N-1} - \mathbb{E}_N^{N-1} F_{N-1}\|_\infty \geq \frac{N^{-\kappa}}{2} \right).$$

But, by a result of concentration of (13) (see Theorem 1.1), we have that there exists a constant $C > 0$ such that for all $N \in \mathbb{N}$,

$$\mathbb{P}_N^{N-1}(\|F_{N-1} - \mathbb{E}_N^{N-1} F_{N-1}\|_\infty \geq N^{-\kappa}) \leq e^{-CN^{2-2\kappa}},$$

so that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_N^{N-1}(\hat{\pi}_N \notin B(\sigma_\beta, N^{-\kappa})) = -\infty.$$

We can now come back to the first term in (9). The same computation as in the proof of Proposition 3.1 in (5), based on Selberg formula, gives that

$$\frac{1}{N} \log C_N^\beta \xrightarrow{N \rightarrow \infty} -\frac{\beta}{2} \log \frac{\beta}{2} + \frac{\beta}{2}.$$

Applying Proposition 2.1 together with Theorem 6 of (12), we can conclude that

$$\limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_N^\theta(x \leq x_N^* \leq x + \delta, \max_{1 \dots N} |x_i| \leq M) \leq -\frac{\beta}{2} \log \frac{\beta}{2} + \frac{\beta}{2} + \sup_{z \in [x, x + \delta]} [\Phi_\beta(z, \sigma_\beta) + I_{\sigma_\beta}^\beta(z, \theta)].$$

One can easily see that $z \mapsto \Phi_\beta(z, \sigma_\beta)$ is continuous on $(\sqrt{2\beta}, +\infty)$ and the continuity of $I_{\sigma_\beta}^\beta(\cdot, \theta)$ will be shown in the proof of Lemma 3.4 below. We therefore get the upper bound :

$$\limsup_{\delta \downarrow 0} \limsup_{N \rightarrow \infty} \frac{1}{N} \log \mathbb{P}_N^\theta(x \leq x_N^* \leq x + \delta, \max_{1 \dots N} |x_i| \leq M) \leq -\frac{\beta}{2} \log \frac{\beta}{2} + \frac{\beta}{2} + \Phi_\beta(x, \sigma_\beta) + I_{\sigma_\beta}^\beta(x, \theta).$$

• We now conclude the proof of Proposition 3.1 by showing the lower bound (3). We proceed as in (4). Let $y > x > r > \sqrt{2\beta}$. Then,

$$\begin{aligned} \mathbb{P}_N^\theta(y \geq x_N^* \geq x) &\geq \mathbb{P}_N^\theta(x_1 \in [x, y], \max_{i=2 \dots N} |x_i| \leq r) \\ &\geq C_N^\beta \exp \left((N-1) \inf_{\substack{z \in [x, y] \\ \mu \in B_r(\sigma_\beta, N^{-\kappa})}} (\Phi_\beta(z, \mu) + I_\mu(z, \theta) - g_\kappa(x-y)) \right) \\ &\quad \cdot \mathbb{P}_N^{N-1}(\hat{\pi}_N \in B_r(\sigma_\beta, N^{-\kappa})), \end{aligned}$$

where $B_r(\sigma_\beta, N^{-\kappa}) = B(\sigma_\beta, N^{-\kappa}) \cap \mathcal{P}([-r, r])$, with $\mathcal{P}([-r, r])$ the set of probability measure whose support is included in $[-r, r]$ and g_κ going to zero at zero by virtue of Proposition 2.1. We proceed as for the upper bound to show that $\mathbb{P}_N^{N-1}(\hat{\pi}_N \in B_r(\sigma_\beta, N^{-\kappa}))$ is going to 1. Knowing the asymptotics of C_N^β , we get

$$\liminf_{N \rightarrow \infty} \mathbb{P}_N^\theta(y \geq x_N^* \geq x) \geq -\frac{\beta}{2} \log \frac{\beta}{2} + \frac{\beta}{2} + \inf_{z \in [x, y]} (\Phi_\beta(z, \sigma_\beta) + I_{\sigma_\beta}^\beta(z, \theta) - g_\kappa(x-y)).$$

We let now y decrease to x . $\Phi_\beta(\cdot, \sigma_\beta)$ and $I_{\sigma_\beta}^\beta(\cdot, \theta)$ are continuous on $(\sqrt{2\beta}, +\infty)$ (see Lemma 3.4 below) so that we have the required lower bound

$$\liminf_{y \rightarrow x} \liminf_{N \rightarrow \infty} \mathbb{P}_N^\theta(y \geq x_N^* \geq x) \geq -\frac{\beta}{2} \log \frac{\beta}{2} + \frac{\beta}{2} + \Phi_\beta(x, \sigma_\beta) + I_{\sigma_\beta}^\beta(x, \theta).$$

This concludes the proof of Proposition 3.1. □

3.2 Proof of Theorem 1.1

We first introduce a few notations that will be useful.

Definition 3.3. For μ a compactly supported measure, we define H_μ its Hilbert transform by

$$H_\mu : \mathbb{R} \setminus \text{co}(\text{supp } \mu) \rightarrow \mathbb{R}$$

$$z \mapsto \int \frac{1}{z - \lambda} d\mu(\lambda).$$

with $\text{co}(\text{supp } \mu)$ the convex envelope of the support of μ .

It is easy to check that H_μ is injective, therefore we can define its inverse G_μ defined on the image of H_μ such that $G_\mu(H_\mu(z)) = z$. The R -transform R_μ is then given, for $z \neq 0$, by $R_\mu(z) = G_\mu(z) - \frac{1}{z}$. Moreover, one can check that $l := \lim_{z \rightarrow 0} R_\mu(z)$ exists and we let $R_\mu(0) = l$, so that R_μ is continuous at 0.

Lemma 3.4. F_θ^β is lower semicontinuous with compact level sets.

Proof. We know from Theorem 6 in (12) that

$$I_\mu^\beta(x, \theta) = \theta v(x, \theta) - \frac{\beta}{2} \int \log \left(1 + \frac{2\theta}{\beta} v(x, \theta) - \frac{2\theta}{\beta} \lambda \right) d\mu(\lambda),$$

$$\text{with } v(x, \theta) := \begin{cases} R_\mu \left(\frac{2\theta}{\beta} \right), & \text{if } H_\mu(x) \geq \frac{2\theta}{\beta}, \\ x - \frac{\beta}{2\theta}, & \text{otherwise,} \end{cases}.$$

Therefore, we have to check that $I_{\sigma_\beta}^\beta(x, \theta)$ is continuous at x^* satisfying $H_{\sigma_\beta}(x^*) = \frac{2\theta}{\beta}$ for a θ such that $x^* > \sqrt{2\beta}$. From Definition 3.3, we see that $v(\cdot, \theta)$ is continuous at x^* . Moreover as $x^* > \sqrt{2\beta}$, it is outside the support of σ_β and $x \mapsto \int \log(x - \lambda) d\sigma_\beta(\lambda)$ is continuous at x^* . Therefore F_θ^β is continuous on $(\sqrt{2\beta}, +\infty)$ and lower semi-continuous on \mathbb{R} .

Moreover, we can check that, for x large enough, $I_{\sigma_\beta}^\beta(x, \theta) \leq \theta x$, so that $F_\theta^\beta(x) \sim_{+\infty} \frac{1}{2}x^2$, its level sets are therefore compact. \square

We now go to the proof of Theorem 1.1. If we define $L_\theta^\beta(x) := F_\theta^\beta(x) - \inf_{x \in \mathbb{R}} F_\theta^\beta(x)$, then L_θ^β is a good rate function and a direct consequence of Proposition 3.1 is that, for

$$Z_N^{\beta, \theta} = \int \dots \int \prod_{i < j} |x_i - x_j|^\beta I_N^\beta(\theta, X_N) e^{-\frac{N}{2} \sum_{i=1}^N x_i^2} dx_1 \dots dx_N,$$

we have that $\lim_{N \rightarrow \infty} \frac{1}{N} \log \frac{Z_N^{\beta, \theta}}{Z_N^\beta} = \inf_{x \in \mathbb{R}} F_\theta^\beta(x)$ so that, under \mathbb{Q}_N^θ , x^* satisfies a large deviation principle with good rate function L_θ^β .

To conclude the proof, we have to study the function F_θ^β and show that L_θ^β coincide with the function K_θ^β as defined in Theorem 1.1.

We recall that, for $x \geq \sqrt{2\beta}$, we have

$$F_\theta^\beta(x) = -\frac{\beta}{2} + \frac{\beta}{2} \log \frac{\beta}{2} - \beta \int \log |x - y| d\sigma_\beta(y) + \frac{1}{2}x^2 - I_{\sigma_\beta}^\beta(x, \theta),$$

where

Relying for example on the proof of Lemma 2.7 in (5), we have that

$$-\frac{\beta}{2} + \frac{\beta}{2} \log \frac{\beta}{2} - \beta \int \log |x - y| d\sigma_\beta(y) + \frac{1}{2}x^2 = \int_{\sqrt{2\beta}}^x \sqrt{z^2 - 2\beta} dz, \quad (10)$$

so that

$$F_\theta^\beta(x) = \int_{\sqrt{2\beta}}^x \sqrt{z^2 - 2\beta} dz - I_{\sigma_\beta}^\beta(x, \theta). \quad (11)$$

From Lemma 2.7 in (5), we also get that

$$H_{\sigma_\beta}(x) = \frac{1}{\beta}(x - \sqrt{x^2 - 2\beta}) \text{ for } x > \sqrt{2\beta}, \quad (12)$$

from which we deduce that $\lim_{x \rightarrow \sqrt{2\beta}} H_{\sigma_\beta}(x) = \frac{\sqrt{2\beta}}{\beta} = \sqrt{\frac{2}{\beta}}$ and H_{σ_β} is decreasing.

• If $\theta \geq \sqrt{\frac{\beta}{2}}$, then for all $x > \sqrt{2\beta}$, $H_{\sigma_\beta}(x) \geq \frac{2\theta}{\beta}$ and

$$I_{\sigma_\beta}^\beta(x, \theta) = \theta x - \frac{\beta}{2} - \frac{\beta}{2} \log \left(\frac{2\theta}{\beta} \right) - \frac{\beta}{2} \int \log |x - y| d\sigma_\beta(y) := S_\theta(x),$$

so that from (10) and (11), we get that

$$F_\theta^\beta(x) = \frac{1}{2} \int_{\sqrt{2\beta}}^x \sqrt{z^2 - 2\beta} dz - \theta x + \frac{1}{4}x^2 + \frac{\beta}{4} - \frac{\beta}{2} \log \frac{\beta}{2} + \frac{\beta}{2} \log \theta.$$

Differentiating this function on $(\sqrt{2\beta}, +\infty)$, we see that it is decreasing on $(\sqrt{2\beta}, \theta + \frac{\beta}{2\theta})$ and then increasing so that its infimum is reached at $\theta + \frac{\beta}{2\theta}$. This gives immediately in this case that $K_\theta^\beta(x) = F_\theta^\beta(x) - \inf_x F_\theta^\beta(x) = L_\theta^\beta(x)$, as defined in Theorem 1.1..

• If $\theta \leq \sqrt{\frac{\beta}{2}}$, then we can check that on $[\sqrt{2\beta}, \theta + \frac{\beta}{2\theta}]$, we have $H_{\sigma_\beta}(x) \geq \frac{2\theta}{\beta}$ and

$I_{\sigma_\beta}^\beta(x, \theta) = \frac{\beta}{2} \int_0^{\frac{2\theta}{\beta}} R_{\sigma_\beta}(u) du$. Moreover, from (12), we get that the inverse of H_{σ_β} is given by $G_{\sigma_\beta}(x) = \frac{\beta}{2}x + \frac{1}{x}$ so that $R_{\sigma_\beta}(x) = \frac{\beta}{2}x$ and $I_{\sigma_\beta}^\beta(x, \theta) = \frac{1}{2}\theta^2$.

In this case, $F_\theta^\beta(x) = \int_{\sqrt{2\beta}}^x \sqrt{z^2 - 2\beta} dz - \frac{1}{2}\theta^2$, which is increasing.

For $x > \theta + \frac{\beta}{2\theta}$, $H_{\sigma_\beta}(x) \leq \frac{2\theta}{\beta}$, so that $I_{\sigma_\beta}^\beta(x, \theta) = S_\theta(x)$ as above.

Therefore, F_θ^β is increasing on $[\sqrt{2\beta}, \theta + \frac{\beta}{2\theta}]$ and on $[\theta + \frac{\beta}{2\theta}, +\infty)$ and is lower-semicontinuous so that its infimum is reached at $\sqrt{2\beta}$ and is equal to $-\frac{1}{2}\theta^2$.

Therefore, $K_\theta^\beta(x) = \int_{\sqrt{2\beta}}^x \sqrt{z^2 - 2\beta} dz$ on $[\sqrt{2\beta}, \theta + \frac{\beta}{2\theta}]$ and coincides with $M_\theta^\beta(x)$ on $[\theta + \frac{\beta}{2\theta}, +\infty)$. This concludes the proof of Theorem 1.1.. \square

4 Proof of Corollary 1.3

In the proof of Theorem 1.1. above, we saw that K_θ^β is increasing on $[\sqrt{2\beta}, +\infty)$ if $\theta \leq \sqrt{\frac{\beta}{2}}$, so that in this case its infimum is reached at $\sqrt{2\beta}$.

We also saw that, differentiating F_θ^β on $(\sqrt{2\beta}, +\infty)$, we got that when $\theta \geq \sqrt{\frac{\beta}{2}}$, it reaches its minimum at $\theta + \frac{\beta}{2\theta}$. This is enough to conclude. \square

Acknowledgments : I would like to thank Sandrine Péché for many fruitful discussions during the preparation of the paper.

References

- [1] Z. D. Bai, *Convergence rate of expected spectral distributions of large random matrices. I. Wigner matrices*, Ann. Probab. **21** (1993), no. 2, 625–648. MR1217559 (95a:60039)
- [2] Z. D. Bai, *Methodologies in spectral analysis of large-dimensional random matrices, a review*, Statist. Sinica **9** (1999), no. 3, 611–677, With comments by G. J. Rodgers and Jack W. Silverstein; and a rejoinder by the author. MR1711663 (2000e:60044)
- [3] J. Baik, G. Ben Arous, and S. Péché, *Phase transition of the largest eigenvalue for non-null complex sample covariance matrices*, Ann. Probab. **33** (2005), no. 5, 1643–1697. MR2165575 (2006g:15046)
- [4] G. Ben Arous, A. Dembo, and A. Guionnet, *Aging of spherical spin glasses*, Probab. Theory Related Fields **120** (2001), no. 1, 1–67. MR1856194 (2003a:82039)
- [5] G. Ben Arous and A. Guionnet, *Large deviations for Wigner’s law and Voiculescu’s non-commutative entropy*, Probab. Theory Related Fields **108** (1997), no. 4, 517–542. MR14656140 (98i:15026)
- [6] E. Brézin and S. Hikami, *Correlations of nearby levels induced by a random potential*, Nuclear Phys. B **479** (1996), no. 3, 697–706. MR1418841 (97j:82080)
- [7] P. Deift, T. Kriecherbauer, K. T.-R. McLaughlin, S. Venakides, and X. Zhou, *Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory*, Comm. Pure Appl. Math. **52** (1999), no. 11, 1335–1425. MR1702716 (2001g:42050)
- [8] P. A. Deift, *Orthogonal polynomials and random matrices: a Riemann-Hilbert approach*, Courant Lecture Notes in Mathematics, vol. 3, New York University Courant Institute of Mathematical Sciences, New York, 1999. MR1677884 (2000g:47048)
- [9] N. El Karoui, *Recent results about the largest eigenvalue of random covariance matrices and statistical application.*, Acta Phys.Polon. B **36** (2005), no. 9, 2681–2697. MR2188088

- [10] N. El Karoui, *Tracy-widom limit for the largest eigenvalue of a large class of complex sample covariance matrices.*, Annals of Probability **35** (2007), no. 2. MR2308592
- [11] D. Féral and S. Péché, *The largest eigenvalue of rank one deformation of large wigner matrices*, Comm.Math.Phys **272** (2007), no. 1, 185–228. MR2291807
- [12] A. Guionnet and M. Maïda, *A Fourier view on the R-transform and related asymptotics of spherical integrals*, J. Funct. Anal. **222** (2005), no. 2, 435–490. MRMR2132396 (2006b:60043)
- [13] A. Guionnet and O. Zeitouni, *Concentration of the spectral measure for large matrices*, Electronic Communicartions in Probability **5** (2000). MR1781846 (2001k:15035)
- [14] A. Guionnet and O. Zeitouni, *Large deviations asymptotics for spherical integrals*, J. Funct. Anal. **188** (2002), no. 2, 461–515. MR1883414 (2003e:60055)
- [15] D. Hoyle and M. Rattray, *Limiting form of the sample covariance eigenspectrum in pca and kernel pca*, Proceedings of Neural Information Processing Systems, 2003.
- [16] K. Johansson, *Universality of the local spacing distribution in certain ensembles of Hermitian Wigner matrices*, Comm. Math. Phys. **215** (2001), no. 3, 683–705. MR1810949 (2002j:15024)
- [17] I. Johnstone, *On the distribution of the largest eigenvalue in principal components analysis.*, Annals of Statistics **29** (2001), 295–327. MR1863961 (2002i:62115)
- [18] L. Laloux, P. Cizeau, M. Potters, and J. Bouchaud, *Random matrix theory and financial correlations*, Intern. J. Theor. Appl. Finance **3** (2000), no. 3, 391–397.
- [19] M. L. Mehta, *Random matrices*, second ed., Academic Press Inc., Boston, MA, 1991. MR1083764 (92f:82002)
- [20] S. Péché, *The largest eigenvalue of small rank perturbations of hermitian random matrices*, Probab. Theory Related Fields **134** (2006), no. 1, 127–173. MR2221787 (2007d:15041)
- [21] C. A. Tracy and H. Widom, *Level-spacing distributions and the Airy kernel*, Comm. Math. Phys. **159** (1994), no. 1, 151–174. MR1257246 (95e:82003)
- [22] E. P. Wigner, *On the distribution of the roots of certain symmetric matrices*, Ann. of Math. (2) **67** (1958), 325–327. MR0095527 (20 #2029)