



Large Magneto-ionic Variations toward the Galactic Center Magnetar, PSR J1745-2900

DOI:
[10.3847/2041-8213/aaa2f8](https://doi.org/10.3847/2041-8213/aaa2f8)

Document Version
Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Desvignes, G., Eatough, R. P., Pen, U. L., Lee, K. J., Mao, S. A., Karuppusamy, R., Schnitzeler, D. H. F. M., Falcke, H., Kramer, M., Wucknitz, O., Spitler, L. G., Torne, P., Liu, K., Bower, G. C., Cognard, I., Lyne, A. G., & Stappers, B. W. (2018). Large Magneto-ionic Variations toward the Galactic Center Magnetar, PSR J1745-2900. *Astrophysical Journal Letters*, 852(1), [L12]. <https://doi.org/10.3847/2041-8213/aaa2f8>

Published in:
Astrophysical Journal Letters

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



LARGE MAGNETO-IONIC VARIATIONS TOWARDS THE GALACTIC CENTRE MAGNETAR,
PSR J1745–2900

G. DESVIGNES,¹ R. P. EATOUGH,¹ U. L. PEN,^{2,3,4,5} K. J. LEE,⁶ S. A. MAO,¹ R. KARUPPUSAMY,¹
D. H. F. M. SCHNITZELER,^{1,7} H. FALCKE,^{8,9,1} M. KRAMER,^{1,10} O. WUCKNITZ,¹ L. G. SPITLER,¹ P. TORNE,¹¹ K. LIU,¹
G. C. BOWER,¹² I. COGNARD,^{13,14} A. G. LYNE,¹⁰ AND B. W. STAPPERS¹⁰

¹*Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany*

²*Canadian Institute for Theoretical Astrophysics, University of Toronto, 60 St. George Street, Toronto, Ontario M5S 3H8, Canada*

³*Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, Ontario M5S 3H4, Canada*

⁴*Canadian Institute for Advanced Research, CIFAR Program in Gravitation and Cosmology, Toronto, Ontario M5G 1Z8, Canada*

⁵*Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario, N2L 2Y5, Canada*

⁶*Kavli institute for astronomy and astrophysics, Peking University, Beijing 100871, P.R. China*

⁷*Bendenweg 51, 53121 Bonn, Germany*

⁸*Department of Astrophysics, Institute for Mathematics, Astrophysics and Particle Physics,*

Radboud University, PO Box 9010, 6500 GL Nijmegen, The Netherlands

⁹*ASTRON, PO Box 2, 7990 AA Dwingeloo, The Netherlands*

¹⁰*Jodrell Bank Centre for Astrophysics, School of Physics and Astronomy, The University of Manchester, Manchester M13 9PL, UK*

¹¹*Instituto de Radioastronomía Milimétrica, Avda. Divina Pastora 7, Núcleo Central, E-18012 Granada, Spain*

¹²*Academia Sinica Institute of Astronomy and Astrophysics, 645 N. A'ohoku Place, Hilo, HI 96720, USA*

¹³*Laboratoire de Physique et Chimie de l'Environnement et de l'Espace LPC2E CNRS-Université d'Orléans, F-45071 Orléans, France*

¹⁴*Station de radioastronomie de Nançay, Observatoire de Paris, CNRS/INSU F-18330 Nançay, France*

(Received; Revised; Accepted)

Submitted to ApJ

Abstract

Polarised radio emission from PSR J1745–2900 has already been used to investigate the strength of the magnetic field in the Galactic Centre, close to Sagittarius A*. Here we report how persistent radio emission from this magnetar, for over four years since its discovery, has revealed large changes in the observed Faraday rotation measure, by up to 3500 rad m⁻² (a five per cent fractional change). From simultaneous analysis of the dispersion measure, we determine that these fluctuations are dominated by variations in either the projected magnetic field or the free electron content within the Galactic Centre, along the changing line of sight to the rapidly moving magnetar. From a structure function analysis of rotation measure variations, and a recent epoch of rapid change of rotation measure, we determine a minimum scale of magneto-ionic fluctuations of size ~ 2 au at the Galactic Centre distance, inferring PSR J1745–2900 is just ~ 0.1 pc behind an additional scattering screen.

Keywords: pulsars: individual (J1745-2900) — Galaxy: center — magnetic fields

1. INTRODUCTION

Measurements of Faraday Rotation in the polarised emission of radio sources can be used to examine the strength and structure of the magnetic field in the interstellar medium (Beck & Wielebinski 2013). A recent notable example is the radio-loud magnetar, PSR J1745–2900, which displays a high rotation measure (RM) of $-66960 \pm 50 \text{ rad m}^{-2}$ as measured in May 2013, second only in the Galaxy to the RM = $(-4.3 \pm 0.1) \times 10^5 \text{ rad m}^{-2}$ of the supermassive black hole candidate, Sagittarius A* (Sgr A*), caused predominantly by the accretion flow on scales smaller than the Bondi-Hoyle radius (Bower et al. 2003). This magnetar therefore allowed first-order estimates of the strength of the magnetic field at the beginnings of the Bondi-Hoyle accretion flow of Sgr A*; $\sim 8 \text{ mG}$ at scales of $\sim 0.1 \text{ pc}$ (Eatough et al. 2013). The magnitude and spatial or time variability of the magnetic field also allows models of the accretion flow to be investigated (Pang et al. 2011).

PSR J1745–2900 has also been used to examine the scattering of radio waves towards the Galactic Centre (GC) by measurements of both the temporal pulse broadening (Spitler et al. 2014) and the angular image broadening (Bower et al. 2014). Combination of the two measurements indicates the principal scattering screen toward the GC is $5.8 \pm 0.3 \text{ kpc}$ in front of the magnetar (Bower et al. 2014).

Atypically for magnetars, PSR J1745–2900 has remained active in the radio band for over four years since its discovery at X-ray wavelengths in 2013 (Kennea et al. 2013; Mori et al. 2013). This has allowed repeated measurements of the RM and dispersion measure (DM). Because the magnetar has a total proper motion of $6.37 \pm 0.16 \text{ mas yr}^{-1}$ relative to Sgr A* (Bower et al. 2015), the time-variations in the measurements of DM and RM presented here occur along different sightlines. The physical scales probed in the GC are therefore directly related to the observing cadence, and the transverse velocity of the magnetar.

In this paper, long term polarimetric observations of PSR J1745–2900 with the Effelsberg and Nançay radio telescopes are presented. Section 2 gives a description of the observational campaign and data analysis techniques used. In Section 3 the results of the analysis are presented, and in Section 4 we turn to physical interpretations of the results.

2. OBSERVATIONS

Following the discovery of radio pulsations from PSR J1745–2900 in April 2013, this source has been monitored with three European radio telescopes oper-

ating at complementary observing frequencies. These are the Effelsberg radio telescope, the Nançay Radio Telescope (NRT) and the Jodrell Bank, Lovell radio telescope. In this work we only refer to measurements from the Effelsberg telescope and the NRT because observations with the Lovell telescope, at a lower frequency of 1.4 GHz, suffer from instrumental depolarisation due to the large RM.

2.1. Effelsberg

PSR J1745–2900 is observed with the Effelsberg telescope typically on a monthly basis, and fortnightly since January 2017. One-hour observations at central frequencies of 8.35 GHz and 4.85 GHz are recorded with the PSRIX backend (Lazarus et al. 2016).

The 500 MHz bandwidth provided by the PSRIX backend is first digitised and split into 512 or 1024 channels when observing at 8.35 GHz and 4.85 GHz, respectively. The channelised data are then coherently dedispersed at a DM of 1778 pc cm^{-3} , the initial DM value measured by Eatough et al. (2013) and finally folded at the period of the magnetar to create ‘single pulse profiles’.

Since March 2017 the new ‘C+’ broadband receiver is available for pulsar observations. Two 2 GHz bands (covering 4 to 8 GHz) are fed into the new PSRIX2 backend, consisting of two CASPER¹ ROACH2 boards. Each board digitises the signal and acts as a full-Stokes spectrometer, creating 2048 frequency channels every $8 \mu\text{s}$. The data are later dedispersed and folded to create single pulses profiles. During the commissioning phase of PSRIX2, the C+ observations replace the 8.35 GHz observations.

2.2. Nançay

Observations with the NRT were carried out with the NUPPI instrumentation on average every 4 days between May 2013 and August 2014, before resuming at a monthly cadence between January and July 2017. The setup of the observations was already presented in Eatough et al. (2013); Spitler et al. (2014); Torne et al. (2015). We briefly summarize it here. A bandwidth of 512 MHz centred at 2.5 GHz is split into 1024 channels and coherently dedispersed using an initial DM value of 1840 pc cm^{-3} (the best DM value derived from timing of the scattered pulse profiles) then folded at the period of the magnetar and written to disk every 30 s.

2.3. Post-processing and calibration

¹ <https://casper.berkeley.edu/>

All the data presented here are corrected for the gain and phase difference between the feeds of the various receivers used. This is achieved by standard pulsar calibration techniques that use observations of a polarised pulsed noise diode. Large $|\text{RM}|$ s, in combination with wide frequency channels, lead to instrumental depolarization. Following the analysis presented in [Schnitzeler & Lee \(2015\)](#), we calculate that an $|\text{RM}|$ of $7 \times 10^4 \text{ rad m}^{-2}$ depolarizes the signal by only 4 per cent in the 2.5 GHz band; at the higher observing frequencies, this effect is negligible. The data reduction is achieved with the standard PSRCHIVE package ([Hotan et al. 2004](#)).

3. RESULTS

Monitoring of the dispersive and polarisation properties of PSR J1745–2900 over a period of ~ 54 months has revealed a rather constant DM of $1762 \pm 11 \text{ cm pc}^{-3}$, while variations in $\text{RM} > 3500 \text{ rad m}^{-2}$ are observed. Because RM is proportional to both projected magnetic field strength B_{\parallel} , and the free electron density n_e , along the line-of-sight s , $\text{RM} \propto \int B_{\parallel}(s)n_e(s)ds$, it is important to disentangle these quantities to understand the physical mechanisms causing the variations in RM. For PSR J1745–2900, which has the highest DM amongst all the known pulsars, measurement of the DM is influenced by pulse scattering. In this section the methods used to measure both the DM and RM are described, as well as the results of our monitoring campaign.

3.1. Dispersion Measure variations

To accurately measure the DM and remove the bias caused by the scattering of the pulse profile (and possibly the variations of it), we modelled both scattering and DM simultaneously over a range of frequency subbands for each NUPPI, PSRIX and PSRIX2 observations. Given the low amount of dispersion across the band of the PSRIX data at 8.35 GHz ($\sim 12 \text{ ms}$), we did not apply this technique to these data.

Following [Spitler et al. \(2014\)](#), we use a scattered Gaussian pulse function to model the single pulses observed between 4 and 8 GHz and the averaged pulse profile at 2.5 GHz. However, in contrast to [Spitler et al. \(2014\)](#), we did not correct for the jittering of the single pulses to create an average ‘de-jittered’ profile. Instead, we model simultaneously some of the brightest single pulses in each observation with different Gaussian widths σ to increase the significance of our results.

The scattered Gaussian pulse profile function for a single channel is given by Equation 3 of [Spitler et al. \(2014\)](#). We extend it here for multiple channels to include the DM as a parameter. We can therefore write

the likelihood Λ as

$$\log \Lambda = -\frac{1}{2} \sum_i^{N_p} \sum_j^{N_c} \sum_k^{N_b} \left\{ \frac{[P_{ijk} - T_{ijk}]^2}{\sigma_{ij}^2} \right\} + \text{const}, \quad (1)$$

where P_{ijk} is the observed single pulse profile i with frequency channel j and profile bin k over N_p profiles included in the modelling with N_c frequency channels and N_b phase bins. σ_{ij}^2 is the variance of the noise in the profile i, j . T_{ijk} is the modelled scattered profile,

$$T_{ijk} = \frac{A_{ij}\sqrt{\pi}}{2} w_i e^{-\frac{w_i^2 - 4\phi_{ijk}\tau}{4\tau^2}} \times \left\{ 1 + \text{erf} \left(\frac{2\tau(\phi_{ijk} - \phi_{0,ij}) - w_i^2}{2\tau w_i} \right) \right\} + b_{ij}. \quad (2)$$

A_{ij} and b_{ij} are, respectively, the amplitude and the baseline offset of channel j from profile i . $\phi_{0,ij}$ is the phase centre of the Gaussian of width w_i , but delayed in each channel because of dispersion. $\phi_{0,ij}$ is given by $\phi_{0,ij} = \phi_{0,i} + k\text{DM} \times (1/f_i^2 - 1/f_0^2)$, where the dispersion constant $k = 4.15 \times 10^3 \text{ MHz}^2 \text{ cm}^3 \text{ pc}^{-1} \text{ s}$. Finally, τ is the scattering time at a reference frequency f_0 . The dimensionality of the model is therefore $2 + N_p \times (2 + 2 \times N_c)$, where N_p is 1 for the NRT averaged profile, typically 2 to 4 for the Effelsberg single pulse observations and N_c range from 8 to 16.

Given the large dimensionality of the model, we implemented the nested sampling tool POLYCHORD ([Handley et al. 2015](#)) to efficiently sample the parameter space instead of using traditional χ^2 -fitting techniques. The priors on all parameters are set to be uniform, except for A_{ij} where we use log-uniform priors. We refer to one-sigma error bars as the 68.3 per cent contours of the one-dimensional marginalized posterior distribution of each parameter. The resulting code can be found online². An example of our modelling results can be seen in Figure 1.

The DM results of the modelling of the Effelsberg and NRT data are shown in Figure 2. A linear least-squares fit to the DM values shows a marginal decrease of approximately 0.6 per cent over four years (two-sigma consistent with no DM change) with $\text{DM} = 1762 \pm 11 \text{ cm pc}^{-3}$ as measured in October 2017. Results from the scattering measurements that come out of this joint analysis will be published elsewhere.

3.2. Rotation Measure variations

To determine the Faraday rotation of the linearly polarised emission of the magnetar, we used the RM

² <https://github.com/gdesvignes/scattering>




Figure 1. Scattered and dispersed observed pulse profiles of PSR J1745–2900 (over a selected phase window) are shown in black color for a set of 16 subbands from a C+ single pulse profile (top panel) and a 2.5 GHz daily averaged profile (bottom panel). The red curves show our model of scattered and dispersed Gaussian profiles taken from the maximum likelihood solution. The MJD of the observation and the estimated DM are indicated at the bottom-left part of each panel.




Figure 2. DM variations of the GC magnetar. The black crosses, red triangles and green stars show the DM values (with one-sigma error bars) from the modelling of the scattered profiles from the 2.5 GHz, 4.85 GHz and C+ data, respectively. The dashed line represents the linear least-squares fit to the data. The MJD and modelled DM are reported in the bottom-left part of each panel.

synthesis method (Brentjens & de Bruyn 2005) and fitted the wrapping of the Stokes vector Q and U under the pulse window as previously done in Eatough et al. (2013). We employed this technique to extract the RM for all our data. In contrast to our DM results, Figure 3 shows the rapid and non-monotonic evolution of the RM during our 4 years of observations.




Figure 3. Variation of RM as a function of MJD. The black crosses, red triangles, blue points and green stars show the RM values estimated from the 2.5 GHz, 4.85 GHz, 8.35 GHz and C+ data, respectively. The orange rectangle shows the value from the ATCA data recorded at 5.4 GHz (Schnitzeler et al. 2016). The black dashed line shows the fitted slope of the recent epoch of large gradient in RM variation.

Since the initial $\text{RM} = -66960 \pm 50 \text{ rad m}^{-2}$ reported in Eatough et al. (2013), the RM changed by over 3500 units to $\text{RM} = -63402 \pm 232 \text{ rad m}^{-2}$ in October 2017 (a relative change of 5.3 per cent). Our measurements are also consistent with the $\text{RM} = -66080 \pm 24 \text{ rad m}^{-2}$ reported by Schnitzeler et al. (2016) using Australian Telescope Compact Array (ATCA) observations in May 2015 at 5.4 GHz. After a steady increase in the first few months, the signed RM increased abruptly around MJD 56566 followed by another steady increase until MJD 57450. Then the signed RM diminished up to MJD 57600 before increasing over the last year of observations. A least-squares fit to the data collected over the last year shows that RM changed by about 7.4 rad m^{-2} per day. It is worth noting that two-weeks of follow-up observations of the magnetar SGR 1806–20 after an outburst phase did not show evidence for RM variations (Gaensler et al. 2005).

3.3. Polarisation fraction

Figure 4 shows the evolution of the linear polarisation fraction L_f as a function of time for all our data. These results show large variations in L_f at all frequencies, with L_f at 2.5 GHz consistently lower than at higher frequencies, by a factor of between 2 and 10.

4. DISCUSSION

The observations presented here show a clear dichotomy between the properties of the DM and RM variations toward PSR J1745–2900; the DM varies marginally while the RM shows very large variations.




Figure 4. Polarisation fraction as a function of time. The black crosses, red triangles and blue points show the values from the 2.5 GHz, 4.85 GHz and 8.35 GHz data, respectively. The polarisation fraction at 2.5 GHz is consistently lower than at the higher frequencies.

This is not entirely unexpected since it has been suggested that the RM is likely caused by local magnetic phenomena, somewhere in the last few parsecs toward the GC, whereas most of the DM is accumulated along the entire line of sight to the magnetar (Eatough et al. 2013). If one considers the strong magnetic field of 8 mG near the Bondi-Hoyle radius, a five per cent change in the local free electron density along 0.1 pc would lead to a large RM variation of $\sim 3500 \text{ rad m}^{-2}$ (assuming $n_e \sim 100 \text{ cm}^{-3}$), but a DM change of only $\sim 0.5 \text{ pc cm}^{-3}$, indistinguishable with our current data set. Therefore our results point to magneto-ionic variations occurring in the GC.

4.1. Structure function analysis

As already mentioned, PSR J1745–2900 has a proper motion (Bower et al. 2015), so our measurements of DM and RM can be used to probe the angular and physical scales on which these fluctuations occur. For sparsely and irregularly sampled data, like these, power spectrum analyses are unsuitable. A frequently used alternative is the structure function of variations, see e.g. Minter & Spangler (1996). Following their notation, the ensemble average second order structure function in DM and RM can be written $\text{SF}_{\text{DM}} = \langle [\text{DM}(\theta) - \text{DM}(\theta + \delta\theta)]^2 \rangle$ and $\text{SF}_{\text{RM}} = \langle [\text{RM}(\theta) - \text{RM}(\theta + \delta\theta)]^2 \rangle$, respectively, where θ is any given line of sight and $\delta\theta$ is the angular separation or ‘lag’ of two measurements. In this work the lag is given by the total magnitude of the proper motion (6.37 mas yr^{-1}) multiplied by the separation in




Figure 5. Structure functions, corrected for measurement errors, for the DM (top panel) and RM (bottom panel). For clarity, the error bars for each lag (grey crosses) are not shown. The red points represent the averaged values over intervals of 0.25 in log scale. The horizontal black line delimits RM variations of 100 rad m^{-2} . The black dashed line shows the linear fit to the average values of SF_{RM} allowing to give an estimate on the angular scale at which the RM variations occur.

time of two DM or RM measurements³. In addition, we corrected each pair of DM and RM used in the structure function for noise bias caused by measurement uncertainties (You et al. 2007). This is achieved by subtracting respectively $(\sigma_{\text{DM}(\theta)}^2 + \sigma_{\text{DM}(\theta+\delta\theta)}^2)$ and $(\sigma_{\text{RM}(\theta)}^2 + \sigma_{\text{RM}(\theta+\delta\theta)}^2)$, where σ_{DM} and σ_{RM} are the individual measurement errors in DM and RM. In Figure 5 we present the results from the structure function analysis. In the top panel, averaged values of SF_{DM} are unchanging across all angular scales from 10^{-10} deg to 10^{-5} deg . This is representative of the flat fit of the DM time series in Section 3.1. The bottom panel of Figure 5 shows SF_{RM} over the same interval in angular scale. Unlike SF_{DM} , SF_{RM} begins rising at angular scales over 10^{-8} deg , equivalent to a physical size of a few au at a GC distance of 8.3 kpc (Gillessen et al. 2009). We define this size as the approximate smallest-scale at which magneto-ionic structure in the GC is observed in our data.

The fitted slope of SF_{RM} (black dashed line), $l = 1.23 \pm 0.13$, is over $3\text{-}\sigma$ away from the expected value in case of Kolmogorov turbulence (where $l = 5/3$) assuming isotropic and stationary fluctuations in RM (Simonetti et al. 1984; Lazio et al. 1990; Haverkorn et al. 2004).

³ Here we assume that the relative proper motion between the magneto-ionic material and the magnetar is of the same order of magnitude as that between the magnetar and Sgr A*

4.2. A physical model based on the depolarisation

Figure 4 shows that PSR J1745–2900 depolarises rapidly with decreasing frequency. This could be intrinsic to the magnetar, or caused by a propagation effect. Because the former has not been observed in magnetars (e.g. Kramer et al. 2007), in this section we model depolarisation due to propagation through the interstellar medium.

For this purpose we invoke a secondary scattering screen, close to the magnetar in the GC, in addition to the distant screen (~ 6 kpc from the GC) responsible for the temporal scatter broadening (Bower et al. 2014; Wucknitz 2014). The distant screen cannot produce the rapid RM variations we observe because of its large size (160 mas at 2.5 GHz) and the fact that spiral arms do not host strong magnetic fields (Haverkorn 2015). Also the size of the distant screen, when scaled to the location of the magnetar, gives a physical scale of ~ 0.01 au, much smaller than the required physical scale (see below). Therefore a secondary scattering screen, close to the GC, is necessary.

Scattering screens located close to the GC (< 700 pc) have been previously indicated for other GC pulsars (Dexter et al. 2017). As the magnetar moves behind or through a magnetized, ionized gas, the variation in RM that we observe over time also imprints itself as a gradient in RM across this secondary scattering disk. Following Schnitzeler et al. (2015), we can show that RM variations of order 100 rad m^{-2} within a scattering disk (Gaussian disk with a FWHM of 0.2–0.5 mas or a uniformly lit circular disk with diameter < 1 mas at 2.5 GHz) is enough to explain little or no depolarization at the highest observing frequencies, and the strong or even complete depolarization at 2.5 GHz. At this frequency an RM gradient of 100 rad m^{-2} across the scattering disk would lead to PA differences up to ~ 80 deg, leading to strong depolarisation.

By considering both measurements from the RM structure function analysis outlined in Section 4.1, and the recently measured large gradient in RM (the fitted slope described in Section 3.2), fluctuations of order 100 rad m^{-2} are readily observable in our data set and occur at angular-scales, or time-scales, of the order tenths of milliarseconds and weeks respectively.

The solid horizontal black line at $\text{SF}_{\text{RM}} = 10000 \text{ rad}^2 \text{ m}^{-4}$ in Figure 5 defines the limit where changes in RM of 100 rad m^{-2} occur. The intercept of this line with the fitted averages of SF_{RM} gives the approximate angular scale at which these fluctuations begin to occur; which is around $6 \times 10^{-8} \text{ deg}$ (~ 0.22 mas or a physical scale of ~ 1.8 au at a GC distance of 8.3 kpc). The fitted slope in RM in Figure 3 indicates a change in RM

of 100 rad m^{-2} occurs on time-scales of just two weeks (angular displacement of ~ 0.24 mas, physical scale of ~ 2.0 au at distance of 8.3 kpc). Much shorter term variations (< 1 hr) have been ruled out. The average value of these two measured physical scales maps to an upper bound on the scattering disk size b of ~ 1.9 au.

The geometric scattering time delay, τ , depends upon both the magnetar’s distance to the screen, z , and its size, b , through $\tau = b^2/2cz$. τ is bounded by the observed scattering time at 2.5 GHz, and is likely substantially shorter since most of the scattering delay occurs at the midway screen (Wucknitz 2014). Following Spitler et al. (2014) we use $\tau < 0.04$ s giving $z \gtrsim 0.1$ pc. Remarkably, this places the secondary scattering screen at least 0.1 pc in front of PSR J1745–2900, the same distance as the projected offset from Sgr A*; therefore placing the screen close to the Bondi-Hoyle accretion radius. The observed magneto-ionic fluctuations are thus potentially occurring within the black hole reservoir. The two main assumptions in this model are the isotropy of the RM variations, and the attribution of depolarisation to propagation effects.

The large RM variations toward PSR J1745–2900, along with the observed depolarisation at 2.5 GHz therefore suggests an additional scattering screen local to the GC and embedded in a strong ambient magnetic field (~ 8 mG; Eatough et al. 2013). Assuming a magnetic origin for the observed RM variations, fluctuations in this field, of $\sim 12 \mu\text{G}$ starting at sizes of ~ 2 au and going up to $\sim 400 \mu\text{G}$ on the largest measured scales of ~ 300 au can occur either due to changes in its strength or orientation. Small changes in the free electron content in this same field can not be ruled out. Our data also suggest the RM fluctuations deviate from Kolmogorov turbulence.

Future observations of PSR J1745–2900 will better constrain the magnetic field structure around Sgr A* and can potentially be used to test if the accretion flow becomes magnetically dominated (Pen et al. 2003).

The authors acknowledge financial support by the European Research Council for the ERC Synergy Grant BlackHoleCam under contract no. 610058. KJL received support from 973 program 2015CB857101, NSFC U15311243, XDB23010200, 11690024, and funding from Max-Planck Partner Group. This work was based on observations with the 100-m telescope of the Max-Planck-Institut für Radioastronomie at Effelsberg. The Nançay radio Observatory is operated by the Paris Observatory, associated to the French Centre National de la Recherche Scientifique (CNRS) and to the Université

d'Orléans. The authors thank D.D. Xu, R. Beck and M. Haverkorn for helpful discussions.

REFERENCES

- Beck, R., & Wielebinski, R. 2013, *Magnetic Fields in Galaxies*, ed. T. D. Oswalt & G. Gilmore, 641
- Bower, G. C., Wright, M. C. H., Falcke, H., & Backer, D. C. 2003, *ApJ*, 588, 331
- Bower, G. C., Deller, A., Demorest, P., et al. 2014, *ApJ*, 780, L2
- . 2015, *ApJ*, 798, 120
- Brentjens, M. A., & de Bruyn, A. G. 2005, *A&A*, 441, 1217
- Dexter, J., Deller, A., Bower, G. C., et al. 2017, *MNRAS*, 471, 3563
- Eatough, R. P., Falcke, H., Karuppusamy, R., et al. 2013, *Nature*, 501, 391
- Gaensler, B. M., Kouveliotou, C., Gelfand, J. D., et al. 2005, *Nature*, 434, 1104
- Gillessen, S., Eisenhauer, F., Fritz, T. K., et al. 2009, *ApJ*, 707, L114
- Handley, W. J., Hobson, M. P., & Lasenby, A. N. 2015, *MNRAS*, 453, 4384
- Haverkorn, M. 2015, in *Astrophysics and Space Science Library*, Vol. 407, *Magnetic Fields in Diffuse Media*, ed. A. Lazarian, E. M. de Gouveia Dal Pino, & C. Melioli, 483
- Haverkorn, M., Gaensler, B. M., McClure-Griffiths, N. M., Dickey, J. M., & Green, A. J. 2004, *ApJ*, 609, 776
- Hotan, A. W., van Straten, W., & Manchester, R. N. 2004, *PASA*, 21, 302
- Kennea, J. A., Burrows, D. N., Kouveliotou, C., et al. 2013, *ApJ*, 770, L24
- Kramer, M., Stappers, B. W., Jessner, A., Lyne, A. G., & Jordan, C. A. 2007, *MNRAS*, 377, 107
- Lazarus, P., Karuppusamy, R., Graikou, E., et al. 2016, *MNRAS*, 458, 868
- Lazio, T. J., Spangler, S. R., & Cordes, J. M. 1990, *ApJ*, 363, 515
- Minter, A. H., & Spangler, S. R. 1996, *ApJ*, 458, 194
- Mori, K., Gotthelf, E. V., Zhang, S., et al. 2013, *ApJ*, 770, L23
- Pang, B., Pen, U.-L., Matzner, C. D., Green, S. R., & Liebendörfer, M. 2011, *MNRAS*, 415, 1228
- Pen, U.-L., Matzner, C. D., & Wong, S. 2003, *ApJ*, 596, L207
- Schnitzeler, D. H. F. M., Banfield, J. K., & Lee, K. J. 2015, *MNRAS*, 450, 3579
- Schnitzeler, D. H. F. M., Eatough, R. P., Ferrière, K., et al. 2016, *MNRAS*, 459, 3005
- Schnitzeler, D. H. F. M., & Lee, K. J. 2015, *MNRAS*, 447, L26
- Simonetti, J. H., Cordes, J. M., & Spangler, S. R. 1984, *ApJ*, 284, 126
- Spitler, L. G., Lee, K. J., Eatough, R. P., et al. 2014, *ApJ*, 780, L3
- Torne, P., Eatough, R. P., Karuppusamy, R., et al. 2015, *MNRAS*, 451, L50
- Wucknitz, O. 2014, in *Proceedings of the 12th European VLBI Network Symposium and Users Meeting PoS(EVN 2014)066*, Cagliari, Italy, 66
- You, X. P., Hobbs, G., Coles, W. A., et al. 2007, *Monthly Notices of the Royal Astronomical Society*, 378, 493