

# Lectins: Proteins with Diverse Applications

Rabia Hamid\*, Akbar Masood, Ishfak H. Wani, and Shaista Rafiq  
Department of Biochemistry, University of Kashmir, Srinagar - 190006, India.

---

## ARTICLE INFO

### Article history:

Received on: 12/04/2013

Revised on: 23/04/2013

Accepted on: 04/05/2013

Available online: 12/05/2013

### Key words:

Lectins, Hemagglutinins,  
Antimicrobial, antitumor,  
antiviral, therapeutic  
applications

---

## ABSTRACT

Lectins are proteins that bind to carbohydrates and sugar containing substances in a specific and reversible way or precipitate glycoconjugates. These heterogeneous class of carbohydrate-binding proteins or glycoproteins of non-immune origin are capable of specific recognition of, and reversible binding to, carbohydrates without altering their covalent structure. Lectins are found in a diversity of organisms and possess the ability to agglutinate erythrocytes with known carbohydrate specificity since they have at least one non-catalytic domain that binds reversibly to specific monosaccharides or oligosaccharides. This review aims to highlight the applications of lectins in various fields of biology. Lectins are isolated from their natural sources by chromatographic procedures with various modulations to increase their production. The yields of animal lectins are usually low compared with the yields of plant lectins such as legume lectins, which form a major source of these proteins. Lectins manifest a diversity of activities including anti-insect activities, antitumor, immunomodulatory, antimicrobial and HIV-1 reverse transcriptase inhibitory, which may find applications in many therapeutic areas. A small number of lectins demonstrate anti-parasitic activities.

---

## INTRODUCTION

“Lectin” has been derived from the Latin word “legere”, which means “to select”, by William Boyd (Boyd and Shapleigh, 1954). This term was generalized to embrace all sugar-specific agglutinins of nonimmune origin, irrespective of source and blood type specificity (Sharon and Lis, 1972). Lectins have the ability to bind carbohydrates and the name “hemagglutinins” is used when the sugar specificity is unknown. Lectins are proteins/glycoproteins, which have at least one non-catalytic domain that exhibits reversible binding to specific monosaccharides or oligosaccharides (Peumans and Van – Damme, 1995 b). They can bind to the carbohydrate moieties on the surface of erythrocytes and agglutinate the erythrocytes, without altering the properties of the carbohydrates. Lectins with specific carbohydrate specificity have been purified from various plant tissues and other organisms. They can be classified on the basis of their carbohydrate specificity. They can also be categorized according to the overall structures into merolectins, hololectins, chimerolectins and superlectins, or be grouped into different families (legume lectins, type II ribosome-inactivating

proteins, monocot mannose-binding lectins, and other lectins). The amount of lectin varies in different organisms. The high yields of lectins from different sources may facilitate mass production. Application of lectins is possible depending on their properties. The antimicrobial and anti-insect activities of lectins can be made use of in the control of pathogens. The production of anti-tumor and antiviral drugs based on lectins may also have a significant utility in therapeutic industry.

### History of lectins

Lectins were first described in 1888 by Stillmark, who observed that crude extracts of castor beans (*Ricinus communis*) contained a toxic substance named ricin that agglutinated human and some animal red blood cells. However, the modern age of lectinology started nearly 100 years later (Bies et al., 2004; Sharon and Lis, 2004).

Lectins were initially found and described in plants, but in subsequent years multiple lectins were isolated from microorganisms and also from animals (Sharon and Lis, 2004). The lectin-induced agglutination of cells has originally served as the most common assay to detect and quantify lectin activity in a variety of organisms (Vlodavsky and Sachs, 1975; Doyle and Keller, 1984; Goldhar, 1995).

---

\* Corresponding Author  
Assistant Professor Department of Biochemistry University of Kashmir  
Mob; 9419548985  
e-mail: [rabeyams@yahoo.co.in](mailto:rabeyams@yahoo.co.in)

Despite their broad applicability as a method for the detection of lectin activity, agglutination assays have considerable limitations because only multivalent lectins can agglutinate. Monovalent lectins, with only one binding site for carbohydrates, are usually not detectable by agglutination assays. Therefore, agglutination assays are mostly applied for lectins that are known to have more than one carbohydrate-binding site. Advances in biophysical and molecular biology techniques as well as the availability of synthetic oligosaccharides have contributed to the identification of many lectins. Most lectins have been purified by affinity chromatography (Agrawal and Goldstein, 1967). For the isolation and characterization of a vast number of lectins, sugar-based polymers like Sephadex (glucose), Sepharose (galactose) or Chitin (N-acetyl-glucosamine) have been used. Glycoprotein-linked matrices are applied to the purification of lectins that recognize more complex saccharides (Goldstein, 2002). Many lectins have, in addition to the carbohydrate-binding domain, another domain with distinct activity. Proteins that carry lectin domain(s) and other domains with quite different properties have been better studied in animals than in plants (Gabijs, 1994).

## PRODUCTION OF LECTINS

Lectins are found in nature. A large number of lectins or hemagglutinins have been purified from different organisms.

### Natural lectins

#### Animal lectins

Lectins are found in different animals. However, the yields are usually extremely low (Table 1). Mass purification of animal lectins necessitates bulk quantities of raw materials which make it not feasible.

#### Mushroom lectins

Yields of lectins from fresh mushrooms are low, e.g., 2.6 mg from 100 g of fresh fruiting bodies of *Pleurocybella porrigens* (Suzuki *et al.* 2009). In fact, the water content in fresh mushrooms is very high. Dried fruiting bodies of the mushrooms *Russula lepida*, *Pholiota adiposa*, and *Inocybe umbrinella* yielded 39, 70, and 15 mg lectin per 100 g fruiting bodies, respectively (Zhang *et al.* 2010, 2009; Zhao *et al.* 2009). Therefore, production from fresh mushroom is also unpractical.

#### Plant lectins

The lectin contents in some parts of plants are higher, e.g., 390 and 75 mg of the purified lectin was recovered from 100 g *Remusatia vivipara* tubers (Bhat *et al.* 2010) and *Astragalus mongholicus* roots (Yan *et al.* 2005), respectively. Lectins are also found in seeds. The lectin content in non legume plants is low, e.g., 3.3 mg lectin from 100 g *Hibiscus mutabilis* seeds (Lam and Ng 2009). Lectins are found in abundance in legume seeds. *Phaseolus vulgaris* is an herbaceous annual plant grown worldwide for its edible beans, popular in both dry and green bean forms. The commercial production of beans is well distributed worldwide.

There are different varieties, including anasazi bean, black beans, cranberry bean, borlotti beans, pink beans, pinto beans, kidney beans, shell beans, white beans, yellow beans and French beans, etc. Lectins or hemagglutinins have been purified from different varieties of *P. vulgaris*. The lectin contents are low in some varieties and high in other varieties (Table 2).

### Purification of lectins or hemagglutinins

Isolation of lectins generally begins with a saline (or buffer) extraction of the finely ground seed meal. Pre – extraction with acids (e.g., acetic acid used by Naeem *et al.* 2007), organic solvents (for example, methanol, diethyl ether or acetone used by Medeiros *et al.* 2010) is often employed to remove lipid or other interfering substances (Sumner and Howell, 1936).

Ammonium sulfate or alcohol fractionation, centrifugation and dissolution of the precipitate yield supernatant liquor containing the lectin(s). Plant lectins or agglutinins may be isolated from saline extracts by conventional protein purification techniques, affinity chromatography or a combination thereof. Virtually all the lectin purification schemes employ affinity chromatography that exploits the specific sugar binding capacity of the lectin (Sharon *et al.*, 1974). Knowledge of the sugar specificity of a lectin which can be obtained from inhibition experiments using simple sugars and crude lectin preparations, permits the design of a suitable purification procedure. Simply stated, a carbohydrate ligand with which the lectin interacts, is insolubilized, the lectin is adsorbed as the extract is percolated slowly over the adsorbent, and displacement of bound lectin is accomplished by elution, either with a sugar that competes for lectin sites with the specific adsorbent or by altering the nature of the eluent (by lowering the pH, increasing the ionic strength, or adding denaturants). The lectin from *Datura stramonium* by affinity chromatography on Sepharose fetuin (Kilpatrick and Yeoman, 1978). Datta and Basu (1983) purified a human erythrocyte specific lectin from the seeds of *Erythrina variegata* Linn. var. *orientalis* Linn. (Leguminosae) by affinity chromatography on acid treated sepharose 4B. An alternative approach involves insolubilized glycoproteins. Felsted *et al.*, 1975 isolated a lectin from saline extracts of red kidney beans (*Phaseolus vulgaris*) by affinity absorption on porcine thyroglobulin-Sepharose. Matsuda *et al.*, 1989 isolated a lectin from winged bean (*Psophocarpus tetragonolobus*) seed extracts by affinity chromatography on Sepharose-6-aminocaproyl-D-galactosamine.

Thus, an affinity system for the isolation of the lectin of red kidney beans (*Phaseolus vulgaris*) involved thyroglobulin – Sepharose (Felsted *et al.*, 1975) and for the isolation of *Limulus polyphemus* lectin, bovine sub maxillary mucin - Sepharose. Fetuin - Sepharose has been employed for the isolation of the agglutinins from wheat germ, jack bean, potato, *Amaranthus hypochondriacus* seeds and several other sources (Owens and Northcote, 1980; Ozeki *et al.*, 1996). The lectin from *Datura stramonium* was isolated by affinity chromatography on Sepharose fetuin (Kilpatrick and Yeoman, 1978).

**Table 1:** Yields of animal lectins obtained by chromatographic isolation from natural sources.

| Natural source   | Chromatography for purification                                     | Lectin yield         | Reference                  |
|--|---|----------------------|----------------------------|
| <i>Acropora millepora</i> (coral) plasma fluid         | Mannose affinity chromatography                                     | 0.7 mg/100 ml plasma | Kvennefors et al. 2008     |
| <i>Aristichthys nobilis</i> (bighead carp) gills       | DEAE-Sepharose, Sephacryl S-200 and Superdex 200                    | 9.4 mg/100 g         | Pan et al. 2010            |
| <i>Bubalus bubalis</i> (Buffalo) heart tissue          | Ammonium sulfate precipitation and Sephadex G50                     | 0.97 mg/100 g        | Ashraf et al. 2010         |
| <i>Holothuria scabra</i> (sea cucumber) coelomic fluid | Ultrafiltration and Phenyl-Sepharose                                | 1.6 mg/100 ml        | Gowda et al. 2008          |
| <i>Macoma birmanica</i> (marine bivalve) foot muscles  | Ammonium sulfate precipitation and N-acetylglucosamine Sepharose 4B | 4.5 mg/100 g         | Adhya et al. 2009          |
| <i>Nemopilema nomurai</i> (jellyfish)                  | SP-Sepharose and BSM-Toyopearl                                      | 0.35 µg/100 g        | Imamichi and Yokoyama 2010 |

**Table 2:** Yields of plant lectins obtained by chromatographic isolation from seeds of different Phaseolus cultivars.

| Phaseolus cultivar           | Chromatography for purification   | Yield(mg/100 g seed)     | Sugar specificity                     | Reference                       |
|------------------------------|---|--------------------------|---------------------------------------|---------------------------------|
| Anasazi bean                 | Affi-gel blue gel, Mono S and Superdex 200  | 13                       | Not found                             | Sharma et al. 2009              |
| Dark red kidney bean         | DEAE-cellulose and Affi-gel blue gel  | 107                      | Not found                             | Xia and Ng 2006                 |
| Escumite bean                | Affinity chromatography (glutaraldehyzed membranes from blood group O erythrocytes) | 163(total of 4 isoforms) | N-acetylglucosamine type glycans      | Castillo Villanueva et al. 2007 |
| Extralong autumn purple bean | Blue-Sepharose, Q-Sepharose, Mono Q and Superdex 75                                 | 35                       | Galactose                             | Fang et al. 2010                |
| French bean 12               | SP-Sepharose, Affi-gel blue, Q-Sepharose, and Superdex 200                          | 4.8                      | Not found                             | Leung et al. 2008               |
| French bean                  | Blue-Sepharose, Q-Sepharose and Superdex 75   | 1100                     | Not found                             | Lam and Ng 2010b                |
| Red kidney bean              | Affi-gel blue gel and CM-Sepharose  | 27.5                     | Lactoferrin, ovalbumin, thyroglobulin | Ye et al. 2001                  |

**Table 3:** Anti-insect activity of lectins.

| Natural source of lectin   | Insect affected                 | Anti-insect effect   | Sugar specificity              | Reference           |
|--|---------------------------------|--|--------------------------------|---------------------|
| <i>Allium sativum</i> (garlic) bulbs                                   | <i>Acyrtosiphon pisum</i>       | Increased mortality  | Mannose                        | Fitches et al. 2008 |
| <i>Arisaema intermedium</i> and <i>Arisaema wallichianum</i> (Araceae) | <i>Bactrocera cucurbitae</i>    | 1. Prolonged period of development<br>2. Inhibited pupation and emergence                                      | Not found                      | Kaur et al., 2009   |
| <i>Gracilaria cornea</i> (red alga)                                    | <i>Boophilus microplus</i>      | Reduced<br>1. body weight of female after oviposition period,<br>2. egg mass weight, and<br>3. hatching period | Fetuin, porcine stomach, mucin | Lima et al. 2005    |
| <i>Gracilaria ormate</i> (red alga)                                    | <i>Callosobruchus maculatus</i> | Delayed development  | Fetuin, porcine stomach, mucin | Leite et al. 2005   |
| <i>Myracrodruon urundeuva</i> (aroeira preta) bark                     | <i>Aedes aegypti</i>            | Increased mortality  | N-acetyl-Dglucosamine          | Sá et al. 2009      |
| <i>Xerocomus chrysenteron</i> fruiting bodies                          | <i>Myzus persicae</i>           | Increased mortality  | Fetuin, porcine stomach mucin  | Jaber et al. 2008   |
| <i>Xerocomus chrysenteron</i> fruiting bodies                          | <i>Myzus persicae</i>           | 1. Increased mortality<br>2. Reduction of body weight, duration of development and fecund                      | Fetuin, porcine stomach mucin  | Jaber et al. 2007   |

**Table 4:** Plant lectins with antimicrobial activity.

| Plant (tissue)                            | Lectin specificity   | Antimicrobial activity   |
|---|----------------------|--|
| <i>Araucaria angustifolia</i> (seed)      | GlcNAc               | <i>Clavibacter michiganensis</i> , <i>Xanthomonas axonopodis</i> pv. <i>Passiflorae</i>  |
| <i>Artocarpus incisa</i> (seed)           | GlcNAc               | <i>Fusarium moniliforme</i> , <i>Saccharomyces cerevisiae</i>  |
| <i>Artocarpus integrifolia</i> (seed)     | GlcNAc               | <i>F. moniliforme</i> , <i>S. cerevisiae</i>   |
| <i>Astragalus mongholicus</i> (root)      | Lactose/D-Gal        | <i>Botrytis cinerea</i> , <i>Fusarium oxysporum</i> , <i>Colletotrichum</i> sp., <i>Drechslera turcia</i>  |
| <i>Eugenia uniflora</i> (seeds)           | Carbohydrate complex | <i>Bacillus subtilis</i> , <i>Corynebacterium bovis</i> , <i>Escherichia coli</i> , <i>Klebsiella</i> sp., <i>Pseudomonas aeruginosa</i> , <i>Streptococcus</i> sp., <i>Staphylococcus aureus</i>  |
| <i>Gastrodia data</i> (corms)             | α-Man/ GlcNAc        | <i>B. cinerea</i> , <i>Ganoderma lucidum</i> , <i>Gibberella zeae</i> , <i>Rhizoctonia solani</i> , <i>Valsa ambiens</i>   |
| <i>Hevea brasiliensis</i> (latex)         | Chitotriose          | <i>B. cinerea</i> , <i>Fusarium culmorum</i> , <i>F. oxysporum</i> f. sp. <i>pisi</i> , <i>Phycomyces blakesleeanus</i> , <i>Pyrenophora trititirepentis</i> , <i>Pyricularia oryzae</i> , <i>Septoria nodorum</i> , <i>Trichoderma hamatum</i>  |
| <i>Myracrodruon urundeuva</i> (heartwood) | GlcNAc               | <i>B. subtilis</i> , <i>Corynebacterium callunae</i> , <i>E. coli</i> , <i>Klebsiella pneumoniae</i> , <i>P. aeruginosa</i> , <i>S. aureus</i> , <i>Streptococcus faecalis</i> , <i>Fusarium solani</i> , <i>F. oxysporum</i> , <i>F. moniliforme</i> , <i>Fusarium decemcellulare</i> , <i>Fusarium lateritium</i> , <i>Fusarium fusarioides</i> , <i>Fusarium verticilloides</i> |
| <i>Ophiopogon japonicus</i> (rhizome)     | Man                  | <i>Gibberella saubinetii</i> , <i>R. solani</i>  |
| <i>Opuntia ficus indica</i> (cladodes)    | Glc/Man              | <i>Colletotrichum gloesporioides</i> , <i>Candida albicans</i> , <i>F. oxysporum</i> , <i>F. solani</i>  |
| <i>Phaseolus coccineus</i> (seeds)        | Sialic acid          | <i>Helminthosporium maydis</i> , <i>Gibberella sanbinetti</i> , <i>R. solani</i> , <i>Sclerotinia sclerotiorum</i>   |
| <i>Phthirusa pyrifolia</i> (leaf)         | Fru-1,6-P2           | <i>B. subtilis</i> , <i>K. pneumoniae</i> , <i>Staphylococcus epidermidis</i> , <i>S. faecalis</i> , <i>F. lateritium</i> , <i>R. solani</i>   |
| <i>Pisum sativum</i> (seed)               | Man                  | <i>Aspergillus flavus</i> , <i>F. oxysporum</i> , <i>Trichoderma viride</i>  |
| <i>Sebastiania jacobinensis</i> (bark)    | Carbohydrate complex | <i>F. moniliforme</i> , <i>F. oxysporum</i>  |
| <i>Talisia esculenta</i> (seeds)          | Man                  | <i>Colletotrichum lindemuthianum</i> , <i>F. oxysporum</i> , <i>S. cerevisiae</i>  |
| <i>Triticum vulgare</i> (seeds)           | GlcNAc               | <i>Fusarium graminearum</i> , <i>F. oxysporum</i>  |
| <i>Urtica dioica</i> (rhizome)            | GlcNAc               | <i>B. cinerea</i> , <i>C. lindemuthianum</i> , <i>Phoma betae</i> , <i>Phycomyces blakesleeanus</i> , <i>Septoria nodorum</i> , <i>Trichoderma hamatum</i> , <i>T. viride</i>  |

Wheat germ agglutinin was also isolated by using insolubilized ovomucoid (Avrameas and Guilbert, 1971). Erythrocytes treated with formaldehyde and glutaraldehyde, have been used as adsorbents for lectin isolation (Avrameas and Guilbert, 1971). Whenever possible commercially available adsorbents are also employed. An increase in the number of purification steps usually results in a lower recovery. In order to produce a large quantity of lectins, the first criterion is a high lectin content in the starting material. The second criterion is the use of a simple purification protocol. Tetrameric escumite lectin was purified by affinity chromatography on a column containing glutaraldehyde membranes from blood group O erythrocytes. Four isoforms were separated on Mono-S (cation exchanger) (Castillo-Villanueva *et al.*, 2007). Dark red kidney bean hemagglutinin was unadsorbed on DEAE-cellulose but adsorbed on Affi-gel blue gel (Xia and Ng, 2006). French bean 35 hemagglutinin with high purity was isolated by chromatography on Blue-Sepharose and Q-Sepharose (Lam and Ng, 2010b).

## APPLICATIONS

Lectins have become the focus of intense interest for biologists and in particular for the research and applications in agriculture and medicine (Movafagh *et al.*, 2013). These proteins with unique characteristics have found use in diverse fields of biology and as more lectins are being isolated and their role in nature elucidated, they continue to occupy an important place in agricultural and therapeutic areas of research.

### Anti-insect activity of lectins

One of the interesting roles of lectins is in host defence against pathogens and predators (Fitches *et al.*, 2010; Hakim *et al.*, 2010; Kaur *et al.*, 2009, 2006a, b). As there is a need to replace conventional insect control measures which cause pollution and disturb the food chain, several alternative measures have been attempted including use of plant lectins. The anti-insect activity of plant lectins against a wide array of insect species have been well documented and represents a potential of using plant lectins as naturally occurring insecticidal agents against pests, which restrain increased crop production (Fitches *et al.*, 2010; Hogervorst *et al.*, 2006). *Bactrocera cucurbitae* is a major pest of cucurbitaceous vegetables and fruits in many parts of the world (Kumar *et al.*, 2006). The pest has so far defied almost all conventional control measures and the damage caused to the standing crop has been reported to be 100% in some cases (Singh *et al.*, 2009).

Lectins have been suggested as one of the promising agents against insect pests and have been engineered successfully into a variety of crops including wheat, rice, tobacco, and potatoes. This approach could be used as a part of integrated pest management strategies and caveat pest attack. In general, it seems that large-scale implementation of transgenic insecticidal and herbicide-tolerant plants does not display considerable negative effects on the environment. Moreover, at least some transgenic plants can improve the corresponding environments and human

health because their production considerably reduces the load of chemical insecticides and herbicides (Velkov *et al.*, 2005). Lectins demonstrate anti-insect activity. They increase the mortality or delay the development of insect (Table 3). When incorporated in an artificial diet, *Arisaema jacquemontii* lectin adversely affected the development of *Bactrocera cucurbitae* larvae (Kaur *et al.*, 2006a). *Arisaema helleborifolium* lectin exhibited anti-insect activity towards the second instar larvae of *B. cucurbitae* (Kaur *et al.*, 2006b). The insecticidal property of lectins may be due to orchestration of enzymatic activity of larvae. After treatment with different lectins, the activity of esterases in larvae was increased whereas the activity of acid phosphatase and alkaline phosphatase decreased. Galectin-1 treatment of *Plutella xylostella* larvae brought about disruption of the microvilli and induced abnormalities in these epithelial cells (Chen *et al.* 2009b). *Dioscorea batatas* lectin inhibited the emergence of *Helicoverpa armigera* larvae into adults by avidly binding to larval brush border and peritrophic membrane (Ohizumi *et al.*, 2009). *Arum maculatum* tuber lectin caused *Lipaphis erysimi* and *Aphis craccivora* to succumb, by binding to the gut brush border membrane vesicle proteins (Majumder *et al.*, 2005). *Olneya tesota* lectin bound to midgut glycoconjugates and microvillae of *Zabrotes subfasciatus* larvae. Diminished oviposition and a failure of emergence of adult beetles were observed (Lagarda-Diaz *et al.*, 2009). *Annona coriacea* lectin displayed toxicity in *Anagasta kuehniella* which apparently resulted from a change in the gut membrane environment and consequent disruption of digestive enzyme recycling mechanisms by binding to midgut proteins (Coelho *et al.*, 2007). *Bauhinia monandra* leaf lectin produced mortality in *Zabrotes subfasciatus* and *Callosobruchus maculatus* when incorporated into an artificial diet. *B. monandra* leaf lectin produced a 40% decrement in weight of *A. kuehniella* larvae. *B. monandra* leaf lectin bound to midgut proteins of the insect *C. maculatus* (Macedo *et al.*, 2007). The detached leaves from transgenic tobacco plants expressing *Allium sativum* lectins reduced the weight gain and development and the metamorphosis of *Spodoptera littoralis* larvae. Furthermore, the larvae were detrimental to the pupal stage resulting in weight reduction and lethal abnormalities (Sadeghi *et al.*, 2008). Production of *Rhopalosiphum maidis* nymphs was significantly reduced on *Galanthus nivalis* agglutinin-expressing plants (Wang *et al.*, 2005). *G. nivalis* agglutinin was also found bound to glycoproteins that can be found in the guts of larvae of *Adalia bipunctata*, *Chrysoperla carnea*, and *Coccinella septempunctata* (Hogervorst *et al.*, 2006). A lectin from *Colocasia esculenta* (L.) Schott corms was shown to have anti-insect potential towards *Bactrocera cucurbitae* (Coquilett) (Thakur *et al.*, 2012) The lectin was found to be specific towards N-acetyl-D-lactosamine (LacNAc), a disaccharide and asialofetuin, a desialylated serum glycoprotein. The lectin significantly decreased the percent pupation and emergence with respect to control. Effect on various enzymes was studied by employing LC50 (51.6 µg ml<sup>-1</sup>) CEA in the artificial diet bioassay of second instar larvae. All the enzymes tested namely esterases, phosphatases (acid and alkaline), superoxide

dismutases, catalase and glutathione-S-transferase showed a significant ( $p < 0.01$ ,  $p < 0.05$ ) increase in their enzyme and specific activities. These results showed that CEA affected normal growth and development and presented stress to the larvae, activating their detoxification and anti-oxidant systems. Thus, the lectin seems to be a useful candidate for the control measures of *Bactrocera cucurbitae*. The lectin gene presents a useful candidate for the integrated pest management. The value of this candidate gene is weighed by the fact that it expresses an edible protein and hence is not expected to pose any serious health threats on human health, if expressed in a transgenic plant.

### Antimicrobial Activity

Many human pathogens utilize cell surface glycans as either receptors or ligands to initiate adhesion and infection (Sharon and Lis, 1989; Sharon and Lis, 2003; Zem et al., 2006; Hyun et al., 2007; Oppenheimer et al., 2008; Magalhaes et al., 2009; Mukhopadhyay et al., 2009). *Escherichia coli* (*E. coli*), for example, binds to host mannosides, while influenza virus binds to host sialic acids (Mukhopadhyay et al., 2009). Other strains of *E. coli* have been discovered that demonstrate specificities towards other host cell surface carbohydrate moieties such as galabiose (Gal- $\alpha$ -4-Gal) and NeuAc- $\alpha$ -2,3-Gal- $\beta$ -3-GalNAc (Khan et al., 2000; Buts et al., 2003). The genital pathogen *Neisseria gonorrhoea* specifically binds N-acetylglucosamine (Gal- $\beta$ -4-GlcNAc, LacNAc), and *Streptococcus pneumoniae* specifically binds the pentasaccharide NeuAc- $\alpha$ -3-Gal- $\beta$ -4-GlcNAc- $\beta$ -3-Gal- $\beta$ -4-Glc as well as the internal tetra- and trisaccharides Gal- $\beta$ -4-GlcNAc- $\beta$ -3-Gal- $\beta$ -4-Glc and GlcNAc- $\beta$ -3-Gal- $\beta$ -4-Glc respectively. *Pseudomonas aeruginosa* specifically binds fucose (L-Fuc) (Barthelson et al., 1998). Bacteria can discriminate between two identical glycans that differ in only one hydroxyl group (Sharon, 2006). Such host-pathogen interactions are multivalent, and therefore the binding events are of high affinity and suited for host invasion (Nimrichter et al., 2004; Mukhopadhyay et al., 2009).

Cytotoxic effects of lectins may be revealed by antitumoral and antiviral activities and also by deleterious effect on microorganisms (Table 4); lectins of different carbohydrate specificities are able to promote growth inhibition or death of fungi and bacteria. Table 4 shows proposed applications of lectins for detection, typing, and control of bacteria and fungi that cause damage to plants and humans. Antibacterial activity on Gram-positive and Gram-negative bacteria occurs through the interaction of lectin with components of the bacterial cell wall including teichoic and teichuronic acids, peptidoglycans and lipopolysaccharides; study revealed that the isolectin I from *Lathyrus ochrus* seeds bind to muramic acid and muramyl dipeptide through hydrogen bonds between ring hydroxyl oxygen atoms of sugar and carbohydrate binding site of lectin and hydrophobic interactions with the side chains of residues Tyr100 and Trp128 of isolectin I (Bourne et al., 1994). Despite the large numbers of lectins and hemagglutinins that have been purified, only a few of them manifested antifungal activity. The inhibition

of fungi growth can occur through lectin binding to hyphae resulting in poor absorption of nutrients as well as by interference on spore germination process (Lis and Sharon, 1981). The polysaccharide chitin is constituent of fungi cell wall and chitin-binding lectins showed antifungal activity; impairment of synthesis and/or deposition of chitin in cell wall may be the reasons of antifungal action (Selitrennikoff, 2001). Probably the carbohydrate-binding property of lectin is involved in the antifungal mechanisms and lectins of different specificities can promote distinct effects. Plant agglutinins are believed to play a role in plant defense mechanism against microorganism phytopathogens (Sa et al., 2009a).

The expression of *Gastrodia elata* lectins in the vascular cells of roots and stems was strongly induced by the fungus *Trichoderma viride*, indicating that lectin is an important defense protein in plants (Sá et al., 2009b). Following insertion of the precursor gene of stinging nettle isolectin I into tobacco, the germination of spores of *Botrytis cinerea*, *Colletotrichum lindemuthianum*, and *T. viride* was significantly reduced (Does et al., 1999). Thus, lectins may be introduced into plants to protect them from fungal attack. Plant lectins can neither bind to glycoconjugates on the fungal membranes nor penetrate the cytoplasm owing to the cell wall barrier. It is not likely lectins directly inhibit fungal growth by modifying fungal membrane structure and/or permeability. However, there may be indirect effects produced by the binding of lectins to carbohydrates on the fungal cell wall surface. Chitinase-free chitin-binding stinging nettle (*Urtica dioica* lectin) impeded fungal growth. Cell wall synthesis was interrupted because of attenuated chitin synthesis and/or deposition (Van Parijs et al., 1991). The effects of nettle lectin on fungal cell wall and hyphal morphology suggest that the nettle lectin regulates endomycorrhizal colonization of the rhizomes. Several other plant lectins inhibit fungal growth. The first group includes small chitin-binding merolectins with one chitin-binding domain, e.g., hevein from rubber tree latex (Van Parijs et al., 1991) and chitin-binding polypeptide from *Amaranthus caudatus* seeds (Broekaert et al., 1992). The only plant lectins that can be considered as fungicidal proteins are the chimerolectins belonging to the class I chitinases. However, the antifungal activity of these proteins is ascribed to their catalytic domain.

### Antitumor activity

Owing to their fine specificity, lectins have various applications in biomedical sciences including cancer research. It is well documented that lectins have an antitumor effect. Plant lectins represent a well-defined and a novel non-traditional source of anticancer compounds. A number of plant lectins (predominantly galactoside and galNAc specific) have been in pre-clinical and clinical trials as potential drugs for treatment of cancer (Ernst et al., 2003). Within the past few years, lectins have become a well-established means for understanding varied aspects of cancer and metastasis. Evidence is now emerging that lectins are dynamic contributors to tumor cell recognition (surface markers), cell

adhesion and localization, signal transduction across membranes, mitogenic stimulation, augmentation of host immune defense, cytotoxicity, and apoptosis. To advance understanding of these lectin-dependent processes, attempts are being made to discover new lectins that have one or more of these functions and to develop lectin- (or glycoconjugate-) based tools that could be used to home in on tumor cells. Legume lectins are one of the most extensively studied plant lectin families for their molecular basis of the protein-carbohydrate interactions for several decades (Damodaran *et al.*, 2008). In recent years, the main interests in this lectin family lay in their potential application as anti-tumour agents that could bind specific cancer cell surface glycoconjugates. Concanavalin A (ConA), a typical legume lectin with a mannose/glucose-binding specificity, was reported to induce apoptosis in murine macrophage PU5-1.8 cells through clustering of mitochondria and release of cytochrome *c*. Recent study has showed that ConA induces apoptosis in human melanoma A375 cells in a caspase-dependent pathway. Subsequently, ConA caused mitochondrial transmembrane potential (MMP) collapse, cytochrome *c* release, activation of caspases and eventually triggering a mitochondria-mediated apoptosis (Liu *et al.*, 2009).

Furthermore, other recent reports have demonstrated that a legume lectin named *S. flavescens* lectin (SFL) can induce tumour cell death through a caspase-dependent apoptotic pathway, and its apoptotic mechanisms is speculated to be the death-receptor pathway (Liu *et al.*, 2008). And, another typical legume lectin with specificity towards sialic acid purified from *Phaseolus coccineus* L. (*Phaseolus. multiflorus* wild) seeds possesses a remarkable anti-proliferative activity. This lectin induced the caspase dependent apoptosis in murine fibrosarcoma L929 cells. Besides, its antineoplastic activity was decreased abruptly when the sialic acid-specific activity was completely inhibited, which indicates that this sugar-binding specificity might be the main reason sparking off the antineoplastic activity and apoptosis (Chen *et al.*, 2009). *Flammulina velutipes* hemagglutinin-inhibited proliferation of leukemia L1210 cells (Ng *et al.*, 2006). *Haliclona crater* lectin displayed a cytotoxic effect on HeLa and FemX cells (Pajic *et al.*, 2002). Dark red kidney bean hemagglutinin exerted an antiproliferative activity toward leukemia L1210 cells (Xia and Ng, 2006). Small glossy black soybean (*Glycine max*) lectin impeded proliferation of breast cancer MCF7 cells and hepatoma HepG2 cells (Lin *et al.*, 2008). Del Monte banana lectin retarded proliferation of (L1210) cells and hepatoma (HepG2) cells (Cheung *et al.*, 2009). Extra long autumn purple bean lectin inhibited the proliferation of hepatoma HepG2 cells by inducing the production of apoptotic bodies (Fang *et al.* 2010). Mistletoe lectin can be used in cancer patients to improve the quality of life (Semiglazov *et al.*, 2006). In order to widen the application of anti-tumor lectins, the mechanism of action was elucidated. Lectins elicit apoptosis in different cancer cell lines. Examples include Korean mistletoe lectin-treated B16-BL6 melanoma cells (Park *et al.*, 2001), Korean mistletoe lectin-treated human A253 cancer cells (Choi *et al.*, 2004), *Agrocybe aegerita* lectin-treated HeLa

cells (Zhao *et al.*, 2009), Abrus agglutinin-treated Dalton's lymphoma cells (Bhutia *et al.* 2008a) and HeLa cells (Bhutia *et al.*, 2008b), *Sophora flavescens* lectin-treated HeLa cells (Liu *et al.*, 2008), *Polygonatum odoratum* lectin treated murine fibrosarcoma L929 cells (Liu *et al.*, 2009b), *Polygonatum cyrtonema* lectin-treated human melanoma A375 cells (Liu *et al.*, 2009a), *Pseudomonas aeruginosa* hemagglutinin-treated breast cancer cells (MDA-MB-468, and MDA-MB-231HM cells; Liu *et al.*, 2009c), French bean hemagglutinin-treated breast cancer MCF-7 cells (Lam and Ng ,2010a), and recombinant protease-resistant galectin-9- treated myeloma cells (Kobayashi *et al.*, 2010). Although the apoptotic pathways look different, activation of different caspases is usually involved. Caspase-3 plays a central role in apoptosis. It interacts with caspase-8 and caspase-9. Therefore, caspase-3 is usually investigated in apoptotic pathways, except in the case of a caspase-3- deficient cell line (e.g., MCF-7 cells) which was used in the study of French bean hemagglutinin (Lam and Ng, 2010a). Caspase-8 and -9 are also activated (Liu *et al.*, 2009a; Liu *et al.*, 2009b, c; Kobayashi *et al.*, 2010; Lam and Ng, 2010a). Apoptosis can be mediated by death receptors initiated by lectins. FAS receptor is the receptor with which lectins often interact (Liu *et al.*, 2009b, c; Lam and Ng, 2010a). The interaction is probably by protein-protein interaction. The Bcl family members (anti-apoptotic factors) were down-regulated (Bhutia *et al.*, 2008a, b; Liu *et al.* 2009a; Lam and Ng, 2010a). The sequestration of cytochrome *c* in mitochondria was interrupted and cytochrome *c* release was observed (Bhutia *et al.* 2008a, b; Liu *et al.*, 2009a, b; Lam and Ng, 2010a). Finally, mitochondrial membrane depolarization was detected (Liu *et al.*, 2009a, b; Lam and Ng, 2010a) and G<sub>0</sub>/G<sub>1</sub> arrest was frequently observed (Bhutia *et al.* 2008a, b; Liu *et al.* 2009c; Lam and Ng 2010a) which seems to be the characteristic of lectin-induced apoptosis although sub G<sub>1</sub> arrest (Park *et al.*, 2001) and G<sub>2</sub>/M arrest (Lam and Ng, 2010a) were found in some cases. Investigations of the anti-tumor effect of lectin in vivo have been reported. *Pleurotus citrinopileatus* lectin (Li *et al.* 2008) and *R. lepida* lectin (Zhang *et al.*, 2010) exerted potent antitumor activity in white Kunming mice bearing sarcoma 180, and caused inhibition of tumor growth when administered intraperitoneally. Lectins have received a lot of attention from cancer biologists due to their remarkable anti-tumor properties. Lectins ConA, ConBr, and CFL are all structurally related and induce apoptosis in the MCF-7 cell line. They have been shown to reduce both proliferation and viability of leukemic cells ConA and ConBr lectins have cytotoxic effects in leukemic cells. Lectins (Con A and Con Br) have been shown to induce internucleosomal DNA fragmentation and alter mitochondrial transmembrane potential in leukemic cells. ConA and ConBr induce apoptosis in leukemic cells by triggering an intrinsic mitochondrial pathway and also increase ROS (Reactive Oxygen Species) (Faheina *et al.*, 2011). The aqueous extract of European mistletoe (*Viscum album*, L.) has been applied in cancer therapy (Lyu *et al.*, 2004). However, in order to make lectin useful practically in the clinical setting, a delivery system is required to lower toxicity, extend exposition, and improve efficacy. Wheat

germ agglutinin and *Ulex europaeus* agglutinin displayed strong interaction with human urinary carcinoma 5,637 cells, which enabled them to target to bladder cancer cells (Plattner *et al.*, 2008). The encapsulation of *Cratylia mollis* lectin with liposomes lowered its tissue toxicity in the liver and kidney, and improved its antitumor activity in Swiss mice inoculated with sarcoma 180 (Andrade *et al.*, 2004). Mistletoe lectin was stabilized with alginate/chitosan microcapsules coated by a biodegradable polymer wall which can be used to protect the lectin from acidic pH in the stomach (Lyu *et al.*, 2004). Immunofluorescence and/or immunohistochemical studies using lectins can reveal the early premalignant stage of prostate carcinogenesis. Expression of glycoconjugates is often altered in tumor cells. Abundant N-acetylglucosamine ( $\alpha$ 1,3) Nacetylglucosamine/ galactose and galactose ( $\beta$ 1,4) Nacetylglucosamine ( $\alpha$ ,2) mannose ( $\alpha$ 1,6) residues were observed in dysplastic epithelium tumor cells as evidenced by labeling by the N-acetylgalactosamine-specific and complex type oligosaccharide-specific lectins. The binding of these lectins to androgen-independent rat prostatic carcinoma was revealed, indicating that these sugar residues are common in some dysplastic and neoplastic prostatic cells (Chan *et al.*, 2001).

Generally speaking, the above-mentioned discoveries of the lectins suggest that they might possess some similar biological activities and anti-tumour mechanisms that are closely correlated with their corresponding molecular structures. Thus, these results would provide new clues for further exploring the anti-tumour mechanisms of the lectins.

### Antiviral Activity

A lectin (D-mannose-specific) from *Gerardia savaglia* was for the first time reported to prevent infection of H9 cells with human immunodeficiency virus (HIV)-1. Furthermore, the lectin inhibited syncytium formation in the HTLV-III<sub>B</sub>/H9-Jurkat cell system and HIV-1/human lymphocyte system by reacting with the oligosaccharide side chains of the HIV-1 gp120 envelop molecule (high-mannose oligosaccharides; Muller *et al.*, 1988). A year later, the lectins concanavalin A, wheat germ agglutinin, *Lens culinaris* agglutinin, *Vicia faba* agglutinin, *Pisum sativum* agglutinin and phytohaem (erythro) agglutinin were found to bind to gp120. They were able to inhibit fusion of HIV-infected cells with CD4 cells by a carbohydrate-specific interaction with the HIV-infected cells (Hansen *et al.*, 1989). Plant lectins displayed anti-coronavirus activity, especially mannose-binding lectins, in severe acute respiratory syndrome coronavirus. They interfered with viral attachment in early stage of replication cycle and suppressed the growth by interacting at the end of the infectious virus cycle (Keyaerts *et al.*, 2007). Banana (*Musa acuminata*) lectin has been shown to inhibit HIV replication (Swanson *et al.* 2010). The treatment of AIDS with lectins is being investigated in many studies. Different lectins have different anti-HIV mechanisms. More recently, lectin from the polychaete marine worm *Chaetopterus variopedatus* inhibited cytopathic effect induced by HIV-1 and the production of viral p24 antigen (Wang *et al.*, 2005). The sea worm (*Serpula vermicularis*) lectin suppressed the

production of viral p24 antigen and cytopathic effect induced by HIV-1 (Molchanova *et al.*, 2007). *P. cyrtonema* Hua lectin inhibited HIV-I- and HIV-II-induced cytopathicity in MT-4 and CEM cells (An *et al.*, 2006). Banana lectin directly bound the HIV-1 envelope protein (gp120) and blocked entry of the virus into the cell, and decreased the levels of the strong-stop product of early reverse transcription (Swanson *et al.*, 2010). Extra long autumn purple bean lectin (Fang *et al.*, 2010) and mushroom *Russula delica* lectin (Zhao *et al.*, 2009) were able to inhibit HIV-1 reverse transcriptase. Hence, lectins are potential drugs for treatment of AIDS. Besides the aforementioned practical applications of lectins, there have been isolated reports of the antibacterial (Ngai and Ng, 2007) and anti-nematode (Wang *et al.*, 2005) activities of lectin.

### CONCLUSION

Lectins are a subject with immense potential and of intense investigations. As more lectins are isolated and further studies are conducted on the biological activities and mechanisms of action of lectins, the production of lectins can be improved and new applications of lectins can be found and explored for significant contributions in various fields of biology.

### REFERENCES

- Adhya M., Singha B., Chatterjee., B. Purification and characterization of an N-acetylglucosamine specific lectin from marine bivalve *Macoma birmanica*. *Fish Shellfish Immunol.* 2009; 27:1–8.
- Agrawal BBL., Goldstein IJ. Protein carbohydrate interaction. VI. Isolation of concanavalin A by specific adsorption on cross-linked dextran gels. *Biochim Biophys Acta.* 1967; 147: 262–271.
- An J., Liu JZ., Wu CF., Li J, Dai L, Van Damme E, Balzarini J, De Clercq E, Chen F, Bao JK. Anti-HIV I/II activity and molecular cloning of a novel mannose/sialic acid-binding lectin from rhizome of *Polygonatum cyrtonema* Hua. *Acta Biochim Biophys Sin* (Shanghai) 2006; 38:70–78.
- Andrade CA., Correia MT., Coelho LC., Nascimento SC., Santos-Magalhães NS. Antitumor activity of *Cratylia mollis* lectin encapsulated into liposomes. *Int J Pharm.* 2004; 278:435–445.
- Ashraf GM., Rizvi S., Naqvi S., Suhail N., Bilal N., Hasan S., Tabish M., Banu N. Purification, characterization, structural analysis and protein chemistry of a buffalo heart galectin-1. *Amino Acids.* 2010; 39(5):1321-32.
- Avrameas S., and Guilbert B. Biologically Active Water-Insoluble Protein Polymers. Their Use for the Isolation of Specifically Interacting Proteins, *Biochimie.* 1971. 53: 603.
- Barthelson R., Mobasser A., Zopf D., Simon P. Adherence of *Streptococcus pneumoniae* to respiratory epithelial cells is inhibited by sialylated oligosaccharides. *Infect Immun.* 1998; 66: 1439–1444.
- Bies C., Lehr CM., Woodley JF. Lectin-mediated drug targeting: history and applications. *Adv Drug Deliv Rev.* 2004; 56: 425–435.
- Bhat GG., Shetty KN., Nagre NN., Neekhra VV., Lingaraju S., Bhat RS., Inamdar SR., Suguna K., Swamy BM. Purification, characterization and molecular cloning of a monocot mannose binding lectin from *Remusatia vivipara* with nematocidal activity. *Glycoconj J.* 2010; 27:309–320.
- Bhutia SK., Mallick SK., Maiti S., Maiti TK. Antitumor and proapoptotic effect of Abrus agglutinin derived peptide in Dalton's lymphoma tumor model. *Chem Biol Interact.* 2008; 174:11–18.
- Bhutia SK., Mallick SK., Stevens SM., Prokai L., Vishwanatha JK. Maiti TK. Induction of mitochondria-dependent apoptosis by Abrus

- agglutinin derived peptides in human cervical cancer cell. *Toxicol In Vitro*. 2008; 22:344–351.
- Bourne Y., Ayoub A., Rouge P., Cambillau C. Interaction of a legume lectin with two components of the bacterial cell wall. *The Journal of Biological Chemistry*. 1994; 269: 9429-9435.
- Boyd WC., Shapleigh E. Specific precipitation activity of plant agglutinins (lectins). *Science*. 1954; 119, 419.
- Broekaert WF., Mariën W., Terras FR., De Bolle MF., Proost P., Van Damme J., Dillen L., Claeys M., Rees SB., Vanderleyden J *et al.* Antimicrobial peptides from *Amaranthus caudatus* seeds with sequence homology to the cysteine/glycine-rich domain of chitin-binding proteins. *Biochemistry*. 1992; 31:4308–4314.
- Broekaert WF., van Parijs J., Leyns F., Joos H., Peumans WJ. A chitin-binding lectin from stinging nettle rhizomes with antifungal properties. *Science*. 1989; 245:1100-1102.
- Buts, L., Bouckaert, J., De Genst, E., Loris, R., Oscarson, S., and Lahmann, M. *et al.* The fimbrial adhesion F17-G of enterotoxigenic *Escherichia coli* has an immunoglobulin-like lectin domain that binds N-acetyl-glucosamine. *Mol Microbiol*. 2003; 49:705–715.
- Castillo-Villanueva A., Caballero-Ortega H., Abdullaev-Jafarova F., Garfias Y., del Carmen Jiménez-Martínez M., Bouquelet S., Martínez G., Mendoza-Hernández G., Zenteno E. Lectin from *Phaseolus acutifolius* var. *escumite*: chemical characterization, sugar specificity, and effect on human T-lymphocytes. *J Agric Food Chem*. 2007; 55:5781–5787.
- Chan FL., Choi HL., Ho SM. Analysis of glycoconjugate patterns of normal and hormone-induced dysplastic Noble rat prostates, and an androgen-independent Noble rat prostate tumor, by lectin histochemistry and protein blotting. *Prostate*. 2001; 46:21–32.
- Chen JB. Liu N., Ji, J., Zhou HJ., Bian CY., Li F., Chen and Bao JK. A novel sialic acid-specific lectin from *Phaseolus coccineus* seeds with potent antineoplastic and antifungal activities. *Phytomedicine*. 2009; 16: 352–360.
- Chen SJ., Chen NT., Wang SH., Hsu JC., Ding WH., Kuo-Huang LL., Huang RN. Insecticidal action of mammalian galectin-1 against diamond back moth (*Plutella xylostella*). *Pest Manag Sci*. 2009; 65:923–930.
- Choi SH., Lyu SY., Park WB. Mistletoe lectin induces apoptosis and telomerase inhibition in human A253 cancer cells through dephosphorylation of Akt. *Arch Pharm Res*. 2004; 27:68–76.
- Ciopraga J., Gozia O., Tudor R., Brezua L., Doyle RJ. *Fusarium* sp. Growth inhibition by wheat germ agglutinin. *Biochimica et Biophysica Acta*. 1999; 1428:424-432.
- Coelho MB., Marangoni S., Macedo ML. Insecticidal action of *Annona coriacea* lectin against the flour moth *Anagasta kuehniella* and the rice moth *Corcyra cephalonica* (Lepidoptera: Pyralidae). *Comp Biochem Physiol C Toxicol Pharmacol*. 2007; 146:406–414.
- Costa RMPB., Vaz AFM., Oliva MLV., Coelho LCBB., Correia MTS., Carneiro-da-Cunha MG. A new mistletoe *Phthirusa pyrifolia* leaf lectin with antimicrobial properties. *Process Biochemistry*. 2010; 45:526-533.
- Damodaran D., Jeyakani J., Chauhan A., Kumar N., Chandra NR., Surolia A. Cancer Lectin DB: a database of lectins relevant to cancer. *Glycoconjugate J* 2008; 25: 191–198.
- Datta TK., Basu PS. Human erythrocyte specific lectin from the seeds of Indian coral tree, *Erythrina variegata* Linn, var. *Orientalis* Linn, Merrill. *J Biosci*. 1983; 5: 25-30.
- Does MP., Houterman PM., Dekker HL., Cornelissen BJ. Processing, targeting, and antifungal activity of stinging nettle agglutinin in transgenic tobacco. *Plant Physiol*. 1999; 120:421–432.
- Doyle RJ., Keller K. Lectins in diagnostic microbiology. *Eur. J. Clin. Microbiol*. 1984; 3: 4-9.
- Ernst E., Schmidt K., Steuer-Vogt MK. Mistletoe for cancer? A systematic review of randomised clinical trials. *International Journal of Cancer*. 2003; 107(2):262-7.
- Faheina-Martins GV., da Silveira AL., Ramos MV., Marques-Santos LF., Araujo DA. Influence of fetal bovine serum on cytotoxic and genotoxic effects of lectins in MCF-7 cells. *J Biochem Mol Toxicol*. 2011;25(5):290-6.
- Fang EF., Lin P., Wong JH., Tsao SW., Ng TB. A lectin with anti- HIV-1 reverse transcriptase, antitumor, and nitric oxide inducing activities from seeds of *Phaseolus vulgaris* cv. extralong autumn purple bean. *J Agric Food Chem*. 2010; 58: 2221–2229.
- Felsted RL., Leavitt RD., Bachur NR. Purification of the phytohemagglutinin family of proteins from red kidney beans *Phaseolus vulgaris* by affinity chromatography. *Biochim. Biophys. Acta*. 1975; 405: 72-72.
- Freire MGM., Gomes VM., Corsini RE., Machado OLT., De Simone SG., Novello JC., Marangoni S., Macedo MLR. Isolation and partial characterization of a novel lectin from *Talisia esculenta* seeds that interferes with fungal growth. *Plant Physiology and Biochemistry*. 2002; 40:61-68.
- Fitches E., Wiles D., Douglas AE., Hinchliffe G., Audsley N., Gatehouse JA. The insecticidal activity of recombinant garlic lectins towards aphids. *Insect Biochem Mol Biol*. 2008; 38:905–915.
- Gabius HJ. Non-carbohydrate binding partners/domains of animal lectins. *Int J Biochem*. 1994; 26:469-477.
- Ghosh M. Purification of a lectin-like antifungal protein from the medicinal herb, *Withania somnifera*. *Fitoterapia*. 2009; 80:91–95.
- Goldstein IJ. Lectin structure-activity: The story is never over. *J Agric Food Chem*. 2002; 50:6583–6585.
- Gowda NM., Goswami U., Khan MI. Purification and characterization of a T-antigen specific lectin from the coelomic fluid of a marine invertebrate, sea cucumber (*Holothuria scabra*). *Fish Shellfish Immunol*. 2008; 24:450–458.
- Hakim RS., Baldwin K., Smaghe G. Regulation of midgut growth, development, and metamorphosis. *Annu. Rev. Entomol*. 2010; 55: 593-608
- Hansen JE., Nielsen CM., Nielsen C., Heegaard P., Mathiesen LR., Nielsen JO. Correlation between carbohydrate structures on the envelope glycoprotein gp120 of HIV-1 and HIV-2 and syncytium inhibition with lectins. *AIDS*. 1989; 3:635–641.
- Hogervorst PA., Ferry N., Gatehouse AM., Wäckers FL., Romeis J. Direct effects of snowdrop lectin (GNA) on larvae of three aphid predators and fate of GNA after ingestion. *J Insect Physiol*. 2006; 52:614–624.
- Hyun S., Kim J., Kwon M., and Yu J. Selection and syntheses of tentacle type peptides as ‘artificial’ lectins against various cell-surface carbohydrates. *Bioorg Med Chem*. 2007; 15: 511–517.
- Imamichi Y., Yokoyama Y. Purification, characterization and cDNA cloning of a novel lectin from the jellyfish *Nemopilema nomurai*. *Comp Biochem Physiol B Biochem Mol Biol*. 2010; 156:12–18.
- Jaber K, Francis F, Paquereau L, Fournier D, Haubruge E Effect of a fungal lectin from *Xeroocomus chrysenteron* (XCL) on the biological parameters of aphids. *Commun Agric Appl Biol Sci*. 2007; 72:629–638.
- Jaber K., Cuartero Diaz G., Haubruge E., Francis F. Investigation of carbohydrate binding property of a fungal lectin from *Xeroocomus chrysenteron* and potential use on *Myzus persicae* aphid. *Commun Agric Appl Biol Sci*. 2008; 73:629–638.
- Kaur M., Singh K., Rup PJ., Kamboj SS., Saxena AK., Sharma M., Bhagat M., Sood SK., Singh J. A tuber lectin from *Arisaema jacquemontii* Blume with anti-insect and antiproliferative properties. *J Biochem Mol Biol*. 2006; 39:432–440.
- Kaur M., Singh K., Rup PJ., Saxena AK., Khan RH., Ashraf MT., Kamboj SS., Singh J. A tuber lectin from *Arisaema helleborifolium* Schott with anti-insect activity against melon fruit fly, *Bactrocera cucurbitae* (Coquillett) and anti-cancer effect on human cancer cell lines. *Arch Biochem Biophys*. 2006; 445:156–165.
- Kaur M., Singh K., Rup PJ., Kamboj SS., Singh J. Anti-insect potential of lectins from *Arisaema* species towards *Bactrocera cucurbitae*. *J Environ Biol*. 2009; 30:1019–1023.
- Keyaerts E., Vijgen L., Pannecouque C., Van Damme E., Peumans W., Egberink H., Balzarini J., Van Ranst M. Plant lectins are potent inhibitors of coronaviruses by interfering with two targets in the viral replication cycle. *Antivir Res*. 2007; 75:179–187.
- Kilpatrick DC., Yeoman MM. Purification of the lectin from *Datura stramonium*. *Biochem J*. 1978; 175(3):1151–1153.



- Khan AS., Kniep B., Oelschlaeger TA., Van Die I., Korhonen T., Hacker, J. Receptor structure for FIC fimbriae of uropathogenic *Escherichia coli*. *Infect Immun.* 2000; 68:3541–3547.
- Kobayashi T., Kuroda J., Ashihara E., Oomizu S., Terui Y., Taniyama A., Adachi S., Takagi T., Yamamoto M., Sasaki N., Horiike S., Hatake K., Yamauchi A., Hirashima M., Taniwaki M. Galectin-9 exhibits anti-myeloma activity through JNK and p38 MAP kinase pathways. *Leukemia.* 2010; 24:843–850.
- Kumar NKK., Verghese A., Shivakumara B., Krishnamoorthy PN., Ranganath HR. Relative incidence of *Bactrocera cucurbitae* (Coquillett) and *Dacus ciliates* Loew on cucurbitaceous vegetables. Fruit flies of economic importance: From basic to applied knowledge. Proceedings of the 7<sup>th</sup> International Symposium on Fruit Flies of Economic Importance, Salvador, Brazil. 2006; 249–253.
- Kvennefors EC., Leggat W., Hoegh-Guldberg O., Degnan BM., Barnes AC. An ancient and variable mannose-binding lectin from the coral *Acropora millepora* binds both pathogens and symbionts. *Dev Comp Immunol.* 2008; 32:1582–1592.
- Lagarda-Diaz I., Guzman-Partida AM., Urbano-Hernandez G., Ortega- Nieblas MM., Robles-Burgueño MR., Winzerling J., Vazquez-Moreno L. Insecticidal action of PF2 lectin from *Olneya tesota* (Palo Fierro) against *Zabrotes subfasciatus* larvae and midgut glycoconjugate binding. *J Agric Food Chem.* 2009; 57:689–694.
- Lam SK., Ng TB. Novel galactonic acid-binding hexameric lectin from *Hibiscus mutabilis* seeds with antiproliferative and potent HIV-1 reverse transcriptase inhibitory activities. *Acta Biochim Pol.* 2009; 56:649–654.
- Lam SK., Ng TB. First report of a haemagglutinin-induced apoptotic pathway in breast cancer cells. *Biosci Rep.* 2010; 30:307–317.
- Lam SK., Ng TB. Isolation and characterization of a French bean hemagglutinin with antitumor, antifungal, and anti-HIV-1 reverse transcriptase activities and an exceptionally high yield. *Phytomedicine.* 2010; 17:457–462.
- Lam SK., Han QF., Ng TB. Isolation and characterization of a lectin with potentially exploitable activities from caper (*Capparis spinosa*) seeds. *Biosci Rep.* 2009; 29:293–299.
- Leite YF., Silva LM., Amorim RC., Freire EA., de Melo Jorge DM., Grangeiro TB., Benevides NM. Purification of a lectin from the marine red alga *Gracilaria ornata* and its effect on the development of the cowpea weevil *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Biochim Biophys Acta.* 2005; 1724:137–145.
- Leung EH., Wong JH., Ng TB. Concurrent purification of two defense proteins from French bean seeds: a defensin-like antifungal peptide and a hemagglutinin. *J Pept Sci.* 2008; 14:349–353.
- Li YR., Liu QH., Wang HX., Ng TB. A novel lectin with potent antitumor, mitogenic and HIV-1 reverse transcriptase inhibitory activities from the edible mushroom *Pleurotus citrinopileatus*. *Biochim Biophys Acta.* 2008; 1780:51–57.
- Lima ME., Carneiro ME., Nascimento AE., Grangeiro TB., Holanda ML., Amorim RC., Benevides NM. Purification of a lectin from the marine red alga *Gracilaria cornea* and its effects on the cattle tick *Boophilus microplus* (Acari: Ixodidae). *J Agric Food Chem.* 2005; 53:6414–6419.
- Lis H and Sharon N. (1981). Lectins in higher plants. In: A. Marcus, (Ed.) *The Biochemistry of Plants* vol. 6. (pp. 371–447). New York, NY: Academic Press.
- Lin P., Ye X., Ng T. Purification of melibiose-binding lectins from two cultivars of Chinese black soybeans. *Acta Biochim Biophys. Sin (Shanghai).* 2008; 40:1029–1038.
- Liu Z., Liu B., Zhang ZT., Zhou TT., Bian HJ., Min MW., Liu YH., Chen J., Bao JK. A mannose-binding lectin from *Sophora flavescens* induces apoptosis in HeLa cells. *Phytomedicine.* 2008; 15:867–875.
- Liu B., Cheng Y., Bian HJ., Bao JK. Molecular mechanisms of *Polygonatum cyrtonema* lectin-induced apoptosis and autophagy in cancer cells. *Autophagy.* 2009; 5:253–255
- Liu B., Zhang B., Min MW., Bian HJ., Chen LF., Liu Q., Bao JK. Induction of apoptosis by *Polygonatum odoratum* lectin and its molecular mechanisms in murine fibrosarcoma L929 cells. *Biochim Biophys Acta.* 2009; 1790:840–844.
- Liu ZB., Hou YF., Di Min-Dong GH., Wu J., Shen ZZ., Shao ZM. PA-MSHA inhibits proliferation and induces apoptosis through the up-regulation and activation of caspases in the human breast cancer cell lines. *J Cell Biochem.* 2009; 108:195–206.
- Liu B., Cheng Y., Zhang B., Bian HJ., Bao JK. *Polygonatum cyrtonema* lectin induces apoptosis and autophagy in human melanoma A375 cells through a mitochondria-mediated ROS-p38-p53 pathway. *Cancer Lett.* 2009; 275: 54–60.
- Lyu SY., Kwon YJ., Joo HJ., Park WB. Preparation of alginate/chitosan microcapsules and enteric coated granules of mistletoe lectin. *Arch Pharm Res.* 2004; 27:118–126.
- Macedo ML., das Graças Machado Freire M., da Silva MB., Coelho LC. Insecticidal action of *Bauhinia monandra* leaf lectin (BmoLL) against *Anagasta kuehniella* (Lepidoptera: Pyralidae), *Zabrotes subfasciatus* and *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Comp Biochem Physiol A Mol Integr Physiol.* 2007; 146:486–498.
- Magalhaes A., Gomes J., Ismail MN., Haslam SM., Mendes N., Osorio H. Fut2-null mice display an altered glycosylation profile and impaired BabA-mediated *Helicobacter pylori* adhesion to gastric mucosa. *Glycobiology.* 2009; 19:1525–36.
- Majumder P., Mondal HA., Das S. Insecticidal activity of *Arum maculatum* tuber lectin and its binding to the glycosylated insect gut receptors. *J Agric Food Chem.* 2005; 53:6725–6729.
- Matsuda T., Kabat EA., Surolia A. Carbohydrate binding specificity of the basic lectin from winged bean (*Psophocarpus tetragonolobus*) *Mol Immunol.* 1989; 26(2):189–95.
- Medeiros DS., Medeiros TL., Ribeiro JK., Monteiro NK., Migliolo L., Uchoa AF., Vasconcelos IM., Oliveira AS., de Sales MP., Santos EA. A lactose specific lectin from the sponge *Cinachyrella apion*: purification, characterization, N-terminal sequences alignment and agglutinating activity on *Leishmania* promastigotes. *Comp Biochem Physiol B Biochem Mol Bio.* 2010; 155:211–216.
- Molchanova V., Chikalovets I., Chernikov O., Belogortseva N., Li W., Wang JH., Yang DY., Zheng YT., Lukyanov P. A new lectin from the sea worm *Serpula vermicularis*: isolation, characterization and anti-HIV activity. *Comp Biochem Physiol C Toxicol Pharmacol.* 2007; 145:184–193.
- Movafagh A., Ghanati K., Amani D., Mahdavi SM., Hashemi M., Davood Zare Abdolahi Z., Darvish H., Gholami M., HaghNejad L., Mosammami S., Safari S., Darehgzani R., Rahimi M., Naini NS., Motlagh MG., Zamani M. The structure Biology and Application of Phytohemagglutinin (PHA) in Phytomedicine: With special up-to-date references to lectins. *Journal of Paramedical Sci.* 2013; 4.
- Mukhopadhyay B., Martins M.B., Karamanska R., Russell DA., Field RA. Bacterial detection using carbohydrate-functionalised CdS quantum dots: a model study exploiting *E. coli* recognition of mannosides. *Tetrahedron Lett.* 2009; 50:886–889.
- Muller WE., Renneisen K., Kreuter MH., Schröder HC., Winkler I. The D-mannose-specific lectin from *Gerardia savaglia* blocks binding of human immunodeficiency virus type I to H9cells and human lymphocytes in vitro. *J Acquir Immune Defic Syndr.* 1988; 1:453–458.
- Naeem A., Haque S., Khan RH. Purification and characterization of a novel beta-D-galactosides-specific lectin from *Clitoria ternatea*. *Protein J.* 2007; 26:403–413.
- Ng TB., Ngai PH., Xia L. An agglutinin with mitogenic and antiproliferative activities from the mushroom *Flammulina velutipes*. *Mycologia.* 2006; 98:167–171.
- Ngai PH., Ng TB. A lectin with antifungal and mitogenic activities from red cluster pepper (*Capsicum frutescens*) seeds. *Appl Microbiol Biotechnol.* 2007; 74:366–371.
- Nimrichter A., Gargir M., Gortler RT., Altstock A., Shtevi Weissshaus O. Intact cell adhesion to glycan microarrays. *Glycobiology.* 2004; 14:197–203.
- Ohizumi Y., Gaidamashvili M., Ohwada S., Matsuda K., Kominami J., Nakamura-Tsuruta S., Hirabayashi J., Naganuma T., Ogawa T., Muramoto K. Mannose-binding lectin from yam (*Dioscorea batatas*) tubers with insecticidal properties against *Helicoverpa armigera* (Lepidoptera: Noctuidae). *J Agric Food Chem.* 2009; 57:2896–2902.

- Oliveira MDL., Andrade CAS., Santos-Magalhaes NS., Coelho LCBB., Teixeira JA., Carneiro-da-Cunha MG., Correia MTS. Purification of a lectin from *Eugenia uniflora* L. seeds and its potential antibacterial activity. *Letters in Applied Microbiology*. 2008; 46:371-376.
- Oppenheimer SB., Alvarez M., Nnoli J. Carbohydrate-based experimental therapeutics for cancer, HIV/AIDS and other diseases. *Acta Histochem*. 2008; 110:6-13.
- Owens RJ., Northcote DH. Purification of potato lectin by affinity chromatography on a fetuin-sepharose matrix. *Phytochemistry*. 1980; 19:1861-1862.
- Pajic I., Kljajic Z., Dogovic N., Sladic D., Juranic Z., Gasic MJ. A novel lectin from the sponge *Haliclona cratera*: isolation, characterization and biological activity. *Comp Biochem Physiol C Toxicol Pharmacol*. 2002; 132:213-221.
- Pan S., Tang J., Gu X. Isolation and characterization of a novel fucose-binding lectin from the gill of bighead carp (*Aristichthys nobilis*). *Vet Immunol Immunopathol*. 2010; 133:154-164.
- Park WB., Lyu SY., Kim JH., Choi SH., Chung HK., Ahn SH., Hong SY., Yoon TJ., Choi MJ. Inhibition of tumor growth and metastasis by Korean mistletoe lectin is associated with apoptosis and antiangiogenesis. *Cancer Biother Radiopharm*. 2001; 16:439-447.
- Peumans W.J., Van Damme EJM. Lectins as plant defense proteins. *Plant Physiol*. 1995(b); 107:347-352.
- Plattner VE., Wagner M., Ratzinger G., Gabor F., Wirth M. Targeted drug delivery: binding and uptake of plant lectins using human 5637 bladder cancer cells. *Eur J Pharm Biopharm*. 2008; 70:572-576.
- Sa RA., Gomes FS., Napoleao TH., Santos ND., Melo CML., Gusmao NB., Coelho LCBB., Paiva PMG., Bieber LW. Antibacterial and antifungal activities of *Myracrodruon urundeuva* heartwood. *Wood Science and Technology*. 2009; 43:85-95.
- Sá RA., Santos ND., da Silva CS., Napoleao TH., Gomes FS., Cavada BS., Coelho LC., Navarro DM., Bieber LW., Paiva PM. Larvicidal activity of lectins from *Myracrodruon urundeuva* on *Aedes aegypti*. *Comp Biochem Physiol C Toxicol Pharmacol*. 2009; 149:300-306.
- Sadeghi A., Smaghe G., Broeders S., Hernalsteens JP., De Greve H., Peumans WJ., Van Damme EJ. Ectopically expressed leaf and bulb lectins from garlic (*Allium sativum* L.) protect transgenic tobacco plants against cotton leafworm (*Spodoptera littoralis*). *Transgenic Res*. 2008; 17: 9-18.
- Santana GMS., Albuquerque LP., Simoes DA., Gusmao NB., Coelho LCBB., Paiva PMG. Isolation of a lectin from *Opuntia ficus indica* cladodes. *Acta Horticulturae*. 2009; 811: 281-286.
- Santi-Gadelha T., Gadelha CAA., Aragao KS., Oliveira CC., Mota MRL., Gomes RC., Pires AF., Toyama MH., Toyama DO., Alencar NMN., Criddle DN., Assrey AMS., Cavada BS. Purification and biological effects of *Araucaria angustifolia* (Araucariaceae) seed lectin. *Biochemical and Biophysical Research Communications* 2006; 350:1050-1055.
- Selitrennikoff CP. Antifungal proteins. *Applied and Environmental Microbiology*. 2001; 67:2883-2894.
- Semiglazov VF., Stepula VV., Dudov A., Schnitker J., Mengs U. Quality of life is improved in breast cancer patients by Standardised Mistletoe Extract PS76A2 during chemotherapy and follow-up: a randomised, placebo-controlled, double-blind, multicentre clinical trial. *Anticancer Res*. 2006; 26:1519-1529.
- Sharma A., Ng TB., Wong JH., Lin P. Purification and characterization of a lectin from *Phaseolus vulgaris* cv. (Anasazi beans). *J Biomed Biotechnol*. 2009; 92-95.
- Sharon N., Lis H. Lectins: cell-agglutinating and sugar-specific proteins. *Science*. 1972; 177: 949-959.
- Sharon N., Lis H., Lotan R. On the structural diversity of lectins. *Coll. Int. CNRS*. 1974; 221: 693-709.
- Sharon N., Lis H. Lectins as cell recognition molecules. *Science*. 1989; 246:227-234.
- Sharon N and Lis H. *Lectins* 2nd ed. Kluwer Scientific Publishers, Amsterdam (2003)
- Sharon N., Lis H. History of lectins: from hemagglutinins to biological recognition molecules. *Glycobiology*. 2004; 53-62.
- Singh K., Kaur M., Rup PJ., Singh J. Effects of Indian coral tree, *Erythrina indica* lectin on eggs and larval development of melon fruit fly, *Bactrocera cucurbitae*. *J. Environ. Biol*. 2009; 318: 509-514.
- Sitohy M., Doheim M., Badr H. Isolation and characterization of a lectin with antifungal activity from Egyptian *Pisum sativum* seeds. *Food Chemistry*. 2007; 104: 971-979.
- Stillmark PH. Doctoral Thesis. 1988.
- Sumner JB., Howell SF. The identification of the hemagglutinin of the jack bean with concanavalin A. *J. Bacteriol*. 1936; 32:227-237.
- Suzuki T., Amano Y., Fujita M., Kobayashi Y., Dohra H., Hirai H., Murata T., Usui T., Morita T., Kawagishi H. Purification, characterization, and cDNA cloning of a lectin from the mushroom *Pleurocybella porrigens*. *Biosci Biotechnol Biochem*. 2009; 73:702-709.
- Swanson MD., Winter HC., Goldstein IJ., Markovitz DM. A lectin isolated from bananas is a potent inhibitor of HIV replication. *J Biol Chem*. 2010; 285:8646-8655.
- Thakur K., Kaur M., Kaur S., Kaur A., Kamboj SS., Singh, J. Purification of *Colocasia esculenta* lectin and determination of its anti-insect potential towards *Bactrocera cucurbitae*. *Journal of Environmental Biol*. 2012; 34: 31-36.
- Tian Q., Wang W., Miao C., Peng H., Liu B., Leng F., Dai L., Chen F., Bao J. Purification, characterization and molecular cloning of a novel mannose-binding lectin from rhizomes of *Ophiopogon japonicus* with antiviral and antifungal activities. *Plant Sci*. 2008; 175: 877-884.
- Trindade MB., Lopes JLS., Soares-Costa A., Monteiro-Moreira AC., Moreira RA., Oliva MLV., Beltrami LM. Structural characterization of novel chitin-binding lectins from the genus *Artocarpus* and their antifungal activity. *Biochimica et Biophysica Acta*. 2006; 1764: 146-152.
- Van Parijs J., Broekaert WF., Goldstein IJ., Peumans WJ. Hevein: an antifungal protein from rubber-tree (*Hevea brasiliensis*) latex. *Planta*. 1999; 183:258-264.
- Vaz AFM., Costa RMPB., Melo AMMA., Oliva MLV., Santana LA., Silva-Lucca RA., Coelho LCBB., Correia MTS. Biocontrol of *Fusarium* species by a novel lectin with low ecotoxicity isolated from *Sebastiania jacobinensis*. *Food Chemistry*. 2010; 119:1507-1513.
- Velkov VV., Medvinsky AB., Sokolov MS., Marchenko AI. Will transgenic plants adversely affect the environment? *J Biosci*. 2005; 30:515-548.
- Vlodavsky I., and Sachs I. Lectin receptors on the cell surface membrane and the kinetics of lectin-induced cell agglutination. *Exp. Cell res*. 1975; 93(1):111-9.
- Wang Z., Zhang K., Sun X., Tang K., Zhang J. Enhancement of resistance to aphids by introducing the snowdrop lectin gene gna into maize plants. *J Biosci*. 2005; 30:627-638.
- Xia L., Ng TB. An antifungal protein from flageolet beans. *Peptides*. 2005; 26:2397-2403.
- Xia L., Ng TB. A hemagglutinin with mitogenic activity from dark red kidney beans. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2006; 844:213-216.
- Xu Q., Liu Y., Wang X., Gu H., Chen Z. Purification and characterization of a novel anti-fungal protein from *Gastrodia elata*. *Plant Physiol and Biochem*. 1998; 36:899-905.
- Yan Q., Jiang Z., Yang S., Deng W., Han L. A novel homodimeric lectin from *Astragalus mongholicus* with antifungal activity. *Arch Biochem Biophys*, 2005; 442:72-81.
- Ye XY., Ng TB., Tsang PW., Wang J. Isolation of a homodimeric lectin with antifungal and antiviral activities from red kidney bean (*Phaseolus vulgaris*) seeds. *J Protein Chem*. 2001; 20:367-375.
- Zem G.C., Badali O., Gaytan M., Hekmatjou H., Alvarez M., Nnoli J. Microbead analysis of cell binding to immobilized lectin: an alternative to microarrays in the development of carbohydrate drugs and diagnostic tests. *Acta Histochem*. 2006; 108:311-317.
- Zhang GQ., Sun J., Wang HX., Ng TB. A novel lectin with antiproliferative activity from the medicinal mushroom *Pholiota adiposa*. *Acta Biochim Pol*. 2009; 56:415-421.
- Zhang G., Sun J., Wang H., Ng TB. First isolation and characterization of a novel lectin with potent antitumor activity from a *Russula* mushroom. *Phytomedicine*. 2010; 17:775-781.

Zhao JK., Wang HX., Ng TB. Purification and characterization of a novel lectin from the toxic wild mushroom *Inocybe umbrinella*. *Toxicon*. 2009; 53:360–366.

**How to cite this article:**

Rabia Hamid, Akbar Masood, Ishfak H. Wani, and Shaista Rafiq., Lectins: Proteins with Diverse Applications. *J App Pharm Sci*. 2013; 3 (4 Suppl 1): S93-S103.