

Lectures on self-avoiding walks

Roland Bauerschmidt, Hugo Duminil-Copin, Jesse Goodman,
and Gordon Slade

ABSTRACT. These lecture notes provide a rapid introduction to a number of rigorous results on self-avoiding walks, with emphasis on the critical behaviour. Following an introductory overview of the central problems, an account is given of the Hammersley–Welsh bound on the number of self-avoiding walks and its consequences for the growth rates of bridges and self-avoiding polygons. A detailed proof that the connective constant on the hexagonal lattice equals $\sqrt{2 + \sqrt{2}}$ is then provided. The lace expansion for self-avoiding walks is described, and its use in understanding the critical behaviour in dimensions $d > 4$ is discussed. Functional integral representations of the self-avoiding walk model are discussed and developed, and their use in a renormalisation group analysis in dimension 4 is sketched. Problems and solutions from tutorials are included.

CONTENTS

Foreword	1
1. Introduction and overview of the critical behaviour	2
2. Bridges and polygons	11
3. The connective constant on the hexagonal lattice	17
4. The lace expansion	26
5. Lace expansion analysis in dimensions $d > 4$	35
6. Integral representation for walk models	43
7. Renormalisation group analysis in dimension 4	49
Appendix A. Solutions to the problems	60
Acknowledgements	71
References	71

Foreword

These notes are based on a course on Self-Avoiding Walks given in Búzios, Brazil, in August 2010, as part of the Clay Mathematics Institute Summer School

2010 *Mathematics Subject Classification*. Primary 82B41; Secondary 60K35
December 9, 2010.

and the XIV Brazilian Probability School. The course consisted of six lectures by Gordon Slade, a lecture by Hugo Duminil-Copin based on recent joint work with Stanislav Smirnov (see Section 3), and tutorials by Roland Bauerschmidt and Jesse Goodman. The written version of Slade's lectures was drafted by Bauerschmidt and Goodman, and the written version of Duminil-Copin's lecture was drafted by himself. The final manuscript was integrated and prepared jointly by the four authors.

1. Introduction and overview of the critical behaviour

These lecture notes focus on a number of rigorous results for self-avoiding walks on the d -dimensional integer lattice \mathbb{Z}^d . The model is defined by assigning equal probability to all paths of length n starting from the origin and without self-intersections. This family of probability measures is not consistent as n is varied, and thus does not define a stochastic process; the model is combinatorial in nature. The natural questions about self-avoiding walks concern the asymptotic behaviour as the length of the paths tends to infinity. Despite its simple definition, the self-avoiding walk is difficult to study in a mathematically rigorous manner. Many of the important problems remain unsolved, and the basic problems encompass many of the features and challenges of critical phenomena. This section gives the basic definitions and an overview of the critical behaviour.

1.1. Simple random walks. The basic reference model is *simple random walk* (SRW). Let $\Omega \subset \mathbb{Z}^d$ be the set of possible steps. The primary examples considered in these lectures are

$$(1.1) \quad \begin{array}{ll} \text{the nearest-neighbour model:} & \Omega = \{x \in \mathbb{Z}^d : \|x\|_1 = 1\}, \\ \text{the spread-out model:} & \Omega = \{x \in \mathbb{Z}^d : 0 < \|x\|_\infty \leq L\}, \end{array}$$

where L is a fixed integer, usually large. An n -step walk is a sequence $\omega = (\omega(0), \omega(1), \dots, \omega(n))$ with $\omega(j) - \omega(j-1) \in \Omega$ for $j = 1, \dots, n$. The n -step simple random walk is the uniform measure on n -step walks. We define the sets

$$(1.2) \quad \mathcal{W}_n(0, x) = \{\omega : \omega \text{ is an } n\text{-step walk with } \omega(0) = 0 \text{ and } \omega(n) = x\}$$

and

$$(1.3) \quad \mathcal{W}_n = \bigcup_{x \in \mathbb{Z}^d} \mathcal{W}_n(0, x).$$

1.2. Self-avoiding walks. The *weakly self-avoiding walk* and the *strictly self-avoiding walk* (the latter also called simply *self-avoiding walk*) are the main subject of these notes. These are random paths on \mathbb{Z}^d , defined as follows. Given an n -step walk $\omega \in \mathcal{W}_n$, and integers s, t with $0 \leq s < t \leq n$, let

$$(1.4) \quad U_{st} = U_{st}(\omega) = -1_{\{\omega(s)=\omega(t)\}} = \begin{cases} -1 & \text{if } \omega(s) = \omega(t), \\ 0 & \text{if } \omega(s) \neq \omega(t). \end{cases}$$

Fix $\lambda \in [0, 1]$. We assign to each path $\omega \in \mathcal{W}_n$ the weighting factor

$$(1.5) \quad \prod_{0 \leq s < t \leq n} (1 + \lambda U_{st}(\omega)).$$

The weights can also be expressed as Boltzmann weights:

$$(1.6) \quad \prod_{0 \leq s < t \leq n} (1 + \lambda U_{st}(\omega)) = \exp\left(-g \sum_{0 \leq s < t \leq n} 1_{\{\omega(s)=\omega(t)\}}\right)$$

with $g = -\log(1 - \lambda) \in [0, \infty)$ for $\lambda \in [0, 1)$. Making the convention $\infty \cdot 0 = 0$, the case $\lambda = 1$ corresponds to $g = \infty$.

The choice $\lambda = 0$ assigns equal weight to all walks in \mathcal{W}_n ; this is the case of the simple random walk. For $\lambda \in (0, 1)$, self-intersections are penalised but not forbidden, and the model is called the *weakly self-avoiding walk*. The choice $\lambda = 1$ prevents any return to a previously visited site, and defines the *self-avoiding walk* (SAW). More precisely, an n -step walk ω is a self-avoiding walk if and only if the expression (1.5) is non-zero for $\lambda = 1$, which happens if and only if ω visits each site at most once, and for such walks the weight equals 1.

These weights give rise to associated partition sums $c_n^{(\lambda)}(x)$ and $c_n^{(\lambda)}$ for walks in $\mathcal{W}_n(0, x)$ and \mathcal{W}_n , respectively:

$$(1.7) \quad c_n^{(\lambda)}(x) = \sum_{\omega \in \mathcal{W}_n(0, x)} \prod_{0 \leq s < t \leq n} (1 + \lambda U_{st}(\omega)), \quad c_n^{(\lambda)} = \sum_{x \in \mathbb{Z}^d} c_n^{(\lambda)}(x).$$

In the case $\lambda = 1$, $c_n^{(1)}(x)$ counts the number of self-avoiding walks of length n ending at x , and $c_n^{(1)}$ counts all n -step self-avoiding walks. The case $\lambda = 0$ reverts to simple random walk, for which $c_n^{(0)} = |\Omega|^n$. When $\lambda = 1$ we will often drop the superscript (1) and write simple c_n instead of $c_n^{(1)}$.

We also define probability measures $\mathbb{Q}_n^{(\lambda)}$ on \mathcal{W}_n with expectations $\mathbb{E}_n^{(\lambda)}$:

$$(1.8) \quad \mathbb{Q}_n^{(\lambda)}(A) = \frac{1}{c_n^{(\lambda)}} \sum_{\omega \in A} \prod_{0 \leq s < t \leq n} (1 + \lambda U_{st}(\omega)) \quad (A \subset \mathcal{W}_n),$$

$$(1.9) \quad \mathbb{E}_n^{(\lambda)}(X) = \frac{1}{c_n^{(\lambda)}} \sum_{\omega \in \mathcal{W}_n} X(\omega) \prod_{0 \leq s < t \leq n} (1 + \lambda U_{st}(\omega)) \quad (X : \mathcal{W}_n \rightarrow \mathbb{R}).$$

The measures $\mathbb{Q}_n^{(\lambda)}$ define the weakly self-avoiding walk when $\lambda \in (0, 1)$ and the strictly self-avoiding walk when $\lambda = 1$. Occasionally we will also consider self-avoiding walks that do not begin at the origin.

1.3. Subadditivity and the connective constant. The sequence $c_n^{(\lambda)}$ has the following submultiplicativity property:

$$(1.10) \quad c_{n+m}^{(\lambda)} \leq \sum_{\omega \in \mathcal{W}_{n+m}} \prod_{0 \leq s < t \leq n} (1 + \lambda U_{st}) \prod_{n \leq s' < t' \leq n+m} (1 + \lambda U_{s't'}) \leq c_n^{(\lambda)} c_m^{(\lambda)}.$$

Therefore, $\log c_n^{(\lambda)}$ is a *subadditive* sequence: $\log c_{n+m}^{(\lambda)} \leq \log c_n^{(\lambda)} + \log c_m^{(\lambda)}$.

LEMMA 1.1. *If $a_1, a_2, \dots \in \mathbb{R}$ obey $a_{n+m} \leq a_n + a_m$ for every n, m , then*

$$(1.11) \quad \lim_{n \rightarrow \infty} \frac{a_n}{n} = \inf_{n \geq 1} \frac{a_n}{n} \in [-\infty, \infty).$$

PROOF. See Problem 1.1. The value $-\infty$ is possible, e.g., for the sequence $a_n = -n^2$. \square

Applying Lemma 1.1 to $c_n^{(\lambda)}$ gives the existence of μ_λ such that $\lim_{n \rightarrow \infty} \frac{1}{n} \log c_n^{(\lambda)} = \log \mu_\lambda \leq \frac{1}{n} \log c_n^{(\lambda)}$ for all n , i.e.,

$$(1.12) \quad \mu_\lambda = \lim_{n \rightarrow \infty} (c_n^{(\lambda)})^{1/n} \text{ exists, and } c_n^{(\lambda)} \geq \mu_\lambda^n \text{ for all } n.$$

In the special case $\lambda = 1$, we write simply $\mu = \mu_1$. This μ , which depends on d (and also on L for the spread-out model), is called the *connective constant*. For the nearest-neighbour model, by counting only walks that move in positive coordinate directions, and by counting walks that are restricted only to prevent immediate reversals of steps, we obtain

$$(1.13) \quad d^n \leq c_n \leq 2d(2d-1)^{n-1} \quad \text{which implies} \quad d \leq \mu \leq 2d-1.$$

For $d = 2$, the following rigorous bounds are known:

$$(1.14) \quad \mu \in [2.625\,622, 2.679\,193].$$

The lower bound is due to Jenson [47] via bridge enumeration (bridges are defined in Section 2.1 below), and the upper bound is due to Pönitz and Tittmann [64] by comparison with finite-memory walks. The estimate

$$(1.15) \quad \mu = 2.638\,158\,530\,31(3)$$

is given in [45]; here the 3 in parentheses represents the subjective error in the last digit. It has been observed that $1/\mu$ is well approximated by the smallest positive root of $581x^4 + 7x^2 - 13 = 0$ [23, 48], though no derivation or explanation of this quartic polynomial is known, and later evidence has raised doubts about its validity [45].

Even though the definition of self-avoiding walks has been restricted to the graph \mathbb{Z}^d thus far, it applies more generally. In 1982, arguments based on a Coulomb gas formalism led Nienhuis [61] to predict that on the hexagonal lattice the connective constant is equal to $\sqrt{2 + \sqrt{2}}$. This was very recently proved by Duminil-Copin and Smirnov [24], whose theorem is the following.

THEOREM 1.2. *The connective constant for the hexagonal lattice is*

$$(1.16) \quad \mu = \sqrt{2 + \sqrt{2}}.$$

The proof of Theorem 1.2 is presented in Section 3 below. Except for trivial cases, this is the only lattice for which the connective constant is known explicitly.

Returning to \mathbb{Z}^d , in 1963, Kesten [50] proved that

$$(1.17) \quad \lim_{n \rightarrow \infty} \frac{c_{n+2}}{c_n} = \mu^2,$$

but it remains an open problem (for $d = 2, 3, 4$) to prove that

$$(1.18) \quad \lim_{n \rightarrow \infty} \frac{c_{n+1}}{c_n} = \mu.$$

Even the proof of $c_{n+1} \geq c_n$ is a non-trivial result, proved by O'Brien [62], though it is not hard to show that $c_{n+2} \geq c_n$.

1.4. $1/d$ expansion. It was proved by Hara and Slade [35] that the connective constant $\mu(d)$ for \mathbb{Z}^d (with nearest-neighbour steps) has an asymptotic expansion in powers of $1/2d$ as $d \rightarrow \infty$: There exist integers $a_i \in \mathbb{Z}$, $i = -1, 0, 1, \dots$ such that

$$(1.19) \quad \mu(d) \sim \sum_{i=-1}^{\infty} \frac{a_i}{(2d)^i}$$

in the sense that $\mu(d) = a_{-1}(2d) + a_0 + \dots + a_{M-1}(2d)^{-(M-1)} + O(d^{-M})$, for each fixed M . In Problem 5.1 below, the first three terms are computed. The constant in the $O(d^{-M})$ term may depend on M . It is expected, though not proved, that the asymptotic series in (1.19) has radius of convergence 0, so that the right-hand side of (1.19) diverges for each fixed d . The values of a_i are known for $i = -1, 0, \dots, 11$ and grow rapidly in magnitude; see Clisby, Liang, and Slade [21].

Graham [26] has proved Borel-type error bounds for the asymptotic expansion of $z_c = z_c(d) = \mu^{-1}$. Namely, writing the asymptotic expansion of z_c as $\sum_{i=1}^{\infty} \alpha_i (2d)^{-i}$, there is a constant C , independent of d and M , such that for each M and for all $d \geq 1$,

$$(1.20) \quad \left| z_c - \sum_{i=1}^{M-1} \frac{\alpha_i}{(2d)^i} \right| \leq \frac{C^M M!}{(2d)^M}.$$

An extension of (1.20) to *complex* values of the dimension d would be needed in order to apply the method of Borel summation to recover the value of z_c , and hence of $\mu(d)$, from the asymptotic series.

1.5. Critical exponents. It is a characteristic feature of models of statistical mechanics at the critical point that there exist *critical exponents* which describe the asymptotic behaviour on the large scale. It is a deep conjecture, not yet properly understood mathematically, that these critical exponents are *universal*, meaning that they depend only on the spatial dimension of the system, but not on details such as the specific lattice in \mathbb{R}^d . For the case of the self-avoiding walk, this conjecture of universality extends to lack of dependence on the constant λ , as soon as $\lambda > 0$. We now introduce the critical exponents, and in Section 1.6 we will discuss what is known about them in more detail.

1.5.1. *Number of self-avoiding walks.* It is predicted that for each d there is a constant γ such that for all $\lambda \in (0, 1]$, and for both the nearest-neighbour and spread-out models,

$$(1.21) \quad c_n^{(\lambda)} \sim A_\lambda \mu_\lambda^n n^{\gamma-1}.$$

Here $f(n) \sim g(n)$ means $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$. The predicted values of the critical exponent γ are:

$$(1.22) \quad \gamma = \begin{cases} 1 & d = 1, \\ \frac{43}{32} & d = 2, \\ 1.16\dots & d = 3, \\ 1 & d = 4, \\ 1 & d \geq 5. \end{cases}$$

In fact, for $d = 4$, the prediction involves a logarithmic correction:

$$(1.23) \quad c_n^{(\lambda)} \sim A_\lambda \mu_\lambda^n (\log n)^{1/4}.$$

This situation should be compared with simple random walk, for which $c_n^{(0)} = |\Omega|^n$, so that μ_0 is equal to the degree $|\Omega|$ of the lattice, and $\gamma = 1$.

In the case of the self-avoiding walk (i.e., $\lambda = 1$), γ has a probabilistic interpretation. Sampling independently from two n -step self-avoiding walks uniformly,

$$(1.24) \quad \mathbb{P}(\omega_1 \cap \omega_2 = \{0\}) = \frac{c_{2n}}{c_n^2} \sim \text{const} \frac{1}{n^{\gamma-1}},$$

so γ is a measure of how likely it is for two self-avoiding walks to avoid each other. The analogous question for SRW is discussed in [53].

Despite the precision of the prediction (1.21), the best rigorously known bounds in dimension $d = 2, 3, 4$ are very far from tight and almost 50 years old. In [29], Hammersley and Welsh proved that, for all $d \geq 2$,

$$(1.25) \quad \mu^n \leq c_n \leq \mu^n e^{\kappa\sqrt{n}}$$

(the lower bound is just subadditivity, the upper bound is nontrivial). This was improved slightly by Kesten [50], who showed that for $d = 3, 4, \dots$,

$$(1.26) \quad \mu^n \leq c_n \leq \mu^n \exp\left(\kappa n^{2/(d+2)} \log n\right).$$

The proof of the Hammersley–Welsh bound is the subject of Section 2.1.

1.5.2. *Mean-square displacement.* Let $|x|$ denote the Euclidean norm of $x \in \mathbb{R}^d$. It is predicted that for $\lambda \in (0, 1]$, and for both the nearest-neighbour and spread-out models,

$$(1.27) \quad \mathbb{E}_n^{(\lambda)} |\omega(n)|^2 \sim D_\lambda n^{2\nu},$$

with

$$(1.28) \quad \nu = \begin{cases} 1 & d = 1, \\ \frac{3}{4} & d = 2, \\ 0.588\dots & d = 3, \\ \frac{1}{2} & d = 4, \\ \frac{1}{2} & d \geq 5. \end{cases}$$

Again a logarithmic correction is predicted for $d = 4$:

$$(1.29) \quad \mathbb{E}_n^{(\lambda)} |\omega(n)|^2 \sim D_\lambda n (\log n)^{1/4}.$$

This should be compared with the SRW, for which $\nu = \frac{1}{2}$ in all dimensions.

Almost nothing is known rigorously about ν in dimensions 2, 3, 4. It is an open problem to show that the mean-square displacement grows at least as rapidly as simple random walk, and grows more slowly than ballistically, i.e., it has not been proved that

$$(1.30) \quad cn \leq \mathbb{E}_n^{(1)} |\omega(n)|^2 \leq Cn^{2-\epsilon},$$

or even that the endpoint is typically as far away as the surface of a ball of volume n , i.e., $cn^{2/d} \leq \mathbb{E}_n^{(1)} |\omega(n)|^2$. Madras (unpublished) has shown $\mathbb{E}_n^{(1)} |\omega(n)|^2 \geq cn^{4/3d}$.

1.5.3. *Two-point function and susceptibility.* The two-point function is defined by

$$(1.31) \quad G_z^{(\lambda)}(x) = \sum_{n=0}^{\infty} c_n^{(\lambda)}(x) z^n,$$

and the susceptibility by

$$(1.32) \quad \chi^{(\lambda)}(z) = \sum_{x \in \mathbb{Z}^d} G_z^{(\lambda)}(x) = \sum_{n=0}^{\infty} c_n^{(\lambda)} z^n.$$

Since $\chi^{(\lambda)}$ is a power series whose coefficients satisfy (1.12), its radius of convergence $z_c^{(\lambda)}$ is given by $z_c^{(\lambda)} = \mu_\lambda^{-1}$. The value $z_c^{(\lambda)}$ is referred to as the *critical point*.

PROPOSITION 1.3. *Fix $\lambda \in [0, 1]$, $z \in (0, z_c^{(\lambda)})$. Then $G_z^{(\lambda)}(x)$ decays exponentially in x .*

PROOF. For simplicity, we consider only the nearest-neighbour model, and we omit λ from the notation. Since $c_n(x) = 0$ if $n < \|x\|_1$,

$$(1.33) \quad G_z(x) = \sum_{n=\|x\|_1}^{\infty} c_n(x) z^n \leq \sum_{n=\|x\|_1}^{\infty} c_n z^n.$$

Fix $z < z_c = 1/\mu$ and choose $\epsilon > 0$ such that $z(\mu + \epsilon) < 1$. Since $c_n^{1/n} \rightarrow \mu$, there exists $K = K(\epsilon)$ such that $c_n \leq K(\mu + \epsilon)^n$ for all n . Hence

$$(1.34) \quad G_z(x) \leq K \sum_{n=\|x\|_1}^{\infty} (z(\mu + \epsilon))^n \leq K'(z(\mu + \epsilon))^{\|x\|_1},$$

as claimed. \square

We restrict temporarily to $\lambda = 1$. Much is known about $G_z(x)$ for $z < z_c$: there is a norm $|\cdot|_z$ on \mathbb{R}^d , satisfying $\|u\|_\infty \leq |u|_z \leq \|u\|_1$ for all $u \in \mathbb{R}^d$, such that $m(z) = \lim_{|x|_z \rightarrow \infty} -\frac{\log G_z(x)}{|x|_z}$ exists and is finite. The *correlation length* is defined by $\xi(z) = 1/m(z)$, and hence approximately

$$(1.35) \quad G_z(x) \approx e^{-|x|_z/\xi(z)}.$$

Indeed more precise asymptotics (Ornstein–Zernike decay) are known [17, 57, 15]:

$$(1.36) \quad G_z(x) \sim \frac{c}{|x|_z^{(d-1)/2}} e^{-|x|_z/\xi(z)} \quad \text{as } x \rightarrow \infty,$$

and the arguments leading to this also prove that

$$(1.37) \quad \lim_{z \nearrow z_c} \xi(z) = \infty.$$

As a refinement of (1.37), it is predicted that as $z \nearrow z_c$,

$$(1.38) \quad \xi(z) \sim \text{const} \left(1 - \frac{z}{z_c}\right)^{-\nu},$$

and that, in addition, as $|x| \rightarrow \infty$ (for $d \geq 2$),

$$(1.39) \quad G_{z_c}(x) \sim \frac{\text{const}}{|x|^{d-2+\eta}}.$$

The exponents γ , η and ν are predicted to be related to each other via *Fisher's relation* (see, e.g., [57]):

$$(1.40) \quad \gamma = (2 - \eta)\nu.$$

There is typically a correspondence between the asymptotic growth of the coefficients in a generating function and the behaviour of the generating function near its dominant singularity. For our purpose we note that, under suitable hypotheses,

$$(1.41) \quad a_n \sim \frac{n^{\gamma-1}}{R^n} \text{ as } n \rightarrow \infty \quad \Leftrightarrow \quad \sum_n a_n z^n \sim \frac{C}{(1 - z/R)^\gamma} \text{ as } z \nearrow R.$$

The easier \implies direction is known as an Abelian theorem, and the more delicate \impliedby direction is known as a Tauberian theorem [36]. With this in mind, our earlier prediction for $c_n^{(\lambda)}$ for $\lambda \in (0, 1]$ corresponds to:

$$(1.42) \quad \chi^{(\lambda)}(z) \sim \frac{\text{const}_\lambda}{(1 - z/z_c)^\gamma}$$

as $z \nearrow z_c$, with an additional factor $|\log(1 - z/z_c)|^{1/4}$ on the right-hand side when $d = 4$.

1.6. Effect of the dimension. Universality asserts that self-avoiding walks on different lattices in a fixed dimension d should behave in the same way, independently of the fine details of how the model is defined. However, the behaviour does depend very strongly on the dimension.

1.6.1. $d = 1$. For the nearest-neighbour model with $\lambda = 1$ it is a triviality that $c_n^{(1)} = 2$ for all $n \geq 1$ and $|\omega(n)| = n$ for all ω , since a self-avoiding walk must continue either in the negative or in the positive direction. Any configuration $\omega \in \mathcal{W}_n$ is possible when $\lambda \in (0, 1)$, however, and it is by no means trivial to prove that the critical behaviour when $\lambda \in (0, 1)$ is similar to the case of $\lambda = 1$. The following theorem of König [52] (extending a result of Greven and den Hollander [27]) proves that the weakly self-avoiding walk measure (1.8) does have ballistic behaviour for all $\lambda \in (0, 1)$.

THEOREM 1.4. *Let $d = 1$. For each $\lambda \in (0, 1)$, there exist $\theta(\lambda) \in (0, 1)$ and $\sigma(\lambda) \in (0, \infty)$ such that for all $u \in \mathbb{R}$,*

$$(1.43) \quad \lim_{n \rightarrow \infty} \mathbb{Q}_n^{(\lambda)} \left(\frac{|\omega(n)| - n\theta}{\sigma\sqrt{n}} \leq u \right) = \int_{-\infty}^u \frac{e^{-t^2/2}}{\sqrt{2\pi}} dt.$$

A similar result is proved in [52] for the 1-dimensional spread-out strictly self-avoiding walk. The result of Theorem 1.4 should be contrasted to the case $\lambda = 0$, which has diffusive rather than ballistic behaviour. It remains an open problem to prove the intuitively appealing statement that θ should be an increasing function of λ . A review of results for $d = 1$ is given in [40].

1.6.2. $d = 2$. Based on non-rigorous Coulomb gas methods, Nienhuis [61] predicted that $\gamma = \frac{43}{32}$, $\nu = \frac{3}{4}$. These predicted values have been confirmed numerically by Monte Carlo simulation, e.g., [55], and exact enumeration of self-avoiding walks up to length $n = 71$ [46].

Lawler, Schramm and Werner [54] have given major mathematical support to these predictions. Roughly speaking, they show that if self-avoiding walk has

a scaling limit, and if this scaling limit has a certain conformal invariance property, then the scaling limit must be $\text{SLE}_{8/3}$ (the Schramm–Loewner evolution with parameter $\kappa = \frac{8}{3}$). The values of γ and ν are then recovered from an $\text{SLE}_{8/3}$ computation. Numerical evidence supporting the statement that the scaling limit is $\text{SLE}_{8/3}$ is given in [49]. However, until now, it remains an open problem to prove the required existence and conformal invariance of the scaling limit.

The result of [54] is discussed in greater detail in the course of Vincent Beffara [1]. Here, we describe it only briefly, as follows. Consider a simply connected domain Ω in the complex plane \mathbb{C} with two points a and b on the boundary. Fix $\delta > 0$, and let $(\Omega_\delta, a_\delta, b_\delta)$ be a discrete approximation of (Ω, a, b) in the following sense: Ω_δ is the largest finite domain of $\delta\mathbb{Z}^2$ included in Ω , a_δ and b_δ are the closest vertices of $\delta\mathbb{Z}^2$ to a and b respectively. When δ goes to 0, this provides an approximation of the domain.

For fixed $z, \delta > 0$, there is a probability measure on the set of self-avoiding walks ω between a_δ and b_δ that remain in Ω_δ by assigning to ω a *Boltzmann weight* proportional to $z^{\ell(\omega)}$, where $\ell(\omega)$ denotes the length of ω . We obtain a random piecewise linear curve denoted by ω_δ .

It is possible to prove that when $z < z_c = 1/\mu$, walks are penalised so much with respect to their length that ω_δ becomes straight when δ goes to 0; this is closely related to the Ornstein–Zernike decay results. On the other hand, it is expected that, when $z > z_c$, the entropy wins against the penalisation and ω_δ becomes space filling when δ tends to 0. Finally, when $z = z_c$, the sequence of measures conjecturally converges to a random continuous curve. It is for this case that we have the following conjecture of Lawler, Schramm and Werner [54].

CONJECTURE 1.5. *For $z = z_c$, the random curve ω_δ converges to $\text{SLE}_{8/3}$ from a and b in the domain Ω .*

It remains a major open problem in 2-dimensional statistical mechanics to prove the conjecture.

1.6.3. $d = 3$. For $d = 3$, there are no rigorous results for critical exponents, and no mathematically well-defined candidate has been proposed for the scaling limit. An early prediction for the values of ν , referred to as the Flory values [25], was $\nu = \frac{3}{d+2}$ for $1 \leq d \leq 4$. This does give the correct answer for $d = 1, 2, 4$, but it is not quite accurate for $d = 3$ —the Flory argument is very remote from a rigorous mathematical proof. Flory’s interest in the problem was motivated by the use of SAWs to model polymer molecules; this application is discussed in detail in the course of Frank den Hollander [42] (see also [43]).

For $d = 3$, there are three methods to compute the exponents approximately. In one method, non-rigorous field theory computations in theoretical physics [28] combine the $n \rightarrow 0$ limit for the $O(n)$ model with an expansion in $\epsilon = 4 - d$ about dimension $d = 4$, with $\epsilon = 1$. Secondly, Monte Carlo studies have been carried out with walks of length 33,000,000 [20], using the pivot algorithm [58, 44]. Finally, exact enumeration plus series analysis has been used; currently the most extensive enumerations in dimensions $d \geq 3$ use the lace expansion [21], and for $d = 3$ walks have been enumerated to length $n = 30$. The exact enumeration estimates for $d = 3$ are $\mu = 4.684043(12)$, $\gamma = 1.1568(8)$, $\nu = 0.5876(5)$ [21]. Monte Carlo estimates are consistent with these values: $\gamma = 1.1575(6)$ [16] and $\nu = 0.587597(7)$ [20].

1.6.4. $d = 4$. Four dimensions is the *upper critical dimension* for the self-avoiding walk. This term encapsulates the notion that for $d > 4$ self-avoiding walk has the same critical behaviour as simple random walk, while for $d < 4$ it does not. The dimension 4 can be guessed by considering the fractal properties of the simple random walk: for $d \geq 2$, the path of a simple random walk is two-dimensional. If $d > 4$, two independent two-dimensional objects should generically not intersect, so that the effect of self-interaction between the past and the future of a simple random walk should be negligible. In $d = 4$, the expected number of intersections between two independent random walks tends to infinity, but only logarithmically in the length. Such considerations are related to the logarithmic corrections that appear in (1.23) and (1.29).

The existence of logarithmic corrections to scaling has been proved for models of weakly self-avoiding walk on a 4-dimensional *hierarchical* lattice, using rigorous renormalisation group methods [5, 9, 10, 32]. The hierarchical lattice is a simplification of the hypercubic lattice \mathbb{Z}^4 which is particularly amenable to the renormalisation group approach. Recently there has been progress in the application of renormalisation group methods to a continuous-time weakly self-avoiding walk model on \mathbb{Z}^4 itself, and in particular it has been proved in this context that the critical two-point function has $|x|^{-2}$ decay [12], which is a statement that the critical exponent η is equal to 0. This is the topic of Section 7 below.

1.6.5. $d \geq 5$. Using the lace expansion, it has been proved that for the nearest-neighbour model in dimensions $d \geq 5$ the critical exponents exist and take their so-called *mean field* values $\gamma = 1$, $\nu = \frac{1}{2}$ [34, 33] and $\eta = 0$ [30], and that the scaling limit is Brownian motion [33]. The lace expansion for self-avoiding walks is discussed in Section 4, and its application to prove simple random walk behaviour in dimensions $d \geq 5$ is discussed in Section 5.

1.7. Tutorial.

PROBLEM 1.1. Let (a_n) be a real-valued sequence that is subadditive, that is, $a_{n+m} \leq a_n + a_m$ holds for all n, m . Prove that $\lim_{n \rightarrow \infty} n^{-1}a_n$ exists in $[-\infty, \infty)$ and equals $\inf_n n^{-1}a_n$.

PROBLEM 1.2. Prove that the connective constant μ for the nearest-neighbour model on the square lattice \mathbb{Z}^2 obeys the strict inequalities $2 < \mu < 3$.

PROBLEM 1.3. A family of probability measures (\mathbb{P}_n) on \mathcal{W}_n is called consistent if $\mathbb{P}_n(\omega) = \sum_{\rho > \omega} \mathbb{P}_m(\rho)$ for all $m > n$ and for all $\omega \in \mathcal{W}_n$, where the sum is over all $\rho \in \mathcal{W}_m$ whose first n steps agree with ω . Show that $\mathbb{Q}_n^{(1)}$, the uniform measure on SAWs, does not provide a consistent family.

PROBLEM 1.4. Show that the Fourier transform of the two-point function of the 1-dimensional strictly self-avoiding walk is given by

$$(1.44) \quad \hat{G}_z(k) = \frac{1 - z^2}{1 + z^2 - 2z \cos k}.$$

Here $\hat{f}(k) = \sum_{x \in \mathbb{Z}^d} f(x) e^{ik \cdot x}$.

PROBLEM 1.5. Suppose that $f(z) = \sum_{n=0}^{\infty} a_n z^n$ has radius of convergence 1. Suppose that $|f(z)| \leq c|1 - z|^{-b}$ uniformly in $|z| < 1$, with $b \geq 1$. Prove that, for

some constant C , $|a_n| \leq Cn^{b-1}$ if $b > 1$, and that $|a_n| \leq C \log n$ if $b = 1$. Hint:

$$(1.45) \quad a_n = \frac{1}{2\pi i} \oint_{\Gamma_n} \frac{f(z)}{z^{n+1}} dz,$$

where $\Gamma_n = \{z \in \mathbb{C} : |z| = 1 - \frac{1}{n}\}$.

PROBLEM 1.6. Consider the nearest-neighbour simple random walk $(X_n)_{n \geq 0}$ on \mathbb{Z}^d started at the origin. Let $D(x) = (2d)^{-1} 1_{\{\|x\|_1=1\}}$ denote its step distribution. The two-point function for simple random walk is defined by

$$(1.46) \quad C_z(x) = \sum_{n \geq 0} c_n^{(0)}(x) z^n = \sum_{n \geq 0} D^{*n}(x) (2dz)^n,$$

where D^{*n} denotes the n -fold convolution of D with itself.

(a) Let u denote the probability that the walk ever returns to the origin. The walk is recurrent if $u = 1$ and transient if $u < 1$. Let N denote the random number of visits to the origin, including the initial visit at time 0, and let $m = \mathbb{E}(N)$. Show that $m = (1 - u)^{-1}$; so the walk is recurrent if and only if $m = \infty$.

(b) Show that

$$(1.47) \quad m = \sum_{n \geq 0} \mathbb{P}(X_n = 0) = \int_{[-\pi, \pi]^d} \frac{1}{1 - \hat{D}(k)} \frac{d^d k}{(2\pi)^d}.$$

Thus transience is characterised by the integrability of $\hat{C}_{z_0}(k)$, where $z_0 = (2d)^{-1}$.

(c) Show that the walk is recurrent in dimensions $d \leq 2$ and transient for $d > 2$.

PROBLEM 1.7. Let $X^1 = (X_i^1)_{i \geq 0}$ and $X^2 = (X_i^2)_{i \geq 0}$ be two independent nearest-neighbour simple random walks on \mathbb{Z}^d started at the origin, and let

$$(1.48) \quad I = \sum_{i \geq 0} \sum_{j \geq 0} 1_{\{X_i^1 = X_j^2\}}$$

be the random number of intersections of the two walks. Show that

$$(1.49) \quad \mathbb{E}(I) = \int_{[-\pi, \pi]^d} \frac{1}{[1 - \hat{D}(k)]^2} \frac{d^d k}{(2\pi)^d}.$$

Thus $\mathbb{E}(I)$ is finite if and only if \hat{C}_{z_0} is square integrable. Conclude that the expected number of intersections is finite if $d > 4$ and infinite if $d \leq 4$.

2. Bridges and polygons

Throughout this section, we consider only the nearest-neighbour strictly self-avoiding walk on \mathbb{Z}^d . We will introduce a class of self-avoiding walks called bridges, and will show that the number of bridges grows with the same exponential rate as the number of self-avoiding walks, namely as μ^n . The analogous fact for the hexagonal lattice \mathbb{H} will be used in Section 3 as an ingredient in the proof that the connective constant for \mathbb{H} is $\sqrt{2 + \sqrt{2}}$. The study of bridges will also lead to the proof of the Hammersley–Welsh bound (1.25) on c_n . Finally, we will study self-avoiding polygons, and show that they too grow in number as μ^n .

2.1. Bridges and the Hammersley–Welsh bound. For a self-avoiding walk ω , denote by $\omega_1(i)$ the first spatial coordinate of $\omega(i)$.

DEFINITION 2.1. An n -step *bridge* is an n -step SAW ω such that

$$(2.1) \quad \omega_1(0) < \omega_1(i) \leq \omega_1(n) \quad \text{for } i = 1, 2, \dots, n.$$

Let b_n be the number of n -step bridges with $\omega(0) = 0$ for $n > 1$, and $b_0 = 1$.

While the number of self-avoiding walks is a *submultiplicative* sequence, the number of bridges is *supermultiplicative*:

$$(2.2) \quad b_{n+m} \geq b_n b_m.$$

Thus, applying Lemma 1.1 to $-\log b_n$, we obtain the existence of the bridge growth constant μ_{Bridge} defined by

$$(2.3) \quad \mu_{\text{Bridge}} = \lim_{n \rightarrow \infty} b_n^{1/n} = \sup_{n \geq 1} b_n^{1/n}.$$

Using the trivial inequality $\mu_{\text{Bridge}} \leq \mu$ we conclude that

$$(2.4) \quad b_n \leq \mu_{\text{Bridge}}^n \leq \mu^n.$$

DEFINITION 2.2. An n -step *half-space walk* is an n -step SAW ω with

$$(2.5) \quad \omega_1(0) < \omega_1(i) \quad \text{for } i = 1, 2, \dots, n.$$

Let $h_0 = 1$, and for $n \geq 1$, let h_n denote the number of n -step half-space walks with $\omega(0) = 0$.

DEFINITION 2.3. The *span* of an n -step SAW ω is

$$(2.6) \quad \max_{0 \leq i \leq n} \omega_1(i) - \min_{0 \leq i \leq n} \omega_1(i).$$

Let $b_{n,A}$ be the number of n -step bridges with span A .

We will use the following result on integer partitions which dates back to 1917, due to Hardy and Ramanujan [37].

THEOREM 2.4. For an integer $A \geq 1$, let $P_D(A)$ denote the number of ways of writing $A = A_1 + \dots + A_k$ with $A_1 > \dots > A_k \geq 1$, for any $k \geq 1$. Then

$$(2.7) \quad \log P_D(A) \sim \pi \left(\frac{A}{3} \right)^{1/2}$$

as $A \rightarrow \infty$.

PROPOSITION 2.5. $h_n \leq P_D(n)b_n$ for all $n \geq 1$.

PROOF. Set $n_0 = 0$ and inductively define

$$(2.8) \quad A_{i+1} = \max_{j > n_i} (-1)^i (\omega_1(j) - \omega_1(n_i))$$

and

$$(2.9) \quad n_{i+1} = \max \{ j > n_i : (-1)^i (\omega_1(j) - \omega_1(n_i)) = A_{i+1} \}.$$

In words, $j = n_1$ maximises $\omega_1(j)$, $j = n_2$ minimises $\omega_1(j)$ for $j > n_1$, n_3 maximises $\omega_1(j)$ for $j > n_2$, and so on in an alternating pattern. In addition $A_1 = \omega_1(n_1) - \omega_1(n_0)$, $A_2 = \omega_1(n_1) - \omega_1(n_2)$ and so on. Moreover, the n_i are chosen to be the last times these extrema are attained.

$A\mu^n n^{\gamma-1}$ from (1.21). It is an open problem to improve Theorem 2.6 in $d = 2, 3, 4$ beyond the result of Kesten [50] shown in (1.26).




FIGURE 2. The decomposition of a self-avoiding walk into two half-space walks.

PROOF OF THEOREM 2.6. We first prove

$$(2.14) \quad c_n \leq \sum_{m=0}^n h_{n-m} h_{m+1},$$

using the decomposition depicted in Figure 2, as follows. Given an n -step SAW ω , let

$$(2.15) \quad x_1 = \min_{0 \leq i \leq n} \omega_1(i), \quad m = \max \{i : \omega_1(i) = x_1\}.$$

Write e_1 for the unit vector in the first coordinate direction of \mathbb{Z}^d . Then (after translating by $\omega(m)$) the walk $(\omega(m), \omega(m+1), \dots, \omega(n))$ is an $(n-m)$ -step half-space walk, and (after translating by $\omega(m) - e_1$) the walk $(\omega(m) - e_1, \omega(m), \omega(m-1), \dots, \omega(1), \omega(0))$ is an $(m+1)$ -step half-space walk. This proves (2.14).

Next we apply Proposition 2.5 in (2.14) and use (2.2) to get

$$(2.16) \quad \begin{aligned} c_n &\leq \sum_{m=0}^n P_D(n-m) P_D(m+1) b_{n-m} b_{m+1} \\ &\leq b_{n+1} \sum_{m=0}^n P_D(n-m) P_D(m+1). \end{aligned}$$

Fix $B > B' > \pi(\frac{2}{3})^{1/2}$. By Theorem 2.4, there is $K > 0$ such that $P_D(A) \leq K \exp(B'(A/2)^{1/2})$ and consequently

$$(2.17) \quad P_D(n-m) P_D(m+1) \leq K^2 \exp \left[B' \left(\sqrt{\frac{n-m}{2}} + \sqrt{\frac{m+1}{2}} \right) \right].$$

The bound $x^{1/2} + y^{1/2} \leq (2x + 2y)^{1/2}$ now gives

$$(2.18) \quad c_n \leq (n+1) K^2 e^{B'\sqrt{n+1}} b_{n+1} \leq e^{B\sqrt{n}} b_{n+1}$$

if $n \geq n_0(B)$. By (2.4), the result follows. \square

COROLLARY 2.7. For $n \geq n_0(B)$,

$$(2.19) \quad b_n \geq c_{n-1} e^{-B\sqrt{n-1}} \geq \mu^{n-1} e^{-B\sqrt{n-1}}.$$

In particular, $b_n^{1/n} \rightarrow \mu$ and so $\mu_{\text{Bridge}} = \mu$.

COROLLARY 2.8. Define the bridge generating function $B(z) = \sum_{n=0}^{\infty} b_n z^n$. Then

$$(2.20) \quad \chi(z) \leq \frac{1}{z} e^{2(B(z)-1)}$$

and in particular $B(1/\mu) = \infty$.

PROOF. In the proof of Proposition 2.5, we decomposed a half-space walk into subwalks on $[n_{i-1}, n_i]$ for $i = 1, \dots, K$. Note that each such subwalk was in fact a bridge of span A_i . With this observation, we conclude that

$$(2.21) \quad h_n \leq \sum_{k=1}^{\infty} \sum_{A_1 > \dots > A_k} \sum_{0=n_0 < n_1 < \dots < n_k=n} \prod_{i=1}^k b_{n_i - n_{i-1}, A_i}$$

(the second sum is over A_1 when $k = 1$). The choice of a descending sequence $A_1 > \dots > A_k$ of arbitrary length is equivalent to the choice of a subset of \mathbb{N} , so that taking generating functions gives

$$(2.22) \quad \sum_{n=0}^{\infty} h_n z^n \leq \prod_{A=1}^{\infty} \left(1 + \sum_{m=1}^{\infty} b_{m,A} z^m \right).$$

Using the inequality $1 + x \leq e^x$, we obtain

$$(2.23) \quad \sum_{n=0}^{\infty} h_n z^n \leq \exp \left(\sum_{A=1}^{\infty} \sum_{m=1}^{\infty} b_{m,A} z^m \right) = e^{B(z)-1}.$$

Now using (2.14) gives

$$(2.24) \quad \begin{aligned} \chi(z) &= \sum_{n=0}^{\infty} c_n z^n \leq \frac{1}{z} \sum_{n=0}^{\infty} \sum_{m=0}^n h_{n-m} z^{n-m} h_{m+1} z^{m+1} \\ &= \frac{1}{z} \left(\sum_{n=0}^{\infty} h_n z^n \right) \left(\sum_{n=1}^{\infty} h_n z^n \right) \\ &\leq \frac{1}{z} e^{2(B(z)-1)}, \end{aligned}$$

as required. \square

2.2. Self-avoiding polygons. A $2n$ -step *self-avoiding return* is a walk $\omega \in \mathcal{W}_{2n}$ with $\omega(2n) = \omega(0) = 0$ and with $\omega(i) \neq \omega(j)$ for distinct pairs i, j other than the pair $0, 2n$. A *self-avoiding polygon* is a self-avoiding return with both the orientation and the location of the origin forgotten. Thus we can count self-avoiding polygons by counting self-avoiding returns up to orientation and translation invariance, and their number is

$$(2.25) \quad q_{2n} = \frac{2dc_{2n-1}(e_1)}{2 \cdot 2n}, \quad n \geq 2,$$

where $e_1 = (1, 0, \dots, 0)$ is the first standard basis vector. Here, the 2 in the denominator cancels the choice of orientation, and the $2n$ cancels the choice of origin in the polygon.

We first observe that two self-avoiding polygons can be concatenated to form a larger self-avoiding polygon. Consider first the case of $d = 2$. The procedure is as in Figure 3, namely we join a “rightmost” bond of one polygon to a “leftmost” bond of the other. This shows that for even integers $m, n \geq 4$, and for $d = 2$,




FIGURE 3. Concatenation of a 10-step polygon and a 14-step polygon to produce a 24-step polygon in \mathbb{Z}^2 .

$q_m q_n \leq q_{m+n}$. With a little thought (see [57] for details), in general dimensions $d \geq 2$ one obtains

$$(2.26) \quad \frac{q_m q_n}{d-1} \leq q_{m+n},$$

and if we set $q_2 = 1$ and make the easy observation that $q_n \leq q_{n+2}$, then (2.26) holds for all even $m, n \geq 2$. It follows from (2.26) that

$$(2.27) \quad q_{2n}^{1/2n} \rightarrow \mu_{\text{Polygon}} \leq \mu, \quad q_{2n} \leq \mu_{\text{Polygon}}^{2n} \leq \mu^{2n} \quad \text{for all } n \geq 2.$$

THEOREM 2.9. *There is a constant $K = K(d)$ such that, for all $n \geq 1$,*

$$(2.28) \quad c_{2n+1}(e_1) \geq \frac{K}{n^{d+2}} b_n^2.$$

PROOF. We first show the inequality

$$(2.29) \quad \sum_{x \in \mathbb{Z}^d} b_n(x)^2 \leq 2d(n+1)^2 c_{2n+1}(e_1)$$

where $b_n(x)$ denotes the number of n -step bridges ending at x . The proof is illustrated in Figure 4. Namely, given n -step bridges ω and v with $\omega(n) = v(n) = x \in \mathbb{Z}^d$, let $\mathbf{v} \in \mathbb{R}^d$ be some non-zero vector orthogonal to x , and fix some unit direction $u \in \mathbb{Z}^d$ with $u \cdot \mathbf{v} > 0$. Let $i \in \{0, 1, \dots, n\}$ be the smallest index maximising $\omega(i) \cdot \mathbf{v}$ and $j \in \{0, 1, \dots, n\}$ the smallest index minimising $v(j) \cdot \mathbf{v}$. Split ω into the pieces before and after i and interchange them to produce a walk $\bar{\omega}$, as in Figure 4(b). Do the same for v and j . Finally combine $\bar{\omega}$ and \bar{v} with an inserted step u to produce a SAW ρ with $\rho(2n+1) = u$, as in Figure 4(c). The resulting map $(\omega, v) \mapsto (\rho, i, j)$ is one-to-one, which proves (2.29).

Now, applying the Cauchy-Schwarz inequality to (2.29) gives

$$(2.30) \quad \begin{aligned} b_n^2 &= \left(\sum_{x \in \mathbb{Z}^d} b_n(x) 1_{\{b_n(x) \neq 0\}} \right)^2 \leq \sum_{x \in \mathbb{Z}^d} b_n(x)^2 \sum_{x \in \mathbb{Z}^d} 1_{\{b_n(x) \neq 0\}} \\ &\leq n(2n+1)^{d-1} \sum_{x \in \mathbb{Z}^d} b_n(x)^2. \end{aligned}$$

Thus $2dc_{2n+1}(e_1) \geq \frac{b_n^2}{n(n+1)^2(2n+1)^{d-1}}$, which completes the proof. \square

COROLLARY 2.10. *There is a $C > 0$ such that*

$$(2.31) \quad \mu^{2n} e^{-C\sqrt{n}} \leq c_{2n+1}(e_1) \leq (n+1)\mu^{2n+2}.$$

In particular, $\mu_{\text{Polygon}} = \mu$.

PROOF. The lower bound follows from Theorem 2.9 and Corollary 2.7. The upper bound follows from (2.25) and (2.27) (using $d \geq 2$). \square




FIGURE 4. Proof of Theorem 2.9. Here $n = 12$. (a) The n -step bridges ω and v , and the vector \mathbf{v} . (b) The derived walks $\bar{\omega}$ and \bar{v} . (c) The $(2n + 1)$ -step walk ρ ; here $u = (1, 0)$. The shaded lines are the hyperplanes orthogonal to \mathbf{v} .

With a little more work, it can be shown that for any fixed $x \neq 0$, $c_n(x)^{1/n} \rightarrow \mu$ as $n \rightarrow \infty$ along the subsequence of integers whose parity agrees with $\|x\|_1$. The details can be found in [57]. Thus the radius of convergence of the two-point function $G_z(x) = \sum_{n=0}^{\infty} c_n(x)z^n$ is equal to $z_c = 1/\mu$ for all x .

3. The connective constant on the hexagonal lattice

Throughout this section, we consider self-avoiding walks on the hexagonal lattice \mathbb{H} . Our first and primary goal is to prove the following theorem from [24]. The proof makes use of a certain observable of broader significance, and following the proof we discuss this in the context of the $O(n)$ models.

THEOREM 3.1. *For the hexagonal lattice \mathbb{H} ,*

$$(3.1) \quad \mu = \sqrt{2 + \sqrt{2}}.$$

As a matter of convenience, we extend walks at their extremities by two half-edges in such a way that they start and end at *mid-edges*, i.e., centres of edges of \mathbb{H} . The set of mid-edges will be called H . We position the hexagonal lattice \mathbb{H} of mesh size 1 in \mathbb{C} so that there exists a horizontal edge e with mid-edge a being 0. We now write c_n for the number of n -step SAWs on the hexagonal lattice \mathbb{H} which start at 0, and $\chi(z) = \sum_{n=0}^{\infty} c_n z^n$ for the susceptibility.

We first point out that it suffices to count bridges. On the hexagonal lattice, a bridge is defined by the following adaptation of Definition 2.1: a *bridge* on \mathbb{H} is a SAW which never revisits the vertical line through its starting point, never visits a vertical line to the right of the vertical line through its endpoint, and moreover starts and ends at the midpoint of a horizontal edge. We now use b_n to denote the number of n -step bridges on \mathbb{H} which start at 0. It is straightforward to adapt the arguments used to prove Corollary 2.7 to the hexagonal lattice, leading to the conclusion that $\mu_{\text{Bridge}} = \mu$ also on \mathbb{H} . Thus it suffices to show that

$$(3.2) \quad \mu_{\text{Bridge}} = \sqrt{2 + \sqrt{2}}.$$

Using notation which anticipates our conclusion but which should not create confusion, we will write

$$(3.3) \quad z_c = \frac{1}{\sqrt{2 + \sqrt{2}}}.$$

We also write $B(z) = \sum_{n=0}^{\infty} b_n z^n$ for $z > 0$. To prove (3.2), it suffices to prove that $B(z_c) = \infty$ or $\chi(z_c) = \infty$, and that $B(z) < \infty$ whenever $z < z_c$. This is what we will prove.

3.1. The holomorphic observable. The proof is based on a generalisation of the two-point function that we call the *holomorphic observable*. In this section, we introduce the holomorphic observable and prove its discrete analyticity. Some preliminary definitions are required.

A *domain* $\Omega \subset H$ is a union of all mid-edges emanating from a given connected collection of vertices $V(\Omega)$; see Figure 5. In other words, a mid-edge x belongs to Ω if at least one end-point of its associated edge is in $V(\Omega)$. The boundary $\partial\Omega$ consists of mid-edges whose associated edge has exactly one endpoint in Ω . We further assume Ω to be simply connected, i.e., having a connected complement.

DEFINITION 3.2. The winding $W_\omega(a, b)$ of a SAW ω between mid-edges a and b (not necessarily the start and end of ω) is the total rotation in radians when ω is traversed from a to b ; see Figure 5.

We write $\omega : a \rightarrow E$ if a walk ω starts at mid-edge a and ends at some mid-edge of $E \subset H$. In the case where $E = \{b\}$, we simply write $\omega : a \rightarrow b$. The *length* $\ell(\omega)$ of the walk is the number of vertices belonging to ω . The following definition provides a generalisation of the two-point function $G_z(x)$.

DEFINITION 3.3. Fix $a \in \partial\Omega$ and $\sigma \in \mathbb{R}$. For $x \in \Omega$ and $z \geq 0$, the *holomorphic observable* is defined to be

$$(3.4) \quad F_z(x) = \sum_{\omega \subset \Omega: a \rightarrow x} e^{-i\sigma W_\omega(a, x)} z^{\ell(\omega)}.$$

In contrast to the two-point function, the weights in the holomorphic observable need not be positive. For the special case $z = z_c$ and $\sigma = \frac{5}{8}$, F_{z_c} satisfies the relation




FIGURE 5. Left: A domain Ω whose boundary mid-edges are pictured by small black squares. Vertices of $V(\Omega)$ correspond to circles. Right: Winding of a SAW ω .

in the following lemma, a relation which can be regarded as a weak form of discrete analyticity, and which will be crucial in the rest of the proof.

LEMMA 3.4. *If $z = z_c$ and $\sigma = \frac{5}{8}$, then, for every vertex $v \in V(\Omega)$,*

$$(3.5) \quad (p - v)F_{z_c}(p) + (q - v)F_{z_c}(q) + (r - v)F_{z_c}(r) = 0,$$

where p, q, r are the mid-edges of the three edges adjacent to v .

PROOF. Let $z \geq 0$ and $\sigma \in \mathbb{R}$. We will specialise later to $z = z_c$ and $\sigma = \frac{5}{8}$. We assume without loss of generality that p, q and r are oriented counter-clockwise around v . By definition, $(p - v)F_z(p) + (q - v)F_z(q) + (r - v)F_z(r)$ is a sum of contributions $c(\omega)$ over all possible SAWs ω ending at p, q or r . For instance, if ω ends at the mid-edge p , then its contribution will be

$$(3.6) \quad c(\omega) = (p - v)e^{-i\sigma W_\omega(a,p)} z^{\ell(\omega)}.$$

The set of walks ω finishing at p, q or r can be partitioned into pairs and triplets of walks as depicted in Figure 6, in the following way:

- If a SAW ω_1 visits all three mid-edges p, q, r , then the edges belonging to ω_1 form a SAW plus (up to a half-edge) a self-avoiding return from v to v . One can associate to ω_1 the walk ω_2 passing through the same edges, but traversing the return from v to v in the opposite direction. Thus, walks visiting the three mid-edges can be grouped in pairs.
- If a walk ω_1 visits only one mid-edge, it can be associated to two walks ω_2 and ω_3 that visit exactly two mid-edges by prolonging the walk one step further (there are two possible choices). The reverse is true: a walk visiting exactly two mid-edges is naturally associated to a walk visiting only one mid-edge by erasing the last step. Thus, walks visiting one or two mid-edges can be grouped in triplets.

We will prove that when $\sigma = \frac{5}{8}$ and $z = z_c$ the sum of contributions for each pair and each triplet vanishes, and therefore the total sum is zero.




FIGURE 6. Left: a pair of walks visiting the three mid-edges and matched together. Right: a triplet of walks, one visiting one mid-edge, the two others visiting two mid-edges, which are matched together.

Let ω_1 and ω_2 be two walks that are grouped as in the first case. Without loss of generality, we assume that ω_1 ends at q and ω_2 ends at r . Note that ω_1 and ω_2 coincide up to the mid-edge p since (ω_1, ω_2) are matched together. Then

$$(3.7) \quad \ell(\omega_1) = \ell(\omega_2) \quad \text{and} \quad \begin{cases} W_{\omega_1}(a, q) = W_{\omega_1}(a, p) + W_{\omega_1}(p, q) = W_{\omega_1}(a, p) - \frac{4\pi}{3} \\ W_{\omega_2}(a, r) = W_{\omega_2}(a, p) + W_{\omega_2}(p, r) = W_{\omega_1}(a, p) + \frac{4\pi}{3}. \end{cases}$$

In evaluating the winding of ω_1 between p and q , we used the fact that $a \in \partial\Omega$ and Ω is simply connected. The term $e^{-i\sigma W_{\omega}(a, x)}$ gives a weight λ or $\bar{\lambda}$ per left or right turn of ω , where

$$(3.8) \quad \lambda = \exp\left(-i\sigma \frac{\pi}{3}\right).$$

Writing $j = e^{i2\pi/3}$, we obtain

$$(3.9) \quad \begin{aligned} c(\omega_1) + c(\omega_2) &= (q - v)e^{-i\sigma W_{\omega_1}(a, q)} z^{\ell(\omega_1)} + (r - v)e^{-i\sigma W_{\omega_2}(a, r)} z^{\ell(\omega_2)} \\ &= (p - v)e^{-i\sigma W_{\omega_1}(a, p)} z^{\ell(\omega_1)} (j\bar{\lambda}^4 + \bar{j}\lambda^4). \end{aligned}$$

Now we set $\sigma = \frac{5}{8}$ so that $j\bar{\lambda}^4 + \bar{j}\lambda^4 = 2\cos(\frac{3\pi}{2}) = 0$, and hence

$$(3.10) \quad c(\omega_1) + c(\omega_2) = 0.$$

Let $\omega_1, \omega_2, \omega_3$ be three walks matched as in the second case. Without loss of generality, we assume that ω_1 ends at p and that ω_2 and ω_3 extend ω_1 to q and r respectively. As before, we easily find that

$$(3.11) \quad \ell(\omega_2) = \ell(\omega_3) = \ell(\omega_1) + 1 \quad \text{and} \quad \begin{cases} W_{\omega_2}(a, r) = W_{\omega_2}(a, p) + W_{\omega_2}(p, q) = W_{\omega_1}(a, p) - \frac{\pi}{3} \\ W_{\omega_3}(a, r) = W_{\omega_3}(a, p) + W_{\omega_3}(p, r) = W_{\omega_1}(a, p) + \frac{\pi}{3}, \end{cases}$$

and thus

$$(3.12) \quad c(\omega_1) + c(\omega_2) + c(\omega_3) = (p - v)e^{-i\sigma W_{\omega_1}(a, p)} z^{\ell(\omega_1)} (1 + zj\bar{\lambda} + z\bar{j}\lambda).$$

Now we choose z such that $1 + zj\bar{\lambda} + z\bar{j}\lambda = 0$. Due to our choice $\sigma = \frac{5}{8}$, we have $\lambda = \exp(-i\frac{5\pi}{24})$. Thus we choose $z_c^{-1} = 2\cos\frac{\pi}{8} = \sqrt{2 + \sqrt{2}}$.

Now the desired identity (3.5) follows immediately by summing over all the pairs and triplets of walks. \square

The last step of the proof of Lemma 3.4 is the *only* place where the choice $z = z_c = 1/\sqrt{2 + \sqrt{2}}$ is used in the proof of Theorem 3.1.

3.2. Proof of Theorem 3.1 completed. Now we will apply Lemma 3.4 to prove Theorem 3.1.

We consider a vertical strip domain S_T composed of the vertices of T strips of hexagons, and its finite version $S_{T,L}$ cut at height L at an angle of $\frac{\pi}{3}$; see Figure 7. We denote the left and right boundaries of S_T by α and β , respectively, and the top and bottom boundaries of $S_{T,L}$ by ϵ and $\bar{\epsilon}$, respectively. We also introduce the positive quantities:

$$(3.13) \quad A_{T,L}(z) = \sum_{\omega \subset S_{T,L}: a \rightarrow \alpha \setminus \{a\}} z^{\ell(\omega)},$$

$$(3.14) \quad B_{T,L}(z) = \sum_{\omega \subset S_{T,L}: a \rightarrow \beta} z^{\ell(\omega)},$$

$$(3.15) \quad E_{T,L}(z) = \sum_{\omega \subset S_{T,L}: a \rightarrow \epsilon \cup \bar{\epsilon}} z^{\ell(\omega)}.$$




FIGURE 7. Domain $S_{T,L}$ and boundary parts α , β , ϵ and $\bar{\epsilon}$.

LEMMA 3.5. For $z = z_c$,

$$(3.16) \quad 1 = c_\alpha A_{T,L}(z_c) + B_{T,L}(z_c) + c_\epsilon E_{T,L}(z_c),$$

where $c_\alpha = \cos\left(\frac{3\pi}{8}\right)$ and $c_\epsilon = \cos\left(\frac{\pi}{4}\right)$.

PROOF. We fix $z = z_c$ and drop it from the notation. We sum the relation (3.5) over all vertices in $V(S_{T,L})$. Contributions at interior mid-edges vanish and we arrive at

$$(3.17) \quad - \sum_{x \in \alpha} F(x) + \sum_{x \in \beta} F(x) + j \sum_{x \in \epsilon} F(x) + \bar{j} \sum_{x \in \bar{\epsilon}} F(x) = 0.$$

The winding of any SAW from a to the bottom part of α is $-\pi$, while the winding to the top part is π . Using this and symmetry, together with the fact that the only SAW from a to a has length 0, we conclude that

$$(3.18) \quad \sum_{x \in \alpha} F(x) = F(a) + \sum_{x \in \alpha \setminus \{a\}} F(x) = 1 + \frac{e^{-i\sigma\pi} + e^{i\sigma\pi}}{2} A_{T,L} = 1 - c_\alpha A_{T,L}.$$

Similarly, the winding from a to any half-edge in β , ϵ or $\bar{\epsilon}$ is respectively 0, $\frac{2\pi}{3}$ or $-\frac{2\pi}{3}$. Therefore, again using symmetry,

$$(3.19) \quad \sum_{x \in \beta} F(x) = B_{T,L}, \quad j \sum_{x \in \epsilon} F(x) + \bar{j} \sum_{x \in \bar{\epsilon}} F(x) = c_\epsilon E_{T,L}.$$

The proof is completed by inserting (3.18)–(3.19) into (3.17). \square

The sequences $(A_{T,L}(z))_{L>0}$ and $(B_{T,L}(z))_{L>0}$ are increasing in L and are bounded for $z \leq z_c$, thanks to (3.16) and the monotonicity in z . Thus they have limits

$$(3.20) \quad A_T(z) = \lim_{L \rightarrow \infty} A_{T,L}(z) = \sum_{\omega \subset S_T: a \rightarrow \alpha \setminus \{a\}} z^{\ell(\omega)},$$

$$(3.21) \quad B_T(z) = \lim_{L \rightarrow \infty} B_{T,L}(z) = \sum_{\omega \subset S_T: a \rightarrow \beta} z^{\ell(\omega)}.$$

When $z = z_c$, via (3.16) again, we conclude that $(E_{T,L}(z_c))_{L>0}$ is decreasing and converges to a limit $E_T(z_c) = \lim_{L \rightarrow \infty} E_{T,L}(z_c)$. Thus, by (3.16),

$$(3.22) \quad 1 = c_\alpha A_T(z_c) + B_T(z_c) + c_\epsilon E_T(z_c).$$

PROOF OF THEOREM 3.1. The bridge generating function is given by $B(z) = \sum_{T=0}^{\infty} B_T(z)$. Recall that it suffices to show that $B(z) < \infty$ for $z < z_c$, and that $B(z_c) = \infty$ or $\chi(z_c) = \infty$.

We first assume $z < z_c$. Since $B_T(z)$ involves only bridges of length at least T , it follows from (3.22) that

$$(3.23) \quad B_T(z) \leq \left(\frac{z}{z_c}\right)^T B_T(z_c) \leq \left(\frac{z}{z_c}\right)^T,$$

and hence $B(z)$ is finite since the right-hand side is summable.

It remains to prove that $B(z_c) = \infty$ or $\chi(z_c) = \infty$. We do this by considering two separate cases. Suppose first that, for some T , $E_T(z_c) > 0$. As noted previously, $E_{T,L}(z_c)$ is decreasing in L . Therefore, as required,

$$(3.24) \quad \chi(z_c) \geq \sum_{L=1}^{\infty} E_{T,L}(z_c) \geq \sum_{L=1}^{\infty} E_T(z_c) = \infty.$$

It remains to consider the case that $E_T^{z_c} = 0$ for every T . In this case, (3.22) simplifies to

$$(3.25) \quad 1 = c_\alpha A_T(z_c) + B_T(z_c).$$

Observe that walks contributing to $A_{T+1}(z_c)$ but not to $A_T(z_c)$ must visit some vertex adjacent to the right edge of S_{T+1} . Cutting such a walk at the first such point (and adding half-edges to the two halves), we obtain two bridges of span $T+1$ in S_{T+1} . We conclude from this that

$$(3.26) \quad A_{T+1}(z_c) - A_T(z_c) \leq z_c (B_{T+1}(z_c))^2.$$

Combining (3.25) for T and $T + 1$ with (3.26), we can write

$$(3.27) \quad \begin{aligned} 0 &= [c_\alpha A_{T+1}(z_c) + B_{T+1}(z_c)] - [c_\alpha A_T(z_c) + B_T(z_c)] \\ &\leq c_\alpha z_c (B_{T+1}(z_c))^2 + B_{T+1}(z_c) - B_T(z_c), \end{aligned}$$

so

$$(3.28) \quad c_\alpha z_c (B_{T+1}(z_c))^2 + B_{T+1}(z_c) \geq B_T(z_c).$$

It is an easy exercise to verify by induction that

$$(3.29) \quad B_T(z_c) \geq \min\{B_1(z_c), 1/(c_\alpha z_c)\} \frac{1}{T}$$

for every $T \geq 1$. This implies, as required, that

$$(3.30) \quad B(z_c) \geq \sum_{T=1}^{\infty} B_T(z_c) = \infty.$$

This completes the proof. \square

3.3. Conjecture 1.5 and the holomorphic observable. Recall the statement of Conjecture 1.5. When formulated on \mathbb{H} , this conjecture concerns a simply connected domain Ω in the complex plane \mathbb{C} with two points a and b on the boundary, with a discrete approximation given by the largest finite domain Ω_δ of $\delta\mathbb{H}$ included in Ω , and with a_δ and b_δ the closest vertices of $\delta\mathbb{H}$ to a and b respectively. A probability measure $\mathbb{P}_{z,\delta}$ is defined on the set of SAWs ω between a_δ and b_δ that remain in Ω_δ by assigning to ω a weight proportional to $z_c^{\ell(\omega)}$. We obtain a random curve denoted ω_δ . We can also define the observable in this context, and we denote it by F_δ . Conjecture 1.5 then asserts that the random curve ω_δ converges to $\text{SLE}_{8/3}$ from a and b in the domain Ω .

A possible approach to proving Conjecture 1.5 might be the following. First, prove a precompactness result for self-avoiding walks. Then, by taking a subsequence, we could assume that the curve γ_δ converges to a continuous curve (in fact, the limiting object would need to be a Loewner chain, see [1]). The second step would consist in identifying the possible limits. The holomorphic observable should play a crucial role in this step. Indeed, if F_δ converges when rescaled to an explicit function, one could use the *martingale technique* introduced in [70] to verify that the only possible limit is $\text{SLE}_{8/3}$.

Regarding the convergence of F_δ , we first recall that in the discrete setting contour integrals should be performed along dual edges. For \mathbb{H} , the dual edges form a triangular lattice, and Lemma 3.4 has the enlightening interpretation that the contour integral vanishes along any elementary dual triangle. Any area enclosed by a discrete closed dual contour is a union of elementary triangles, and hence the integral along any discrete closed contour also vanishes. This is a discrete analogue of Morera's theorem. It implies that if the limit of F_δ (properly rescaled) exists and is continuous, then it is automatically holomorphic. By studying the boundary conditions, it is even possible to identify the limit. This leads to the following conjecture, which is based on ideas in [70].

CONJECTURE 3.6. *Let Ω be a simply connected domain (not equal to \mathbb{C}), let $z \in \Omega$, and let a, b be two distinct points on the boundary of Ω . We assume that the boundary of Ω is smooth near b . For $\delta > 0$, let F_δ be the holomorphic observable*

in the domain $(\Omega_\delta, a_\delta, b_\delta)$ approximating (Ω, a, b) , and let z_δ be the closest point in Ω_δ to z . Then

$$(3.31) \quad \lim_{\delta \rightarrow 0} \frac{F_\delta(a_\delta, z_\delta)}{F_\delta(a_\delta, b_\delta)} = \left(\frac{\Phi'(z)}{\Phi'(b)} \right)^{5/8},$$

where Φ is a conformal map from Ω to the upper half-plane mapping a to ∞ and b to 0 .

The right-hand side of (3.31) is well-defined, since the conformal map Φ is unique up to multiplication by a real factor.

3.4. Loop models and holomorphic observables. The original motivation for the introduction of the holomorphic observable stems from a more general context, which we now discuss. The *loop $O(n)$ model* is a lattice model on a domain Ω . We restrict attention in this discussion to the hexagonal lattice \mathbb{H} . A configuration ω is a family of self-avoiding loops, and its probability is proportional to $z^{\#\text{edges}} n^{\#\text{loops}}$. The *loop parameter* n is taken in $[0, 2]$. There are other variants of the model, for instance, one can introduce an interface going from one point a on the boundary to the inside, or one interface between two points of the boundary. The case $n = 1$ corresponds to the Ising model, while the case $n = 0$ corresponds to the self-avoiding walk (when allowing one interface).

Fix $n \in [0, 2]$. It is a non-rigorous prediction of [61] that the model has the following three phases distinguished by the value of z :

- If $z < 1/\sqrt{2 + \sqrt{2 - n}}$, the loops are sparse (typically of logarithmic size in the size of the domain). This phase is subcritical.
- If $z = 1/\sqrt{2 + \sqrt{2 - n}}$, the loops are dilute (there are loops of the size of the domain which are typically separated by a distance of the size of the domain). This phase is critical.
- If $z > 1/\sqrt{2 + \sqrt{2 - n}}$, the loops are dense (there are loops of the size of the domain which are typically separated by a distance much smaller than the size of the domain). This phase is critical as well.

Consider the special case of the Ising model at its critical value $z_c = 1/\sqrt{3}$. Let E denote the set of configurations consisting only of self-avoiding loops, and let $E(a, x)$ denote the set of configurations with self-avoiding loops plus an interface γ from a to x . Then, ignoring the issue of boundary conditions, the Ising spin-spin correlation is given in terms of the loop model by

$$(3.32) \quad \langle \sigma(a)\sigma(x) \rangle = \frac{\sum_{\omega \in E(a, x)} z_c^{\#\text{edges}}}{\sum_{\omega \in E} z_c^{\#\text{edges}}}.$$

A natural operation in physics consists in flipping the sign of the coupling constant of the Ising model along a path from a to x , in such a way that a monodromy is introduced: if we follow a path turning around x , spins are reversed after one whole turn. See, e.g., [65]. In terms of the loop representation, the spin-spin correlation $\langle \sigma(a)\sigma(x) \rangle_{\text{monodromy}}$ in this new Ising model is

$$(3.33) \quad \langle \sigma(a)\sigma(x) \rangle_{\text{monodromy}} = \frac{\sum_{\omega \in E(a, x)} (-1)^{\#\text{turns of } \gamma \text{ around } x} z_c^{\#\text{edges}}}{\sum_{\omega \in E} z_c^{\#\text{edges}}}$$

where γ is the interface between a and x .

The numerator of the right-hand side of (3.33) can be rewritten as

$$(3.34) \quad \sum_{\omega \in E(a,x)} e^{-i\frac{1}{2}W_\gamma(a,x)} z^{\#\text{edges}} n^{\#\text{loops}}$$

with $n = 1$. This is of the same form as the holomorphic observable (3.4). With general values of n , and with the freedom to choose the value of $\sigma \in [0, 1]$, we obtain the observable

$$(3.35) \quad F_z(x) = \sum_{\omega \in E(a,x)} e^{-i\sigma W_\gamma(a,x)} z^{\#\text{edges}} n^{\#\text{loops}}.$$

The values of σ and z need to be chosen according to the value of n . If $\sigma = \sigma(n)$ satisfies $2 \cos[(1 + 2\sigma)2\pi/3] = -n$ and $z = z(n) = 1/\sqrt{2 + \sqrt{2 - n}}$, then the proof of Lemma 3.4 can be modified to yield its conclusion in this more general context.

To conclude this discussion, consider the loop $O(n)$ model with a family of self-avoiding loops and a single interface between two boundary points a and b . For $n = 1$ and $z = 1/\sqrt{3}$, it has been proved that the interface converges to SLE_3 [18]. For other values of z and n , the following behaviour is conjectured [70].

CONJECTURE 3.7. Fix $n \in [0, 2]$. For $z = 1/\sqrt{2 + \sqrt{2 - n}}$, the interface between a and b converges, as the lattice spacing goes to zero, to

$$(3.36) \quad \text{SLE}_\kappa \quad \text{with} \quad \kappa = \frac{4\pi}{2\pi - \arccos(-n/2)}.$$

For $z > 1/\sqrt{2 + \sqrt{2 - n}}$, the interface between a and b converges, as the lattice spacing goes to zero, to

$$(3.37) \quad \text{SLE}(\kappa) \quad \text{with} \quad \kappa = \frac{4\pi}{\arccos(-n/2)}.$$

Conjecture 3.7 is summarised in Figure 8. The value of \arccos is in $[0, \pi]$, so the first regime corresponds to $\kappa \in [\frac{8}{3}, 4]$ and the second to $\kappa \in [4, 8]$. These two critical regimes do not belong to the same universality class, in the sense that the scaling




FIGURE 8. Phase diagram for $O(n)$ models.

limit of the interface is not the same. In particular, since SLE_κ curves are simple for $\kappa \leq 4$ but not for $\kappa > 4$ (see [1]), in the dilute phase the interface is conjectured to be simple in the scaling limit, but not in the dense phase. In addition, all the SLE_κ models for $\frac{8}{3} \leq \kappa \leq 8$ arise in these $O(n)$ models. This rich behaviour is at the heart of the mathematical interest in $O(n)$ models. To prove the conjecture remains a major challenge in 2-dimensional statistical mechanics.

4. The lace expansion

4.1. Main results. In dimensions $d \geq 5$, it has been proved that SAW has the same scaling behaviour as SRW. The following two theorems, due to Hara and Slade [33, 34] and to Hara [30], respectively, show that the critical exponents γ, ν, η exist and take the values $\gamma = 1, \nu = \frac{1}{2}, \eta = 0$, and that the scaling limit is Brownian motion.

THEOREM 4.1. *Fix $d \geq 5$, and consider the nearest-neighbour SAW on \mathbb{Z}^d . There exist constants $A, D, \epsilon > 0$ such that, as $n \rightarrow \infty$,*

$$(4.1) \quad c_n = A\mu^n[1 + O(n^{-\epsilon})],$$

$$(4.2) \quad \mathbb{E}_n |\omega(n)|^2 = Dn[1 + O(n^{-\epsilon})].$$

Also,

$$(4.3) \quad \left(\frac{\omega(\lfloor nt \rfloor)}{\sqrt{Dn}} \right)_{t \geq 0} \rightarrow (B_t)_{t \geq 0},$$

where B_t denotes Brownian motion and the convergence is in distribution.

THEOREM 4.2. *Fix $d \geq 5$, and consider the nearest-neighbour SAW on \mathbb{Z}^d . There are constants $c, \epsilon > 0$ such that, as $x \rightarrow \infty$,*

$$(4.4) \quad G_{z_c}(x) = \frac{c}{|x|^{d-2}} \left[1 + O(|x|^{-\epsilon}) \right].$$

The proofs are based on the lace expansion, a technique that was introduced by Brydges and Spencer [14] to study the weakly SAW in dimensions $d > 4$. Since 1985, the method of lace expansion has been highly developed and extended to several other models: percolation ($d > 6$), oriented percolation ($d > 4$ spatial dimensions), the contact process ($d > 4$), lattice trees and lattice animals ($d > 8$), the Ising model ($d > 4$), and to random subgraphs of high-dimensional transitive graphs such as the Boolean cube. For a review and references, see [69].

Versions of Theorems 4.1–4.2 have been proved also for spread-out models; see [57, 31]. More recently, the above two theorems have been extended also to study long-range SAWs based on simple random walks which take steps of length r with probability proportional to $r^{-d-\alpha}$ for some α . For $\alpha \in (0, 2)$, the upper critical dimension (recall Section 1.6.4) is reduced from 4 to 2α , and the Brownian limit is replaced by a stable law in dimensions $d > 2\alpha$ [38]. Further results in this direction can be found in [39, 19].

Our goal now is modest. In this section, we will derive the lace expansion. In Section 5, we will sketch a proof of how it can be used to prove that $\gamma = 1$, in the sense that

$$(4.5) \quad \chi(z) \asymp (1 - z/z_c)^{-1} \quad \text{as } z \nearrow z_c,$$

both for the nearest-neighbour model with $d \geq d_0 \gg 4$, and for the spread-out model with $L \geq L_0(d) \gg 1$ and any $d > 4$. Here, the notation $f(z) \asymp g(z)$ means that there exist positive c_1, c_2 such that $c_1 g(z) \leq f(z) \leq c_2 g(z)$ holds uniformly in z . The lower bound in (4.5) holds in all dimensions and follows immediately from the elementary observation in (1.12) that $c_n \geq \mu^n = z_c^{-n}$, since

$$(4.6) \quad \chi(z) = \sum_{n=0}^{\infty} c_n z^n \geq \sum_{n=0}^{\infty} (\mu z)^n = \frac{1}{1 - z/z_c}$$

for $z < z_c$. It therefore suffices to prove that in high dimensions we have the complementary upper bound

$$(4.7) \quad \chi(z) \leq \frac{C}{1 - z/z_c}$$

for some finite constant C .

4.2. The differential inequality for $\chi(z)$. We prove (4.7) by means of a *differential inequality*—an inequality relating $\frac{d}{dz}\chi(z)$ to $\chi(z)$. The derivation of the differential inequality and its implication for (4.7) first appeared in [3].

The differential inequality is expressed in terms of the quantity

$$(4.8) \quad \mathbf{B}(z) = \sum_{x \in \mathbb{Z}^d} G_z(x)^2$$

for $z \leq z_c$. Proposition 1.3 ensures that $\mathbf{B}(z)$ is finite for $z < z_c$. If we assume, as usual, that $G_{z_c} \sim c|x|^{-(d-2+\eta)}$, then $\mathbf{B}(z_c)$ will be finite precisely when $d > 4 - 2\eta$. With Fisher's relation (1.40) and the predicted values of γ and ν from (1.22) and (1.28), this inequality can be expected to hold, and correspondingly $\mathbf{B}(z_c) < \infty$, only for $d > 4$ (this is a prediction, not a theorem). We refer to $\mathbf{B}(z)$ as the *bubble diagram* because we express (4.8) diagrammatically as

$$(4.9) \quad \mathbf{B}(z) = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \quad \quad \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array}$$


In this diagram, each line represents a factor $G_z(x)$ and the unlabelled vertex is summed over $x \in \mathbb{Z}^d$. The condition that $\mathbf{B}(z_c) < \infty$ will be referred to as the *bubble condition*.

We now derive the differential inequality

$$(4.10) \quad \frac{d}{dz} (z\chi(z)) \geq \frac{\chi(z)^2}{\mathbf{B}(z)}.$$

Assuming (4.10), we obtain (4.7) as if we were solving a differential equation. Namely, using the monotonicity of \mathbf{B} , we first replace $\mathbf{B}(z)$ by $\mathbf{B}(z_c)$ in (4.10). We then rearrange and integrate from z to z_c , using the terminal value $\chi(z_c) = \infty$

inequality by relaxing the avoidance pattern to [14], keeping the requirement that the walk 23 should be non-empty:



[14]

$$(4.15) \quad Q(z) \geq \chi(z)^2 - Q(z)(B(z) - 1).$$

Rearranging gives the inequality (4.10).

4.3. The lace expansion by inclusion-exclusion. The proof of the bubble condition is based on the lace expansion. The original derivation of the lace expansion by Brydges and Spencer [14] made use of a certain graphical construction called a *lace*. Later, it was realised that repeated inclusion-exclusion leads to the same expansion [68]. We present the inclusion-exclusion approach now; the approach via laces is treated in the problems of Section 4.4. The underlying graph plays little role in the derivation, and the following discussion pertains to either nearest-neighbour or spread-out SAWs. Indeed, with minor modifications, the discussion also applies on general graphs [22].

We use the convolution $(f * g)(x) = \sum_{y \in \mathbb{Z}^d} f(y)g(x - y)$ of two functions f, g on \mathbb{Z}^d . The lace expansion gives rise to a formula for $c_n(x)$, for $n \geq 1$, of the form

$$(4.16) \quad \begin{aligned} c_n(x) &= (c_1 * c_{n-1})(x) + \sum_{m=2}^n (\pi_m * c_{n-m})(x) \\ &= \sum_{y \in \mathbb{Z}^d} c_1(y)c_{n-1}(x - y) + \sum_{m=2}^n \sum_{y \in \mathbb{Z}^d} \pi_m(y)c_{n-m}(x - y), \end{aligned}$$

in which the coefficients $\pi_m(y)$ are certain combinatorial integers that we will define below. Note that the identity (4.16) would hold for SRW with $\pi \equiv 0$. The quantity $\pi_m(y)$ can therefore be understood as a correction factor determining to what degree SAWs fail to behave like SRWs. In this sense, the lace expansion studies the SAW as a perturbation of the SRW.

Our starting point is similar to that of the derivation of the differential inequality (4.10), but now we will work with identities rather than inequalities. Also, rather than working with generating functions, we will work instead with walks with a fixed number of steps and without factors z : diagrams now arise from walks of fixed length. We begin by dividing an n -step SAW ($n \geq 1$) into its first step and the remainder of the walk. Because of self-avoidance, these two parts must be

as

$$\begin{aligned}
\pi_m(y) &= - \begin{array}{c} \text{loop} \\ 0 \end{array} \delta_{0y} + \begin{array}{c} 1 \quad y \\ \text{diagram} \\ 0 \quad 3 \\ [123] \end{array} - \begin{array}{c} 1 \quad 5 \\ \text{diagram} \\ 0 \quad 3 \quad y \\ [1234][345] \end{array} + \dots \\
(4.21) \quad &= \sum_{N=1}^{\infty} (-1)^N \pi_m^{(N)}(y),
\end{aligned}$$

where $\delta_{xy} = 1_{\{x=y\}}$ denotes the Kronecker delta, then (4.20) becomes (4.16), namely

$$(4.22) \quad c_n(x) = \sum_{y \in \mathbb{Z}^d} c_1(y) c_{n-1}(x-y) + \sum_{m=2}^n \sum_{y \in \mathbb{Z}^d} \pi_m(y) c_{n-m}(x-y).$$

By definition, $\pi_m^{(1)}(y)$ counts the number of m -step self-avoiding returns if $y = 0$, and is otherwise 0. Also, $\pi_m^{(2)}(y)$ counts the number of m -step “ θ -diagrams” with vertices 0 and y , i.e., the number of m -step walks which start at zero, end at y , and are self-avoiding apart from a required return to 0 and a visit to y before terminating at y . With more attention to the inclusion-exclusion procedure, it can be seen that in the three diagrams on the right-hand side of (4.21) all the individual subwalks must have length at least 1 except for subwalk 3 of the third term which may have length 0. As noted above, the inclusion-exclusion procedure terminates after a finite number of steps, so the terms in the series (4.21) are eventually all zero, but as m increases more and more terms are non-zero. If the diagrams make you uncomfortable, formulas for $\pi_m(y)$ are given in Section 4.4. This completes the derivation of the lace expansion.

Our next task is to relate $\pi_m(y)$ to our goal of proving the bubble condition. Equation (4.16) contains two convolutions: a convolution in space given by the sum over y , and a convolution in time given by the sum over m . To eliminate these and facilitate analysis, we pass to generating functions and Fourier transforms. By definition of the two-point function,

$$(4.23) \quad G_z(x) = \sum_{n=0}^{\infty} c_n(x) z^n = \delta_{0x} + \sum_{n=1}^{\infty} c_n(x) z^n,$$

and we define

$$(4.24) \quad \Pi_z(x) = \sum_{m=2}^{\infty} \pi_m(x) z^m.$$

From (4.16), we obtain

$$\begin{aligned}
(4.25) \quad G_z(x) &= \delta_{0x} + \sum_{y \in \mathbb{Z}^d} z c_1(y) G_z(x-y) + \sum_{y \in \mathbb{Z}^d} \Pi_z(y) G_z(x-y) \\
&= \delta_{0x} + z(c_1 * G_z)(x) + (\Pi_z * G_z)(x).
\end{aligned}$$

Given an absolutely summable function $f : \mathbb{Z}^d \rightarrow \mathbb{C}$, we write its Fourier transform as

$$(4.26) \quad \hat{f}(k) = \sum_{x \in \mathbb{Z}^d} f(x) e^{ik \cdot x},$$

with $k = (k_1, \dots, k_d) \in [-\pi, \pi]^d$. Then (4.25) gives

$$(4.27) \quad \hat{G}_z(k) = 1 + z\hat{c}_1(k)\hat{G}_z(k) + \hat{\Pi}_z(k)\hat{G}_z(k).$$

We solve for $\hat{G}_z(k)$ to obtain

$$(4.28) \quad \hat{G}_z(k) = \frac{1}{1 - z\hat{c}_1(k) - \hat{\Pi}_z(k)}.$$

It is convenient to express $c_1(y)$ in terms of the probability distribution for the steps of the corresponding SRW model:

$$(4.29) \quad D(y) = \frac{c_1(y)}{|\Omega|}, \quad \hat{c}_1(k) = |\Omega| \hat{D}(k),$$

where $|\Omega|$ denotes the cardinality of either option for the set Ω defined in (1.1). For the nearest-neighbour model, $|\Omega| = 2d$ and

$$(4.30) \quad \hat{D}(k) = \frac{1}{2d} \sum_{j=1}^d (e^{ik_j} + e^{-ik_j}) = \frac{1}{d} \sum_{j=1}^d \cos k_j.$$

To simplify the notation, we define $\hat{F}_z(k)$ by

$$(4.31) \quad \hat{G}_z(k) = \frac{1}{1 - z|\Omega|\hat{D}(k) - \hat{\Pi}_z(k)} = \frac{1}{\hat{F}_z(k)}.$$

Notice that $\hat{G}_z(0) = \sum_{x \in \mathbb{Z}^d} \sum_{n=0}^{\infty} c_n(x) z^n = \chi(z)$, so that $\hat{G}_z(0)$ will have a singularity at $z = z_c$. To emphasise this, we will write

$$(4.32) \quad \begin{aligned} \hat{F}_z(k) &= \hat{F}_z(0) + (\hat{F}_z(k) - \hat{F}_z(0)) \\ &= \chi(z)^{-1} + z|\Omega|(1 - \hat{D}(k)) + (\hat{\Pi}_z(0) - \hat{\Pi}_z(k)). \end{aligned}$$

Now we can make contact with our goal of proving the bubble condition. By Parseval's relation,

$$(4.33) \quad \mathbf{B}(z) = \sum_{x \in \mathbb{Z}^d} G_z(x)^2 = \int_{[-\pi, \pi]^d} |\hat{G}_z(k)|^2 \frac{d^d k}{(2\pi)^d}$$

(this includes the case where one side of the equality, and hence both, are infinite). The issue of whether $\mathbf{B}(z_c) < \infty$ or not boils down to the question of whether the singularity of the integrand is integrable or not, so we will need to understand the asymptotics of the terms in (4.32) as $k \rightarrow 0$ and $z \nearrow z_c$. In principle there could be other singularities when $z = z_c$, but for the nearest-neighbour and spread-out models $1 - \hat{D}(k) > 0$ for non-zero k , and one of the goals of the analysis will be to prove that the term $\hat{\Pi}_z(0) - \hat{\Pi}_z(k)$ cannot create a cancellation.

The term $1 - \hat{D}(k)$ is explicit, and for the nearest-neighbour model has asymptotic behaviour

$$(4.34) \quad 1 - \hat{D}(k) = \frac{1}{d} \sum_{j=1}^d (1 - \cos k_j) \sim \frac{|k|^2}{2d}$$

as $k \rightarrow 0$. We need to see that the term $\hat{\Pi}_z(0) - \hat{\Pi}_z(k)$ is relatively small in high dimensions. By symmetry, we can write this term as

$$(4.35) \quad \hat{\Pi}_z(0) - \hat{\Pi}_z(k) = \sum_{x \in \mathbb{Z}^d} (1 - e^{ik \cdot x}) \Pi_z(x) = \sum_{x \in \mathbb{Z}^d} (1 - \cos k \cdot x) \Pi_z(x).$$

Finally, we note that the equation $\chi(z_c) = \infty$ can be rewritten as $0 = \chi(z_c)^{-1} = 1 - z_c |\Omega| - \hat{\Pi}_{z_c}(0)$, from which we see that the critical point z_c is given implicitly by

$$(4.36) \quad z_c = \frac{1}{|\Omega|} (1 - \hat{\Pi}_{z_c}(0)).$$

This equation has been the starting point for the study of z_c , in particular for the derivation of the $1/d$ expansion for the connective constant discussed in Section 1.4. Problem 5.1 below indicates how the first terms are obtained.

4.4. Tutorial. These problems develop the original derivation of the lace expansion by Brydges and Spencer [14]. All this material can also be found in [69].

We require a notion of graphs on integer intervals, and connectivity of these graphs. We emphasise in advance that the notion of connectivity is *not* the usual graph theoretic one, but that it is the right notion in this context.

DEFINITION 4.3. (i) Let $I = [a, b]$ be an interval of non-negative integers. An *edge* is a pair $st = \{s, t\}$ with $s, t \in \mathbb{Z}$ and $a \leq s < t \leq b$. A *graph* on $[a, b]$ is a set of edges. We denote the set of all graphs on $[a, b]$ by $\mathcal{B}[a, b]$.

(ii) A graph $\Gamma \in \mathcal{B}[a, b]$ is *connected* if a, b are endpoints of edges, and if for any $c \in (a, b)$, there are $s, t \in [a, b]$ such that $c \in (s, t)$ and $st \in \Gamma$. Equivalently, Γ is connected if $(a, b) = \cup_{st \in \Gamma} (s, t)$. The set of all connected graphs on $[a, b]$ is denoted $\mathcal{G}[a, b]$.

PROBLEM 4.1. Give an example of a graph which is connected in the above sense, but not path-connected in the usual graph theoretic sense, and give an example which is path-connected, but not connected in the above sense.

Let $U_{st}(\omega) = -1_{\{\omega(s) \neq \omega(t)\}}$, and for $a < b$ define

$$(4.37) \quad K[a, b](\omega) = \prod_{a \leq s < t \leq b} (1 + U_{st}(\omega)), \quad K[a, a](\omega) = 1,$$

so that

$$(4.38) \quad c_n(x) = \sum_{\omega \in \mathcal{W}_n(0, x)} K[0, n](\omega).$$

PROBLEM 4.2. Show that

$$(4.39) \quad K[a, b](\omega) = \sum_{\Gamma \in \mathcal{B}[a, b]} \prod_{st \in \Gamma} U_{st}(\omega).$$

PROBLEM 4.3. For $a < b$, let

$$(4.40) \quad J[a, b](\omega) = \sum_{\Gamma \in \mathcal{G}[a, b]} \prod_{st \in \Gamma} U_{st}(\omega).$$

Show that

$$(4.41) \quad K[a, b] = K[a + 1, b] + \sum_{j=a+1}^b J[a, j]K[j, b].$$

PROBLEM 4.4. Define

$$(4.42) \quad \pi_m(x) = \sum_{\omega \in \mathcal{W}_m(0, x)} J[0, m](\omega)$$

for $m \geq 1$. Use Problem 4.3 to show that, for $n \geq 1$,

$$(4.43) \quad c_n(x) = (c_1 * c_{n-1})(x) + \sum_{m=1}^n (\pi_m * c_{n-m})(x).$$

(Compared to (4.16), the sum here starts at $m = 1$ instead of $m = 2$. In fact, we will see that $\pi_1(x) = 0$ for the self-avoiding walk, since walks cannot self-intersect in 1 step.)

DEFINITION 4.4. A *lace* is a minimally connected graph, that is, a connected graph for which the removal of any edge would result in a disconnected graph. The set of laces on $[a, b]$ is denoted $\mathcal{L}[a, b]$.

PROBLEM 4.5. Let $L = \{s_1 t_1, \dots, s_N t_N\}$, where $s_l < t_l$ and $s_l \leq s_{l+1}$ for all l (and all the edges are different). Show that L is a lace if and only if

$$(4.44) \quad a = s_1 < s_2, \quad s_N < t_{N-1} < t_N = b, \quad s_{l+1} < t_l \leq s_{l+2} \quad (1 \leq l \leq N-2),$$

or $L = \{ab\}$ if $N = 1$. In particular, for $N > 1$, L divides $[a, b]$ into $2N - 1$ subintervals,

$$(4.45) \quad [s_1, s_2], [s_2, t_1], [t_1, s_3], [s_3, t_2], \dots, [t_{N-2}, s_N], [s_N, t_{N-1}], [t_{N-1}, t_N].$$

Determine which of these intervals must have length at least 1, and which can have length 0.

Let $\Gamma \in \mathcal{G}[a, b]$ be a connected graph. We associate a unique lace L_Γ to Γ as follows: Let

$$(4.46) \quad \begin{aligned} t_1 &= \max\{t : at \in \Gamma\}, \quad s_1 = a, \\ t_{i+1} &= \max\{t : \exists s < t_i \text{ such that } st \in \Gamma\}, \quad s_{i+1} = \min\{s : st_{i+1} \in \Gamma\}. \end{aligned}$$

The procedure terminates when $t_N = b$ for some N , and we then define $L_\Gamma = \{s_1 t_1, \dots, s_N t_N\}$. We define the set of edges *compatible* with a lace $L \in \mathcal{L}[a, b]$ to be

$$(4.47) \quad \mathcal{C}(L) = \{st : L_{L \cup \{st\}} = L, st \notin L\}.$$

PROBLEM 4.6. Show that $L_\Gamma = L$ if and only if $L \subset \Gamma$ and $\Gamma \setminus L \subset \mathcal{C}(L)$.

PROBLEM 4.7. Show that

$$(4.48) \quad J[a, b](\omega) = \sum_{L \in \mathcal{L}[a, b]} \prod_{st \in L} U_{st}(\omega) \sum_{\Gamma: L_\Gamma = L} \prod_{s't' \in \Gamma \setminus L} U_{s't'}(\omega).$$

Conclude from the previous exercise that

$$(4.49) \quad \sum_{\Gamma: L_\Gamma = L} \prod_{s't' \in \Gamma \setminus L} U_{s't'}(\omega) = \prod_{s't' \in \mathcal{C}(L)} (1 + U_{s't'}(\omega)),$$

and thus

$$(4.50) \quad J[a, b](\omega) = \sum_{L \in \mathcal{L}[a, b]} \prod_{st \in L} U_{st}(\omega) \prod_{s't' \in \mathcal{C}(L)} (1 + U_{s't'}(\omega)).$$

PROBLEM 4.8. Let $\mathcal{L}^{(N)}[a, b]$ denote the set of laces on $[a, b]$ which consist of exactly N edges. Define

$$(4.51) \quad J^{(N)}[a, b](\omega) = \sum_{L \in \mathcal{L}^{(N)}[a, b]} \prod_{st \in L} U_{st}(\omega) \prod_{s't' \in \mathcal{C}(L)} (1 + U_{s't'}(\omega))$$

and

$$(4.52) \quad \pi_m^{(N)}(x) = (-1)^N \sum_{\omega \in \mathcal{W}_m(0,x)} J^{(N)}[0,m](\omega).$$

(a) Prove that

$$(4.53) \quad \pi_m(x) = \sum_{N=1}^{\infty} (-1)^N \pi_m^{(N)}(x)$$

with $\pi_m^{(N)}(x) \geq 0$.

- (b) Describe the walk configurations that correspond to non-zero terms in $\pi_m^{(N)}(x)$, for $N = 1, 2, 3, 4$. What parts of the walk must be mutually avoiding?
- (c) What is the interpretation of the possibly empty intervals in Problem 4.5?

5. Lace expansion analysis in dimensions $d > 4$

In this section, we outline a proof that the bubble condition holds for the nearest-neighbour model in sufficiently high dimensions, and for the spread-out model in dimensions $d > 4$ provided L is large enough. As noted above, the bubble condition implies that $\gamma = 1$ in the sense that the susceptibility diverges linearly at the critical point as in (4.5). Proving the bubble condition will require control of the generating function $\hat{\Pi}_z(k)$ at the critical value $z = z_c$. According to (4.21) (see also Problem 4.8), Π_z is given by an infinite series

$$(5.1) \quad \Pi_z(x) = \sum_{N=1}^{\infty} (-1)^N \Pi_z^{(N)}(x), \quad \Pi_z^{(N)}(x) = \sum_{m=2}^{\infty} \pi_m^{(N)}(x) z^m.$$

The lace expansion is said to converge if $\Pi_z(x)$ is absolutely summable when $z = z_c$, in the strong sense that

$$(5.2) \quad \sum_{x \in \mathbb{Z}^d} \sum_{N=1}^{\infty} \Pi_{z_c}^{(N)}(x) < \infty.$$

There are now several different approaches to proving convergence of the lace expansion. In particular, a powerful but technically demanding method involves the study of (4.16) by induction on n [41]. Here we will follow the relatively simple approach of [69], which was inspired by a similar argument for percolation in [2]. Some details are omitted below; these can all be found in [69].

We will make use of the usual ℓ^p norms on functions on \mathbb{Z}^d , for $p = 1, 2, \infty$. In addition, when dealing with functions on the torus $[-\pi, \pi]^d$, we will use the usual L^p norms with respect to the probability measure $(2\pi)^{-d} d^d k$ on the torus, for $p = 1, 2$. To simplify the notation, we will sometimes omit the measure, and write, e.g., $\mathbf{B}(z) = \int \hat{G}_z^2 = \|\hat{G}_z\|_2^2$.

5.1. Diagrammatic estimates. We will obtain bounds on $\Pi_z(x)$ in terms of $G_z(x)$ and the closely related quantity $H_z(x)$ defined by

$$(5.3) \quad H_z(x) = G_z(x) - \delta_{0x} = \sum_{n=1}^{\infty} c_n(x) z^n.$$

The trivial term $c_0(x) = \delta_{0x}$ in $G_z(x)$ gives rise to a contribution 1 in the bubble diagram, and it will be important in the following that this contribution sometimes be omitted. It is for this reason that we use H_z as well as G_z .

The following *diagrammatic estimates* bound Π_z in terms of H_z and G_z . Once this theorem has been proved, the details of the definition of Π_z are no longer needed—the rest of the argument is analysis that uses the diagrammatic estimates.

THEOREM 5.1. *For any $z \geq 0$,*

$$(5.4) \quad \sum_{x \in \mathbb{Z}^d} \Pi_z^{(1)}(x) \leq z |\Omega| \|H_z\|_\infty,$$

$$(5.5) \quad \sum_{x \in \mathbb{Z}^d} \Pi_z^{(N)}(x) \leq \|H_z\|_\infty \|G_z * H_z\|_\infty^{N-1} \quad (N \geq 2),$$

$$(5.6) \quad \sum_{x \in \mathbb{Z}^d} (1 - \cos k \cdot x) \Pi_z^{(1)}(x) = 0,$$

$$(5.7) \quad \sum_{x \in \mathbb{Z}^d} (1 - \cos k \cdot x) \Pi_z^{(N)}(x) \leq N^2 \|(1 - \cos k \cdot x) H_z\|_\infty \|G_z * H_z\|_\infty^{N-1} \quad (N \geq 2).$$

PROOF. We prove just the cases $N = 1, 2$ here; the complete proof can be found in [69, Theorem 4.1].

For $N = 1$, since $\pi_m^{(1)}(y)$ is equal to $-\delta_{0y}$ times the number $\sum_{e \in \Omega} c_{m-1}(e)$ of self-avoiding returns, we have

$$(5.8) \quad \sum_{x \in \mathbb{Z}^d} \Pi_z^{(1)}(x) = - \sum_{y \in \Omega} \sum_{m=2}^{\infty} c_{m-1}(y) z^m = - \sum_{y \in \Omega} z H_z(y),$$

which implies (5.4). Also, (5.6) follows from

$$(5.9) \quad \sum_{x \in \mathbb{Z}^d} (1 - \cos k \cdot x) \Pi_z^{(1)}(x) = (1 - \cos k \cdot 0) \Pi_z^{(1)}(0) = 0.$$

For $N = 2$, dropping the mutual avoidance constraint between the three lines in $\pi_m^{(2)}(x)$ in (4.21) gives

$$(5.10) \quad \begin{aligned} \sum_{x \in \mathbb{Z}^d} \Pi_z^{(2)}(x) &\leq \sum_{x \in \mathbb{Z}^d} H_z(x)^3 \leq \|H_z\|_\infty (H_z * H_z)(0) \\ &\leq \|H_z\|_\infty \|H_z * H_z\|_\infty \end{aligned}$$

and

$$(5.11) \quad \sum_{x \in \mathbb{Z}^d} (1 - \cos k \cdot x) \Pi_z^{(2)}(x) \leq \|(1 - \cos k \cdot x) H_z\|_\infty \|H_z * H_z\|_\infty.$$

Since $0 \leq H_z(x) \leq G_z(x)$, this is stronger than (5.5) and (5.7). \square

5.2. The small parameter. Theorem 5.1 shows that the sum over N in (5.1) can be dominated by the sum of a geometric series with ratio $\|G_z * H_z\|_\infty$. Ideally, we would like this ratio to be small. A Cauchy–Schwarz estimate gives

$$(5.12) \quad \begin{aligned} \|H_z * G_z\|_\infty &\leq \|H_z\|_\infty + \|H_z * H_z\|_\infty \leq \|H_z\|_\infty + \|H_z\|_2^2 \\ &\leq \|H_z\|_\infty + \|G_z\|_2^2 = \|H_z\|_\infty + \mathbf{B}(z), \end{aligned}$$

but this looks problematic because the upper bound involves the bubble diagram—the very quantity we are trying to prove is finite at the critical point! So we will need some insight to make good use of the diagrammatic estimates.

An important idea will be to use not just the finiteness, but also the smallness of H_z . Specifically, we might hope that $\|H_{z_c}\|_2^2 = \|\hat{H}_{z_c}\|_2^2 = \|\hat{G}_{z_c} - 1\|_2^2$ should be small when the corresponding quantity for SRW is small.

Let $C_z(x) = \sum_{n=0} c_n^{(0)}(x)z^n$ be the analogue of $G_z(x)$ for the SRW model. Its critical value is $z_0 = |\Omega|^{-1}$, and

$$(5.13) \quad \hat{C}_z(k) = \frac{1}{1 - z|\Omega|\hat{D}(k)}, \quad \hat{C}_{z_0}(k) = \frac{1}{1 - \hat{D}(k)}.$$

The SRW analogue of $\|\hat{G}_{z_c} - 1\|_2^2$ is

$$(5.14) \quad \|\hat{C}_{z_0} - 1\|_2^2 = \int \left(\frac{1}{1 - \hat{D}} - 1 \right)^2 = \int \frac{\hat{D}^2}{(1 - \hat{D})^2}.$$

The following elementary proposition shows that the above integral is small for the models we are studying. The hypothesis $d > 4$ is needed for convergence, due to the $(|k|^{-2})^2$ singularity at the origin.

PROPOSITION 5.2. *Let $d > 4$. Then*

$$(5.15) \quad \int \frac{\hat{D}^2}{(1 - \hat{D})^2} \leq \beta$$

where, for some constant K ,

$$(5.16) \quad \beta = \begin{cases} \frac{K}{d-4} & \text{for the nearest-neighbour model,} \\ \frac{K}{L^d} & \text{for the spread-out model.} \end{cases}$$

PROOF. This is a calculus problem. For the nearest-neighbour model, see [57, Lemma A.3], and for the spread-out model see [69, Proposition 5.3]. \square

We will prove the following theorem.

THEOREM 5.3. *There are constants β_0 and C , independent of d and L , such that when (5.15) holds with $\beta \leq \beta_0$ we have $\mathbf{B}(z_c) \leq 1 + C\beta$.*

Theorem 5.3 achieves our goal of proving the bubble condition for the nearest-neighbour model in sufficiently high dimensions, and for the spread-out model with L sufficiently large in dimensions $d > 4$. As noted previously, this gives the following corollary that $\gamma = 1$ in high dimensions.

COROLLARY 5.4. *When (5.15) holds with $\beta \leq \beta_0$, then as $z \nearrow z_c$,*

$$(5.17) \quad \chi(z) \asymp \frac{1}{1 - z/z_c}.$$

5.3. Proof of Theorem 5.3. We begin with the following elementary lemma, which will be a principal ingredient in the proof.

LEMMA 5.5. *Let $a < b$ be real numbers and let f be a continuous real-valued function on $[z_1, z_2]$ such that $f(z_1) \leq a$. Suppose that, for each $z \in (z_1, z_2)$, we have the implication*

$$(5.18) \quad f(z) \leq b \quad \implies \quad f(z) \leq a.$$

Then $f(z) \leq a$ for all $z \in [z_1, z_2]$.

PROOF. The result is a straightforward application of the Intermediate Value Theorem. \square




FIGURE 9. The definition of $p(z)$.

We will apply Lemma 5.5 to a carefully chosen function f , based on a *coupling* between \hat{G} on the parameter range $[0, z_c)$, and the SRW analogue \hat{C} on the parameter range $[0, z_0)$. To define the coupling, let $z \in [0, z_c)$ and define $p(z) \in [0, z_0)$ by

$$(5.19) \quad \hat{G}_z(0) = \chi(z) = \hat{C}_{p(z)}(0) = \frac{1}{1 - p(z) |\Omega|},$$

i.e.,

$$(5.20) \quad p(z) |\Omega| = 1 - \chi(z)^{-1} = z |\Omega| + \hat{\Pi}_z(0).$$

See Figure 9. We expect (or hope!) that $\hat{G}_z(k) \approx \hat{C}_{p(z)}(k)$ for *all* k , not just for $k = 0$, as well as an additional condition that expresses another form of similarity between $\hat{G}_z(k)$ and $\hat{C}_{p(z)}(k)$. For the latter, we define

$$(5.21) \quad -\frac{1}{2} \Delta_k \hat{G}_z(l) = \hat{G}_z(l) - \frac{1}{2} (\hat{G}_z(l+k) + \hat{G}_z(l-k));$$

this is the Fourier transform of $(1 - \cos k \cdot x) G_z(x)$ with l as the dual variable. We aim to apply Lemma 5.5 with $z_1 = 0$, $z_2 = z_c$, $a = 1 + \text{const} \beta$ (with a constant whose value is determined in (5.27) below), $b = 4$ (in fact, any fixed $b > 1$ will do here), and

$$(5.22) \quad f(z) = \max \{f_1(z), f_2(z), f_3(z)\}$$

where

$$(5.23) \quad f_1(z) = z|\Omega|, \quad f_2(z) = \sup_{k \in [-\pi, \pi]^d} \frac{|\hat{G}_z(k)|}{|\hat{C}_{p(z)}(k)|},$$

and

$$(5.24) \quad f_3(z) = \sup_{k, l \in [-\pi, \pi]^d} \frac{\frac{1}{2} |\Delta_k \hat{G}_z(l)|}{|U_{p(z)}(k, l)|},$$

with

$$(5.25) \quad U_{p(z)}(k, l) = 16\hat{C}_{p(z)}(k)^{-1} \left(\hat{C}_{p(z)}(l-k)\hat{C}_{p(z)}(l) + \hat{C}_{p(z)}(l+k)\hat{C}_{p(z)}(l) \right. \\ \left. + \hat{C}_{p(z)}(l-k)\hat{C}_{p(z)}(l+k) \right).$$

The choice of $U_{p(z)}(k, l)$ is made for technical reasons not explained here, and should be regarded as a useful replacement for the more natural choice $\frac{1}{2} |\Delta_k \hat{C}_{p(z)}(l)|$.

The conclusion from Lemma 5.5 would be that $f(z) \leq a$ for all $z \in [0, z_c)$. The inequality (5.15) can be used to show that $\|(1 - \hat{D})^{-1}\|_2^2 \leq 1 + 3\beta$ (see [69, (5.10)]), and hence we may assume that $\|(1 - \hat{D})^{-1}\|_2^2 \leq 2$. Using $f_2(z) \leq a$ we therefore conclude that

$$(5.26) \quad B(z_c) = \lim_{z \nearrow z_c} B(z) = \lim_{z \nearrow z_c} \|\hat{G}_z\|_2^2 \\ \leq a^2 \lim_{z \nearrow z_c} \|\hat{C}_{p(z)}\|_2^2 = a^2 \|(1 - \hat{D})^{-1}\|_2^2 \\ \leq 2a^2 < \infty$$

which is our goal. Thus it suffices to verify the hypotheses on $f(z)$ in Lemma 5.5. This is the content of the following lemma.

LEMMA 5.6. *The function $f(z)$ defined by (5.22)–(5.24) is continuous on $[0, z_c)$, with $f(0) = 1$, and for each $z \in (0, z_c)$,*

$$(5.27) \quad f(z) \leq 4 \quad \implies \quad f(z) \leq 1 + O(\beta).$$

PROOF. It is relatively easy to verify the continuity of f , and we omit the details. To see that $f(0) = 1 \leq a$, we observe that $f_1(0) = 0$, $p(0) = 0$ and hence $f_2(0) = 1/1 = 1$, and $f_3(0) = 0$. The difficult step is to prove the implication (5.27), and the remainder of the proof concerns this step. We assume throughout that $f(z) \leq 4$.

We consider first $f_1(z) = z|\Omega|$. Our goal is to prove that $f_1(z) \leq 1 + O(\beta)$, and for this we will only use the assumptions $f_1(z) \leq 4$ and $f_2(z) \leq 4$; we do not yet need f_3 . Since $0 < \chi(z) < \infty$, we have $\chi(z)^{-1} = 1 - z|\Omega| - \hat{\Pi}_z(0) > 0$, i.e.,

$$(5.28) \quad f_1(z) = z|\Omega| < 1 - \hat{\Pi}_z(0) \leq 1 + |\hat{\Pi}_z(0)|.$$

The required bound for $f_1(z)$ will follow once we show that for all $z \in (0, z_c)$ and for all $k \in [-\pi, \pi]^d$,

$$(5.29) \quad |\hat{\Pi}_z(k)| \leq O(\beta).$$

To prove (5.29) we use Theorem 5.1 (more precisely, (5.4)–(5.5)), to obtain

$$(5.30) \quad \begin{aligned} |\hat{\Pi}_z(k)| &\leq \sum_{N=1}^{\infty} \sum_{x \in \mathbb{Z}^d} \Pi_z^{(N)}(x) \\ &\leq \|H_z\|_{\infty} \left(f_1(z) + \sum_{N=2}^{\infty} \|G_z * H_z\|_{\infty}^{N-1} \right). \end{aligned}$$

For the first term, we use $f_1(z) \leq 4$. For the second term, we need a bound on $\|G_z * H_z\|_{\infty}$ in order to bound the sum. By definition,

$$(5.31) \quad \|G_z * H_z\|_{\infty} \leq \|H_z\|_{\infty} + \|H_z * H_z\|_{\infty} \leq \|H_z\|_{\infty} + \|H_z\|_2^2.$$

Now H_z is the generating function for SAWs which take at least one step. By omitting the avoidance constraint between the first step and subsequent steps, we obtain

$$(5.32) \quad H_z(x) \leq z |\Omega| (D * G_z)(x) \leq 4(D * G_z)(x).$$

Thus we can bound the second term in (5.31), using $f_2(z) \leq 4$ and Proposition 5.2, as

$$(5.33) \quad \begin{aligned} \|H_z\|_2^2 &\leq 4^2 \|D * G_z\|_2^2 = 4^2 \|\hat{D}\hat{G}_z\|_2^2 \\ &\leq 4^4 \|\hat{D}\hat{C}_{p(z)}\|_2^2 = 4^4 \|D * C_{p(z)}\|_2^2 \\ &\leq 4^4 \|D * C_{z_0}\|_2^2 = 4^4 \|\hat{D}(1 - \hat{D})^{-1}\|_2^2 \\ &\leq 4^4 \beta. \end{aligned}$$

Similar estimates show $\|H_z\|_{\infty} \leq O(\beta)$. If we substitute these estimates into (5.30), we obtain

$$(5.34) \quad |\hat{\Pi}_z(k)| \leq C\beta \left(4 + \sum_{N=2}^{\infty} (C\beta)^{N-1} \right)$$

for some constant C , so that (5.29) will hold for β sufficiently small. This completes the proof for $f_1(z)$.

We next sketch the proof that $f_2(z) \leq 1 + O(\beta)$. Recalling the notation $\hat{F}_z(k) = \hat{G}_z(k)^{-1}$ introduced in (4.31), and using the formulas (5.19) and (5.20) for $p(z)$, we obtain

$$(5.35) \quad \begin{aligned} \frac{\hat{G}_z(k)}{\hat{C}_{p(z)}(k)} - 1 &= \frac{1 - p(z) |\Omega| \hat{D}(k)}{\hat{F}_z(k)} - 1 \\ &= \frac{1 - (z |\Omega| + \hat{\Pi}_z(0)) \hat{D}(k) - \hat{F}_z(k)}{\hat{F}_z(k)} \\ &= \frac{-\hat{\Pi}_z(0) \hat{D}(k) + \hat{\Pi}_z(k)}{\hat{F}_z(k)} \\ &= \frac{\hat{\Pi}_z(0)(1 - \hat{D}(k)) - (\hat{\Pi}_z(0) - \hat{\Pi}_z(k))}{\hat{F}_z(k)}. \end{aligned}$$

The bound $f_3(z) \leq 4$ and (5.7) can be used to show that $|\hat{\Pi}_z(0) - \hat{\Pi}_z(k)| \leq O(\beta)(1 - \hat{D}(k))$ (see [69] for details); it is precisely at this point that the need to include f_3

in the definition of f arises. Together with (5.29), this shows that the numerator of (5.35) is $O(\beta)(1 - \hat{D}(k))$.

For the denominator, we recall the formula (4.32):

$$(5.36) \quad \hat{F}_z(k) = \chi(z)^{-1} + z |\Omega| (1 - \hat{D}(k)) + (\hat{\Pi}_z(0) - \hat{\Pi}_z(k)).$$

To bound $\hat{F}_z(k)$ from below, we consider two parameter ranges for z . If $z \leq \frac{1}{2} |\Omega|^{-1}$, we can make the trivial estimate $\chi(z)^{-1} \geq \hat{C}_z(0)^{-1} = 1 - z |\Omega| \geq \frac{1}{2}$, so that $\hat{F}_z(k) \geq \frac{1}{2} + 0 - O(\beta) \geq \frac{1}{4}$ for small β . Since the numerator of (5.35) is itself $O(\beta)$, this proves that $f_2(z) \leq 1 + O(\beta)$ for this range of z .

It remains to consider $\frac{1}{2} |\Omega|^{-1} \leq z \leq z_c$. Now we estimate

$$(5.37) \quad \hat{F}_z(k) \geq 0 + \frac{1}{2} (1 - \hat{D}(k)) - O(\beta)(1 - \hat{D}(k)) \geq \frac{1}{4} (1 - \hat{D}(k)).$$

The factors $1 - \hat{D}(k)$ in the numerator and denominator of (5.35) cancel, leaving $O(\beta)$ as desired.

Finally the proof for $f_3(z)$ is similar to the proof for $f_2(z)$, and we refer to [69] for the details. \square

5.4. Tutorial. For simplicity, we restrict our attention now to the nearest-neighbour model of SAWs in dimensions sufficiently high that the preceding arguments and conclusions apply. In Lemma 5.6, we found that $f_2(z) \leq a = 1 + O(d^{-1})$, since $\beta \leq O((d-4)^{-1}) = O(d^{-1})$. This estimate, which states that

$$(5.38) \quad \hat{G}_z(k) \leq a \hat{C}_{p(z)}(k) \quad k \in [\pi, \pi]^d, \quad z \in (0, z_c),$$

is most important for $k \approx 0$, the *small frequencies*, and it is referred to as the *infrared bound*. Other bounds obtained in Lemma 5.6 can be framed as follows: there is a constant c , independent of $z \leq z_c$, such that

$$(5.39) \quad \|H_z\|_2^2 \leq cd^{-1}, \quad \|H_z\|_\infty \leq cd^{-1}, \quad \|\Pi_z\|_1 \leq cd^{-1},$$

and

$$(5.40) \quad \|\Pi_z^{(N)}\|_1 \leq (Cd^{-1})^N, \quad \sum_{N=M}^{\infty} \|\Pi_z^{(N)}\|_1 \leq O(Cd^{-M}).$$

We also recall that the Fourier transform of the two-point function can be written as

$$(5.41) \quad \hat{G}_z(k) = \frac{1}{1 - z|\Omega|\hat{D}(k) - \hat{\Pi}_z(k)}.$$

Since $\hat{G}_z(0) \rightarrow \infty$ as $z \rightarrow z_c$, we obtain the equation

$$(5.42) \quad 1 - z_c |\Omega| - \hat{\Pi}_{z_c}(0) = 0.$$

This equation provides a starting point to study the connective constant $\mu = z_c^{-1}$.

PROBLEM 5.1. In this problem, we show that the connective constant obeys

$$(5.43) \quad \mu = 2d - 1 - (2d)^{-1} + O((2d)^{-2}) \quad \text{as } d \rightarrow \infty.$$

This special case of the results discussed in Section 1.4 was first proved by Kesten [51], by very different means.

(a) Let $m \geq 1$ be an integer. Show that $\|(1 - \hat{D})^{-m}\|_1$ is non-increasing in $d > 2m$. In particular, it follows that $\|\hat{C}_{z_0}\|_2$ is bounded uniformly in $d > 4$.

Hint: $A^{-m} = \Gamma(m)^{-1} \int_0^\infty u^{m-1} e^{-uA} du$.

(b) Let $H_z^{(j)}(x) = \sum_{m=j}^{\infty} c_m(x)z^m$ be the generating function for SAWs that take at least j steps. By relaxing the condition of mutual self-avoidance for the first j steps, show that

$$(5.44) \quad \|H_{z_c}^{(j)}\|_{\infty} \leq O((2d)^{-j/2}), \quad j > 1.$$

Hint: Use the infrared bound for the two-point function (5.38), and that the probability that a $2j$ -step simple random walk which starts at 0 also ends at 0 is

$$(5.45) \quad \|\hat{D}^{2j}\|_1 \leq O((2d)^{-j}).$$

(c) Recall that $\pi_n^{(1)}(x) = 0$ if $x \neq 0$, so that $\hat{\Pi}_z^{(1)}(0) = \sum_{x \in \mathbb{Z}^d} \Pi_z^{(1)}(x) = \Pi_z^{(1)}(0)$ is the generating function for all self-avoiding returns. Prove that

$$(5.46) \quad \hat{\Pi}_{z_c}^{(1)}(0) = (2d)^{-1} + 3(2d)^{-2} + O((2d)^{-3})$$

(d) Note that $\hat{\Pi}_z^{(2)}(0)$ is the generating function for all θ -walks: paths that visit their eventual endpoint, return to the origin, then return to their endpoint, and are otherwise self-avoiding. Prove that

$$(5.47) \quad \hat{\Pi}_z^{(2)}(0) = (2d)^{-2} + O((2d)^{-3}).$$

(e) Conclude from (c) and (d) that

$$(5.48) \quad \hat{\Pi}_z(0) = -(2d)^{-1} - 2(2d)^{-2} + O((2d)^{-3}),$$

and use this to show

$$(5.49) \quad \mu = 2d - 1 - (2d)^{-1} + O((2d)^{-2}).$$

We have seen in Section 4.2 that $\chi(z) \asymp (1 - z/z_c)^{-1}$ in high dimensions, assuming the bubble condition. The next problem shows that this bound can be improved to an asymptotic formula.

PROBLEM 5.2. (a) Show that

$$(5.50) \quad \frac{d[z\chi(z)]}{dz} = V(z)\chi(z)^2, \quad \text{where } V(z) = 1 - \hat{\Pi}_z(0) + z \frac{d\hat{\Pi}_z(0)}{dz}.$$

Hint: Let $\hat{F}_z(0) = \chi(z)^{-1} = 1 - z|\Omega| - \hat{\Pi}_z(0)$ and express the left-hand side in terms of $\hat{F}_z(0)$.

(b) Show that $\hat{\Pi}_{z_c}(0)$, $\frac{d}{dz}\hat{\Pi}_{z_c}(0)$ and thus $V(z_c)$ are finite. It follows that

$$(5.51) \quad \frac{d[z\chi(z)]}{dz} = V(z)\chi(z)^2 \sim V(z_c)\chi(z)^2 \quad \text{as } z \nearrow z_c,$$

where $f(z) \sim g(z)$ means $\lim_{z \nearrow z_c} f(z)/g(z) = 1$.

(c) Prove that $\chi(z) \sim A(1 - z/z_c)^{-1}$ as $z \nearrow z_c$, where the constant A is given by $A = z_c^{-1}[2d + \frac{d}{dz}|_{z=z_c}\hat{\Pi}_z(0)]^{-1}$.

6. Integral representation for walk models

It has long been understood by physicists that it is sometimes possible to represent random fields by random walks. Ideas in this direction due to Symanzik [71] were influential among mathematicians, and inspired, e.g., the analysis of [6, 7] who showed how to use random walks to represent and analyse ferromagnetic lattice spin systems. In this section, we develop representations of two random walk models in terms of random fields, via functional integrals. Our ultimate goal is rather the opposite to that of [6, 7], namely we wish to study models of random walks via studying their integral representations. This will be the topic of Section 7.

We begin in Section 6.1 with some background material about Gaussian integrals. In Section 6.2, we use these Gaussian integrals to represent a model of SAWs in a background of self-avoiding loops, a model closely related to the $O(n)$ loop model discussed in Section 3.4. The random field in these Gaussian integrals is called a *boson* field in physics. It was realised in the physics literature [59, 63] that the loops in the loop model could be eliminated by the use of anti-commuting variables, referred to as a *fermion* field, thereby providing a representation for models of SAWs. The anti-commuting variables can be understood in terms of differential forms with their anti-commuting wedge product, and in Sections 6.3–6.4 we provide the relevant background on differential forms and their integration. Finally, in Section 6.5, we obtain an integral representation for SAWs. The ideas in this section are developed in further detail in [11].

6.1. Gaussian integrals. Fix a positive integer M . Later, we identify the set $\{1, \dots, M\}$ with a finite set Λ on which the walks related to the fields take place, e.g., $\Lambda \subset \mathbb{Z}^d$. Consider a two-component real field

$$(6.1) \quad (u, v) = (u_x, v_x)_{x \in \{1, \dots, M\}} \in \mathbb{R}^M \times \mathbb{R}^M.$$

From this, we obtain the associated complex field $(\varphi, \bar{\varphi}) = (\varphi_x, \bar{\varphi}_x)_{x \in \{1, \dots, M\}}$, where

$$(6.2) \quad \varphi_x = u_x + iv_x, \quad \bar{\varphi}_x = u_x - iv_x;$$

this is the so-called *boson field*. We wish to integrate with respect to the variables $(\varphi_x, \bar{\varphi}_x)$, and for this we will use the differentials $d\varphi_x = du_x + i dv_x$ and $d\bar{\varphi}_x = du_x - i dv_x$. As we will discuss in more detail in Section 6.3, differentials are multiplied using an anti-commuting product, so in particular $du_x dv_x = -dv_x du_x$, $du_x du_x = dv_x dv_x = 0$, and $d\bar{\varphi}_x d\varphi_x = 2i du_x dv_x$.

Let $C = (C_{xy})_{x, y \in \{1, \dots, M\}}$ be an $M \times M$ complex matrix with positive Hermitian part, meaning that

$$(6.3) \quad \sum_{x, y=1}^M \varphi_x (C_{xy} + \bar{C}_{yx}) \bar{\varphi}_y > 0 \quad \text{for all } \varphi \neq 0 \text{ in } \mathbb{C}^M.$$

It is not difficult to see that this implies that $A = C^{-1}$ exists. The (complex) Gaussian measure with covariance C is defined by

$$(6.4) \quad d\mu_C(\varphi, \bar{\varphi}) = \frac{1}{Z_C} e^{-\varphi A \bar{\varphi}} d\bar{\varphi} d\varphi,$$

where $\varphi A \bar{\varphi} = \sum_{x, y=1}^M \varphi_x A_{xy} \bar{\varphi}_y$, and

$$(6.5) \quad d\bar{\varphi} d\varphi = d\bar{\varphi}_1 d\varphi_1 \cdots d\bar{\varphi}_M d\varphi_M = (2i)^M du_1 dv_1 \cdots du_M dv_M$$

is a multiple of the Lebesgue measure on \mathbb{R}^{2M} . The normalisation constant

$$(6.6) \quad Z_C = \int_{\mathbb{R}^{2M}} e^{-\varphi A \bar{\varphi}} d\bar{\varphi} d\varphi$$

can be computed explicitly.

LEMMA 6.1. *For C with positive Hermitian part, the normalisation of the Gaussian integral is given by*

$$(6.7) \quad Z_C = \frac{(2\pi i)^M}{\det A}.$$

PROOF. In this proof, we make the simplifying assumption that C and thus also A are Hermitian, though the result holds more generally; see [11]. By the spectral theorem for Hermitian matrices, there is a positive diagonal matrix $D = \text{diag}(d_x)$ and a unitary matrix U such that $A = U^{-1}DU$. Then, $\varphi A \bar{\varphi} = \rho D \bar{\rho}$ where $\rho = \bar{U}\varphi$ (\bar{U} is the complex conjugate of U). By a change of variables in the integral and explicit computation of the resulting 1-dimensional integral,

$$(6.8) \quad Z_C = \prod_{x=1}^M \int_{\mathbb{R}^2} e^{-d_x(u_x^2 + v_x^2)} 2i du_x dv_x = \frac{(2\pi i)^M}{\prod_{x=1}^M d_x} = \frac{(2\pi i)^M}{\det A}. \quad \square$$

We define the differential operators

$$(6.9) \quad \frac{\partial}{\partial \varphi_x} = \frac{1}{2} \left(\frac{\partial}{\partial u_x} - i \frac{\partial}{\partial v_x} \right), \quad \frac{\partial}{\partial \bar{\varphi}_x} = \frac{1}{2} \left(\frac{\partial}{\partial u_x} + i \frac{\partial}{\partial v_x} \right).$$

It is easy to check that

$$(6.10) \quad \frac{\partial \varphi_y}{\partial \varphi_x} = \frac{\partial \bar{\varphi}_y}{\partial \bar{\varphi}_x} = \delta_{xy}, \quad \frac{\partial \bar{\varphi}_y}{\partial \varphi_x} = \frac{\partial \varphi_y}{\partial \bar{\varphi}_x} = 0.$$

The following integration by parts formula will be useful.

LEMMA 6.2. *For C with positive Hermitian part, and for nice functions F ,*

$$(6.11) \quad \int \bar{\varphi}_a F d\mu_C(\varphi, \bar{\varphi}) = \sum_{x=1}^M C_{ax} \int \frac{\partial F}{\partial \varphi_x} d\mu_C(\varphi, \bar{\varphi}).$$

PROOF. Integrating by parts, we obtain

$$(6.12) \quad \begin{aligned} \int \frac{\partial F}{\partial \varphi_x} e^{-\varphi A \bar{\varphi}} d\bar{\varphi} d\varphi &= - \int F \frac{\partial}{\partial \varphi_x} e^{-\varphi A \bar{\varphi}} d\bar{\varphi} d\varphi \\ &= \int F \sum_y A_{xy} \bar{\varphi}_y e^{-\varphi A \bar{\varphi}} d\bar{\varphi} d\varphi. \end{aligned}$$

It follows from the fact that $C = A^{-1}$ that

$$(6.13) \quad \sum_{x=1}^M C_{ax} \int \frac{\partial F}{\partial \varphi_x} d\mu_C = \int \sum_{x,y} C_{ax} A_{xy} \bar{\varphi}_y F d\mu_C = \int \bar{\varphi}_a F d\mu_C. \quad \square$$

The following application of Lemma 6.2 is a special case of Wick's Theorem. The quantity appearing on the right-hand side of (6.14) is the *permanent* of the submatrix of C indexed by $(x_i, y_j)_{i,j=1}^k$.

LEMMA 6.3. Let $\{x_1, \dots, x_k\}$ and $\{y_1, \dots, y_k\}$ each be sets with k distinct elements from $\{1, \dots, M\}$. Then

$$(6.14) \quad \int \prod_{l=1}^k \bar{\varphi}_{x_l} \varphi_{y_l} d\mu_C = \sum_{\sigma \in S_k} \prod_{l=1}^k C_{x_l, y_{\sigma(l)}},$$

where the sum is over the set S_k of permutations of $\{1, \dots, k\}$.

PROOF. This follows by repeated application of the integration by parts formula in Lemma 6.2. Each time the formula is applied, one factor of $\bar{\varphi}$ disappears on the right-hand side of (6.11), and the partial differentiation eliminates one factor φ as well. \square

6.2. Integral representation for a loop model. Let Λ be a finite set of cardinality M . Fix $a, b \in \Lambda$ and a subset $X \subset \Lambda \setminus \{a, b\}$. An example we have in mind is $\Lambda \subset \mathbb{Z}^d$ and $X = \Lambda \setminus \{a, b\}$. We define the integral

$$(6.15) \quad G_{ab, X} = \int \bar{\varphi}_a \varphi_b \prod_{x \in X} (1 + \varphi_x \bar{\varphi}_x) d\mu_C.$$

As we now explain, this can be interpreted as a loop model whose configurations consist of a self-avoiding walk from a to b whose intermediate steps lie in X , together with a background of closed loops in X . We denote by $\mathcal{S}_{ab}(X)$ the set of sequences $(a, x_1, \dots, x_{n-1}, b)$ with $n \geq 1$ arbitrary and the $x_i \in X$ distinct—these are SAWs with rather general steps.

Repeated integration by parts gives

$$(6.16) \quad G_{ab, X} = \sum_{\omega \in \mathcal{S}_{ab}(X)} C^\omega \int \prod_{x \in X \setminus \omega} (1 + \varphi_x \bar{\varphi}_x) d\mu_C,$$

where $C^\omega = \prod_{i=1}^{\ell(\omega)} C_{w(i-1), w(i)}$. Also, by expanding the product and applying Lemma 6.3, we obtain

$$(6.17) \quad \begin{aligned} \int \prod_{x \in X \setminus \omega} (1 + \varphi_x \bar{\varphi}_x) d\mu_C &= \sum_{Z \subset X \setminus \omega} \int \prod_{x \in Z} \varphi_x \bar{\varphi}_x d\mu_C \\ &= \sum_{Z \subset X \setminus \omega} \sum_{\sigma \in S(Z)} \prod_{z \in Z} C_{z, \sigma(z)}, \end{aligned}$$

with $S(Z)$ is the set of permutations of the set Z . Altogether, this gives

$$(6.18) \quad G_{ab, X} = \sum_{\omega \in \mathcal{S}_{ab}(X)} C^\omega \sum_{Z \subset X \setminus \omega} \sum_{\sigma \in S(Z)} \prod_{z \in Z} C_{z, \sigma(z)}.$$

Thus, by decomposing the permutation σ into cycles, we can interpret (6.15) as the generating function for self-avoiding walks from a to b in a background of loops with weight C_{xy} for every step between x and y (with each loop corresponding to a cycle of σ). See Figure 10.

6.3. Differential forms. Our next goal is to modify the example of Section 6.2 with the help of differential forms, which are versions of what physicists call *fermions*, to obtain an integral representation for the generating function for self-avoiding walks *without* the loop background. A gentle introduction to differential forms can be found in [66].




FIGURE 10. Self-avoiding walk from a to b with loop background. Loops can have length zero. The loops will be eliminated by the use of differential forms.

The *Grassmann algebra* \mathcal{N} of differential forms is generated by the one-forms $du_1, dv_1, \dots, du_M, dv_M$, with anticommutative product \wedge . A p -form (a differential form of degree p) is a function of the variables (u, v) times a product of p differentials or sum of these. Because of anticommutativity, $du_x \wedge du_x = dv_x \wedge dv_x = 0$, and any p -form with $p > 2M$ must be zero. A form of maximal degree can thus be written uniquely as

$$(6.19) \quad K = f(u, v) du_1 \wedge dv_1 \wedge \cdots \wedge du_M \wedge dv_M,$$

where $du_1 \wedge dv_1 \wedge \cdots \wedge du_M \wedge dv_M$ is the standard volume form on \mathbb{R}^{2M} . A general differential form is a linear combination of p -forms, where different terms in the sum can have different values of p . Together, the differential forms constitute the algebra \mathcal{N} .

We will omit the wedge \wedge from the notation from now on, and write simply $du_1 dv_1$ for $du_1 \wedge dv_1$, but it should be borne in mind that order is significant in such an expression: $du_x dv_y = -dv_y du_x$. On the other hand, two forms of *even* degree commute.

We again use complex variables, and write

$$(6.20) \quad \begin{aligned} \varphi_x &= u_x + i v_x, & \bar{\varphi}_x &= u_x - i v_x, \\ d\varphi_x &= du_x + i dv_x, & d\bar{\varphi}_x &= du_x - i dv_x. \end{aligned}$$

Then

$$(6.21) \quad d\bar{\varphi}_x d\varphi_x = 2i du_x dv_x.$$

Given any fixed choice of the complex square root, we introduce the notation

$$(6.22) \quad \psi_x = \frac{1}{\sqrt{2\pi i}} d\varphi_x, \quad \bar{\psi}_x = \frac{1}{\sqrt{2\pi i}} d\bar{\varphi}_x.$$

The collection of differential forms

$$(6.23) \quad (\psi, \bar{\psi}) = (\psi_x, \bar{\psi}_x)_{x \in \{1, \dots, M\}}$$

is called the *fermion* field. It follows that

$$(6.24) \quad \bar{\psi}_x \psi_x = \frac{1}{\pi} du_x dv_x.$$

Let $\Lambda = \{1, \dots, M\}$. Given an $M \times M$ matrix A , we define the differential form

$$(6.25) \quad S_A = \varphi A \bar{\varphi} + \psi A \bar{\psi} = \sum_{x,y \in \Lambda} \varphi_x A_{xy} \bar{\varphi}_y + \sum_{x,y \in \Lambda} \psi_x A_{xy} \bar{\psi}_y.$$

An example of special interest is the case where $A_{uv} = \delta_{ux} \delta_{vx}$ for some fixed $x \in \Lambda$. In this case, we write τ_x in place of S_A , i.e.,

$$(6.26) \quad \tau_x = \varphi_x \bar{\varphi}_x + \psi_x \bar{\psi}_x.$$

6.4. Functions of forms and integrals of forms. The following definition tells us how to integrate a differential form.

DEFINITION 6.4. Let F be a differential form whose term K of maximal degree is as in (6.19). The *integral of F* is then defined to be

$$(6.27) \quad \int F = \int K = \int_{\mathbb{R}^{2M}} f(u, v) du_1 dv_1 \cdots du_M dv_M.$$

In particular, if F contains no term of degree $2M$ then its integral is zero.

We also need to define functions of even differential forms.

DEFINITION 6.5. Let $K = (K_j)_{j \in J}$ be a finite collection of differential forms, with each K_j even (a sum of forms of even degrees). Let $K_j^{(0)}$ be the degree zero part of K_j . Given a C^∞ function $F : \mathbb{R}^J \rightarrow \mathbb{C}$, we define $F(K)$ to be the form given by the Taylor polynomial (a polynomial in ψ and $\bar{\psi}$)

$$(6.28) \quad F(K) = \sum_{\alpha} \frac{1}{\alpha!} F^{(\alpha)}(K^{(0)})(K - K^{(0)})^\alpha$$

where $\alpha = (\alpha_1, \dots, \alpha_j)$ is a multi-index and

$$(6.29) \quad \alpha! = \prod_{j \in J} \alpha_j!, \quad (K - K^{(0)})^\alpha = \prod_{j \in J} (K_j - K_j^{(0)})^{\alpha_j}.$$

The sum in (6.28) is finite due to anticommutativity, and the product in (6.29) is well-defined because all factors are even and thus commute.

EXAMPLE 6.6. A simple but important example is $J = 1$ and $F(t) = e^{-t}$, for which we obtain, e.g.,

$$(6.30) \quad e^{-\tau_x} = e^{-\varphi_x \bar{\varphi}_x - \psi_x \bar{\psi}_x} = e^{-\varphi_x \bar{\varphi}_x} (1 - \psi_x \bar{\psi}_x),$$

$$(6.31) \quad e^{-S_A} = e^{-\varphi A \bar{\varphi} - \psi A \bar{\psi}} = e^{-\varphi A \bar{\varphi}} \sum_{n=0}^M \frac{(-1)^n}{n!} (\psi A \bar{\psi})^n.$$

The following lemma displays a remarkable self-normalisation property of these integrals.

LEMMA 6.7. *If A is a complex $M \times M$ matrix with positive Hermitian part, then*

$$(6.32) \quad \int e^{-S_A} = 1.$$

PROOF. Using (6.31) and Definition 6.4,

$$(6.33) \quad \begin{aligned} \int e^{-S_A} &= \int_{\mathbb{R}^{2M}} e^{-\varphi A \bar{\varphi}} \frac{1}{M!} (-1)^M (\psi A \bar{\psi})^M \\ &= \frac{1}{M!} \left(\frac{-1}{2\pi i} \right)^M \int_{\mathbb{R}^{2M}} e^{-\varphi A \bar{\varphi}} (d\varphi A d\bar{\varphi})^M. \end{aligned}$$

By definition,

$$(6.34) \quad (d\varphi A d\bar{\varphi})^M = \sum_{x_1, y_1} \cdots \sum_{x_M, y_M} A_{x_1 y_1} \cdots A_{x_M y_M} d\varphi_{x_1} d\bar{\varphi}_{y_1} \cdots d\varphi_{x_M} d\bar{\varphi}_{y_M}.$$

Due to the antisymmetry, non-zero contributions to the above sum require that x_1, \dots, x_M and y_1, \dots, y_M each be a permutation of $\{1, \dots, M\}$. Thus, by interchanging the (commuting) pairs $d\varphi_{x_i} d\bar{\varphi}_{y_i}$ so as to place the x_i in the order $1, \dots, M$, and then relabelling the y_i , we obtain

$$(6.35) \quad \begin{aligned} (d\varphi A d\bar{\varphi})^M &= M! \sum_{y_1, \dots, y_M} A_{1y_1} \cdots A_{My_M} d\varphi_1 d\bar{\varphi}_{y_1} \cdots d\varphi_M d\bar{\varphi}_{y_M} \\ &= M! \sum_{y_1, \dots, y_M} \epsilon_{y_1, \dots, y_M} A_{1y_1} \cdots A_{My_M} d\varphi_1 d\bar{\varphi}_{y_1} \cdots d\varphi_M d\bar{\varphi}_{y_M} \\ &= M! (-1)^M (\det A) d\bar{\varphi} d\varphi, \end{aligned}$$

where $\epsilon_{y_1, \dots, y_M}$ is the sign of the permutation (y_1, \dots, y_M) of $\{1, \dots, M\}$. With Lemma 6.1, it follows that

$$(6.36) \quad \int e^{-S_A} = \frac{\det A}{(2\pi i)^M} \int_{\mathbb{R}^{2M}} e^{-\varphi A \bar{\varphi}} d\bar{\varphi} d\varphi = 1. \quad \square$$

REMARK 6.8. More generally, the calculation in the previous proof also shows that for a function $f = f(\varphi, \bar{\varphi})$, a form of degree zero,

$$(6.37) \quad \int e^{-S_A} f = \int f d\mu_C \quad (C = A^{-1}),$$

provided f is such that the integral on the right-hand side converges. In our present setup, we have defined $\int e^{-S_A} F$ for more general forms F , so this provides an extension of the Gaussian integral of Section 6.1.

The self-normalisation property of Lemma 6.7 has the following beautiful extension. The precise hypotheses needed on F can be found in [11, Proposition 4.4].

LEMMA 6.9. *If A is a complex $M \times M$ matrix with positive Hermitian part, and $F : \mathbb{R}^M \rightarrow \mathbb{C}$ is a nice function (exponential growth at infinity is permitted), then*

$$(6.38) \quad \int e^{-S_A} F(\tau) = F(0),$$

where we regard τ as the vector (τ_1, \dots, τ_M) .

PROOF (SKETCH). If F is Schwartz class, e.g., then it can be expressed in terms of its Fourier transform as

$$(6.39) \quad F(t) = \frac{1}{(2\pi)^M} \int_{\mathbb{R}^M} \hat{F}(k) e^{-ik \cdot t} dk_1 \cdots dk_M.$$

It then follows that

$$(6.40) \quad \int e^{-S_A} F(\tau) = \frac{1}{(2\pi)^M} \int \hat{F}(k) \left(\int e^{-S_A - ik \cdot \tau} \right) dk = F(0)$$

because $S_A + ik \cdot \tau = S_{A+iK}$ with $K = \text{diag}(k_x)_{x=1}^M$, and thus $\int e^{-S_{A+iK}} = 1$ by Lemma 6.7. \square

It is not difficult to extend the integration by parts formula for Gaussian measures, Lemma 6.2, to the present more general setting; see [11] for details. The result is the following.

LEMMA 6.10. *For $a \in \Lambda$, for $C = A^{-1}$ with positive Hermitian part, and for forms F for which the integrals exist,*

$$(6.41) \quad \int e^{-S_A} \bar{\varphi}_a F = \sum_{x \in \Lambda} C_{ax} \int e^{-S_A} \frac{\partial F}{\partial \varphi_x}.$$

6.5. Integral representation for self-avoiding walk. Let Λ be a finite set and let $a, b \in \Lambda$. In Section 6.2, we showed that the integral

$$(6.42) \quad \int \bar{\varphi}_a \varphi_b \prod_{x \neq a, b} (1 + \varphi_x \bar{\varphi}_x) d\mu_C$$

is the generating function for SAWs in a background of self-avoiding loops. The following theorem shows that the loops are eliminated if we replace the factors $(1 + \varphi_x \bar{\varphi}_x)$ by $(1 + \tau_x) = (1 + \varphi_x \bar{\varphi}_x + \psi_x \bar{\psi}_x)$ and replace the Gaussian measure $d\mu_C$ by e^{-S_A} with $A = C^{-1}$.

THEOREM 6.11. *For $C = A^{-1}$ with positive Hermitian part, and for $a, b \in \Lambda$,*

$$(6.43) \quad \sum_{\omega \in \mathcal{S}_{a,b}(\Lambda)} C^\omega = \int e^{-S_A} \bar{\varphi}_a \varphi_b \prod_{x \neq a, b} (1 + \tau_x).$$

PROOF. Exactly as in Section 6.2, but now using the integration by parts formula of Lemma 6.10, we obtain

$$(6.44) \quad \int e^{-S_A} \bar{\varphi}_a \varphi_b \prod_{x \neq a, b} (1 + \tau_x) = \sum_{\omega \in \mathcal{S}_{a,b}(\Lambda)} C^\omega \int e^{-S_A} \prod_{x \in \Lambda \setminus \omega} (1 + \tau_x).$$

However, the integral on the right-hand side, which formerly generated loops, is now equal to 1 by Lemma 6.9. \square

7. Renormalisation group analysis in dimension 4

The integral representation of Theorem 6.11 opens up the following possibility for studying SAWs on \mathbb{Z}^d : approximate \mathbb{Z}^d by a large finite set Λ , rewrite the SAW two-point function as an integral as in (6.43), and apply methods of analysis to compute the asymptotic behaviour of the integral uniformly in the limit $\Lambda \nearrow \mathbb{Z}^d$. In this section, we sketch how such a program can be carried out for a particular model of continuous-time weakly SAW on the 4-dimensional lattice \mathbb{Z}^4 , using a variant of Theorem 6.11. In this approach, once the integral representation has been invoked, the original SAWs no longer appear and play no further role in the analysis. The method of proof is a rigorous renormalisation group method [12, 13]. There is work in progress, not discussed further here, to attempt to extend this

program to a particular spread-out version of the discrete-time strictly SAW model on \mathbb{Z}^d using Theorem 6.11.

We begin in Section 7.1 with the definition of the continuous-time weakly SAW and a statement of the main result for its two-point function, followed by some commentary on related results. The approximation of the two-point function on \mathbb{Z}^d by a two-point function on a d -dimensional finite torus Λ is discussed in Section 7.2, and the integral representation of the two-point function on Λ is explained in Section 7.3. The discussion of integration of differential forms from Section 6.4 is developed further in Section 7.4. At this point, the stage is set for the application of the renormalisation group method, and this is described briefly in Sections 7.5–7.7. A more extensive account of all this can be found in [12, 13].

7.1. Continuous-time weakly self-avoiding walk. The definition of the discrete-time weakly self-avoiding walk was given in Section 1.2. With an unimportant change in our conventions, and writing $z = e^{-\nu}$ and using the parameter $g > 0$ of (1.6) rather than λ , the two-point function (1.31) can be rewritten as

$$(7.1) \quad G_\nu^{(g),\text{DT}}(x) = \sum_{n=0}^{\infty} \sum_{\omega \in \mathcal{W}_n(0,x)} \exp\left(-g \sum_{i,j=0}^n 1_{\{\omega(i)=\omega(j)\}}\right) e^{-\nu n},$$

where “DT” emphasises the fact that the walks are in discrete time. The *local time* at $v \in \mathbb{Z}^d$ is defined as the number of visits to v up to time n , i.e.,

$$(7.2) \quad L_{v,n} = L_{v,n}(\omega) = \sum_{i=0}^n 1_{\{\omega(i)=v\}}.$$

Note that $\sum_{v \in \mathbb{Z}^d} L_{v,n} = n$ is independent of the walk ω , and that

$$(7.3) \quad \sum_{v \in \mathbb{Z}^d} L_{v,n}^2 = \sum_{v \in \mathbb{Z}^d} \sum_{i,j=0}^n 1_{\{\omega(i)=v\}} 1_{\{\omega(j)=v\}} = \sum_{i,j=0}^n 1_{\{\omega(i)=\omega(j)\}}.$$

Thus, writing $z = e^{-\nu}$, the two-point function can be rewritten as

$$(7.4) \quad G_\nu^{(g),\text{DT}}(x) = \sum_{n=0}^{\infty} \sum_{\omega \in \mathcal{W}_n(0,x)} e^{-g \sum_{v \in \mathbb{Z}^d} L_{v,n}^2} e^{-\nu n}.$$

The two-point function of the continuous-time weakly SAW is a modification of (7.4) in which the underlying random walk model has *continuous*, rather than discrete, time. To define the modification, we consider the *continuous-time* random walk X which takes nearest-neighbour steps like the usual SRW, but whose jumps occur after independent $\text{Exp}(2d)$ holding times at each vertex. In other words, the steps occur at the events of a rate- $2d$ Poisson process, rather than at integer times. We write \mathbb{E}_0 for the expectation associated to the process X started at $X(0) = 0 \in \mathbb{Z}^d$. The local time of X at v up to time T is now defined by

$$(7.5) \quad L_{v,T} = \int_0^T 1_{\{X(s)=v\}} ds.$$

The probabilistic structure of (1.7)–(1.9) extends naturally to the continuous-time setting. With this in mind, we define the *two-point function* of continuous-time weakly SAW by

$$(7.6) \quad G_\nu^{(g)}(x) = \int_0^\infty \mathbb{E}_0(e^{-g \sum_v L_{v,T}^2} 1_{\{X(T)=x\}}) e^{-\nu T} dT;$$

this is a natural modification of (7.1). The continuous-time SAW is predicted to lie in the same universality class as the discrete-time SAW.

Using a subadditivity argument as in Section 1.3, it is not difficult to see that the limit

$$(7.7) \quad \lim_{T \rightarrow \infty} \left(\mathbb{E}_0(e^{-g \sum_v L_{v,T}^2}) \right)^{1/T} = e^{\nu_c(g)}$$

exists, for some $\nu_c(g) \leq 0$. We leave it as an exercise to show that $\nu_c(g) > -\infty$. In particular, $G_\nu^{(g)}(x)$ is well-defined for $\nu > \nu_c(g)$. The following theorem of Brydges and Slade [12, 13] shows that the critical exponent η is equal to 0 for this model, in dimensions $d \geq 4$.

THEOREM 7.1. *Let $d \geq 4$. For $g \geq 0$ sufficiently small, there exists $c_g > 0$ such that*

$$(7.8) \quad G_{\nu_c(g)}^{(g)}(x) = \frac{c_g}{|x|^{d-2}}(1 + o(1)) \quad \text{as } |x| \rightarrow \infty.$$

Theorem 7.1 should be compared with the result of Theorem 4.2 for $d \geq 5$. The main point in Theorem 7.1 is the inclusion of the upper critical dimension $d = 4$. In particular, there is no logarithmic correction to the leading asymptotic behaviour of the critical two-point function when $d = 4$. The case $g = 0$ is the classical result that the SRW Green function obeys $G_0^{(0)}(x) \sim c_0|x|^{-(d-2)}$, which in fact holds in all dimensions $d > 2$.

The proof of Theorem 7.1 is based on an integral representation combined with a rigorous renormalisation group method, and is inspired by the methods used in [5, 9, 10] for the continuous-time weakly self-avoiding walk on the 4-dimensional *hierarchical lattice*. The hierarchical lattice is a modification of the lattice \mathbb{Z}^d that is particularly amenable to a renormalisation group approach. It is predicted that the models on the hierarchical lattice and \mathbb{Z}^d lie in the same universality class. Strong evidence for this is the result of Brydges and Imbrie [9] that on the 4-dimensional hierarchical lattice the typical end-to-end distance after time T is given, for small $g > 0$ and as $T \rightarrow \infty$, by

$$(7.9) \quad \frac{\mathbb{E}_0(|\omega(T)| e^{-g \sum_v L_{v,T}^2})}{\mathbb{E}_0(e^{-g \sum_v L_{v,T}^2})} = c T^{1/2} (\log T)^{1/8} \left[1 + \frac{\log \log T}{32 \log T} + O\left(\frac{1}{\log T}\right) \right].$$

This matches the prediction (1.29) for \mathbb{Z}^4 . There are related results by Hara and Ohno [32], proved with a completely different renormalisation group approach, for the critical two-point function, susceptibility and correlation length of the *discrete-time* weakly self-avoiding walk on the d -dimensional hierarchical lattice for $d \geq 4$.

Recently, Mitter and Scoppola [60] used the integral representation and renormalisation group analysis to study a continuous-time weakly self-avoiding walk with long-range steps. In the model of [60], each step of length r has a weight decaying like $r^{-d-\alpha}$, with $\alpha = \frac{1}{2}(3 + \epsilon)$ for small $\epsilon > 0$, in dimension $d = 3$. This is *below* the upper critical dimension $2\alpha = 3 + \epsilon$ (recall the discussion below Theorem 4.2). The main result is a control of the renormalisation group trajectory, a first step towards the computation of the asymptotic behaviour of the critical two-point function below the upper critical dimension. This is a rigorous version, for the weakly self-avoiding walk, of the expansion in $\epsilon = 4 - d$ discussed in [72].

7.2. Finite-volume approximation. Integral representations of the type discussed in Section 6.5 are for walks on a finite set. In preparation for the integral representation, we first discuss the approximation of the two-point function $G_{\nu_c}^{(g)}(x)$ on \mathbb{Z}^d by a two-point function on the finite torus $\Lambda = \mathbb{Z}^d/R\mathbb{Z}^d$ with side length $R \in \mathbb{Z}_+$. For later convenience, we will always take $R = L^N$ with L a large dyadic integer. The parameter g is regarded as a fixed positive number and will sometimes be omitted in what follows, to simplify the notation. We denote by G^Λ the natural modification of (7.6) in which the random walk on \mathbb{Z}^d is replaced by the random walk on Λ .

THEOREM 7.2. *Let $d \geq 1$, $g > 0$, and $x \in \mathbb{Z}^d$. Then for all $\nu \geq \nu_c$,*

$$(7.10) \quad G_\nu(x) = \lim_{\nu' \searrow \nu} \lim_{N \rightarrow \infty} G_{\nu'}^\Lambda(x),$$

where, on the right-hand side, x is the canonical representative of x in Λ for L^N large compared to x .

PROOF. This follows from a version of the Simon–Lieb inequality [67, 56] for the continuous-time weakly self-avoiding walk. In the problems of Section 7.8 below, we develop the corresponding argument in the discrete-time setting. With a little more work, the same approach can be adapted to continuous time. \square

We are most interested in the case $\nu = \nu_c$ in Theorem 7.2. The theorem allows for the study of the critical two-point function on \mathbb{Z}^d via the subcritical two-point function in finite volume, provided sufficient control is maintained to take the limits. Since SRW is recurrent in finite volume, its Green function is infinite, and the flexibility of taking ν slightly larger than ν_c helps bypass this concern.

7.3. Integral representation. We recall the introduction of the boson field $(\varphi_x, \bar{\varphi}_x)$ in (6.20) and the fermion field $(\psi_x, \bar{\psi}_x)$ in (6.22), and now index these fields with x in the torus $\Lambda = \mathbb{Z}^d/L^N\mathbb{Z}^d$. We also recall from (6.26) the definition, for $x \in \Lambda$, of the differential form

$$(7.11) \quad \tau_x = \varphi_x \bar{\varphi}_x + \psi_x \bar{\psi}_x.$$

The Laplacian Δ applies to the boson and fermion fields according to

$$(7.12) \quad (\Delta\varphi)_x = \sum_{y:y \sim x} (\varphi_y - \varphi_x), \quad (\Delta\psi)_x = \sum_{y:y \sim x} (\psi_y - \psi_x),$$

where the sum is over the neighbours y of x in the torus Λ . We also define the differential forms

$$(7.13) \quad \tau_{\Delta,x} = \frac{1}{2}(\varphi_x(-\Delta\bar{\varphi})_x + (-\Delta\varphi)_x\bar{\varphi}_x + \psi_x(-\Delta\bar{\psi})_x + (-\Delta\psi)_x\bar{\psi}_x).$$

The following theorem is proved in [9]; see also [11, Theorem 5.1] for a self-contained proof. Its requirement that $G_\nu^\Lambda(x) < \infty$ for large Λ is a consequence of Theorem 7.2.

THEOREM 7.3. *For $\nu > \nu_c$ and $0, x \in \Lambda$, and for Λ large enough that $G_\nu^\Lambda(x) < \infty$, the finite-volume two-point function has the integral representation*

$$(7.14) \quad G_\nu^\Lambda(x) = \int e^{-\sum_{v \in \Lambda} (\tau_{\Delta,v} + g\tau_v^2 + \nu\tau_v)} \bar{\varphi}_0 \varphi_x.$$

It is the goal of the method to show that the infinite-volume critical two-point function is asymptotically equal to a multiple of the inverse Laplacian on \mathbb{Z}^d , for $d \geq 4$. To exhibit an explicit factor to account for this multiple, we introduce a parameter $z_0 > -1$ by making the change of variables $\varphi_x \mapsto (1 + z_0)^{1/2} \varphi_x$. With this change of variables, the integral representation (7.14) becomes

$$(7.15) \quad G_\nu^\Lambda(x) = (1 + z_0) \int e^{-S(\Lambda)} e^{-\tilde{V}_0(\Lambda)} \bar{\varphi}_0 \varphi_x,$$

where

$$(7.16) \quad S(\Lambda) = \sum_{v \in \Lambda} (\tau_{\Delta, v} + m^2 \tau_v),$$

$$(7.17) \quad \tilde{V}_0(\Lambda) = \sum_{v \in \Lambda} (g_0 \tau_v^2 + \nu_0 \tau_v + z_0 \tau_{\Delta, v}),$$

with

$$(7.18) \quad g_0 = (1 + z_0)^2 g, \quad \nu_0 = (1 + z_0) \nu_c, \quad m^2 = (1 + z_0)(\nu - \nu_c).$$

In particular, the limit $\nu \searrow \nu_c$ corresponds to $m^2 \searrow 0$.

It is often convenient in statistical mechanics to obtain a correlation function by differentiation of a partition function with respect to an external field, and we will follow this approach here. Introducing an *external field* $\sigma \in \mathbb{C}$, we define

$$(7.19) \quad V_0(\Lambda) = \tilde{V}_0(\Lambda) + \sigma \bar{\varphi}_0 + \bar{\sigma} \varphi_x.$$

Then the two-point function is given by

$$(7.20) \quad G_\nu^\Lambda(x) = (1 + z_0) \frac{\partial^2}{\partial \sigma \partial \bar{\sigma}} \Big|_{\sigma = \bar{\sigma} = 0} \int_{\mathbb{C}^\Lambda} e^{-S(\Lambda) - V_0(\Lambda)}.$$

Our goal now is the evaluation of the large- x asymptotic behaviour of

$$(7.21) \quad G_{\nu_c}(x) = \lim_{m^2 \searrow 0} \lim_{N \rightarrow \infty} (1 + z_0) \frac{\partial^2}{\partial \sigma \partial \bar{\sigma}} \Big|_{\sigma = \bar{\sigma} = 0} \int_{\mathbb{C}^\Lambda} e^{-S(\Lambda) - V_0(\Lambda)}.$$

For the case $\tilde{V}_0 = 0$ (so in particular $z_0 = 0$), in view of Remark 6.8 the right-hand side becomes

$$(7.22) \quad \lim_{m^2 \searrow 0} \lim_{N \rightarrow \infty} \int_{\mathbb{C}^\Lambda} e^{-S(\Lambda)} \bar{\varphi}_0 \varphi_x = \lim_{m^2 \searrow 0} \lim_{\Lambda \nearrow \mathbb{Z}^d} \int \bar{\varphi}_0 \varphi_x d\mu_{(-\Delta_\Lambda + m^2)^{-1}},$$

and by Lemma 6.3 this is equal to

$$(7.23) \quad \lim_{m^2 \searrow 0} \lim_{\Lambda \nearrow \mathbb{Z}^d} (-\Delta_\Lambda + m^2)_{0x}^{-1} = (-\Delta_{\mathbb{Z}^d})_{0x}^{-1} \sim c_0 |x|^{-(d-2)}$$

(we have added subscripts to the Laplacians to emphasise where they act). The goal of the forthcoming analysis is to show that for small $g > 0$, and with the correct choice of z_0 , the effect of \tilde{V}_0 is a small perturbation in the sense that its presence does not change the power in this $|x|^{-(d-2)}$ decay.

7.4. Superexpectation. We will need some further development of the theory of integration of differential forms discussed in Section 6.4. As before, we denote the algebra of differential forms, now with index set Λ , by \mathcal{N} . Let C be a $\Lambda \times \Lambda$ matrix, with positive-definite Hermitian part, and with inverse $A = C^{-1}$. The *Gaussian superexpectation* with covariance matrix C is defined by

$$(7.24) \quad \mathbb{E}_C F = \int e^{-S_A} F \quad \text{for } F \in \mathcal{N}.$$

The name ‘‘superexpectation’’ comes from the fact that the integral representation for the two-point function is actually a *supersymmetric* field theory; supersymmetry is discussed in [11].

Note that, by Lemma 6.7 and Remark 6.8, $\mathbb{E}_C 1 = 1$, and more generally $\mathbb{E}_C f = \int f d\mu_C$ if f is a zero-form. The latter property shows that the Gaussian superexpectation extends the ordinary Gaussian expectation, and we wish to take this further. Recall the elementary fact that if $X_1 \sim N(0, \sigma_1^2)$ and $X_2 \sim N(0, \sigma_2^2)$ are independent normal random variables, then $X_1 + X_2 \sim N(0, \sigma_1^2 + \sigma_2^2)$. In particular, if $X \sim N(0, \sigma_1^2 + \sigma_2^2)$ then we can evaluate $\mathbb{E}(f(X))$ in stages as

$$(7.25) \quad \mathbb{E}(f(X)) = \mathbb{E}(\mathbb{E}(f(X_1 + X_2) | X_2)).$$

It will be a crucial ingredient of the following analysis that this has an extension to the superexpectation, as we describe next.

By definition, any form $F \in \mathcal{N}$ is a linear combination of products of factors ψ_{x_i} and $\bar{\psi}_{\bar{x}_i}$, with $x_i, \bar{x}_i \in \Lambda$ and with coefficients given by functions of φ and $\bar{\varphi}$. The coefficients may also depend on the external field $(\sigma, \bar{\sigma})$, but we leave the dependence on $\sigma, \bar{\sigma}$ implicit in the notation. We also define an algebra \mathcal{N}^\times with twice as many fields as \mathcal{N} , namely with boson fields (ϕ, ξ) and fermion fields (ψ, η) , where $\phi = (\varphi, \bar{\varphi})$, $\xi = (\zeta, \bar{\zeta})$, $\psi = \frac{1}{\sqrt{2\pi i}}(d\varphi, d\bar{\varphi})$, $\eta = \frac{1}{\sqrt{2\pi i}}(d\zeta, d\bar{\zeta})$. For a form $F = f(\varphi, \bar{\varphi})\psi^x \bar{\psi}^y$ (where ψ^x denotes a product $\psi_{x_1} \cdots \psi_{x_j}$), we define

$$(7.26) \quad \theta F = f(\varphi + \xi, \bar{\varphi} + \bar{\xi})(\psi + \eta)^x (\bar{\psi} + \bar{\eta})^y,$$

and we extend this to a map $\theta : \mathcal{N} \rightarrow \mathcal{N}^\times$ by linearity. Then we understand the map $\mathbb{E}_C \circ \theta : \mathcal{N} \rightarrow \mathcal{N}$ as the integration with respect to the *fluctuation fields* ξ and η , with the fields ϕ and ψ left fixed. This is like a conditional expectation. However, this is not standard probability theory, since \mathbb{E}_C does not arise from a probability measure and takes values in the (non-commutative) algebra of forms.

The superexpectation has the following important convolution property, analogous to (7.25) (see [9, 13]).

PROPOSITION 7.4. *Let $F \in \mathcal{N}$, and suppose that C_1 and C' have positive-definite Hermitian parts. Then*

$$(7.27) \quad \mathbb{E}_{C'+C_1} F = \mathbb{E}_{C'}(\mathbb{E}_{C_1} \theta F).$$

Suppose C and C_j , $j = 1, \dots, N$, are $\Lambda \times \Lambda$ matrices with positive-definite Hermitian parts, such that

$$(7.28) \quad C = \sum_{j=1}^N C_j.$$

Then, by the above proposition,

$$(7.29) \quad \mathbb{E}_C F = (\mathbb{E}_{C_N} \circ \mathbb{E}_{C_{N-1}} \theta \circ \cdots \circ \mathbb{E}_{C_1} \theta) F.$$

In the next section, we describe a particular choice of the decomposition (7.28), which will allow us to control the progressive integration in (7.29).

7.5. Decomposition of the covariance. Our goal is to compute the large- x asymptotic behaviour of the two-point function using (7.20), which we can now rewrite as

$$(7.30) \quad G_\nu^\Lambda(x) = (1 + z_0) \frac{\partial^2}{\partial \sigma \partial \bar{\sigma}} \Big|_{\sigma = \bar{\sigma} = 0} \mathbb{E}_C e^{-V_0(\Lambda)},$$

with $C = (-\Delta + m^2)^{-1}$. The Laplacian is on the torus Λ , and we must take the limits as Λ approaches \mathbb{Z}^d and m^2 approaches zero, so C is an approximation to $(-\Delta_{\mathbb{Z}^d})^{-1}$. The operator $(-\Delta_{\mathbb{Z}^d})^{-1}$ decays as $|x|^{-2}$ in dimension $d = 4$, and such long-range correlations make the analysis difficult. The renormalisation group approach takes the long-range correlations into account progressively, by making a good decomposition of the covariance C into a sum of terms with *finite* range, together with progressive integration as in (7.29). The particular decomposition used is given in the following theorem, which extends a result of Brydges, Guadagni and Mitter [8]; see also [4, 13]. In its statement, $\nabla_x^\alpha = \nabla_{x_1}^{\alpha_1} \cdots \nabla_{x_d}^{\alpha_d}$ for a multi-index $\alpha = (\alpha_1, \dots, \alpha_d)$, where ∇_{x_k} denotes the finite-difference operator $\nabla_{x_k} f(x, y) = f(x + e_k, y) - f(x, y)$.

THEOREM 7.5. *Let $d > 2$ and $N \in \mathbb{Z}_+$, and let Λ be the torus $\mathbb{Z}^d / L^N \mathbb{Z}^d$, with L a sufficiently large dyadic integer. Let $m^2 > 0$ and let $C = (-\Delta + m^2)^{-1}$ on Λ . There exist positive-definite $\Lambda \times \Lambda$ matrices C_1, \dots, C_N such that:*

- (a) $C = \sum_{j=1}^N C_j$,
- (b) $C_j(x, y) = 0$ if $|x - y| \geq \frac{1}{2}L^j$,
- (c) for multi-indices α, β with ℓ^1 norms $|\alpha|_1, |\beta|_1$ at most some fixed value p , and for $j < N$,

$$(7.31) \quad |\nabla_x^\alpha \nabla_y^\beta C_j(x, y)| \leq cL^{-(j-1)(2[\phi] - (|\alpha|_1 + |\beta|_1))},$$

where $[\phi] = \frac{1}{2}(d - 2)$, and c is independent of j and N .

The decomposition in Theorem 7.5(a) is called a *finite-range* decomposition because of item (b): the covariance C_j has range $\frac{1}{2}L^j$, and fields at points separated beyond that range are uncorrelated under \mathbb{E}_{C_j} .

To compute the important expectation $\mathbb{E}_C e^{-V_0(\Lambda)}$ in (7.30), we use Theorem 7.5 and Proposition 7.4 to evaluate it progressively. Namely, if we define

$$(7.32) \quad Z_0 = e^{-V_0(\Lambda)}, \quad Z_{j+1} = \mathbb{E}_{C_{j+1}} \theta Z_j \quad (j + 1 < N), \quad Z_N = \mathbb{E}_{C_N} Z_{N-1},$$

then the desired expectation is equal to $Z_N = \mathbb{E}_C e^{-V_0(\Lambda)}$. Thus we are led to study the recursion $Z_j \mapsto Z_{j+1}$.

In the expectation $Z_{j+1} = \mathbb{E}_{C_{j+1}} \theta Z_j$, on the right-hand side we write $\varphi_j = \varphi_{j+1} + \zeta_{j+1}$, as in (7.26), and similarly for $\bar{\varphi}_j, d\varphi_j, d\bar{\varphi}_j$. The expectation $\mathbb{E}_{C_{j+1}} \theta$ integrates out $\zeta_{j+1}, \bar{\zeta}_{j+1}, d\zeta_{j+1}, d\bar{\zeta}_{j+1}$ leaving dependence of Z_{j+1} on $\varphi_{j+1}, \bar{\varphi}_{j+1}, d\varphi_{j+1}, d\bar{\varphi}_{j+1}$. This process is repeated. The ζ_j fields that are integrated out are the fluctuation fields.

It follows from Remark 6.8 and Lemma 6.3 that $\mathbb{E}_{C_{j+1}} |\zeta_{j,x}|^2 = C_{j+1}(x, x)$. With Theorem 7.5(c), this indicates that the typical size of the fluctuation field ζ_j is of order $L^{-j[\phi]}$; the number $[\phi] = \frac{1}{2}(d - 2)$ is referred to as the *scaling dimension* or *engineering dimension* of the field. Moreover, Theorem 7.5(c) also indicates that

the derivative of $\zeta_{j,x}$ is typically smaller than the field itself by a factor L^{-j} , so that the fluctuation field remains approximately constant over a distance L^j .

To make systematic use of this behaviour of the fields, we introduce nested pavings of Λ by sets of *blocks* \mathcal{B}_j on scales $j = 0, \dots, N$. The blocks in \mathcal{B}_0 are simply the points in Λ . The blocks in \mathcal{B}_1 form a disjoint paving of Λ by boxes of side L . More generally, each block in \mathcal{B}_j has side L^j and consists of L^d disjoint blocks in \mathcal{B}_{j-1} . A *polymer* on scale j is any union of blocks in \mathcal{B}_j , and we denote the set of scale- j polymers by \mathcal{P}_j . (This terminology is standard but these polymers have nothing to do with physical polymers or random walks, they merely provide a means of organising subsets in the pavings of the torus.)




FIGURE 11. The four small shaded squares represent a polymer in \mathcal{P}_0 , and the three larger shaded squares represent its closure in \mathcal{P}_1 .

For a block $B \in \mathcal{B}_j$, the above considerations concerning the typical size of the fluctuation field suggest that, at each of the L^{dj} points $x \in B$, $\zeta_{j,x}$ has typical size $L^{-j[\phi]}$, and hence

$$(7.33) \quad \sum_{x \in B} \zeta_{j,x}^p \approx L^{dj} L^{-pj[\phi]} = L^{(d-p[\phi])j}.$$

The above sum is *relevant* (growing exponentially in j) for $p[\phi] < d$, *irrelevant* (decaying exponentially in j) for $p[\phi] > d$, and *marginal* (neither growing or decaying) for $p[\phi] = d$. Since $\tau_x = \varphi_x \bar{\varphi}_x + \psi_x \bar{\psi}_x$ is quadratic in the fields, it corresponds to $p = 2$. Thus $p[\phi] = 2[\phi] = d - 2 < d$ and τ_x is relevant in all dimensions. Similarly, τ_x^2 corresponds to $p = 4$ with $p[\phi] = 4[\phi] = 2d - 4$, so that τ_x^2 is irrelevant for $d > 4$, marginal for $d = 4$, and relevant for $d < 4$. The monomial $\tau_{\Delta,x}$ is marginal in all dimensions. In fact, the three monomials τ_x^2 , τ_x and $\tau_{\Delta,x}$, which constitute the initial potential \tilde{V}_0 , are precisely the marginal and relevant local monomials that are Euclidean invariant and obey an additional symmetry between bosons and fermions called *supersymmetry* (see [11]).

7.6. The map $Z_0 \mapsto Z_1$. For an idea of how the recursion $Z_j \mapsto Z_{j+1}$ might be studied, let us take $j = 0$ and consider the map $Z_0 \mapsto Z_1 = \mathbb{E}_{C_1} \theta Z_0$.

For simplicity, we set $\sigma = \bar{\sigma} = 0$, so that $V_0 = g_0 \tau^2 + \nu_0 \tau + z_0 \tau_\Delta$ is translation invariant. As usual, the monomials in V_0 depend on the fields $\varphi, \bar{\varphi}, \psi, \bar{\psi}$. As discussed above, we decompose the field φ as $\varphi = \varphi_1 + \zeta_1$, and similarly for $\bar{\varphi}, \psi, \bar{\psi}$. The operation $\mathbb{E}_{C_1} \theta$ integrates out the fields $\zeta_1, \bar{\zeta}_1, d\zeta_1, d\bar{\zeta}_1$. Recall that, by definition, \mathcal{P}_0 is the set of subsets of Λ . We write $I_0(x) = e^{-V_0(x)}$, and, for $X \in \mathcal{P}_0$, write $I_0^X = \prod_{x \in X} I_0(x) = e^{-V_0(X)}$ where $V_0(X) = \sum_{x \in X} V_0(x)$. In this notation, the dependence on the fields is left implicit. Let

$$(7.34) \quad V_1 = g_1 \tau^2 + \nu_1 \tau + z_1 \tau_\Delta$$

denote a modification of V_0 in which the coupling constants in V_0 have been adjusted, or *renormalised*, to some new values g_1, ν_1, z_1 . This is the origin of the term “renormalisation” in the renormalisation group. We set $I_1^X = e^{-V_1(X)}$, but with the fields in V_1 given by $\varphi_1, \bar{\varphi}_1, d\varphi_1, d\bar{\varphi}_1$. Let $\delta I_1^X = \prod_{x \in X} (I_1(x) - \theta I_0(x))$; this is an element of \mathcal{N}^\times since I_1 depends on the fields φ_1 and so on, while θI_0 depends on $\varphi_1 + \zeta_1$ and so on.

Then we obtain

$$(7.35) \quad \begin{aligned} Z_1(\Lambda) &= \mathbb{E}_{C_1} \theta I_0(\Lambda) = E_{C_1} \prod_{x \in \Lambda} (I_1(x) + \delta I_1(x)) \\ &= \mathbb{E}_{C_1} \sum_{X \in \mathcal{P}_0} I_1^{\Lambda \setminus X} \delta I_1^X = \sum_{X \in \mathcal{P}_0} I_1^{\Lambda \setminus X} \mathbb{E}_{C_1} \delta I_1^X. \end{aligned}$$

Here we have expressed Z_1 as a sum over a polymer on scale 0; we wish to express it as a sum over a polymer on scale 1. To this end, for a polymer X on scale 0, we define the *closure* \bar{X} to be the smallest polymer on scale 1 containing X : see Figure 11. We can now write

$$(7.36) \quad Z_1(\Lambda) = \sum_{U \in \mathcal{P}_1} I_1^{\Lambda \setminus U} K_1(U),$$

where

$$(7.37) \quad K_1(U) = \sum_{X \in \mathcal{P}_0: \bar{X}=U} I_1^{U \setminus X} \mathbb{E}_{C_1} \delta I_1^X.$$

DEFINITION 7.6. For $j = 0, 1, 2, \dots, N$, and for $F, G : \mathcal{P}_j \rightarrow \mathcal{N}_{\text{even}}$, where $\mathcal{N}_{\text{even}}$ denotes the forms of even degree, the *circle product* of F, G is

$$(7.38) \quad (F \circ G)(\Lambda) = \sum_{U \in \mathcal{P}_j(\Lambda)} F(\Lambda \setminus U) G(U).$$

Note that the circle product depends on the scale j .

The circle product is associative and commutative (the latter due to the restriction to forms of even degree). With the circle product, we can encode the formula (7.36) compactly as $Z_1(\Lambda) = (I_1 \circ K_1)(\Lambda)$, with the convention that $I_1(U) = I_1^U$. The identity element for the circle product is $1_{\{U=\emptyset\}}$. Thus, if we define $K_0(X) = 1_{\{X=\emptyset\}}$, then $Z_0(\Lambda) = I_0(\Lambda) = (I_0 \circ K_0)(\Lambda)$.

All later stages of the recursion proceed inductively from $Z_j = (I_j \circ K_j)(\Lambda)$. The interaction I_j continues to be defined by a potential V_j , but the form of the dependence will not, in general, be as simple as $I = e^{-V}$. The interaction does, however,

obey $I_j(X) = \prod_{B \in \mathcal{B}_j(X)} I_j(B)$, for all $X \in \mathcal{P}_j$ and for all j . The following factorisation property of K_1 , which can be verified from (7.37), allows the induction to proceed. If $U \in \mathcal{P}_1$ has connected components U_1, \dots, U_k , then $K_1(U) = \prod_{i=1}^k K_1(U_i)$; the notion of connectivity here includes blocks touching at a corner. The induction will preserve this key property for K_j and \mathcal{P}_j , for all j .

7.7. Remaining steps in the proof. Our goal is to prove Theorem 7.1. According to (7.21), we need to show that there is a choice of z_0 such that, for g small and positive,

$$(7.39) \quad G_{\nu_c}(x) = \lim_{m^2 \searrow 0} \lim_{N \rightarrow \infty} (1 + z_0) \frac{\partial^2}{\partial \sigma \partial \bar{\sigma}} \Big|_{\sigma = \bar{\sigma} = 0} Z_N(\Lambda) \sim c_g |x|^{-(d-2)}.$$

In particular, we see from this that the correct choice of z_0 will appear in the value of the constant c_g . The remaining steps in the proof of (7.39) are summarised, imprecisely, as follows. Much is left unsaid here, and details can be found in [13].

THEOREM 7.7. *Let $d \geq 4$, and let $g > 0$ be sufficiently small. There is a choice of V_1, \dots, V_N given, for $X \subset \Lambda$, by*

$$(7.40) \quad V_j(X) = \sum_{v \in X} (g_j \tau_v^2 + \nu_j \tau_v + z_j \tau_{\Delta, v}) + \lambda_j (\sigma \bar{\varphi}_0 + \bar{\sigma} \varphi_x) + q_j^2 \sigma \bar{\sigma},$$

with V_j determining I_j , and a choice of K_1, \dots, K_N with $K_j : \mathcal{P}_j \rightarrow \mathcal{N}$ obeying the key factorisation property mentioned above, such that

$$(7.41) \quad Z_j(\Lambda) = (I_j \circ K_j)(\Lambda)$$

obeys the recursion $Z_{j+1} = \mathbb{E}_{C_{j+1}} \theta Z_j$. Moreover, $(V_j, K_j)_{0 \leq j \leq N}$ obeys the flow equations

$$(7.42) \quad g_{j+1} = g_j - c g_j^2 + r_{g,j}$$

$$(7.43) \quad \nu_{j+1} = \nu_j + 2g_j C_{j+1}(0, 0) + r_{\nu,j}$$

$$(7.44) \quad z_{j+1} = z_j + r_{z,j}$$

$$(7.45) \quad K_{j+1} = r_{K,j}$$

where the r terms represent error terms. Further equations define the evolution of λ_j and q_j .

The previous theorem represents the recursion $Z_j \mapsto Z_{j+1}$ as a dynamical system. A fixed-point theorem is used to make the correct choice of the initial value z_0 so that the r terms remain small on all scales, and so that (g_j, ν_j, z_j, K_j) flows to $(0, 0, 0, 0)$. The latter is referred to as *infrared asymptotic freedom*, and is the effect anticipated below (7.23). This final ingredient is summarised in the following theorem.

THEOREM 7.8. *If $g > 0$ is sufficiently small (independent of N and m^2), there exists z_0 such that*

$$(7.46) \quad \lim_{m^2 \searrow 0} \lim_{N \rightarrow \infty} V_N = \lambda_\infty (\sigma \bar{\varphi}_0 + \bar{\sigma} \varphi_x) + q_\infty \sigma \bar{\sigma},$$

with $\lambda_\infty > 0$ and, as $x \rightarrow \infty$, $q_\infty \sim \lambda_\infty^2 (-\Delta_{\mathbb{Z}^d})_{0x}^{-1}$. Moreover, in an appropriately defined Banach space,

$$(7.47) \quad \lim_{m^2 \searrow 0} \lim_{N \rightarrow \infty} K_N(\Lambda) = 0.$$

At scale N there are only two polymers, namely the single block Λ and the empty set \emptyset . By definition, $I_N(\emptyset) = K_N(\emptyset) = 1$. Also, the field has been entirely integrated out at scale N , and from Theorem 7.8 and the definition of the circle product, we obtain

$$(7.48) \quad Z_N(\Lambda) = I_N(\Lambda) + K_N(\Lambda) \approx I_N(\Lambda) \approx e^{-q_N \sigma \bar{\sigma}}.$$

Let $z_0^* = \lim_{m^2 \searrow 0} z_0$. With (7.39) and $q_N \rightarrow q_\infty$, this gives

$$(7.49) \quad G_{\nu_c}(x) = (1 + z_0^*)q_\infty \sim (1 + z_0^*)\lambda_\infty^2 (-\Delta_{\mathbb{Z}^d})_{0x}^{-1} \sim (1 + z_0^*)\lambda_\infty^2 c_0 |x|^{-(d-2)}.$$

This is the desired conclusion of Theorem 7.1.

7.8. Tutorial. These problems develop a proof of the discrete-time version of Theorem 7.2. The proof makes use of a Simon–Lieb inequality—this is now a generic term for inequalities of the sort introduced in [67, 56] for the Ising model. The approach developed here can be adapted to prove Theorem 7.2.

Let Γ represent either $\Gamma = \mathbb{Z}^d$ or the discrete torus $\Gamma = \mathbb{Z}^d/R\mathbb{Z}^d$. Let \mathbb{E}_x denote the expectation for the usual discrete-time SRW on Γ , which we denote now by $(X_n)_{n \geq 0}$, starting at x . Let $I_{m,n}$ denote the number of self-intersections of X between times m and n :

$$(7.50) \quad I_{m,n} = \sum_{m \leq i < j \leq n} 1_{\{X_i = X_j\}}, \quad I_n = I_{0,n}.$$

We define the two-point function of the weakly SAW in the domain $D \subset \Gamma$ by

$$(7.51) \quad G_{\nu,D}(x,y) = \sum_{n \geq 0} \mathbb{E}_x(e^{-gI_n} 1_{\{X_n = y, n < T_D\}}) e^{-\nu n}, \quad x, y \in \Gamma, \nu \in \mathbb{R},$$

where $T_D = \inf\{n \geq 0 : X_n \notin D\}$ is the exit time of D . We define the boundary $\partial D = \{x \notin D : \exists y \in D \text{ s.t. } x \sim y\}$, and the closure $\bar{D} = D \cup \partial D$. The two-point function on the entire graph is written as G_ν rather than $G_{\nu,\Gamma}$. Let $c_n(x,y) = \mathbb{E}_x(e^{-gI_n} 1_{\{X_n = y\}})$, let $c_n = \sum_{y \in \Gamma} c_n(0,y)$, and define the susceptibility by

$$(7.52) \quad \chi(\nu) = \sum_{y \in \Gamma} G_\nu(0,y) = \sum_{n \geq 0} c_n e^{-\nu n}.$$

PROBLEM 7.1. Verify that $(c_n)_{n \geq 0}$ is a submultiplicative sequence, i.e. $c_{n+m} \leq c_n c_m$, and conclude that $\frac{1}{n} \log(c_n)$ converges to its infimum, which is ν_c by definition. In particular, notice that for $\nu < \nu_c$, $\chi(\nu) = \infty$ and for $\nu > \nu_c$, $\chi(\nu) < \infty$.

PROBLEM 7.2. Let $\chi^R(\nu)$ be the susceptibility for $\mathbb{Z}^d/R'\mathbb{Z}^d$ where $R' = 2R + 1$, and let $\chi(\nu)$ be the susceptibility for \mathbb{Z}^d . Prove that $\chi^R(\nu) \leq \chi(\nu)$ for $R' \geq 3$, and, in particular, that $\nu_c(\mathbb{Z}^d) \geq \nu_c(\mathbb{Z}^d/R\mathbb{Z}^d)$. Here, $\nu_c(\Gamma)$ denotes the critical point of the weakly SAW on Γ .

PROBLEM 7.3. Prove the following version of the Simon–Lieb inequality for the discrete-time weakly SAW on Γ . Given $D \subset \Gamma$, show that

$$(7.53) \quad G_\nu(x,y) - G_{\nu,D}(x,y) \leq \sum_{z \in \partial D} G_{\nu,\bar{D}}(x,z) G_\nu(z,y).$$

Note that if $x \in D$ and $y \in D^c$, then $G_{\nu,D}(x,y) = 0$.

The following problem provides an approach to proving exponential decay of a subcritical two-point function which, unlike Proposition 1.3, adapts well to the continuous-time setting.

PROBLEM 7.4. Let $\Lambda_R = \{-R+1, \dots, R\}^d \subset \mathbb{Z}^d$. For $\nu > \nu_c$, $\sum_{y \in \mathbb{Z}^d} G_\nu(0, y)$ is finite, and thus $\theta = \sum_{y \in \partial \Lambda_R} G_\nu(0, y) < 1$ for R sufficiently large. Conclude from Problem 7.3 with $D = \Lambda_R$ that for $y \notin \Lambda_R$,

$$(7.54) \quad G_\nu(0, y) \leq \theta^{\lfloor |y|_\infty / (R+1) \rfloor} \sup_{x \in \mathbb{Z}^d} G_\nu(0, x).$$

PROBLEM 7.5. Let $(T_R)_{R \in \mathbb{N}}$ be a sequence of discrete tori with the vertex sets V_R embedded in \mathbb{Z}^d by $V_R = \Lambda_R$ where Λ_R is as in Problem 7.4; in particular, $V_R \subset V_{R+1}$. Let G_ν^R be the two-point function on T_R , and G_ν be the two-point function on \mathbb{Z}^d . Use Problem 7.2 and Problem 7.4 to prove that for all $\nu > \nu_c = \nu_c(\mathbb{Z}^d)$, $x, y \in \mathbb{Z}^d$,

$$(7.55) \quad G_\nu^R(x, y) \rightarrow G_\nu(x, y) \quad \text{as } R \rightarrow \infty.$$

Conclude that

$$(7.56) \quad G_{\nu_c}(x, y) = \lim_{\nu \searrow \nu_c} \lim_{R \rightarrow \infty} G_\nu^R(x, y).$$

Appendix A. Solutions to the problems

A.1. Solutions for Tutorial 1.7.

PROBLEM 1.1. Let M be an integer, and for every $n \in \mathbb{N}$, write $n = Mk + r$ with $0 \leq r < M$. Then,

$$(A.1) \quad \frac{1}{n} a_n \leq \frac{k}{n} a_M + \frac{1}{n} a_r, \quad \text{and, thus,} \quad \limsup_{n \rightarrow \infty} \frac{1}{n} a_n \leq \frac{1}{M} a_M.$$

In particular,

$$(A.2) \quad \limsup_{n \rightarrow \infty} \frac{1}{n} a_n \leq \inf_{M \in \mathbb{N}} \frac{1}{M} a_M \leq \liminf_{M \rightarrow \infty} \frac{1}{M} a_M,$$

which implies both statements of the claim. \square

PROBLEM 1.2. The number of n -step walks with steps only in positive coordinate directions is d^n . The number of walks which do not reverse direction is $2d(2d-1)^{n-1}$. Thus,

$$(A.3) \quad d^n \leq c_n \leq 2d(2d-1)^{n-1} \quad \text{and therefore} \quad d \leq \mu \leq 2d-1.$$

The upper bound can easily be improved by excluding more patterns that lead to self-intersecting walks than merely reversals of steps. For example, by considering walks which do not contain anti-clockwise ‘‘unit squares’’ (see Figure 12), we obtain

$$(A.4) \quad c_{3n+1} \leq 2d((2d-1)^3 - 1)^n = 4(26^{1/3})^{3n},$$

giving $\mu \leq 26^{1/3} < 3$. Similarly, the lower bound can be improved by considering walks that take steps either in positive coordinate directions, i.e., north or east, or in an east-north-west-north pattern: see Figure 12. It follows that

$$(A.5) \quad c_{4n} \geq (d^4 + 1)^n = (17^{1/4})^{4n},$$

where $17^{1/4} > 2$. In particular, $2 < 17^{1/4} \leq \mu \leq 26^{1/3} < 3$. \square

PROBLEM 1.3. SAWs can get trapped: see Figure 13. A trapped walk ω of length n does not arise as the restriction of a walk ρ of length $m > n$ to the first n steps. Thus, under $\mathbb{Q}_n^{(1)}$, ω has positive probability, while $\sum_{\rho > \omega} \mathbb{Q}_m^{(1)}(\rho) = 0$. \square




FIGURE 12. Left: The walk does contain a unit square. Right: The walk only takes steps east, north, or in east-north-west-north patterns (thick line).




FIGURE 13. Trapped walk.

PROBLEM 1.4. $c_n(x) = 1_{\{|x|=n\}}$, so $G_z(x) = \sum_{n \geq 0} c_n(x)z^n = z^{|x|}$, and

$$\begin{aligned} \hat{G}_z(k) &= \sum_{x \in \mathbb{Z}} z^{|x|} e^{ikx} = -1 + \sum_{n \geq 0} z^n (e^{ikn} + e^{-ikn}) \\ (A.6) \quad &= -1 + (1 - ze^{ik})^{-1} + (1 - ze^{-ik})^{-1} = \frac{1 - z^2}{1 - 2z \cos k + z^2}, \end{aligned}$$

as claimed. \square

PROBLEM 1.5. The assumption implies

$$(A.7) \quad |f((1 - 1/n)e^{i\varphi})| \leq c |1 - (1 - 1/n)e^{i\varphi}|^{-b}.$$

Note that for $\varphi \in [0, \pi/2]$,

$$(A.8) \quad |\operatorname{Re}(1 - (1 - 1/n)e^{i\varphi})| = 1 - (1 - 1/n) \cos \varphi \geq 1/n,$$

$$(A.9) \quad |\operatorname{Im}(1 - (1 - 1/n)e^{i\varphi})| = |(1 - 1/n) \sin \varphi| \geq (1 - 1/n) \frac{2\varphi}{\pi}.$$

Suppose $b > 1$. The integral is estimated using $|z|^n \geq Ce^{-1}$ for $|z| = 1 - 1/n$,

$$\begin{aligned} (A.10) \quad \frac{1}{2\pi} \int_0^{\pi/2} |f((1 - 1/n)e^{i\varphi})| d\varphi &\leq c \int_0^{\pi/2} \left(\frac{1}{n} + (1 - 1/n) \frac{2\varphi}{\pi} \right)^{-b} d\varphi \\ &= (1 - 1/n)^{-1} c \int_{1/n}^1 t^{-b} dt = (1 - 1/n)^{-1} cb(n^{b-1} - 1) \leq cn^{b-1}, \end{aligned}$$

and, since $|f(z)|$ is bounded for z bounded away from 1,

$$(A.11) \quad \frac{1}{2\pi} \int_{\pi/2}^{\pi} |f((1 - 1/n)e^{i\varphi})| d\varphi \leq c.$$

Likewise, the contributions for the interval $[\pi, 2\pi]$ are estimated and we obtain

$$(A.12) \quad |a_n| \leq cn^{b-1}.$$

The above assumed $b > 1$ but the extension to $b = 1$ is easy. \square

PROBLEM 1.6. (a) Let $T_0 = 0$ and $T_k = \inf\{n > T_{k-1} : X_n = 0\}$. Then $u = P(T_1 < \infty)$, and by induction and the strong Markov property, $P(T_k < \infty) = u^k$. It follows that

$$(A.13) \quad m = \mathbb{E}(N) = \sum_{k \geq 0} \mathbb{P}(T_k < \infty) = (1 - u)^{-1}.$$

(b) The solution relies on the formula

$$(A.14) \quad \mathbb{P}(X_n = 0) = \int_{[-\pi, \pi]^d} \hat{D}(k)^n \frac{d^d k}{(2\pi)^d}.$$

Some care is required when performing the sum over n since the best uniform bound on \hat{D}^n is 1 which is not summable. A solution is to make use of monotone convergence first, and then apply the dominated convergence theorem, as follows,

$$(A.15) \quad m = \lim_{t \nearrow 1} \sum_{n \geq 0} \mathbb{P}(X_n = 0)t^n = \lim_{t \nearrow 1} \int_{[-\pi, \pi]^d} \frac{1}{1 - t\hat{D}(k)} \frac{d^d k}{(2\pi)^d}.$$

Note that \hat{D} is a real-valued function and that

$$(A.16) \quad \frac{1}{1 - t\hat{D}(k)} \leq \frac{2}{1 - \hat{D}(k)} \quad \text{for } t \in [1/2, 1],$$

so that if $(1 - \hat{D})^{-1} \in L^1$, then the claim follows by dominated convergence. In the case that $(1 - \hat{D})^{-1} \notin L^1$, the claim follows from Fatou's lemma.

(c) $\hat{D}(k) = \sum_{j=1}^d (e^{ik_j} + e^{-ik_j}) = 2 \sum_{j=1}^d \cos(k_j)$ and thus $1 - \hat{D}(k) = O(1)|k|^2$ as $k \rightarrow 0$. Note further that

$$(A.17) \quad \int_{\mathbb{R}^d} f(|k|) dk = V_{d-1} \int_0^\infty f(r) r^{d-1} dr,$$

where V_{d-1} is the volume of the $(d-1)$ -dimensional sphere, and in particular,

$$(A.18) \quad \int_{[-\epsilon, \epsilon]^d} |k|^{-p} dk \text{ is integrable if and only if } d > p. \quad \square$$

PROBLEM 1.7. Note that

$$(A.19) \quad I = \sum_{x \in \mathbb{Z}^d} \left(\sum_{i \geq 0} 1_{\{X_i^1 = x\}} \right) \left(\sum_{j \geq 0} 1_{\{X_j^2 = x\}} \right),$$

and thus, by Parseval's theorem, if $f \in L^2(\mathbb{Z}^d)$,

$$(A.20) \quad \mathbb{E}(I) = \sum_{x \in \mathbb{Z}^d} f(x)^2 = \int_{[-\pi, \pi]^d} |\hat{f}(k)|^2 \frac{d^d k}{(2\pi)^d},$$

where

$$(A.21) \quad f(x) = \sum_{j \geq 0} \mathbb{P}\{X_j^1 = x\} = \sum_{j \geq 0} D^{*j}(x), \quad \hat{f}(k) = \sum_{j \geq 0} \hat{D}(k)^j = \frac{1}{1 - \hat{D}(k)}.$$

If $f \notin L^2(\mathbb{Z}^d)$, then both sides must be infinite. \square

A.2. Solutions for Tutorial 4.4.

PROBLEM 4.1. The graph $\{0n\}$ is connected on $[0, n]$ in the above sense but not path-connected. Also, $\{01, 12, \dots, (n-1)n\}$ is path-connected but not connected in the above sense since the open intervals $(i-1, i)$ do not overlap. \square

PROBLEM 4.2. This is an application of the identity

$$(A.22) \quad \prod_{i \in I} (1 + u_i) = \sum_{S \subset I} \prod_{i \in S} u_i$$

with I being the set of edges on $[a, b]$. \square

PROBLEM 4.3. The identity corresponds to a decomposition of $\mathcal{B}[a, b]$ by connected components. The term $K[a+1, b]$ corresponds to graphs Γ for which $a \notin \Gamma$.

So assume $a \in \Gamma$. We shall show that Γ can be written uniquely as $\Gamma = \Gamma' \cup \Gamma''$ where $\Gamma' \in \mathcal{G}[a, j]$ and $\Gamma'' \in \mathcal{B}[j, b]$ for some $j \in (a, b]$. Informally, Γ' is the connected component of Γ containing a , though we must verify that this notion is well-defined. Conversely it is clear that if $\Gamma' \in \mathcal{G}[a, j]$, $\Gamma'' \in \mathcal{B}[j, b]$ for some $j \in (a, b]$, then $\Gamma = \Gamma' \cup \Gamma'' \in \mathcal{B}[a, b]$ with $a \in \Gamma$. Then the result will follow since

$$(A.23) \quad \prod_{st \in \Gamma' \cup \Gamma''} U_{st} = \prod_{st \in \Gamma'} U_{st} \prod_{st \in \Gamma''} U_{st}.$$

Let

$$(A.24) \quad j = \min \{i \in (a, b] : i \notin (s, t) \text{ for some } st \in \Gamma\}.$$

The minimum is well defined since there can be no $st \in \Gamma$ for which $b \in (s, t)$. By construction, every edge $st \in \Gamma$ satisfies $t \leq j$ or $s \geq j$, so that we can write $\Gamma = \Gamma' \cup \Gamma''$ where $\Gamma' \in \mathcal{B}[a, j]$, $\Gamma'' \in \mathcal{B}[j, b]$. We must show that $\Gamma' \in \mathcal{G}[a, j]$, i.e., that Γ' is connected. But $\cup_{st \in \Gamma'} (s, t) = (a, j) \cap \cup_{st \in \Gamma} (s, t) = (a, j)$ by the minimality of j .

Finally we check that the decomposition $\Gamma = \Gamma' \cup \Gamma''$ is unique: this follows because if $\Gamma' \in \mathcal{G}[a, j']$ and $\Gamma'' \in \mathcal{B}[j', b]$ then the formula (A.24) recovers $j = j'$. \square

PROBLEM 4.4. The convolutions correspond to summing over the values of $\omega(1)$ and $\omega(m)$. Namely, noting that a walk ω on $[0, n]$ is equivalent to a pair of walks ω_0 on $[0, m]$ and ω_1 on $[m, n]$ with $\omega_0(m) = \omega_1(m)$, we have

$$(A.25) \quad \begin{aligned} c_n(x) &= \sum_{\omega \in \mathcal{W}_n(0, x)} K[1, n](\omega) + \sum_{m=1}^n \sum_{\omega \in \mathcal{W}_n(0, x)} J[0, m](\omega) K[m, n](\omega) \\ &= \sum_{y \in \mathbb{Z}^d} \sum_{\omega_0 \in \mathcal{W}_1(0, y)} \sum_{\substack{\omega_1: [1, n] \rightarrow \mathbb{Z}^d, \\ \omega_1(1)=y, \omega_1(n)=x}} K[1, n](\omega_1) \\ &\quad + \sum_{m=1}^n \sum_{y \in \mathbb{Z}^d} \sum_{\omega_0 \in \mathcal{W}_m(0, y)} \sum_{\substack{\omega_1: [m, n] \rightarrow \mathbb{Z}^d, \\ \omega_1(m)=y, \omega_1(n)=x}} J[0, m](\omega_0) K[m, n](\omega_1) \\ &= \sum_{y \in \mathbb{Z}^d} 1_{\{y \in \Omega\}} c_{n-1}(x-y) + \sum_{m=1}^n \sum_{y \in \mathbb{Z}^d} \pi_m(y) c_{n-m}(x-y) \end{aligned}$$

where we use the translation invariance (in time and space) of K . Since $c_1(y) = 1_{\{y \in \Omega\}}$, this is the desired equation. \square




FIGURE 14. Laces in $\mathcal{L}^{(N)}[a, b]$ for $N = 1, 2, 3, 4$, with $s_1 = a$ and $t_N = b$.

PROBLEM 4.5. Figure 14 is helpful. Note first that if L is a lace, then $s_l < s_{l+1}$ for each l . Indeed, if $s_l = s_{l+1}$, we may assume that $t_l < t_{l+1}$. But then $(s_l, t_l) \subset (s_{l+1}, t_{l+1})$ so that $L \setminus \{s_l t_l\}$ is still connected. A similar argument gives $t_l < t_{l+1}$. The requirement that L is connected implies that $a = s_1$ and $b = t_N$.

Suppose to the contrary that (1) $s_{l+1} \geq t_l$ ($1 \leq l \leq N-1$) or (2) $s_{l+2} < t_l$ ($1 \leq l \leq N-2$). In case (1), L is not connected, since $s_i \geq t_l$ for $i \geq l+1$ while $t_i \leq t_l$ for $i \leq l$. In case (2), the edge $s_{l+1} t_{l+1}$ is redundant since $(s_{l+1}, t_{l+1}) \subset (s_l, t_l) \cup (s_{l+2}, t_{l+2}) = (s_l, t_{l+2})$.

For the converse, the hypotheses imply that $\cup_{st \in L} (s, t) = (a, b)$, so L is connected. Neither $s_1 t_1$ nor $s_N t_N$ can be removed from L since they are the only edges containing the endpoints. If $s_l t_l$ is removed, $2 \leq l \leq N-1$, then $t_{l-1} \leq s_{l+1}$ implies that $\cup_{st \in L} (s, t) = (a, t_{l-1}) \cup (s_{l+1}, b) \neq (a, b)$. So $L \setminus \{st\}$ is not connected. Since connectedness is a monotone property, no strict subset of L can be connected, so L is minimally connected, i.e., a lace.

Finally the intervals are as follows: the first and last intervals are $[s_1, s_2]$ and $[t_{N-1}, t_N]$; the $2i^{\text{th}}$ interval is $[s_{i+1}, t_i]$ ($1 \leq i \leq N-1$); and the $(2i+1)^{\text{st}}$ interval is $[t_i, s_{i+2}]$, $1 \leq i \leq N-2$. The inequalities above show that the points $\{s_i, t_i\}$ do indeed form the intervals claimed, and that the intervals $[t_i, s_{i+2}]$ can be empty while the other intervals must be non-empty. \square

PROBLEM 4.6. Figure 15 is helpful. First, since necessarily $L_\Gamma \subset \Gamma$, we may assume that $L \subset \Gamma$, and we write $\Gamma = L \cup A$ with $A \cap L = \emptyset$.

Next, we reformulate the inductive procedure for selecting the edges of L_Γ . At each step, the edge $s_{i+1} t_{i+1}$ is, among all edges $st \in \Gamma$ satisfying $s < t_i$, the one




FIGURE 15. (a) A connected graph Γ and its associated lace $L = L_\Gamma$. (b) The dotted edges are compatible with the lace L . (c) The dotted edge is not compatible with the lace L .

that is *maximal* with respect to the following order relation: $st \succ s't'$ if and only if $t > t'$ or $t = t'$ and st is longer than $s't'$ (i.e., $t - s > t' - s'$).

The result follows at once from this observation. Indeed, $L_\Gamma = L$ means that at each inductive step, $s_{i+1}t_{i+1} \in L$ is the maximal edge st satisfying $s < t_i$, among all edges of $L \cup A$. This is equivalent to saying that for each $s't' \in A$, at each inductive step, $s_{i+1}t_{i+1}$ is the maximal edge among all edges of $L \cup \{s't'\}$. But this is precisely the condition that $A \subset \mathcal{C}(L)$. \square

PROBLEM 4.7. The first equation is simply a decomposition of $\Gamma \in \mathcal{G}[a, b]$ according to the value of L_Γ . The second equation follows using (A.22) because Problem 4.6 shows that Γ for which $L_\Gamma = L$ can be identified as L together with an arbitrary subset of edges from $\mathcal{C}(L)$. The last equation is immediate from the preceding ones. \square

PROBLEM 4.8. (a) By definition,

$$\begin{aligned}
 \sum_{N=1}^{\infty} (-1)^N \pi_m^{(N)}(x) &= \sum_{\omega \in \mathcal{W}_m(0,x)} \sum_{N=1}^{\infty} J^{(N)}[0, m](\omega) \\
 \text{(A.26)} \qquad \qquad \qquad &= \sum_{\omega \in \mathcal{W}_m(0,x)} J[0, m](\omega) = \pi_m(x).
 \end{aligned}$$

Each of the N factors U_{st} , $st \in L$, contributes -1 , so $\pi_m^{(N)}(x) \geq 0$.

(b) $N = 1$: The only lace with 1 edge is $L = \{0m\}$, and every edge except $0m$ is compatible with L . So $J^{(1)}[0, m]$ contains the single factor U_{0m} , and the factor $1 + U_{s't'}$ for each $s't' \neq 0m$. So a contributing ω must have $\omega(s') \neq \omega(t')$ whenever $s't' \neq 0m$, as well as $0 = \omega(0) = \omega(m) = x$. Hence $\pi_m^{(1)}(x) = 0$ for $x \neq 0$, and $\pi_m^{(1)}(0)$ is the number of m -step self-avoiding returns.

$N = 2$: For $L = \{0t_1, s_1m\}$, the factors U_{st} , $st \in L$, require that ω should start at 0, visit x (at step s_1), return to 0 (at step t_1), then return to x . The compatible edges consist of every edge except the edges of L and the edges $0t$, $t > t_1$ and sm , $s < s_1$. This implies that each of the three intervals in ω must be self-avoiding

and mutually avoiding, except for the intersections required above. (In particular, $x \neq 0$.) (Intersections of the form $\omega(0) = \omega(t)$, $t > t_1$, might not appear to be forbidden, but actually they are impossible since we require $\omega(t) \neq \omega(t_1) = \omega(0)$.)

$N = 3, 4, \dots$: As for $N = 2$, ω must have self-intersections corresponding to the edges of the lace and self-avoidance corresponding to each compatible edge. It is convenient to recall the $2N - 1$ intervals from Problem 4.5. Because of compatible edges, ω is required to be self-avoiding on each of these intervals. In addition, certain of these intervals are required to be mutually avoiding, but not all of them need be, corresponding to the fact that an edge spanning too many intervals cannot be compatible. The pattern of mutual avoidance is described as follows: for $N = 3$,

$$(A.27) \quad [1234][345]$$

and for $N = 4$,

$$(A.28) \quad [1234][3456][567]$$

where, for instance [3456] indicates that the third to sixth interval must be mutually self-avoiding, except for the required intersections. These intersections require that at the endpoints of the intervals, ω must visit the following points (for the case $N = 4$):

$$(A.29) \quad 0, x_1, 0, x_2, x_1, x_3, x_2, x_3$$

where $x_3 = x$, corresponding to the intervals $[s_1, s_2]$, $[s_2, t_1]$, $[t_1, s_3]$, $[s_3, t_2]$, $[t_2, s_4]$, $[s_4, t_3]$, $[t_3, t_4]$.

To prove the avoidance patterns amounts to analysing exactly which edges are compatible. For instance, it is easy to verify that if $s_{i+1} \leq s < t_i$ for $1 \leq i \leq N - 1$, then $st \in \mathcal{C}(L)$ if and only if $t \leq t_{i+1}$ (assuming $st \notin L$).

(c) The possibly empty intervals indicate that the 3rd, 5th, \dots , $(2N - 3)$ rd segments of the diagrams above can be empty, whereas all other segments must have non-zero length. The picture for $N = 11$ is



where the lines that are slashed are exactly the lines that are permitted to have length zero. \square

A.3. Solutions for Tutorial 5.4.

PROBLEM 5.1. (a) Using the hint and Fubini's theorem (which is applicable because $d > 2m$),

$$(A.30) \quad \begin{aligned} \int_{[-\pi, \pi]^d} \frac{1}{[1 - \hat{D}(k)]^m} \frac{d^d k}{(2\pi)^d} &= \Gamma(m)^{-1} \int_0^\infty \left(\int_{[-\pi, \pi]^d} e^{-u[1 - \hat{D}(k)]} \frac{d^d k}{(2\pi)^d} \right) u^{m-1} du \\ &= \Gamma(m)^{-1} \int_0^\infty \left(\int_{-\pi}^\pi e^{-u(1 - \cos k)/d} \frac{dk}{2\pi} \right)^d u^{m-1} du. \end{aligned}$$

The inner integral is decreasing as a function of d by Hölder's inequality.

(b) Relaxing the self-avoidance for the first j steps gives the inequality

$$(A.31) \quad H_z^{(j)}(x) \leq (z|\Omega|D)^{*j} * G_z(x),$$

and thus, by Cauchy-Schwarz,

$$(A.32) \quad \|H_z^{(j)}\|_\infty \leq \|\hat{H}_z^{(j)}\|_1 \leq (z|\Omega|)^j \|\hat{D}^j\|_2 \|\hat{G}_z\|_2.$$

The claim now follows from $z_c|\Omega| \leq a$,

$$(A.33) \quad \|\hat{D}^j\|_2 = \|\hat{D}^{2j}\|_1^{1/2} \leq O((2d)^{-j/2}),$$

and

$$(A.34) \quad \|\hat{G}_z\|_2 \leq a \|\hat{C}_{p(z)}\|_2 \leq a \|\hat{C}_{1/|\Omega|}\|_2 \leq O(1)$$

by the infrared bound (5.38) and (a).

(c) Calculating the first two terms explicitly, we obtain

$$(A.35) \quad \begin{aligned} \hat{\Pi}_z^{(1)}(0) &= (2d)z^2 + (2d)(2d-2)z^4 + \sum_{m \geq 6} \hat{\pi}_m^{(1)}(0)z^m \\ &= (2d)z^2 + (2d)(2d-2)z^4 + O((2d)^{-3}), \end{aligned}$$

where the remainder was estimated as in (b), and using the symmetry of D :

$$(A.36) \quad \begin{aligned} \sum_{m \geq j} \hat{\pi}_m^{(1)}(0)z^m &= (H_z^{(j-1)} * z|\Omega|D)(0) \\ &\leq (z|\Omega|)^j \|D^{*j} * G_z\|_\infty \leq O((2d)^{-j/2}). \end{aligned}$$

Using (5.40), we obtain

$$(A.37) \quad \hat{\Pi}_z(0) = -\hat{\Pi}_z^{(1)}(0) + O((2d)^{-2}).$$

Equation (5.42) gives $z_c = (2d)^{-1} - (2d)^{-1} \hat{\Pi}_{z_c}^{(1)}(0) = (2d)^{-1} + O((2d)^{-2})$, from which we obtain

$$(A.38) \quad \begin{aligned} z_c &= (2d)^{-1} + (2d)^{-1} \hat{\Pi}_{z_c}^{(1)}(0) + O((2d)^{-3}) \\ &= (2d)^{-1} + (2d)^{-2} + O((2d)^{-3}). \end{aligned}$$

Finally, we obtain

$$(A.39) \quad \hat{\Pi}_{z_c}^{(1)}(0) = [(2d)^{-1} + 2(2d)^{-2}] + (2d)^{-2} + O((2d)^{-3}).$$

(d) The generating function for θ -walks from 0 to x can be written as

$$(A.40) \quad \hat{\Pi}_z^{(2)}(0) = (2d)z^3 + 3(2d)(2d-2)z^5 + \sum_{m \geq 7} \hat{\pi}_m^{(2)}(0)z^m.$$

Using $z_c = (2d)^{-1} + (2d)^{-2} + O((2d)^{-3})$ from part (c), we obtain

$$(A.41) \quad (2d)z_c^3 + 3(2d)(2d-2)z_c^5 = (2d)^{-2} + O((2d)^{-3}).$$

The remainder is estimated using the fact that in a θ -walks from 0 to x of length $m \geq 7$, either two of the subwalks take just one step and the other takes at least

5 steps, or at least two of the subwalks take at least 3 steps. Thus there is a combinatorial constant K such that

$$(A.42) \quad \sum_{m \geq 7} \hat{\pi}_m^{(2)}(0) z^m \leq K \sum_e H_z^{(5)}(e) z |\Omega| D(e) z |\Omega| D(e) \\ + K \sum_x H_z^{(1)}(x) H_z^{(3)}(x) H_z^{(3)}(x).$$

The first term can be estimated by an L^∞ bound (use $z|\Omega| \leq a$ and $|\Omega| = 2d$):

$$(A.43) \quad \sum_e H_z^{(5)}(e) z |\Omega| D(-e) z |\Omega| D(e) \leq \|H_z^{(5)}\|_\infty a^2 (2d)^{-1} \\ \leq O((2d)^{-7/2}) \leq O((2d)^{-3}).$$

The second term is estimated in the spirit of (b):

$$(A.44) \quad \sum_x H_z^{(1)}(x) H_z^{(3)}(-x) H_z^{(3)}(x) \leq O((2d)^{-1/2}) (H_z^{(3)} * H_z^{(3)})(0) \\ \leq O((2d)^{-1/2}) ((z|\Omega|D)^{*6} * G_z^{*2})(0) \\ \leq O((2d)^{-1/2}) \|\hat{D}^6 \hat{G}_z^2\|_1 \\ \leq O((2d)^{-1/2}) \|\hat{D}^6\|_2 \|\hat{G}_z^2\|_2 \leq O((2d)^{-7/2}).$$

(e) Using (5.40), we obtain

$$(A.45) \quad \hat{\Pi}_{z_c}(0) = -(2d)^{-1} - 2(2d)^{-2} + O((2d)^{-3}).$$

From (5.42), it then follows that

$$(A.46) \quad z_c = (2d)^{-1} + (2d)^{-2} + 2(2d)^{-3} + O((2d)^{-4}).$$

Inverting this finally yields

$$(A.47) \quad \mu = 2d - 1 - (2d)^{-1} + O((2d)^{-2}). \quad \square$$

PROBLEM 5.2. (a) This is a straightforward calculation.

(b) This requires an extension of the diagrammatic estimates. The argument is sketched in [69, Section 5.4].

(c) Note that

$$(A.48) \quad \frac{d[z\chi(z)]^{-1}}{dz} = -[z\chi(z)]^{-2} \frac{d[z\chi(z)]}{dz} \sim -\frac{V(z_c)}{z^2} \sim -\frac{V(z_c)}{z_c^2}.$$

Integrating this asymptotic relation, we obtain

$$(A.49) \quad \lim_{z \rightarrow z_c} [z\chi(z)]^{-1} - [z\chi(z)]^{-1} \sim -\frac{V(z_c)}{z_c^2} (z_c - z).$$

The limit vanishes and thus

$$(A.50) \quad \chi(z)^{-1} \sim V(z_c)^{-1} (1 - z/z_c)^{-1}.$$

The claim then follows from the definition of $V(z_c)$ and (5.42). \square

A.4. Solutions for Tutorial 7.8.

PROBLEM 7.1. For $m, n \geq 0$,

$$\begin{aligned}
 I_{0,n+m} &= \sum_{0 \leq i < j \leq n+m} 1_{\{X_i=X_j\}} \\
 (A.51) \quad &\geq \sum_{0 \leq i < j \leq m} 1_{\{X_i=X_j\}} + \sum_{m \leq i < j \leq n+m} 1_{\{X_i=X_j\}} = I_{0,m} + I_{m,n+m}.
 \end{aligned}$$

By translation invariance and the Markov property, $I_{m,n+m}$ is independent of $I_{0,m}$ and has the same law as $I_{0,n}$. Therefore

$$(A.52) \quad c_{n+m} \leq \mathbb{E}_0(e^{-gI_{0,m}} e^{-gI_{m,m+n}}) = \mathbb{E}_0(e^{-gI_{0,m}}) \mathbb{E}_0(e^{-gI_{0,n}}) = c_m c_n$$

as claimed. The remaining statements follow since (A.52) implies that $(\log c_n)_{n \geq 0}$ is a subadditive sequence, and Lemma 1.1 can be applied. \square

PROBLEM 7.2. Note that there is a one-to-one correspondence between nearest-neighbour walks on \mathbb{Z}^d and such walks on the torus $\mathbb{Z}^d/R\mathbb{Z}^d$, $R \geq 3$, by folding a walk on \mathbb{Z}^d (the image under the canonical projection $\mathbb{Z}^d \rightarrow \mathbb{Z}^d/R\mathbb{Z}^d$), and corresponding unfolding of walks on $\mathbb{Z}^d/R\mathbb{Z}^d$ (unique for the nearest-neighbour step distribution provided $R \geq 3$). Given a walk $X = (X_n)_{n \geq 0}$ on \mathbb{Z}^d starting at 0, we denote the folded (or projected) walk by X' . Write $\Lambda_R = \{-R+1, \dots, R\}^d$ and $R' = 2R+1$; then

$$\begin{aligned}
 I_n(X) &= \sum_{0 \leq i < j \leq n} \sum_{x \in \mathbb{Z}^d} 1_{\{X_i=X_j=x\}} = \sum_{0 \leq i < j \leq n} \sum_{x \in \Lambda_R} \sum_{y \in \mathbb{Z}^d} 1_{\{X_i=X_j=x+yR'\}} \\
 &\leq \sum_{0 \leq i < j \leq n} \sum_{x \in \Lambda_R} \sum_{y_1, y_2 \in \mathbb{Z}^d} 1_{\{X_i=x+y_1R'\}} 1_{\{X_j=x+y_2R'\}} \\
 (A.53) \quad &= I_n(X'),
 \end{aligned}$$

and thus

$$(A.54) \quad \mathbb{E}(e^{-gI_n}) \geq \mathbb{E}^R(e^{-gI_n}).$$

The desired inequalities both follow from this one. \square

PROBLEM 7.3. Note that

$$(A.55) \quad G_\nu(x, y) - G_{\nu, D}(x, y) = \sum_{n=0}^{\infty} \mathbb{E}_x(e^{-gI_n} 1_{\{X_n=y, n \geq T_D\}}) e^{-\nu n},$$

and, by partitioning in T_D and X_{T_D} , we obtain

$$(A.56) \quad \mathbb{E}_x(e^{-gI_n} 1_{\{X_n=y, n \geq T_D\}}) = \sum_{z \in \partial D} \sum_{m=0}^n \mathbb{E}_x(e^{-gI_n} 1_{\{X_n=y\}} 1_{\{X_{T_D}=z\}} 1_{\{T_D=m\}}).$$

Using $I_n \geq I_m + I_{m,n}$ and the Markov property, it follows that

$$\begin{aligned}
 &\mathbb{E}_x(e^{-gI_n} 1_{\{X_n=y\}} 1_{\{X_{T_D}=z\}} 1_{\{T_D=m\}}) \\
 &\leq \mathbb{E}_x(e^{-gI_m} 1_{\{X_m=z\}} 1_{\{T_D=m\}}) e^{-gI_{m,n}} 1_{\{X_n=y\}} \\
 (A.57) \quad &= \mathbb{E}_x(e^{-gI_m} 1_{\{X_m=z\}} 1_{\{T_D=m\}}) \mathbb{E}_z(e^{-gI_{n-m}} 1_{\{X_{n-m}=y\}}).
 \end{aligned}$$

Thus, because $\{T_D = m, X_m = z\} = \{m \leq T_D, X_m = z\}$ for $z \in \partial D$,

$$\begin{aligned}
G_\nu(x, y) - G_{\nu, D}(x, y) &\leq \sum_{z \in \partial D} \sum_{n=0}^{\infty} \sum_{m=0}^n \mathbb{E}_x(e^{-gI_m} \mathbf{1}_{\{X_m=z\}} \mathbf{1}_{\{T_D=m\}}) \\
&\quad \cdot \mathbb{E}_z(e^{-gI_{n-m}} \mathbf{1}_{\{X_{n-m}=y\}}) e^{-\nu n} \\
\text{(A.58)} \qquad \qquad \qquad &\leq \sum_{z \in \partial D} G_{\nu, \bar{D}}(x, z) G_\nu(z, y),
\end{aligned}$$

as claimed. \square

PROBLEM 7.4. Let $m = \lfloor |y|_\infty / (R+1) \rfloor$. By the Simon-Lieb inequality (7.53), translation invariance, and the bound $G_{\nu, \bar{D}}(x, z) \leq G_\nu(x, z)$, we have

$$\begin{aligned}
G_\nu(x, y) &\leq \sum_{z_1 \in x + \partial \Lambda_R} G_\nu(x, z_1) G_\nu(z_1, y) \\
&\leq \cdots \leq \sum_{z_1 \in z_0 + \partial \Lambda_R} \cdots \sum_{z_m \in z_{m-1} + \partial \Lambda_R} G_\nu(x, z_1) G_\nu(z_1, z_2) \cdots G_\nu(z_m, y) \\
\text{(A.59)} \qquad &\leq \theta^m \sup_{x \in \mathbb{Z}^d} G_\nu(0, x).
\end{aligned}$$

Note that we applied (7.53) in such a manner that the term $G_{\nu, D}(x, y)$ vanishes. \square

PROBLEM 7.5. Fix $\nu > \nu_c$, and let $D_R = \{-R+2, \dots, R-1\}^d$ be the interior of Λ_R . By monotone convergence,

$$\text{(A.60)} \qquad \qquad \qquad G_\nu(x, y) = \lim_{R \rightarrow \infty} G_{\nu, D_R}(x, y).$$

Hence, to prove (7.55), it suffices to show that $\lim_{R \rightarrow \infty} G_\nu^R(x, y) - G_{\nu, D_R}(x, y) = 0$. Now,

$$\text{(A.61)} \qquad \qquad \qquad G_\nu^R(x, y) - G_{\nu, D_R}(x, y) = G_\nu^R(x, y) - G_{\nu, D_R}^R(x, y),$$

and thus, from the Simon-Lieb inequality (Problem 7.3), it follows that

$$\begin{aligned}
\text{(A.62)} \qquad \qquad \qquad G_\nu^R(x, y) - G_{\nu, D_R}(x, y) &\leq \sum_{z \in \partial D_R} G_{\nu, \bar{D}_R}^R(x, z) G_\nu^R(z, y) \\
&\leq \left(\sup_{z \in \partial D_R} G_{\nu, \bar{D}_R}^R(x, z) \right) \left(\sum_{z \in \partial D_R} G_\nu^R(z, y) \right).
\end{aligned}$$

By Problem 7.2,

$$\text{(A.63)} \qquad \qquad \qquad \sum_{z \in \partial D_R} G_\nu^R(z, y) \leq \sum_{z \in \Lambda_R} G_\nu^R(z, y) = \chi^R(\nu) \leq \chi(\nu) < \infty,$$

and, by Problem 7.4 and the fact that $G_\nu(0, x)$ is uniformly bounded since the susceptibility is finite,

$$\begin{aligned}
\text{(A.64)} \qquad \qquad \qquad \sup_{z \in \partial D_R} G_{\nu, \bar{D}_R}^R(x, z) &= \sup_{z \in \partial D_R} G_{\nu, \bar{D}_R}(x, z) \leq \sup_{z \in \partial D_R} G_\nu(x, z) \\
&\leq \sup_{z \in \partial D_R} C e^{-\gamma|z-x|} \leq C e^{-\gamma(R-|x|)} \rightarrow 0
\end{aligned}$$

as $R \rightarrow \infty$. Therefore $\lim_{R \rightarrow \infty} G_\nu^R(x, y) - G_{\nu, \bar{D}_R}(x, y) = 0$, proving (7.55). Finally (7.56) follows since $G_{\nu_c}(x, y) = \lim_{\nu \searrow \nu_c} G_\nu(x, y)$ by monotone convergence. \square

Acknowledgements

The authors are grateful to the organisers of both the Clay Mathematics Institute Summer School and the XIV Brazilian Probability School for having delivered such a stimulating and energetic event. We are particularly grateful to Vlas Sidoravicius for his tireless efforts on behalf of the School and its participants.

Hugo Duminil-Copin thanks Stanislav Smirnov for very stimulating discussions and encouragements related to their joint work presented in Section 3. Gordon Slade thanks David Brydges for many discussions related to their joint work presented in Section 7.

The work of Hugo Duminil-Copin was supported by the ANR grant BLAN06-3-134462, the EU Marie-Curie RTN CODY, the ERC AG CONFRA, and the Swiss FNS. The work of Gordon Slade was supported in part by NSERC of Canada.

References

- [1] V. Beffara, *SLE and other conformally invariant objects*, Lectures in this Summer School.
- [2] C. Borgs, J.T. Chayes, R. van der Hofstad, G. Slade, and J. Spencer, *Random subgraphs of finite graphs: II. The lace expansion and the triangle condition*, Ann. Probab. **33** (2005), 1886–1944.
- [3] A. Bovier, G. Felder, and J. Fröhlich, *On the critical properties of the Edwards and the self-avoiding walk model of polymer chains*, Nucl. Phys. B **230** [FS10] (1984), 119–147.
- [4] D.C. Brydges, *Lectures on the renormalisation group*, Statistical Mechanics (S. Sheffield and T. Spencer, eds.), American Mathematical Society, Providence, (2009), IAS/Park City Mathematics Series, Volume 16, pp. 7–93.
- [5] D. Brydges, S.N. Evans, and J.Z. Imbrie, *Self-avoiding walk on a hierarchical lattice in four dimensions*, Ann. Probab. **20** (1992), 82–124.
- [6] D.C. Brydges, J. Fröhlich, and A.D. Sokal, *The random walk representation of classical spin systems and correlation inequalities. II. The skeleton inequalities*, Commun. Math. Phys. **91** (1983), 117–139.
- [7] D.C. Brydges, J. Fröhlich, and T. Spencer, *The random walk representation of classical spin systems and correlation inequalities*, Commun. Math. Phys. **83** (1982), 123–150.
- [8] D.C. Brydges, G. Guadagni, and P.K. Mitter, *Finite range decomposition of Gaussian processes*, J. Stat. Phys. **115** (2004), 415–449.
- [9] D.C. Brydges and J.Z. Imbrie, *End-to-end distance from the Green’s function for a hierarchical self-avoiding walk in four dimensions*, Commun. Math. Phys. **239** (2003), 523–547.
- [10] ———, *Green’s function for a hierarchical self-avoiding walk in four dimensions*, Commun. Math. Phys. **239** (2003), 549–584.
- [11] D.C. Brydges, J.Z. Imbrie, and G. Slade, *Functional integral representations for self-avoiding walk*, Probab. Surveys **6** (2009), 34–61.
- [12] D. Brydges and G. Slade, *Renormalisation group analysis of weakly self-avoiding walk in dimensions four and higher*, To appear in *Proceedings of the International Congress of Mathematicians*, Hyderabad, 2010, ed. R. Bhatia, Hindustan Book Agency, Delhi.
- [13] ———, *Weakly self-avoiding walk in dimensions four and higher: a renormalisation group approach*, In preparation.
- [14] D.C. Brydges and T. Spencer, *Self-avoiding walk in 5 or more dimensions*, Commun. Math. Phys. **97** (1985), 125–148.
- [15] M. Campanino, D. Ioffe, and Y. Velenik, *Random path representation and sharp correlations asymptotics at high-temperatures*, Adv. Stud. Pure Math. **39** (2004), 29–52.
- [16] S. Caracciolo, M.S. Causo, and A. Pelissetto, *High-precision determination of the critical exponent γ for self-avoiding walks*, Phys. Rev. E **57** (1998), 1215–1218.
- [17] J.T. Chayes and L. Chayes, *Ornstein-Zernike behavior for self-avoiding walks at all noncritical temperatures*, Commun. Math. Phys. **105** (1986), 221–238.
- [18] D. Chelkak and S. Smirnov, *Universality in the 2D Ising model and conformal invariance of fermionic observables*, arXiv:0910.2045, (2009).

- [19] L.-C. Chen and A. Sakai, *Asymptotic behavior of the gyration radius for long-range self-avoiding walk and long-range oriented percolation*, to appear in *Ann. Probab.* Preprint, (2010).
- [20] N. Clisby, *Accurate estimate of the critical exponent ν for self-avoiding walks via a fast implementation of the pivot algorithm*, *Phys. Rev. Lett.* **104** (2010), 055702.
- [21] N. Clisby, R. Liang, and G. Slade, *Self-avoiding walk enumeration via the lace expansion*, *J. Phys. A: Math. Theor.* **40** (2007), 10973–11017.
- [22] N. Clisby and G. Slade, *Polygons and the lace expansion*, *Polygons, Polyominoes and Polycubes* (A.J. Guttmann, ed.), Springer, Dordrecht, (2009), Lecture Notes in Physics Vol. 775, pp. 117–142.
- [23] A.R. Conway, I.G. Enting, and A.J. Guttmann, *Algebraic techniques for enumerating self-avoiding walks on the square lattice*, *J. Phys. A: Math. Gen.* **26** (1993), 1519–1534.
- [24] H. Duminil-Copin and S. Smirnov, *The connective constant of the hexagonal lattice equals $\sqrt{2} + \sqrt{2}$* , arXiv:1007.0575, (2010).
- [25] P.J. Flory, *The configuration of a real polymer chain*, *J. Chem. Phys.* **17** (1949), 303–310.
- [26] B.T. Graham, *Borel-type bounds for the self-avoiding walk connective constant*, *J. Phys. A: Math. Theor.* **43** (2010), 235001.
- [27] A. Greven and F. den Hollander, *A variational characterization of the speed of a one-dimensional self-repellent random walk*, *Ann. Appl. Probab.* **3** (1993), 1067–1099.
- [28] R. Guida and J. Zinn-Justin, *Critical exponents of the N -vector model*, *J. Phys. A: Math. Gen.* **31** (1998), 8103–8121.
- [29] J.M. Hammersley and D.J.A. Welsh, *Further results on the rate of convergence to the connective constant of the hypercubical lattice*, *Quart. J. Math. Oxford* (**2**), **13** (1962), 108–110.
- [30] T. Hara, *Decay of correlations in nearest-neighbor self-avoiding walk, percolation, lattice trees and animals*, *Ann. Probab.* **36** (2008), 530–593.
- [31] T. Hara, R. van der Hofstad, and G. Slade, *Critical two-point functions and the lace expansion for spread-out high-dimensional percolation and related models*, *Ann. Probab.* **31** (2003), 349–408.
- [32] T. Hara and M. Ohno, *Renormalization group analysis of hierarchical weakly self-avoiding walk in four dimensions*, In preparation.
- [33] T. Hara and G. Slade, *Self-avoiding walk in five or more dimensions. I. The critical behaviour*, *Commun. Math. Phys.* **147** (1992), 101–136.
- [34] ———, *The lace expansion for self-avoiding walk in five or more dimensions.*, *Reviews in Math. Phys.* **4** (1992), 235–327.
- [35] ———, *The self-avoiding-walk and percolation critical points in high dimensions*, *Combin. Probab. Comput.* **4** (1995), 197–215.
- [36] G.H. Hardy, *Divergent series*, Oxford University Press, Oxford, (1949).
- [37] G.H. Hardy and S. Ramanujan, *Asymptotic formulae for the distribution of integers of various types*, *Proc. Lond. Math. Soc.* (**2**) **16** (1917), 112–132.
- [38] M. Heydenreich, *Long-range self-avoiding walk converges to alpha-stable processes*, To appear in *Ann. I. Henri Poincaré Probab. Statist.* Preprint, (2008).
- [39] M. Heydenreich, R. van der Hofstad, and A. Sakai, *Mean-field behavior for long- and finite range Ising model, percolation and self-avoiding walk*, *J. Stat. Phys.* **132** (2008), 1001–1049.
- [40] R. van der Hofstad and W. König, *A survey of one-dimensional random polymers*, *J. Stat. Phys.* **103** (2001), 915–944.
- [41] R. van der Hofstad and G. Slade, *A generalised inductive approach to the lace expansion*, *Probab. Theory Related Fields* **122** (2002), 389–430.
- [42] F. den Hollander, *Random polymers*, Lectures in this Summer School.
- [43] ———, *Random polymers*, Springer, Berlin, (2009), Lecture Notes in Mathematics Vol. 1974. Ecole d’Été de Probabilités de Saint-Flour XXXVII–2007.
- [44] E. J. Janse van Rensburg, *Monte Carlo methods for the self-avoiding walk*, *J. Phys. A: Math. Theor.* **42** (2009), 323001.
- [45] I. Jensen, *A parallel algorithm for the enumeration of self-avoiding polygons on the square lattice*, *J. Phys. A: Math. Gen.* **36** (2003), 5731–5745.
- [46] ———, *Enumeration of self-avoiding walks on the square lattice*, *J. Phys. A: Math. Gen.* **37** (2004), 5503–5524.
- [47] ———, *Improved lower bounds on the connective constants for two-dimensional self-avoiding walks*, *J. Phys. A: Math. Gen.* **37** (2004), 11521–11529.

- [48] I. Jensen and A.J. Guttmann, *Self-avoiding polygons on the square lattice*, J. Phys. A: Math. Gen. **32** (1999), 4867–4876.
- [49] T. Kennedy, *Conformal invariance and stochastic Loewner evolution predictions for the 2D self-avoiding walk—Monte Carlo tests*, J. Stat. Phys. **114** (2004), 51–78.
- [50] H. Kesten, *On the number of self-avoiding walks*, J. Math. Phys. **4** (1963), 960–969.
- [51] ———, *On the number of self-avoiding walks. II*, J. Math. Phys. **5** (1964), 1128–1137.
- [52] W. König, *A central limit theorem for a one-dimensional polymer measure*, Ann. Probab **24** (1996), 1012–1035.
- [53] G.F. Lawler, *Intersections of random walks*, Birkhäuser, Boston, (1991).
- [54] G.F. Lawler, O. Schramm, and W. Werner, *On the scaling limit of planar self-avoiding walk*, Proc. Symposia Pure Math. **72** (2004), 339–364.
- [55] B. Li, N. Madras, and A.D. Sokal, *Critical exponents, hyperscaling, and universal amplitude ratios for two- and three-dimensional self-avoiding walks*, J. Stat. Phys. **80** (1995), 661–754.
- [56] E.H. Lieb, *A refinement of Simon’s correlation inequality*, Commun. Math. Phys. **77** (1980), 127–136.
- [57] N. Madras and G. Slade, *The self-avoiding walk*, Birkhäuser, Boston, (1993).
- [58] N. Madras and A.D. Sokal, *The pivot algorithm: A highly efficient Monte Carlo method for the self-avoiding walk*, J. Stat. Phys. **50** (1988), 109–186.
- [59] A.J. McKane, *Reformulation of $n \rightarrow 0$ models using anticommuting scalar fields*, Phys. Lett. A **76** (1980), 22–24.
- [60] P.K. Mitter and B. Scoppola, *The global renormalization group trajectory in a critical supersymmetric field theory on the lattice \mathbb{Z}^3* , J. Stat. Phys. **133** (2008), 921–1011.
- [61] B. Nienhuis, *Exact critical exponents of the $O(n)$ models in two dimensions*, Phys. Rev. Lett. **49** (1982), 1062–1065.
- [62] G.L. O’Brien, *Monotonicity of the number of self-avoiding walks*, J. Stat. Phys. **59** (1990), 969–979.
- [63] G. Parisi and N. Sourlas, *Self-avoiding walk and supersymmetry*, J. Phys. Lett. **41** (1980), L403–L406.
- [64] A. Pönitz and P. Tittmann, *Improved upper bounds for self-avoiding walks in \mathbb{Z}^d* , Electron. J. Combin. **7** (2000), Paper R21.
- [65] V. Riva and J. Cardy, *Holomorphic parafermions in the Potts model and stochastic Loewner evolution*, J. Stat. Mech.: Theory Exp. (2006), P12001.
- [66] W. Rudin, *Principles of mathematical analysis*, 3rd ed., McGraw–Hill, New York, (1976).
- [67] B. Simon, *Correlation inequalities and the decay of correlations in ferromagnets*, Commun. Math. Phys. **77** (1980), 111–126.
- [68] G. Slade, *The lace expansion and the upper critical dimension for percolation*, Lectures in Applied Mathematics **27** (1991), 53–63, (Mathematics of Random Media, eds. W.E. Kohler and B.S. White, A.M.S., Providence).
- [69] ———, *The lace expansion and its applications.*, Springer, Berlin, (2006), Lecture Notes in Mathematics Vol. 1879. Ecole d’Eté de Probabilités de Saint–Flour XXXIV–2004.
- [70] S. Smirnov, *Towards conformal invariance of 2D lattice models*, International Congress of Mathematicians (Zürich), vol. II, Eur. Math. Soc., (2006), pp. 1421–1451.
- [71] K. Symanzik, *Euclidean quantum field theory*, Local Quantum Field Theory (New York) (R. Jost, ed.), Academic Press, (1969).
- [72] K. Wilson and J. Kogut, *The renormalization group and the ϵ expansion*, Phys. Rep. **12** (1974), 75–200.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, 1984 MATHEMATICS
ROAD, VANCOUVER, BC, CANADA V6T 1Z2

E-mail address: `brt@math.ubc.ca`

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE GENÈVE, 2-4 RUE DU LIÈVRE, 1211
GENÈVE, SWITZERLAND

E-mail address: `hugo.duminil@unige.ch`

EURANDOM, TECHNISCHE UNIVERSITEIT EINDHOVEN, PO Box 513, 5600 MB EINDHOVEN,
THE NETHERLANDS

E-mail address: `j.a.goodman@tue.nl`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, 1984 MATHEMATICS
ROAD, VANCOUVER, BC, CANADA V6T 1Z2

E-mail address: `slade@math.ubc.ca`