



This is a repository copy of *Left localizations of left Artinian rings*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/124659/>

Version: Submitted Version

Article:

Bavula, V.V. (2016) Left localizations of left Artinian rings. *Journal of Algebra and Its Applications*, 15 (09). 1650165. ISSN 0219-4988

<https://doi.org/10.1142/S0219498816501656>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Left localizations of left Artinian rings

V. V. Bavula

Abstract

For an arbitrary left Artinian ring R , explicit descriptions are given of all the left denominator sets S of R and left localizations $S^{-1}R$ of R . It is proved that, up to R -isomorphism, there are only finitely many left localizations and each of them is an idempotent localization, i.e. $S^{-1}R \simeq S_e^{-1}R$ and $\text{ass}(S) = \text{ass}(S_e)$ where $S_e = \{1, e\}$ is a left denominator set of R and e is an idempotent. Moreover, the idempotent e is unique up to a conjugation. It is proved that the number of maximal left denominator sets of R is finite and does not exceed the number of isomorphism classes of simple left R -modules. The set of maximal left denominator sets of R and the left localization radical of R are described.

Key Words: Goldie's Theorem, the left quotient ring of a ring, the largest left quotient ring of a ring, a maximal left denominator set, the left localization radical of a ring, a maximal left localization of a ring, a left localization maximal ring, a left Artinian ring.

Mathematics subject classification 2010: 16P50, 16P60, 16P20, 16U20.

Contents

1. Introduction.
2. Preliminaries.
3. Idempotent left denominator sets.
4. Left localizations of left Artinian rings.
5. Structure of left Artinian rings with zero left localization radical.
6. Characterization of the left localization radical of a left Artinian ring.
7. Description of left denominator sets of a left Artinian ring.
8. Localizations of Artinian rings.
9. Rings with left Artinian left quotient ring.

1 Introduction

In this paper, module means a left module, and the following notation is fixed:

- R is a ring with 1, R^* is its group of units and $\text{Inn}(R) := \{\omega_u \mid u \in R^*\}$ is the group of inner automorphisms of R where $\omega_u(r) := uru^{-1}$ for $r \in R$, $\text{rad}(R)$ is the Jacobson radical of R ;
- $\text{Ore}_l(R) := \{S \mid S \text{ is a left Ore set in } R\}$;
- $\text{Den}_l(R) := \{S \mid S \text{ is a left denominator set in } R\}$;
- $\text{Loc}_l(R) := \{[S^{-1}R] \mid S \in \text{Den}_l(R)\}$ is the set of R -isomorphism classes $[S^{-1}R]$ of left localizations of the ring R ; $[S^{-1}R] = [S'^{-1}R]$ iff the map $S^{-1}R \rightarrow S'^{-1}R$, $s^{-1}r \mapsto s'^{-1}r$ is a well-defined isomorphism. We identify $S^{-1}R$ with $[S^{-1}R]$. So $S^{-1}R = S'^{-1}R$ iff the rings $S^{-1}R$ and $S'^{-1}R$ are R -isomorphic;
- $\text{Ass}_l(R) := \{\text{ass}(S) \mid S \in \text{Den}_l(R)\}$ where $\text{ass}(S) := \{r \in R \mid sr = 0 \text{ for some } s = s(r) \in S\}$;
- $\text{Den}_l(R, \mathfrak{a}) := \{S \in \text{Den}_l(R) \mid \text{ass}(S) = \mathfrak{a}\}$.

In brief, for an arbitrary left Artinian R , this paper presents a complete picture of how left localizations and left denominator sets of R look like, and the situation is so ‘simple’ and natural that one cannot image a better/simpler one, see below.

Every left localization of left Artinian ring is an idempotent left localization. We say that a left localization $A = S^{-1}R$ of a ring R is an *idempotent left localization* of R if there is an idempotent $e \in R$ such that $S_e = \{1, e\} \in \text{Den}_l(R)$ and the rings $S_e^{-1}R$ and A are R -isomorphic (equivalently, the map $A = S^{-1}R \rightarrow S_e^{-1}R$, $s^{-1}r \mapsto s^{-1}r$, is an isomorphism).

The following theorem shows that every left localization of a left Artinian ring is an idempotent left localization.

- **(Theorem 4.5)** *Let R be a left Artinian ring and $S \in \text{Den}_l(R, \mathfrak{a})$. Then there exists a nonzero idempotent $e \in R$ such that $S_e := \{1, e\} \in \text{Den}_l(R, \mathfrak{a})$ and the rings $S^{-1}R$ and $S_e^{-1}R$ are R -isomorphic. The idempotent e is unique up to conjugation.*

There are only finitely many left localizations for a left Artinian ring. Let R be a left Artinian ring, $\text{rad}(R)$ be its radical, $\overline{R} := R/\text{rad}(R) = \prod_{i=1}^s \overline{R}_i$ – a direct product of simple Artinian rings \overline{R}_i , $\overline{1}_i$ be the identity element of the ring \overline{R}_i . So, $1 = \sum_{i=1}^s \overline{1}_i$ is the sum of orthogonal central idempotents of \overline{R} , $1 = \sum_{i=1}^s 1_i$ is a sum of orthogonal idempotents of R such that 1_i is a lifting of $\overline{1}_i$ (see (13)). For each non-empty set I of $\{1, \dots, s\}$, let $e_I := \sum_{i \in I} 1_i$,

$$\mathcal{I}'_I := \mathcal{I}'_I(R) := \{e_I \mid e_I R(1 - e_I) = 0\} \text{ and } |\mathcal{I}'_I| < \infty.$$

The following theorem shows that for a left Artinian ring R there are only finitely many left localizations. Moreover, it gives explicit descriptions of the sets $\text{Loc}_l(R)$ and $\text{Ass}_l(R)$.

- **(Theorem 4.6)** *Let R be a left Artinian ring. Then the map $\mathcal{I}'_I(R) \rightarrow \text{Loc}_l(R)$, $e \mapsto S_e^{-1}R = R/(1 - e)R$, is a bijection. The map $\mathcal{I}'_I(R) \rightarrow \text{Ass}_l(R)$, $e \mapsto (1 - e)R$, is a bijection.*

Classification of denominator sets of a left Artinian ring. A subset S of a ring R is called a *multiplicative set* if $1 \in S$, $SS \subseteq S$ and $0 \notin S$. The next theorem gives a criterion for a multiplicative set of a left Artinian ring to be a left denominator set and provides an explicit description/classification of all the left denominator sets of R .

- **(Theorem 7.6)** *Let R be a left Artinian ring and S be a multiplicative set of R . The following statements are equivalent.*

1. $S \in \text{Den}_l(R)$.
2. *There is a nonzero idempotent $e \in R$ such that $eR(1 - e) = 0$, $S \subseteq \begin{pmatrix} R_{11}^* & 0 \\ R_{21} & R_{22} \end{pmatrix}$ and there is an element $s \in S$ such that $s = \begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix}$ where $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ is the matrix ring associated with the idempotent e (see (6)).*
3. *There is a unit $\lambda \in R^*$, and idempotent $e \in \mathcal{I}'_I(R)$ and an element $s \in S$ such that $\lambda S \lambda^{-1} \subseteq \begin{pmatrix} R_{11}^* & 0 \\ R_{21} & R_{22} \end{pmatrix}$ and $\lambda s \lambda^{-1} = \begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix}$ where $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ is the matrix ring associated with the idempotent e .*
4. *There is an element $s \in S$ such that $S_s \in \text{Den}_l(R)$ where $S_s = \{s^i \mid i \in \mathbb{N}\}$ and the images of all the elements of S in the ring $S_s^{-1}R$ are units.*

If one of the equivalent conditions holds then $\text{ass}(S) = \ker_R(s \cdot)$ in all three cases regardless of the choice of s .

So, in order to obtain all the left denominator sets of a left Artinian ring R we have to choose an element $e \in \mathcal{I}'_I(R)$, a multiplicative set S of $\begin{pmatrix} R_{11}^* & 0 \\ R_{21} & R_{22} \end{pmatrix}$ that contains an element of the form

$\begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix}$. Then S is a left denominator set of R and an arbitrary left denominator set of R is of the type $\lambda S \lambda^{-1}$ for some $\lambda \in R^*$.

The maximal left denominator sets of a left Artinian ring R . In [2], the concept of maximal left denominator set of ring was introduced and it was shown that the set $\max.\text{Den}_l(R)$ of maximal left denominator sets of a ring R is a non-empty set. For a left Artinian ring R , the finite set (\mathcal{I}'_l, \geq) is a partially ordered set where $e_I \geq e_J$ iff $I \supseteq J$. Let $\min \mathcal{I}'_l$ be the set of minimal elements of \mathcal{I}'_l . The next theorem provides a description of the maximal left denominator sets of a left Artinian ring.

• **(Theorem 4.10)** *Let R be a left Artinian ring. Then*

1. $\max.\text{Den}_l(R) = \{T_e \mid e \in \min \mathcal{I}'_l(R)\}$ where $T_e = \{u \in R \mid u + (1-e)R \in (R/(1-e)R)^*\}$.
2. $|\max.\text{Den}_l(R)| \leq s$ where s is the number of isomorphism classes of left simple R -modules.
3. $|\max.\text{Den}_l(R)| = s$ iff R is a semisimple ring.

The maximal left denominator sets of a ring with left Artinian left quotient ring. The next theorem shows that a ring with left Artinian left quotient ring has only finitely many maximal left denominator sets.

• **(Theorem 9.1)** *Let R be a ring such that $Q_l(R)$ is a left Artinian ring and s be the number of iso-classes of simple left $Q_l(R)$ -modules. Then*

1. the map $\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q_l(R))$, $S \mapsto SQ_l(R)^*$, is a bijection with the inverse $T \mapsto T \cap R$. In particular, $|\max.\text{Den}_l(R)| = |\max.\text{Den}_l(Q_l(R))| \leq s < \infty$.
2. $|\max.\text{Den}_l(R)| = s$ iff $Q_l(R)$ is a semisimple ring iff R is a semiprime left Goldie ring.

Recall that the largest left quotient ring $Q_l(R)$ of R is a left Artinian ring iff the (classical) left quotient ring $Q_{l,cl}(R) := \mathcal{C}_R^{-1}R$ is a left Artinian ring, and in this case $S_l(R) = \mathcal{C}_R$, [3].

Criterion for the powers of an element to be a left denominator set. For a left Artinian ring R , the following theorem is an explicit criterion for the powers of a non-nilpotent element of R to be a left denominator set.

• **(Theorem 7.4)** *Let R be a left Artinian ring, $\mathcal{I}'_l(R)$ be as above, $s \in R$ be a non-nilpotent element of R , $e = e(s)$ be the idempotent associated with the element s (see (9)), $S_e = \{1, e\}$ and $S_s = \{s^i \mid i \in \mathbb{N}\}$. The following statements are equivalent.*

1. $S_s \in \text{Den}_l(R)$.
2. $S_e \in \text{Den}_l(R)$ and $(1-e)s(1-e)$ is a nilpotent element.
3. $eR(1-e) = 0$ and $(1-e)s(1-e)$ is a nilpotent element.

If one of the equivalent conditions holds then $\text{ass}(S_s) = (1-e)R$ and $S_s^{-1}R \simeq S_e^{-1}R \simeq R/(1-e)R$, the core $S_{s,c}$ of the left denominator set S_s is equal to $\{s^i \mid i \geq 1, (1-e)s^i(1-e) = 0\}$.

Duality between left and right localizations of an Artinian ring. The sets of left and right localizations of a ring R , $(\text{Loc}_l(R), \rightarrow)$ and $(\text{Loc}_r(R), \rightarrow)$, are partially ordered sets where $[A] \rightarrow [B]$ if there is a ring R -homomorphism $A \rightarrow B$. In general, the left and right localizations of a ring R are almost unrelated but for each Artinian ring R there is a duality between the partially ordered sets $(\text{Loc}_l(R), \rightarrow)$ and $(\text{Loc}_r(R), \rightarrow)$.

• **(Theorem 4.12)** *Let R be an Artinian ring. Then the map*

$$\text{Loc}_l(R) \setminus [R] \rightarrow \text{Loc}_r(R) \setminus [R], \quad [R/(1-e_I)R] \rightarrow [R/R(1-e_{CI})],$$

is an anti-isomorphism of posets (i.e. an order reversing bijection). In particular, $|\text{Loc}_l(R)| = |\text{Loc}_r(R)|$.

2 Preliminaries

In this section, we collect necessary results that are used in the proofs of this paper. More results on localizations of rings (and some of the missed standard definitions) the reader can find in [7], [9] and [8]. In this paper the following notation will remained fixed.

Notation:

- $S_{\mathfrak{a}} = S_{\mathfrak{a}}(R) = S_{l,\mathfrak{a}}(R)$ is the *largest element* of the poset $(\text{Den}_l(R, \mathfrak{a}), \subseteq)$ and $Q_{\mathfrak{a}}(R) := Q_{l,\mathfrak{a}}(R) := S_{\mathfrak{a}}^{-1}R$ is the *largest left quotient ring associated to \mathfrak{a}* , $S_{\mathfrak{a}}$ exists (Theorem 2.1, [2]);
- in particular, $S_0 = S_0(R) = S_{l,0}(R)$ is the largest element of the poset $(\text{Den}_l(R, 0), \subseteq)$ and $Q_l(R) := S_0^{-1}R$ is the *largest left quotient ring of R* , [2];
- $\text{Loc}_l(R, \mathfrak{a}) := \{S^{-1}R \in \text{Loc}_l(R) \mid S \in \text{Den}_l(R, \mathfrak{a})\}$.

In [2], we introduce the following new concepts and prove their existence for an *arbitrary* ring: *the largest left quotient ring of a ring, the largest regular left Ore set of a ring, the left localization radical of a ring, a maximal left denominator set, a maximal left quotient ring of a ring, a (left) localization maximal ring*. Using an analogy with rings, the counter parts of the three concepts: a maximal left denominator set, the left localization radical and a maximal left quotient ring, for rings would be a left maximal ideal, the Jacobson radical and a simple factor ring, respectively.

The largest regular left Ore set and the largest left quotient ring of a ring. Let R be a ring. A *multiplicatively closed subset* S of R or a *multiplicative subset* of R (i.e. a multiplicative sub-semigroup of (R, \cdot) such that $1 \in S$ and $0 \notin S$) is said to be a *left Ore set* if it satisfies the *left Ore condition*: for each $r \in R$ and $s \in S$, $Sr \cap Rs \neq \emptyset$. Let $\text{Ore}_l(R)$ be the set of all left Ore sets of R . For $S \in \text{Ore}_l(R)$, $\text{ass}(S) := \{r \in R \mid sr = 0 \text{ for some } s \in S\}$ is an ideal of the ring R .

A left Ore set S is called a *left denominator set* of the ring R if $rs = 0$ for some elements $r \in R$ and $s \in S$ implies $tr = 0$ for some element $t \in S$, i.e. $r \in \text{ass}(S)$. Let $\text{Den}_l(R)$ be the set of all left denominator sets of R . For $S \in \text{Den}_l(R)$, let $S^{-1}R = \{s^{-1}r \mid s \in S, r \in R\}$ be the *left localization* of the ring R at S (the *left quotient ring of R at S*). Let us stress that in Ore's method of localization one can localize *precisely* at left denominator sets.

In general, the set \mathcal{C} of regular elements of a ring R is neither left nor right Ore set of the ring R and as a result neither left nor right classical quotient ring ($Q_{l,cl}(R) := \mathcal{C}^{-1}R$ and $Q_{r,cl}(R) := R\mathcal{C}^{-1}$) exists. Remarkably, there exists the largest regular left Ore set $S_0 = S_{l,0} = S_{l,0}(R)$, [2]. This means that the set $S_{l,0}(R)$ is an Ore set of the ring R that consists of regular elements (i.e., $S_{l,0}(R) \subseteq \mathcal{C}$) and contains all the left Ore sets in R that consist of regular elements. Also, there exists the largest regular (left and right) Ore set $S_{l,r,0}(R)$ of the ring R . In general, all the sets \mathcal{C} , $S_{l,0}(R)$, $S_{r,0}(R)$ and $S_{l,r,0}(R)$ are distinct, for example, when $R = \mathbb{I}_1 = K\langle x, \partial, \int \rangle$ is the ring of polynomial integro-differential operators over a field K of characteristic zero, [1]. In [1], these four sets are found for $R = \mathbb{I}_1$.

Definition, [1], [2]. The ring

$$Q_l(R) := S_{l,0}(R)^{-1}R$$

(respectively, $Q_r(R) := RS_{r,0}(R)^{-1}$ and $Q(R) := S_{l,r,0}(R)^{-1}R \simeq RS_{l,r,0}(R)^{-1}$) is called the *largest left* (respectively, *right and two-sided*) *quotient ring* of the ring R .

In general, the rings $Q_l(R)$, $Q_r(R)$ and $Q(R)$ are not isomorphic, for example, when $R = \mathbb{I}_1$, [1]. The next theorem gives various properties of the ring $Q_l(R)$. In particular, it describes its group of units.

Theorem 2.1 [2]

1. $S_0(Q_l(R)) = Q_l(R)^*$ and $S_0(Q_l(R)) \cap R = S_0(R)$.

2. $Q_l(R)^* = \langle S_0(R), S_0(R)^{-1} \rangle$, i.e. the group of units of the ring $Q_l(R)$ is generated by the sets $S_0(R)$ and $S_0(R)^{-1} := \{s^{-1} \mid s \in S_0(R)\}$.
3. $Q_l(R)^* = \{s^{-1}t \mid s, t \in S_0(R)\}$.
4. $Q_l(Q_l(R)) = Q_l(R)$.

The maximal left denominator sets and the maximal left localizations of a ring. The set $(\text{Den}_l(R), \subseteq)$ is a poset (partially ordered set). In [2], it is proved that the set $\text{max.Den}_l(R)$ of its maximal elements is a *non-empty* set.

Definition, [2]. An element S of the set $\text{max.Den}_l(R)$ is called a *maximal left denominator set* of the ring R and the ring $S^{-1}R$ is called a *maximal left quotient ring* of the ring R or a *maximal left localization ring* of the ring R . The intersection

$$\mathfrak{l}_R := \text{l.rad}(R) := \bigcap_{S \in \text{max.Den}_l(R)} \text{ass}(S) \quad (1)$$

is called the *left localization radical* of the ring R , [2].

For a ring R , there is the canonical exact sequence

$$0 \rightarrow \mathfrak{l}_R \rightarrow R \xrightarrow{\sigma} \prod_{S \in \text{max.Den}_l(R)} S^{-1}R, \quad \sigma := \prod_{S \in \text{max.Den}_l(R)} \sigma_S, \quad (2)$$

where $\sigma_S : R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$. For a ring R with a semisimple left quotient ring, the left localization radical \mathfrak{l}_R coincides with the prime radical \mathfrak{n}_R of R , [2]. In general, $\mathfrak{l}_R \neq \mathfrak{n}_R$ and $\mathfrak{l}_R \neq \text{rad}(R)$, Theorem 4.13.(4).

The maximal elements of $\text{Ass}_l(R)$. Let $\text{max.Ass}_l(R)$ be the set of maximal elements of the poset $(\text{Ass}_l(R), \subseteq)$ and

$$\text{ass.max.Den}_l(R) := \{\text{ass}(S) \mid S \in \text{max.Den}_l(R)\}. \quad (3)$$

These two sets are equal (Proposition 2.4), a proof is based on Lemma 2.2 and Corollary 2.3. For an non-empty set X or R , let $\text{r.ass}(X) := \{r \in R \mid rx = 0 \text{ for some } x = x(r) \in X\}$.

Lemma 2.2 [2] *Let $S \in \text{Den}_l(R, \mathfrak{a})$ and $T \in \text{Den}_l(R, \mathfrak{b})$ be such that $\mathfrak{a} \subseteq \mathfrak{b}$. Let ST be the multiplicative semigroup generated by S and T in (R, \cdot) . Then*

1. $\text{r.ass}(ST) \subseteq \mathfrak{b}$.
2. $ST \in \text{Den}_l(R, \mathfrak{c})$ and $\mathfrak{b} \subseteq \mathfrak{c}$.

Corollary 2.3 *Let R be a ring, $S \in \text{max.Den}_l(R)$ and $T \in \text{Den}_l(R)$. Then $T \subseteq S$ iff $\text{ass}(T) \subseteq \text{ass}(S)$.*

Proof. (\Rightarrow) If $T \subseteq S$ then $\text{ass}(T) \subseteq \text{ass}(S)$.

(\Leftarrow) If $\text{ass}(T) \subseteq \text{ass}(S)$. then, by Lemma 2.2, $ST \in \text{Den}_l(R)$ and $S \subseteq ST$, hence $S = ST$, by the maximality of S . Then $T \subseteq S$. \square

Proposition 2.4 [2] $\text{max.Ass}_l(R) = \text{ass.max.Den}_l(R) \neq \emptyset$. *In particular, the ideals of this set are incomparable (i.e. neither $\mathfrak{a} \subseteq \mathfrak{b}$ nor $\mathfrak{a} \supseteq \mathfrak{b}$).*

Properties of the maximal left quotient rings of a ring. The next theorem describes various properties of the maximal left quotient rings of a ring, in particular, their groups of units and their largest left quotient rings.

Theorem 2.5 [2] Let $S \in \max.\text{Den}_l(R)$, $A = S^{-1}R$, A^* be the group of units of the ring A ; $\mathfrak{a} := \text{ass}(S)$, $\pi_{\mathfrak{a}} : R \rightarrow R/\mathfrak{a}$, $a \mapsto a + \mathfrak{a}$, and $\sigma_{\mathfrak{a}} : R \rightarrow A$, $r \mapsto \frac{r}{1}$. Then

1. $S = S_{\mathfrak{a}}(R)$, $S = \pi_{\mathfrak{a}}^{-1}(S_0(R/\mathfrak{a}))$, $\pi_{\mathfrak{a}}(S) = S_0(R/\mathfrak{a})$ and $A = S_0(R/\mathfrak{a})^{-1}R/\mathfrak{a} = Q_l(R/\mathfrak{a})$.
2. $S_0(A) = A^*$ and $S_0(A) \cap (R/\mathfrak{a}) = S_0(R/\mathfrak{a})$.
3. $S = \sigma_{\mathfrak{a}}^{-1}(A^*)$.
4. $A^* = \langle \pi_{\mathfrak{a}}(S), \pi_{\mathfrak{a}}(S)^{-1} \rangle$, i.e. the group of units of the ring A is generated by the sets $\pi_{\mathfrak{a}}(S)$ and $\pi_{\mathfrak{a}}^{-1}(S) := \{\pi_{\mathfrak{a}}(s)^{-1} \mid s \in S\}$.
5. $A^* = \{\pi_{\mathfrak{a}}(s)^{-1}\pi_{\mathfrak{a}}(t) \mid s, t \in S\}$.
6. $Q_l(A) = A$ and $\text{Ass}_l(A) = \{0\}$. In particular, if $T \in \text{Den}_l(A, 0)$ then $T \subseteq A^*$.

Let $\max.\text{Loc}_l(R)$ be the set of maximal elements of the poset $(\text{Loc}_l(R), \rightarrow)$ where $A \rightarrow B$ for $A, B \in \text{Loc}_l(R)$ means that there exist $S, T \in \text{Den}_l(R)$ such that $S \subseteq T$, $A = S^{-1}R$ and $B = T^{-1}R$ (then there exists a natural ring homomorphism $A \rightarrow B$, $s^{-1}r \mapsto s^{-1}r$). Then (see [2]),

$$\max.\text{Loc}_l(R) = \{S^{-1}R \mid S \in \max.\text{Den}_l(R)\} = \{Q_l(R/\mathfrak{a}) \mid \mathfrak{a} \in \text{ass}.\max.\text{Den}_l(R)\}. \quad (4)$$

The maximal left quotient rings of a finite direct product of rings.

Theorem 2.6 [4] Let $R = \prod_{i=1}^n R_i$ be the direct product of rings R_i . Then for each $i = 1, \dots, n$, the map

$$\max.\text{Den}_l(R_i) \rightarrow \max.\text{Den}_l(R), \quad S_i \mapsto R_1 \times \dots \times S_i \times \dots \times R_n, \quad (5)$$

is an injection. Moreover, $\max.\text{Den}_l(R) = \prod_{i=1}^n \max.\text{Den}_l(R_i)$ in the sense of (5), i.e.

$$\max.\text{Den}_l(R) = \{S_i \mid S_i \in \max.\text{Den}_l(R_i), i = 1, \dots, n\},$$

$S_i^{-1}R \simeq S_i^{-1}R_i$, $\text{ass}_R(S_i) = R_1 \times \dots \times \text{ass}_{R_i}(S_i) \times \dots \times R_n$. The core of the left denominator set S_i in R coincides with the core of the left denominator set S_i in R_i , i.e.

$$(R_1 \times \dots \times S_i \times \dots \times R_n)_c = 0 \times \dots \times S_{i,c} \times \dots \times 0.$$

Corollary 2.7 Let $R = \prod_{i=1}^n R_i$ be the direct product of rings R_i . Then $\text{t}_R = \prod_{i=1}^n \text{t}_{R_i}$.

A bijection between $\max.\text{Den}_l(R)$ and $\max.\text{Den}_l(Q_l(R))$.

Proposition 2.8 [4] Let R be a ring, S_l be the largest regular left Ore set of the ring R , $Q_l := S_l^{-1}R$ be the largest left quotient ring of the ring R , and \mathcal{C} be the set of regular elements of the ring R . Then

1. $S_l \subseteq S$ for all $S \in \max.\text{Den}_l(R)$. In particular, $\mathcal{C} \subseteq S$ for all $S \in \max.\text{Den}_l(R)$ provided \mathcal{C} is a left Ore set.
2. Either $\max.\text{Den}_l(R) = \{\mathcal{C}\}$ or, otherwise, $\mathcal{C} \notin \max.\text{Den}_l(R)$.
3. The map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q_l), \quad S \mapsto SQ_l^* = \{c^{-1}s \mid c \in S_l, s \in S\},$$

is a bijection with the inverse $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$ where $\sigma : R \rightarrow Q_l$, $r \mapsto \frac{r}{1}$, and SQ_l^* is the sub-semigroup of (Q_l, \cdot) generated by the set S and the group Q_l^* of units of the ring Q_l , and $S^{-1}R = (SQ_l^*)^{-1}Q_l$.

4. If \mathcal{C} is a left Ore set then the map

$$\max.\text{Den}_l(R) \rightarrow \max.\text{Den}_l(Q := \mathcal{C}^{-1}R), \quad S \mapsto SQ^* = \{c^{-1}s \mid c \in \mathcal{C}, s \in S\},$$

is a bijection with the inverse $\mathcal{T} \mapsto \sigma^{-1}(\mathcal{T})$ where $\sigma : R \rightarrow Q$, $r \mapsto \frac{r}{1}$, and SQ^* is the sub-semigroup of (Q, \cdot) generated by the set S and the group Q^* of units of the ring Q , and $S^{-1}R = (SQ^*)^{-1}Q$.

3 Idempotent left denominator sets

Theorem 4.5 states that every left localization of a left Artinian ring is an idempotent localization. In this section, several results on idempotent left denominator sets are given that are used in proofs of the subsequent sections.

Let R be a ring and $e \in R$ be a *nonzero* idempotent. Then $1 = e_1 + e_2$ is the sum of orthogonal idempotents where $e_1 = e$ and $e_2 = 1 - e_1$. The ring R can be seen as the *matrix ring associated with the idempotent e* ,

$$R = \bigoplus_{i,j=1}^2 R_{ij} = \begin{pmatrix} R_{11} & R_{12} \\ R_{21} & R_{22} \end{pmatrix} \quad \text{where } R_{ij} := e_i R e_j. \quad (6)$$

For an element $r \in R$, let $r \cdot : R \rightarrow R$, $x \mapsto rx$, and $\cdot r : R \rightarrow R$, $x \mapsto xr$. The next proposition is a criterion for an *idempotent multiplicative set* $S_e = \{1, e\}$ to be a left denominator set of the ring R .

Proposition 3.1 *Let e be a nonzero idempotent of a ring R . We keep the notation as above. Then $S_e = \{1, e\} \in \text{Den}_l(R)$ iff $R_{12} = 0$. In this case, $\text{ass}(S_e) = \ker(e \cdot) = (1 - e)R$ and $S_e^{-1}R \simeq R/\text{ass}(S_e) \simeq R/(1 - e)R \simeq R_{11}$.*

Proof. (\Rightarrow) If $S_e \in \text{Den}_l(R)$ then $\mathfrak{a} := \text{ass}(S_e) = \ker(e_1 \cdot) = e_2 R$. Since $R_{12}e_1 = 0$, we must have $R_{12} \subseteq \mathfrak{a} = e_2 R = R_{21} + R_{22}$, hence $R_{12} = 0$. Clearly, $S_e^{-1}R \simeq R/\mathfrak{a} \simeq R_{11}$ since $e_2 \in \mathfrak{a}$ and $1 \equiv e_1 \pmod{\mathfrak{a}}$.

(\Leftarrow) Suppose that $R_{12} = 0$, i.e. $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$. Then $S_e \in \text{Ore}_l(R)$ since for any element $r = \begin{pmatrix} r_{11} & 0 \\ r_{21} & r_{22} \end{pmatrix} \in R$, $er = \begin{pmatrix} r_{11} & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} r_{11} & 0 \\ 0 & 0 \end{pmatrix} e$. The inclusion $\ker(\cdot e) = R_{22} \subseteq \mathfrak{a}$ implies that $S_e \in \text{Den}_l(R)$. \square

Proposition 3.1 means that the idempotent multiplicative set S_e is a left denominator set of R iff the ring R is *left triangular* (as the matrix ring associated with the idempotent e), i.e.

$$R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}.$$

This fact is the most vivid demonstration of the fact that very often a left denominator set fails to be a right denominator set. Recall that $\text{Den}(R)$ is the set of (left and right) denominator sets of a ring R . The next corollary demonstrates that the condition being a *left and right* denominator set is a strong one.

Corollary 3.2 *Let e be a nonzero idempotent of a ring R . Then $S_e = \{1, e\} \in \text{Den}(R)$ iff $R_{12} = 0$ and $R_{21} = 0$ iff e is a central idempotent. In this case, $\text{ass}(S_e) = \ker(e \cdot) = (1 - e)R$ and $S_e^{-1}R \simeq R/(1 - e)R \simeq R_{11}$.*

Proof. The first ‘iff’ is due to Proposition 3.1. The second ‘iff’ is obvious. \square

Let $\text{Aut}(R)$ be the group of automorphisms of the ring R and $\text{Inn}(R) := \{\omega_u \mid u \in R^*\}$ be the group of inner automorphisms of the ring R where $\omega_u(r) := uru^{-1}$ for $r \in R$. The group $\text{Inn}(R)$ is a normal subgroup of $\text{Aut}(R)$ (since for $\sigma \in \text{Aut}(R)$ and $\omega_u \in \text{Inn}(R)$, $\sigma\omega_u\sigma^{-1} = \omega_{\sigma(u)}$).

Let $e \in R$ be an idempotent. If $S_e = \{1, e\} \in \text{Den}_l(R)$ then the set S_e is called an *idempotent left denominator set* of the ring R and the idempotent e is called a *left denominator idempotent* of R . Let $\text{IDen}_l(R)$ be the set of all the idempotent left denominator sets of the ring R and let $\mathcal{I}_l = \mathcal{I}_l(R)$ be the set of all left denominator idempotents of the ring R . By Proposition 3.1,

$$\mathcal{I}_l(R) = \{e \in R \mid e^2 = e, eR(1 - e) = 0\}. \quad (7)$$

The map

$$\mathcal{I}_l(R) \rightarrow \text{IDen}_l(R), \quad e \mapsto S_e = \{1, e\}, \quad (8)$$

is a bijection. The groups $\text{Aut}(R)$ and $\text{Inn}(R)$ act in the obvious way on the sets $\mathcal{I}_l(R)$ and $\text{IDen}_l(R)$. A ring R is called a *local ring* if the factor ring $R/\text{rad}(R)$ is a division ring.

Corollary 3.3 *Let R be a ring and $e \in R \setminus \{0, 1\}$ be an idempotent.*

1. *The following statements are equivalent.*
 - (a) *The idempotents e and $1 - e$ are left denominator idempotents.*
 - (b) *The idempotent e is a central idempotent.*
 - (c) *The idempotents e and $1 - e$ are right denominator idempotents.*
2. *The following statements are equivalent.*
 - (a) *All the idempotents of R are left denominator idempotents.*
 - (b) *All the idempotents of R are central idempotents.*
 - (c) *All the idempotents of R are right denominator idempotents.*
3. *Let R be a left Artinian ring. Then the following statements are equivalent.*
 - (a) *All the idempotents of R are left denominator idempotents.*
 - (b) *The ring R is a direct product of finitely many local left Artinian rings.*
 - (c) *All the idempotents of R are right denominator idempotents.*

Proof. 1. Statement 1 follows from Proposition 3.1.

2. Statement 2 follows from statement 1.

3. Statement 3 follows from statement 2 and the fact that 1 is the only nonzero idempotent of a local left Artinian ring (in such a ring, every nonzero idempotent e is a primitive one, hence is conjugate to 1, i.e. $e = 1$). \square

A set (X, \geq) is called a *pre-ordered set* if

- (i) $x \geq x$,
- (ii) $x \geq y$ and $y \geq z$ implies $x \geq z$.

In general, the conditions $x \geq y$ and $y \geq x$ do not imply $x = y$. If this property holds the pre-ordered set X is called a *partially ordered set*, a *poset*, for short. The set \mathcal{I}_l is a pre-ordered set (\mathcal{I}_l, \geq) where

$$e_1 \geq e_2 \text{ iff } e_2 e_1 = e_2 \text{ iff } (1 - e_2)(1 - e_1) = 1 - e_1 \text{ iff } (1 - e_1)R \subseteq (1 - e_2)R.$$

The last ‘iff’ follows from the fact that the inclusion $(1 - e_1)R \subseteq (1 - e_2)R$ implies the inclusion $e_2(1 - e_1)R \subseteq e_2(1 - e_2)R = 0$, and so $e_2 e_1 = e_2$. Via the bijection (8), the set $\text{IDen}_l(R)$ is a pre-ordered set $(\text{IDen}_l(R), \geq)$ where $S_{e_1} \geq S_{e_2}$ iff $e_1 \geq e_2$.

Definition. We say that a left localization $A = S^{-1}R$ of a ring R is an *idempotent left localization* of R if there is an idempotent $e \in R$ such that $S_e = \{1, e\} \in \text{Den}_l(R)$ and $S_e^{-1}R = A$ in $\text{Loc}_l(R)$, i.e. the map $A = S^{-1}R \rightarrow S_e^{-1}R$, $s^{-1}r \mapsto s^{-1}r$, is an isomorphism, i.e. $\text{ass}(S) = \text{ass}(S_e)$ and $s + \text{ass}(S_e) \in (R/\text{ass}(S_e))^*$ for all $s \in S$. Let $\text{ILoc}_l(R) := \{S_e^{-1}R \mid e \in \mathcal{I}_l(R)\}$, the set of all the idempotent left localizations of R .

Remark. In general, even for Artinian rings, it is not true that the condition $S^{-1}R \simeq S_e^{-1}R$ or even $S^{-1}R = S_e^{-1}R$ in $\text{Loc}_l(R)$ for some $S \in \text{Den}_l(R)$ and $S_e \in \text{IDen}_l(R)$ implies that the set S contains an idempotent element distinct from 1.

Example: Let $R = \begin{pmatrix} \mathbb{Q} & 0 \\ \mathbb{Q} & \mathbb{Q} \end{pmatrix}$, $e = E_{11}$, $\mathfrak{a} = (1-e)R$, $s = \begin{pmatrix} 2 & 0 \\ 1 & 0 \end{pmatrix}$, $S = \{1, s^i = \begin{pmatrix} 2^i & 0 \\ 2^{i-1} & 0 \end{pmatrix} \mid i \geq 1\}$. Then $S_e = \{1, e\} \in \text{Den}_l(R, \mathfrak{a})$ (by Proposition 3.1) and $S \in \text{Den}_l(R, \mathfrak{a})$ (by Corollary 4.4), $S_e^{-1}R \simeq R/\mathfrak{a} \simeq S^{-1}R$ (by Proposition 3.1 and Corollary 4.4). Clearly, 1 is the only idempotent of the set S and $S_e^{-1}R = S^{-1}R$ in $\text{Loc}_l(R)$. For all $i \geq 1$, $s^i \mathfrak{a} = 0$ and $\mathfrak{a} s^i \neq 0$.

Lemma 3.4 *Let $e_1, e_2 \in \mathcal{I}_l(R)$. The following statements are equivalent.*

1. $e_1 \geq e_2$.
2. $\frac{e_1}{1}$ is a unit in the ring $S_{e_2}^{-1}R$; equivalently, $\frac{e_1}{1} = 1 \in S_{e_2}^{-1}R$.
3. The map $S_{e_1}^{-1}R \rightarrow S_{e_2}^{-1}R$, $e_1^i r \rightarrow e_2^i r$, where $i = 0, -1$, is well-defined.
4. $\text{ass}(S_{e_1}) \subseteq \text{ass}(S_{e_2})$.

Proof. (1 \Rightarrow 2) If $e_1 \geq e_2$, i.e. $e_2 e_1 = e_2$ then $\frac{e_2}{1} \frac{e_1}{1} = \frac{e_2}{1}$ in $S_{e_2}^{-1}R$, and so $\frac{e_1}{1} = 1 = \frac{e_2}{1}$ since $\frac{e_2}{1}$ is the identity element of the ring $S_{e_2}^{-1}R$.

(2 \Rightarrow 3) If $\frac{e_1}{1}$ is a unit of the ring $S_{e_2}^{-1}R$ then the map $e_1^i r \rightarrow e_2^i r$ exists by the universal property of left localizations.

(3 \Rightarrow 1) If the map in statement 3 is well-defined then $\frac{e_1}{1} = 1 = \frac{e_2}{1}$ in $S_{e_2}^{-1}R$ and so $e_1 - e_2 \in \text{ass}(S_{e_2}) = \ker_R(e_2 \cdot)$, i.e. $e_2(e_1 - e_2) = 0$. This means that $e_1 \geq e_2$.

(1 \Rightarrow 4) The equality $e_2 e_1 = e_2$ implies that $\text{ass}(S_{e_1}) = \ker(e_1 \cdot) \subseteq \ker(e_2 \cdot) = \text{ass}(S_{e_2})$.

(4 \Rightarrow 2) Notice that $S_{e_1}^{-1}R = R/\text{ass}(S_{e_1})$, $S_{e_2}^{-1}R = R/\text{ass}(S_{e_2})$ and $e_1 \in 1 + \text{ass}(S_{e_1})$. If $\text{ass}(S_{e_1}) \subseteq \text{ass}(S_{e_2})$ then $S_{e_2}^{-1}R \ni \frac{e_1}{1} = 1 + \text{ass}(S_{e_1})$ is a unit. \square

Let us define an equivalence relation \sim on $\mathcal{I}_l(R)$ by the rule $e_1 \sim e_2$ iff $e_1 \geq e_2$ and $e_2 \geq e_1$. Let $[e] := \{f \in \mathcal{I}_l(R) \mid f \sim e\}$ be the equivalence class of $e \in \mathcal{I}_l(R)$. Then set $\mathcal{I}_l(R)/\sim := \{[e] \mid e \in \mathcal{I}_l(R)\}$ of equivalence classes is a poset $(\mathcal{I}_l(R)/\sim, \geq)$ where $[e_1] \geq [e_2]$ if $e_1 \geq e_2$.

Lemma 3.5 *Let $e_1, e_2 \in \mathcal{I}_l(R)$. The following statements are equivalent.*

1. $e_1 \sim e_2$.
2. $e_1 \equiv 1 \pmod{\text{ass}(S_{e_2})}$ and $e_2 \equiv 1 \pmod{\text{ass}(S_{e_1})}$ (i.e. $S_{e_1}^{-1}R = S_{e_2}^{-1}R$ in $\text{Loc}_l(R)$).
3. The map $S_{e_1}^{-1}R \rightarrow S_{e_2}^{-1}R$, $e_1^i r \mapsto e_2^i r$, where $i = 0, -1$, is an isomorphism.
4. $\text{ass}(S_{e_1}) = \text{ass}(S_{e_2})$.

Proof. The lemma is an easy corollary of Proposition 3.1 and Lemma 3.4. \square

Notice that every ideal is invariant under the inner automorphisms. By Lemma 3.5, the group of inner automorphisms $\text{Inn}(R)$ of the ring R acts on the set $\mathcal{I}_l(R)/\sim$ by the rule: for any $u \in R^*$ and $[e] \in \mathcal{I}_l(R)/\sim$, $u[e]u^{-1} := [ueu^{-1}]$.

Example. Let $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ be any triangular ring where R_{11} and R_{22} are arbitrary rings and R_{21} be an arbitrary (R_{22}, R_{11}) -bimodule. Let $e_1 = E_{11}$. By Proposition 3.1, $S_{e_1} = \{1, e_1\} \in \text{Den}_l(R, \mathfrak{a})$ where $\mathfrak{a} = E_{22}R$. Clearly, $R^* = \begin{pmatrix} R_{11}^* & 0 \\ R_{21} & R_{22}^* \end{pmatrix}$ and $\text{Inn}(R) \cdot e_1 = \begin{pmatrix} 1 & 0 \\ R_{21} & 0 \end{pmatrix}$ since for all units $u = \begin{pmatrix} \alpha & 0 \\ \beta & \gamma \end{pmatrix} \in R^*$,

$$ue_1u^{-1} = \begin{pmatrix} 1 & 0 \\ \beta\alpha^{-1} & 0 \end{pmatrix}.$$

Therefore, for any element $a \in R_{21}$, $e_a := \begin{pmatrix} 1 & 0 \\ a & 0 \end{pmatrix}$ is an idempotent of R , $S_{e_a} \in \text{Den}_l(R, \mathfrak{a})$ and $S_{e_a}^{-1}R \simeq R/\mathfrak{a} = S_{e_1}^{-1}R$.

4 Left localizations of left Artinian rings

Throughout this section, R is a left Artinian ring if it is not stated otherwise. The aim of this section is to prove, for a left Artinian ring R , that every left localization of R is an idempotent left localization (Theorem 4.5), there are only finitely many left localization rings of R and to give a classification of all of them (Theorem 4.6), to classify the maximal left denominator sets of R (Theorem 4.10), to give an explicit description of the left localization radical \mathfrak{l}_R of R (Theorem 4.13). The ideals in $\text{Ass}_l(R)$ have many interesting/unexpected properties (Corollary 4.11).

An element r of a ring R is called a *left regular* if the map $\cdot r : R \rightarrow R$ is an injection, i.e. $xr = 0$ implies $x = 0$. Recall that every left Artinian ring is left Noetherian.

Lemma 4.1 *Let R be a left Artinian ring.*

1. *Every left regular element of the ring R is a unit.*
2. *Let $S \in \text{Den}_l(R, \mathfrak{a})$. Then $S^{-1}R \simeq R/\mathfrak{a}$ (an R -isomorphism).*
3. *Let $s \in R$ and $\cdot s : R \rightarrow R, r \mapsto rs$. Then $R = Rs \oplus \ker(\cdot s)$ iff $Rs \cap \ker(\cdot s) = 0$.*
4. *Let $S \in \text{Den}_l(R, \mathfrak{a})$ and $T \in \text{Den}_l(R, \mathfrak{b})$. Then the rings $S^{-1}R$ and $T^{-1}R$ are R -isomorphic iff $\mathfrak{a} = \mathfrak{b}$.*

Proof. 1. Trivial.

2. Statement 2 follows from statement 1. Let $\pi : R \rightarrow R/\mathfrak{a}, r \mapsto \bar{r} = r + \mathfrak{a}$. Then $\pi(S) \in \text{Den}_l(R/\mathfrak{a}, 0)$ and $S^{-1}R \simeq \pi(S)^{-1}(R/\mathfrak{a})$. By statement 1, the set of regular elements $\pi(S)$ of the left Artinian ring R/\mathfrak{a} consists of units, and so $\pi(S)^{-1}(R/\mathfrak{a}) \simeq R/\mathfrak{a}$.

3. (\Rightarrow) Trivial.

(\Leftarrow) if $Rs \cap \ker(\cdot s) = 0$ then $Rs \oplus \ker(\cdot s) \subseteq R$. It follows from the short exact sequence of R -modules $0 \rightarrow \ker(\cdot s) \rightarrow R \xrightarrow{\cdot s} Rs \rightarrow 0$ that the left R -modules $Rs \oplus \ker(\cdot s)$ and R have the same length. Then, $Rs \oplus \ker(\cdot s) = R$.

4. Statement 4 follows from statement 2. \square

Statement 4 does not hold for non-Artinian rings, eg, $R = \mathbb{Z}, S_1 = \{1\}$ and $S_2 = \{2^i \mid i \in \mathbb{N}\}$. Statement 4 does not hold if the condition ‘ R -isomorphic’ is replaced by ‘isomorphic’, eg, if $R = K \times K, 1 = e_1 + e_2, K$ is a field then $S_{e_1}^{-1}R \simeq K \simeq S_{e_2}^{-1}R$ but $\text{ass}(S_{e_1}) = (1 - e_1)R = \{0\} \times K \neq K \times \{0\} = (1 - e_2)R = \text{ass}(S_{e_2})$.

The idempotent $e(s)$ associated with $s \in R$. Suppose that R is a left Artinian ring. Then for each element $s \in R$ there is the least natural number $n = n(s)$ such that $Rs^n = Rs^i$ for all $i \geq n$. By Lemma 4.1.(3), the number $n = n(s)$ is the least natural number n such that $Rs^i \cap \ker(\cdot s^i) = 0$ for all $i \geq n$ or equivalently $R = Rs^i \oplus \ker(\cdot s^i)$ for all $i \geq n$ (equivalently, $\ker(\cdot s^i) = \ker(\cdot s^n)$ for all $i \geq n$). So, for all $i \geq n, Rs^n = Rs^i$ and $\ker(\cdot s^n) = \ker(\cdot s^i)$. For each $i \geq n$, let

$$1 = e[s^i] + e'[s^i] \tag{9}$$

be the sum of orthogonal idempotents that corresponds to the decomposition $R = Rs^i \oplus \ker(\cdot s^i)$. Then $e[s^n] = e[s^i]$ and $e'[s^n] = e'[s^i]$ for all $i \geq n$. These common values are denoted by $e(s)$ and $e'(s)$, respectively. The idempotent $e(s)$ of the ring R is called the *idempotent associated with the element s* . The element s is a nilpotent iff $e(s) = 0$. The element s is a unit iff $e(s) = 1$. The element s is neither a nilpotent element nor a unit iff $e(s) \neq 0, 1$.

Orthogonal idempotents. The ring R is a left Artinian ring. Its radical $\text{rad}(R)$ is a nilpotent ideal, and so it coincides with the prime radical \mathfrak{n}_R of the ring R , and

$$\bar{R} := R/\text{rad}(R) \simeq \prod_{i=1}^s M_{n_i}(D_i) \tag{10}$$

where $\bar{R}_i := M_{n_i}(D_i)$ is the ring of $n_i \times n_i$ matrices with entries from a division ring D_i . Let $\{\bar{E}_{pq}(i) \mid p, q = 1, \dots, n_i\}$ be the matrix units of the ring \bar{R}_i . Since the radical $\text{rad}(R)$ is a nil

ideal (even a nilpotent ideal) of R , the decomposition of 1 in \overline{R} as a sum of primitive orthogonal idempotents

$$1 = \sum_{i=1}^s \sum_{j=1}^{n_i} \overline{E}_{jj}(i) \quad (11)$$

can be lifted to a decomposition of $1 \in R$ as a sum of primitive orthogonal idempotents

$$1 = \sum_{i=1}^s \sum_{j=1}^{n_i} E_{jj}(i) \quad (12)$$

and any such a lift is unique up to conjugation (i.e. up to inner automorphism) and order of idempotents (i.e. permutation of the idempotents), (Proposition 18.23.5, [6]). The sum $\overline{1}_i := \sum_{j=1}^{n_i} \overline{E}_{jj}(i)$ is the identity of the ring \overline{R}_i and let $1_i := \sum_{j=1}^{n_i} E_{jj}(i)$. Then $1 = \sum_{i=1}^s \overline{1}_i$ is the sum of central orthogonal idempotents of the ring \overline{R} , and

$$1 = \sum_{i=1}^s 1_i \quad (13)$$

is the sum of orthogonal idempotents of the ring R (in general, not necessarily central). For each non-empty subset I of the set $\{1, \dots, s\}$, let

$$e_I := 1_I := \sum_{i \in I} 1_i. \quad (14)$$

Since the idempotents $\overline{1}_i$ are central, the sum (13) is *unique up to inner automorphism* of R , i.e. if $1 = \sum_{i=1}^s 1'_i$ is another lifting with $\overline{1}'_i = \overline{1}_i$ then $1'_i = u 1_i u^{-1}$ for all i and some unit $u \in R^*$. Let \mathfrak{a} be an ideal of the ring R such that $\mathfrak{a} \not\subseteq \text{rad}(R)$. Then $(\mathfrak{a} + \text{rad}(R))/\text{rad}(R)$ is a nonzero ideal of the semisimple ring \overline{R} , and so

$$(\mathfrak{a} + \text{rad}(R))/\text{rad}(R) = \prod_{i \in I(\mathfrak{a})} M_{n_i}(D_i) = \overline{R} \overline{1}_{I(\mathfrak{a})} \quad (15)$$

for a unique non-empty subset $I(\mathfrak{a})$ of the set $\{1, \dots, s\}$ where $\overline{1}_{I(\mathfrak{a})} := \sum_{i \in I(\mathfrak{a})} \overline{1}_i$ is the central idempotent of the ring \overline{R} . The set $I(\mathfrak{a})$ is a proper subset of $\{1, \dots, s\}$ iff $\mathfrak{a} \neq R$.

Every left localization of a left Artinian ring is an idempotent left localization. Proposition 4.2 and Corollary 4.3 are about lifting (in many different ways) left denominator sets of a factor ring of a ring to the ring (under certain conditions).

Proposition 4.2 *Let R be an arbitrary ring, \mathfrak{a} be its ideal, $\overline{R} := R/\mathfrak{a}$ and S be a multiplicative set of the ring R such that for each element $a \in \mathfrak{a}$ there is an element $s \in S$ such that $sa = 0$. Let $\pi : R \rightarrow \overline{R}$, $r \mapsto \overline{r} = r + \mathfrak{a}$, and $\overline{S} := \pi(S)$.*

1. *If $\overline{S} \in \text{Ore}_l(\overline{R}, \overline{\mathfrak{b}})$ then $S \in \text{Ore}_l(R, \mathfrak{b})$ where $\mathfrak{b} = \pi^{-1}(\overline{\mathfrak{b}})$.*
2. *If $\overline{S} \in \text{Den}_l(\overline{R}, \overline{\mathfrak{b}})$ then $S \in \text{Den}_l(R, \mathfrak{b})$ where $\mathfrak{b} = \pi^{-1}(\overline{\mathfrak{b}})$ and $S^{-1}R \simeq \overline{S}^{-1}\overline{R}$.*

Proof. 1. (i) $S \in \text{Ore}_l(R)$: For given elements $s \in S$ and $r \in R$, we have to find elements $s' \in S$ and $r' \in R$ such that $s'r = r's$. Since $\overline{S} \in \text{Ore}_l(\overline{R})$, $\overline{s}_1 \overline{r} = \overline{r}_1 \overline{s}$ for some elements $s_1 \in S$ and $r_1 \in R$. Then $s_1 r - r_1 s \in \mathfrak{a}$, and so $s_2 (s_1 r - r_1 s) = 0$ for some element $s_2 \in S$. It suffices to take $s' = s_2 s_1$ and $r = s_2 r_1$.

(ii) $\text{ass}(S) = \mathfrak{b}$: Let $sr = 0$ for some elements $s \in S$ and $r \in R$. Then $\overline{s} \overline{r} = 0$ in \overline{R} , and so $\overline{r} \in \overline{\mathfrak{b}}$, hence $r \in \mathfrak{b}$, i.e. $\text{ass}(S) \subseteq \mathfrak{b}$.

Let $b \in \mathfrak{b}$. Then $\overline{s} \overline{b} = 0$ for some element $s \in S$. Then $sb \in \mathfrak{a}$, and so $s_1 sb = 0$ for some element $s_1 \in S$. Hence $\mathfrak{b} \subseteq \text{ass}(S)$.

2. (i) $S \in \text{Den}_l(R, \mathfrak{b})$: In view of statement 1, it suffices to show that $rs = 0$ for some elements $s \in S$ and $r \in R$ implies $s'r = 0$ for some element $s' \in S$. We have the equality $\overline{r}\overline{s} = 0$ in the ring \overline{R} . Then $\overline{s}_1\overline{r} = 0$ for some element $s_1 \in S$ since $\overline{S} \in \text{Den}_l(R, \overline{\mathfrak{b}})$. Then $s_1r \in \mathfrak{a}$, hence $s_2s_1r = 0$ for some element $s_2 \in S$. It suffices to take $s' = s_2s_1$.

(ii) $S^{-1}R \simeq \overline{S}^{-1}\overline{R}$: By the universal property of left localizations, the map $S^{-1}R \rightarrow \overline{S}^{-1}\overline{R}$, $s^{-1}r \mapsto \overline{s}^{-1}\overline{r}$, is a ring homomorphism which is obviously an epimorphism. Suppose that an element $s^{-1}r \in S^{-1}R$ belongs to the kernel of the epimorphism. Then $\overline{r} = 0$ in $\overline{S}^{-1}\overline{R}$, and so $\overline{s}_1\overline{r} = 0$ in \overline{R} for some element $s_1 \in S$. This means that $s_1r \in \mathfrak{a}$, and so $s_2s_1r = 0$ for some element $s_2 \in S$. Therefore, $\frac{r}{1} = 0$ and $s^{-1}r = 0$. The epimorphism is an isomorphism. \square

Corollary 4.3 *Let R be a ring and $S \in \text{Den}_l(R, \mathfrak{a})$ (respectively, $S \in \text{Ore}_l(R, \mathfrak{a})$). Then $S + \mathfrak{a} \in \text{Den}_l(R, \mathfrak{a})$ (respectively, $S + \mathfrak{a} \in \text{Ore}_l(R, \mathfrak{a})$) and $(S + \mathfrak{a})^{-1}R \simeq S^{-1}R$.*

Proof. Since $S \cap \mathfrak{a} = \emptyset$, the set $S + \mathfrak{a}$ is a multiplicative set. We keep the notation of Proposition 4.2. Since $S \in \text{Den}_l(R, \mathfrak{a})$ (respectively, $S \in \text{Ore}_l(R, \mathfrak{a})$), we have that $\overline{S + \mathfrak{a}} = \overline{S} \in \text{Den}_l(\overline{R}, 0)$ (respectively, $\overline{S} \in \text{Ore}_l(\overline{R}, 0)$). By Proposition 4.2, $S + \mathfrak{a} \in \text{Den}_l(R, \mathfrak{a})$ (respectively, $S + \mathfrak{a} \in \text{Ore}_l(R, \mathfrak{a})$) and $(S + \mathfrak{a})^{-1}R \simeq \overline{S}^{-1}\overline{R} \simeq S^{-1}R$. \square

Corollary 4.4 *Let R be a ring, \mathfrak{a} be an ideal of R such that $\mathfrak{a} \neq R$, an element $s \in R$ be such that the element $s + \mathfrak{a}$ is a unit of the ring R/\mathfrak{a} and $\bigcup_{i \geq 1} \ker_R(s^i \cdot) = \mathfrak{a}$. Then $S_s := \{s^i \mid i \in \mathbb{N}\} \in \text{Den}_l(R, \mathfrak{a})$ and $S_s^{-1}R \simeq R/\mathfrak{a}$.*

Proof. We keep the notation of Proposition 4.2. The set $\overline{S}_s := \{\overline{s}^i \mid i \in \mathbb{N}\}$ (where $\overline{s} = s + \mathfrak{a}$) consists of units of the ring $\overline{R} = R/\mathfrak{a}$, and so $\overline{S}_s \in \text{Den}_l(\overline{R}, 0)$. By Proposition 4.2, $S_s \in \text{Den}_l(R, \mathfrak{a})$. Now, it is obvious that $S_s^{-1}R \simeq R/\mathfrak{a}$ (since $\overline{S}_s \subseteq \overline{R}^*$). \square

Let S be a nonempty subset of a ring R . The set $\ker_l(S) = \{\ker(\cdot s) \mid s \in S\}$ of left ideals of the ring R is a poset with respect to \subseteq . Let $\max.\ker_l(S)$ be the set of maximal elements of the poset $\ker_l(S)$. The set $\max.\ker_l(S)$ is a non-empty set provided the ring R satisfies the a.c.c. for left annihilators. Similarly, the set $\ker_r(S) = \{\ker(s \cdot) \mid s \in S\}$ of right ideals of the ring R is a poset with respect to \subseteq . Let $\max.\ker_r(S)$ be the set of maximal elements of the poset $\ker_r(S)$. The set $\max.\ker_r(R)$ is a non-empty set provided the ring R satisfies the a.c.c. for right annihilators.

The first statement of the following theorem shows that every left localization of a left Artinian ring is an idempotent left localization.

Theorem 4.5 *Let R be a left Artinian ring and $S \in \text{Den}_l(R, \mathfrak{a})$. Then*

1. *There exists a nonzero idempotent $e \in R$ such that $S_e := \{1, e\} \in \text{Den}_l(R, \mathfrak{a})$ and the rings $S^{-1}R$ and $S_e^{-1}R$ are R -isomorphic.*
2. (a) *If $\mathfrak{a} = 0$ then $e = 1$.*
 (b) *If $\mathfrak{a} \neq 0$ then $\mathfrak{a} = (1 - e)R = 1_{I(\mathfrak{a})}R \not\subseteq \text{rad}(R)$ and the idempotent e is conjugate to $1_{CI(\mathfrak{a})} = 1 - 1_{I(\mathfrak{a})}$ where the set $I(\mathfrak{a})$ is defined in (15).*
 (c)

$$R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}, \quad \mathfrak{a} = \begin{pmatrix} 0 & 0 \\ R_{21} & R_{22} \end{pmatrix} \quad \text{and} \quad S \subseteq \begin{pmatrix} R_{11}^* & 0 \\ R_{21} & R_{22} \end{pmatrix}, \quad (16)$$

where $R_{ij} = e_i R e_j$, $e_1 = e$ and $e_2 = 1 - e_1$.

Proof. 1. If $\mathfrak{a} = 0$ then $S \subseteq R^*$ (Lemma 4.1.(1)) and so $S^{-1}R = R$. It suffices to take $e = 1$.

We can assume that $\mathfrak{a} \neq 0$. Then necessarily $\mathfrak{a} \neq R$, i.e. \mathfrak{a} is a proper ideal of the ring R and $S \not\subseteq R^*$. By Corollary 4.3, $S + \mathfrak{a} \in \text{Den}_l(R, \mathfrak{a})$ and $(S + \mathfrak{a})^{-1}R \simeq S^{-1}R$ (an R -isomorphism). Without loss of generality we may assume that $S + \mathfrak{a} \subseteq S$ (replacing S by $S + \mathfrak{a}$, if necessary).

The ring R is a left Artinian ring. Therefore, we can fix an element $s \in S$ such that $\ker(\cdot s) \in \max.\ker_l(S)$, and so $\ker(\cdot s) \neq 0$ (since $S \not\subseteq R^*$ and by Lemma 4.1.(1)). Then necessarily $Rs \cap \ker(\cdot s) = 0$, and so

$$R = Rs \oplus \ker(\cdot s) \quad (17)$$

is the direct sum of left ideals of the ring R , by Lemma 4.1.(3). Let $1 = e_1 + e_2$ be the corresponding decomposition of 1 as a sum of orthogonal idempotents. Then

$$R = \bigoplus_{i,j=1}^2 R_{ij} = \begin{pmatrix} R_{11} & R_{12} \\ R_{21} & R_{22} \end{pmatrix} \text{ where } R_{ij} := e_i R e_j,$$

$Rs = Re_1 = \begin{pmatrix} R_{11} & 0 \\ R_{21} & 0 \end{pmatrix}$ and $\ker(\cdot s) = Re_2 = \begin{pmatrix} 0 & R_{12} \\ 0 & R_{22} \end{pmatrix}$. Since $s \in Rs$, the element s has the form

$$s = \begin{pmatrix} s_{11} & 0 \\ s_{21} & 0 \end{pmatrix}, \quad (18)$$

the ring R_{11} is a left Artinian ring and the map $\cdot s : R_{11} \rightarrow R_{11}$ is an injection, by (17). By Lemma 4.1, s_{11} is a unit of the ring R_{11} , i.e. $s_{11} \in R_{11}^*$.

Since $s \cdot \begin{pmatrix} 0 & 0 \\ s_{21} & 0 \end{pmatrix} = 0$, the element $\begin{pmatrix} 0 & 0 \\ s_{21} & 0 \end{pmatrix}$ belongs to the ideal \mathfrak{a} . Then $s_{11} = \begin{pmatrix} s_{11} & 0 \\ 0 & 0 \end{pmatrix} \in S$ since $S + \mathfrak{a} \subseteq S$, and $\ker(\cdot s_{11}) = \begin{pmatrix} 0 & R_{12} \\ 0 & R_{22} \end{pmatrix}$ since $s_{11} \in R_{11}^*$. Since $s_{11}R_{21} = 0$, we have the inclusion

$$\begin{pmatrix} 0 & R_{12} \\ R_{21} & R_{22} \end{pmatrix} \subseteq \mathfrak{a}.$$

Hence, $R_{12}R_{21} \subseteq \mathfrak{a}$.

(i) $R_{12}R_{21} = 0$: Suppose that $R_{12}R_{21} \neq 0$, we seek a contradiction. Fix a nonzero element $a = \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} \in R_{12}R_{21}$. Then, for some element $t = \begin{pmatrix} t_{11} & t_{12} \\ t_{21} & t_{22} \end{pmatrix} \in S$,

$$0 = ta = \begin{pmatrix} t_{11} & t_{12} \\ t_{21} & t_{22} \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} t_{11}a & 0 \\ t_{21}a & 0 \end{pmatrix}.$$

In particular, $t_{11}a = 0$. Since $\begin{pmatrix} 0 & t_{12} \\ t_{21} & t_{22} \end{pmatrix} \in \mathfrak{a}$, the element $t_{11} \in S$. The kernel $\ker_{R_{11}}(\cdot t_{11})$ of the map $\cdot t_{11} : R_{11} \rightarrow R_{11}$ is nonzero (otherwise $t_{11} \in R_{11}^*$ since R_{11} is a left Artinian ring but $t_{11}a = 0$ and $0 \neq a \in R_{11}$, a contradiction). Then

$$\ker_R(\cdot t_{11}) \supseteq \ker_{R_{11}}(\cdot t_{11}) \bigoplus \begin{pmatrix} 0 & R_{12} \\ 0 & R_{22} \end{pmatrix} \supsetneq \begin{pmatrix} 0 & R_{12} \\ 0 & R_{22} \end{pmatrix} = \ker_R(\cdot s).$$

This contradicts to the maximality of $\ker_R(\cdot s)$. Therefore, $R_{12}R_{21} = 0$.

(ii) $S \subseteq \begin{pmatrix} R_{11}^* & R_{12} \\ R_{21} & R_{22} \end{pmatrix}$: Suppose that this inclusion does not hold, i.e. there exists an element $s' = \begin{pmatrix} s'_{11} & s'_{12} \\ s'_{21} & s'_{22} \end{pmatrix} \in S$ with $s'_{11} \notin R_{11}^*$, we seek a contradiction. Then $s'_{11} \in S$ since $S + \mathfrak{a} \subseteq S$ and

$$\begin{pmatrix} 0 & s'_{12} \\ s'_{21} & s'_{22} \end{pmatrix} \in \begin{pmatrix} 0 & R_{12} \\ R_{21} & R_{22} \end{pmatrix} \subseteq \mathfrak{a}.$$

By Lemma 4.1.(3), $\ker_{R_{11}}(\cdot s'_{11}) \neq 0$ (since $s'_{11} \notin R_{11}^*$). Then

$$\ker_R(\cdot s'_{11}) \supseteq \ker_{R_{11}}(\cdot s'_{11}) \bigoplus \begin{pmatrix} 0 & R_{12} \\ 0 & R_{22} \end{pmatrix} \supsetneq \begin{pmatrix} 0 & R_{12} \\ 0 & R_{22} \end{pmatrix} = \ker_R(\cdot s).$$

This contradicts to the maximality of $\ker_R(\cdot s)$.

(iii) $R_{12} = 0$: Let $a_{12} = \begin{pmatrix} 0 & a_{12} \\ 0 & 0 \end{pmatrix} \in R_{12}$. Since $R_{12} \subseteq \mathfrak{a}$, there is an element $t = \begin{pmatrix} t_{11} & * \\ * & * \end{pmatrix} \in S$ such that $0 = ta_{12} = \begin{pmatrix} 0 & t_{11}a_{12} \\ 0 & * \end{pmatrix}$. We must have $a_{12} = 0$ since $t_{11} \in R_{11}^*$, by (ii).

(iv) $\mathfrak{a} = \begin{pmatrix} 0 & 0 \\ R_{21} & R_{22} \end{pmatrix} = (1 - e_1)R$: The second equality is obvious. By (ii) and (iii), $\mathfrak{a} \subseteq \begin{pmatrix} 0 & 0 \\ R_{21} & R_{22} \end{pmatrix}$. By (18), $s \begin{pmatrix} 0 & 0 \\ R_{21} & R_{22} \end{pmatrix} = 0$. Hence, $\mathfrak{a} = \begin{pmatrix} 0 & 0 \\ R_{21} & R_{22} \end{pmatrix}$.

(v) $S_{e_1} = \{1, e_1\} \in \text{Den}_l(R, \mathfrak{a})$ and $S_{e_1}^{-1}R \simeq R/(1 - e_1)R = R/\mathfrak{a} \simeq S^{-1}R$ are R -isomorphisms: By (iii), (iv) and Proposition 3.1, $S_{e_1} \in \text{Den}_l(R, \mathfrak{a})$ and $S_{e_1}^{-1}R \simeq R/\mathfrak{a}$ (an R -isomorphism). By Lemma 4.1.(2), $S^{-1}R \simeq R/\mathfrak{a}$, an R -isomorphism.

2. (a) Trivial.

(c) The first two equalities of the statement (c) follow from statement 1 and Proposition 3.1, the last equality follows from (ii).

(b) Suppose that $\mathfrak{a} \neq 0$. Let $e = e_1$ and $e_2 = 1 - e_1$ be as above. By Lemma 3.1, $\mathfrak{a} = (1 - e_1)R$, hence $\mathfrak{a} \not\subseteq \text{rad}(R)$ (otherwise, $0 \neq e_2 = 1 - e_1 \in \text{rad}(R)$, a contradiction). Notice that $\text{rad}(R) = \begin{pmatrix} \text{rad}(R_{11}) & 0 \\ R_{21} & \text{rad}(R_{22}) \end{pmatrix}$ and

$$(\mathfrak{a} + \text{rad}(R))/\text{rad}(R) = \begin{pmatrix} 0 & 0 \\ 0 & R_{22}/\text{rad}(R_{22}) \end{pmatrix} \subset \begin{pmatrix} R_{11}/\text{rad}(R_{11}) & 0 \\ 0 & R_{22}/\text{rad}(R_{22}) \end{pmatrix} = R/\text{rad}(R).$$

Therefore, $\bar{e}_2 = e_2 + \text{rad}(R) = 1_{I(\mathfrak{a})} + \text{rad}(R)$ and $\bar{e}_1 = e_1 + \text{rad}(R) = 1 - 1_{I(\mathfrak{a})} + \text{rad}(R) = 1_{CI(\mathfrak{a})} + \text{rad}(R)$. Hence, $e_2 = u1_{I(\mathfrak{a})}u^{-1}$ and $e_1 = u1_{CI(\mathfrak{a})}u^{-1}$ for some unit $u \in R^*$. Then $\mathfrak{a} = u^{-1}\mathfrak{a}u = u^{-1}e_2Ru = 1_{I(\mathfrak{a})}R$. \square

An idempotent e of a ring R is called a *left triangular idempotent* if $eR(1 - e) = 0$. So, an idempotent e of R is a left triangular iff the ring $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ is *left triangular* iff $S_e = \{1, e\} \in \text{Den}_l(R)$, by Proposition 3.1 iff e is a left denominator idempotent of R .

Let R be a left Artinian ring with (13). Let

$$\mathcal{I}'_I := \mathcal{I}'_I(R) := \{e_I := \sum_{i \in I} 1_i \mid e_I R(1 - e_I) = 0\} \text{ and } \mathcal{I}''_I(R) := \mathcal{I}'_I(R) \setminus \{1\} \quad (19)$$

where I is a *nonempty* subset of $\{1, \dots, s\}$. Notice that $1 - e_I = e_{CI}$ where $CI := \{1, \dots, s\} \setminus I$ and $e_\emptyset := 0$. Clearly, $1 \in \mathcal{I}'_I$. The set \mathcal{I}'_I is a finite nonempty set that consists of left triangular idempotents of the ring R . The set $\mathcal{I}'_I(R)$ is one of the most important objects as far as left denominator sets and left localizations of R are concerned (Theorem 4.6).

Let, for a moment, R be an arbitrary ring. For a ring $A \in \text{Loc}_l(R)$, let $[A]$ be the isomorphism class of A . We usually drop the brackets. Let $\text{Loc}_l(R)/\simeq$ be the set of isomorphism classes of left localizations of the ring R . The groups $\text{Aut}(R)$ and $\text{Inn}(R)$ act on the sets $\text{Den}_l(R)$ and $\text{Loc}_l(R)$ in the obvious way: the action of an automorphism $\sigma \in \text{Aut}(R)$ on $S \in \text{Den}_l(R)$ and $S^{-1}R$ is defined as $\sigma(S)$ and $\sigma(S)^{-1}\sigma(R) = \sigma(S)^{-1}R$. Let $\text{Den}_l(R)/G$ and $\text{Loc}_l(R)/G$ be the sets of G -orbits of the groups $G = \text{Aut}(R)$, $\text{Inn}(R)$ in $\text{Den}_l(R)$ and $\text{Loc}_l(R)$, respectively. Every $\text{Aut}(R)$ -orbit is a union of $\text{Inn}(R)$ -orbits. Every isomorphism class $[A] \in \text{Loc}_l(R)/\simeq$ is a union of $\text{Aut}(R)$ -orbits and of $\text{Inn}(R)$ -orbits.

By Lemma 4.1.(4), for a left Artinian ring R the group $\text{Inn}(R)$ acts *trivially* on $\text{Loc}_l(R)$, i.e. each element of $\text{Loc}_l(R)$ is an $\text{Inn}(R)$ -orbit. Recall that the set $\text{Loc}_l(R)$ consists of R -isomorphism classes of left localizations of the ring R and if two left localizations $S^{-1}R$ and $S'^{-1}R$ are R -isomorphic we write $S^{-1}R = S'^{-1}R$.

The following theorem shows that, for a left Artinian ring R , up to isomorphism, there are only finitely many left localizations. Moreover, there are only finitely many left localizations up to R -automorphism, i.e. the set $\text{Loc}_l(R)$ is finite. The set $\text{Ass}_l(R)$ is explicitly described and it is also a finite set.

Theorem 4.6 *Let R be a left Artinian ring. Then*

1. The map $\mathcal{I}'_l(R) \rightarrow \{\text{ass}(S_e) \mid e \in \mathcal{I}'_l(R)\}$, $e \mapsto \text{ass}(S_e) = (1-e)R$ is a bijection.
2. The map $\mathcal{I}'_l(R) \rightarrow \mathcal{I}_l(R)/\sim$, $e \mapsto [e]$ is a bijection.
3. The map $\mathcal{I}'_l(R) \rightarrow \text{Loc}_l(R)$, $e \mapsto S_e^{-1}R = R/(1-e)R$, is a bijection. So, $\text{Loc}_l(R) = \{S_e^{-1}R \mid e \in \mathcal{I}'_l(R)\}$, $|\text{Loc}_l(R)| = |\mathcal{I}'_l(R)| < \infty$ and up to isomorphism there are only finitely many left localizations of the ring R .
4. Let $e, f \in \mathcal{I}'_l(R)$. Then $S_e^{-1}R = S_f^{-1}R$ iff $\text{ass}(S_e) = \text{ass}(S_f)$ iff $e = f$.
5. Let $e, f \in \mathcal{I}'_l(R)$. Then $e \geq f$ iff $[e] \geq [f]$ iff the map $S_e^{-1}R \rightarrow S_f^{-1}R$, $e^i r \mapsto e^i r = \frac{r}{1}$, is well-defined iff $\text{ass}(S_e) \subseteq \text{ass}(S_f)$.
6. The map $\mathcal{I}'_l(R) \rightarrow \text{Ass}_l(R)$, $e \mapsto (1-e)R$, is a bijection, i.e. $\text{Ass}_l(R) = \{(1-e)R \mid e \in \mathcal{I}'_l(R)\}$ is a finite set, $|\text{Ass}_l(R)| = |\mathcal{I}'_l(R)| < \infty$.

Proof. 1. By the very definition, if the idempotents e_1 and e_2 are distinct elements of the set $\mathcal{I}'_l(R)$ then $\text{ass}(S_{e_1}) \neq \text{ass}(S_{e_2})$.

2. Statement 2 follows from Theorem 4.5, Lemma 3.5 and statement 1.

3. Statement 3 follows from Theorem 4.5 and statement 1.

4. Statement 4 follows from statements 1 and 3.

5. It is obvious that $e \geq f$ iff $[e] \geq [f]$. By Lemma 3.4, $e \geq f$ iff the map $S_e^{-1}R \rightarrow S_f^{-1}R$, $e^i r \mapsto e^i r$, is well-defined. By Lemma 3.4, $e \geq f$ iff $\text{ass}(S_e) \subseteq \text{ass}(S_f)$.

6. Statement 6 follows from statement 1 and Theorem 4.5.(1). \square

Remark. Theorem 4.6 shows that not every ideal of a left Artinian ring R belongs to $\text{Ass}_l(R)$.

The core of a left denominator set of a left Artinian ring.

Definition. [4]. Let R be a ring and $S \in \text{Ore}_l(R)$. The *core* S_c of the left Ore set S is the set of all the elements $s \in S$ such that $\ker(s \cdot) = \text{ass}(S)$ where $s \cdot : R \rightarrow R$, $r \mapsto sr$. If $S_c \neq \emptyset$ then $SS_c \subseteq S_c$.

The next theorem is an explicit description of the core of a left denominator set of a left Artinian ring. In particular, it is a non-empty set.

Theorem 4.7 *Let R be a left Artinian ring, $S \in \text{Den}_l(R, \mathfrak{a})$ and $\mathfrak{a} \neq 0$. We keep the notation of Theorem 4.5. Then $S_c = \{s \in S \mid (1-e)s(1-e) = 0\} \neq \emptyset$, i.e. $S_c = \{s = \begin{pmatrix} s_{11} & 0 \\ s_{21} & 0 \end{pmatrix} \in S\}$, see (16).*

Proof. Let $s = \begin{pmatrix} s_{11} & 0 \\ s_{21} & s_{22} \end{pmatrix} \in S$. Then $s_{22} \in \mathfrak{a}$ and so $s's_{22} = 0$ for some element $s' = \begin{pmatrix} s'_{11} & 0 \\ s'_{21} & s'_{22} \end{pmatrix} \in S$. Then

$$S \ni s's = \begin{pmatrix} s'_{11} & 0 \\ s'_{21} & s'_{22} \end{pmatrix} \begin{pmatrix} s_{11} & 0 \\ s_{21} & 0 \end{pmatrix} + s's'_{22} = \begin{pmatrix} s'_{11}s_{11} & 0 \\ * & 0 \end{pmatrix}.$$

Therefore, $S'_c := \{s \in S \mid (1-e)s(1-e) = 0\} \neq \emptyset$ as $s's \in S$. Clearly, $S'_c \subseteq S_c$ since $\mathfrak{a} = \begin{pmatrix} 0 & 0 \\ * & * \end{pmatrix}$.

Conversely, if $s = \begin{pmatrix} s_{11} & 0 \\ s_{21} & s_{22} \end{pmatrix} \in S_c$ then $0 = s \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & s_{22} \end{pmatrix}$, i.e. $s \in S'_c$. Therefore, $S_c = S'_c$. \square

The maximal left denominator sets of a left Artinian ring. By Theorem 4.6, the posets (\mathcal{I}'_l, \geq) and $(\mathcal{I}_l/\sim, \geq)$ are isomorphic via the map $e \mapsto [e]$. If $e_I, e_J \in \mathcal{I}'_l$ then $e_I \geq e_J$ iff $I \supseteq J$ iff $(1-e_I)R \subseteq (1-e_J)R$ (Lemma 3.4). Let $\min \mathcal{I}'_l$ be the set of minimal elements of the poset \mathcal{I}'_l .

Lemma 4.8 *Let R be a left Artinian ring and \mathcal{I}'_i be as above.*

1. *If $e_I, e_J \in \mathcal{I}'_i$ and $e_I e_J \neq 0$ then $e_I e_J = e_{I \cap J} = e_J e_I \in \mathcal{I}'_i$.*
2. *If $e_I, e_J \in \mathcal{I}'_i$ with $I \cap J = \emptyset$ then $e_{I \cup J} = e_I + e_J \in \mathcal{I}'_i$.*
3. *If $e_I \in \min \mathcal{I}'_i$ and $e_J \in \mathcal{I}'_i$ then either $I \subseteq J$ or otherwise $I \cap J = \emptyset$.*
4. *If $e_I, e_J \in \min \mathcal{I}'_i$ and $e_I \neq e_J$ then $e_I e_J = 0$ and $I \cap J = \emptyset$.*
5. *If $e_I, e_J \in \min \mathcal{I}'_i$ and $e_I \neq e_J$ then $e_I R e_J = 0$.*

Proof. 1. It is obvious that $e_I e_J = e_{I \cap J} = e_J e_I$. In view of Proposition 3.1, we have to show that the equalities $e_I R(1 - e_I) = 0$ and $e_J R(1 - e_J) = 0$ imply the equality $e_I e_J R(1 - e_I e_J) = 0$:

$$e_I e_J R(1 - e_I e_J) = e_I e_J R(1 - e_J + e_J - e_I e_J) = e_J e_I R(1 - e_I) e_J = 0.$$

2. Since $I \cap J = \emptyset$, we have $e_I e_J = e_J e_I = 0$ and $1 - e_I - e_J = (1 - e_I)(1 - e_J)$. Then

$$(e_I + e_J)R(1 - e_I - e_J) = (e_I + e_J)R(1 - e_I)(1 - e_J) = 0.$$

By Proposition 3.1, $e_I + e_J \in \mathcal{I}'_i$.

3. Suppose that $I \cap J \neq \emptyset$. Then, by statement 1, $e_{I \cap J} = e_I e_J \in \mathcal{I}'_i$, and $e_I \geq e_{I \cap J}$. By the minimality of e_I , we must have the equality $e_I = e_{I \cap J}$, i.e. $I \subseteq J$.

4. Statement 4 follows from statement 3.

5. Using statement 4 and the fact that $e_I R(1 - e_I) = 0$, we see that $0 = e_I R(1 - e_I) e_J = e_I R e_J$.

□

Changing, if necessary, the order of the idempotents $1_1, \dots, 1_s$ we may assume that $\min \mathcal{I}'_i(R) = \{e_{I_1}, \dots, e_{I_t}\}$ where

$$I_1 = \{1, \dots, d_1\}, I_2 = \{d_1 + 1, \dots, d_1 + d_2\}, \dots, I_t = \left\{ \sum_{i=1}^{t-1} d_i + 1, \dots, \sum_{i=1}^t d_i \right\} \quad (20)$$

for some positive natural numbers d_1, \dots, d_t . The set

$$\{e_1 := e_{I_1}, \dots, e_t := e_{I_t}, e_{t+1} = 1 - \sum_{i=1}^t e_i\}$$

is the set of orthogonal idempotents of the ring R such that $1 = e_1 + \dots + e_t + e_{t+1}$ (it is possible that $e_{t+1} = 0$). By Lemma 4.8.(5), the ring R can be seen as a matrix ring

$$R = \bigoplus_{i,j=1}^{t+1} R_{ij} \quad \text{where } R_{ij} := e_i R e_j.$$

By Lemma 4.8.(5), $R_{ij} = 0$ for all $i \neq j$ such that $1 \leq i, j \leq t$. Hence, $R_{i,t+1} = e_i R(1 - e_i - \sum_{j \neq i} e_j) = 0$ for $i = 1, \dots, t$. So, the ring R has the form

$$R = \begin{pmatrix} R_{11} & 0 & 0 & \dots & 0 \\ 0 & R_{22} & 0 & \dots & 0 \\ & & \dots & & \\ 0 & \dots & 0 & R_{tt} & 0 \\ R_{t+1,1} & \dots & R_{t+1,t-1} & R_{t+1,t} & R_{t+1,t+1} \end{pmatrix} \quad (21)$$

Proposition 4.9 *Let R be a left Artinian ring and \mathcal{I}'_i be as above. Let I_1, \dots, I_t be non-empty subsets of $\{1, \dots, s\}$ such that $I_i \cap I_j = \emptyset$ for all $i \neq j$. Then $\min \mathcal{I}'_i(R) = \{e_{I_1}, \dots, e_{I_t}\}$ iff the ring R has the form (21) and none of the rings R_{11}, \dots, R_{tt} is a left triangular and, for each nonempty subset $I \subseteq \{1, \dots, s\} \setminus \bigcup_{j=1}^t I_j$, $e_I R(1 - e_I) \neq 0$.*

Proof. (\Rightarrow) Obvious.

(\Leftarrow) The conditions that the ring R has the form (21) and none of the rings R_{11}, \dots, R_{tt} are left triangular show that $\min \mathcal{I}'_l(R) \supseteq \{e_{I_1}, \dots, e_{I_s}\}$. Then the last condition of the proposition means that $\min \mathcal{I}'_l(R) = \{e_{I_1}, \dots, e_{I_s}\}$ (if the equality does not hold then there exists an element $e_I \in \min \mathcal{I}'_l(R) \setminus \{e_{I_1}, \dots, e_{I_s}\}$, and so $e_I R(1 - e_I) = 0$. By Lemma 4.8.(4), $I \subseteq \{1, \dots, s\} \setminus \bigcup_{j=1}^s I_j$, but this is impossible since $e_I R(1 - e_I) \neq 0$, by the assumption). \square

The next theorem provides a description of the maximal left denominator sets of a left Artinian ring.

Theorem 4.10 *Let R be a left Artinian ring. Then*

1. $\max.\text{Den}_l(R) = \{T_e \mid e \in \min \mathcal{I}'_l(R)\}$ where $T_e = \{u \in R \mid u + (1 - e)R \in (R/(1 - e)R)^*\}$.
2. $|\max.\text{Den}_l(R)| \leq s$ (s is the number of isomorphism classes of left simple R -modules).
3. $|\max.\text{Den}_l(R)| = s$ iff R is a semisimple ring.

Proof. 1. By Proposition 2.4, $\max.\text{Ass}_l(R) = \text{ass}.\max.\text{Den}_l(R)$. By Theorem 4.6.(1,5), $\text{ass}.\max.\text{Den}_l(R) = \{(1 - e)R \mid e \in \min \mathcal{I}'_l\}$ and for each element $e \in \min \mathcal{I}'_l$, $S_e^{-1}R \simeq R/(1 - e)R$ is a largest left quotient ring $S_{(1-e)R}^{-1}R$ where $S_{(1-e)R}$ is the largest left denominator set with $\text{ass}(S_{(1-e)R}) = (1 - e)R$. By Theorem 2.5.(3), $S_{(1-e)R} = T_e$.

2. Statement 2 follows from Lemma 4.8.(4).

3. Statement 3 follows from statement 1 and Lemma 4.8.(5). \square

Corollary 4.11 *Let R be a left Artinian ring, $S \in \text{Den}_l(R, \mathfrak{a})$, $e \in R$ be an idempotent such that $S_e \in \text{Den}_l(R, \mathfrak{a})$ and so $S_e^{-1}R \simeq R/\mathfrak{a} \simeq S^{-1}R$ (by Theorem 4.5.(1) and Proposition 3.1), and M be an R -module. Then*

1. $\mathfrak{a}^2 = \mathfrak{a}$.
2. \mathfrak{a} is a projective right R -module.
3. $S^{-1}M = 0$ iff $\mathfrak{a}M = M$ iff $(1 - e)M = M$ iff $eM = 0$.
4. For all $\mathfrak{a}, \mathfrak{a}' \in \text{Ass}_l(R)$, $\mathfrak{a}\mathfrak{a}' = \mathfrak{a} \cap \mathfrak{a}' = \mathfrak{a}'\mathfrak{a}$.

Proof. 1. We keep the notation of Theorem 4.5. Then $\mathfrak{a} = R_{21} + R_{22}$ and $\mathfrak{a}^2 = R_{22}R_{21} + R_{22}^2 = R_{21} + R_{22} = \mathfrak{a}$.

2. Since $\mathfrak{a} = (1 - e)R$, we have $R = \mathfrak{a} \oplus eR$, and so the ideal \mathfrak{a} is a projective right R -module.

3. $S^{-1}M = 0$ iff $S_e^{-1}M = 0$ iff $M = \text{tor}_{S_e}(M)$ iff $eM = 0$ iff $M = (1 - e)M = (1 - e)RM = \mathfrak{a}M$.

4. By Theorem 4.6.(6), $\mathfrak{a} = (1 - e)R$ and $\mathfrak{a}' = (1 - e')R$ for some idempotents $e, e' \in \mathcal{I}'_l(R)$. The idempotents $1 - e$ and $1 - e'$ commute. So, $\mathfrak{a} \cap \mathfrak{a}' = (1 - e)(1 - e')R = (1 - e)\mathfrak{a}' = (1 - e)R\mathfrak{a}' = \mathfrak{a}\mathfrak{a}'$. \square

Duality. Let R be an Artinian ring and R^{op} be its opposite ring. Then $\text{Den}_r(R) = \text{Den}_l(R^{op})$. Using this fact we have analogous results for right denominator sets of R . We replace the subscript 'l' by 'r' everywhere when dealing with 'right' concepts. For example, $\mathcal{I}'_r(R) := \mathcal{I}'_l(R^{op})$ and $\mathcal{I}''_r(R) := \mathcal{I}''_l(R^{op})$. Recall that $\mathcal{I}''_l(R) = \{e_I \mid e_I R e_{CI} = 0 \text{ where } \emptyset \neq I \subsetneq \{1, \dots, s\}\}$. Similarly, $\mathcal{I}''_r(R) = \{e_J \mid e_{CJ} R e_J = 0 \text{ where } \emptyset \neq J \subsetneq \{1, \dots, s\}\}$. Then the map

$$\mathcal{I}''_l(R) \rightarrow \mathcal{I}''_r(R), \quad e_I \mapsto 1 - e_I = e_{CI}, \quad (22)$$

is an order-reversing bijection of posets. Let $\min \mathcal{I}''_*(R)$ and $\max \mathcal{I}''_*(R)$ be the sets of minimal and maximal elements of the poset $\mathcal{I}''_*(R)$ where $*$ = l, r . The map (22) induces the bijections

$$\min \mathcal{I}''_l(R) \rightarrow \max \mathcal{I}''_r(R), \quad e_I \mapsto 1 - e_I = e_{CI}, \quad (23)$$

$$\max \mathcal{I}''_l(R) \rightarrow \min \mathcal{I}''_r(R), \quad e_I \mapsto 1 - e_I = e_{CI}. \quad (24)$$

Clearly,

$$\begin{aligned}\text{Loc}_l(R) \setminus [R] &= \{[R/(1 - e_I)R] = [R/e_{CI}R] \mid I \in \mathcal{I}_l''(R)\} = \{[R/e_JR] \mid J \in \mathcal{I}_r''(R)\}, \\ \text{Loc}_r(R) \setminus [R] &= \{[R/R(1 - e_J)] = [R/Re_{CJ}] \mid J \in \mathcal{I}_r''(R)\} = \{[R/Re_I] \mid I \in \mathcal{I}_l''(R)\}.\end{aligned}$$

These equalities and the bijections (23) and (24) imply the next theorem.

Theorem 4.12 *Let R be an Artinian ring. Then the map*

$$\text{Loc}_l(R) \setminus [R] \rightarrow \text{Loc}_r(R) \setminus [R], \quad [R/(1 - e_I)R] \rightarrow [R/R(1 - e_{CI})],$$

is an anti-isomorphism of posets (i.e. an order reversing bijection). In particular, $|\text{Loc}_l(R)| = |\text{Loc}_r(R)|$.

The left localization radical \mathfrak{l}_R of a left Artinian ring R . The next theorem gives an explicit description of the left localization radical $\mathfrak{l} = \mathfrak{l}_R$ of a left Artinian ring R and a criterion for $\mathfrak{l} \neq 0$.

Theorem 4.13 *Let R be a left Artinian ring. Then*

1. $\mathfrak{l} = \bigcap_{e \in \min \mathcal{I}_l'(R)} (1 - e)R = \prod_{e \in \min \mathcal{I}_l'(R)} (1 - e)R = (1 - \sum_{e \in \min \mathcal{I}_l'(R)} e)R$. If, in addition, R is a right Artinian ring then $R = \prod_{f_j \in \max \mathcal{I}_r''(R)} f_j R$.
2. $\mathfrak{l} = \begin{cases} \bigoplus_{i=1}^{t+1} R_{t+1,i} \neq 0 & \text{if } 1 \neq \sum_{e \in \min \mathcal{I}_l'(R)} e, \\ 0 & \text{otherwise.} \end{cases}$
3. $\mathfrak{l}^2 = \mathfrak{l}$ and \mathfrak{l} is a projective right R -module.
4. $\mathfrak{l} \subseteq \text{rad}(R)$ iff $\mathfrak{l} = 0$.

Proof. 1. The first equality follows from Theorem 4.10.(1). The minimal idempotents $e \in \min \mathcal{I}_l'(R)$ are orthogonal idempotents. Hence, $\bigcap_{e \in \min \mathcal{I}_l'(R)} (1 - e)R = (\prod_{e \in \min \mathcal{I}_l'(R)} (1 - e))R = (1 - \sum_{e \in \min \mathcal{I}_l'(R)} e)R$. The last equality in statement 1 follows from (23).

2. Statement 2 follows from statement 1 and (21) since $e_{t+1} = 1 - \sum_{e \in \min \mathcal{I}_l'(R)} e$.

3. Statement 3 follows from statement 1.

4. The radical $\text{rad}(R)$ is a nilpotent ideal. So, statement 4 follows from statement 3. \square

Corollary 4.14 *Let R be a left Artinian ring and $e := \sum_{e' \in \min \mathcal{I}_l'(R)} e'$. Then*

1. $S_e \in \text{Den}_l(R)$.
2. $\text{ass}(S_e) = \mathfrak{l}_R$.
3. e is the least upper bound of the set $\min \mathcal{I}_l'(R)$ in $\mathcal{I}_l'(R)$.

Proof. 1. By Lemma 4.8.(4), $e'e'' = 0$ for all distinct idempotents $e', e'' \in \min \mathcal{I}_l'(R)$. Hence, $1 - e = \prod_{e' \in \min \mathcal{I}_l'(R)} (1 - e')$ and so $eR(1 - e) = (\sum_{e' \in \min \mathcal{I}_l'(R)} e') \cdot R \cdot \prod_{e'' \in \min \mathcal{I}_l'(R)} (1 - e'') = 0$. Hence, $S_e \in \text{Den}_l(R)$, by Proposition 3.1.

2. By statement 1 and Proposition 3.1, $\text{ass}(S_e) = (1 - e)R = \mathfrak{l}_R$ (Theorem 4.13.(1)).

3. Clearly, $e \geq e'$ for all $e' \in \min \mathcal{I}_l'(R)$. Given $f \in \mathcal{I}_l'(R)$ such that $f \geq e'$ for all $e' \in \min \mathcal{I}_l'(R)$, then $f \geq e$, by Lemma 4.8.(3,4), and statement 3 follows. \square

5 Structure of left Artinian rings with zero left localization radical

In this section, a characterization of left Artinian rings with zero left localization radical is given (Theorem 5.5). A criterion is given for a left Artinian ring to be a left localization maximal ring (Theorem 5.3). For an Artinian ring, it is shown that if the left localization radical is zero then so is the right localization radical, and vice versa (Theorem 5.6).

The left localization maximal rings. These are precisely the rings in which we cannot invert anything on the left.

Definition, [2]. A ring A is called a *left localization maximal ring* if $A = Q_l(A)$ and $\text{Ass}_l(A) = \{0\}$. A ring A is called a *right localization maximal ring* if $A = Q_r(A)$ and $\text{Ass}_r(A) = \{0\}$. A ring A is called a *localization maximal ring* (or a *two-sided localization maximal ring*) if $A = Q_{l,r}(A)$ and $\text{Ass}_{l,r}(A) = \{0\}$.

Example. Let A be a simple ring. Then $Q_l(A)$ is a left localization maximal ring and $Q_r(A)$ is a right localization maximal ring. In particular, a division ring is a (left; right; and two-sided) localization maximal ring. More generally, a simple Artinian ring (i.e. the matrix ring over a division ring) is a (left; right; and two-sided) localization maximal ring.

The next theorem is a criterion for a left quotient ring of a ring to be a maximal left quotient ring.

Theorem 5.1 [2] *Let a ring A be a left localization of a ring R , i.e. $A \in \text{Loc}_l(R, \mathfrak{a})$ for some $\mathfrak{a} \in \text{Ass}_l(R)$. Then $A \in \text{max.Loc}_l(R)$ iff $Q_l(A) = A$ and $\text{Ass}_l(A) = \{0\}$, i.e. A is a left localization maximal ring.*

Corollary 5.2 *The left localization maximal rings are precisely the localizations of all the rings at their maximal left denominators sets.*

Proof. The statement follows from Theorem 2.5.(6) and Theorem 5.1. \square

Criterion for a left Artinian ring to be a left localization maximal ring. The next theorem is a criterion for a left Artinian ring to be a left localizable maximal ring.

Theorem 5.3 *Let R be a left Artinian ring. Then the following statement are equivalent.*

1. R is a left localization maximal ring.
2. $\mathcal{I}_l'(R) = \{1\}$.
3. Either $s = 1$ or, otherwise, for every proper subset I of $\{1, \dots, s\}$, $e_I R(1 - e_I) \neq 0$.

Proof. (1 \Leftrightarrow 2) Theorem 4.6.(3).

(2 \Leftrightarrow 3) Proposition 3.1. \square

Theorem 5.3 shows that ‘generically’ every left Artinian ring is left localization maximal.

Left-right symmetry of localization maximality for Artinian rings. For Artinian rings, the concept of ‘localization maximality’ is left-right symmetric as the next theorem shows.

Theorem 5.4 *An Artinian ring is left localization maximal iff it is right localization maximal.*

Proof. The result is obvious if $s = 1$. If $s > 1$ then, by Theorem 5.3, R is a left localization maximal iff $e_I R e_{CI} \neq 0$ for all proper subsets I of $\{1, \dots, s\}$ iff $e_{CI} R e_I \neq 0$ for all proper subsets I of $\{1, \dots, s\}$ iff R is a right localization maximal ring. \square

Structure of left Artinian rings with zero left localization radical.

Theorem 5.5 *Let R be a left Artinian ring and \mathfrak{l} be its left localization radical. Then*

1. *The ring R is a direct product of left localization maximal rings iff $\mathfrak{l} = 0$ iff \mathfrak{l} is a nilpotent ideal of R .*
2. *If R is a direct product of left localization maximal rings then the direct product is unique up to order, i.e. if $R = \prod_{i=1}^t A_i = \prod_{j=1}^{t'} A'_j$ are two such direct products then $t = t'$ and $A_i = A'_{\sigma(i)}$ for all $i = 1, \dots, t$ where σ is a permutation of the set $\{1, \dots, t\}$.*

Proof. 1. By Theorem 4.13, $\mathfrak{l} = 0$ iff \mathfrak{l} is a nilpotent ideal. It remains to establish the first ‘iff’ in statement 1.

(\Leftarrow) If $\mathfrak{l} = 0$ then R is a finite direct product of localization maximal rings, by Theorem 4.13.(2), Theorem 5.3 and (21).

(\Rightarrow) If $R = \prod_{i=1}^t A_i$ is a direct product of left localization maximal (necessarily left Artinian) rings A_i . Then $\mathfrak{l}_{A_i} = 0$ and $\mathfrak{l} = \prod_{i=1}^t \mathfrak{l}_{A_i} = 0$, by Corollary 2.7.

2. Let $1 = e_1 + \dots + e_t$ and $1 = e'_1 + \dots + e'_{t'}$ be the sums of central idempotents that correspond to the direct products $R = \prod_{i=1}^t A_i$ and $R = \prod_{j=1}^{t'} A'_j$. By Theorem 5.3, for each $i = 1, \dots, t$ there is a unique $j = \sigma(i)$ such that $e_i = e_j e'_j$. Then, by symmetry, $t = t'$ and $e_i = e'_{\sigma(i)}$ for some permutation σ of the set $\{1, \dots, t\}$. \square

Theorem 5.6 *Let R be an Artinian ring, \mathfrak{l} and \mathfrak{r} be the left and right localization radicals of R , respectively. Then $\mathfrak{l} = 0$ iff $\mathfrak{r} = 0$.*

Proof. $\mathfrak{l} = 0$ iff $R = \prod_{i=1}^s A_i$ is a direct product of left localization maximal rings (Theorem 5.5.(1)) iff $R = \prod_{i=1}^s A_i$ is a direct product of right localization maximal rings (Theorem 5.4) iff $\mathfrak{r} = 0$ (Theorem 5.5). \square

Remark. In general, for an Artinian ring R , $\mathfrak{l} \neq \mathfrak{r}$ (Corollary 7.13).

6 Characterization of the left localization radical of a left Artinian ring

The aim of this section is to introduce the little left localization radical of a ring and to give a characterization of it and of the left localization radical of a left Artinian ring (Theorem 6.5). A right ideal \mathfrak{a} of a ring R is called an *idempotent right ideal* if $\mathfrak{a} = fR$ for some idempotent $f \in R$. A left ideal \mathfrak{b} of a ring R is called an *idempotent left ideal* if $\mathfrak{b} = Re$ for some idempotent $e \in R$. Let $\mathcal{IL}_l(R)$ and $\mathcal{IL}_r(R)$ be the sets of left and right idempotent ideals of the ring R , respectively, and $\mathcal{IT}_l(R)$ and $\mathcal{IT}_r(R)$ be the sets of left and right idempotent ideals that in addition are *two-sided* ideals, respectively.

Lemma 6.1 *Let R be a ring and f, f' be idempotents of R . Then $fR = f'R$ iff $R(1 - f) = R(1 - f')$.*

Proof. In view of left-right symmetry it suffices to prove that the implication (\Rightarrow) holds. The equality $fR = f'R$ yields the equality $f = f'f$ ($f = f'r$ for some $r \in R$, hence $f'f = f'^2e = f'r = f$). Then $(1 - f')(1 - f) = 1 - f' - f + f'f = 1 - f'$, and so $R(1 - f') \subseteq R(1 - f)$. By symmetry, $R(1 - f) = R(1 - f')$. \square

Lemma 6.2 *Let R be a ring and f be an idempotent of R . Then fR is an ideal iff $(1 - f)Rf = 0$ iff $1 - f \in \mathcal{I}_l(R)$ iff $R(1 - f)$ is an ideal.*

Proof. Notice that $R = fR \oplus (1 - f)R$. Then fR is an ideal of R iff $(1 - f)RfR \subseteq fR$ iff $(1 - f)Rf = 0$ iff $(1 - f) \in \mathcal{I}_l(R)$, by (7). Similarly, $R = Rf \oplus R(1 - f)$. So, $R(1 - f)$ is an ideal iff $R(1 - f)Rf \subseteq R(1 - f)$ iff $(1 - f)Rf = 0$. \square

Corollary 6.3 *Let R be a ring.*

1. *The map $\mathcal{IL}_r(R) \rightarrow \mathcal{IL}_l(R)$, $fR \mapsto R(1-f)$, is a bijection with the inverse $Re \mapsto (1-e)R$.*
2. *The map $\mathcal{IT}_r(R) \rightarrow \mathcal{IT}_l(R)$, $fR \mapsto R(1-f)$, is a bijection with the inverse $Re \mapsto (1-e)R$.*

Proof. 1. By Lemma 6.1, the maps $fR \mapsto R(1-f)$ and $Re \mapsto (1-e)R$ are well-defined. Clearly, they are mutually inverse.

2. Statement 2 follows from statement 1 and Lemma 6.2. \square

Corollary 6.4 *Let R be a ring.*

1. *The map $\mathcal{I}_l(R)/\sim \rightarrow \mathcal{IT}_r(R)$, $[e] \mapsto (1-e)R$, is a bijection with the inverse $fR \mapsto [1-f]$.*
2. $\mathcal{IT}_r(R) = \{\text{ass}(S_e) \mid S_e \in \text{IDen}_l(R)\}$.

Proof. 1. Statement 1 follows from Lemma 3.5 and Lemma 6.2.

2. Statement 2 follows from statement 1. \square

Let R be an arbitrary ring. The intersection

$$\mathfrak{l}' := \mathfrak{l}'_R := \bigcap_{\mathfrak{a} \in \text{Ass}_l(R) \setminus \{0\}} \mathfrak{a}$$

is called the *little left localization radical* provided $\text{Ass}_l(R) \setminus \{0\} \neq \emptyset$ and $\mathfrak{l}' := 0$, otherwise. The set $(\mathcal{IT}_r(R), \leq)$ is a poset. For an arbitrary poset (P, \leq) , we denote by $\text{Max}(P)$ and $\text{Min}(P)$ the sets of maximal and minimal elements of P , respectively.

Characterization of the (little) left localization radical of a left Artinian ring.

Theorem 6.5 *Let R be a left Artinian ring and \mathfrak{l} be its left localization radical of R . Then*

1. $\mathfrak{l} = \bigcap_{\mathfrak{a} \in \text{Max}(\mathcal{IT}_r(R)) \setminus R} \mathfrak{a} = \prod_{\mathfrak{a} \in \text{Max}(\mathcal{IT}_r(R)) \setminus R} \mathfrak{a}$.
2. *Suppose that $\mathcal{IT}_r(R) \neq \{0\}$. Then $\mathfrak{l}' = \prod_{\mathfrak{a} \in \mathcal{IT}_r(R) \setminus \{0\}} \mathfrak{a} = \bigcap_{\mathfrak{a} \in \mathcal{IT}_r(R) \setminus \{0\}} \mathfrak{a} = \bigcap_{\mathfrak{a} \in \text{Min}(\mathcal{IT}_r(R)) \setminus \{0\}} \mathfrak{a} = \prod_{\mathfrak{a} \in \text{Min}(\mathcal{IT}_r(R)) \setminus R} \mathfrak{a}$.*

Proof. Statements 1 and 2 follow from Corollary 6.4.(1) and Corollary 4.11. \square

7 Description of left denominator sets of a left Artinian ring

In this section, a description of left denominator sets of a left Artinian ring R is given (Theorem 7.6), the sets of left localizable and non-localizable elements are described (Proposition 7.2 and Proposition 7.3). Theorem 7.4 is a criterion for the powers of a non-nilpotent element to be a left denominator set. Theorem 7.7 describes the set $\mathcal{C}_l(R)$ of completely left localizable elements of R . Theorem 7.9 is a criterion for $\mathcal{C}_l(R) = R^*$.

The set of left localizable elements of a left Artinian ring.

Lemma 7.1 *Let $S \in \text{Den}_l(R, \mathfrak{a})$ (respectively, $S \in \text{Den}(R, \mathfrak{a})$), $\sigma : R \rightarrow S^{-1}R$, $r \mapsto \frac{r}{1}$, and $G := (S^{-1}R)^*$ be the group of units of the ring $S^{-1}R$. Then $S' := \sigma^{-1}(G) \in \text{Den}_l(R, \mathfrak{a})$ (respectively, $S' := \sigma^{-1}(G) \in \text{Den}(R, \mathfrak{a})$) and $S^{-1}R \simeq S'^{-1}R$.*

Proof. It suffices to prove the statement for $S \in \text{Den}_l(R, \mathfrak{a})$. Clearly, S' is a multiplicative set.

(i) $S' \in \text{Ore}_l(R)$: We have to show that for given elements $s' \in S'$ and $r \in R$ there are elements $s'' \in S'$ and $r' \in R$ such that $s''r = r's'$.

$$\sigma(r) = \sigma(r)\sigma(s')^{-1}\sigma(s') = \sigma(s_1)^{-1}\sigma(r_1)\sigma(s')$$

for some elements $s_1 \in S$ and $r_1 \in R$. Hence, $s_2(s_1r - r_1s') = 0$ for some element $s_2 \in S$. It suffices to take $s'' = s_2s_1$ and $r' = s_2r_1$.

(ii) $\text{ass}(S') = \mathfrak{a}$: If $s'a = 0$ for some elements $s' \in S'$ and $a \in R$ then $\sigma(s')\sigma(a) = 0$ and so $a \in \ker(\sigma) = \mathfrak{a}$ since $\sigma(s') \in G$.

(iii) $S' \in \text{Den}_l(R, \mathfrak{a})$: If $as' = 0$ for some elements $a \in R$ and $s' \in S'$ then $\sigma(a)\sigma(s') = 0$ and so $a \in \ker(\sigma) = \mathfrak{a}$ since $\sigma(s') \in G$.

(iv) $S'^{-1}R \simeq S^{-1}R$: By the universal property of left localizations, the map $S'^{-1}R \rightarrow S^{-1}R$, $s'^{-1}r \mapsto s^{-1}r$, is a well-defined ring homomorphism which is obviously an epimorphism. Its kernel is equal to zero: if $s^{-1}r = 0$ then $r \in \mathfrak{a} = \text{ass}(S) = \text{ass}(S')$, and so $s'^{-1}r = 0$. \square

Definition, [4]. An element r of a ring R is called a *left localizable element* if there exists a left denominator set S of R such that $r \in S$ (and so the element $\frac{r}{1} \neq 0$ is invertible in the ring $S^{-1}R$), equivalently, if there exists a left denominator set T of R such that the element $\frac{r}{1}$ is invertible in the ring $T^{-1}R$ (Lemma 7.1). The set of left localizable elements is denoted $\mathcal{L}_l(R)$.

Clearly,

$$\mathcal{L}_l(R) = \bigcup_{S \in \text{max.Den}_l(R)} S. \quad (25)$$

Similarly, a *right localizable element* is defined and let $\mathcal{L}_r(R)$ be the set of right localizable elements of the ring R . The elements of the *set of left and right localizable elements*,

$$\mathcal{L}_{l,r}(R) = \mathcal{L}_l(R) \cap \mathcal{L}_r(R),$$

are called *left and right localizable elements*. An element $r \in R$ is called a *localizable element* if there exists a (left and right) denominator set $S \in \text{Den}(R)$ such that $r \in S$, equivalently, if there exists a (left and right) denominator set $T \in \text{Den}(R)$ such that the element $\frac{r}{1}$ is invertible in the ring $T^{-1}R$ (Lemma 7.1). The set of all localizable elements of the ring R is denoted by $\mathcal{L}(R)$. Clearly,

$$\mathcal{L}(R) \subseteq \mathcal{L}_{l,r}(R).$$

The sets

$$\mathcal{NL}_l(R) := R \setminus \mathcal{L}_l(R), \quad \mathcal{NL}_r(R) := R \setminus \mathcal{L}_r(R), \quad \mathcal{NL}_{l,r}(R) := R \setminus \mathcal{L}_{l,r}(R), \quad \mathcal{NL}(R) := R \setminus \mathcal{L}(R),$$

are called the *sets of left non-localizable; of right non-localizable; of left and right non-localizable; of non-localizable elements*, respectively, [4]. The elements of these sets are called correspondingly (eg, an element $r \in \mathcal{NL}_l(R)$ is called a *left non-localizable element*).

The next corollary is an explicit description of the set $\mathcal{L}_l(R)$ of left localizable elements for a left Artinian ring.

Proposition 7.2 *Let R be a left Artinian ring. Then $\mathcal{L}_l(R) = \{r \in R \mid r + (1-e)R \in (R/(1-e)R)^*$ for some $e \in \min \mathcal{I}'_l(R)\}$.*

Proof. Let \mathcal{R} be the RHS of the equality of the corollary. Notice that

$$\mathcal{L}_l(R) = \bigcup_{S \in \text{max.Den}_l(R)} S = \bigcup_{e \in \min \mathcal{I}'_l(R)} T_e = \mathcal{R},$$

by Theorem 4.10.(1). \square

The set of left non-localizable elements of a left Artinian ring. The next proposition describes the set of left left non-localizable elements of a left Artinian ring and gives a criterion when it is an ideal.

Proposition 7.3 *Let R be a left Artinian ring. Then*

1. *The set $\mathcal{NL}_l(R)$ of left non-localizable elements of R is equal to the set $\{r \in R \mid r + (1-e)R \notin (R/(1-e)R)^* \text{ for all } e \in \min \mathcal{I}'_l(R)\}$.*
2. *$R \cdot \mathcal{NL}_l(R) \cdot R \subseteq \mathcal{NL}_l(R)$. In particular, $\mathcal{NL}_l(R) \cdot \mathcal{NL}_l(R) \subseteq \mathcal{NL}_l(R)$.*
3. *$\mathcal{NL}_l(R)$ is an ideal of the ring R iff the ring $R/(1-e)R$ is a division ring for all $e \in \min \mathcal{I}'_l(R)$.*

Proof. 1. Statement 1 follows from Proposition 7.2.

2. Statement 2 follows from statement 1 and the fact that in a left Artinian ring every one-sided invertible element is a unit.

3. (\Leftarrow) Trivial.

(\Rightarrow) In view of statement 2, we have to show that $\mathcal{NL}_l(R) + \mathcal{NL}_l(R) \subseteq \mathcal{NL}_l(R)$. Suppose that the ring $A = R/(1-e)R$ is not a division ring for some idempotent $e \in \min \mathcal{I}'_l(R)$, we seek a contradiction. Then, up to order, the ring $\bar{A} := A/\text{rad}(A) = \prod_{i=1}^m \bar{R}_i$ is a direct product of matrix rings $\bar{R}_i = M_{n_i}(D_i)$ where $m \leq n$. Notice that an element $r \in R$ is a unit iff $r + \text{rad}(R) \in \bar{R}$ is a unit.

Case $m = 1$ and $n_1 \geq 2$. Then the elements $a = E_{11}(1)$ and $b = \sum_{i=2}^{n_1} E_{ii}(1)$ belong to $\mathcal{NL}_l(R)$ but their sum does not, $a + b \notin \mathcal{NL}_l(R)$, a contradiction.

Case $m \geq 2$. Then the elements $a = 1_1$ and $b = \sum_{i=2}^m 1_i$ belong to $\mathcal{NL}_l(R)$ but their sum, $a + b$, does not, a contradiction. \square

Criterion for the powers of an element to be a left denominator set. For a left Artinian ring R , the following theorem is an explicit criterion for the powers of a non-nilpotent element of R to be a left denominator set.

Theorem 7.4 *Let R be a left Artinian ring, $\mathcal{I}'_l(R)$ be as above, $s \in R$ be a non-nilpotent element of R , $e = e(s)$ be the idempotent associated with the element s , $S_e = \{1, e\}$ and $S_s = \{s^i \mid i \in \mathbb{N}\}$. The following statements are equivalent.*

1. $S_s \in \text{Den}_l(R)$.
2. $S_e \in \text{Den}_l(R)$ and $(1-e)s(1-e)$ is a nilpotent element.
3. $eR(1-e) = 0$ and $(1-e)s(1-e)$ is a nilpotent element.

If one of the equivalent conditions holds then $\text{ass}(S_s) = (1-e)R$ and $S_s^{-1}R \simeq S_e^{-1}R \simeq R/(1-e)R$, the core $S_{s,c}$ of the left denominator set S_s is equal to $\{s^i \mid i \in \mathbb{N}, (1-e)s^i(1-e) = 0\}$.

Proof. (2 \Leftrightarrow 3) Statements 2 and 3 are equivalent by Lemma 3.1.

(1 \Rightarrow 2) Suppose that $S_s \in \text{Den}_l(R)$. By Theorem 4.5 there exists $S_{e_1} \in \text{Den}_l(R)$ such that $S_s^{-1}R \simeq S_{e_1}^{-1}R \simeq R/(1-e_1)R$ and $\text{ass}(S_s) = \text{ass}(S_{e_1}) = (1-e_1)R =: \mathfrak{a}$. With respect to the matrix decomposition $R = \bigoplus_{i,j=1}^2 R_{ij}$ associated with the idempotent e_1 , the ring R is left triangular, i.e.

$R_{12} = 0$ and the element s has the form $s = \begin{pmatrix} u & 0 \\ v & w \end{pmatrix}$ where $u \in R_{11}^*$ (Theorem 4.5). For all $i \geq 1$,

$s^i = \begin{pmatrix} u^i & 0 \\ v_i & w^i \end{pmatrix}$ where $v_i \in R_{21}$. Since $e_2 := 1 - e_1 \in \mathfrak{a} = e_2R$, $0 = s^i e_2 = \begin{pmatrix} 0 & 0 \\ 0 & w^i \end{pmatrix}$ for some $i \geq 1$, and so $w^i = 0$.

(i) $Re_1 = Rs^i$ for all $i \geq 1$ such that $w^i = 0$: Since $s^i = \begin{pmatrix} u^i & 0 \\ v_i & 0 \end{pmatrix} \in Re_1$, we see that $Re_1 \supseteq Rs^i$.

Since $s^i = \begin{pmatrix} u^i & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ v_i & 0 \end{pmatrix}$ and $u^i \in R_{11}^*$, $Rs^i = R \begin{pmatrix} 1 & 0 \\ v_i & 0 \end{pmatrix} = Re_1$ since $\begin{pmatrix} a & 0 \\ b & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ v_i & 0 \end{pmatrix} = \begin{pmatrix} a & 0 \\ b & 0 \end{pmatrix}$.

(ii) If $w^i = 0$ then $Rs^i = Rs^{i+1}$: Since $s^{i+1} = ss^i = \begin{pmatrix} u & 0 \\ v & w \end{pmatrix} \begin{pmatrix} u^i & 0 \\ v_i & 0 \end{pmatrix} = \begin{pmatrix} u^{i+1} & 0 \\ v_{i+1} & 0 \end{pmatrix}$, have $Rs^{i+1} = Re_1 = Rs^i$, by (i).

(iii) $Re_1 = Re$: By (i) and (ii), $Re_1 = Rs^i = Re$.

(iv) $S_e \in \text{Den}_l(R, \mathfrak{a})$: By (iii) and Lemma 6.1, $\mathfrak{a} = (1 - e_1)R = (1 - e)R$. Since $(1 - e)R$ is an ideal of R , we must have $eR(1 - e) \subseteq eR \cap (1 - e)R = 0$, i.e. $eR(1 - e) = 0$. By Proposition 3.1, $S_e \in \text{Den}_l(R, \mathfrak{a})$.

Clearly, $S_s^{-1}R = R/\mathfrak{a} \simeq S_e^{-1}R$. The ring R can be seen as the matrix ring associated with the idempotent e , see (6), $R = \begin{pmatrix} R'_{11} & 0 \\ R'_{21} & R'_{22} \end{pmatrix}$. Then $s = \begin{pmatrix} u' & 0 \\ v' & w' \end{pmatrix}$ and $s^i = \begin{pmatrix} u'^i & 0 \\ v'_i & w'^i \end{pmatrix}$ for all $i \geq 1$ where $v'_i \in R'_{21}$. Since $1 - e \in \mathfrak{a}$ we must have $0 = s^i(1 - e) = \begin{pmatrix} 0 & 0 \\ 0 & w'^i \end{pmatrix}$ for some $i \geq 1$, i.e. $w' = (1 - e)s(1 - e)$ is a nilpotent element.

(1 \Leftrightarrow 2) This implication and all the statements of the theorem is a particular case of Lemma 7.5. \square

Lemma 7.5 *Let R be a left triangular matrix ring $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ where R_{11} and R_{22} are arbitrary rings and R_{21} is an arbitrary (R_{22}, R_{11}) -bimodule. Let $s = \begin{pmatrix} u & 0 \\ v & w \end{pmatrix} \in R$ where $u \in R_{11}^*$ and w a nilpotent element of the ring R_{22} . Then the set $S_s = \{s^i \mid i \in \mathbb{N}\}$ is a left denominator set of the ring R such that $\text{ass}(S_s) = \begin{pmatrix} 0 & 0 \\ R_{21} & R_{22} \end{pmatrix}$, $S_s^{-1}R \simeq R/\text{ass}(S_s) \simeq R_{11}$ and the core $S_{s,c}$ of S_s is equal to $\{s^i \mid i \in \mathbb{N}, w^i = 0\}$.*

Proof. First, we prove all the statements but the one about the core. Then, the statement about the core will follow, see (v). Fix a natural number j such that $w^j = 0$, and so $s^j = \begin{pmatrix} u^j & 0 \\ * & 0 \end{pmatrix}$. Notice that for each natural number $k \geq 1$, $S_s \in \text{Den}_l(R)$ iff $S_{s^k} \in \text{Den}_l(R)$; and if $S_s \in \text{Den}_l(R)$ then $\text{ass}(S_s) = \text{ass}(S_{s^k})$ and $S_s^{-1}R \simeq S_{s^k}^{-1}R$. We may assume that $w = 0$ from the very beginning, i.e. $s = \begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix}$. Notice that $t = \begin{pmatrix} 1 & 0 \\ -vu^{-1} & 1 \end{pmatrix} \in R^*$ and

$$\omega_t(s) = \begin{pmatrix} 1 & 0 \\ -vu^{-1} & 1 \end{pmatrix} \begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -vu^{-1} & 1 \end{pmatrix}^{-1} = \begin{pmatrix} u & 0 \\ 0 & 0 \end{pmatrix}. \quad (26)$$

Every ideal of the ring R is invariant under the inner automorphisms. So, by replacing the element s by $\omega_t(s)$ we may assume that $s = \begin{pmatrix} u & 0 \\ 0 & 0 \end{pmatrix}$.

(i) $S_s \in \text{Ore}_l(R)$: We have to show that for given elements $s^i \in S_s$ and $r = \begin{pmatrix} a & 0 \\ b & c \end{pmatrix} \in R$, there are elements $s^j \in S_s$ and $r' = \begin{pmatrix} a' & 0 \\ b' & c' \end{pmatrix} \in R$ such that $s^j r = r' s^i$. Notice that

$$sr = \begin{pmatrix} ua & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} uau^{-i} & 0 \\ 0 & 0 \end{pmatrix} s^i.$$

So, it suffices to take $j = 1$.

(ii) $\text{ass}(S_s) = \ker(\cdot s) = \mathfrak{a}$ where $\mathfrak{a} := \begin{pmatrix} 0 & 0 \\ R_{21} & R_{22} \end{pmatrix}$ since $u \in R_{11}^*$.

(iii) $S_s \in \text{Den}_l(R, \mathfrak{a})$: $\ker(\cdot s) = \begin{pmatrix} 0 & 0 \\ 0 & R_{22} \end{pmatrix} \subseteq \mathfrak{a}$.

(iv) $S_s^{-1}R \simeq R/\mathfrak{a} \simeq R_{11}$ since $u \in R_{11}^*$.

(v) Coming back to the most general situation, then $s^i = \begin{pmatrix} u^i & 0 \\ * & w^i \end{pmatrix} \in S_{s,c}$ iff $w^i R_{22} = 0$ iff $w^i = 0$. \square

Classification of all the denominator sets of a left Artinian ring. The next theorem gives a criterion for a multiplicative set of a left Artinian ring to be a left denominator set, also it is an explicit description of all the left denominator sets of R .

Theorem 7.6 *Let R be a left Artinian ring and S be a multiplicative set of R . The following statements are equivalent.*

1. $S \in \text{Den}_l(R)$.
2. There is a nonzero idempotent $e \in R$ such that $eR(1-e) = 0$, $S \subseteq \begin{pmatrix} R_{11}^* & 0 \\ R_{21} & R_{22} \end{pmatrix}$ and there is an element $s \in S$ such that $s = \begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix}$ where $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ is the matrix ring associated with the idempotent e .
3. There is a unit $\lambda \in R^*$, an idempotent $e \in \mathcal{I}'_l(R)$ and an element $s \in S$ such that $\lambda S \lambda^{-1} \subseteq \begin{pmatrix} R_{11}^* & 0 \\ R_{21} & R_{22} \end{pmatrix}$ and $\lambda s \lambda^{-1} = \begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix}$ where $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ is the matrix ring associated with the idempotent e .
4. There is an element $s \in S$ such that $S_s \in \text{Den}_l(R)$ and the images of all the elements of S in the ring $S_s^{-1}R$ are units.

If one of the equivalent conditions holds then $\text{ass}(S) = \ker_R(s \cdot)$ in cases 2 and 3 regardless of the choice of s .

Proof. (1 \Rightarrow 2) The existence of the idempotent e follows from Theorem 4.5.(2). Since $e_2 := 1 - e \in \text{aass}(S)$, there exists an element $s = \begin{pmatrix} u & 0 \\ v & w \end{pmatrix} \in S$ such that $0 = se_2 = \begin{pmatrix} 0 & 0 \\ 0 & w \end{pmatrix}$, i.e. $w = 0$.

(2 \Leftrightarrow 3) This implication is obvious due to the fact that every idempotent $e \in R$ such that $eR(1-e) = 0$ is conjugate to an idempotent of the set $\mathcal{I}_e(R)'$.

(2 \Rightarrow 4) Using the inner automorphism ω_t as in (26), and replacing the element s by $\omega_t(s)$ we may assume that $s = \begin{pmatrix} u & 0 \\ 0 & 0 \end{pmatrix}$, i.e. $v = 0$ in statement 2. By Theorem 7.4, $S_s \in \text{Den}_l(R)$ and $S_s^{-1}R \simeq R_{11}$. So, the images of all the elements of S in the ring $S_s^{-1}R$ are units.

(4 \Rightarrow 1) This implication is a particular case of Proposition 4.2.(2) where $\mathfrak{a} = \text{ass}(S_s) = \cup_{i \geq 1} \ker(s^i \cdot)$. In particular, $S \in \text{Den}_l(R, \mathfrak{a})$. \square

Theorem 7.6 gives the following algorithm of obtaining all the left denominator sets S of a left Artinian ring R . Choose an idempotent e of $\mathcal{I}'_l(R)$, there are only finitely many of them. Then we have the triangular decomposition of the ring $R = \begin{pmatrix} R_{11} & 0 \\ R_{21} & R_{22} \end{pmatrix}$ associated with the idempotent e . Choose an arbitrary element $s = \begin{pmatrix} u & 0 \\ v & 0 \end{pmatrix}$ with $u \in R_{11}^*$. Choose an arbitrary set of elements $s_i = \begin{pmatrix} u_i & 0 \\ v_i & w_i \end{pmatrix}$, $i \in I$, where $u_i \in R_{11}^*$ for all $i \in I$. Consider the monoid S' generated by the set $\{s, s_i \mid i \in I\}$. Then an arbitrary left denominator set S of R is of the type $\lambda S' \lambda^{-1}$ where $\lambda \in R^*$.

The set of completely left localizable elements of a left Artinian ring.

Definition, [4]. For an arbitrary ring R , the intersection

$$C_l(R) := \bigcap_{S \in \text{max.Den}_l(R)} S$$

is called the set of *completely left localizable elements* of R and an element of the set $\mathcal{C}_l(R)$ is called a *completely localizable element*.

By Proposition 2.8, $S_l(R) \subseteq \mathcal{C}_l(R)$. In general, this inclusion is strict, see Theorem 7.9. Moreover, Theorem 7.9 is a criterion for $S_l(R) = \mathcal{C}_l(R)$ for a left Artinian ring R (notice that $S_l(R) = \mathcal{C}_R = R^*$). The next theorem describes the set of completely left localizable elements of a left Artinian ring.

Theorem 7.7 *Let R be a left Artinian ring. Then $\mathcal{C}_l(R) = \{r \in R \mid r + (1 - e)R \in (R/(1 - e)R)^*$ for all $e \in \min \mathcal{I}'_l(R)\}$.*

Proof. The theorem follows from Theorem 4.10. \square

A left Artinian ring R is a *strongly left triangular* if there is an idempotent $e_I = \sum_{i \in I} 1_i$ for some proper subset I of $\{1, \dots, s\}$ such that $e_I R (1 - e_I) = 0$. This definition does not depend on the choice of the idempotents $1_1, \dots, 1_s$ since any two sets of them are conjugate. A left Artinian ring R which is not a strongly left triangular is called a *non-strongly-left-triangular*.

Lemma 7.8 *Let R be a left Artinian ring. Then*

1. *The ring R is a non-strongly-left-triangular ring iff $\mathcal{I}_l(R) = \{1\}$ iff $\mathcal{I}'_l(R) = \{1\}$ iff R is a left localization maximal ring.*
2. *If the ring R is non-strongly-left-triangular then $\mathcal{C}_l(R) = R^*$.*

Proof. 1. Theorem 5.3.

2. This follows from statement 1, Theorem 7.7 and Theorem 5.3 as $\mathcal{I}_l(R) = \{1\}$. \square

The next theorem is a criterion for $\mathcal{C}_l(R) = R^*$.

Theorem 7.9 *Let R be a left Artinian ring. We assume that (20) and (21) hold. In particular, $\min \mathcal{I}'_l(R) = \{e_{I_1}, \dots, e_{I_t}\}$. Then the following statements are equivalent.*

1. $\mathcal{C}_l(R) = R^*$.
2. $\bigcup_{i=1}^t I_i = \{1, \dots, s\}$.
3. $R \simeq \prod_{i=1}^t R_{ii}$.
4. R is a direct product $\prod_{i=1}^{t'} R_i$ of (necessarily left Artinian) non-strongly-left-triangular rings R_i .
5. $\mathfrak{l}_R = 0$.

If one of the equivalent conditions holds then $t = t'$ and, up to order, $R_i \simeq R_{ii}$ for $i = 1, \dots, t$.

Proof. (2 \Leftrightarrow 3) Obvious, see (21).

(3 \Rightarrow 4) Lemma 7.8.

(1 \Rightarrow 2) Suppose that statement 2 does not hold, we seek a contradiction. In view of the decomposition (21), an element $r = (r_{ij}) \in R = \bigoplus_{i,j=1}^{t+1} R_{ij}$ is a unit iff $r_{ii} \in R_{ii}^*$ for $i = 1, \dots, t+1$. By Theorem 7.7, the element $a = (a_{ij})$ which is a diagonal matrix with $a_{11} = 1, \dots, a_{tt} = 1$ and $a_{t+1,t+1} = 0$ is a non-zero element which is not a unit but belongs to $\mathcal{C}_l(R)$, a contradiction.

(4 \Rightarrow 1) By Theorem 2.6,

$$\max.\text{Den}_l\left(\prod_{i=1}^{t'} R_i\right) = \prod_{i=1}^{t'} \max.\text{Den}_l(R_i).$$

By Lemma 7.8.(2), $\max.\text{Den}_l(R_i) = \{R_i^*\}$. Hence, $\mathcal{C}_l(R) = R^*$, by Theorem 2.6. Therefore, statements 1–4 are equivalent.

The fact that $t = t'$ follows from Lemma 7.8.(1) and Lemma 7.10. Since none of the rings R_{ii} and R_i where $i = 1, \dots, t$, is a product of rings then it is well-known and easy to prove that, up to order, $R_i \simeq R_{ii}$ (even the equality hold) for all i .

(3 \Leftrightarrow 5) This equivalence follows from Theorem 5.5, Lemma 7.8.(1) and the equivalence 1 \Leftrightarrow 3. \square

Lemma 7.10 *Let $R = \prod_{i=1}^n R_i$ be a product of left Artinian rings R_i . Then*

1. $\mathcal{I}_l(R) = \{e_{i_1} + \dots + e_{i_\nu} \mid e_i \in \mathcal{I}_l(R_i), i, \nu = 1, \dots, n\}$.
2. $\min \mathcal{I}_l(R) = \prod_{i=1}^n \min \mathcal{I}_l(R_i)$.

Proof. Straightforward. \square

The rings of lower and upper triangular matrices over a division ring. Let D be a division ring, $L_n = L_n(D)$ and $U_n = U_n(D)$ be the rings of lower and upper triangular matrices, respectively, when $n \geq 2$. Let E_{ij} be the matrix units where $i, j = 1, \dots, n$. The decomposition (13) takes the form $1 = 1_1 + \dots + 1_n$ where $1_i := E_{ii}$.

Lemma 7.11 *Let $n \geq 2$ and $1_{[s]} = E_{11} + \dots + E_{ss}$. Then*

1. $\mathcal{I}_l(L_n) = \{1_{[s]} \mid [s] = \{1, \dots, s\}, s = 1, \dots, n\}$ and $\min \mathcal{I}'_l(L_n) = \{E_{11}\}$.
2. $\max.\text{Den}_l(L_n) = \{T_{E_{11}}\}$ and $T_{E_{11}}^{-1} L_n \simeq S_{E_{11}}^{-1} L_n \simeq D$ (see Theorem 4.10.(1)).
3. $\mathfrak{l}_{L_n} = \text{ass}(T_{E_{11}}) = (1 - E_{11})L_n$.

Proof. 1. Statement 1 is obvious.

2. Statement 2 follows from statement 1, Theorem 4.10 and Proposition 3.1.

3. Statement 3 follows from statements 1 and 2. \square

Corollary 7.12 *Let $n \geq 2$. Then*

1. $\mathcal{I}_l(U_n) = \{1_{[s]'} \mid [s] := \{s, s+1, \dots, n\}, s = 1, \dots, n\}$ and $\min \mathcal{I}'_l(U_n) = \{E_{nn}\}$.
2. $\max.\text{Den}_l(U_n) = \{T_{E_{nn}}\}$ and $T_{E_{nn}}^{-1} U_n \simeq S_{E_{nn}}^{-1} U_n \simeq D$.
3. $\mathfrak{l}_{U_n} = (1 - E_{nn})U_n$.

Proof. The D -homomorphism

$$U_n \rightarrow L_n, \quad E_{ij} \mapsto E_{n+1-i, n+1-j}, \quad i = 1, \dots, n,$$

is a ring isomorphism, and the results follows from Lemma 7.11. \square

Let $D^{op} := \{d^0 \mid d \in D\}$ be the division ring *opposite* to the division ring D , i.e. $d_1^0 d_2^0 := (d_2 d_1)^0$. The map $D \rightarrow D^0, d \mapsto d^0$, is a ring anti-isomorphism. The map

$$T_0 : L_n(D) \rightarrow U_n(D^0), \quad \sum d_{ij} E_{ij} \mapsto \sum d_{ij}^0 E_{ji}, \quad (27)$$

is an anti-isomorphism. Let $\mathcal{I}'_r(R)$ be the right analogue of $\mathcal{I}'_l(R)$:

$$\mathcal{I}'_r(R) := \{e_I \mid (1 - e_I)R e_I = 0\}. \quad (28)$$

Let \mathfrak{r}_R be the right localization radical of a ring R .

Corollary 7.13 *Let $n \geq 2$. Then*

1. $\mathcal{I}_r(L_n) = \{1_{[s]'} \mid [s]' := \{s, s+1, \dots, n\}, s = 1, \dots, n\}$ and $\min \mathcal{I}_r(L_n) = \{E_{nn}\}$.
2. $\max.\text{Den}_r(L_n) = \{T'_{E_{nn}}\}$ and $L_n T'_{E_{nn}}{}^{-1} \simeq L_n S_{E_{nn}}^{-1} \simeq D$ where $T'_{E_{nn}}$ is the right analogue of T_e , see Theorem 4.10.(1).
3. $\mathfrak{r}_{L_n} = L_n(1 - E_{nn})$ and $\mathfrak{l}_{L_n} \neq \mathfrak{r}_{L_n}$.

Proof. These follows from Corollary 7.12 and (27). \square

Corollary 7.14 *Let $n \geq 2$. Then*

1. $\mathcal{I}_r(U_n) = \{1_{[s]} \mid [s] = \{1, \dots, s\}, s = 1, \dots, n\}$ and $\min \mathcal{I}_r(U_n) = \{E_{11}\}$.
2. $\max.\text{Den}_r(R_n) = \{T'_{E_{11}}\}$ and $U_n T'_{E_{11}}{}^{-1} \simeq U_n S_{E_{11}}^{-1} \simeq D$.
3. $\mathfrak{r}_{U_n} = U_n(1 - E_{11})$ and $\mathfrak{l}_{U_n} \neq \mathfrak{r}_{U_n}$.

Proof. These follows from Lemma 7.11 and (27). \square

8 Localizations of Artinian rings

This section is about (left and right) denominator sets, $\text{Den}(R)$, and localizations $\text{Loc}(R) := \{S^{-1}R = RS^{-1} \mid S \in \text{Den}(R)\}$ of an Artinian ring R . The results of this section are analogous to their left versions but much more simpler due to Corollary 3.2. Their proofs follow from the left analogues in a straightforward manner and are left for the reader as an easy exercise. For the two-sided (i.e. left and right) concepts we use the same notations but the subscript ‘1’ is dropped, eg $\max.\text{Den}(R)$ is the set of maximal (left and right) denominator sets of R and $\text{Ass}(R) = \{\text{ass}(S) \mid S \in \max.\text{Den}(R)\}$. Briefly, for (left and right) denominator sets and localizations of Artinian rings, the central idempotents play a crucial role.

Let R be a left Artinian ring. It can be uniquely (up to permutation) presented as a direct product of rings

$$R = \prod_{i=1}^t R_i \quad (29)$$

where R_i are necessarily left Artinian rings none of which is a direct product of two rings. Let

$$1 = \sum_{i=1}^t e_i \quad (30)$$

be the corresponding sum of central idempotents and let

$$\mathcal{I}(R) = \{e_i \mid i = 1, \dots, t\} \text{ and } \mathcal{I}'(R) = \{e_I := \sum_{i \in I} e_i \mid \emptyset \neq I \subseteq \{1, \dots, t\}\}. \quad (31)$$

Clearly, $R_i = e_i R$ and none of e_i is a sum of two nonzero central idempotents of R .

The first statement of the following theorem shows that every localization of an Artinian ring is a (central) idempotent localization.

Theorem 8.1 *Let R be an Artinian ring and $S \in \text{Den}(R, \mathfrak{a})$. Then*

1. *There exists a central idempotent $e \in R$ such that $S_e := \{1, e\} \in \text{Den}(R, \mathfrak{a})$ and the rings $S^{-1}R$ and $S_e^{-1}R$ are R -isomorphic.*
2. (a) *If $\mathfrak{a} = 0$ then $e = 1$.*
 (b) *If $\mathfrak{a} \neq 0$ then $\mathfrak{a} = (1 - e)R \not\subseteq \text{rad}(R)$.*

$$(c) \quad R = \begin{pmatrix} R_{11} & 0 \\ 0 & R_{22} \end{pmatrix}, \quad \mathfrak{a} = \begin{pmatrix} 0 & 0 \\ 0 & R_{22} \end{pmatrix} \quad \text{and} \quad S \subseteq \begin{pmatrix} R_{11}^* & 0 \\ 0 & R_{22} \end{pmatrix}, \quad (32)$$

where $R_{ij} = e_i R e_j$, $e_1 = e$ and $e_2 = 1 - e_1$.

The following theorem shows that, for an Artinian ring R there are only finitely many left localizations. Moreover, there are only finitely many left localizations up to R -automorphism, i.e. the set $\text{Loc}(R)$ is finite. The set $\text{Ass}(R)$ is explicitly described and it is also a finite set.

Theorem 8.2 *Let R be an Artinian ring. Then*

1. *The map $\mathcal{I}'(R) \rightarrow \{\text{ass}(S_e) \mid e \in \mathcal{I}(R)\}$, $e \mapsto \text{ass}(S_e) = (1 - e)R$ is a bijection.*
2. *The map $\mathcal{I}'(R) \rightarrow \text{Loc}(R)$, $e \mapsto S_e^{-1}R = R/(1 - e)R$, is a bijection. So, $\text{Loc}(R) = \{S_e^{-1}R \mid e \in \mathcal{I}'(R)\}$, $|\text{Loc}(R)| = |\mathcal{I}'(R)| = 2^t - 1 < \infty$ and up to isomorphism there are only finitely many left localizations of the ring R .*
3. *The map $\mathcal{I}'(R) \rightarrow \text{Ass}(R)$, $e \mapsto (1 - e)R$, is a bijection, i.e. $\text{Ass}(R) = \{(1 - e)R \mid e \in \mathcal{I}'(R)\}$ is a finite set, $|\text{Ass}(R)| = |\mathcal{I}'(R)| = 2^t - 1 < \infty$.*

Let R be a ring and $S \in \text{Ore}(R)$. The *core* S_c of the Ore set S is the set of all the elements $s \in S$ such that $\ker(\cdot s) = \ker(\cdot s) = \text{ass}(S)$ where $\cdot s : R \rightarrow R$, $r \mapsto rs$. If $S_c \neq \emptyset$ then $SS_cS \subseteq S_c$. The next theorem is an explicit description of the core of a denominator set of an Artinian ring. In particular, it is a non-empty set.

Theorem 8.3 *Let R be an Artinian ring, $S \in \text{Den}(R, \mathfrak{a})$ and $\mathfrak{a} \neq 0$. We keep the notation of Theorem 8.1. Then $S_c = \{s \in S \mid (1 - e)s = 0\} \neq \emptyset$, i.e. $S_c = \{s = \begin{pmatrix} s_{11} & 0 \\ 0 & 0 \end{pmatrix} \in S\}$, see (32).*

Theorem 8.4 *Let R be an Artinian ring and $\mathcal{I}'(R)$ be as above. Then*

1. *there are precisely $2^t - 1$ idempotent (left and right) denominator sets. Moreover, the map*

$$\mathcal{I}'(R) \rightarrow \text{IDen}(R), \quad e_I \mapsto S_{e_I} = \{1, e_I\},$$

is a bijection where $\text{ass}(S_{e_I}) = R(1 - e_I)$ and $S_{e_I}^{-1}R \simeq R/R(1 - e_I) \simeq \prod_{i \in I} R_i$.

2. *$\max.\text{Den}(R) = \{S_i := R_1 \times \cdots \times R_{i-1} \times R_i^* \times R_{i+1} \times \cdots \times R_t \mid i = 1, \dots, t\}$ and $\max.\text{Loc}(R) = \{R_i \mid i = 1, \dots, t\}$.*
3. *The (left and right) localization radical $\cap_{S \in \max.\text{Den}(R)} \text{ass}(S)$ of R is equal to zero.*
4. *The set $\mathcal{C}(R) := \cap_{i=1}^t S_i$ of completely (left and right) localizable elements of R is the group R^* of units of R .*
5. *The set of (left and right) localizable elements $\mathcal{L}(R) = \bigcup_{i=1}^t S_i$ is equal to $\{r = (r_1, \dots, r_t) \in \prod_{i=1}^t R_i \mid r_i \in R_i^* \text{ for some } i\}$.*
6. *The set of (left and right) non-localizable elements $\mathcal{NL}(R) := R \setminus \mathcal{L}(R)$ is equal to $\{(r_1, \dots, r_n) \in \prod_{i=1}^t R_i \mid r_i \text{ is a zero divisor of } R_i \text{ for } i = 1, \dots, t\}$.*
7. (a) *Every (left and right) denominator set $S \in \text{Den}(R)$ contains precisely one central idempotent e_I where $\emptyset \neq I \in 2^{\mathcal{I}'(R)}$ such that $\text{ass}(S) = R(1 - e_I)$.*
 (b) *Every (left and right) denominator set $S \in \text{Den}(R)$ contains precisely one central idempotent e_I where $\emptyset \neq I \in 2^{\mathcal{I}'(R)}$ such that $S^{-1}R \simeq S_e^{-1}R$ is an R -isomorphism.*
8. *Every (left and right) denominator set S of R is obtained in the following way: fix $e_I \in \mathcal{I}'(R)$ and take a multiplicative submonoid S of (R, \cdot) such that for all elements $r = (r_1, \dots, r_n) \in S$, $r_i \in R_i^*$ for $i \in I$, and there exists an element $r' = (r'_1, \dots, r'_m) \in S$ with $r'_j = 0$ for all $j \notin I$. Then $S \in \text{Den}(R)$.*

Theorem 8.5 *Let R be an Artinian ring and $r = (r_1, \dots, r_t) \in \prod_{i=1}^t R_i$ be a non-nilpotent element. Then $S_r = \{r_i \mid i \in \mathbb{N}\} \in \text{Den}(R)$ iff each r_i is either a unit or a nilpotent element.*

Corollary 8.6 *All the denominator sets of R consists of units iff 1 is the only central idempotent of R .*

9 Rings with left Artinian left quotient ring

The aim of this section is to show that if the left quotient ring $Q_l(R)$ of a ring R is a left Artinian ring then $|\text{max.Den}_l(R)| < \infty$ (Theorem 9.1.(1)). Recall that the largest left quotient ring $Q_l(R)$ of R is a left Artinian ring iff the (classical) left quotient ring $Q_{l,c}(R) := \mathcal{C}_R^{-1}R$ is a left Artinian ring, and in this case $S_l(R) = \mathcal{C}_R$, [3].

Let R be a ring, $S \in \text{Den}_l(R)$, M be an R -module and N, L be submodules of M . We say that N and L are S -equal in M and write $N \stackrel{S}{=} L$ if $S^{-1}N = S^{-1}L$ in $S^{-1}M$. Clearly, N and L are S -equal iff, for each pair of elements $n \in N$ and $l \in L$, $sn \in L$ and $tl \in N$ for some elements $s, t \in S$.

Theorem 9.1 *Let R be a ring such that $Q_l(R)$ is a left Artinian ring and s be the number of iso-classes of simple left $Q_l(R)$ -modules. Then*

1. *the map $\text{max.Den}_l(R) \rightarrow \text{max.Den}_l(Q_l(R))$, $S \mapsto SQ_l(R)^*$, is a bijection with the inverse $T \mapsto T \cap R$. In particular, $|\text{max.Den}_l(R)| = |\text{max.Den}_l(Q_l(R))| \leq s < \infty$.*
2. *$|\text{max.Den}_l(R)| = s$ iff $Q_l(R)$ is a semisimple ring iff R is a semiprime left Goldie ring.*
3. *The map $\text{max.Ass}_l(R) \rightarrow \text{max.Ass}_l(Q_l(R))$, $\mathfrak{a} \mapsto S_l(R)^{-1}\mathfrak{a}$, is a bijection with the inverse $\mathfrak{b} \mapsto \mathfrak{b} \cap R$. In particular, $|\text{max.Ass}_l(R)| = |\text{max.Ass}_l(Q_l(R))| \leq s < \infty$.*
4. *For all ideals $\mathfrak{a}, \mathfrak{a}' \in \text{max.Ass}_l(R)$, $\mathfrak{a}\mathfrak{a}' \stackrel{S}{=} \mathfrak{a} \cap \mathfrak{a}' \stackrel{S}{=} \mathfrak{a}'\mathfrak{a}$ and $\mathfrak{a}^2 \stackrel{S}{=} \mathfrak{a}$.*

Proof. 1. Statement 1 follows from Proposition 2.8 and Theorem 4.10.

2. Statement 2 follows from Theorem 4.10.(3).

3. Statement 2 follows from the equality $\text{max.Ass}_l(R) = \{\text{ass}_l(S) \mid S \in \text{max.Den}_l(R)\}$ (Proposition 2.4), statement 1 and the fact that the largest left quotient ring $Q_l(R)$ is a left Artinian ring.

4. Statement 3 follows from the fact that for all ideals $\mathfrak{b}, \mathfrak{b}' \in \text{Ass}_l(Q_l(R))$, $\mathfrak{b}\mathfrak{b}' = \mathfrak{b} \cap \mathfrak{b}' = \mathfrak{b}'\mathfrak{b}$ (Corollary 4.11). \square

Lemma 9.2 *Let R be a ring such that $Q_l(R)$ is a left Artinian ring, $S_l = S_l(R)$ and $\text{Ass}_l(R, S_l) := \{\mathfrak{a} \in \text{Ass}_l(R) \mid \mathfrak{a} \cap S_l = \emptyset\}$. Then for all ideals $\mathfrak{a}, \mathfrak{a}' \in \text{Ass}_l(R)$,*

1. $\mathfrak{a}^2 \stackrel{S_l}{=} \mathfrak{a}$.
2. $\mathfrak{a}\mathfrak{a}' \stackrel{S_l}{=} \mathfrak{a} \cap \mathfrak{a}' \stackrel{S_l}{=} \mathfrak{a}'\mathfrak{a}$

Proof. The ring $Q_l := Q_l(R)$ is a left Artinian ring. So, for all ideals \mathfrak{a} of R , $S_l^{-1}\mathfrak{a}$ are ideals in Q_l . Clearly, $S_l^{-1}\mathfrak{a} = Q_l$ iff $\mathfrak{a} \cap S_l \neq \emptyset$. So, if one of the ideals in statements 1 and 2 meets S_l , then statements 1 and 2 hold. We can assume that $\mathfrak{a}, \mathfrak{a}' \in \text{Ass}_l(R, S_l)$. Choose $S \in \text{Den}_l(R, \mathfrak{a})$ and $S' \in \text{Den}_l(R, \mathfrak{a}')$. Let Q_l^* be the group of units of the ring Q_l . Then $SQ_l^* \in \text{Den}_l(Q_l, \mathfrak{b} := S_l^{-1}\mathfrak{a})$ and $S'Q_l^* \in \text{Den}_l(Q_l, \mathfrak{b}' := S_l^{-1}\mathfrak{a}')$. By Corollary 4.11, $\mathfrak{b}\mathfrak{b}' = \mathfrak{b} \cap \mathfrak{b}' = \mathfrak{b}'\mathfrak{b}$. Then statements 1 and 2 follow. \square

Acknowledgements

The work is partly supported by the Royal Society and EPSRC.

References

- [1] V. V. Bavula, The algebra of integro-differential operators on an affine line and its modules, *J. Pure Appl. Algebra*, **217** (2013) 495-529. (Arxiv:math.RA: 1011.2997).
- [2] V. V. Bavula, The largest left quotient ring of a ring, Arxiv:math.RA:1101.5107.
- [3] V. V. Bavula, Characterizations of left orders in left Artinian rings, *J. of Algebra and its Appl.*, Arxiv:math.RA:1212.3529.
- [4] V. V. Bavula, New criteria for a ring to have a semisimple left quotient ring, Arxiv:math.RA.
- [5] V. V. Bavula, Left localizable rings and their characterizations.
- [6] C. Faith, Algebra. II. Ring theory. Grundlehren der Mathematischen Wissenschaften, No. 191. Springer-Verlag, Berlin-New York, 1976.
- [7] A. V. Jategaonkar, Localization in Noetherian Rings, Londo Math. Soc. LMS 98, Cambridge Univ. Press, 1986.
- [8] J. C. McConnell and J. C. Robson, Noncommutative Noetherian rings. With the cooperation of L. W. Small. Revised edition. Graduate Studies in Mathematics, 30. American Mathematical Society, Providence, RI, 2001. 636 pp.
- [9] B. Stenström, Rings of Quotients, Springer-Verlag, Berlin, Heidelberg, New York, 1975.

Department of Pure Mathematics
University of Sheffield
Hicks Building
Sheffield S3 7RH
UK
email: v.bavula@sheffield.ac.uk