



HAL
open science

Leftmost derivations of propagating scattered context grammars: a new proof

Tomáš Masopust, Jiří Techet

► **To cite this version:**

Tomáš Masopust, Jiří Techet. Leftmost derivations of propagating scattered context grammars: a new proof. *Discrete Mathematics and Theoretical Computer Science, DMTCS*, 2008, Vol. 10 no. 2 (2), pp.39–46. 10.46298/dmtcs.446 . hal-00972321

HAL Id: hal-00972321

<https://hal.inria.fr/hal-00972321>

Submitted on 3 Apr 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Leftmost Derivations of Propagating Scattered Context Grammars: A New Proof

Tomáš Masopust[†] and Jiří Techet[‡]

Faculty of Information Technology

Brno University of Technology

Božetěchova 2, Brno 61266

Czech Republic

masopust@fit.vutbr.cz, techet@fit.vutbr.cz

received September 17, 2007, revised April 8, 2008, accepted April 11, 2008.

In 1973, V. Virkkunen proved that propagating scattered context grammars which use leftmost derivations are as powerful as context-sensitive grammars. This paper brings a significantly simplified proof of this result.

Keywords: formal languages, propagating scattered context grammars, leftmost derivations, generative power

1 Introduction

Propagating scattered context grammars, introduced in [3], represent an important type of semi-parallel rewriting systems. Since their introduction, however, the exact relationship of the family of languages they generate to the family of context-sensitive languages is unknown. The language family generated by these grammars is included in the family of context-sensitive languages; on the other hand, the question of whether this inclusion is proper represents an open problem in formal language theory. There have been several attempts to modify the definition of propagating scattered context grammars to obtain the family of context-sensitive languages (see [1, 2, 7, 9, 11]). The approach discussed in [11] allows the productions to be applied only in a leftmost way and, thereby, obtain the family of context-sensitive languages generated by these grammars. This result is of some interest as the use of context-free, context-sensitive, and unrestricted productions in a leftmost way in the corresponding grammars of the Chomsky hierarchy does not have any impact on their generative power.

The proof in [11] consists of two parts; first, two preliminary lemmas (Lemma 2 and Lemma 3) are given and then the main result, stated in Theorem 2, is presented as a straightforward corollary of these

[†]Supported by the Czech Ministry of Education under the Research Plan No. MSM 0021630528.

[‡]Supported by the Czech Grant Agency project No. 102/05/H050.

two lemmas. In Lemma 2 it is demonstrated how any sentence of some context-sensitive language can be derived by a propagating scattered context grammar which uses leftmost derivations. Every sentence generated in such a way contains, however, some additional symbols. Lemma 3 shows how these symbols can be removed. Together, the proof consists of six-page-long construction part and not even one-page-long basic idea of the construction which makes it extremely hard to follow. A more formal proof of the correctness of the construction is missing.

This paper aims to present the proof of this result in a much simpler and more readable way. The main difference of our proof lies (1) in the way how the symbols to be rewritten are selected and (2) the way how context-sensitive productions are simulated. Furthermore, the proof is based on a single construction instead of two. All this leads to a significantly simpler and more transparent construction.

2 Preliminaries and definitions

We assume that the reader is familiar with formal language theory (see [10]). For an alphabet V , $|V|$ denotes the cardinality of V . V^* represents the free monoid generated by V . The unit of V^* is denoted by ε . Set $V^+ = V^* - \{\varepsilon\}$. For $w \in V^*$, $|w|$ and $\text{alph}(w)$ denote the length of w and the set of symbols occurring in w , respectively.

A *grammar* is a quadruple $G = (V, T, P, S)$, where V is the *total alphabet*, $T \subset V$ is the set of *terminals*, P is a finite set of *productions* of the form $x \rightarrow y$, where $x \in V^*(V - T)V^*$, $y \in V^*$, and $S \in V - T$ is the *start symbol* of G . If $u = z_1xz_2$, $v = z_1yz_2$, and $x \rightarrow y \in P$, where $z_1, z_2 \in V^*$, then G makes a *derivation step* from u to v according to $x \rightarrow y$, symbolically written as $u \Rightarrow_G v [x \rightarrow y]$ or, simply, $u \Rightarrow_G v$. Let \Rightarrow_G^+ and \Rightarrow_G^* denote the transitive closure of \Rightarrow_G and the reflexive and transitive closure of \Rightarrow_G , respectively. If $S \Rightarrow_G^* w$, where $w \in T^*$, $S \Rightarrow_G^* w$ is said to be a *successful derivation* of G . The *language of G* , denoted by $L(G)$, is defined as $L(G) = \{w \in T^* : S \Rightarrow_G^* w\}$. If each production of G is of the form $xAy \rightarrow xuy$, where $x, y \in V^*$, $A \in V - T$, $u \in V^+$, then G is a *context-sensitive grammar*. The family of context-sensitive languages is denoted by $\mathcal{L}(CS)$. If each production of G is of one of the following forms: $AB \rightarrow CD$, $A \rightarrow BC$, $A \rightarrow a$, where $A, B, C, D \in V - T$, and $a \in T$, then G is a grammar in the *Kuroda normal form*.

Lemma 1 ([4]) *For every context-sensitive grammar there exists an equivalent grammar in the Kuroda normal form.*

A *scattered context grammar* (see [1, 2, 3, 5, 6, 7, 8, 9, 11]) is a quadruple $G = (V, T, P, S)$, where V is the total alphabet, $T \subset V$ is the set of terminals, $S \in V - T$ is the start symbol of G , and P is a finite set of productions such that each production has the form $(A_1, A_2, \dots, A_n) \rightarrow (x_1, x_2, \dots, x_n)$, for some $n \geq 1$, where $A_i \in V - T$, and $x_i \in V^*$, for all $1 \leq i \leq n$. If each production $(A_1, A_2, \dots, A_n) \rightarrow (x_1, x_2, \dots, x_n) \in P$ satisfies $x_i \in V^+$ for all $1 \leq i \leq n$, then G is a *propagating scattered context grammar*. If $(A_1, A_2, \dots, A_n) \rightarrow (x_1, x_2, \dots, x_n) \in P$, $u = u_1A_1u_2A_2 \dots u_nA_nu_{n+1}$, and $v = u_1x_1u_2x_2 \dots u_nx_nu_{n+1}$, where $u_i \in V^*$ for all $1 \leq i \leq n+1$, then G makes a *derivation step* from u to v according to $p = (A_1, A_2, \dots, A_n) \rightarrow (x_1, x_2, \dots, x_n)$, symbolically written as $u \Rightarrow_G v [p]$ or, simply, $u \Rightarrow_G v$. In addition, if $A_i \notin \text{alph}(u_i)$ for all $1 \leq i \leq n$, then the direct derivation is *leftmost*, and we write $u \text{ lm} \Rightarrow_G v [p]$; if $A_i \notin \text{alph}(u_{i+1})$ for all $1 \leq i \leq n$, then the direct derivation is *rightmost*, and we write $u \text{ rm} \Rightarrow_G v [p]$. The language of G , denoted by $L(G)$, is defined as $L(G) = \{w \in T^* : S \Rightarrow_G^* w\}$. A propagating scattered context grammar $G = (V, T, P, S)$ uses *leftmost* or *rightmost derivations* if its language is defined as $L(G, \text{lm}) = \{w \in T^* : S \text{ lm} \Rightarrow_G^* w\}$ or $L(G, \text{rm}) = \{w \in T^* : S \text{ rm} \Rightarrow_G^* w\}$.

$w\}$, respectively. The family of languages generated by propagating scattered context grammars which use leftmost or rightmost derivations is denoted by $\mathcal{L}(PSC, \text{lm})$ or $\mathcal{L}(PSC, \text{rm})$, respectively.

3 Main Results

The following theorem and its proof, which represent the main result of this paper, demonstrate that propagating scattered context grammars which use leftmost derivations are equivalent to context-sensitive grammars.

Theorem 1 $\mathcal{L}(PSC, \text{lm}) = \mathcal{L}(CS)$.

Proof: As propagating scattered context grammars do not contain erasing productions, their derivations can be simulated by linear bounded automata. As a result, $\mathcal{L}(PSC, \text{lm}) \subseteq \mathcal{L}(CS)$. In what follows, we demonstrate that also $\mathcal{L}(CS) \subseteq \mathcal{L}(PSC, \text{lm})$ holds true by demonstrating that for every grammar in the Kuroda normal form there exists an equivalent propagating scattered context grammar which uses leftmost derivations.

Let $G = (V, T, P, S)$ be a grammar in the Kuroda normal form. Set $N_1 = (V - T) \cup \{\bar{a} : a \in T\}$ (suppose that $(V - T) \cap \{\bar{a} : a \in T\} = \emptyset$), $\hat{N}_1 = \{\hat{A} : A \in N_1\}$. Let $n = |N_1|$; then, we denote the elements of N_1 as $\{A_1, A_2, \dots, A_n\}$. Define the homomorphism α from V^* to N_1^* as $\alpha(A) = A$ for each $A \in V - T$, and $\alpha(a) = \bar{a}$ for each $a \in T$. Set $N'_2 = \{A' : A \in V - T\}$, $N_3 = \{\langle ab \rangle : a, b \in V\}$, $N'_4 = \{\langle Aa \rangle' : A \in V - T, a \in V\}$, and

$$\begin{aligned} N_5 = & \{\langle a, 0 \rangle, \langle ab, 0 \rangle : a, b \in V\} \\ & \cup \{\langle a, i, j \rangle : a \in V - T, 1 \leq i \leq 3, 1 \leq j \leq n\} \\ & \cup \{\langle ab, 4 \rangle : a, b \in T\}. \end{aligned}$$

Without loss of generality, assume that the sets $N_1, \hat{N}_1, N'_2, N_3, N'_4, N_5, \{\bar{S}, X\}$, and T are pairwise disjoint. Define the propagating scattered context grammar

$$\bar{G} = (N_1 \cup \hat{N}_1 \cup N'_2 \cup N_3 \cup N'_4 \cup N_5 \cup \{\bar{S}, X\} \cup T, T, \bar{P}, \bar{S}),$$

where \bar{P} is constructed as follows:

1. (a) For each $a \in L(G)$, where $a \in T$, add $(\bar{S}) \rightarrow (a)$ to \bar{P} ;
- (b) For each $S \Rightarrow_G ab$, where $a, b \in V$, add $(\bar{S}) \rightarrow (\langle ab, 0 \rangle X)$ to \bar{P} ;
2. For each $a, b, c \in V$, add
 - (a) $(\langle a, 0 \rangle, \alpha(b)) \rightarrow (\alpha(a), \langle b, 0 \rangle)$,
 - (b) $(\alpha(a), \langle b, 0 \rangle) \rightarrow (\langle a, 0 \rangle, \alpha(b))$,
 - (c) $(\langle a, 0 \rangle, \langle bc \rangle) \rightarrow (\alpha(a), \langle bc, 0 \rangle)$,
 - (d) $(\alpha(a), \langle bc, 0 \rangle) \rightarrow (\langle a, 0 \rangle, \langle bc \rangle)$ to \bar{P} ;
3. For each $A \rightarrow a \in P$ and $b \in V$, add

- (a) $(\langle A, 0 \rangle) \rightarrow (\langle a, 0 \rangle)$,
- (b) $(\langle Ab, 0 \rangle) \rightarrow (\langle ab, 0 \rangle)$,
- (c) $(\langle bA, 0 \rangle) \rightarrow (\langle ba, 0 \rangle)$ to \bar{P} ;
4. For each $A \rightarrow BC \in P$ and $a \in V$, add
- (a) $(\langle A, 0 \rangle) \rightarrow (B\langle C, 0 \rangle)$,
- (b) $(\langle Aa, 0 \rangle) \rightarrow (B\langle Ca, 0 \rangle)$,
- (c) $(\langle aA, 0 \rangle) \rightarrow (\alpha(a)\langle BC, 0 \rangle)$ to \bar{P} ;
5. For each $AB \rightarrow CD \in P$, $a \in V$, $E \in N_3 \cup N'_4$, $F' \in \{B', \langle Ba \rangle'\}$, $1 \leq i \leq n$, and $1 \leq j \leq n - 1$, add
- (a) $(\langle AB, 0 \rangle) \rightarrow (\langle CD, 0 \rangle)$,
- (b) i. $(\langle A, 0 \rangle, B, X) \rightarrow (\langle A, 1, 1 \rangle, B', A_1)$,
- ii. $(\langle A, 0 \rangle, \langle Ba \rangle', X) \rightarrow (\langle A, 1, 1 \rangle, \langle Ba \rangle', A_1)$,
- (c) i. $(\langle A, 1, i \rangle, A_i) \rightarrow (\langle A, 2, i \rangle, \hat{A}_i)$,
- ii. $(\langle A, 2, i \rangle, F', \hat{A}_i) \rightarrow (\langle A, 3, i \rangle, F', A_i)$,
- iii. $(\langle A, 3, j \rangle, E, A_j) \rightarrow (\langle A, 1, j + 1 \rangle, E, A_{j+1})$,
- (d) i. $(\langle A, 3, n \rangle, B', E, A_n) \rightarrow (\langle C, 0 \rangle, D, E, X)$,
- ii. $(\langle A, 3, n \rangle, \langle Ba \rangle', A_n) \rightarrow (\langle C, 0 \rangle, \langle Da \rangle', X)$ to \bar{P} ;
6. For each $a, b, c \in T$, add
- (a) $(\langle ab, 0 \rangle) \rightarrow (\langle ab, 4 \rangle)$,
- (b) $(\bar{c}, \langle ab, 4 \rangle) \rightarrow (c, \langle ab, 4 \rangle)$,
- (c) $(\langle ab, 4 \rangle, X) \rightarrow (a, b)$ to \bar{P} .

In short, productions introduced in (1) initiate the derivation, productions from (2) are used to select the nonterminal to be rewritten, productions from (3), (4), and (5) simulate G 's productions of the form $A \rightarrow a$, $A \rightarrow BC$, and $AB \rightarrow CD$, respectively, and, finally, productions from (6) finish the derivation. In the following paragraphs, we describe the derivation of \bar{G} in greater detail.

Every derivation starts either by a production introduced in (1a) to generate sentences $a \in L(G)$, where $a \in T$, or by a production introduced in (1b) to generate sentences $x \in L(G)$, where $|x| \geq 2$. As \bar{S} does not occur on the right-hand side of any production, productions from (1) are not used during the rest of the derivation.

Consider G 's sentential form $a_1 a_2 \dots a_k$, where $a_1, a_2, \dots, a_k \in V$, for some $k \geq 2$. In \bar{G} , this sentential form corresponds to

$$b_1 b_2 \dots b_{r-1} \langle a_r, 0 \rangle b_{r+1} b_{r+2} \dots b_{k-2} \langle a_{k-1} a_k \rangle X,$$

where $b_i = \alpha(a_i)$ for all $i \in \{1, 2, \dots, r-1, r+1, r+2, \dots, k-2\}$, for some $1 \leq r \leq k-2$, or to

$$b_1 b_2 \dots b_{k-2} \langle a_{k-1} a_k, 0 \rangle X,$$

where $b_i = \alpha(a_i)$ for all $1 \leq i \leq k-2$ (observe that every right-hand side of a production from (1b) represents a sentential form of this kind). To simulate a G 's production, the leftmost nonterminal from its left-hand side has to be selected in the sentential form of \bar{G} . This is done by appending 0 to the symbol to be selected by productions from (2). Specifically, for a symbol $a \in V$, (2a) selects the leftmost symbol a immediately following the currently selected symbol and (2b) selects the leftmost symbol a preceding the currently selected symbol. Productions from (2c) and (2d) are used to select and unselect the penultimate nonterminal in \bar{G} 's sentential form which is composed of two symbols from V . Observe that in this way, any symbol (except for the final X) in every sentential form of \bar{G} can be selected. Further, observe that during a derivation, always one symbol is selected.

After the nonterminal is selected, the use of the G 's production can be simulated. Productions of the form $A \rightarrow a$ are simulated by (3a) for every selected nonterminal a_1, a_2, \dots, a_{k-2} and by (3b), (3c) if the penultimate nonterminal (which contains a_{k-1}, a_k) of the \bar{G} 's sentential form is selected. Analogously, productions of the form $A \rightarrow BC$ are simulated by productions from (4).

Productions from (5a) are used to simulate an application of productions of the form $AB \rightarrow CD$ within the penultimate nonterminal of \bar{G} 's sentential form. In what follows, we demonstrate how productions from (5b), (5c), and (5d) are used if this production is simulated within $a_1 a_2 \dots a_{k-2}$. Suppose that the sentential form in \bar{G} is of the form

$$b_1 b_2 \dots b_{r-1} \langle a_r, 0 \rangle b_{r+1} b_{r+2} \dots b_{k-2} \langle a_{k-1} a_k \rangle X$$

and we simulate the application of $a_r a_{r+1} \rightarrow c_r c_{r+1} \in P$. Recall that $N_1 = \{A_1, A_2, \dots, A_n\}$ denotes the set of all symbols which may appear in $b_{r+1} b_{r+2} \dots b_{k-2}$. First, to select $b_{r+1} = \alpha(a_{r+1})$, the production

$$\langle \langle a_r, 0 \rangle, b_{r+1}, X \rangle \rightarrow \langle \langle a_r, 1, 1 \rangle, b'_{r+1}, A_1 \rangle$$

from (5(b)i) is applied in a successful derivation, so

$$\begin{aligned} & b_1 b_2 \dots b_{r-1} \langle a_r, 0 \rangle b_{r+1} b_{r+2} \dots b_{k-2} \langle a_{k-1} a_k \rangle X \\ \text{lm} \Rightarrow_{\bar{G}} & b_1 b_2 \dots b_{r-1} \langle a_r, 1, 1 \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_1. \end{aligned}$$

Observe that if b_{r+1} does not immediately follow $\langle a_r, 0 \rangle$, the leftmost $b \in \text{alph}(b_{r+2} b_{r+3} \dots b_{k-2})$ satisfying $b = b_{r+1}$ is selected by the production from (5(b)i). The purpose of productions from (5c) is to verify that the nonterminal immediately following $\langle a_r, 0 \rangle$ has been selected. First, the production

$$\langle \langle a_r, 1, 1 \rangle, A_1 \rangle \rightarrow \langle \langle a_r, 2, 1 \rangle, \hat{A}_1 \rangle$$

from (5(c)i) is applied to tag the first A_1 following $\langle a_r, 1, 1 \rangle$, so

$$\begin{aligned} & b_1 b_2 \dots b_{r-1} \langle a_r, 1, 1 \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_1 \\ \text{lm} \Rightarrow_{\bar{G}} & b_1 b_2 \dots b_{r-1} \langle a_r, 2, 1 \rangle b'_{r+1} y_1 \langle a_{k-1} a_k \rangle d_1, \end{aligned}$$

where either

$$y_1 = b_{r+2} b_{r+3} \dots b_{m-1} \hat{A}_1 b_{m+1} b_{m+2} \dots b_{k-2}, \quad d_1 = A_1,$$

satisfying $A_1 \notin \text{alph}(b_{r+2} b_{r+3} \dots b_{m-1})$, for some $1 \leq m \leq k-2$, or

$$y_1 = b_{r+2} b_{r+3} \dots b_{k-2}, \quad d_1 = \hat{A}_1,$$

satisfying $A_1 \notin \text{alph}(y_1)$. Then, the production

$$\langle \langle a_r, 2, 1 \rangle, b'_{r+1}, \hat{A}_1 \rangle \rightarrow \langle \langle a_r, 3, 1 \rangle, b'_{r+1}, A_1 \rangle$$

from (5(c)ii) is applied to untag the first symbol \hat{A}_1 following b'_{r+1} , so

$$\begin{aligned} & b_1 b_2 \dots b_{r-1} \langle a_r, 2, 1 \rangle b'_{r+1} y_1 \langle a_{k-1} a_k \rangle d_1, \\ \text{lm} \Rightarrow_{\bar{G}} & b_1 b_2 \dots b_{r-1} \langle a_r, 3, 1 \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_1. \end{aligned}$$

This means that if A_1 occurs between $\langle a_r, 2, 1 \rangle$ and b'_{r+1} , it is tagged by the production from (5(c)i) but it cannot be untagged by any production from (5(c)ii), so the derivation is blocked. Finally, the production

$$\langle \langle a_r, 3, 1 \rangle, \langle a_{k-1} a_k \rangle, A_1 \rangle \rightarrow \langle \langle a_r, 1, 2 \rangle, \langle a_{k-1} a_k \rangle, A_2 \rangle$$

from (5(c)iii) is applied, so

$$\begin{aligned} & b_1 b_2 \dots b_{r-1} \langle a_r, 3, 1 \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_1 \\ \text{lm} \Rightarrow_{\bar{G}} & b_1 b_2 \dots b_{r-1} \langle a_r, 1, 2 \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_2, \end{aligned}$$

and the same verification continues for A_2 . This verification proceeds for all symbols from N_1 so this part of the derivation can be expressed as

$$\begin{array}{ccc} u_1 & \text{lm} \Rightarrow_{\bar{G}} v_1 [p_{11}] & \text{lm} \Rightarrow_{\bar{G}} w_1 [p_{12}] \\ \text{lm} \Rightarrow_{\bar{G}} u_2 [p_{13}] & \text{lm} \Rightarrow_{\bar{G}} v_2 [p_{21}] & \text{lm} \Rightarrow_{\bar{G}} w_2 [p_{22}] \\ \vdots & & \\ \text{lm} \Rightarrow_{\bar{G}} u_n [p_{(n-1)3}] & \text{lm} \Rightarrow_{\bar{G}} v_n [p_{n1}] & \text{lm} \Rightarrow_{\bar{G}} w_n [p_{n2}] \end{array}$$

with

$$\begin{aligned} u_i &= b_1 b_2 \dots b_{r-1} \langle a_r, 1, i \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_i, \\ v_i &= b_1 b_2 \dots b_{r-1} \langle a_r, 2, i \rangle b'_{r+1} y_i \langle a_{k-1} a_k \rangle d_i, \\ w_i &= b_1 b_2 \dots b_{r-1} \langle a_r, 3, i \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_i, \end{aligned}$$

where p_{i1} , p_{i2} , and p_{i3} are productions from (5(c)i), (5(c)ii), and (5(c)iii), respectively, for all $1 \leq i \leq n$, $1 \leq j \leq n-1$, and either

$$y_i = b_{r+2} b_{r+3} \dots b_{i_{m-1}} \hat{A}_i b_{i_{m+1}} b_{i_{m+2}} \dots b_{k-2}, \quad d_i = A_i,$$

satisfying $A_i \notin \text{alph}(b_{r+2} b_{r+3} \dots b_{i_{m-1}})$, for some $1 \leq i_m \leq k-2$, or

$$y_i = b_{r+2} b_{r+3} \dots b_{k-2}, \quad d_i = \hat{A}_i,$$

satisfying $A_i \notin \text{alph}(y_i)$. After the verification is finished, the application of $a_r a_{r+1} \rightarrow c_r c_{r+1} \in P$ is simulated by

$$\langle \langle a_r, 3, n \rangle, b'_{r+1}, \langle a_{k-1} a_k \rangle, A_n \rangle \rightarrow \langle \langle c_r, 0 \rangle, c_{r+1}, \langle a_{k-1} a_k \rangle, X \rangle$$

from (5(d)i), so

$$\begin{aligned} & b_1 b_2 \dots b_{r-1} \langle a_r, 3, n \rangle b'_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle A_n \\ \text{lm} \Rightarrow_{\bar{G}} & b_1 b_2 \dots b_{r-1} \langle c_r, 0 \rangle c_{r+1} b_{r+2} b_{r+3} \dots b_{k-2} \langle a_{k-1} a_k \rangle X. \end{aligned}$$

Observe that in order to simulate a production of the form $AB \rightarrow CD$ within $a_{k-2}a_{k-1}$, productions from (5(b)ii) and (5(d)ii) have to be used instead of productions from (5(b)i) and (5(d)i) in the simulation described above. The details are left to the reader.

Finally, consider a G 's sentence $a_1a_2 \dots a_k \in T^+$. This corresponds to

$$\bar{a}_1\bar{a}_2 \dots \bar{a}_{r-1}\langle a_r, 0 \rangle \bar{a}_{r+1}\bar{a}_{r+2} \dots \bar{a}_{k-2}\langle a_{k-1}a_k \rangle X,$$

or

$$\bar{a}_1\bar{a}_2 \dots \bar{a}_{k-2}\langle a_{k-1}a_k, 0 \rangle X$$

in \bar{G} after finishing the simulation. To enter the final phase in \bar{G} , we need the sentential form to be in the second above described form. This can be achieved by applying a production from (2c) to the first sentential form. The rest of the derivation can be expressed as

$$\begin{array}{l} \bar{a}_1\bar{a}_2 \dots \bar{a}_{k-2}\langle a_{k-1}a_k, 0 \rangle X \\ \text{lm} \Rightarrow_{\bar{G}} \bar{a}_1\bar{a}_2 \dots \bar{a}_{k-2}\langle a_{k-1}a_k, 4 \rangle X \quad [p_{6a}] \\ \text{lm} \Rightarrow_{\bar{G}}^{k-2} a_1a_2 \dots a_{k-2}\langle a_{k-1}a_k, 4 \rangle X \quad [\Xi_{6b}] \\ \text{lm} \Rightarrow_{\bar{G}} a_1a_2 \dots a_{k-2}a_{k-1}a_k \quad [p_{6c}], \end{array}$$

where p_{6a} and p_{6c} are productions introduced in steps (6a) and (6c), respectively, and Ξ_{6b} is a sequence of $k-2$ productions from (6b). As a result, $x \in L(\bar{G}, \text{lm})$ if and only if $x \in L(G)$. Therefore, $\mathcal{L}(CS) \subseteq \mathcal{L}(PSC, \text{lm})$.

As $\mathcal{L}(PSC, \text{lm}) \subseteq \mathcal{L}(CS)$ and $\mathcal{L}(CS) \subseteq \mathcal{L}(PSC, \text{lm})$, we obtain $\mathcal{L}(PSC, \text{lm}) = \mathcal{L}(CS)$, so the theorem holds. \square

Next, we state the following corollary.

Corollary 1 $\mathcal{L}(PSC, \text{rm}) = \mathcal{L}(CS)$.

Proof: This corollary can be proved by a straightforward modification of the proof of Theorem 1 and its proof is, therefore, left to the reader. \square

References

- [1] H. Fernau. Scattered context grammars with regulation. *Annals of Bucharest University, Mathematics-Informatics Series*, 45(1):41–49, 1996.
- [2] J. Gonczarowski and M. K. Warmuth. Scattered versus context-sensitive rewriting. *Acta Informatica*, 27:81–95, 1989.
- [3] S. Greibach and J. Hopcroft. Scattered context grammars. *Journal of Computer and System Sciences*, 3:233–247, 1969.
- [4] S. Y. Kuroda. Classes of languages and linear-bounded automata. *Information and Control*, 7(2):207–223, 1964.
- [5] T. Masopust. Scattered context grammars can generate the powers of 2. In *Proceedings of the 13th Conference Student EEICT 2007, Volume 4*, pages 401–404, Brno, 2007. Faculty of Electrical Engineering and Communication BUT.
- [6] A. Meduna and J. Techet. Canonical scattered context generators of sentences with their parses. *Theoretical Computer Science*, 389:73–81, 2007.
- [7] A. Meduna and J. Techet. Maximal and minimal scattered context rewriting. In *FCT 2007 Proceedings*, volume 2007, pages 412–423. Springer Verlag, 2007.
- [8] A. Meduna and J. Techet. Reduction of scattered context generators of sentences preceded by their leftmost parses. In *Proceedings of 9th International Workshop on Descriptive Complexity of Formal Systems*, pages 178–185. University of Pavol Jozef Šafárik, 2007.
- [9] D. Milgram and A. Rosenfeld. A note on scattered context grammars. *Information Processing Letters*, 1:47–50, 1971.
- [10] A. Salomaa. *Formal Languages*. Academic Press, New York, 1973.
- [11] V. Virkkunen. On scattered context grammars. *Acta Universitatis Ouluensis, Series A, Mathematica* 6:75–82, 1973.