

## LEMMA ON LOGARITHMIC DERIVATIVES AND HOLOMORPHIC CURVES IN ALGEBRAIC VARIETIES<sup>1)</sup>

JUNJIRO NOGUCHI

Nevanlinna's lemma on logarithmic derivatives played an essential role in the proof of the second main theorem for meromorphic functions on the complex plane  $C$  (cf., e.g., [17]). In [19, Lemma 2.3] it was generalized for entire holomorphic curves  $f: C \rightarrow M$  in a compact complex manifold  $M$  (Lemma 2.3 in [19] is still valid for non-Kähler  $M$ ). Here we call, in general, a holomorphic mapping from a domain of  $C$  or a Riemann surface into  $M$  a holomorphic curve in  $M$ , and sometimes use it in the sense of its image if no confusion occurs. Applying the above generalized lemma on logarithmic derivatives to holomorphic curves  $f: C \rightarrow V$  in a complex projective algebraic smooth variety  $V$  and making use of Ochiai [22, Theorem A], we had an inequality of the second main theorem type for  $f$  and divisors on  $V$  (see [19, Main Theorem] and [20]). Other generalizations of Nevanlinna's lemma on logarithmic derivatives were obtained by Nevanlinna [16], Griffiths-King [10, § 9] and Vitter [23].

In this paper we first deal with holomorphic curves  $f: \Delta^* \rightarrow M$  from the punctured disc  $\Delta^* = \{|z| \geq 1\}$  with center at the infinity  $\infty$  of the Riemann sphere into a compact Kähler manifold  $M$ . Our first aim is to prove the following lemma on logarithmic derivatives which is a generalization of Nevanlinna [16, III, p. 370] and will play a crucial role in §§ 3 and 4 (see § 1 as to the notation):

**MAIN LEMMA (2.2).** *Let  $f: \Delta^* \rightarrow M$  be a holomorphic curve in  $M$ ,  $\omega \in H^0(M, \mathfrak{X}_M^1)$  a  $d$ -closed meromorphic 1-form with logarithmic poles and put  $f^*\omega = \zeta(z)dz$ . Then we have*

$$m(r, \zeta) \leq O(\log^+ T_f(r)) + O(\log r)$$

as  $r \rightarrow \infty$  except for  $r \in E$ , where  $E$  is a subset of  $[1, \infty)$  with finite linear

---

Received January 24, 1980.

1) Work supported in part by the Sakkokai Foundation.

*measure.*

The difficulty of the present case comes from the fact that the domain  $\Delta^*$  is not simply connected. In the proof we shall apply the negative curvature method introduced by Griffiths-King [10, Propositions (6.9) and (9.3)] as in Vitter [23].

In § 3 we shall be concerned with the value distribution of holomorphic curves  $f: \Delta^* \rightarrow V$  in a complex projective algebraic smooth variety  $V$ . Let  $D$  be an effective reduced divisor on  $V$ . Combining Main Lemma (2.2) with Ochiai [22, Theorem A] as in [19, § 3] and [20], we shall obtain an inequality of the second main theorem type

$$(3.2) \quad KT_f(r) \leq N(r, \text{Supp}(f^*D)) + S(r),$$

where  $K$  is a positive constant independent of  $f$  and  $S(r)$  is a small term such as

$$S(r) \leq O(\log^+ T_f(r)) + O(\log r)$$

as  $r \rightarrow \infty$  outside a set of  $r$  with finite linear measure (see Theorem (3.1)). As a corollary, we shall see that an inequality similar to (3.2) holds for a holomorphic curve from a compact Riemann surface minus a finite number of points into  $V$  (Corollary (3.3)).

In § 4 we shall study the extension problem of big Picard type for holomorphic curves  $f: \Delta^* \rightarrow X$  in an algebraic subvariety  $X$  of general type in a quasi-Abelian variety  $A$  (cf. § 4). Let  $W$  be the union of subvarieties of  $X$  which are translations of non-trivial closed algebraic subgroups of  $A$ . Then  $W$  is a proper algebraic subvariety of  $X$  such that each irreducible component of  $W$  is foliated by translations of a non-trivial closed algebraic subgroup of  $A$  (see Lemma (4.1) whose proof is essentially due to Kawamata [13]). Using Lemma (4.4) due to M. Green by which he completed Ochiai's work [22] on Bloch's conjecture [2], and applying Main Lemma (2.2), we shall prove the following extension theorem of big Picard type:

**THEOREM (4.5).** *Any holomorphic curve  $f: \Delta^* \rightarrow X$  has a holomorphic extension  $\tilde{f}: \Delta = \Delta^* \cup \{\infty\} \rightarrow \bar{X}$  unless  $f(\Delta^*) \subset W$ , where  $\bar{X}$  is a completion of  $X$ .*

As a corollary of Theorem (4.5) we will see that any holomorphic mapping  $f: N - S \rightarrow X$  from a complex manifold  $N$  minus a thin analytic set  $S$  into  $X$  extends meromorphically over  $N$  unless  $f(N - S) \subset W$

(Corollary (4.7)). Fujimoto ([3], [5]) and Green ([8]) obtained extension theorems of big Picard type for holomorphic mappings into projective space omitting hyperplanes in general position or intersecting them with positive defects (cf. also [4] and [7]). We will discuss the relationship between our results and those of Fujimoto and Green.

§1. Preliminaries

We set

$$\begin{aligned} \Delta^* &= \{z \in \mathbb{C}; |z| \geq 1\}, & \Delta^*(r) &= \{1 \leq |z| < r\}, \\ \Gamma(r) &= \{|z| = r\}, & d &= \partial + \bar{\partial}, & d^c &= \frac{i}{4\pi}(\bar{\partial} - \partial). \end{aligned}$$

In this paper we assume that functions on  $\Delta^*$  and mappings from  $\Delta^*$  are defined in neighborhoods of  $\Delta^*$  in  $\mathbb{C}$ . Let  $\xi$  be a function on  $\Delta^*$  satisfying

- (i)  $\xi$  is differentiable outside a discrete set of points,
- (ii)  $\xi$  is locally written as a difference of two subharmonic functions.

Then we have

$$(1.1) \quad \int_1^r \frac{dt}{t} \int_{\Delta^*(t)} dd^c \xi = \frac{1}{4\pi} \int_{\Gamma(r)} \xi(re^{i\theta}) d\theta - \frac{1}{4\pi} \int_{\Gamma(1)} \xi(e^{i\theta}) d\theta - (\log r) \int_{\Gamma(1)} d^c \xi,$$

where  $dd^c \xi$  is taken in the sense of currents (cf., e.g., [10]). Let  $F$  be a multiplicative meromorphic function on  $\Delta^*$ , i.e.,  $F$  is a many-valued meromorphic function such that the modulus  $|F|$  is one-valued. We set

$$m(r, F) = \frac{1}{2\pi} \int_{\Gamma(r)} \log^+ |F(re^{i\theta})| d\theta,$$

where  $\log^+ |F| = \max\{0, \log |F|\}$ . Let  $D = \sum_{i=1}^{\infty} \nu_i a_i$  be a divisor with integral coefficients  $\nu_i \in \mathbb{Z}$  on  $\Delta^*$  and set

$$\begin{aligned} n(t, D) &= \sum_{1 \leq |a_i| < t} \nu_i, \\ N(r, D) &= \int_1^r \frac{n(t, D)}{t} dt. \end{aligned}$$

Since  $|F|$  is one-valued, the divisor  $(F)$  determined by  $F$  is defined on  $\Delta^*$  and so is the divisor  $(F)_0$  (resp.  $(F)_\infty$ ) of zeros (resp. poles) of  $F$ . We put

$$(1.2) \quad T(r, F) = N(r, (F)_\infty) + m(r, F).$$

Applying (1.1) to  $\xi = \log |F|^2$ , we get

$$(1.3) \quad T\left(r, \frac{1}{F}\right) = T(r, F) - \frac{1}{2\pi} \int_{r(1)} \log |F| d\theta - (\log r) \int_{r(1)} d^c \log |F|^2$$

(cf. [16, I, p. 369]).

Let  $M$  be a compact Kähler manifold and  $\Omega$  a  $(1, 1)$ -form on  $M$ . We set

$$T_f(r, \Omega) = \int_1^r \frac{dt}{t} \int_{A^*(t)} f^* \Omega$$

for a holomorphic curve  $f: A^* \rightarrow M$ . Let  $D$  be an effective divisor on  $M$  and  $f: A^* \rightarrow M$  a holomorphic curve such that  $f(A^*)$  is not contained in the support  $\text{Supp}(D)$  of  $D$ . We take a metric  $\|\cdot\|$  in the line bundle  $[D]$  determined by  $D$  and denote by  $\Omega_0$  the curvature form of the metric. Letting  $\sigma \in H^0(M, [D])$  be a global holomorphic section of  $[D]$  such that the divisor  $(\sigma)$  determined by  $\sigma$  equals  $D$  and  $\|\sigma\| \leq 1$ , we put

$$m_f(r, D) = \frac{1}{2\pi} \int_{r(1)} \log \frac{1}{\|\sigma \circ f\|} d\theta.$$

Applying (1.1) to  $\xi = f^* \log \|\sigma\|^2$ , we obtain

$$(1.4) \quad \begin{aligned} T_f(r, \Omega_0) &= N(r, f^* D) + m_f(r, D) - m_f(1, D) \\ &\quad + (\log r) \int_{r(1)} d^c \log \|\sigma \circ f\|^2, \end{aligned}$$

where  $f^* D$  denotes the pull-backed divisor of  $D$  by  $f$  (cf. [10]). Let  $\mathfrak{M}_M^*$  be the sheaf of germs of meromorphic functions which do not identically vanish, and define a sheaf  $\mathfrak{A}_M^1$  by

$$(1.5) \quad \begin{array}{ccccccc} 0 & \longrightarrow & C^* & \longrightarrow & \mathfrak{M}_M^* & \xrightarrow{d \log} & \mathfrak{A}_M^1 & \longrightarrow & 0, \\ & & & & \omega & & \omega & & \\ & & & & & & \gamma & \mapsto & d \log \gamma \end{array}$$

where  $C^*$  denotes the multiplicative group of non-zero complex numbers (cf. [19, § 1(b)]). Let  $\omega \in H^0(M, \mathfrak{A}_M^1)$ . Then we have the residue  $\text{Res}(\omega)$  which is a divisor homologous to zero such that the line bundle  $[\text{Res}(\omega)]$  equals  $\delta\omega$ , where  $\delta: H^0(M, \mathfrak{A}_M^1) \rightarrow H^1(M, C^*)$  is the coboundary operator associated with (1.5) (cf. [19, § 1(b)]). By Weil [24, p. 101] there is a multiplicative meromorphic function  $\theta$  on  $M$  such that the divisor  $(\theta)$  equals  $\text{Res}(\omega)$ . Since  $d \log \theta \in H^0(M, \mathfrak{A}_M^1)$  and  $\omega - d \log \theta$  is holomorphic every-

where on  $M$ , we have the decomposition

$$(1.6) \quad \omega = d \log \theta + \omega_1 ,$$

where  $\omega_1$  is a holomorphic 1-form on  $M$ .

**§ 2. Lemma on logarithmic derivatives**

Let  $f: \Delta^* \rightarrow M$  be a holomorphic curve in a compact Kähler manifold  $M$  with Kähler metric  $h$  and the associated form  $\Omega$ , and set

$$T_f(r) = T_f(r, \Omega) .$$

Let  $\omega \in H^0(M, \mathfrak{A}_M^1)$  and  $\omega = d \log \theta + \omega_1$  be the decomposition as (1.6). We set

$$\text{Res}^+(\omega) = (\theta)_0 , \quad \text{Res}^-(\omega) = (\theta)_\infty .$$

Then by [24, p. 101] there is respectively a metric  $\|\cdot\|$  in each of  $[\text{Res}^+(\omega)]$  and  $[\text{Res}^-(\omega)]$  such that both metrics have the same curvature form  $\Omega_0$ ; furthermore there are sections  $\sigma_1 \in H^0(M, [\text{Res}^-(\omega)])$  and  $\sigma_2 \in H^0(M, [\text{Res}^+(\omega)])$  such that  $(\sigma_1) = \text{Res}^-(\omega)$ ,  $(\sigma_2) = \text{Res}^+(\omega)$ ,  $\|\sigma_i\| \leq 1$  and

$$(2.1) \quad |\theta| = \frac{\|\sigma_2\|}{\|\sigma_1\|} .$$

We put  $f^*\omega = \zeta(z)dz$ .

**MAIN LEMMA (2.2).** *Let the notation be as above. Assume that  $\text{Supp}(\text{Res}(\omega)) \not\supset f(\Delta^*)$ . Then*

$$(2.3) \quad m(r, \zeta) \leq 18 \log^+ T_f(r) + O(\log r)$$

for  $r \geq 1$  outside a set of  $r$  with finite linear measure.

*Proof.* Set  $f^*d \log \theta = \zeta_0 dz$  and  $f^*\omega_1 = \zeta_1 dz$ . Then we have

$$(2.4) \quad m(r, \zeta) \leq m(r, \zeta_0) + m(r, \zeta_1) + \log 2 .$$

We first estimate the term  $m(r, \zeta_1)$ . Take a positive constant  $C_1$  so that

$$|\omega_1(v)|^2 \leq C_1 h(v, v)$$

for every holomorphic tangent vector  $v \in T(M)$ . Setting  $f^*\Omega = s(z)(i/2) dz \wedge d\bar{z}$ , we get

$$(2.5) \quad |\zeta_1(z)|^2 \leq C_1 s(z) ,$$

so that

$$(2.6) \quad \begin{aligned} m(r, \zeta_1) &\leq \frac{1}{4\pi} \int_{r(r)} \log(1 + |\zeta_1|^2) d\theta \leq \frac{1}{2} \log \left( 1 + \frac{C_1}{2\pi} \int_{r(r)} s d\theta \right) \\ &\leq \frac{1}{2} \log \left( 1 + \frac{C_1}{2\pi r} \frac{d}{dr} \int_{d^*(r)} f^* \Omega \right). \end{aligned}$$

Since  $\int_{d^*(r)} f^* \Omega$  is a monotone increasing function in  $r \geq 1$ , the inequality

$$\frac{d}{dr} \int_{d^*(r)} f^* \Omega \leq \left( \int_{d^*(r)} f^* \Omega \right)^2$$

holds for  $r \geq 1$  outside a set  $E_1$  of  $r$  with finite linear measure. Combining this with (2.6), we have

$$(2.7) \quad m(r, \zeta_1) \leq \frac{1}{2} \log \left( 1 + \frac{C_1}{2\pi r} \left( \int_{d^*(r)} f^* \Omega \right)^2 \right)$$

for  $r \notin E_1$ ; moreover we have

$$(2.8) \quad \int_{d^*(r)} f^* \Omega = r \frac{d}{dr} \int_1^r \frac{dt}{t} \int_{d^*(t)} f^* \Omega = r \frac{d}{dr} T_f(r) \leq r(T_f(r))^2$$

for  $r \notin E_2$ , where  $E_2$  is a set similar to  $E_1$ . It follows from (2.7) and (2.8) that

$$(2.9) \quad m(r, \zeta_1) \leq 2 \log^+ T_f(r) + \frac{1}{2} \log r + \frac{1}{2} \log^+ \frac{C_1}{2\pi} + \frac{1}{2} \log 2$$

for  $r \notin E_1 \cup E_2$ .

Now we estimate the term  $m(r, \zeta_0)$  in (2.4). Set  $F = f^* \theta$ . Then  $F$  is a multiplicative meromorphic function on  $d^*$  and by (2.1),  $|F| = \|\sigma_2 \circ f\| / \|\sigma_1 \circ f\|$ , so that

$$m(r, F) \leq \frac{1}{2\pi} \int_{r(r)} \log \frac{1}{\|\sigma_1 \circ f\|} d\theta = m_f(r, \text{Res}^-(\omega)).$$

On the other hand,  $N(r, (F)_\infty) \leq N(r, f^* \text{Res}^-(\omega))$ . Thus we see, taking into account (1.4), that

$$(2.10) \quad T(r, F) \leq T_f(r, \Omega_0) + C_2 \log r + C_3,$$

where  $C_2$  and  $C_3$  are some non-negative constants. Letting  $C_4$  be a positive constant such that  $\Omega_0 \leq C_4 \Omega$ , we have

$$(2.11) \quad T_f(r, \Omega_0) \leq C_4 T_f(r).$$

We complete the proof by combining (2.9) with (2.10), (2.11) and the following one variable lemma.

LEMMA (2.12). *Let  $G$  be a multiplicative meromorphic function on  $\Delta^*$ . Then the inequality*

$$m(r, G'/G) \leq 16 \log^+ T(r, G) + O(\log r)$$

holds for  $r \geq 1$  outside a set  $E$  of  $r$  with finite linear measure.

*Proof.* Let  $w$  be an inhomogeneous coordinate of the 1-dimensional complex projective space  $P^1$ . Then the standard Kähler form  $\psi_0$  on  $P^1$  is written as

$$\psi_0 = \frac{1}{(1 + |w|^2)^2} \frac{i}{2\pi} dw \wedge d\bar{w}.$$

By Griffiths-King [10, Proposition (6.9)] we see that the singular form

$$\Psi = \frac{a_0(|w| + |w|^{-1})^{2+2\epsilon}}{(\log b_0(1 + |w|^2))^2 (\log b_0(1 + |w|^{-2}))^2} \psi_0$$

satisfies

$$(2.13) \quad \text{Ric } \Psi \geq (|w| + |w|^{-1})^{-2\epsilon} \Psi$$

for suitably chosen positive constants  $a_0$ ,  $b_0$  and  $\epsilon$  ( $\epsilon < 1$ ). Since  $\Psi$  is invariant by transformations,  $w \rightarrow e^{i\theta} w$ , with real  $\theta \in \mathbf{R}$  and  $G$  is multiplicative, the pull-backed form  $G^*\Psi$  of  $\Psi$  by  $G$  is well-defined. We set

$$(2.14) \quad \left\{ \begin{array}{l} g = \frac{G'}{G}, \\ G^*\Psi = \xi \frac{i}{2\pi} dz \wedge d\bar{z} = \frac{a_0(|G| + |G|^{-1})^{2\epsilon}}{(\log b_0(1 + |G|^2))^2 (\log b_0(1 + |G|^{-2}))^2} \\ \quad \times |g|^2 \frac{i}{2\pi} dz \wedge d\bar{z}. \end{array} \right.$$

Then by (2.13) we have

$$(2.15) \quad G^*\text{Ric } \Psi = dd^c \log \xi \geq (|G| + |G|^{-1})^{-2\epsilon} \xi \frac{i}{2\pi} dz \wedge d\bar{z}.$$

Furthermore, taking  $dd^c \log \xi$  in the sense of currents, we get

$$(2.16) \quad dd^c \log \xi = G^* \text{Ric } \Psi - \varepsilon((G)_0 + (G)_\infty) + (g)_0 - (g)_\infty .$$

Noting that  $(g)_\infty = \text{Supp}((G)_0 + (G)_\infty) \leq (G)_0 + (G)_\infty$ , we deduce from (2.15) and (2.16) that

$$(2.17) \quad (|G| + |G|^{-1})^{-2\varepsilon} \xi \frac{i}{2\pi} dz \wedge d\bar{z} \leq (1 + \varepsilon)((G)_0 + (G)_\infty) + dd^c \log \xi .$$

We infer from (1.1) and (2.17) that

$$(2.18) \quad \int_1^r \frac{dt}{t} \int_{A^*(t)} \frac{\xi}{(|G| + |G|^{-1})^{2\varepsilon}} \frac{i}{2\pi} dz \wedge d\bar{z} \leq (1 + \varepsilon)(N(r, (G)_0) + N(r, (G)_\infty)) \\ + \frac{1}{4\pi} \int_{\Gamma(r)} \log \xi d\theta - (\log r) \int_{\Gamma(1)} d^c \log \xi - \frac{1}{4\pi} \int_{\Gamma(1)} \log \xi d\theta .$$

We have by the definition of  $\xi$  in (2.14)

$$(2.19) \quad \frac{1}{4\pi} \int_{\Gamma(r)} \log \xi d\theta \leq m(r, g) + \varepsilon \left( m(r, G) + m\left(r, \frac{1}{G}\right) \right) \\ + \log^+ a_0 + \log^+(\log b_0)^{-2} + \varepsilon \log 2 .$$

We put

$$(2.20) \quad \begin{cases} A(t) = \int_{A^*(t)} \frac{\xi}{(|G| + |G|^{-1})^{2\varepsilon}} \frac{i}{2\pi} dz \wedge d\bar{z} , \\ B(r) = \int_1^r \frac{A(t)}{t} dt . \end{cases}$$

Then inequalities (2.18), (2.19), (1.3) and  $\varepsilon < 1$  yield

$$(2.21) \quad B(r) \leq m(r, g) + 4T(r, G) + O(\log r) + O(1) .$$

Let us compute  $m(r, g)$ :

$$(2.22) \quad m(r, g) = \frac{1}{4\pi} \int_{\Gamma(r)} \log^+ \left( \xi (|G| + |G|^{-1})^{-2\varepsilon} \frac{1}{a_0} \right. \\ \left. \times (\log b_0 (1 + |G|^2))^2 (\log b_0 (1 + |G|^{-2}))^2 \right) d\theta \\ \leq \frac{1}{4\pi} \int_{\Gamma(r)} \log(1 + \xi (|G| + |G|^{-1})^{-2\varepsilon}) d\theta \\ + \frac{1}{2\pi} \int_{\Gamma(r)} \log(1 + \log^+ b_0 + 2 \log^+ |G|) d\theta \\ + \frac{1}{2\pi} \int_{\Gamma(r)} \log\left(1 + \log^+ b_0 + 2 \log^+ \frac{1}{|G|}\right) d\theta + \log^+ \frac{1}{a_0}$$



$$\begin{aligned} &\leq \frac{1}{2} \log \left( 1 + \frac{1}{2\pi} \int_{r(r)} \xi(|G| + |G|^{-1})^{-2\epsilon} d\theta \right) \\ &\quad + \log(1 + \log^+ b_0 + 2m(r, G)) \\ &\quad + \log \left( 1 + \log^+ b_0 + 2m \left( r, \frac{1}{G} \right) \right) + \log^+ \frac{1}{a_0} \\ &\hspace{15em} \text{(by the concavity of "log")} \\ &\leq \frac{1}{2} \log \left( 1 + \frac{1}{2r} \frac{d}{dr} A(r) \right) + 2 \log^+ T(r, G) + O(\log r) + O(1). \end{aligned}$$

Since  $A(r)$  and  $B(r)$  are monotone increasing, we see that the inequalities

$$(2.23) \quad \left\{ \begin{aligned} \frac{d}{dr} A(r) &\leq (A(r))^2, \\ \frac{d}{dr} B(r) &\leq (B(r))^2 \end{aligned} \right.$$

hold for  $r \geq 1$  outside a set  $E$  of  $r$  with finite linear measure. Using the identity,  $dB(r)/dr = A(r)/r$ , and combining (2.22) with (2.21) and (2.23), we have

$$\begin{aligned} m(r, g) &\leq \frac{1}{2} \log \left( 1 + \frac{1}{2} r(B(r))^4 \right) + 2 \log^+ T(r, G) + O(\log r) + O(1) \\ &\leq 2 \log^+ m(r, g) + 4 \log^+ T(r, G) + O(\log r) + O(1) \end{aligned}$$

for  $r \notin E$ . Note that  $2 \log^+ m(r, g) \leq 2m(r, g)/e$  and  $1 - 2/e > 1/4$ . Hence we infer that

$$(2.24) \quad m(r, g) \leq 16 \log^+ T(r, G) + O(\log r) + O(1)$$

for  $r \notin E$ . This completes the proof.

*Remark 1.* In the above proof we used the metric form (cf. (2.14)) due to Griffiths-King [10, Proposition (6.9)] as in Vitter [23], whose curvature behaves nicely. If we use the following metric form due to Grauert-Reckziegel [6] which is simpler than (2.14)

$$\Phi = (1 + |G|^{2\epsilon}) |G|^{2\epsilon} |g|^2 \frac{i}{2\pi} dz \wedge d\bar{z}$$

with any  $\epsilon > 0$ , we have

$$\text{Ric } \Phi = \epsilon^2 (|G|^\epsilon + |G|^{-\epsilon})^{-2} |g|^2 \frac{i}{2\pi} dz \wedge d\bar{z}$$

and obtain the following estimate:

$$(2.25) \quad m(r, g) \leq 8\varepsilon T(r, G) + 4 \log^+ \frac{1}{\varepsilon} + 8 \log^+ T(r, G) \\ + (\varepsilon C_1 + 2) \log r + \varepsilon C_2 + C_3$$

for  $r \geq 1$  outside a set  $E$  of  $r$  with finite linear measure, where  $C_i$ ,  $i = 1, 2, 3$ , are non-negative constants independent of  $r$  and  $\varepsilon$ , and  $E$  is independent of  $\varepsilon$ . Because of the presence of the term  $8\varepsilon T(r, G)$  in (2.25), inequality (2.24) is better than (2.25), but inequality (2.25) is also sufficient for the later use in §§ 3 and 4.

*Remark 2.* It is hoped that Main Lemma (2.2) can be applied to the study of holomorphic curves in compact Kähler manifolds.

**EXAMPLE.** We give an example of  $f: \mathbb{A}^* \rightarrow M$  and  $\theta$  such that  $f^*\theta$  is really infinitely many-valued. Let  $M = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$  be an elliptic curve with  $\text{Im } \tau > 0$  and  $\pi: \mathbb{C} \rightarrow M$  the universal covering. Take any two points  $a, b$  of  $M$  so that  $n(a - b) \neq 0$  for all  $n \in \mathbb{Z}$ . Then there is a multiplicative meromorphic function  $\theta$  on  $M$  such that  $(\theta)_0 = a$  and  $(\theta)_\infty = b$ . Since  $n(a - b) \neq 0$  for all  $n \in \mathbb{Z}$ ,  $\theta$  is infinitely many-valued. Let  $\gamma_1$  (resp.  $\gamma_2$ ) be the cycle in  $M$  defined by  $\gamma_1: [0, 1] \ni t \rightarrow \pi(t) \in M$  (resp.  $\gamma_2: [0, 1] \ni t \rightarrow \pi(t\tau) \in M$ ). Then  $\{\gamma_1, \gamma_2\}$  is a basis of the first homology group  $H_1(M, \mathbb{Z})$ . One of the periods  $\frac{1}{2\pi i} \int_{\gamma_j} d \log \theta$ ,  $j = 1, 2$ , is irrational. Suppose that  $\frac{1}{2\pi i} \int_{\gamma_1} d \log \theta$  is irrational. The covering  $\mathbb{C} \xrightarrow{\pi} M$  is decomposed as

$$\mathbb{C} \xrightarrow{\pi_0} \mathbb{C}/\mathbb{Z} \xrightarrow{\pi_1} \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z}) = M.$$

Set  $\gamma: [0, 1] \ni t \rightarrow \pi_0(t) \in \mathbb{C}/\mathbb{Z} = \mathbb{C}^*$ , which is a cycle around  $\infty$  (or 0). Then  $\pi_1 \gamma = \gamma_1$ , so that the period  $\frac{1}{2\pi i} \int_\gamma d \log \theta \circ \pi_1$  is irrational. Let  $i: \mathbb{A}^* \rightarrow \mathbb{C}^*$  be the natural inclusion mapping and put  $f = \pi_1 \circ i: \mathbb{A}^* \rightarrow M$ . Then  $f^*\theta$  is infinitely many-valued.

Let  $\zeta^{(k)}$  denote the  $k$ -th derivative of  $\zeta$ . Using Main Lemma (2.2) inductively, one easily see the following:

**COROLLARY (2.26).** *Let the notation be as above. Then the inequality*

$$T(r, \zeta^{(k)}) \leq (k + 1)N(r, \text{Supp}(f^*\text{Res}(\omega))) + O(\log^+ T_f(r)) + O(\log r)$$

*holds for  $r \geq 1$  outside a set  $E$  with finite linear measure.*

§ 3. Inequality of the second main theorem type

Let  $V$  be a complex projective algebraic smooth variety of dimension  $n$ ,  $D$  an effective reduced divisor on  $V$  and  $\Omega_V^1(\log D)$  the sheaf of logarithmic 1-forms along  $D$  (cf., e.g., [12], [19]). Then  $\{\omega \in H^0(V, \Omega_V^1); \text{Supp}(\text{Res}(\omega)) \subset D\}$  spans  $H^0(V, \Omega_V^1(\log D))$  over  $C$  (see [19, Proposition 1.2]). Assume that there is a system  $\{\omega_i\}_{i=1}^{n+1}$  in  $H^0(V, \Omega_V^1(\log D))$  such that  $\phi_i = \omega_1 \wedge \cdots \wedge \omega_{i-1} \wedge \omega_{i+1} \wedge \cdots \wedge \omega_{n+1}, 1 \leq i \leq n+1$ , are linearly independent over  $C$ . Let  $f: \Delta^* \rightarrow V$  be a holomorphic curve such that  $f(\Delta^*) \not\subset D$ . Assume that  $f$  is non-degenerate with respect to  $\{\omega_i\}_{i=1}^{n+1}$ , i.e.,  $f(\Delta^*) \not\subset \{\sum c_i \phi_i = 0\}$  for any  $(c_i) \in C^{n+1} - \{O\}$ . Let  $\Omega$  be a Kähler form on  $V$  and set  $T_f(r) = T_f(r, \Omega)$ . Making use of Corollary (2.26) and Ochiai [22, Theorem A] as in [19, § 3] and [20], we have the following theorem.

**THEOREM (3.1).** *Let  $\{\omega_i\}_{i=1}^{n+1} \subset H^0(V, \Omega_V^1(\log D))$  and  $f: \Delta^* \rightarrow V$  be as above. Then there is a positive constant  $K$  depending only on  $\Omega$  and  $\{\omega_i\}_{i=1}^{n+1}$ , such that*

$$(3.2) \quad KT_f(r) < N(r, \text{Supp}(f^*D)) + S(r),$$

where  $S(r) = O(\log^+ T_f(r)) + O(\log r)$  as  $r \rightarrow \infty$  outside a set of  $r$  with finite linear measure.

Let  $\bar{R}$  be a compact Riemann surface,  $R = \bar{R} - \{a_i\}_{i=1}^q$  with distinct  $a_i \in \bar{R}$  and  $q < \infty$ , and  $a_0 \in R$  any point. Then there is a multiplicative meromorphic function  $\alpha$  such that  $(\alpha) = qa_0 - \sum a_i$ . The modulus  $|\alpha|$  turns out to be an exhaustion function of  $R$ . Set

$$R(t) = \{|\alpha| < t\}.$$

Let  $f: R \rightarrow V$  be a holomorphic curve. Put

$$T_f(r) = \int_1^r \frac{dt}{t} \int_{R(t)} f^* \Omega$$

for  $f$  and

$$n\left(t, \sum_{i=1}^{\infty} \nu_i b_i\right) = \sum_{|\alpha(b_i)| < t} \nu_i, \quad N\left(r, \sum_{i=1}^{\infty} \nu_i b_i\right) = \int_1^r \frac{n(t, \sum \nu_i b_i)}{t} dt$$

for a divisor  $\sum_{i=1}^{\infty} \nu_i b_i$  on  $R$  (cf. § 1 and [10, § 2]). For  $r_0$  large enough,  $R - R(r_0)$  is a union of  $\Delta_i^*$ ,  $i = 1, \dots, q$ , where  $\Delta_i^* \cap \Delta_j^* = \emptyset$  for  $i \neq j$  and  $\Delta_i = \Delta_i^* \cup \{a_i\}$  are a neighborhood of  $a_i$  in  $\bar{R}$ . Moreover the restriction

$1/z_i = 1/(\alpha|_{\Delta_i})$  of  $1/\alpha$  on every  $\Delta_i$  gives rise to a local coordinate in  $\Delta_i$  and  $\Delta_i^*$  is written as  $\Delta_i^* = \{r_0 \leq |z_i| < \infty\}$ . Therefore we have the following corollary of Theorem (3.1):

**COROLLARY (3.3).** *Let  $\{\omega_i\}_{i=1}^{n+1} \subset H^0(V, \Omega_V^1(\log D))$  be as in Theorem (3.1). Let  $f: R \rightarrow V$  be a holomorphic curve which is non-degenerate with respect to  $\{\omega_i\}_{i=1}^{n+1}$ . Then there is a positive constant  $K$  depending only on  $\Omega$  and  $\{\omega_i\}$  such that*

$$KT_f(r) \leq N(r, \text{Supp}(f^*D)) + S(r),$$

where  $S(r)$  is a small quantity as in (3.2).

*Remark.* Assume that  $\dim V = 1$ , and let us calculate sharp  $K$  in (3.2) in the way of the proof. The higher dimensional case will be discussed in § 4. Set  $T_f(r) = T_f(r, \Omega)$  for  $\Omega$  such that  $\int_V \Omega = 1$ .

(1) Let  $V = P^1$ . If the assumption of Theorem (3.1) for  $D$  is satisfied,  $D$  must consist of at least three points. Let  $D = \sum_{i=1}^q w_i$  be an effective reduced divisor on  $P^1$  with inhomogeneous coordinate  $w$  such that  $w_1 = 0, w_2 = \infty$  and  $q \geq 3$ . Let  $w_0 \in P^1 - D$  and set

$$\begin{aligned} \omega_1 &= d \log w \in H^0(P^1, \Omega_{P^1}^1(\log D)) \\ \omega_2 &= d \log \frac{\prod_{i=3}^q (w - w_i)}{(w - w_0)^{q-2}} \in H^0(P^1, \Omega_{P^1}^1(\log(D + w_0))). \end{aligned}$$

Then  $\phi = \omega_2/\omega_1$  is a rational function such that the degree  $\deg(\phi)_\infty$  of the divisor  $(\phi)_\infty$  is  $q - 1$ . We have by [18, Theorem 1]

$$(3.4) \quad T(r, f^*\phi) = (q - 1)T_f(r) + O(1).$$

Setting  $f^*\omega_i = \zeta_i dz$  for  $i = 1, 2$ , we obtain

$$\begin{aligned} (3.5) \quad T(r, f^*\phi) &= T\left(r, \frac{\zeta_1}{\zeta_2}\right) \leq T(r, \zeta_1) + T(r, \zeta_2) + O(\log r) + O(1) \\ &= N(r, f^{-1}(w_0)) + \sum_{i=1}^q N(r, f^{-1}(w_i)) + S(r). \end{aligned}$$

Hence we have by (3.4), (3.5) and the first main theorem (1.4)

$$(q - 2)T_f(r) \leq \sum_{i=1}^q N(r, f^{-1}(w_i)) + S(r),$$

which is the famous second main theorem for meromorphic functions on  $C$ .

(2) Let  $V$  be an elliptic curve. Then inequality (3.2) holds if  $D$

consists of one point  $a_0 \in V$ . On the other hand,  $H^0(V, \Omega_V^1(\log a_0)) = H^0(V, \Omega_V^1)$  is of dimension 1, where  $\Omega_V^1$  denotes the sheaf of germs of holomorphic 1-forms over  $V$ , so that the assumption of Theorem (3.1) is not fulfilled, but we can derive (3.2) for  $D = a_0$  by the method of the proof of Theorem (3.1) as follows. Take any point  $a_1 \in V - \{a_0\}$ . Then there is a multiplicative meromorphic function  $\theta$  such that  $(\theta) = a_0 - a_1$ . Set  $\omega_1 = d \log \theta \in H^0(V, \Omega_V^1(\log(a_0 + a_1)))$  and let  $\omega_2 \in H^0(V, \Omega_V^1)$  and  $\omega_2 \neq 0$ . We put  $\phi = \omega_1/\omega_2$ . Then  $\phi$  is a rational function on  $V$  such that  $\deg(\phi)_\infty = \deg(a_0 + a_1) = 2$ , so that by [18, Theorem 1] we have

$$(3.6) \quad T(r, f^*\phi) = 2T_f(r) + O(1).$$

Letting  $f^*\omega_i = \zeta_i dz$ ,  $i = 1, 2$ , we see that

$$(3.7) \quad \begin{aligned} T(r, f^*\phi) &= T\left(r, \frac{\zeta_1}{\zeta_2}\right) \leq T(r, \zeta_1) + T(r, \zeta_2) + O(\log r) + O(1) \\ &= N(r, f^{-1}(a_0)) + N(r, f^{-1}(a_1)) + S(r). \end{aligned}$$

Therefore it follows from (3.6) and (3.7) that

$$T_f(r) \leq N(r, f^{-1}(a_0)) + S(r).$$

(3) Let  $V$  be a compact Riemann surface of genus  $\geq 2$ . Then  $\dim H^0(V, \Omega_V^1) \geq 2$ , so that the condition of Theorem (3.1) is satisfied with  $D = 0$ . This implies the well-known fact that the isolated singularity of a holomorphic curve in  $V$  of genus  $\geq 2$  is removable.

**§ 4. Extension theorem of big Picard type**

Let  $A$  be a quasi-Abelian variety (see [11] and [12]), i.e.,  $A$  is an algebraic group which is commutative and admits the exact sequence

$$0 \longrightarrow (C^*)^l \longrightarrow A \xrightarrow{\rho} A_0 \longrightarrow 0,$$

where  $A_0$  is an Abelian variety. Taking the natural embedding  $(C^*)^l \subset (P^1)^l$ , we have a smooth completion  $\bar{A} = (P^1)^l \times_{(C^*)^l} A$  of  $A$  with boundary divisor  $D$  which has only normal crossings, and the canonical projection  $\bar{\rho}: \bar{A} \rightarrow A_0$ . One may regard  $\bar{\rho}: \bar{A} \rightarrow A_0$  as a fibre bundle over  $A_0$  with fibre  $(P^1)^l$  and structure group  $(C^*)^l$ . Let  $X$  be an algebraic subvariety of  $A$  which is of general type or equally of hyperbolic type (cf. [11]). In the present case,  $X$  is of general type if and only if the group  $\{a \in A; X + a = X\}$  of translations which preserve  $X$  is finite (see [11] and [12]). Let

$W$  be the union of subvarieties of  $X$  which are translations of non-trivial closed algebraic subgroups of  $A$ .

LEMMA (4.1). *Let  $X$  and  $W$  be as above. Then  $W$  is a proper algebraic subvariety of  $X$ , of which each irreducible component is foliated by translations of a non-trivial closed algebraic subgroup of  $A$ .*

*Remark.* This lemma was proved in [21] when  $\dim X = 2$ . In [13], Kawamata proved it in the case when  $A$  is an Abelian variety. To prove it in the present form, we need further consideration. The idea of the following proof is due to Kawamata.

*Proof.* Let  $\pi: C^m \rightarrow A$  be the universal covering with  $m = \dim A$ ,  $A = C^m/\Lambda$  with a discrete subgroup  $\Lambda$  (cf. [12]), and  $\lambda: C^m - \{O\} \rightarrow P^{m-1}$  the natural mapping into the projective space  $P^{m-1}$  of lines in  $C^m$  through the origin  $O$ . Let  $U$  be a small open set in  $P^{m-1}$  and set

$$s(\bar{X}) = \bigcup_{x \in U} (\bar{X} + \pi(s(x)), x) \subset \bar{A} \times U$$

for a holomorphic section  $s \in \Gamma(U, C^m - \{O\})$ , where  $\bar{X}$  is the Zariski closure of  $X$  in  $\bar{A}$  and “ $+ \pi(s(x))$ ” stands for the natural action of  $A$  on  $\bar{A}$ . Hence  $s(\bar{X})$  is an analytic subset of  $\bar{A} \times U$ . We set

$$Y_U = \bigcap_{s \in \Gamma(U, C^m - \{O\})} s(\bar{X}) \subset \bar{A} \times U.$$

Then  $Y_U$  is again an analytic subset of  $\bar{A} \times U$  and we see that a point  $(a, x) \in \bar{A} \times U$  belongs to  $Y_U$  if and only if  $a + \phi(t) \in \bar{X}$  for every  $t \in C$ , where  $\phi(t)$  is the analytic 1-parameter subgroup of  $A$  such that  $d\phi/dt(0) = x$ . Let  $B_x$  denote the Zariski closure in  $A$  of the analytic 1-parameter subgroup of  $A$  associated with the vector  $x$ . Then we have that

$$(4.2) \quad (a, x) \in Y_U \iff a + B_x \subset \bar{X}.$$

Let  $U'$  be another small open set in  $P^{m-1}$ . Then it follows from (4.2) that  $Y_U$  coincides with  $Y_{U'}$  in  $\bar{A} \times (U \cap U')$ , so that  $Y = \bigcup_U Y_U$  is a well-defined analytic subset of  $\bar{A} \times P^{m-1}$  and so algebraic in  $\bar{A} \times P^{m-1}$ . Let  $Y_0 = Y \cap (A \times P^{m-1})$  and  $p: A \times P^{m-1} \rightarrow A$  be the projection. Then by (4.2) and the definition of  $W$ ,  $p(Y_0) = W$ . Since  $p$  is proper and rational,  $W$  is a closed algebraic subvariety of  $X$ . Now we must show that  $W \neq X$  and each irreducible component of  $W$  is foliated by translations of a non-trivial closed algebraic subgroup of  $A$ . Since there are only countably many

non-trivial closed algebraic subgroups in  $A$  as in the case of an Abelian variety (cf. [12]), we denote them by  $\{B_i\}_{i=1}^\infty$ . We see by (4.2) that

$$(4.3) \quad a \in W \iff a + B_i \subset W \text{ for some } B_i .$$

Let  $h_i: X \rightarrow A/B_i$  be the restriction of the natural morphism from  $A$  onto the quotient  $A/B_i$  on  $X$  and put

$$W_i = \{x \in X; \dim_x h_i^{-1}(h_i(x)) = \dim B_i\} .$$

Then  $W_i$  is a proper algebraic subvariety of  $X$  because  $X$  is of general type, and  $W = \bigcup_i W_i$  by (4.3). Let  $W_i = \bigcup_j W_{ij}$  be the irreducible decomposition of  $W_i$ . We get a countable covering  $W = \bigcup_{i,j} W_{ij}$ . It is clear that every  $W_{ij} \neq X$ . By virtue of Baire's theorem we see that  $W \neq X$  and that an irreducible component of  $W$  must be one  $W_{ij}$  which is foliated by translations of  $B_i$ .

Let  $Z$  be an algebraic subvariety of  $A$  and  $Z_{\text{reg}}$  the set of regular points of  $Z$  with the inclusion mapping  $i: Z_{\text{reg}} \rightarrow A$ . Let  $J_\nu(Z_{\text{reg}})$  (resp.  $J_\nu(A)$ ) be the  $\nu$ -th holomorphic jet bundle over  $Z_{\text{reg}}$  (resp.  $A$ ) (see [22]). Then the mapping  $i$  naturally induces a bundle homomorphism  $i_*: J_\nu(Z_{\text{reg}}) \rightarrow J_\nu(A)$ . Since  $A$  is a quasi-Abelian variety, there is a regular isomorphism  $J_\nu(A) \cong A \times C^{\nu m}$ . Let  $q: A \times C^{\nu m} \rightarrow C^{\nu m}$  be the projection and set

$$I_\nu = q \circ i_*: J_\nu(Z_{\text{reg}}) \rightarrow C^{\nu m} \quad (\text{cf. [22]}).$$

We denote by  $j_\nu g$  the  $\nu$ -th jet of a holomorphic curve  $g: (C, 0) \rightarrow Z_{\text{reg}}$  from a neighborhood of the origin  $0$  of  $C$  into  $Z_{\text{reg}}$ .

LEMMA (4.4). *Let  $X$  and  $W$  be as in Lemma (4.1). Let  $g: (C, 0) \rightarrow X$  be a holomorphic curve such that  $g(0) \notin W$  and  $g(0) \in Z_{\text{reg}}$ , where  $Z$  is the Zariski closure of the image of  $g$  in  $X$ . Then the differential*

$$dI_\nu: T(J_\nu(Z_{\text{reg}})) \rightarrow T(C^{\nu m})$$

*is injective at  $j_\nu g$  for all large  $\nu$ , where  $T(\cdot)$  denotes the holomorphic tangent bundle.*

This lemma is a refined version of a lemma due to M. Green by which he completed the work of Ochiai [22] on Bloch's conjecture [2]<sup>3</sup>. M. Green showed it in case  $A$  is complete, i.e.,  $A$  is an Abelian variety, but his proof works in the non-complete case.

2) M. Green gave the proof of the lemma at "Conference on Geometric Function Theory" held at Katata, Sept. 1-6, 1978.

Let  $\bar{X}$  be the Zariski closure of  $X$  in  $\bar{A}$ .

**THEOREM (4.5)** (big Picard theorem). *Let  $X$  and  $W$  be as above. Then any holomorphic curve  $f: \Delta^* \rightarrow X$  has a holomorphic extension  $\tilde{f}: \Delta = \Delta^* \cup \{\infty\} \rightarrow \bar{X}$  unless  $f(\Delta^*) \subset W$ .*

*Proof.* We fix a Kähler form  $\Omega$  on  $\bar{A}$  and set  $T_f(r) = T_f(r, \Omega)$ . By (2.10), (2.11) and [16, I, p. 369], it suffices to prove that  $T_f(r)/\log r$  is bounded as  $r \rightarrow \infty$ . Let  $Z$  be the Zariski closure of  $f(\Delta^*)$  in  $X$ . Then  $f(z) \notin W$  and  $f(z) \in Z_{\text{reg}}$  for  $z \in \Delta^*$  except for some discrete set of points. Making use of Lemma (4.4) and Main Lemma (2.2) (more precisely, Corollary (2.26)) as in [19], we have

$$(4.6) \quad T_f(r) \leq K_1 \log^+ T_f(r) + K_2 \log r$$

for  $r \geq 1$  outside a set  $E$  of  $r$  with finite linear measure, where  $K_1$  and  $K_2$  are non-negative constants independent of  $r$ . We may assume that  $f$  is not a constant curve. Then we see that  $T_f(r) \uparrow \infty$  as  $r \uparrow \infty$ . Since  $T_f(r)$  is a convex increasing function in  $\log r$ ,  $T_f(r)/\log r$  is monotone increasing. Therefore we have by (4.6)

$$\lim_{r \rightarrow \infty} \frac{T_f(r)}{\log r} \leq K_2,$$

which completes the proof.

**COROLLARY (4.7).** *Let  $f: N - S \rightarrow X$  be a holomorphic mapping from a complex manifold  $N$  minus a thin analytic set  $S$  into  $X$ . If  $f(N - S) \not\subset W$ , then  $f$  extends to a meromorphic mapping  $\tilde{f}: N \rightarrow \bar{X}$ .*

*Proof.* We take an embedding  $\bar{X} \subset P^N$  into some projective space  $P^N$  with a homogeneous coordinate system  $(w_0, \dots, w_N)$  such that  $f(N - S) \not\subset \{w_0 = 0\}$ . Let  $f_i = f^*(w_i/w_0)$ . It is enough to prove that every  $f_i$  extends to a meromorphic function on  $N$ . By virtue of Hartogs' theorem, we may assume that  $N = \Delta \times \Delta^{k-1}$  and  $S = \{\infty\} \times \Delta^{k-1}$  ( $k = \dim N$ ). Put  $S' = \{z' \in \Delta^{k-1}; \Delta^* \times \{z'\} \subset f^{-1}(W)\}$ , which is a thin analytic set of  $\Delta^{k-1}$ . By Hartogs' theorem, it suffices to show that  $f_i$  extends meromorphically over  $\Delta \times (\Delta^{k-1} - S')$ . For each  $z'_0 \in \Delta^{k-1} - S'$ , the holomorphic curve  $f(\cdot, z'_0): \Delta^* \ni z_1 \mapsto f(z_1, z'_0) \in X$  does not lie in  $W$ . By Theorem (4.5),  $f$  is extendable over  $\Delta$ , so that  $f_i(\cdot, z'_0)$  is meromorphic in  $\Delta$ . We put  $f_i(z_1, z'_0) = z_1^{\mu(z'_0)} \cdot g_i(z_1, z'_0)$ , where  $\mu(z'_0) \in Z$  and  $g_i(\infty, z'_0) \neq 0, \infty$ . Take a small neighborhood  $U$  of  $z'_0$ . Then we see that  $\mu(z')$  is bounded in  $z' \in U$ . Therefore  $f_i(z_1, z')$



is meromorphic in  $\Delta \times U$ , and so is in  $\Delta \times (\Delta^{k-1} - S)$ .


*Remark.* Fujimoto ([3], [5]) and Green ([8]) proved extension theorems of big Picard type for holomorphic mappings into  $P^n$  omitting more than  $n + 1$  hyperplanes in general position. Their results will be discussed in Example 1 below. Here, let us give a simple and new observation to another theorem of Green [8, Parts 4 and 5] from the viewpoint of this paper. He proved the following interesting theorem:

*Let  $f: C \rightarrow V \subset P^n$  be a holomorphic curve into a subvariety  $V$  of  $P^n$  omitting  $\dim V + 2$  non-redundant hyperplane sections of  $V$ . Then  $f$  is algebraically degenerate, i.e.,  $f(C)$  is contained in a proper subvariety of  $V$ .*

Here "non-redundant" means that no one of the hyperplane sections is contained in the union of the others. Let  $D$  be the sum of the  $\dim V + 2$  hyperplane sections of  $V$ . Let  $\pi: V' \rightarrow V - D$  be a desingularization of  $V - D$  and  $\bar{V}'$  a smooth completion of  $V'$  with boundary divisor  $D'$  of normal crossing type. Setting  $\bar{q}(V') = \dim H^0(\bar{V}', \Omega_{\bar{V}'}^1(\log D'))$  which is called the logarithmic irregularity of  $V'$  ([12]), we have by the assumption for  $D$

$$(4.8) \quad \bar{q}(V') < \dim V'.$$

We may assume that  $f$  can be lifted to a holomorphic curve  $f': C \rightarrow V'$  such that  $\pi \circ f' = f$ . Let  $\alpha: V' \rightarrow A$  be the quasi-Albanese mapping (see [12]),  $X = \overline{\alpha(V')}$  the Zariski closure of  $\alpha(V')$  in  $A$ ,  $G$  the identity component of the group  $\{a \in A; X + a = X\}$ ,  $h: A \rightarrow A/G = A_1$  the canonical mapping onto the quotient  $A/G = A_1$  and  $X_1 = \overline{h(X)}$ . Then (4.8) implies that  $X_1$  is of positive dimension and of general type. Let  $W_1$  be the union of subvarieties of  $X_1$  which are translations of non-trivial closed algebraic subgroups of  $A_1$ . By Lemma (4.1),  $W_1$  is a proper algebraic subvariety of  $X_1$ . Put  $f_1 = h \circ \alpha \circ f'$ :



Then we have  $f_1(C) \subset W_1$  by Theorem (4.5) if  $f_1$  is not a constant curve, so that  $f$  is algebraically degenerate. Thus inequality (4.8) implies the

algebraic degeneracy of  $f'$ ; this is just a non-complete version of Bloch's conjecture (see [2], [22]).

EXAMPLE 1. Let  $D_i, 0 \leq i \leq n + k$ , be  $n + k + 1$  distinct hyperplanes of  $P^n$  and set  $V = P^n - \sum_0^{n+k} D_i$ . Then we have

$$\bar{q}(V) = \dim H^0(P^n, \Omega_{P^n}^1(\log \sum_0^{n+k} D_i)) = n + k.$$

Assume that  $k \geq 1$ . Then  $\bar{q}(V) > \dim V$ . Let  $\alpha: V \rightarrow A = (C^*)^{n+k}$  be the quasi-Albanese mapping and  $f: C \rightarrow V$  a holomorphic curve. As in Remark above, we see that  $\alpha \circ f(C)$  lies in a translation of a closed algebraic subgroup of  $A$ , so that  $f(C)$  lies in a proper linear subspace of  $P^n$ . This fact was proved in Green [7, Theorem 2].

Suppose that  $k = 1$  and the  $D_i$ 's are in general position. We take a system  $(w_0, w_1, \dots, w_n)$  of homogeneous coordinates of  $P^n$  so that  $D_i = \{w_i = 0\}$  for  $i = 0, 1, \dots, n$  and  $D_{n+1} = \{w_0 + \dots + w_n = 0\}$ . Put  $x_i = w_i/w_0$  for  $i = 1, \dots, n$ . Then the quasi-Albanese mapping  $\alpha: V \rightarrow (C^*)^{n+1}$  is written as

$$\alpha: V \ni (x_1, \dots, x_n) \mapsto \left( x_1, \dots, x_n, \frac{1 + x_1 + \dots + x_n}{n} \right) \in (C^*)^{n+1}.$$

Set  $X = \{(y_1, \dots, y_{n+1}) \in (C^*)^{n+1}; ny_{n+1} = 1 + y_1 + \dots + y_n\}$ . Then  $\alpha: V \rightarrow X$  is biregular and so  $X$  is of general type. Let  $\Pi$  denotes the union of diagonal hyperplanes of  $\sum_1^{n+1} D_i$  (see [15, Example 16, p. 395] and [4, p. 243]). Let  $W$  be the proper algebraic subvariety of  $X$  as in Lemma (4.1). Then  $W = \alpha(\Pi)$ . In this case, Fujimoto [4, Theorem 5.5] and Green [8, Part 3] showed Theorem (4.5) (cf. also [1], [5] and [7]). In case  $n = 2$ , the figure of  $W$  in  $X$  is as follows:




Fig. 1

Here each  $W_i \cong C^*$  and  $W = W_1 \cup W_2 \cup W_3$ .

EXAMPLE 2 ([14, Example 1, p. 92]). Let  $Q = \sum_{i=0}^4 L_i$  be a complete

quadrilateral in  $P^2$  as in Kobayashi [14, Example 1, p. 92], and set  $V = P^2 - Q$ . Take a homogeneous coordinate system  $(w_0, w_1, w_2)$  of  $P^2$  such that

$$\begin{aligned} L_0 &= \{w_0 = 0\}, & L_1 &= \{w_1 = 0\}, & L_2 &= \{w_0 - w_1 = 0\}, \\ L_3 &= \{w_2 = 0\}, & L_4 &= \{w_0 - w_2 = 0\}. \end{aligned}$$

Then we have the quasi-Albanese mapping

$$\alpha: V \ni (x_1, x_2) \mapsto \left(\frac{1}{2}x_1, x_1 - 1, \frac{1}{2}x_2, x_2 - 1\right) \in (C^*)^4,$$

where  $x_i = w_i/w_0, i = 1, 2$ . Thus  $\alpha(V) = X = \{(y_1, \dots, y_4) \in (C^*)^4; y_2 = 2y_1 - 1, y_4 = 2y_3 - 1\}$  and  $\alpha: V \rightarrow X$  is biregular. Since there is no  $C^*$  in  $X, W = \emptyset$ . Therefore any holomorphic curve  $f: \Delta^* \rightarrow V$  is extendable to a holomorphic curve  $\tilde{f}: \Delta \rightarrow P^2$ . Kobayashi [14, p. 92] proved this fact by showing that  $V$  is hyperbolically embedded in  $P^2$ .

EXAMPLE 3 ([19, § 4(b)]). Let  $X = \{(x_1, \dots, x_{n+2}) \in (C^*)^{n+2}; x_{n+1} = 1 + x_1 + \dots + x_{n-1}, x_{n+2} = x_1 + \dots + x_n\}$  and  $n \geq 3$ . Then  $X$  is of general type. For the simplicity, let  $n = 3$ . Let  $W$  be the proper algebraic subvariety of  $X$  as in Lemma (4.1). Then we see that

$$W = W_1 \cup W_2 \cup \dots \cup W_5,$$

where  $W_1 \cong (C^*)^2$  and  $W_i \cong C^*$  for  $i = 2, 3, 4, 5$ . The figure of  $W$  in  $X$  is illustrated as follows:




Fig. 2

EXAMPLE 4 ([22, § 5]). Let  $A = E_1 \times \dots \times E_4$  be a product of four elliptic curves  $E_i$  belonging to distinct isogeny classes. Let  $X$  be the hypersurface of  $A$  as defined in Ochiai [22, § 5]. Then the algebraic sub-

variety  $W$  of  $X$  as in Lemma (4.1) consists of several elliptic curves which are mutually disjoint.

Lastly we pose a problem and a conjecture related to Theorems (4.5) and (3.1).

**PROBLEM.** *What can we say of the Kobayashi hyperbolicity of  $X$  or  $X - W$  in Theorem (4.5)?*

*Remark.* Green [9] gave a nice criterion of the Kobayashi hyperbolicity, but in the present case his criterion does not work since an irreducible component  $W'$  of  $W$  may admit a non-constant holomorphic curve  $f: C \rightarrow W'$  omitting the other components of  $W$  (see Examples 3 and 4).

The case (2) of Remark to Theorem (3.1) suggests that the following conjecture may be true:

**CONJECTURE.** Let  $A$  be an Abelian variety and  $D$  an effective reduced divisor on  $A$ . Let  $\Omega \in c_1([D])$  be a semi-positive definite  $(1, 1)$ -form in the first Chern class  $c_1([D]) \in H^{1,1}(A, \mathbb{C})$  of  $[D]$ . Then we have

$$T_r(r, \Omega) \leq N(r, f^*D) + S(r)$$

for algebraically non-degenerate holomorphic curves  $f: \Delta^*$  (or  $C$ )  $\rightarrow A$ , where  $S(r) = O(\log^+ T_r(r, \Omega)) + O(\log r)$  as  $r \rightarrow \infty$  outside a set of  $r$  with finite linear measure.

#### REFERENCES

- [ 1 ] A. Bloch, Sur les systèmes de fonctions holomorphes à variétés linéaires lacunaires, Ann. Sci. École Norm. Sup., **43** (1926), 309–362.
- [ 2 ] —, Sur les systèmes de fonctions uniformes satisfaisant à l'équation d'une variété algébrique dont l'irrégularité dépasse la dimension, J. Math. Pures Appl., **5** (1926), 19–66.
- [ 3 ] H. Fujimoto, Extensions of the big Picard's theorem, Tôhoku Math. J., **24** (1972), 415–422.
- [ 4 ] —, Families of holomorphic maps into the projective space omitting some hyperplanes, J. Math. Soc. Japan, **25** (1973), 235–249.
- [ 5 ] —, On meromorphic maps into the complex projective space, J. Math. Soc. Japan, **26** (1974), 272–288.
- [ 6 ] H. Grauert and H. Reckziegel, Hermitesche Metriken und normale Familien holomorpher Abbildungen, Math. Z., **89** (1965), 108–125.
- [ 7 ] M. Green, Holomorphic maps into complex projective space omitting hyperplanes, Trans. Amer. Math. Soc., **169** (1972), 89–103.
- [ 8 ] —, Some Picard theorems for holomorphic maps to algebraic varieties, Amer. J. Math., **97** (1975), 43–75.
- [ 9 ] —, The hyperbolicity of the complement of  $2n+1$  hyperplanes in general posi-

- tion in  $P^n$ , and related results, Proc. Amer. Math. Soc., **66** (1977), 103–113.
- [10] P. Griffiths and J. King, Nevanlinna theory and holomorphic mappings between algebraic varieties, Acta Math., **130** (1973), 145–220.
  - [11] S. Iitaka, Logarithmic forms of algebraic varieties, J. Fac. Sci. Univ. Tokyo Sect. IA, **23** (1976), 525–544.
  - [12] —, On logarithmic Kodaira dimension of algebraic varieties, Complex Analysis and Algebraic Geometry, pp. 175–189, Iwanami, Tokyo, 1977.
  - [13] Y. Kawamata, On Bloch's conjecture, Invent. Math., **57** (1980), 97–100.
  - [14] S. Kobayashi, Hyperbolic Manifolds and Holomorphic Mappings, Pure and Appl. Math., **2**, Dekker, New York, 1970.
  - [15] —, Intrinsic distances, measures and geometric function theory, Bull. Amer. Math. Soc., **82** (1976), 357–416.
  - [16] R. Nevanlinna, Einige Eindeutigkeitsätze in der Theorie der meromorphen Funktionen, Acta Math., **48** (1926), 367–391.
  - [17] —, Le Théorème de Picard-Borel et la théorie des fonctions méromorphes, Gauthier-Villars, Paris, 1939.
  - [18] J. Noguchi, Holomorphic mappings into closed Riemann surfaces, Hiroshima Math. J., **6** (1976), 281–291.
  - [19] —, Holomorphic curves in algebraic varieties, Hiroshima Math. J., **7** (1977), 833–853.
  - [20] —, Supplement to “Holomorphic curves in algebraic varieties”, Hiroshima Math. J., **10** (1980), 229–231.
  - [21] —, Rigidity of holomorphic curves in some surfaces of hyperbolic type, unpublished notes.
  - [22] T. Ochiai, On holomorphic curves in algebraic varieties with ample irregularity, Invent. Math., **43** (1977), 83–96.
  - [23] A. L. Vitter, The lemma of the logarithmic derivative in several complex variables, Duke Math. J., **44** (1977), 89–104.
  - [24] A. Weil, Introduction à l'Étude des Variétés kählériennes, Hermann, Paris, 1958.

*Department of Mathematics  
College of General Education  
Osaka University*