

Essay

Letters, numbers, symbols and words: how to communicate grades of evidence and recommendations

Holger J. Schünemann, Dana Best, Gunn Vist, Andrew D. Oxman, for the GRADE Working Group

§ See related article page 672

Abstract

THE GRADE WORKING GROUP IS DEVELOPING and evaluating a common, sensible approach to grading quality of evidence and strength of recommendations in health care. In this article, we discuss the advantages and disadvantages of using letters, numbers, symbols or words to represent grades of evidence and recommendations. Using multiple strategies, we searched for comparative studies of alternative ways of representing ordered categories in any context. In addition, we contacted experts and reviewed theoretical work and qualitative research on how best to communicate grades of any kind quickly and clearly. We were unable to identify health care research that addressed, either directly or indirectly, the best way to present grades of evidence and recommendations. We found examples of symbols used by government, commercial and consumer organizations to communicate quality of evidence or strength of recommendations, but no comparative studies. Although a number of grading systems are used in health care and other fields, there is little or no evidence of how well various presentations are understood. Before promoting the use of specific symbols, numbers, letters or words, the extent to which the intended message is comprehended should be evaluated.

Organizations such as the Canadian Task Force on Preventive Health Care¹ and more than 100 other groups² use various systems of codes to communicate grades of evidence and recommendations. The codes fall primarily into 3 categories: letters (e.g., A, B, C, etc.), numbers (e.g., I, II, III, etc.) and mixed letters and numbers (e.g., Ia, Ib, IIa, etc.).

Health care practitioners, especially students, are often puzzled by the message a grade conveys. For example, the administration of oral anticoagulation in patients with atrial fibrillation and rheumatic mitral valve disease receives various grades of recommendation from different organizations: Class I based on level B evidence by the American Heart Association,³ grade C recommendation based on level IV evidence by SIGN² and grade 1C+ (where the 1 indicates the balance between benefit and harm and C+ the methodological quality of the underlying evidence) by the American College of Chest Physicians.⁴ Thus, the various grading systems may not be fulfilling their intended func-

tion: to communicate a clear message, quickly and concisely. Indeed, if the same code, used by different systems, represents different meanings, bewilderment and incomprehension may result.

We formed the GRADE Working Group with the hope of reaching agreement on a common, sensible approach to grading quality of evidence and strength of recommendations. We consider here the advantages and disadvantages of using letters, numbers, symbols or words to represent different grades.

Evidence

We conducted a search of MEDLINE and PsychLit databases for the period 1966 to April 2002 (strategy available on request). In addition, we searched for theoretical work and qualitative research addressing how best to communicate grades of any kind quickly and clearly. Furthermore, we searched relevant texts⁵⁻⁸ and bibliographies and contacted researchers from other fields (e.g., psychology, marketing and graphic design). Because empirical evidence on the use of symbols comes from grading schemes unrelated to health care — such as *Consumer Reports*⁹ and restaurant¹⁰ and hotel guides¹¹ — we contacted organizations responsible for popular grading schemes (list available on request).

Based on the information derived from the literature searches, reflection on the target audience and the messages that grades of evidence and recommendations are intended to communicate, we established criteria for assessing alternative grading schemes. See Table 1 on the *CMAJ* Web site.

We did not find any studies comparing different systems of communicating grades in health care. A number of studies have compared alternative ways of presenting information about risk,^{2,12-16} but none addressed the use of codes or grades. We also did not find comparative studies of alternative ways of presenting grades or evaluations of how well grading systems are understood or used by target audiences.

We identified several examples of symbols used by gov-

ernment, commercial and consumer organizations.^{9,11,14,17} Among the best known guides that use symbols are the Michelin restaurant and hotel guides,¹¹ which use spoons and stars to communicate hotel quality where a larger number of symbols indicates higher quality. However, neither Michelin nor any other organization or agency we contacted was aware of a comparative study or evaluation of the effectiveness of their own or other schemes for communicating the intended information.

Considerations in developing grading systems

Although health care professionals are the main target audience for systematic reviews and practice guidelines, policymakers, insurers and consumers also use them. As consumers (patients) are the ultimate beneficiaries of reviews and guidelines, some authors argue that these tools should be accessible to the general public, often via the Internet.¹⁸ To ensure that such information is comprehensible to this wide range of users, grades of evidence and recommendations should be understandable to people from different cultures with varying levels of literacy and different languages.

By definition, grades are ordered; from 2 to as many as 20 levels have been used for grading evidence and recommendations.² A small number of levels may be easier to understand and interpret than many levels.¹⁵ In a landmark article, Miller¹⁹ described how humans' capacity to perceive differences, for example, in the loudness of a sound or the saltiness of a solution, failed beyond 7 categories.

The number of levels used and the upper and lower limits in the grading scheme should be as intuitively obvious as possible and clearly described. If numbers are used, it should be clear whether higher numbers indicate a better grade than lower numbers and what the upper limit of the system is.

Because grades of evidence and recommendations represent multiple dimensions, the system displaying this information should convey more than 1 dimension. At a minimum, the presentation should distinguish between 2 basic concepts: the quality of evidence (i.e., the extent to which one can be confident that an estimate of effect is correct) and the strength of the recommendation (i.e., the extent to which one can be confident that adherence to the recommendation will do more good than harm).¹⁶ Many current grading systems do not make this distinction adequately.²

Associations that people may make between a system presenting grades of evidence and recommendations and other types of grades may help or hinder interpretation. For example, letters are commonly used for grades in schools, where they have a specific meaning, e.g., "C" represents average and "F" represents failure.

We identified potential problems associated with using letters, numbers, symbols and words. See Table 2 on the *CMAJ* Web site.

Letters are commonly used for grades, easily communi-

cated verbally and are likely to be understood intuitively in many cultures. However, because there is more than 1 alphabet, the use of letters may be limited across cultures and languages.

Letters and numbers can be used together to represent 2 dimensions, but this may create confusion about which represents the quality of evidence and which represents the balance between benefits and harms. In addition, many practice guideline schemes already use letters and numbers with varying definitions, which may cause misunderstandings.

Numbers are intuitively communicated and understood, succinct, do not require a high degree of literacy and may have the same meaning across cultures and languages to a larger extent than letters.

The extent to which symbols and words are easily understood is likely to vary greatly, as is the risk of problems with associations, recognition of the number of levels and limits and the ability to convey 2 dimensions. Symbols are succinct and likely to be understood across different levels of literacy and across different languages, but they may have different meanings in different cultures. Symbols have the advantage of being quickly and easily recognized, easy to identify in scanned text and, once learned, may convey understanding better than words through a strong association with the concept they represent. However, symbols may be difficult or inappropriate to communicate verbally. For example, a smiling face would be hard to explain in association with a strong recommendation for chemotherapy in end-stage cancer.

Summary

We were unable to identify research that addresses either directly or indirectly the presentation of grades of evidence and recommendations. An argument against continuing to use letters and numbers in this context without comparative studies is the confusion that currently exists due to inconsistent use of letters and numbers by many organizations. If symbols are used, they should be easily understood across different cultures; their limits, direction and number of levels must be intuitively clear; and it must be possible to convey 2 dimensions easily. Examples of numbers, letters and symbols that appear to meet most of these criteria are presented in Fig. 1.

Before promoting the use of specific symbols or words, the extent to which their intended message is comprehended in a particular grading scheme should be evaluated in a comparative, cross-cultural study including clinicians and consumers as participants.

The GRADE Working Group has developed a system for grading evidence and recommendations and is evaluating its reliability and sensibility. We are also developing guidelines for considering costs and issues of equity when making recommendations, how to accommodate questions about diagnostic tests and how best to present grades of evidence and recommendations.

Quality of evidence	Numbers	Letters	Circles	Stars	Multiple
High	1	A	●	☆☆☆☆	⊕⊕⊕⊕
Moderate	2	B	◐	☆☆☆	⊕⊕⊕👤
Low	3	C	◑	☆☆	⊕⊕👤👤
Very low	4	D	◒	☆	⊕👤👤👤

Action based on balance between benefit and harm	Numbers	Letters	Traffic lights	Thumbs	Arrows
Do	1	A			↑↑
Probably do	2	B			↑?
Probably don't do	3	C			↓?
Don't do	4	D			↓↓

Fig. 1: Examples of possible symbols for representing quality of evidence and the balance between benefits and harm in health care recommendations. See Tables 1 and 2 on the CMAJ Web site for selection criteria (see www.cmaj.ca).

This article has been peer reviewed.

From the Departments of Medicine and of Social and Preventive Medicine, University of Buffalo, Buffalo, NY (Schünemann); the Department of Clinical Epidemiology and Biostatistics, McMaster University, Hamilton, Ont. (Schünemann); the Division of General Pediatrics, Children's National Medical Center, Washington, DC (Best); and the Department of Health Services Research, Norwegian Directorate for Health and Social Welfare, Oslo, Norway (Vist, Oxman)

Competing interests: None declared.

Contributors: All authors contributed substantially to the concept, design, data analysis, acquisition and interpretation. Dana Best and Holger Schünemann drafted the first version. Gunn Vist and Andrew Oxman revised it critically for important intellectual content. All of the authors approved the final version.

Opinions expressed in this paper do not necessarily represent those of the institutions with which the authors are affiliated.

References

1. Canadian Task Force on Preventive Health Care. History and methods. Available: www.ctfphc.org (accessed 2003 Aug 27).
2. National Guideline Clearinghouse. NGC Browse — Organizations. Available: www.guidelines.gov/browse/browse.aspx (accessed 2003 Jul 17).
3. Fuster V, Rydén LE, Asinger RW, Cannon DS, Crijns HJ, Frye RL, et al. ACC/AHA/ESC guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and the European Society of Cardiology Committee to develop guidelines for the management of patients with arterial fibrillation. *J Am Coll Cardiol* 2001;38:1266.

4. Albers G, Dalen JE, Laupacis A, Manning WJ, Petersen P, Singer DE. Antithrombotic therapy in atrial fibrillation. *Chest* 2001;119:194S-206S.
5. Harris R. *Information graphics: a comprehensive illustrated reference*. Oxford: Oxford University Press; 1999.
6. Tufte E. *Visual explanations*. Cheshire (CT): Graphics Press; 1997.
7. Tufte E. *The visual display of quantitative information*. 2nd ed. Cheshire (CT): Graphics Press; 2001.
8. Wroldstad M, Fisher DF, editors. *Toward a new understanding of literacy*. Westport (CT):Praeger; 1986.
9. *Consumer Reports* [Web site]. Available: www.consumerreports.org (accessed 2003 Jul 17).
10. Zagat Survey. Available: www.zagat.com/ (accessed 2003 Jul 17).
11. Michelin. *Key for hotels and restaurants*. Available: www.michelin-travel.com/eng/aide/p1342.htm (accessed 2003 Jul 17).
12. McGettigan P, Sly K, O'Connell D, Hill S, Henry D. The effects of information framing on the practices of physicians. *J Gen Intern Med* 1999;14:633-42.
13. Edwards A, Hood K, Matthews E., Russell D, Russell I, Barker J, et al. The effectiveness of one-to-one risk communication interventions in health care: a systematic review. *Med Decis Making* 2000;20:290-7.
14. Film Advisory Board. FAB rating system. Available: http://www.filmadvisoryboard.org/ratings/Default.asp (accessed 2003 Jul 17).
15. DeVellis B. *Scale development: theory and applications*. Newbury Park (CA): Sage Publications; 1991.
16. Wainer H. Graphical data analysis. *Annu Rev Psychol* 1981;32:191-241.

17. National Highway Traffic Safety Administration. Available: www.nhtsa.dot.gov (accessed 2003 Jul 15).
18. DISCERN: Health information on the Internet. Available: www.discrim.org.uk/HOTI.htm (accessed 2003 Jul 17).
19. Miller G. The magic number seven plus or minus two: some limits on our capacity for processing information. *Psychol Bull* 1956;63:81-97.

Correspondence to: Dr. Holger J. Schünemann, Departments of Medicine and of Social and Preventive Medicine, University of Buffalo, 270 Farber Hall, 3435 Main St., Buffalo NY 14214, USA; fax 716-898-4493; hjs@buffalo.edu

Members of the GRADE Working Group: David Atkins, Chief Medical Officer, Center for Practice and Technology Assessment, Agency for Healthcare Research and Quality, USA; Dana Best, Assistant Professor, Department of General Pediatrics and Adolescent Medicine, George Washington University, Children's National Medical Center, USA; Peter A Briss, Acting Chief Community Guide Branch, Centers for Disease Control and Prevention, USA; Martin Eccles, Professor, and James Mason, Professor, Centre for Health Services Research, University of Newcastle upon Tyne, U.K.; Yngve Falck-Ytter, Associate Director, German Cochrane Centre, Institute for Medical Biometry and Medical Informatics, University Hospital Freiburg, Germany; Gunn E. Vist, Researcher, Signe Flottorp, Researcher, and Andrew D. Oxman, Director, Department of Health Services Research, Norwegian Directorate for Health and Social Welfare, Norway; Gordon H. Guyatt, Professor, and Roman Jaeschke, Associate Clinical Professor, Departments of Clinical Epidemiology and Biostatistics and Medicine, McMaster University, Canada; Robin T. Harbour, Quality and Information

Director, Scottish Intercollegiate Guidelines Network, United Kingdom; Margaret C. Haugh, Methodologist, Fédération Nationale des Centres de Lutte Contre le Cancer, France; David Henry, Professor and Suzanne Hill, Senior Lecturer, Department of Clinical Pharmacology, Faculty of Medicine and Health Sciences, University of Newcastle, Australia; Gillian Leng, Guidelines Programme Director, National Institute for Clinical Excellence, United Kingdom; Alessandro Liberati, Professor, Università di Modena e Reggio Emilia and Centro per la Valutazione della Efficacia della Assistenza Sanitaria, Italy; Nicola Magrini, Director, Centro per la Valutazione della Efficacia della Assistenza Sanitaria, Italy; Philippa Middleton, Honorary Research Fellow, Australasian Cochrane Centre, Australia; Jacek Mrukowicz, Executive Director, Polish Institute for Evidence Based Medicine, Poland; Dianne O'Connell, Senior Epidemiologist, Cancer Epidemiology Research Unit, Cancer Research and Registers Division, The Cancer Council, Australia; Bob Phillips, Associate Fellow, Centre for Evidence-based Medicine, University Department of Psychiatry, Warneford Hospital, United Kingdom; Holger J Schünemann, Assistant Professor, Departments of Medicine and of Social & Preventive Medicine, University of Buffalo, USA; Tessa Tan-Torres Edejer, Medical Officer/Scientist, Global Programme on Evidence for Health Policy, World Health Organisation, Switzerland; Helena Varonen, Associate Editor, Finnish Medical Society Duodecim, Finland; John W. Williams Jr., Associate Professor, The Center for Health Services Research in Primary Care, Health Services Research and Development, Department of Veterans Affairs Medical Center and Duke University Medical Center, USA; Stephanie Zaza, Acting Associate Director for Science, Epidemiology Program Office, Centers for Disease Control and Prevention, USA

CMAJ Essay Prize Deadline: Feb. 1, 2004

CMAJ is offering a single open-category prize of \$1000 for the best piece of writing submitted to the 2004 CMAJ Essay Prize contest. We welcome entries from physicians, students, residents and non-physicians. We are looking for reflective essays of up to 2000 words on topics of interest to a general medical readership.

The winner will be selected by a committee appointed from CMAJ's Editorial Board. The judges will make their decision on the basis of originality of thought and quality of writing. The winning manuscript must be suitable for publication in CMAJ. All papers submitted will be considered for publication. The journal reserves the right not to award a prize. Prospective entrants are encouraged to read the description of the selection of winners for the 2000 Essay Prize in our June 26, 2001, issue (available at www.cmaj.ca/cgi/content/full/164/13/1859).

Authors should submit their papers with a cover letter stating that they would like their manuscript to be considered for the 2004 CMAJ Essay Prize. We welcome submissions by mail (CMAJ, 1867 Alta Vista Drive, Ottawa ON K1G 3Y6), fax (613 565-5471) or email (pubs@cma.ca) until the deadline of Feb. 1, 2004.

