

Level Set Method

For Motion by Mean Curvature

Tobias Holck Colding and William P. Minicozzi II

ABSTRACT. Modeling of a wide class of physical phenomena, such as crystal growth and flame propagation, leads to tracking fronts moving with curvature-dependent speed. When the speed is the curvature this leads to one of the classical degenerate nonlinear second-order differential equations on Euclidean space. One naturally wonders, "What is the regularity of solutions?" A priori solutions are only defined in a weak sense, but it turns out that they are always twice differentiable classical solutions. This result is optimal; their second derivative is continuous only in very rigid situations that have a simple geometric interpretation. The proof weaves together analysis and geometry. Without deeply understanding the underlying geometry, it is impossible to prove fine analytical properties.



Figure 1. Oil droplets in water can be modeled by the level set method.

Tobias Holck Colding is Cecil and Ida B. Green Distinguished Professor of Mathematics at MIT. His email address is colding@math.mit.edu.

William P. Minicozzi II is Professor of Mathematics at MIT. His email address is minicozz@math.mit.edu.

For permission to reprint this article, please contact: reprint-permission@ams.org.

DOI: <http://dx.doi.org/10.1090/noti1439>

The spread of a forest fire, the growth of a crystal, the inflation of an airbag, and a droplet of oil floating in water can all be modeled by the level set method. See Figure 1. One of the challenges for modeling is the presence of discontinuities. For example, two separate fires can expand over time and eventually merge to one, as in Figure 2, or a droplet of liquid can split as in Figure 3. The level set method allows for this.




Figure 2. After two fires merge the evolving front is connected.



Figure 3. Water separates into droplets.

The level set method has been used with great success over the last thirty years in both pure and applied mathematics. Given an initial interface or front M_0 bounding a region in \mathbf{R}^{n+1} , the level set method is used to analyze its subsequent motion under a velocity field. The idea is to represent the evolving front as a level set of a function $v(x, t)$, where x is in \mathbf{R}^{n+1} and t is time. The initial front M_0 is given by

$$M_0 = \{x \mid v(x, 0) = 0\},$$

and the evolving front is described for all later time t as the set where $v(x, t)$ vanishes, as in Figure 4. There are many functions that have M_0 as a level set, but the evolution of the level set does not depend on the choice of the function $v(x, 0)$.

In mean curvature flow, the velocity vector field is the mean curvature vector, and the evolving front is the level




Figure 4. The gray areas represent trees that a forest fire has not yet reached. The edge is the burning fire front that is moving inward as time propagates left to right in the upper part of the figure. The propagation of the fire front is given as a level set of an evolving function in the second line.




Figure 5. The red curves represent the evolving front at three different times. Each is the zero level set of the evolving function. The front here is moving outward as in Figure 2. In Figures 3 and 4, the front is moving inward.

set of a function that satisfies a nonlinear degenerate parabolic equation. Solutions are defined in a weak, so-called “viscosity,” sense; in general, they may not even be differentiable (let alone twice differentiable). However, it turns out that for a monotonically advancing front viscosity solutions are in fact twice differentiable and satisfy the equation in the classical sense. Moreover, the situation becomes very rigid when the second derivative is continuous.

Suppose $\Sigma \subset \mathbf{R}^{n+1}$ is an embedded hypersurface and \mathbf{n} is the unit normal of Σ . The **mean curvature** is given by $H = \text{div}_\Sigma(\mathbf{n})$. Here

$$\text{div}_\Sigma(\mathbf{n}) = \sum_{i=1}^n \langle \nabla_{e_i} \mathbf{n}, e_i \rangle,$$

where e_i is an orthonormal basis for the tangent space of Σ . For example, at a point where \mathbf{n} points in the x_{n+1} direction and the principal directions are in the other axis directions,

$$\text{div}_\Sigma(\mathbf{n}) = \sum_{i=1}^n \frac{\partial \mathbf{n}_i}{\partial x_i}$$

is the sum (n times the mean) of the principal curvatures. If $\Sigma = u^{-1}(s)$ is the level set of a function u on \mathbf{R}^{n+1} and

s is a regular value, then $\mathbf{n} = \frac{\nabla u}{|\nabla u|}$ and

$$H = \sum_{i=1}^n \langle \nabla_{e_i} \mathbf{n}, e_i \rangle = \text{div}_{\mathbf{R}^{n+1}} \left(\frac{\nabla u}{|\nabla u|} \right).$$

The last equality used that $\langle \nabla_{\mathbf{n}} \mathbf{n}, \mathbf{n} \rangle$ is automatically 0 because \mathbf{n} is a unit vector.

A one-parameter family of smooth hypersurfaces $M_t \subset \mathbf{R}^{n+1}$ flows by the **mean curvature flow** if the speed is equal to the mean curvature and points inward:

$$x_t = -H \mathbf{n},$$

where H and \mathbf{n} are the mean curvature and unit normal of M_t at the point x . Our flows will always start at a smooth embedded connected hypersurface, even if it becomes disconnected and nonsmooth at later times. The earliest reference to the mean curvature flow we know of is in the work of George Birkhoff in the 1910s, where he used a discrete version of this, and independently in the material science literature in the 1920s.

Two Key Properties

- H is the gradient of area, so mean curvature flow is the negative gradient flow for volume ($\text{Vol} M_t$ decreases most efficiently).
- Avoidance property: If M_0 and N_0 are disjoint, then M_t and N_t remain disjoint.

The avoidance principle is simply a geometric formulation of the maximum principle. An application of it is illustrated in Figure 6, which shows that if one closed hypersurface (the red one) encloses another (the blue one), then the outer one can never catch up with the inner one. The reason for this is that if it did there would be a first point of contact and right before that the inner one would contract faster than the outer one, contradicting that the outer was catching up.

Curve Shortening Flow

When $n = 1$ and the hypersurface is a curve, the flow is the curve shortening flow. Under the curve shortening flow, a round circle shrinks through round circles to a point in finite time. A remarkable result of Matthew Grayson from 1987 (using earlier work of Richard Hamilton and Michael Gage) shows that any simple closed curve in the plane remains smooth under the flow until it disappears




Figure 6. Disjoint surfaces avoid each other; contact leads to a contradiction.




Figure 7. Grayson proved that even a tightly wound region becomes round under curve shortening flow.

in finite time in a point. Right before it disappears, the curve will be an almost round circle.

The evolution of the snake-like simple closed curve in Figure 7 illustrates this remarkable fact. (The eight figures are time shots of the evolution.)

Level Set Flow

The analytical formulation of the flow is the level set equation that can be deduced as follows. Given a closed embedded hypersurface $\Sigma \subset \mathbf{R}^{n+1}$, choose a function $v_0 : \mathbf{R}^{n+1} \rightarrow \mathbf{R}$ that is zero on Σ , positive inside the domain bounded by Σ , and negative outside. (Alternatively, choose a function that is negative inside and positive outside.)

- If we simultaneously flow $\{v_0 = s_1\}$ and $\{v_0 = s_2\}$ for $s_1 \neq s_2$, then avoidance implies they stay disjoint.
- In the level set flow, we look for $v : \mathbf{R}^{n+1} \times [0, \infty) \rightarrow \mathbf{R}$ so that each level set $t \rightarrow \{v(\cdot, t) = s\}$ flows by mean curvature and $v(\cdot, 0) = v_0$.
- If $\nabla v \neq 0$ and the level sets of v flow by mean curvature, then

$$v_t = |\nabla v| \operatorname{div} \left(\frac{\nabla v}{|\nabla v|} \right).$$

This is degenerate parabolic and undefined when $\nabla v = 0$. It may not have classical solutions.

In a paper cited more than 12,000 times from 1988, Stanley Osher and James Sethian studied this equation numerically. The analytical foundation was provided by Craig Evans and Joel Spruck in a series of four papers in the early 1990s and, independently and at the same time, by Yun Gang Chen, Yoshikazu Giga, and Shunichi Goto. Both of these two groups constructed (continuous) viscosity solutions and showed uniqueness. The notion of viscosity solutions had been developed by Pierre-Louis Lions and Michael G. Crandall in the early 1980s. The work of these two groups on the level set flow was one of the significant applications of this theory.

Examples of Singularities

Under mean curvature flow a round sphere remains round but shrinks and eventually becomes extinct in a point. A round cylinder remains round and eventually becomes extinct in a line. The marriage ring is the example of a thin torus of revolution in \mathbf{R}^3 . Under the flow the marriage ring shrinks to a circle, then disappears. See Figure 8.

Dumbbell

Figure 9 shows the evolution of a rotationally symmetric mean convex dumbbell in \mathbf{R}^3 . If the neck is sufficiently thin, then the neck pinches off first, and the surface disconnects into two components (bell). Later each component (bell) shrinks to a round point. This example falls into a larger category of surfaces that are rotationally symmetric around an axis. Because of the symmetry, the solution reduces to a one-dimensional heat equation. This was analyzed in the early 1990s by Sigurd Angenent, Steven Altschuler, and Giga; cf. also with work of Halil Mete Soner and Panagiotis Souganidis from around the same time. A key tool in the arguments of Angenent-Altschuler-Giga was a parabolic Sturm-Liouville theorem of Angenent that holds in one spatial dimension.

Singular Set

Under mean curvature flow closed hypersurfaces contract, develop singularities, and eventually become extinct. The **singular set** S is the set of points in space and time where the flow is not smooth.

In the first three examples—the sphere, the cylinder, and the marriage ring— S is a point, a line, and a closed curve, respectively. In each case, the singularities occur only at a single time. In contrast, the dumbbell has two singular times with one singular point at the first time and two at the second.

Mean Convex Flows

A hypersurface is convex if every principal curvature is positive. It is mean convex if $H > 0$, i.e., if the sum of the principal curvatures is positive at every point. Under the mean curvature flow, a mean convex hypersurface moves inward, and, since mean convexity is preserved, it will continue to move inward and eventually sweep out the entire compact domain bounded by the initial hypersurface.

In many applications, the speed of the moving interface does not change sign and the front moves monotonically. This corresponds to positive mean curvature in our case. Monotone movement can be modeled particularly



Figure 8. The marriage ring shrinks to a circle under mean curvature flow.




Figure 9. A narrow neck on a dumbbell pinches off, and then each half shrinks to a point.

efficiently numerically by the Fast Marching Method of James Sethian.

Level Set Flow for Mean Convex Hypersurfaces

When the hypersurfaces are mean convex, the equation can be rewritten as a degenerate elliptic equation for a function u defined by

$$u(x) = \{t \mid x \in M_t\}.$$

We say that u is the **arrival time**, since it is the time the hypersurfaces M_t arrive at x as the front sweeps through the compact domain bounded by the initial hypersurface. The arrival time has a game theoretic interpretation by work of Robert Kohn and Sylvia Serfaty. It follows easily that if we set $v(x, t) = u(x) - t$, then v satisfies the level set flow. Now the level set equation $v_t = |\nabla v| \operatorname{div}(\nabla v / |\nabla v|)$ becomes

$$-1 = |\nabla u| \operatorname{div} \left(\frac{\nabla u}{|\nabla u|} \right).$$

This is a degenerate elliptic equation that is undefined when $\nabla u = 0$. Note that if u satisfies this equation, then so does u plus a constant. This just corresponds to shifting the time when the flow arrives by a constant. A

particular example of a solution to this equation is the function $u = -\frac{1}{2}(x_1^2 + x_2^2)$, which is the arrival time for shrinking round cylinders in \mathbf{R}^3 . In general, Evans-Spruck (cf. Chen-Giga-Goto) constructed Lipschitz solutions to this equation.

Singular Set of Mean Convex Level Set Flow

The singular set of the flow is the critical set of u . Namely, $(x, u(x))$ is singular if and only if $\nabla_x u = 0$. For instance, in the example of the shrinking round cylinders in \mathbf{R}^3 , the arrival time is given by $u = -\frac{1}{2}(x_1^2 + x_2^2)$, and the flow is singular in the line $x_1 = x_2 = 0$, that is, exactly where $\nabla u = 0$.

We will next see that even though the arrival time was only a solution to the level set equation in a weak sense, it turns out to be always a twice differentiable classical solution.

Differentiability

From [CM2]:

- u is twice differentiable everywhere, with bounded second derivatives, and smooth away from the critical set.
- u satisfies the equation everywhere in the classical sense.
- At each critical point the Hessian is symmetric and has only two eigenvalues, 0 and $-\frac{1}{k}$; $-\frac{1}{k}$ has multiplicity $k + 1$.

This result is equivalent to saying that at a critical point, say $x = 0$ and $u(x) = 0$, the function u is (after possibly a rotation of \mathbf{R}^{n+1}) up to higher-order terms equal to the quadratic polynomial

$$-\frac{1}{k} (x_1^2 + \dots + x_{k+1}^2).$$

This second-order approximation is simply the arrival time of the shrinking round cylinders. It suggests that the level sets of u right before the critical value and near the origin should be approximately cylinders (with an $n - k$ dimensional axis). This has indeed been known for a long time and is due to Huisken, White, Sinestrari, Andrews, and Haslhofer-Kleiner. It also suggests that those cylinders should be nearly the same (after rescaling to unit size). That is, the axis of the cylinders should not depend on the value of the level set. This last property, however, was only very recently established in [CM1] and is the key to proving that the function is twice differentiable.¹ The proof that the axis is unique, independent on the level set, relies on a key new inequality that draws

This kind of uniqueness is a famously difficult problem in geometric analysis

¹Uniqueness of the axis is parallel to the fact that a function is differentiable at a point precisely if on all sufficiently small scales at that point it looks like the same linear function.




Figure 10. The figure illustrates a situation that turns out to be impossible, where the axes of the cylinders for three different level sets near the critical point are radically different. Instead, as the level sets approach the critical point, the axes go to a unique limit.

its inspiration from real algebraic geometry, although the proof is entirely new. This kind of uniqueness is a famously difficult problem in geometric analysis, and no general case had previously been known.

Regularity of Solutions

We have seen that the arrival time is always twice differentiable, and one may wonder whether there is even more regularity. Gerhard Huisken showed already in 1990 that the arrival time is C^2 for convex M_0 . However, in 1992 Tom Ilmanen gave an example of a rotationally symmetric mean convex M_0 in \mathbf{R}^3 where u is not C^2 . This result of Ilmanen shows that the above theorem about differentiability cannot be improved to C^2 . We will see later that in fact one can entirely characterize when the arrival time is C^2 . In the plane, Kohn and Serfaty (2006) showed that u is C^3 , and for $n > 1$ Natasa Sesum (2008) gave an example of a convex M_0 where u is not C^3 . Thus Huisken's result is optimal for $n > 1$.

The next result shows that one can entirely characterize when the arrival time is C^2 .

Continuous Differentiability

[CM3]: u is C^2 if and only if:

- There is exactly one singular time (where the flow becomes extinct).
- The singular set S is a k -dimensional closed connected embedded C^1 submanifold of cylindrical singularities.

Moreover, the axis of each cylinder is the tangent plane to S .

When u is C^2 in \mathbf{R}^3 , the singular set S is either:

- (1) a single point with a spherical singularity or
- (2) a simple closed C^1 curve of cylindrical singularities.

The examples of the sphere and the marriage ring show that each of these phenomena can happen, whereas the example of the dumbbell does not fall into either case, showing that in that case the arrival time is not C^2 .

We can restate this result for \mathbf{R}^3 in terms of the structure of the critical set and Hessian: u is C^2 if and only if u has exactly one critical value and the critical set is either:

- (1) a single point where Hess_u is $-\frac{1}{2}$ times the identity or
- (2) a simple closed C^1 curve where Hess_u has eigenvalues 0 and -1 with multiplicities 1 and 2, respectively.

In case (2), the kernel of Hess_u is tangent to the curve.

Concluding Remarks

We have seen that for one of the classical differential equations, in order to understand the analysis it is necessary to understand the underlying geometry. There are many tantalizing parallels to other differential equations, both elliptic and parabolic.

For references about mean curvature flow, see the survey [CMP] and [CM1]–[CM3].

References

- [CM1] T. H. COLDING and W. P. MINICOZZI II, Uniqueness of blowups and Łojasiewicz inequalities, *Annals of Math.* **182** (1) (2015), 221–285. MR 3374960
- [CM2] ———, Differentiability of the arrival time, *Comm. Pure Appl. Math.*, DOI: 10.1002/cpa.21635.
- [CM3] ———, Regularity of the level set flow, submitted, <https://arxiv.org/abs/1606.05185>.
- [CMP] T. H. COLDING, W. P. MINICOZZI II, and E. K. PEDERSEN, Mean curvature flow, *Bull. Amer. Math. Soc. (N.S.)* **52** (2015), no. 2, 297–333. MR 3312634

Credits

Figure 1, by Lee Netherton, is used under the terms of the Creative Commons License.

<https://creativecommons.org/licenses/by-sa/2.0/>.

Figure 2 was created by James Sethian and is used with permission.

Figure 3 is courtesy of Adam Hart-Davis, www.adam-hart-davis.org.

Figure 5 was created by James Sethian and is used with permission.

Figures 7 and 9 were created via computer simulation by Uwe Mayer and are used with permission.

Photos of Tobias Holck Colding and William P. Minicozzi II are courtesy of Bryce Vickmark/MIT.



Tobias Holck Colding

ABOUT THE AUTHORS

Tobias Holck Colding completed the PhD under Christopher Croke at the University of Pennsylvania in 1992. Prior to joining MIT he was a professor at Courant Institute–NYU. He is a foreign member of the Royal Danish Academy of Science and Letters and Fellow of the American Academy of Arts & Sciences. He received the 2010 Oswald Veblen Prize in Geometry and the 2016 Carlsberg Foundation Research Prize.



William P. Minicozzi

William P. Minicozzi II received a PhD in mathematics from Stanford University under Rick Schoen in 1994. In 2010 he received the Oswald Veblen Prize in Geometry. He was elected Fellow of the American Academy of Arts & Sciences in 2015.

MathJobs.Org

The automated job application database sponsored by the AMS

Free for Applicants



MathJobs.Org offers a paperless application process for applicants and employers in mathematics

Registered Applicants Can:

- Create their own portfolio of application documents
- Make applications online to participating employers
- Choose to make a cover sheet viewable by all registered employers

Registered Employers Can:

- Post up to seven job ads
- Set all criteria for required documents, and add specific questions
- Receive and upload reference letters
- Manage applicant information and correspondence quickly and easily
- Set limited access permissions for faculty and EOE administrators
- Search for and sort additional applicants in the database
- Choose an advertising-only account, or a discounted single ad account

Visit mathjobs.org for pricing information

Contact: Membership and Programs Department

American Mathematical Society
201 Charles Street
Providence, RI 02904-2294 USA

800.321.4267, ext. 4105
Email: mathjobs@ams.org

