

# Lichen flora of Rodnei Mountains National Park (Eastern Carpathians, Romania) including new records for the Romanian mycoflora

Ioana Violeta Ardelean<sup>1,2</sup>, Christine Keller<sup>1</sup> & Christoph Scheidegger<sup>1</sup>

<sup>1</sup>WSL, Swiss Federal Research Institute, 8903 Birmensdorf, Switzerland.

E-mail: ioana.ardelean@wsl.ch

<sup>2</sup>Faculty of Biology and Geology, Babeş-Bolyai University, Cluj-Napoca, Romania

**Abstract:** We investigated the lichen flora of the main habitats existing in Rodnei Mountains identifying 283 lichen species, and one subspecific taxon. Of these, 67 taxa are new records for the lichen flora of Romania, and 182 species are reported for the first time in Rodnei Mountains. Considering previous reports and our results, 442 lichen taxa are reported in Rodnei Mountains region in total, accounting for approx. 35% of the total lichen flora of Romania. When comparing the Red Lists of Romania and surrounding Carpathian countries, our data revealed the presence of a high number of threatened species in the region.

## INTRODUCTION

During the last years, various lichenological studies reported new species for the Romanian lichen flora (Crişan, 2006; Çobanoğlu et al., 2009, 2011; Yavuz & Çobanolu, 2008; Vondrák & Šoun, 2008), but at the same time, species that had been recorded before, such as *Anzina carneonivea* (Scheidegger, 1985) and *Lepraria incana* (Bartók, 1999) are missing in the current checklist (Ciurchea, 2004).

260 lichen species from 11 locations were previously reported in Rodnei Mountains (Ciurchea, 2004), and reflected the high species richness of the national park. However, the knowledge of lichen species distribution is still scattered in Romania and some of the species records have not been rechecked since more than a century (Bartok & Crişan, personal communication).

Conserved areas are well known sanctuaries for threatened species including various groups of lichens (Goward, 1995; Zoller et al., 2000; Nascimbene et al., 2013; Ignatov et al., 2004; Lackovičová & Guttová, 2006). Furthermore, lichen habitats such as old-growth forest stands and veteran trees in extensively managed meadows with their important lichen microhabitats are often lost in managed and perturbed areas (Wolseley, 1995; Thor, 1995; Scheidegger & Werth, 2009).

The aim of this study was to assess the lichen flora of characteristic habitats of the Rodnei

Mountains in a replicated design with a standarized lichen diversity assessment. This study should also reveal the importance of conserved areas for the maintenance of lichen diversity in the Rodnei Mountains and thus contribute to decisions about future conservation strategies within this biosphere reserve.

## MATERIALS AND METHODS

### Study area

The Rodnei Mountains are located in the northern part of the Eastern Carpathians reaching their highest elevation at Pietrosul Mare Peak (2303 m). Most of the study area is part of the Rodnei Mountains National Park, established in 1932 and declared as a UNESCO Biosphere Reserve in 1979.

### Climate

Due to the position and orientation (East–West) of the mountains, the climate is characterised by the Baltic and the Oceanic influences. The mean annual temperature decreases with altitude, ranging between 6 °C at the base of the mountains and -1.5 °C at the highest altitudes. Mean annual precipitations range from 1300 to 1400 mm (Gorduza, 1983).

### Sampling

The sampling method was structured according to the sampling design described by Scheidegger

et al. (2002). Four main substrates were considered in each circular plot of 1 ha: trees, dead wood, soil and rock, and for each we applied 6 relevés. If the substrate was not available or it was not colonised with lichens, it was substituted with other available substrates, thus achieving 24 relevés in each plot. All lichen species within a relevé surface of 0.2 m<sup>2</sup> were collected (except the crustose lichens from rocks).

The investigated habitats are well represented in the Rodnei Mountains and follow an altitudinal gradient from wooded meadows to mixed and coniferous forests, to *Pinus mugo* shrubs and alpine vegetation with bare rocks. Each of the five habitats was analysed at two levels: conserved and managed (except *Pinus mugo* shrubs and wooded meadows, for which only one type of management was found in the study area). There were seven replicated circular plots of 1 ha for each habitat with its corresponding levels of conservation (i.e., 56 plots in total). The minimum distance between the plots was 100 m. The selection of the 9 sampling localities (Fig. 1)

aimed at covering the types of habitat and stand characteristics (Table 1) of the investigated area.


The lichen specimens were identified based on morphological and chemical characteristics using mostly the keys of Smith et al. (2009), Wirth (1995) and Tønsberg (1992).

For the crustose sterile species and the specimens that needed chemical analyses for identification, thin layer chromatography (TLC) was used according to the methods described in White & James (1985) with solvents A, B and C.

The specimens are stored in the Herbaria of "Alexandru Borza" Botanical Garden, Babeş-Bolyai University, Cluj-Napoca, Romania.

### Data assessment

The importance of lichen flora in our results was determined by the total number of species, the number of new species for the region, and the new species for Romanian lichen flora. We also took into account the number of species in the Red List of macrolichens from Romania (Bartok & Crişan, personal communication)



**Fig. 1.** Location of the Rodnei Mts National Park in Romania, and study area with sampling localities. 1 – Cascada Cailor, 2 – Rotunda Pass, 3 – Repedea Valley, 4 – Borşa 1, 5 – Borşa 2, 6 – Izvorul Dragoş Valley, 7 – Pietrosul Mare, 8 – Gropile, 9 – Bătrâna.

**Table 1.** 56 investigated plots (within the 9 sampling localities, see Fig. 1). Abbreviations: AC – conserved alpine vegetation with bare rocks, AM – managed alpine vegetation with bare rocks, PM – *Pinus mugo* shrubs, CC – conserved coniferous forest, CM – managed coniferous forest, MC – conserved mixed forest, MM – managed mixed forest, WM – wooded meadows, C – conserved, M – managed.

Plot code	Location name	Coordinates x	Coordinates y	Altitude (m)	Exposition	Slope [°]	Conservation status	Habitat type
AC1	Petrosul Mare	47,5941	24,6365	2143	W	33	C	alpine vegetation
AC2	Petrosul Mare	47,5915	24,6379	2125	E	10	C	alpine vegetation
AC3	Petrosul Mare	47,5955	24,6505	2136	E	5	C	alpine vegetation
AC4	Petrosul Mare	47,5935	24,6407	2171	S	39	C	alpine vegetation
AC5	Petrosul Mare	47,6004	24,6231	1916	W	39	C	alpine vegetation
AC6	Petrosul Mare	47,5994	24,6304	2193	S	12	C	alpine vegetation
AC7	Petrosul Mare	47,5996	24,6279	2173	S	43	C	alpine vegetation
AM1	Gropile	47,579	24,6325	2085	E	7	M	alpine vegetation
AM2	Gropile	47,5703	24,6468	2020	S	24	M	alpine vegetation
AM3	Gropile	47,5715	24,6396	2004	S	5	M	alpine vegetation
AM4	Gropile	47,5739	24,6344	1085	S	29	M	alpine vegetation
AM5	Bătrâna	47,5699	24,6124	1915	S	24	M	alpine vegetation
AM6	Bătrâna	47,568	24,6116	1816	S	26	M	alpine vegetation
AM7	Bătrâna	47,5782	24,6032	1719	W	24	M	alpine vegetation
PC1	Petrosul Mare	47,6018	24,6479	1785	E	14	C	<i>Pinus mugo</i> shrubs
PC2	Petrosul Mare	47,5886	24,6472	1882	S	36	C	<i>Pinus mugo</i> shrubs
PC3	Petrosul Mare	47,6019	24,6506	1788	N	28	C	<i>Pinus mugo</i> shrubs
PC4	Petrosul Mare	47,5913	24,6437	2040	S	32	C	<i>Pinus mugo</i> shrubs
PC5	Petrosul Mare	47,6014	24,6229	2019	W	33	C	<i>Pinus mugo</i> shrubs
PC6	Petrosul Mare	47,6071	24,6113	1821	W	26	C	<i>Pinus mugo</i> shrubs
PC7	Bătrâna	47,5692	24,596	1781	S	30	C	<i>Pinus mugo</i> shrubs
CC1	Repedea Valley	47,5964	24,6918	1164	E	26	C	spruce forest
CC2	Repedea Valley	47,5952	24,679	1326	S	38	C	spruce forest
CC3	Petrosul Mare	47,6059	24,6067	1596	S	32	C	spruce forest
CC4	Petrosul Mare	47,6028	24,6054	1477	S	33	C	spruce forest
CC5	Petrosul Mare	47,6053	24,6026	1470	S	37	C	spruce forest
CC6	Petrosul Mare	47,6036	24,6002	1353	W	25	C	spruce forest
CC7	Petrosul Mare	47,6014	24,6013	1316	S	37	C	spruce forest
CM1	Borșa1	47,62	24,6589	1148	N	27	M	spruce forest
CM2	Repedea Valley	47,5938	24,7012	1151	N	31	M	spruce forest
CM3	Repedea Valley	47,5982	24,7051	1375	W	36	M	spruce forest
CM4	Repedea Valley	47,6018	24,7087	1357	N	25	M	spruce forest
CM5	Cascada Cailor	47,6062	24,795	1275	W	14	M	spruce forest
CM6	Izvorul Dragoș Valley	47,5982	24,6022	1081	E	35	M	spruce forest
CM7	Izvorul Dragoș Valley	47,5989	24,6016	932	N	19	M	spruce forest
MC1	Repedea Valley	47,5972	24,6907	966	N	40	C	mixed forest
MC2	Cascada Cailor	47,5951	24,797	1207	N	35	C	mixed forest
MC3	Rotunda Pass	47,5244	24,9998	1044	W	21	C	mixed forest
MC4	Rotunda Pass	47,5284	25,0041	1089	W	14	C	mixed forest
MC5	Izvorul Dragoș Valley	47,6083	24,5874	863	E	35	C	mixed forest
MC6	Repedea Valley	47,5943	24,6815	1180	E	44	C	mixed forest
MC7	Izvorul Dragoș Valley	47,6016	24,5935	1019	N	15	C	mixed forest
MM1	Repedea Valley	47,5899	24,7002	1196	W	42	M	mixed forest
MM2	Repedea Valley	47,6149	24,6977	866	W	14	M	mixed forest

Table 1 (continued)

Plot code	Location name	Coordinates x	Coordinates y	Altitude (m)	Exposition	Slope [°]	Conservation status	Habitat type
MM3	Repedea Valley	47,6128	24,6871	1257	E	35	M	mixed forest
MM4	Repedea Valley	47,6146	24,6922	1058	E	33	M	mixed forest
MM5	Izvorul Dragoş Valley	47,6227	24,5758	903	N	32	M	mixed forest
MM6	Izvorul Dragoş Valley	47,6228	24,5858	930	N	41	M	mixed forest
MM7	Izvorul Dragoş Valley	47,6242	24,5957	915	S	5	M	mixed forest
WM1	Borşa1	47,6216	24,6494	1095	N	18	M	wooded meadows
WM2	Borşa1	47,6224	24,6396	1048	N	13	M	wooded meadows
WM3	Borşa1	47,6228	24,6325	1048	N	6	M	wooded meadows
WM4	Borşa1	47,6287	24,6635	1988	N	16	M	wooded meadows
WM5	Borşa2	47,6504	24,6356	783	N	10	M	wooded meadows
WM6	Izvorul Dragoş Valley	47,6209	24,6176	1036	N	23	M	wooded meadows
WM7	Izvorul Dragoş Valley	47,6211	24,6125	957	N	33	M	wooded meadows

and the Red Lists of the surrounding countries which harbour the Carpathian mountain ridge: Hungary (Lökös & Tóth, 1996), Ukraine (Didukh, 2009), Slovakia (Pisút et al., 2001), and Poland (Cieslinski et al., 2003), with detailed information in Table 2.

## RESULTS AND DISCUSSIONS

We found 283 lichens species and one subspecific taxon for the Rodnei Mountains in 56 plots (Table 3). We confirmed 102 species that were reported previously, plus 182 taxa that were new records for the region, of which 67 species have never been reported in Romania. Poorly developed specimens that could not be unambiguously assigned to listed species are not

reported as new. Considering previous reports and our results, 442 lichen taxa are reported in Rodnei Mountains region, accounting for approx. 35% of the total lichen flora of Romania. Out of 284 taxa, 13 are listed in the Red List of macrolichens of Romania (Bartok & Crişan, personal communication), 8 species in the Red List of Ukraine (Didukh, 2009), 65 in the Red List of Hungary (Lökös & Tóth, 1996), 96 in the Red List of Slovakia (Pisút et al., 2001), and 125 in the Red List of Poland (Cieslinski et al., 2003).

The most species-rich genus is *Cladonia*, of which we found 43 species in our study area. Even though most species are common, this genus contributed substantially to the lichen species richness of Rodnei Mountains.

**Table 2.** Red Lists (RL) including the number of lichen species according to IUCN categories: RE – Regionally extinct, CR – Critically Endangered, EN – Endangered, VU – Vulnerable, NT – Near Threatened, LC – Least Concern, DD – Data Deficient, additional category in RL of Hungary: R – rare (problematic species including literature data without voucher specimens and new findings with no further information on the distribution), and in RL of Ukraine: R – rare (species known from few locations, with relatively stable populations but low rates), and Total no of sp. – total number of species in the Red List.

RL	IUCN Categories							R	Total no of sp.
	RE	CR	EN	VU	NT	LC	DD		
Romania		20	27	26					73
Hungary	30	24	44	45				260	403
Ukraine			5	26				21	52
Slovakia	88	140	48	169	114		14		573
Poland	141	179	201	165	68	22	110		886

**Table 3.** Lichen species recorded in the study area. Abbreviations: New R Mts – new records for the Rodnei Mountains, New RO – new records for Romania; RL RO – Red List of Macrolichens from Romania (Bartok & Crișan, personal communication), RL UK – Red List of lichens from Ukraine (Didukh, 2009), RL HU – Red List of lichens from Hungary (Lökös & Tóth, 1996), RL PL – Red List of lichens from Poland (Cieslinski et al., 2003), RL SL – Red List of lichens from Slovakia (Pisút et al., 2001); Obs – number of plots where the species was observed; categories of red lists: RE – Regionally extinct, CR – Critically Endangered, EN – Endangered, VU – Vulnerable, NT – Near Threatened, LC – Least Concern, DD – Data Deficient, R – rare; poorly developed specimens are marked with \*

Table 3 (continued)

Lichen taxa	New R Mts	New RO	RL RO	RL UK	RL HU	RL PL	RL SL	Obs	Plot codes
<i>Calicium cf. salicinum</i> * Pers.					EN	VU	EN	1	MC6
<i>Candelariella reflexa</i> (Nyl.) Lettau	1				R			4	MC6, MM4, WM5, 7
<i>Candelariella vitellina</i> (Ehrh.) Müll. Arg.	1							3	MM5, WM5, 6
<i>Catillaria nigroclavata</i> (Nyl.) J. Steiner	1				R			1	WM7
<i>Cetraria aculeata</i> (Schreb.) Fr.					CR			2	AC3,5
<i>Cetraria ericetorum</i> Opiz						NT	VU	6	AC3,5, 6,7, AM2, 5
<i>Cetraria islandica</i> (L.) Ach.			VU		EN	VU	VU	24	PM1, 2, 3, 4, 5, 6, 7, PM2, 3, 4, 5, 6, 7, WM5, AC1, 2, 3, 4, 5, 6, 7, CC3, 4, AM11, 2, 3, 4, 5, 6, 7
<i>Cetraria muricata</i> (Ach.) Eckfeldt	1	1				NT		1	AC3
<i>Cetraria sepincola</i> (Ehrh.) Ach.	1		EN		R	EN	VU	1	PM7
<i>Cetrelia olivetorum</i> (Nyl.) W.L. Culb. & C.F. Culb.	1				VU	EN		2	MC2, 3
<i>Chaenotheca brachypoda</i> (Ach.) Tibell	1	1				EN	CR	1	MC1
<i>Chaenotheca brunneola</i> (Ach.) Müll. Arg.	1				CR	EN	CR	2	CC1, CM7
<i>Chaenotheca chrysoccephala</i> (Ach.) Th. Fr.	1				VU		VU	5	MC1, CC1, 4, 5, 6
<i>Chaenotheca ferruginea</i> (Turner ex Sm.) Mig.	1				EN			1	CC4
<i>Chaenotheca furfuracea</i> (L.) Tibell	1				EN	NT	NT	4	MC6, MM3, 7, CC1
<i>Chaenotheca gracilenta</i> (Ach.) Mattsson & Middelb.	1						CR	1	MM3
<i>Chaenotheca stemonea</i> (Ach.) Müll. Arg.	1	1				EN	CR	1	CC6
<i>Chaenotheca trichialis</i> (Ach.) Hellb.	1				EN	NT	CR	3	MM4, CC1,5
<i>Chrysothrix candelaris</i> (L.) J.R. Laundon	1						CR	2	MC2, 3
<i>Cladonia arbuscula</i> (Wallr.) Flot.					VU			12	AC1, 2, 3, 4, 5, 6, 7, AM2, 3, 4, 5, 6
<i>Cladonia bellidiflora</i> (Ach.) Schaer.	1					EN		8	AC4, 7, AM1, 2, 3, 4, 5, 6
<i>Cladonia caespiticia</i> (Pers.) Flörke						EN		3	AC1, CC6, CM5
<i>Cladonia callosa</i> Delise ex Harm.	1	1						1	AC2
<i>Cladonia carneola</i> (Fr.) Fr.	1					CR	EN	5	PM6, WM5, CC2, 3, AM2
<i>Cladonia cenotea</i> (Ach.) Schaer.					VU			11	MC2, MM2, 3, PM1, 2, WM1, 3, 6, CC1, 2, CM5
<i>Cladonia cervicornis</i> (Ach.) Flot.								8	AC1, 2, 3, 4, 6, 7, AM1,3
<i>Cladonia chlorophaea</i> (Flörke ex Sommerf.) Spreng.	1							34	MC5, 6, 7, MM2, 6, PM1, 2, 3, 4, 5, 6, 7, WM1, 3, 4, 5, 6, 7, AC1, 2, 3, 4, 5, 6, 7, CC2, 3, 5, 6, 7, CM5, AM1, 3, 4, 7
<i>Cladonia coccifera</i> (L.) Willd.					EN			16	AC1, 2, 3, 4, 5, 6, 7, CC3, AM1, 2, 3, 4, 5, 6, 7, PM7
<i>Cladonia coniocraea</i> (Flörke) Spreng.								28	MC1, 2, 3, 4, 5, 6, 7, MM2, 3, 4, 5, 6, 7, PM3, WM1, 3, 4, 6, 7, CC1, 3, 5, 6, 7, CM5, 6, AC5, AM3
<i>Cladonia cornuta</i> (L.) Hoffm.					VU		VU	13	MC2, 4, 5, 7, MM2, 5, 6, PM1, 3, WM3, CC1, CM5, AM7
<i>Cladonia deformis</i> (L.) Hoffm.								4	PM1, CC3, CC5, AM2
<i>Cladonia digitata</i> (L.) Hoffm.								41	MC1, 2, 3, 4, 5, 6, 7, MM2, 3, 4, 6, PM1, 2, 3, 4, 5, 6, WM1, 2, 3, 6, AC1, 2, 3, 4, 5, 6, CC1, 2, 3, 4, 5, 6, 7, CM1, 2, 5, AM1, 2, 3, 7
<i>Cladonia diversa</i> Asperges	1	1						3	AC3, AC4, AM1
<i>Cladonia fimbriata</i> (L.) Fr.								15	MC4, 5, 7, MM2, 5, 6, WM1, 2, 4, 5, 6, 7, CC3, 7, CM5,

Table 3 (continued)

Table 3 (continued)

Lichen taxa	New R Mts	New RO	RL RO	RL UK	RL HU	RL PL	RL SL	Obs	Plot codes
<i>Cladonia uncialis</i> (L.) Weber ex F.H. Wigg.	1				VU			7	AC3, AC4, AM1, AM2, AM4, AM5, AM6
<i>Collema flaccidum</i> (Ach.) Ach.					EN	EN		3	MC3, 4, 6
<i>Cornicularia normoerica</i> (Gunnerus) Du Rietz					VU			7	PM5, AC1, 3, 4, 5, 7, AM1
<i>Dibaeis baeomyces</i> (L. f.) Rambold & Hertel					NT			1	AC3
<i>Dimerella pineti</i> (Ach.) Vězda	1							6	MM2, 4, CC6, CM4, 6, 7
<i>Ephabe lanata</i> (L.) Vain.					EN			1	AC4
<i>Evernia divaricata</i> (L.) Ach.					CR	CR		4	MC4, MM3, 5, CC4
<i>Evernia prunastri</i> (L.) Ach.					NT	EN		11	MC2, 3, 4, 5, 6, 7, MM1, WM2, 3, CC1, 6
<i>Fellhaneropsis vezdae</i> (Coppins & P. James Sérus. & Coppins	1	1			LC			1	MM4
<i>Flavocetraria cucullata</i> (Bellardi) Kärnefelt & A. Thell					VU			2	AM2, 4
<i>Fuscidea arboricola</i> Coppins & Tønsberg	1	1						8	MM2, 3, CC2, 5, 6, 7, CM6
<i>Fuscidea pusilla</i> Tønsberg	1	1						7	MC6, MM2, WM5, CC5, 7, CM3, 4
<i>Graphis pulverulenta</i> (Pers.) Ach.	1							5	MC1, 3, 7, MM2, 7,
<i>Graphis scripta</i> (L.) Ach.					NT	EN		5	MC1, 3, 4, 5, 6
<i>Heterodermia speciosa</i> (Wulfen) Trevis.	1			R	CR	CR		1	MC5
<i>Hypocenomyce scalaris</i> (Ach. ex Lilj.) M. Choisy								1	WM7
<i>Hypogymnia farinacea</i> Zopf	1				VU	VU		4	WM3, 7, CC6, CM5,
<i>Hypogymnia physodes</i> (L.) Nyl.								39	MC1, 2, 3, 4, 5, 6, 7, MM1, 2, 3, 4, 5, 6, 7, PM2, 3, 4, 6, WM1, 2, 3, 4, 5, 6, 7, AC3, CC1, 2, 3, 4, 5, 6, 7, CM1, 2, 3, 4, 5, 6
<i>Hypogymnia tubulosa</i> (Schaer.) Hav.					VU	NT	NT	14	MC3, 5, 7, MM3, PM4, WM1, 2, 3, 4, 6, 7, CC3, 4, CM6
<i>Hypogymnia vittata</i> (Ach.) Parrique					CR	VU		1	AC6
<i>Icmadophila ericetorum</i> (L.) Zahlbr.	1				EN	CR		2	AC3, CC3
<i>Imshaugia aleurites</i> (Ach.) S.L.F. Mey.						VU		1	WM4
<i>Japewia subaurifera</i> Muhr & Tønsberg	1	1						2	MM1, CC2
<i>Lecanactis abietina</i> (Ehrh. ex Ach.) Körb.	1					CR		1	CC1
<i>Lecanora albella</i> (Pers.) Ach.					EN	CR		1	MC4
<i>Lecanora albellula</i> (Nyl.) Th.	1	1						2	WM2, 4
<i>Lecanora argentata</i> (Ach.) Malme	1							4	MC3, 4, 5, 6
<i>Lecanora carpinea</i> (L.) Vain.	1							8	MC2, 3, 4, 5, 6, 7, MM5, WM2
<i>Lecanora cf. hybocarpa</i> (Tuck.) Brodo								1	MC5
<i>Lecanora chlarotera</i> Nyl.	1							17	MC1, 2, 6, 7, MM2, 3, 4, 5, 6, 7, WM2, 3, 4, 5, 6, 7, CC4
<i>Lecanora cinereofusca</i> H. Magn.	1				CR			3	MC3, 4, 5
<i>Lecanora circumborealis</i> Brodo & Vitik.	1	1						4	MC6, WM3, 4, 5
<i>Lecanora compallens</i> Herk & Aptroot	1	1						2	AC3, 5
<i>Lecanora conizaeoides</i> Nyl. ex Cromb.	1							10	MM3, WM1, CC3, 4, 5, 6, CM3, 4, 5, 6
<i>Lecanora expallens</i> Ach.	1							4	MC1, 4, CC1, 5
<i>Lecanora glabrata</i> (Ach.) Malme	1							1	WM5
<i>Lecanora intumescens</i> (Rebent.) Rabenh.					EN	EN		6	MC1, 2, 4, 6, 7, CC2

Table 3 (continued)

Lichen taxa	New Mts	R	New RO	RL RO	RL UK	RL HU	RL PL	RL SL	Obs	Plot codes
<i>Lecanora pulicaris</i> (Pers.) Ach.	1								23	MC1, 2, 5, MC6, 7, PM5, 6, 7, WM1, 2, 3, 4, 5, 6, 7, CC2, 3, 6, 7, CM3, 4, 5, 6
<i>Lecanora saligna</i> (Schrad.) Zahlbr.	1								1	WM3
<i>Lecanora strobilina</i> (Spreng.) Kieff.	1	1							7	MC6, MM2, 4, 5, PM5, WM2, 7
<i>Lecanora subintricata</i> (Nyl.) Th. Fr.	1								4	PM6, WM2, 4, 7
<i>Lecanora subrugosa</i> Nyl.	1								LC	1 MM4
<i>Lecanora symmicta</i> (Ach.) Ach.	1								5	PM2, 5, 6, WM2, 4
<i>Lecanora umbrina</i> (Ehrh.) Röhl	1									WM7
<i>Lecanora varia</i> (Hoffm.) Ach.	1						R	VU	4	WM2, 3, 4, 5
<i>Lecidea berengeriana</i> (A. Massal.) Nyl.	1								1	AC5
<i>Lecidea leprariooides</i> Tønsberg	1	1							1	CC1
<i>Lecidea nylanderi</i> (Anzi) Th. Fr.	1	1							24	MC1, 2, 3, 4, 5, 7, MM1, 2, 4, 5, 7, PM1, WM5, CC1, 2, 3, 4, 5, 6, 7, CM1, 2, 3, 4
<i>Lecidella cf. albida</i> Hafellner									1	MC3
<i>Lecidella elaeochroma</i> (Ach.) M. Choisy									9	MC1, 2, 3, 4, 5, 6, 7, MM1, 4
<i>Lecidella elaeochroma f. soralifera</i> (Erichsen) D. Hawksw.	1	1							1	CC2
<i>Lecidella subviridis</i> Tønsberg	1	1							10	MC5, 6, 7, MM1, 2, 7, CC2, 3, 6, 7
<i>Lecidoma demissum</i> (Rutstr.) Gotth. Schneid. & Hertel							EN		5	AC2, 4, 7, AM1, 3
<i>Lepraria eburnea</i> J.R. Laundon	1	1							11	MC1, 6, 7, MM1, 2, 4, 5, 7, CM1, 4, 7
<i>Lepraria ecorticata</i> (J.R. Laundon) Kukwa	1	1							2	PM4, CC1
<i>Lepraria elobata</i> Tønsberg	1	1							24	MC2, 3, 4, 5, 6, MM6, 7, PM1, 2, 3, WM6, AC2, 4, CC2, 3, 4, 5, 6, 7, CM2, 3, 4, 5, 6
<i>Lepraria incana</i> (L.) Ach.	1								1	MC3
<i>Lepraria jackii</i> Tønsberg	1	1							17	MC1, 2, MM1, 3, 7, PM5, WM5, AC5, CC2, 3, 4, 5, 6, CM1, 3, 4, 7
<i>Lepraria lobificans</i> Nyl.	1					R			32	MC1, 2, 3, 4, 5, 6, 7, MM1, 2, 3, 4, 5, 6, 7, PM2, 3, 7, WM1, 4, AC2, CC1, 2, 3, 4, 5, 6, 7, CM2, 3, 4, 6, 7
<i>Lepraria membranacea</i> (Dicks.) Vain.	1								2	MM2, CC1
<i>Lepraria neglecta</i> Vain.	1	1							1	AC4
<i>Lepraria nivalis</i> J.R. Laundon	1	1							11	PM2, 5, WM2, AC2, 3, 4, 5, 6, 7, CC2, AM7
<i>Lepraria rigidula</i> (B. de Lesd.) Diederich	1	1							7	MC2, 6, MM1, WM2, CC1, 5, CM3
<i>Lepraria toensbergiana</i> Slav.-Bay. & Kukwa	1	1							8	MC7, MM3, PM3, CC1, 3, 7, CM3, 4,
<i>Lepraria umbricola</i> Tønsberg	1	1							3	MM1, CM2, 7
<i>Lepraria vousauxii</i> (Hue) RC. Harris	1	1				R			3	MM2, CM1, 7
<i>Leprocaulon microscopicum</i> (Vill.) Gams	1								2	MC5, CC4
<i>Leptogium lichenoides</i> (L.) Zahlbr.							LC		5	MC3, 6, MM5, 6, 7
<i>Lichenomphalia hudsoniana</i> (H.S. Jenn.) Redhead et al.	1				R		NT	VU	5	PM5, AC3, CC6, CM6, AM2
<i>Lichenomphalia umbellifera</i> (L.) Redhead et al.	1	1					NT		3	CM3, 6

Table 3 (continued)

Lichen taxa	New R Mts	New RO	RL RO	RL UK	RL HU	RL PL	RL SL	Obs	Plot codes
<i>Lichenomphalia velutina</i> (Quél.) Redhead et al.	1	1						1	MC1
<i>Lobaria pulmonaria</i> (L.) Hoffm.			VU	VU	RE	EN	CR	2	MC3, MM1
<i>Loxospora cismonica</i> (Beltr.) Hafellner	1					CR	RE	2	MC1, 5,
<i>Loxospora elatina</i> (Ach.) A. Massal.	1					EN	NT	16	MC1, 2, 3, 4, 5, 6, 7, MM1, 2, CC1, 4, 5, 6, CM2, 4, 5
<i>Megalospora tuberculosa</i> (Fée) Sipman	1	1						1	MC7
<i>Melanelia hepatizon</i> (Ach.) A. Thell							NT	4	PM6, AC4, 6, AM5
<i>Melanelia stygia</i> (L.) Essl.	1						NT	3	PM2, AC3, 5
<i>Melanelixia fuliginosa</i> subsp. <i>glabratula</i> (Lamy) J.R. Laundon								23	MC1, 2, 3, 4, 5, 7, MM1, 2, 3, 4, 5, 7, WM1, 2, 4, 5, 6, 7, CC4, 7, CM6, 7
<i>Melanelixia subargentifera</i> (Nyl.) O. Blanco et al.	1					VU	VU	2	WM2, 3
<i>Melanelixia subaurifera</i> (Nyl.) O. Blanco et al.	1		VU				NT	11	MC2, 3, 4, 5, MM2, 5, WM2, 3, 5, 6, 7,
<i>Melanohalea elegantula</i> (Zahlbr.) O. Blanco et al.	1	1		R		VU		2	MC6, MM1
<i>Melanohalea exasperatula</i> (Nyl.) O. Blanco et al.	1							6	MC6, WM2, 3, 4, 5, 7
<i>Melaspilea ochrothalamia</i> Nyl.	1	1						2	PM1, PM5
<i>Melaspilea proximella</i> (Nyl.) Nyl.	1							1	PM4
<i>Menegazzia terebrata</i> (Hoffm.) A. Massal.	1			CR	CR	CR		1	MC4
<i>Micarea elachista</i> (Körb.) Coppins & R. Sant.	1	1				EN		2	WM3
<i>Micarea farinosa</i> Coppins & Aptroot	1	1						1	CM2
<i>Micarea lignaria</i> (Ach.) Hedl.				RE				5	PM2, AC3, 5, 6, AM1
<i>Micarea melaena</i> (Nyl.) Hedl.	1			R	NT			1	CM4
<i>Micarea micrococca</i> (Körb.) Gams	1	1						1	MM4,
<i>Micarea misella</i> (Nyl.) Hedl.	1			R				2	MC2, MM5
<i>Micarea prasina</i> Fr.	1							23	MC1, 4, 5, 7, MM1, 3, 4, 5, 6, 7, PM7, CC1, 2, 3, 4, 6, 7, CM2, 3, 4, 5, 6, 7
<i>Micarea turfosa</i> (A. Massal.) Du Rietz	1	1				DD		1	AC2
<i>Micarea viridileprosa</i> Coppins & Van den Boom	1	1						1	MC1
<i>Micarea xanthonica</i> Coppins & Tønsberg	1	1						1	MM5
<i>Mycobilimbia epixanthoides</i> (Stirt.) Zahlbr.	1	1						10	MC5, 7, MM4, 5, 6, 7, AC2, CC6, 7, CM7
<i>Mycoblastus fucatus</i> (Stirt.) Zahlbr.	1	1						6	MC2, 4, 5, 7, MM3, CC5,
<i>Mycoblastus sanguinarius</i> (L.) Norman	1			VU	CR			1	MC2
<i>Nephroma parile</i> (Ach.) Ach.	1		VU	CR	CR			1	CC2
<i>Ochrolechia androgyna</i> (Hoffm.) Arnold	1			VU				3	MC6, CC2, AM4
<i>Opegrapha niveoatra</i> (Borrer) J.R. Laundon	1							1	MM2
<i>Opegrapha varia</i> Pers.					NT	VU		4	MC1, 3, 5, 6
<i>Opegrapha viridis</i> Pers.			R	VU	VU			3	MC1, 3, 4,
<i>Opegrapha vulgata</i> (Ach.) Ach.				VU	VU			2	MC1, 5,
<i>Parmelia omphalodes</i> (L.) Ach.	1				EN	NT		4	PM2, AC3, CC2, AM1,
<i>Parmelia saxatilis</i> (L.) Ach.						NT		12	PM6, WM2, 4, AC3, 4, 5, CC2, 4, AM2, 4, 5, 6
<i>Parmelia submontana</i> Nádv.		1			VU	CR		6	MC2, 3, 6, WM4, 5, 7,

Table 3 (continued)

Lichen taxa	New Mts	R	New RO	RL RO	RL UK	RL HU	RL PL	RL SL	Obs	Plot codes	
<i>Parmelia sulcata</i> Taylor									10	MC2, 4, 6, PM6, WM2, 3, 4, 6, 7, CC5	
<i>Parmelia quercina</i> (Willd.) Hale	1						CR	CR	2	WM2, 3	
<i>Parmeliopsis ambigua</i> (Wulfen) Nyl.									18	MC2, PM1, 2, 3, 4, 6, 7, WM1, 3, 4, 6, 7, CC1, 2, 3, 6, 7, CM5	
<i>Parmeliopsis hyperopta</i> (Ach.) Vain.							VU	VU	6	MC4, PM1, 4, 6, 7, CC4	
<i>Peltigera canina</i> (L.) Willd.							VU		3	WM3, 5, 6	
<i>Peltigera degenerii</i> Gyeln.	1						EN	VU	3	MC3, 5, MM5	
<i>Peltigera horizontalis</i> (Huds.) Baumg.							EN		6	MC2, 3, 5, MM5, WM3, 4,	
<i>Peltigera hymenina</i> (Ach.) Delise	1	1					EN	DD	10	MC4, 5, 6, MM1, 3, 4, WM2, 3, CC6, CM2	
<i>Peltigera lepidophora</i> (Nyl.) Bitter	1			CR			EN	CR	1	MM2	
<i>Peltigera membranacea</i> (Ach.) Nyl.	1						DD		1	WM5	
<i>Peltigera polydactylon</i> (Neck.) Hoffm.	1						EN	DD	4	MC1, MM2, 4, WM5,	
<i>Peltigera praetextata</i> (Flörke ex Sommerf.) Zopf	1						VU		8	MC1, 2, 3, 4, MM4, 5, 6, 7,	
<i>Peltigera rufescens</i> (Weiss) Humb.	1								3	WM3, 5, 7	
<i>Pertusaria amara</i> (Ach.) Nyl.								NT	2	MC2, 5	
<i>Pertusaria coccodes</i> (Ach.) Nyl.	1	1					R	NT	1	MM4	
<i>Pertusaria leioplaca</i> DC.							R	NT	4	MC1, 2, 5, 6,	
<i>Pertusaria oculata</i> (Dicks.) Th. Fr.	1								1	AC5	
<i>Pertusaria pertusa</i> (L.) Tuck.							VU	NT	1	MC3	
<i>Pertusaria pupillaris</i> (Nyl.) Th. Fr.	1	1						NT	1	MC6	
<i>Phaeophyscia endophoenicea</i> (Harm.) Moberg	1	1					R	EN	1	MC6	
<i>Phaeophyscia orbicularis</i> (Neck.) Moberg	1								3	MM4, WM3, 4	
<i>Pblyctis argena</i> (Ach.) Flot.	1								6	MC2, 3, 4, 6, MM1, 6,	
<i>Physcia adscendens</i> (Fr.) H. Olivier	1								6	MM4, WM2, 3, 4, 6, 7,	
<i>Physcia aipolia</i> (Ehrh. ex Humb.) Fürnr.	1							NT	3	WM2, 5, 7	
<i>Physcia dubia</i> (Hoffm.) Lettau	1						R		1	WM2	
<i>Physcia leptalea</i> (Ach.) DC.	1	1							1	MM5	
<i>Physcia stellaris</i> (L.) Nyl.									5	MC7, WM2, 3, 4, 6	
<i>Physcia tenella</i> (Scop.) DC.	1								4	WM2, 3, 4, 6,	
<i>Placynthiella dasaea</i> (Stirt.) Tønsberg	1	1					R		12	MC1, 2, 7, MM2, 3, 4, 7, WM7, CC2, 4, CM5, AM5	
<i>Placynthiella icmalea</i> (Ach.) Coppins & P. James	1	1					R		16	MC2, 5, 7, MM3, 5, 6, 7, PM4, 5, WM1, 3, CM5, 6, AM5, 6, 7	
<i>Placynthiella uliginosa</i> (Schrad.) Coppins & P. James	1						R		3	PM3, WM6, AM7	
<i>Platismatia glauca</i> (L.) W.L. Culb. & C.F. Culb.							R	NT	15	MC2, 3, 4, MM1, 3, WM2, 6, AC3, CC1, 4, 5, 6, CM2, 5, 6	
<i>Porina aenea</i> (Wallr.) Zahlbr.	1						R		4	MC1, MM1, 6, 7	
<i>Porina lectissima</i> (Fr.) Zahlbr.	1								1	MM1	
<i>Pseudephbe pubescens</i> (L.) M. Choisy							EN		11	PM2, 6, AC1, 3, 4, 5, 6, 7, AM1, 2, 3	
<i>Pseudevernia furfuracea</i> (L.) Zopf								NT	37	MC2, 3, 4, 5, 6, 7, MM1, 2, 3, 5, 6, 7, PM1, 2, 3, 4, 5, 6, WM1, 2, 3, 4, 5, 6, 7, AC3, CC1, 2, 3, 4, 6, 7, CM1, 2, 4, 5, AM1	
<i>Pycnothelia papillaria</i> (Ehrh.) L.M. Dufour							EN	EN	VU	1	AC4
<i>Pyrenula nitida</i> (Weigel) Ach.							VU	EN	5	MC1, 2, 3, 4, MM2	

Table 3 (continued)

Lichen taxa	New R Mts	New RO	RL RO	RL UK	RL HU	RL PL	RL SL	Obs	Plot codes
<i>Pyrenula nitidella</i> (Flörke ex Schaer.) Müll. Arg.	1				R	EN	CR	1	MC3
<i>Ramalina canariensis</i> J. Steiner	1				R			3	MC2, 3, 6
<i>Ramalina farinacea</i> (L.) Ach.					VU	EN		3	MC2, 4, 5
<i>Ramalina fastigiatoides</i> (Pers.) Ach.					EN	EN		1	MC2
<i>Ramalina pollinaria</i> (Westr.) Ach.					VU	VU		1	MC3
<i>Rinodina sophodes</i> (Ach.) A. Massal.					EN	VU		3	WM2, 3, 5
<i>Ropalospora viridis</i> (Tønsberg) Tønsberg								2	MC2, 5
<i>Scoliciosporum chlorococcum</i> (Graewe ex Stenf.) Věžda	1							12	MC7, MM2, 4, 5, 7, WM3, 5, 6, 7, CC7, CM3, 5
<i>Scoliciosporum sarothonii</i> (Vain.) Věžda	1	1						7	MC6, MM3, 5, 7, WM5, 6, 7
<i>Scoliciosporum umbrinum</i> (Ach.) Arnold								4	MM2, 4, 5, WM5,
<i>Sphaerophorus fragilis</i> (L.) Pers.					EN	VU		4	PM2, AC3, 4, 6
<i>Stereocaulon alpinum</i> Laurer			VU		EN	VU		1	AC3
<i>Strigula jamesii</i> (Swinscow) R.C. Harris	1	1			VU	LC		1	MC5
<i>Thamnolia vermicularis</i> (Sw.) Schaer.								14	PM5, 7, AC1, 2, 3, 4, AC5, 6, 7, AM1, 2, 3, 4, 5
<i>Thelotrema lepadinum</i> (Ach.) Ach.	1					EN	CR	2	MC3, 4
<i>Trapelia corticola</i> Coppins & P. James	1	1						1	PM1
<i>Trapeliopsis flexuosa</i> (Fr.) Coppins & P. James								13	MM2, 3, 6, WM5, 6, AC5, 7, CC3, 4, 5, 7, CM6, 7
<i>Trapeliopsis gelatinosa</i> (Flörke) Coppins & P. James	1					NT		3	MC1, CC2, AM5
<i>Trapeliopsis granulosa</i> (Hoffm.) Lumbsch	1							22	MC7, PM1, 3, 4, 5, 6, AC1, 2, 3, 4, 5, 6, 7, CC3, CM2, AM1, 2, 3, 4, 5, 6, 7
<i>Trapeliopsis pseudogranulosa</i> Coppins & P. James	1	1						2	PM5, CC1
<i>Trapeliopsis viridescens</i> (Schrad.) Coppins & P. James	1				NT			4	MC1, 7, AC7, CC6
<i>Tuckermannopsis chlorophylla</i> (Willd.) Hale			VU		EN	VU	NT	2	MC2, WM1
<i>Umbilicaria crustulosa</i> (Ach.) Lamy	1				EN	NT		2	AC7, CC4
<i>Umbilicaria cylindrica</i> (L.) Delise					RE			28	PM1, 2, 3, 4, 5, 6, 7, WM1, 4, 6, AC1, 2, 3, 4, 5, 6, 7, CC2, 3, 4, 5, AM1, 2, 3, 4, 5, 6, 7
<i>Umbilicaria deusta</i> (L.) Baumg.					CR	LC		11	AC1, 2, 4, 6, CC4, AM1, 2, 4, 5, 6, 7
<i>Umbilicaria polyphylla</i> (L.) Baumg.					CR	LC		6	PM6, WM1, AC6, AM1, 3, 6
<i>Umbilicaria vellea</i> (L.) Ach.					EN	VU		1	CC4
<i>Usnea chaetophora</i> Stirt.	1	1						2	MC2, CC4
<i>Usnea diplotypus</i> Vain.	1	1			CR	RE		2	CC4, 5
<i>Usnea filipendula</i> Stirt.					CR	VU	CR	5	MC4, CC1, 3, 4, 5, CM2,
<i>Usnea florida</i> (L.) Weber ex F.H. Wigg.			VU		CR	CR	CR	1	MC3
<i>Usnea fulvoreagens</i> (Räsänen) Räsänen	1		EN		CR	CR	RE	1	MC4
<i>Usnea glabrescens</i> (Nyl. ex Vain.) Vain.	1							1	CC4
<i>Usnea hirta</i> (L.) Weber ex F.H. Wigg.					VU	VU		5	WM2, 3, 4, 5, CC6
<i>Usnea lapponica</i> Vain.	1							1	MC4
<i>Usnea scabrata</i> Nyl.	1							1	MM1
<i>Usnea subfloridana</i> Stirt.					CR	EN	CR	4	MC2, 4, 6, WM5
<i>Vulpicida pinastri</i> (Scop.) J.-E. Mattsson	1				NT	NT		11	PM4, 6, 7, WM2, 4, 6, 7, CC4, 5, 6, 7

Table 3 (continued)

Lichen taxa	New Mts	R	New RO	RL RO	RL UK	RL HU	RL PL	RL SL	Obs	Plot codes
<i>Xanthoparmelia conspersa</i> (Ehrh. ex Ach.) Hale									1	WM4
<i>Xanthoparmelia somloensis</i> (Gyeln.) Hale	1								1	WM2
<i>Xanthoria candelaria</i> (L.) Th. Fr.	1								2	WM4, 7
<i>Xanthoria cf. ulophylloides</i> * Räsänen				VU	VU				1	WM2
<i>Xanthoria parietina</i> (L.) Beltr.					NT				5	WM2, 3, 4, 5, 6
<i>Xanthoria polycarpa</i> (Hoffm.) Rieber	1				NT				1	WM5, 7
<i>Xylographa parallela</i> (Ach.) Fr.					EN				3	MC2, WM4, CC1
<i>Xylographa vitiligo</i> (Ach.) J.R. Laundon	1				DD	EN			2	MM3, PM7
Totals	182	67	12	8	65	125	96			

Several inconspicuous taxa (*Micarea* sp., *Placynthiella* sp., *Scoliosporum* sp., *Lichenomphalia umbelifera*, *Lichenomphalia velutina*) possibly overlooked in previous surveys, were found mainly because of the systematic sampling protocol used here, which forced us to consider all lichens present in small sampling plots.

Another group of neglected species in Romania includes taxa that only recently have received a thorough taxonomic treatment that makes identification possible in routine lichen diversity assessments. Among these is the genera *Lepraria* for which we found 13 species, all new for the Rodnei Mountains, and 10 species new for the country.

The numerous new species for the region is mainly due to the fact that the previous studies focused on rocks, whereas substrates such as trees, soil, and wood, were poorly investigated. For example, there are only 49 previously reported species on soil substrate versus the 108 taxa that we report here. We only confirm 158 out of the 260 species present in the previous checklist, which was represented mainly by saxicolous lichens (ca 57 % of total) (APNMR, 2013), including crustose, foliose and fruticose species, whereas in our survey the crustose saxicolous lichens were not sampled.

We found a relatively high number of species that are restricted to conserved habitats in Rodnei Mountains and which are known as indicators of ancient woodland and ecological continuity in forest landscapes (Rose, 1976; Goward, 1995; Wolseley, 1995; Thor, 1995; Gauslaa & Solhaug, 1996): *Arthonia caesia*, *Bryoria lanestris*, *Cetrelia olivetorum*, *Chaenotheca brachypoda*, *Ch. brunneola*, *Heteroder-*

*mia speciosa*, *Lecanora cinereofusca*, *Lobaria pulmonaria*, *Loxospora cismonica*, *Megalospora tuberculosa*, *Menegazzia terebrata*, *Pertusaria coccodes*, *Thelotrema lepadinum*, *Usnea florida* and *U. subfloridana*.

Also species from Red Lists, which depend on rare and often threatened habitats (Thor, 1995), such as *Anisomeridium biforme* and *Usnea fulvoreagens*, are extinct in some regions of the Carpathians (Pisút et al., 2001) but were recorded (once each) in conserved mixed forests in the present study of the Rodnei Mountains. Other examples of species limited to conserved areas, having Critically Endangered status in the Red Lists and being recorded once or twice in our study include: *Arthonia vinosa*, *Bellemerea cinereofuscens*, *Bryoria capillaris*, *B. chalibiformis*, *Cetraria aculeata*, *Cladonia magyarica*, *Hypogymnia vittata*, *Icmadophila ericetorum*, *Lecanactis abietina*, *Lecanora albella*, *Mycoblastus sanguinarius*, *Nephroma parile*, *Phaeophyscia endophoenicea* and *Pyrenula nitidella* (Table 3). All these species stress the importance of maintaining their suitable habitats, which are currently restricted to protected areas.

The present study, with its large number of new species for Romanian lichen flora and for the Rodnei Mountains region, suggests the need of more detailed inventories. To this day, there is no official Red List of lichens for Romania, except a manuscript dealing with macrolichens only (Bartok & Crișan, personal communication). In this context lichen diversity and richness inventories are important for better estimating the degree of threat for each species. This will enable the compilation of a comprehensive Red List of Lichens from Romania that will be an invaluable

able tool for promoting lichen conservation in the country (Thor, 1995; Dietrich & Scheidegger, 1997; Scheidegger et al., 2000).

The high total number of lichen species, the large number of species that appear in Red Lists, together with the indicator species for ancient woodlands and ecological continuity of forest landscapes, underline the high value of conserved habitats in the Rodnei Mountains. The existing National Park here has the potential of conserving a considerable part of the lichen diversity and the lichen's characteristic habitats. Their presence is also important for other groups of organisms, which depend on lichens or their habitats, thus emphasizing the biocomplexity of microbial and invertebrate communities. It is important that at least these „islands” such as Rodnei Mountains to be conserved if otherwise natural resources are still exploited in a non-sustainable manner that is leading to a substantial loss of biodiversity.

## ACKNOWLEDGEMENTS

This work was possible with the financial support of the Sectoral Operational Programme for Human Resources Development 2007-2013, co-financed by the European Social Fund, under the project number POSDRU/107/1.5/S/76841 with the title „Modern Doctoral Studies: Internationalization and Interdisciplinarity”. We also acknowledge financial support from the Federal Office for the Environment (FOEN) to Christoph Scheidegger.

## REFERENCES

- APNMR (Rodnei Mountains National Park Administration). *Management Plan 2013 Rodnei Mountains National Park, Biosphere Reserve (NATURA 2000 SAC and SPA)* (In Romanian). <http://www.parcrodna.ro/>.
- Bartók, K. 1999. Pesticide usage and epiphytic lichen diversity in Romanian orchards. *The Lichenologist* 31(1): 21–25.
- Cieslinski, S., Czyzewska K. & Fabiszewski, J. 2003. Czerwona lista porostów wymarlych i zagrożonych w Polsce. Red List of extinct and threatened lichens in Poland. – In: Czyzewska, K. (ed.). *Zagrożenie Porostów w Polsce. The threat to lichens in Poland. Monographiae Botanicae* 91: 13–49.
- Ciurchea, M. 2004. *Key to Lichens from Romania* (In Romanian). Ed. Bit, Iași. 471 pp.
- Çobanoğlu, G., Yavuz, M., Costache, I. & Radu, I. 2011. Additional and new lichen records from Cozia National Park, Romania. *Mycotaxon* 114: 193–196. <http://dx.doi.org/10.5248/114.193>
- Çobanoğlu, G., Yavuz, M., Costache, I., Radu, I., Açıkgöz, B. & Baloniu, L. 2009. Epiphytic and terricolous lichens diversity in Cozia National Park (Romania). *Muzeul Olteniei Craiova. Oltenia. Studii și comunicări. Științele Naturii* 25: 17–22.
- Crișan, F. 2006. Five new species for the Romanian lichen flora. In: Gafta, D. & Akeroyd, J. (eds). *Nature Conservation, Concepts and Practice. Series: Environmental Science and Engineering*, pp. 131–136.
- Didukh, Y. A. (ed.) 2009. *Red data book of Ukraine, Vegetable kingdom* (In Ukrainian). Globalconsalt-ing. 912 pp.
- Dietrich , M. & Scheidegger, C. 1997. A representative survey of frequency of epiphytic lichens at the regional and national levels and its use for the Red List of Switzerland. In: Türk, R. & Zorrer, R. (eds). *Progress and Problems in Lichenology in the Nineties – IAL3. Bibliotheca Lichenologica* 68: 145–154.
- Gauslaa, Y. & Solhaug, K. A. 1996. Differences in the susceptibility to light stress between epiphytic lichens of ancient and young boreal forest stands. *Functional Ecology* 10(3): 344–354. <http://dx.doi.org/10.2307/2390282>
- Gorduza V. 1983. Physico-geographical caracterisation of the Pietrosu Rodnei Nature Reserve. In: Pietrosul Rodnei at 50 years (In Romanian). *Academia RSR Cluj Napoca-Baia Mare*, pp: 56–66.
- Goward, T. 1995. *Nephroma occultum* and the maintenance of lichen diversity in British Columbia. *Mitteilungen der Eidgenössischen Forschungsanstalt für Wald, Schnee und Landschaft* 70: 11–27.
- Ignatov, M.S., Ignatova, E.A., Konstantinova, N. A., Pronkina, G. A. & Urbanavichus, G. P. 2004. Lichens / The present-day state of biological diversity within protected areas. Issue 3. *Lichens and Bryophytes*. Moscow, pp 6–7.
- Lackovičová, A. & Guttová, A. 2006. Lichen diversity—history, contemporary occurrence and trends in Slovakia: *Gyalecta ulmi* and *Leptogium saturninum*. In: Lackovičová, A., Guttová, A., Lisická, E. & Lizoň, P. (eds). *Central European Lichens – Diversity and Threat. Mycotaxon Ltd., Ithaca, and Institute of Botany, Slovak Academy of Sciences*, Bratislava, pp. 219–240.
- Lökös, L. & Tóth, E. 1996. Red List of lichens of Hungary (a proposal). *Vegetation Mapping and Botanical Research On Nature Conservation Purpose* 7: 337–343.
- Nascimbene, J., Thor, G. & Nimis, P. L. 2013. Effects of forest management on epiphytic lichens in temperate deciduous forests of Europe—A review. *Forest Ecology and Management* 298: 27–38. <http://dx.doi.org/10.1016/j.foreco.2013.03.008>
- Pisút, I., Guttová, A., Lackovicová, A. & Lisická, E. 2001. Cerveny zoznam lisajníkov Slovenska (December 2001) [Red List of lichens of Slovakia

- (December 2001)] *Ochrana Prírody*, Supplement 20: 23–30.
- Rose, F. 1976. Lichenological indicators of age and environment continuity in woodlands. In: Brown, D. H., Hawksworth, D. L. & Bailey, R. H. (eds). *Lichenology: Progress and Problems*. Academic Press, London, pp. 279–307.
- Scheidegger, C. 1985. Systematische Studien zur Krustenflechte *Anzina carneonivea* (Trapeliaceae, Lecanorales). *Nova Hedwigia* 41:191–218.
- Scheidegger, C., Groner, U., Keller, C. & Stofer, S. 2002. Biodiversity Assessment Tools – Lichens. In: Nimis, P.L., Scheidegger, C. & Wolseley, P.A. (eds.). *Monitoring with Lichens – Monitoring Lichens*. Dordrecht, Boston, London, Kluwer Academic, pp. 359–365. [http://dx.doi.org/10.1007/978-94-010-0423-7\\_35](http://dx.doi.org/10.1007/978-94-010-0423-7_35)
- Scheidegger, C., Stofer, S., Dietrich, M., Groner, U., Keller, C. & Roth, I. 2000. Estimating regional extinction probabilities and reduction in populations of rare epiphytic lichen-forming fungi. *Forest Snow and Landscape Research* 75(3): 415–433.
- Scheidegger, C. & Werth, S. 2009. Conservation strategies for lichens: insights from population biology. *Fungal biology reviews* 23: 55–66. <http://dx.doi.org/10.1016/j.fbr.2009.10.003>
- Smith, C. W., Aproot, A., Coppins, B. J., Fletcher, A., Gilbert, O. L., James, P. W. & Wolseley, P. A. 2009. *The Lichens of Great Britain and Ireland*. Ed. British Lichen Society, London. 1046 pp.
- Thor, G. 1995. Red Lists – Aspects of their compilation and use in Lichen conservation. *Mitteilungen der Eidgenössischen Forschungsanstalt für Wald, Schnee und Landschaft* 70: 11–27.
- Tønsberg, T. 1992. The sorediate and isidiate, corticolous, crustose lichens in Norway. *Sommerfeltia* 14: 1–331.
- Vondrák, J. & Šoun, J. 2008. Some newly recorded and noteworthy lichen-forming and lichenicolous fungi from Romania. *Acta Botanica Hungarica* 50(1–2): 215–221. <http://dx.doi.org/10.1556/ABot.50.2008.1-2.16>
- White, F. J. & James, P. W. 1985. A new guide to microchemical theniques for the identification of lichen substances. *British Lichen Society Bulletin Supplement*. 27 pp.
- Wirth, V. 1995. *Die Flechten Baden-Württembergs*. (2 vols) Ed. Ulmer, Stuttgart. 1006 pp.
- Wolseley, P. A. 1995. A global perspective on the status of lichens and their conservation. *Mitteilungen der Eidgenössischen Forschungsanstalt für Wald, Schnee und Landschaft* 70: 11–27.
- Yavuz, M. & Çobanolu, G., 2008. Lichen records from Dobrogea, Romania. *Muzeul Olteniei Craiova. Oltenia. Studii și comunicări. Științele Naturii* 24: 17–21.
- Zoller, S., Frey, B. & Scheidegger, C., 2000. Juvenile development and diaspore survival in the threatened epiphytic lichen species *Sticta fuliginosa*, *Leptogium saturninum* and *Menegazzia terebrata*: conclusions for in situ conservation. *Plant Biology* 2: 496–504. <http://dx.doi.org/10.1055/s-2000-5954>

