

Life-threatening complications of adult-onset Still's disease

Petros Efthimiou · Sabeeda Kadavath · Bella Mehta

Received: 29 November 2013 / Revised: 27 December 2013 / Accepted: 3 January 2014 / Published online: 17 January 2014
© Clinical Rheumatology 2014

Abstract Adult-onset Still's Disease (AOSD) since its description in 1971 has proven to be a very complex and challenging disease entity. This rare auto-inflammatory disease is classically described by the “Still's triad” of fever, rash, and arthritis, although the atypical cases frequently outnumber the typical ones. The exact pathogenesis and etiologic factors responsible for the clinical features remain largely obscure, despite recent suggestive cytokine biology findings. Diagnosis is made on clinical grounds, following the exclusion of mimickers of infectious, autoimmune or neoplastic etiology, with the additional consideration of non-specific laboratory abnormalities such as peripheral leukocytosis and elevation of serum ferritin and other acute phase reactants. The disease manifestations are protean and can include diverse complications, affecting multiple organ systems. Moreover, the severity of the organ involvement can vary considerably, representing a wide spectrum from the self-limited to severe. The mainstay of therapy has evolved from the traditional use of corticosteroids and oral immunosuppressants to the newer targeted treatments with biologic agents. The scope of this review is to alert the clinician to the existence of life-threatening AOSD complications, namely the macrophage activation syndrome, disseminated intravascular coagulopathy, thrombotic thrombocytopenic purpura, diffuse alveolar hemorrhage, and pulmonary arterial hypertension. Such knowledge may lead in earlier

recognition, prompt treatment, and, ideally, improved patient outcomes.

Keywords Adult-onset still's disease · Complications · Etiology · Macrophage activation syndrome (MAS) · Pathogenesis · Treatment

Introduction

Adult-onset Still's disease (AOSD) is a systemic inflammatory disorder analogous to the systemic form of juvenile idiopathic arthritis (sJIA). The clinical manifestations of this disease range from spiking fevers, arthritis, evanescent rash, elevated liver enzymes, lymphadenopathy, hepatosplenomegaly, and serositis [1–3]. While it is not possible to make an accurate prognosis during the initial presentation of the disease, several clinical characteristics, identified in large patient cohorts, may hint toward a more complicated course (Table 1).

The pathogenetic mechanism for this condition has yet to be ascertained. Involvement of an infectious agent, as suggested by the reported temporal relationship between disease onset and prior exposure to viral, bacterial, and other pathogenic organisms has been suspected. This hypothesis is consistent with the pre-eminent role played by innate immunity in this disease. A predisposing genetic background has also been suggested, although no consistent associations with HLA haplotypes or other non-HLA genes has been confirmed [4]. Several proinflammatory cytokines have been implicated in the pathogenesis; high levels of interleukin (IL)-1, IL-6, and IL-18; macrophage colony-stimulating factor; interferon- γ ; and tumor necrosis factor (TNF- α) have been detected in sera from patients with AOSD [5]. The clinical response to the empirical use of biologic medications that were originally developed for the treatment of rheumatoid arthritis (RA;

P. Efthimiou (✉) · S. Kadavath
Rheumatology Division, Lincoln Medical and Mental Health Center,
234 E. 149th Street, New York, NY 10451, USA
e-mail: pe53@cornell.edu

P. Efthimiou
Weill Cornell Medical College, New York, NY, USA

B. Mehta
Westchester Medical Center, New York, NY, USA

Table 1 Unfavorable prognostic factors for AOSD patients

• Rash
• Polyarthritis
• Root joint arthritis (hips and shoulders)
• Pleuritis
• Interstitial pneumonia
• Elevated ferritin levels
• Failure of fever to subside after 3 days of systemic corticosteroid treatment

Table 2) and target these inflammatory mediators has shown promise, thus pointing towards their role in the disease pathogenesis [6–9]. Prognosis depends on the course of the disease and tends to be more favorable when systemic symptoms predominate and the disease does not evolve into its chronic articular sub-type. Laboratory abnormalities including neutrophil leukocytosis, abnormal liver function tests, and increased ferritin levels.

Diseases that often need to be excluded, before the diagnosis of AOSD can be safely made, include infectious, neoplastic, and autoimmune disorders. However, lack of disease specificity for the symptoms and the laboratory markers hinders the diagnosis and the absence of established disease activity markers makes determination of treatment efficacy problematic [10–13]. More than one-third of patients' progress into chronic disease where complications such as pericarditis, cardiac tamponade, disseminated intravascular coagulation, and hepatic and respiratory failure are associated with untoward morbidity and mortality [14, 15].

AOSD treatment has evolved and largely abandoned the use of non-steroidal anti-inflammatory drugs (NSAIDs). Systemic corticosteroids are used, predominantly during the initial presentation, where the systemic symptoms predominate and often in combination with immunosuppressants (e.g., methotrexate) [16]. Biological agents (mainly IL-1 and IL-6 inhibitors) have been successfully used in refractory cases [5].

Table 2 Biological treatments for refractory AOSD patients

IL-1 receptor inhibitors
• Anakinra [47]
• Rilonacept [106]
• Canakinumab [6]
TNF-alpha inhibitors
• Infliximab [107]
• Etanercept [108]
• Adalimumab [109]
IL-6 receptor inhibitors
• Tocilizumab [110]
B-cell depletors
• Rituximab [111]

The goal of this article is to discuss some of the most severe AOSD-related complications and review the literature for the collective experience relating to their clinical presentation, management, and prognosis.

Macrophage activation syndrome

Macrophage activation syndrome or reactive hemophagocytic syndrome (RHS) is a life-threatening complication of AOSD, with a reported mortality rate ranging between 10 and 22 % [17–20]. It is characterized by an uncontrollable activation of the reticulo-endothelial system within the bone marrow, reticuloendothelial system and central nervous system, with subsequent phagocytosis of hematopoietic cells by tissue macrophages (histiocytes) [20, 21]. Its clinical picture consists of acute high fever, lymphadenopathy, and hepatosplenomegaly (Table 3). Laboratory findings include pancytopenia and high serum levels of ferritin, triglycerides, and liver enzymes, while, paradoxically, demonstrating a normal erythrocyte sedimentation rate (ESR). Although macrophage activation syndrome (MAS) is, overall, a very rare condition, it is not uncommon in AOSD, with an incidence of 12–14 % in two recent series, a rate higher than other rheumatic diseases [22, 23]. Clinical and biological features of MAS closely resemble reactive hemophagocytic lymphohistocytosis (HLH) and the disease is in fact considered today as a subclass of HLH (secondary), which is induced by heterogeneous disorders including infections, malignant tumors, and medications [24]. HLH criteria include a decrease in white blood cell count, platelets, or fibrinogen and the demonstration of low or absent natural killer cell activity or soluble IL-2 receptor. However, the minimum threshold level for hyperferritinemia required for the diagnosis of HLH (500 µg/l) is not useful in detecting MAS, with the ferritin level generally peaking to more than 5,000 µg/l in the latter condition [25].

Hypofibrinogenemia is one of the most important clues for the diagnosis of MAS since patients usually have high fibrinogen levels due to their underlying inflammatory disease. An increased fibrinolysis is most probably due to the uncontrolled activation of the macrophages and plasminogen overproduction. A decrease in factor II and factor VII+X values has also been observed in some reports [26].

The most frequently implicated triggering factors include infections, medications, and disease flare ups [24, 27]. Patients with MAS have a decreased natural capacity to eradicate infected cells and to eliminate antigen stimulation. This in turn induces T cell activation and proliferation with cytokine secretion (interferon-gamma and granulocyte macrophage colony-stimulating factor) and macrophage hyper-activation. The end result is an uncontrollable increase in TNF α , interleukin-1, and interleukin-6 production with a severe systemic inflammatory reaction, i.e., “cytokine storm” [20].

Table 3 MAS diagnostic factors

Clinical features
High fever—non-remitting
Hepatomegaly, splenomegaly
Lymphadenopathy
CNS dysfunction
Hemorrhages
Laboratory features
Abnormal liver function tests
Decreased ESR
Cytopenia
Coagulopathy
Hypertriglyceridemia
Hyponatremia
Hypoalbuminemia
Hyperferritinemia
Elevated sCD25 and sCD163
Histopathological features
BM—macrophage hemophagocytosis
Increased CD163 staining BM

ESR erythrocyte sedimentation rate, BM bone marrow

The immunodeficiency state induced by the treatment of AOSD may lead to reactivation of latent viruses, such as Epstein-Barr virus or cytomegalovirus, which could, theoretically, trigger MAS. There is also a suggestion that certain therapeutic agents, such as non-steroidal anti-inflammatory drugs, methotrexate, sulfasalazine, penicillamine, and lately TNF- α , IL-1, and IL-6 inhibitors may be capable of provoking MAS, often complicating their therapeutic use [23].

Preliminary diagnostic guideline used in MAS complicating sJIA was based on a set of laboratory and clinical criteria [27]. The clinical criteria include central nervous system dysfunction, hemorrhage, and hepatomegaly while the laboratory criteria consisted of a low platelet and white cell count, hypofibrinogenemia and elevated AST (Table 4). Early suspicion of MAS is most commonly raised by the detection of subtle laboratory changes, whereas clinical symptoms may be delayed. A recent international effort to identify candidate markers using an expert consensus process identified nine criteria that included a falling platelet count, hyperferritinemia, evidence of macrophage hemophagocytosis in the bone marrow, increased liver enzymes, falling leukocyte count, a persistent continuous fever ≥ 38 °C, a falling erythrocyte sedimentation rate, hypofibrinogenemia, and hypertriglyceridemia [25]. Hemophagocytosis, seen in bone marrow aspiration and biopsy establishes the diagnosis, even though hemophagocytosis could be seen more frequently in biopsies

Table 4 AOSD vs. MAS: clinical features and laboratory investigations

	AOSD	MAS
1. Fever pattern	Quotidian	Non-remitting
2. Rash	Evanescant maculopapular	Urticaria
3. Arthritis	+	–
4. RES involvement	+	++
5. CNS involvement	Rarely	+
6. ESR	High	Low
7. WBC	High	Low/normal
8. Platelets	High	Low/normal
9. Ferritin	High	High/very high
10. AST/ALT	Normal	High/normal
11. Fibrinogen	Normal	Low/normal
12. Triglycerides	Normal	High

ALT alanine aminotransferase, AST aspartate aminotransferase, CNS central nervous system, ESR erythrocyte sedimentation rate, RES reticulo-endothelial system, WBC white blood cells

from the liver, spleen, and/or lymph nodes. Bone marrow aspiration is considered the gold standard and is usually required in atypical cases where there may be a diagnostic dilemma. There is significant overlap between AOSD and MAS and these two conditions are often thought to be anchoring the same disease spectrum, with AOSD representing the milder form. Moreover, it may be a two-way street, with the individual patients' clinical manifestations fluctuating between those two phenotypes, throughout the disease's lifespan [22]. It is not unusual for a patient to simultaneously carry both diagnoses, i.e., AOSD and concurrent mild MAS. The similarities in clinical presentation may reflect overlapping pathophysiologic mechanisms. Less severe or "subclinical" cases of MAS may not be diagnosed if they respond to immunosuppressive treatments given for a presumed flare of AOSD [28–30]. In 2005, Mazodier et al. published the results of a study showing significantly elevated serum IL-18 concentrations and decreased levels of IL-18 binding protein (IL-18BP) in the population affected with MAS, when compared to healthy controls [31]. Interestingly, IL-18 has been repeatedly shown to play a major role in the pathogenesis of AOSD and IL-18 inhibition may prove to be a desirable future approach for both entities. A case of AOSD-associated thrombotic microangiopathy (TMA) has also been reported suggesting the association of IL-18, high levels were seen at onset of disease which dramatically decreased with TMA resolution suggesting the antiangiogenic effect of IL18 [32].

Empiric treatment of MAS traditionally consisted of the following agents alone or in combination: intravenous pulse corticosteroids, immune gamma globulin (IVIG), and cyclosporine A (CyA). Mizrahi et al. described a patient with relapsing MAS successfully treated with CyA and mycophenolate mofetil (MMF) [33].

There are several reports of the use of IL-1 inhibitors in the management of MAS. Kelly et al. described a case of successful use of anakinra in management of MAS in a 13-year-old patient with sJIA [34]. Two more cases of MAS associated with sJIA reported symptoms promptly resolving after anakinra initiation. In addition to the reported efficacy, the authors marveled at the avoidance of the potentially toxic immunosuppressant cyclosporine A and/or the antineoplastic agent etoposide [35]. SJIA-associated MAS has required anakinra doses as high as 10 mg/kg/day and, in one published report, combination with abatacept [36]. Recently, a case of AOSD-associated MAS that was complicated by a history of active spinal tuberculosis responded favorably within 48 h to the addition of anakinra to her corticosteroid regimen. Furthermore, there was no tuberculosis reactivation after 10 months of anakinra treatment [37].

Anti-cytokine therapies are finding a niche in the treatment of MAS that complicates sJIA. TNF inhibitors, despite some early anecdotal reports of efficacy, are currently considered ineffective or even harmful. Temporal association of anti-TNF α therapy and bacterial infection has been suggested to play a catalytic role in the deregulation of macrophage–lymphocyte interactions in MAS [38]. There are several cases of MAS following the use of etanercept in patients with AOSD [37, 39]. In such cases, the observed decreases in ESR and WBC counts may be erroneously interpreted as response to treatment, while in reality they may signify the beginning of MAS. Therefore, MAS should always be considered in the differential diagnosis of AOSD patients with low WBC and platelet counts lacking clinical improvement after etanercept use.

Tocilizumab, a humanized anti-interleukin-6 receptor monoclonal antibody, was reported to be effective in the management of AOSD that involved MAS during the induction therapy, although its efficacy is unclear for the highly active phase of the disease [40].

Future therapeutic protocols for the treatment of MAS may include different combinations of pulse corticosteroids, CyA or etoposide, and anti-cytokine treatment, such as blockade of IL-1 and/or IL-6.

Disseminated intravascular coagulopathy

Disseminated intravascular coagulopathy is a life-threatening complication caused by an uninhibited activation of the coagulation system. This severe complication has been reported to occur in AOSD and its diagnosis can be challenging, even for seasoned clinicians [41]. Eight disseminated intravascular coagulopathy (DIC) cases, some associated with hepatopathy have been reported in the literature to date [42]. The clinical presentation of a patient with AOSD complicated by coagulopathy and the patient becoming acutely ill may

closely mimic sepsis, leading to treatment with antimicrobials only and delaying management with immunomodulators.

To further complicate diagnosis, DIC and multi-organ failure may also occur as a complication of AOSD-related MAS [43]. However, patterns of non-remitting fever and purpuric or petechial rash that accompany DIC can be differentiated from the remitting high-spiking fever and evanescent maculopapular rash of AOSD. The precipitous fall of ESR in the context of worsening clinical condition and persistent hyper-ferritinemia should be a red flag and raise the index of suspicion for the presence of DIC. High titers of serum-soluble adhesion molecules and soluble thrombomodulin have also been noted during the first episode of DIC which indicate endothelial cell damage in AOSD complicated by DIC [44].

Interestingly, there have been reports of patients developing acute myocarditis during DIC attacks in AOSD, potentially implicating a, yet unidentified, infectious trigger [45]. Treatment with immunomodulators, together with supportive measures, can be effective in even some of the most severe cases, as it was in a case complicated with concomitant acute respiratory distress syndrome (ARDS) and DIC successfully managed with prednisolone therapy [46].

The efficacy of anakinra has also been advocated patients with severe flare of AOSD and DIC [47]. Some experts have even suggested it should be first line therapy in life-threatening DIC [43, 48].

More recently, a case of glucocorticoid and cyclosporine refractory AOSD complicated by DIC was successfully managed with tocilizumab, a humanized anti-IL-6 receptor monoclonal antibody. IL-6 inhibition led to a rapid response, symptomatic improvement, and corticosteroid sparing [49].

Thrombotic thrombocytopenic purpura

Thrombotic thrombocytopenic purpura (TTP) associated with AOSD has been documented with nine published cases in the literature and is invariably associated with significant morbidity and mortality [50]. TTP is more common and more severe in African Americans compared to other races and ethnicities [51]. ADAMTS13 activity and inhibition have been found to be important predictors of disease outcome [52]. An early sign that should raise the index of suspicion for TTP is development of acute vision impairment. In extreme cases, multiple thrombi formation and massive cytokine release causing progressive vascular permeability, worsening brain edema and, ultimately, rapid death may also occur [53].

The dual diagnosis of AOSD and TTP has interested scientists and physicians alike since its first description, since there has always been hope of discovery of a common pathogenetic path. However, no mechanism, so far, has been able to sufficiently explain their coexistence. It has been suggested

that autoantibodies to CD36 (glycoprotein IV) may play a dual role in the pathogenesis of TTP and AOSD, since CD36 is important in the clearance of inflammatory cells and T-cell activation [54]. Alternatively, decreased activity of von Willebrand factor-cleaving protease (a disintegrin and metalloproteinase with thrombospondin type 1 motif), member 13 (ADAMTS-13) may play a prominent role in the pathogenesis of TTP [55]. In the presence of ADAMTS-13 inhibitors (e.g., antibodies against ADAMTS-13), intravascular platelet thrombi develop [56]. The characteristic pathologic findings in TTP include hyalinized thrombi in the vascular beds of the brain, pancreas, heart, and kidneys. TTP is a medical emergency, since the outcome of untreated TTP is renal failure and death [57]. There have been at least 11 published cases of TTP/HUS associated with AOSD, five of which with presenting renal failure [56]. Masuyama et al. reported a case of AOSD complicated by TTP that resulted in retinal microangiopathy and rapidly fatal cerebral edema [53].

TTP is often diagnosed on the basis of thrombocytopenia and microangiopathic hemolytic anemia that cannot be explained otherwise, so that treatment can be begun early to lower mortality rates in this often fatal disease [58]. Lack of treatment results in a mortality rate of about 90 %, but even with treatment the mortality rate can be as high as 20 %. The current mainstay of treatment is plasma exchange because of its efficacy in randomized clinical trials, and relapsing episodes can be managed successfully with additional plasma exchange, aspirin, vincristine, hemodialysis, corticosteroids, intravenous immunoglobulins, azathioprine, cyclophosphamide, rituximab, and/or splenectomy [50, 56]. There are only rare reports of treatment of patients with TTP and AOSD with biologic agents. In 2010, Sumida et al. presented a case of TTP in multi-drug-resistant Still's successfully treated with tocilizumab that seemed to be successful in treating both the TTP and the underlying treatment-refractory AOSD [59].

Diffuse alveolar hemorrhage

Pulmonary involvement is well-known in AOSD and is seen in up to 53 % of AOSD cases, with the most common pulmonary diseases being pleural effusion and transient pulmonary infiltrates [60]. Life-threatening pulmonary complications of AOSD include respiratory distress syndrome, and diffuse alveolar hemorrhage. It is still a matter of speculation whether the association between AOSD and diffuse alveolar hemorrhage (DAH) is coincidental or if it can be attributed to some common pathogenic link. The first case of chronic AOSD complicated with diffuse alveolar hemorrhage during the acute flare of the disease has been reported by Sari et al. in 2009 [61]. Several studies suggest that alterations in pro-

inflammatory cytokine production play an important role in the pathogenesis of AOSD.

Interleukin-18 (IL-18) appears to be an anchor cytokine in AOSD pathogenesis, as it is overproduced in the acute phase of the disease and is believed to be the cytokine initiating the inflammatory cascade that includes interferon gamma, IL-6, and TNF- α [62]. DAH is a medical emergency characterized by the accumulation of red blood cells in the alveolar spaces. Hemoptysis, coughing, and progressive dyspnea are common initial symptoms, and the disease may progress to acute respiratory failure. Hemoptysis is not a *sine qua non* symptom, and may be even absent. Chest radiography is nonspecific and commonly shows new patchy or diffuse alveolar opacities. Characteristic laboratory features of DAH include decreasing hemoglobin and hemorrhagic bronchoalveolar lavage fluid [63, 64]. Treatment modalities in DAH include pulse steroids administered at a dose of 1,000 mg/day for 5 days [61], plasma exchange, and intravenous cyclophosphamide.

Pulmonary arterial hypertension

Cardiopulmonary disease has been described in 30–40 % of patients with AOSD. Pulmonary manifestations include aseptic pneumonitis, interstitial lung disease, serositis with accompanying pleural effusion, and rarely ARDS [65]. Affected patients may complain of cough, pleuritic chest pain, and dyspnea. The association of pulmonary arterial hypertension (PAH) with connective tissue diseases has been well characterized in autoimmune diseases such as systemic sclerosis and systemic lupus erythematosus. We are aware of only six published cases to date [66].

It was in 1990 that the first case was reported in Japan of a 29-year-old woman who had developed progressive, pre-capillary severe hypertension 2.5 years after the diagnosis of AOSD was established [67]. Two additional cases were reported in northern Taiwan [68], in a single-center case series of 19 patients with co-prevalence of PAH and various connective tissue diseases. Another case reported from the USA was that of a 29-year-old woman with a 9-year history of AOSD who developed the characteristic clinical and hemodynamic findings of PAH and despite treatment, including anakinra 100 mg/day SQ, died 2.5 months after the diagnosis of PAH [69]. In 2009, a case was reported in an 18-year-old female from India with new onset AOSD who quickly developed severe PAH (mean PAP of 65 mmHg) in the absence of other causes [70]. Campos et al. reported a case of PAH responsive to anakinra in a 27-year-old woman with a 7-year history of AOSD [66].

PAH is diagnosed when the resting mean pulmonary artery pressure (PAPm) is greater than 25 mmHg, in the setting of pulmonary capillary wedge pressure <15 mmHg, based on right heart catheterization and after exclusion of conditions

like left heart and thromboembolic disease [71]. Pulmonary microangiopathy, which may also cause renal involvement and thrombotic thrombocytopenic purpura in AOSD patients, has been implicated. Inflammatory mechanisms appear to play a significant role in autoimmune rheumatic-disease-associated PAH, although the mechanisms may differ significantly in AOSD where innate immunity mechanisms have a predominant role. In autoimmune-disease-associated PAH, the presence of antinuclear antibodies, rheumatoid factor, IgG, and complement fraction deposits in the walls of pulmonary vessels are characteristic [72]. It is believed that the regulatory T-cells play a role in preventing B-cell activity, and lead to the severe angioproliferative changes. Venocclusive disease, formation of microthrombi, and pulmonary fibrosis are also frequently seen. The key histological findings are intimal thickening and media hypertrophy of the small- and medium-sized pulmonary arteries and formation of plexiform lesions. Both the sporadic forms of PAH and those associated with rheumatic diseases are characterized by endothelial proliferation [73].

The treatment modality for PAH in AOSD aims to target the inflammatory component. Survival in CTD-associated PAH (CTD-PAH) has been observed to be shorter than in idiopathic PAH (IPAH) [74]. The dearth of available data on this complication, the rarity of reported cases and the subsequent absence of randomized control trials, has led to management of this condition with empirical therapy centered around the use of immunosuppressive therapy in conjunction with vasodilator therapy. Several case reports describe regression of PAH with immunosuppressive therapy in patients with rheumatic diseases [75]. All three groups of advanced therapies; prostanoids, endothelin receptor antagonists, and phosphodiesterase-5 inhibitors used in idiopathic PAH have also been shown to improve pulmonary hemodynamics and functional status in patients with CTD-PAH [76–80]. Biologic agents have been explored as an important avenue in patients, especially in steroid-resistant cases [81]. Anakinra, a recombinant human IL-1 receptor antagonist, has been shown to reduce the systemic inflammatory reactions [82]. Newer approaches, including tocilizumab, an anti-IL-6 receptor monoclonal antibody have also been proposed to target the IL6 dysregulation [83–85].

Early diagnosis, classification, and initiation of a disease-specific treatment algorithm might improve survival in these patients.

Liver involvement

Liver involvement in AOSD patients can range from mild to severe. They can present with no symptom or with hepatomegaly, icterus, loss of appetite, fatigue, and right abdominal pain. In a series studying liver involvement in 77 patients with

AOSD, clinical and laboratory features were compared with previously published data [86–88]. Hepatomegaly was reported in nearly half of the AOSD patients while transient elevation of liver enzymes was seen in three quarter of the cases. Hepatic failure, however, is a rare complication of adult Still's disease and it has been suggested that it may be associated with hemophagocytic syndrome [89].

In a 10-year retrospective study of 12 patients who satisfied the diagnostic criteria for adult Still's disease, severe liver abnormalities were noted. One hundred percent of patients presented with fever, and 41 % with hepatomegaly. Abnormalities in liver function tests were seen in 92 % of patients, wherein 17 % of patients presented with serum aminotransferase levels five times the normal level and 83 % of patients with levels that were between two and five times normal [90].

The pathophysiology of liver involvement is not known. Cytokine production and sustained macrophage activation have been implicated to play a role in the liver manifestations [87, 91]. Previous descriptions of hepatic histopathology in AOSD consist of portal inflammatory infiltrates in most cases [92], mild interstitial hepatitis [93], mild portal fibrosis, focal hepatitis with necrosis [94], and mild chronic necro-inflammatory change [95].

NSAIDs and MTX may also contribute to the liver damage of AOSD. Remission with the use of these drugs has been observed to even normalize the level of transaminase [85–87]. Severe liver disease however may not respond to anti-inflammatory or immunosuppressive therapy. There are reports of two patients with adult Still's disease-related liver failure who required liver transplantation, one of whom died [96, 97].

Discussion

AOSD is a systemic auto-inflammatory disease that may be infrequently associated with significant morbidity and life-threatening manifestations. Although the prognosis of AOSD is generally favorable, death may occur in some patients due to overwhelming infection, hepatic failure, amyloidosis, adult respiratory distress syndrome, heart failure, macrophage activation syndrome, and pulmonary hypertension [69, 98–101]. Clinicians should be aware of those complications, since early recognition and prompt management in a monitored setting can significantly decrease morbidity and mortality. Treatment efforts should include both supportive measure and coordinated efforts to control the underlying systemic inflammation. High doses of systemic glucocorticoids combined with immunosuppressive agents are usually recommended [102–104]. Targeted therapy with biologic agents, in particular with anti-IL1 and anti-IL6 inhibitors has enhanced our ability to manage refractory AOSD and its associated severe complications [105]. However, the rarity of the disease,

diversity of clinical presentation, and lack of controlled studies (especially in AOSD) has limited our knowledge of the exact efficacy and safety of our interventions. For example, novel treatments for AOSD and its pediatric counterpart sJIA have been reported to both effectively treat and potentially induce severe complications such as MAS. This gap in knowledge could be potentially filled by information gathered by prospective, disease specific, registries, and an international collaborative effort.

Disclosures None.

References

1. Jaqua NT, Finger D, Hawley JS. Adult-onset Still's disease masquerading as sepsis in an asplenic active duty soldier. *Case reports in medicine*. 2012;2012:349521. PubMed PMID: 23251171. Pubmed Central PMCID: PMC3522482. Epub 2012/12/20. eng.
2. Duburcq T, Delannoy PY, Sivova N, Leroy O (2013) Adult onset Still's disease revealed by a myocarditis. *Ann Fr Anesth Reanim* 32(1):50–52, PubMed PMID: 23200908. Epub 2012/12/04. Myocardite revelant une maladie de Still de l'adulte. fre
3. Fautrel B (2008) Adult-onset Still disease. *Best Pract Res Clin Rheumatol* 22(5):773–792, PubMed PMID: 19028363. Epub 2008/11/26. eng
4. Efthimiou P, Paik PK, Bielory L (2006) Diagnosis and management of adult onset Still's disease. *Ann Rheum Dis* 65(5):564–572, PubMed PMID: 16219707. Epub 2005/10/13. eng
5. Kontzias A, Efthimiou P (2008) Adult-onset Still's disease: pathogenesis, clinical manifestations and therapeutic advances. *Drugs* 68(3):319–337, PubMed PMID: 18257609. Epub 2008/02/09. eng
6. Kontzias A, Efthimiou P (2012) The use of Canakinumab, a novel IL-1beta long-acting inhibitor, in refractory adult-onset Still's disease. *Semin Arthritis Rheum* 42(2):201–205, PubMed PMID: 22512815. Epub 2012/04/20. eng
7. Yokota S, Tanaka T, Kishimoto T (2012) Efficacy, safety and tolerability of tocilizumab in patients with systemic juvenile idiopathic arthritis. *Ther Adv Musculoskelet Dis* 4(6):387–397, PubMed PMID: 23227116. Pubmed Central PMCID: PMC3512171. Epub 2012/12/12. eng
8. De Benedetti F, Brunner HI, Ruperto N, Kenwright A, Wright S, Calvo I et al (2012) Randomized trial of tocilizumab in systemic juvenile idiopathic arthritis. *N Engl J Med* 367(25):2385–2395, PubMed PMID: 23252525. Epub 2012/12/21. eng
9. Ogata A, Tanaka T (2012) Tocilizumab for the treatment of rheumatoid arthritis and other systemic autoimmune diseases: current perspectives and future directions. *Int J Rheumatol* 2012:946048, PubMed PMID: 22315615. Pubmed Central PMCID: PMC3270395. Epub 2012/02/09. en
10. Pouchot J, Sampalis JS, Beaudet F, Carette S, Decary F, Salusinsky-Sternbach M et al (1991) Adult Still's disease: manifestations, disease course, and outcome in 62 patients. *Medicine (Baltimore)* 70(2):118–136, PubMed PMID: 2005777. Epub 1991/03/01. eng
11. Ohta A, Yamaguchi M, Kaneoka H, Nagayoshi T, Hiida M (1987) Adult Still's disease: review of 228 cases from the literature. *J Rheumatol* 14(6):1139–1146, PubMed PMID: 3325642
12. Efthimiou P, Kontzias A, Ward CM, Ogden NS (2007) Adult-onset Still's disease: can recent advances in our understanding of its pathogenesis lead to targeted therapy? *Nat Clin Pract Rheumatol* 3(6):328–335, PubMed PMID: 17538564. Epub 2007/06/01. eng
13. Cagatay Y, Gul A, Cagatay A, Kamali S, Karadeniz A, Inanc M et al (2009) Adult-onset Still's disease. *Int J Clin Pract* 63(7):1050–1055, PubMed PMID: 17511792. Epub 2007/05/22. eng
14. Najam F, Fuchs HA (2001) Cardiac tamponade: a life-threatening complication of Still's disease. *J Clin Rheumatol* 7(2):97–101, PubMed PMID: 17039104. Epub 2006/10/14. eng
15. Biron C, Chambellan A, Agard C, Denis M, Germaud P, Hamidou MA (2006) Acute respiratory failure revealing adult-onset Still's disease: diagnostic value of low glycosylated ferritin level. *Clin Rheumatol* 25(5):766–768, PubMed PMID: 16283419. Epub 2005/11/12. eng
16. Bagnari V, Colina M, Ciancio G, Govoni M, Trotta F (2010) Adult-onset Still's disease. *Rheumatol Int* 30(7):855–862, PubMed PMID: 20020138. Epub 2009/12/19. eng
17. Silva CA, Silva CH, Robazzi TC, Lotito AP, Mendroni Junior A, Jacob CM et al (2004) Macrophage activation syndrome associated with systemic juvenile idiopathic arthritis. *J Pediatr* 80(6):517–522, PubMed PMID: 15622430. Epub 2004/12/29. Síndrome de ativacao macrofagica associada com artrite idiopatica juvenil sistêmica. por
18. Ramanan AV, Schneider R (2003) Macrophage activation syndrome following initiation of etanercept in a child with systemic onset juvenile rheumatoid arthritis. *J Rheumatol* 30(2):401–403, PubMed PMID: 12563702. Epub 2003/02/04. eng
19. Sawar H, Espinoza LR, Gedalia A (2004) Macrophage activation syndrome and etanercept in children with systemic juvenile rheumatoid arthritis. *J Rheumatol* 31(3):623, author reply 4. PubMed PMID: 14994420. Epub 2004/03/03. eng
20. Tristano AG (2008) Macrophage activation syndrome: a frequent but under-diagnosed complication associated with rheumatic diseases. *Medical science monitor : international medical journal of experimental and clinical research* 14(3):RA27–RA36, PubMed PMID: 18301366. Epub 2008/02/28. eng
21. Durand M, Troyanov Y, Laflamme P, Gregoire G (2010) Macrophage activation syndrome treated with anakinra. *J Rheumatol* 37(4):879–880, PubMed PMID: 20360206. Epub 2010/04/03. eng
22. Arlet JB, D Leth, Marinho A, Amoura Z, Wechsler B, Papo T, et al. Reactive haemophagocytic syndrome in adult onset Still's disease: report of 6 patients and review of the literature. *Ann Rheum Dis*. 2006 Mar 15. PubMed PMID: 16540551.
23. Singh S, Samant R, Joshi VR (2008) Adult onset Still's disease: a study of 14 cases. *Clin Rheumatol* 27(1):35–39, PubMed PMID: 17701269. Epub 2007/08/19. eng
24. Sawhney S, Woo P, Murray KJ (2001) Macrophage activation syndrome: a potentially fatal complication of rheumatic disorders. *Arch Dis Child* 85(5):421–426, PubMed PMID: 11668110. Pubmed Central PMCID: 1718981. Epub 2001/10/23. eng
25. Davi S, Consolaro A, Guseinova D, Pistorio A, Ruperto N, Martini A et al (2011) An international consensus survey of diagnostic criteria for macrophage activation syndrome in systemic juvenile idiopathic arthritis. *J Rheumatol* 38(4):764–768, PubMed PMID: 21285158. Epub 2011/02/03. eng
26. de Saint Basile G, Fischer A, Dautzenberg MD, Durandy A, Le Deist F, Angles-Cano E et al (1985) Enhanced plasminogen-activator production by leukocytes in the human and murine Chediak-Higashi syndrome. *Blood* 65(5):1275–1281, PubMed PMID: 4039611. Epub 1985/05/01. eng
27. Ravelli A, Magni-Manzoni S, Pistorio A, Besana C, Foti T, Ruperto N et al (2005) Preliminary diagnostic guidelines for macrophage activation syndrome complicating systemic juvenile idiopathic arthritis. *J Pediatr* 146(5):598–604, PubMed PMID: 15870661. Epub 2005/05/05. eng
28. Fall N, Barnes M, Thornton S, Luyrink L, Olson J, Ilowite NT et al (2007) Gene expression profiling of peripheral blood from patients

- with untreated new-onset systemic juvenile idiopathic arthritis reveals molecular heterogeneity that may predict macrophage activation syndrome. *Arthritis Rheum* 56(11):3793–3804, PubMed PMID: 17968951. Epub 2007/10/31. eng
29. Blessing J, Prada A, Siegel DM, Villanueva J, Olson J, Ilowite NT et al (2007) The diagnostic significance of soluble CD163 and soluble interleukin-2 receptor alpha-chain in macrophage activation syndrome and untreated new-onset systemic juvenile idiopathic arthritis. *Arthritis Rheum* 56(3):965–971, PubMed PMID: 17328073. Epub 2007/03/01. eng
 30. Behrens EM, Beukelman T, Paessler M, Cron RQ (2007) Occult macrophage activation syndrome in patients with systemic juvenile idiopathic arthritis. *J Rheumatol* 34(5):1133–1138, PubMed PMID: 17343315. Epub 2007/03/09. eng
 31. Mazodier K, Marin V, Novick D, Farnarier C, Robitail S, Schleinitz N et al (2005) Severe imbalance of IL-18/IL-18BP in patients with secondary hemophagocytic syndrome. *Blood* 106(10):3483–3489, PubMed PMID: 16020503. Pubmed Central PMCID: 1895045. Epub 2005/07/16. eng
 32. El Karoui K, Karras A, Lebrun G, Charles P, Arlet J-B, Jacquot C, Orssaud C, Nochy D, Pouchot J (2009) Thrombotic microangiopathy and Purtscher-like retinopathy associated with adult-onset Still's disease: a role for glomerular vascular endothelial growth factor? *Arthritis Rheum* 61:1609–1613. doi:10.1002/art.24826
 33. Mizrahi M, Ben-Chetrit E (2009) Relapsing macrophage activating syndrome in a 15-year-old girl with Still's disease: a case report. *J Med Case Reports* 3:138, PubMed PMID: 20062775. Pubmed Central PMCID: 2803809. Epub 2010/01/12. eng
 34. Kelly A, Ramanan AV (2008) A case of macrophage activation syndrome successfully treated with anakinra. *Nat Clin Pract Rheumatol* 4(11):615–620, PubMed PMID: 18825135. Epub 2008/10/01. eng
 35. Bruck N, Suttorp M, Kabus M, Heubner G, Gahr M, Pessler F (2011) Rapid and sustained remission of systemic juvenile idiopathic arthritis-associated macrophage activation syndrome through treatment with anakinra and corticosteroids. *J Clin Rheumatol* 17(1):23–27, PubMed PMID: 21169853. Epub 2010/12/21. eng
 36. Record JL, Beukelman T, Cron RQ (2011) Combination therapy of abatacept and anakinra in children with refractory systemic juvenile idiopathic arthritis: a retrospective case series. *J Rheumatol* 38(1):180–181, PubMed PMID: 21196588. Epub 2011/01/05. eng
 37. Pamuk ON, Pamuk GE, Usta U, Cakir N (2007) Hemophagocytic syndrome in one patient with adult-onset Still's disease. Presentation with febrile neutropenia. *Clin Rheumatol* 26(5):797–800, PubMed PMID: 16550302. Epub 2006/03/22. eng
 38. Nadia EA, Carvalho JF, Bonfa E, Lotito AP, Silva CA (2009) Macrophage activation syndrome associated with etanercept in a child with systemic onset juvenile idiopathic arthritis. *The Israel Medical Association journal : IMAJ* 11(10):635–636, PubMed PMID: 20077954. Epub 2010/01/19. eng
 39. Stern A, Riley R, Buckley L (2001) Worsening of macrophage activation syndrome in a patient with adult onset Still's disease after initiation of etanercept therapy. *J Clin Rheumatol* 7(4):252–256, PubMed PMID: 17039144. Epub 2006/10/14. eng
 40. Kobayashi M, Takahashi Y, Yamashita H, Kaneko H, Mimori A (2011) Benefit and a possible risk of tocilizumab therapy for adult-onset Still's disease accompanied by macrophage-activation syndrome. *Modern rheumatology/the Japan Rheumatism Association* 21(1):92–96, PubMed PMID: 20737186. Epub 2010/08/26. eng
 41. Arai Y, Handa T, Mitani K (2004) Adult-onset Still disease presenting with disseminated intravascular coagulation. *Rinsho Ketsueki* 45(4):316–318
 42. Aellen P, Raccaud O, Waldburger M, Chamot AM, Gerster JC (1991) Still's disease in adults with disseminated intravascular coagulation. *Schweizerische Rundschau fur Medizin Praxis= Revue suisse de medecine Praxis* 80(15):376–378, PubMed PMID: 2031109. Epub 1991/04/09. Maladie de Still de l'adulte avec coagulation intravasculaire disseminee (CIVD). fre
 43. Park JH, Bae JH, Choi YS, Lee HS, Jun JB, Jung S et al (2004) Adult-onset Still's disease with disseminated intravascular coagulation and multiple organ dysfunctions dramatically treated with cyclosporine A. *J Korean Med Sci* 19(1):137–141, PubMed PMID: 14966357
 44. Sanada I, Kawano F, Tsukamoto A, Kiyokawa T, Shido T, Koga S (1997) Disseminated intravascular coagulation in a case of adult onset Still's disease. [Rinsho ketsueki. The Japanese journal of clinical hematology 38(11):1194–1198, PubMed PMID: 9423337. Epub 1998/01/10. jpn
 45. Colina M, Govoni M, Trotta F (2009) Fatal myocarditis in adult-onset Still disease with diffuse intravascular coagulation. *Rheumatol Int* 29(11):1355–1357, PubMed PMID: 19048255. Epub 2008/12/03. eng
 46. Yokoyama M, Suwa A, Shinozawa T, Fujii T, Mimori T, Akizuki M et al (1995) A case of adult onset Still's disease complicated with adult respiratory distress syndrome and disseminated intravascular coagulation. *Nihon Rinsho Men'eki Gakkai kaishi=Japanese journal of clinical immunology* 18(2):207–214, PubMed PMID: 7553055. Epub 1995/04/01. jpn
 47. Kotter I, Wacker A, Koch S, Henes J, Richter C, Engel A et al (2007) Anakinra in patients with treatment-resistant adult-onset Still's disease: four case reports with serial cytokine measurements and a review of the literature. *Semin Arthritis Rheum* 37(3):189–197, PubMed PMID: 17583775. Epub 2007/06/23. eng
 48. Mori T, Tanigawa M, Iwasaki E, Tamaki S, Ono T, Wada H et al (1993) Cyclosporine therapy of adult onset Still's disease with disseminated intravascular coagulation. [Rinsho ketsueki. The Japanese journal of clinical hematology 34(2):147–152, PubMed PMID: 8492412. Epub 1993/02/01. jpn
 49. Matsumoto K, Nagashima T, Takatori S, Kawahara Y, Yagi M, Iwamoto M et al (2009) Glucocorticoid and cyclosporine refractory adult onset Still's disease successfully treated with tocilizumab. *Clin Rheumatol* 28(4):485–487, PubMed PMID: 19184270. Epub 2009/02/03. eng
 50. Perez MG, Rodwig FR Jr (2003) Chronic relapsing thrombotic thrombocytopenic purpura in adult onset Still's disease. *South Med J* 96(1):46–49, PubMed PMID: 12602713. Epub 2003/02/27. eng
 51. Gopal M, Cohn CD, McEntire MR, Alperin JB (2009) Thrombotic thrombocytopenic purpura and adult onset Still's disease. *Am J Med Sci* 337(5):373–376, PubMed PMID: 19322066. Epub 2009/03/27. eng
 52. Zheng XL, Kaufman RM, Goodnough LT, Sadler JE (2004) Effect of plasma exchange on plasma ADAMTS13 metalloprotease activity, inhibitor level, and clinical outcome in patients with idiopathic and nonidiopathic thrombotic thrombocytopenic purpura. *Blood* 103(11):4043–4049, PubMed PMID: 14982878. Epub 2004/02/26. eng
 53. Masuyama A, Kobayashi H, Kobayashi Y, Yokoe I, Sugimura Y, Maniwa K et al (2013) A case of adult-onset Still's disease complicated by thrombotic thrombocytopenic purpura with retinal microangiopathy and rapidly fatal cerebral edema. *Modern rheumatology/the Japan Rheumatism Association* 23(2):379–385, PubMed PMID: 22623015. Epub 2012/05/25. eng
 54. Tandon NN, Rock G, Jamieson GA (1994) Anti-CD36 antibodies in thrombotic thrombocytopenic purpura. *Br J Haematol* 88(4):816–825, PubMed PMID: 7529543. Epub 1994/12/01. eng
 55. Furlan M, Robles R, Galbusera M, Remuzzi G, Kyrle PA, Brenner B et al (1998) von Willebrand factor-cleaving protease in thrombotic thrombocytopenic purpura and the hemolytic-uremic syndrome. *N Engl J Med* 339(22):1578–1584, PubMed PMID: 9828245. Epub 1998/11/26. eng
 56. Sayarlioglu M, Sayarlioglu H, Ozkaya M, Balakan O, Ucar MA (2008) Thrombotic thrombocytopenic purpura-hemolytic uremic

- syndrome and adult onset Still's disease: case report and review of the literature. *Modern rheumatology/the Japan Rheumatism Association* 18(4):403–406, PubMed PMID: 18427722. Epub 2008/04/23. eng
57. Ruggenti P, Noris M, Remuzzi G (2001) Thrombotic microangiopathy, hemolytic uremic syndrome, and thrombotic thrombocytopenic purpura. *Kidney Int* 60(3):831–846, PubMed PMID: 11532079. Epub 2001/09/05. eng
 58. Onundarson PT, Rowe JM, Heal JM, Francis CW (1992) Response to plasma exchange and splenectomy in thrombotic thrombocytopenic purpura. A 10-year experience at a single institution. *Arch Intern Med* 152(4):791–796
 59. Sumida K, Ubara Y, Hoshino J, Suwabe T, Hiramatsu R, Hasegawa E et al (2010) Etanercept-refractory adult-onset Still's disease with thrombotic thrombocytopenic purpura successfully treated with tocilizumab. *Clin Rheumatol* 29(10):1191–1194, PubMed PMID: 20225049. Epub 2010/03/13. eng
 60. Cheema GS, Quismorio FP Jr (1999) Pulmonary involvement in adult-onset Still's disease. *Curr Opin Pulm Med* 5(5):305–309, PubMed PMID: 10461535
 61. Sari I, Birlık M, Binicier O, Akar S, Yilmaz E, Onen F et al (2009) A case of adult-onset Still's disease complicated with diffuse alveolar hemorrhage. *J Korean Med Sci* 24(1):155–157, PubMed PMID: 19270830. Pubmed Central PMCID: PMC2650997. Epub 2009/03/10. eng
 62. Efthimiou P, Georgy S (2006) Pathogenesis and management of adult-onset Still's disease. *Semin Arthritis Rheum* 36(3):144–152, PubMed PMID: 16949136. Epub 2006/09/05. eng
 63. Collard HR, Schwarz MI (2004) Diffuse alveolar hemorrhage. *Clin Chest Med* 25(3):583–592, vii. PubMed PMID: 15331194. Epub 2004/08/28. eng
 64. Schwarz MI, Fontenot AP (2004) Drug-induced diffuse alveolar hemorrhage syndromes and vasculitis. *Clin Chest Med* 25(1):133–140, PubMed PMID: 15062605. Epub 2004/04/06. eng
 65. Yamaguchi M, Ohta A, Tsunematsu T, Kasukawa R, Mizushima Y, Kashiwagi H et al (1992) Preliminary criteria for classification of adult Still's disease. *J Rheumatol* 19(3):424–430, PubMed PMID: 1578458
 66. Campos M, Schiopu E (2012) Pulmonary arterial hypertension in adult-onset Still's disease: rapid response to anakinra. Case reports in rheumatology 2012:537613, PubMed PMID: 22973530. Pubmed Central PMCID: PMC3437612. Epub 2012/09/14. eng
 67. Zen A, Yamashita N, Ueda M, Asakawa Y, Yoshikawa Y, Funai T et al (1990) A case of adult Still's disease with pulmonary hypertension. *Ryumachi* 30(1):45–52
 68. Chen CH, Chen HA, Wang HP, Liao HT, Chou CT, Huang DF (2006) Pulmonary arterial hypertension in autoimmune diseases: an analysis of 19 cases from a medical center in northern Taiwan. *Journal of microbiology, immunology, and infection = Wei mian yu gan ran za zhi* 39(2):162–168
 69. Mubashir E, Ahmed MM, Hayat S, Heldmann M, Berney SM (2007) Pulmonary hypertension in a patient with adult-onset Still's disease. *Clin Rheumatol* 26(8):1359–1361, PubMed PMID: 16871352. Epub 2006/07/28. eng
 70. Thakare M, Habibi S, Agrawal S, Narsimulu G (2013) Pulmonary arterial hypertension complicating adult-onset Still's disease. *Clin Rheumatol* 32(Suppl 1):S1–S2, PubMed PMID: 19669855. Epub 2009/08/12. eng
 71. Chatterjee S (2011) Pulmonary hypertension in systemic sclerosis. *Semin Arthritis Rheum* 41(1):19–37, PubMed PMID: 21047671. Epub 2010/11/05. eng
 72. Sanchez O, Humbert M, Sitbon O, Jais X, Simonneau G (2010) Guidelines for the diagnosis and treatment of pulmonary hypertension. *Rev Mal Respir* 27(2):141–150
 73. Hoeper MM (2002) Pulmonary hypertension in collagen vascular disease. *Eur Respir J* 19(3):571–576, PubMed PMID: 11936539. Epub 2002/04/09. eng
 74. Ngian GS, Stevens W, Prior D, Gabbay E, Roddy J, Tran A et al (2012) Predictors of mortality in connective tissue disease-associated pulmonary arterial hypertension: a cohort study. *Arthritis Res Ther* 14(5):R213, PubMed PMID: 23039366. Pubmed Central PMCID: PMC3580525. Epub 2012/10/09. Eng
 75. Sanchez O, Humbert M, Sitbon O, Simonneau G (1999) Treatment of pulmonary hypertension secondary to connective tissue diseases. *Thorax* 54(3):273–277, PubMed PMID: 10325906. Pubmed Central PMCID: PMC1745447. Epub 1999/05/18. eng
 76. Badesch DB, Hill NS, Burgess G, Rubin LJ, Barst RJ, Galie N et al (2007) Sildenafil for pulmonary arterial hypertension associated with connective tissue disease. *J Rheumatol* 34(12):2417–2422, PubMed PMID: 17985403. Epub 2007/11/07. eng
 77. Girgis RE, Frost AE, Hill NS, Hom EM, Langleben D, McLaughlin VV et al (2007) Selective endothelin A receptor antagonism with sitaxsentan for pulmonary arterial hypertension associated with connective tissue disease. *Ann Rheum Dis* 66(11):1467–1472, PubMed PMID: 17472992. Pubmed Central PMCID: PMC2111639. Epub 2007/05/03. eng
 78. Denton CP, Humbert M, Rubin L, Black CM (2006) Bosentan treatment for pulmonary arterial hypertension related to connective tissue disease: a subgroup analysis of the pivotal clinical trials and their open-label extensions. *Ann Rheum Dis* 65(10):1336–1340, PubMed PMID: 16793845. Pubmed Central PMCID: PMC1798307. Epub 2006/06/24. eng
 79. Oudiz RJ, Schilz RJ, Barst RJ, Galie N, Rich S, Rubin LJ et al (2004) Treprostinil, a prostacyclin analogue, in pulmonary arterial hypertension associated with connective tissue disease. *Chest* 126(2):420–427, PubMed PMID: 15302727. Epub 2004/08/11. eng
 80. Badesch DB, Tapson VF, McGoon MD, Brundage BH, Rubin LJ, Wigley FM et al (2000) Continuous intravenous epoprostenol for pulmonary hypertension due to the scleroderma spectrum of disease. A randomized, controlled trial. *Ann Intern Med* 132(6):425–434
 81. Franchini S, Dagna L, Salvo F, Aiello P, Baldissera E, Sabbadini MG (2010) Efficacy of traditional and biologic agents in different clinical phenotypes of adult-onset Still's disease. *Arthritis Rheum* 62(8):2530–2535, PubMed PMID: 20506370. Epub 2010/05/28. eng
 82. Fitzgerald AA, Leclercq SA, Yan A, Homik JE, Dinarello CA (2005) Rapid responses to anakinra in patients with refractory adult-onset Still's disease. *Arthritis Rheum* 52(6):1794–1803, PubMed PMID: 15934079
 83. Kishimoto T (1989) The biology of interleukin-6. *Blood* 74(1):1–10, PubMed PMID: 2473791. Epub 1989/07/01. eng
 84. Akira S, Tanga T, Kishimoto T (1993) Interleukin-6 in biology and medicine. *Adv Immunol* 54:1–78, PubMed PMID: 8379461. Epub 1993/01/01. eng
 85. Kishimoto T (2005) Interleukin-6: from basic science to medicine—40 years in immunology. *Annu Rev Immunol* 23:1–21, PubMed PMID: 15771564. Epub 2005/03/18. eng
 86. Chen DY, Lan JL, Hsieh TY et al (2004) Clinical manifestations, disease course, and complications of adult-onset Still's disease in Taiwan. *J Formos Med Assoc* 103:844–852
 87. Pouchot J, Sampalis JS, Beaudet F et al (1991) Adult Still's disease: manifestations, disease course, and outcome in 62 patients. *Medicine* 70:118–136
 88. Andres E, Kurtz JE, Perrin AE et al (2003) Retrospective monocentric study of 17 patients with adult Still's disease, with special focus on liver abnormalities. *Hepato-gastroenterology* 50:192–195
 89. Ishikawa M, Masumoto T, Oguni T, Hyoudou N, Michitaka K, Horiike N, Onji M (1997) A case of hemophagocytic syndrome with severe liver injury manifesting adult Still's disease. *Nihon Rinsho Meneki Gakkai Kaishi* 20:447–452

90. Walker NJ, Zurier RB (2002) Liver abnormalities in rheumatic diseases. *Clin Liver Dis* 6:933–946
91. Arlet JB, Le TH, Marinho A et al (2006) Reactive haemophagocytic syndrome in adult-onset Still's disease: a report of six patients and a review of the literature. *Ann Rheum Dis* 65:1596–1601
92. Ohta A, Yamaguchi M, Kaneoka H, Nagayoshi T, Hiida M (1987) Adult Still's disease: Review of 228 cases from the literature. *J Rheumatol* 14:1139
93. Wouters JMGW, van Rijswijk MH, van de Putte LBA (1985) Adult onset Still's disease in the elderly: a report of two cases. *J Rheumatol* 12:1791
94. Esdaile JM, Tannenbaum H (1980) Hawkins adult Still's disease. *Am J Med* 68:825
95. Tesser JRP, Pisko EJ, Hartz JW, Weinblatt ME (1982) Chronic liver disease and Still's disease. *Arthritis Rheum* 25:579
96. Dino O, Provenzano G, Giannuoli E, Sciarrino E, Pouyet M, Pagliaro L (1992) Fulminant hepatic failure in adult onset Still's disease. *J Rheum* 23:784–785
97. McFarlane M, Harth M, Wall WJ (1997) Liver transplant in adult Still's disease. *J Rheum* 24:2038–2041
98. Cush JJ, Medsger TA Jr, Christy WC, Herbert DC, Cooperstein LA (1987) Adult-onset Still's disease. Clinical course and outcome. *Arthritis Rheum* 30(2):186–194, PubMed PMID: 3827959
99. Reginato AJ, Schumacher HR Jr, Baker DG, O'Connor CR, Ferreiros J (1987) Adult onset Still's disease: experience in 23 patients and literature review with emphasis on organ failure. *Semin Arthritis Rheum* 17(1):39–57, PubMed PMID: 3306931
100. Taillan B, Fuzibet JG, Vinti H, Castela J, Ferrari E, Pesce A et al (1990) Non-infectious mitral endocarditis during Still's disease in adults. A case with histologic study of the endocardium. *Ann Med Interne (Paris)* 14(1):78–79
101. Cabane J, Michon A, Ziza JM, Bourgeois P, Bletry O, Godeau P et al (1990) Comparison of long term evolution of adult onset and juvenile onset Still's disease, both followed up for more than 10 years. *Ann Rheum Dis* 49(5):283–285, PubMed PMID: 2344206
102. Sayarlioglu M, Ucmak H, Ozkaya M, Ucar MA, Sayarlioglu H (2007) Leflunomide and methotrexate combination in refractory adult-onset Still's disease. *Ann Pharmacother* 41(7):1319–1320, PubMed PMID: 17535840. Epub 2007/05/31. eng
103. Cefle A (2005) Leflunomide and azathioprine combination in refractory adult-onset Still's disease. *Ann Pharmacother* 39(4):764–767, PubMed PMID: 15741418
104. Meaux-Ruault N, Magy N, Gil H, Dupond JL (2003) Efficacy of thalidomide in refractory adult Still's disease: a new case report. *Clin Exp Rheumatol* 21(2):272, PubMed PMID: 12747295
105. Pouchot J, Arlet J-B (2012) *Best Pract Res Clin Rheumatol* 26:477–87
106. Efthimiou P P, O. Rilonacept treatment of refractory adult onset Stills disease. *Arthritis & Rheumatism*. 2009 October 2009; 60.
107. Kraetsch HGAC, Kalden JR, Manger B (2002) Successful treatment of a small cohort of patients with adult onset of Still's disease with infliximab: first experiences. *Ann Rheum Dis* 61(9):859–860
108. Yang DH, Chang DM, Lai JH, Kuo SY, Ho TY, Lin KM et al (2008) Etanercept as a rescue agent in patient with adult onset Still's disease complicated with congestive heart failure. *Rheumatol Int* 29(1):95–98, PubMed PMID: 18500457. Epub 2008/05/27. eng
109. Benucci M, Li GF, Del Rosso A, Manfredi M (2005) Adalimumab (anti-TNF-alpha) therapy to improve the clinical course of adult-onset Still's disease: the first case report. *Clin Exp Rheumatol* 23(5):733, PubMed PMID: 16173267
110. Iwamoto M, Nara H, Hirata D, Minota S, Nishimoto N, Yoshizaki K (2002) Humanized monoclonal anti-interleukin-6 receptor antibody for treatment of intractable adult-onset Still's disease. *Arthritis and rheumatism* 46(12):3388–3389, PubMed PMID: 12483747
111. Ahmadi-Simab K, Lamprecht P, Jankowiak C, Gross WL (2006) Successful treatment of refractory adult onset Still's disease with rituximab. *Ann Rheum Dis* 65(8):1117–1118, PubMed PMID: 16837497. Pubmed Central PMCID: 1798247. Epub 2006/07/14. eng