

LIFETIME ASYMPTOTICS OF ITERATED BROWNIAN MOTION IN \mathbb{R}^n *

ERKAN NANE^{1,2}

Abstract. Let $\tau_D(Z)$ be the first exit time of iterated Brownian motion from a domain $D \subset \mathbb{R}^n$ started at $z \in D$ and let $P_z[\tau_D(Z) > t]$ be its distribution. In this paper we establish the exact asymptotics of $P_z[\tau_D(Z) > t]$ over bounded domains as an improvement of the results in DeBlassie (2004) [12] and Nane (2006) [24], for $z \in D$

$$\lim_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3}\lambda_D^{2/3}t^{1/3}\right) P_z[\tau_D(Z) > t] = C(z),$$

where $C(z) = (\lambda_D 2^{7/2})/\sqrt{3\pi} (\psi(z) \int_D \psi(y) dy)^2$. Here λ_D is the first eigenvalue of the Dirichlet Laplacian $\frac{1}{2}\Delta$ in D , and ψ is the eigenfunction corresponding to λ_D . We also study lifetime asymptotics of Brownian-time Brownian motion, $Z_t^1 = z + X(|Y(t)|)$, where X_t and Y_t are independent one-dimensional Brownian motions, in several unbounded domains. Using these results we obtain partial results for lifetime asymptotics of iterated Brownian motion in these unbounded domains.

Mathematics Subject Classification. 60J65, 60K99.

Received May 5, 2006. Revised September 8, 2006.

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

Iterated Brownian motion (IBM) has attracted the interest of several authors [1–3, 7–10, 12, 15, 18, 23, 24, 27, 29]. Several other iterated processes including Brownian-time Brownian motion (BTBM) have also been studied [1, 2, 19, 25, 26]. One of the main differences between these iterated processes and Brownian motion is that they are not Markov processes. However, these processes have many properties similar to that of Brownian motion (see [2, 3, 12, 23], and references therein).

To define iterated Brownian motion Z_t started at $z \in \mathbb{R}$, let X_t^+ , X_t^- and Y_t be three independent one-dimensional Brownian motions, all started at 0. Two-sided Brownian motion is defined by

$$X_t = \begin{cases} X_t^+, & t \geq 0 \\ X_{(-t)}^-, & t < 0. \end{cases}$$

Keywords and phrases. Iterated Brownian motion, Brownian-time Brownian motion, exit time, bounded domain, twisted domain, unbounded convex domain.

* Supported in part by NSF Grant # 9700585-DMS.

¹ Department of Mathematics, Purdue University, West Lafayette, IN 47906, USA; enane@math.purdue.edu

² Current address: Department of Statistics and Probability, Michigan State University, A413 Wells Hall, East Lansing, MI 48824-1027, USA; nane@stt.msu.edu

© EDP Sciences, SMAI 2007

Then iterated Brownian motion started at $z \in \mathbb{R}$ is

$$Z_t = z + X(Y_t), \quad t \geq 0.$$

In \mathbb{R}^n , one requires X^\pm to be independent n -dimensional Brownian motions. This is the version of the iterated Brownian motion due to Burdzy, see [7].

We next define another closely related process, the so called Brownian-time Brownian motion. Let X_t and Y_t be two independent one-dimensional Brownian motions, all started at 0. Brownian-time Brownian motion started at $z \in \mathbb{R}$ is

$$Z_t^1 = z + X(|Y_t|) \quad t \geq 0.$$

In \mathbb{R}^n one requires X to be an n -dimensional Brownian motion.

Let τ_D be the first exit time of Brownian motion from a domain $D \subset \mathbb{R}^n$. The large time behavior of $P_z[\tau_D > t]$ has been studied for several types of domains, including general cones [5, 11], parabola-shaped domains [4, 22], twisted domains [13], unbounded convex domains [21] and bounded domains [28]. Our aim in this article is to do the same for the first exit time of IBM from bounded domains in \mathbb{R}^n , and for the first exit time of BTBM from several domains in \mathbb{R}^n . See Bañuelos and DeBlassie [3], Li [21], Lifshits and Shi [22] and Nane [23] for a survey of results obtained for Brownian motion and iterated Brownian motion in these domains.

For many bounded domains $D \subset \mathbb{R}^n$ the asymptotics of $P_z[\tau_D > t]$ are well-known. (See [28] for a more precise statement of this.) For $z \in D$,

$$\lim_{t \rightarrow \infty} e^{\lambda_D t} P_z[\tau_D > t] = \psi(z) \int_D \psi(y) dy, \tag{1.1}$$

where λ_D is the first eigenvalue of $\frac{1}{2}\Delta$ with Dirichlet boundary conditions and ψ is its corresponding eigenfunction.

DeBlassie [12] proved that for iterated Brownian motion in bounded domains, for $z \in D$,

$$\lim_{t \rightarrow \infty} t^{-1/3} \log P_z[\tau_D(Z) > t] = -\frac{3}{2}\pi^{2/3} \lambda_D^{2/3}. \tag{1.2}$$

The limits (1.1) and (1.2) are very different in that the latter involves taking the logarithm which may kill many unwanted multipliers of the exponential. It is then natural to ask if it is possible to obtain an analogue of (1.1) for IBM. That is, can one remove the log in (1.2)? In [24], we improved the limit in (1.2) as follows, for $z \in D$,

$$\begin{aligned} 2C(z) &\leq \liminf_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3} \lambda_D^{2/3} t^{1/3}\right) P_z[\tau_D(Z) > t] \\ &\leq \limsup_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3} \lambda_D^{2/3} t^{1/3}\right) P_z[\tau_D(Z) > t] \leq \pi C(z), \end{aligned}$$

where $C(z) = \lambda_D \sqrt{2\pi/3} (\psi(z) \int_D \psi(y) dy)^2$.

In this paper we prove the following theorem which improves both limits above.

Theorem 1.1. *Let $D \subset \mathbb{R}^n$ be a domain for which (1.1) holds pointwise and let λ_D and ψ be as above. Then for $z \in D$,*

$$\lim_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3} \lambda_D^{2/3} t^{1/3}\right) P_z[\tau_D(Z) > t] = \frac{(\lambda_D 2^{7/2})}{\sqrt{3\pi}} \left(\psi(z) \int_D \psi(y) dy\right)^2.$$

Remark 1. Observe that $2\lambda_D \sqrt{2\pi/3} \leq (\lambda_D 2^{7/2})/\sqrt{3\pi} \leq \pi \lambda_D \sqrt{2\pi/3}$, so Theorem 1.1 is in agreement with the results obtained previously in [12, 24].

In [13], DeBlassie and Smits studied the tail behavior of the first exit time of Brownian motion in twisted domains in the plane. Let $D \subset \mathbb{R}^2$ be a domain whose boundary consists of three curves (in polar coordinates)

$$\begin{aligned} C_1 : \quad & \theta = f_1(r), \quad r \geq r_1 \\ C_2 : \quad & \theta = f_2(r), \quad r \geq r_1 \\ C_3 : \quad & r = r_1, \quad f_2(r) \leq \theta \leq f_1(r) \end{aligned}$$

where f_1 and f_2 are smooth and the curves C_1 and C_2 do not cross:

$$0 < f_1(r) - f_2(r) < \pi, \quad r \geq r_1.$$

DeBlassie and Smits call D a twisted domain if there are constants $r_0 > 0$, $\gamma > 0$ and $p \in (0, 1]$ and a smooth function $f(r)$ such that the curves $f_1(r)$ and $f_2(r)$, $r \geq r_0$, are obtained from $f(r)$ by moving out $\pm\gamma r^p$ units along the normal to the curve $\theta = f(r)$ at the point whose polar coordinates are $(r, f(r))$. They call γr^p the growth radius and $\theta = f(r)$ the generating curve. DeBlassie and Smits [13] (Th. 1.1) have the following tail behavior of the first exit time of Brownian motion in twisted domains $D \subset \mathbb{R}^2$ with growth radius γr^p , $\gamma > 0$, $0 < p < 1$

$$\lim_{t \rightarrow \infty} t^{-\left(\frac{1-p}{1+p}\right)} \log P_z[\tau_D > t] = -l_1 = - \left[\frac{\pi^{2p-1}}{\gamma^{2p}(1-p)^{2p}} \right]^{\frac{2}{p+1}} C_p \tag{1.3}$$

where

$$C_p = (1+p) \left[\frac{\pi^{2+p}}{8^p p^{2p} (1-p)^{1-p}} \frac{\Gamma^{2p}\left(\frac{1-p}{2p}\right)}{\Gamma^{2p}\left(\frac{1}{2p}\right)} \right]^{\frac{1}{p+1}}.$$

For these domains, Nane [23] obtained the following result for IBM: for all $z \in D$,

$$\lim_{t \rightarrow \infty} t^{-\left(\frac{1-p}{3+p}\right)} \log P_z[\tau_D(Z) > t] = - \left(\frac{3+p}{2+2p} \right) \left(\frac{1+p}{1-p} \right)^{\left(\frac{1-p}{3+p}\right)} \pi^{\left(\frac{2-2p}{3+p}\right)} l_1^{\left(\frac{2+2p}{3+p}\right)},$$

where l_1 is the limit given by (1.3).

We obtained in [24], the following for BTBM in twisted domains, for $z \in D$,

$$\lim_{t \rightarrow \infty} t^{-\left(\frac{1-p}{p+3}\right)} \log P_z[\tau_D(Z^1) > t] = -2^{\left(\frac{2p-2}{3+p}\right)} \left(\frac{3+p}{2+2p} \right) \left(\frac{1+p}{1-p} \right)^{\left(\frac{1-p}{3+p}\right)} \pi^{\left(\frac{2-2p}{3+p}\right)} l_1^{\left(\frac{2+2p}{3+p}\right)},$$

where l_1 is the limit given by the limit given by (1.3).

DeBlassie and Smits [13] also obtained similar results for $p = 1$. Let $D \subset \mathbb{R}^2$ be a twisted domain with growth radius γr , $\gamma > 0$. Then for $z \in D$,

$$\lim_{t \rightarrow \infty} [\log t]^{-1} \log P_z[\tau_D > t] = -C(\gamma) = -\pi \left[4 \arccos \frac{1}{\sqrt{1+\gamma^2}} \right]^{-1}. \tag{1.4}$$

We obtain the following lifetime asymptotics of BTBM in twisted domains.

Theorem 1.2. *Let $D \subset \mathbb{R}^2$ be a twisted domain with growth radius γr , $\gamma > 0$. Then for $z \in D$,*

$$\lim_{t \rightarrow \infty} [\log t]^{-1} \log P_z[\tau_D(Z^1) > t] = -C(\gamma)/2,$$

with $C(\gamma)$ as in (1.4).

Using Theorem 1.3. from [24], which says that for all $z \in D$ and all $t > 0$,

$$P_z[\tau_D(Z) > t] \leq 2P_z[\tau_D(Z^1) > t],$$

we obtain the following for IBM in twisted domains.

Corollary 1.1. *Let $D \subset \mathbb{R}^2$ be a twisted domain with growth radius γr , $\gamma > 0$. Then for $z \in D$,*

$$\limsup_{t \rightarrow \infty} [\log t]^{-1} \log P_z[\tau_D(Z) > t] \leq -C(\gamma)/2,$$

with $C(\gamma)$ as in (1.4).

In [21], using Gaussian techniques, Li studied lifetime asymptotics of Brownian motion in domains of the following form

$$P_f = \{(x, y) \in \mathbb{R}^{n+1} : y > f(x), x \in \mathbb{R}^n\}$$

for $f(x) = \exp(|x|^p)$, $p > 0$. Li established that for $z \in P_f$,

$$\lim_{t \rightarrow \infty} t^{-1} (\log t)^{2/p} \log P_z[\tau_{P_f} > t] = -j_\nu^2, \quad (1.5)$$

where $\nu = (n-2)/2$ and j_ν is the smallest positive zero of the Bessel function J_ν .

We obtain the following theorem in these domains

Theorem 1.3. *Let P_f be as above with $f(x) = \exp(|x|^p)$, $p > 0$. Then for $z \in P_f$,*

$$\lim_{t \rightarrow \infty} t^{-1/3} (\log t)^{4/3p} \log P_z[\tau_{P_f}(Z^1) > t] = -C(p),$$

where

$$C(p) = (3/2)^{(4+3p)/3p} 2^{1/3} (j_\nu^2 2^{2/p})^{2/3} (\pi^2/8)^{1/3}.$$

Using Theorem 1.3. from [24], we obtain the following for IBM in these domains.

Corollary 1.2. *Let P_f be as above with $f(x) = \exp(|x|^p)$, $p > 0$. Then for $z \in P_f$,*

$$\limsup_{t \rightarrow \infty} t^{-1/3} (\log t)^{4/3p} \log P_z[\tau_{P_f}(Z) > t] \leq -C(p).$$

For $f(x) = h(|x|)$ and $h^{-1}(x) = Ax^\alpha (\log x)^\beta$, $x > 1$. Li obtained the following: let $\epsilon > 0$. For t large, $z \in P_f$

$$-(1+\epsilon)C_{\alpha,\beta,1} \leq t^{-\frac{(1-\alpha)}{(1+\alpha)}} (\log t)^{\frac{2\beta}{(1+\alpha)}} \log P_z[\tau_{P_f} > t] \leq -(1-\epsilon)C_{\alpha,\beta,2}, \quad (1.6)$$

where

$$C_{\alpha,\beta,1} = 2^{-1} (1-\alpha)^{-1} (\alpha^{-\alpha} (1+\alpha)^{2\beta+2} A^{-2} j_\nu^2) 1/(1+\alpha)$$

and

$$C_{\alpha,\beta,2} = (1+\alpha) (2\alpha)^{-\alpha/(1+\alpha)} (2^{-1} (1+\alpha))^{2\beta/(1+\alpha)} C^{1/(1+\alpha)}$$

where $C = (1-\alpha)^{-1} 2^{2\beta-1} A^{-2} j_\nu^2$.

We have the following for BTBM in these domains.

Theorem 1.4. *For $0 < \alpha < 1$ and $\beta \in \mathbb{R}$,*

$$\begin{aligned} -C(1) &\leq \liminf_{t \rightarrow \infty} t^{-(1-\alpha)/(3+\alpha)} (\log t)^{4\beta(1+\alpha)/(3+\alpha)} \log P_z[\tau_{P_f}(Z^1) > t] \\ &\leq \limsup_{t \rightarrow \infty} t^{-(1-\alpha)/(3\alpha+1)} (\log t)^{4\beta(1+\alpha)/(3+\alpha)} \log P_z[\tau_{P_f}(Z^1) > t] \\ &\leq -C(2) \end{aligned}$$

where

$$C(1) = \left(\frac{3 + \alpha}{2(1 + \alpha)} \right)^{\frac{3 + \alpha + 4\beta}{3 + \alpha}} \left(\frac{1 - \alpha}{3 + \alpha} \right) (\pi^2 / 8)^{\frac{1 - \alpha}{(3 + \alpha)}} (C_{\alpha, \beta, 1})^{\frac{2(1 + \alpha)}{(3 + \alpha)}} 2^{\frac{4\beta}{(3 + \alpha)}},$$

and

$$C(2) = \left(\frac{3 + \alpha}{2(1 + \alpha)} \right)^{\frac{3 + \alpha + 4\beta}{3 + \alpha}} \left(\frac{1 - \alpha}{3 + \alpha} \right) (\pi^2 / 8)^{\frac{1 - \alpha}{(3 + \alpha)}} (C_{\alpha, \beta, 2})^{\frac{2(1 + \alpha)}{(3 + \alpha)}} 2^{\frac{4\beta}{(3 + \alpha)}}.$$

Using Theorem 1.3. from [24], we obtain the following for IBM in these domains.

Corollary 1.3. For $0 < \alpha < 1$ and $\beta \in \mathbb{R}$. Let P_f be as above with $f(x) = h(|x|)$, $h^{-1}(x) = Ax^\alpha(\log x)^\beta$, $x > 1$. Then for $z \in P_f$,

$$\limsup_{t \rightarrow \infty} t^{-(1-\alpha)/(3\alpha+1)} (\log t)^{4\beta(1+\alpha)/(3+\alpha)} \log P_z[\tau_{P_f}(Z^1) > t] \leq -C(2)$$

where $C(2)$ is as above.

The paper is organized as follows. In Section 2, we give some preliminary lemmas to be used in the proof of main results. We also recall several asymptotic results from Nane [24] to be used in the proof of main results. Theorem 1.1 is proved in Section 3. Section 4 is devoted to prove Theorems 1.2, 1.3 and 1.4.

2. PRELIMINARIES

In this section we state some preliminary facts that will be used in the proof of main results.

In what follows we will write $f \approx g$ and $f \lesssim g$ to mean that for some positive C_1 and C_2 , $C_1 \leq f/g \leq C_2$ and $f \leq C_1g$, respectively. We will also write $f(t) \sim g(t)$, as $t \rightarrow \infty$, to mean that $f(t)/g(t) \rightarrow 1$, as $t \rightarrow \infty$.

The main fact is the following Tauberian theorem ([14, Laplace transform method, 1958, Chapter 4]). Laporte [20] also studied these types of integrals. Let h and f be continuous functions on \mathbb{R} . Suppose f is non-positive and has a global max at x_0 , $f'(x_0) = 0$, $f''(x_0) < 0$ and $h(x_0) \neq 0$ and $\int_{-\infty}^{\infty} h(x) \exp(\lambda f(x)) < \infty$ for all $\lambda > 0$. Then as $\lambda \rightarrow \infty$,

$$\int_0^{\infty} h(x) \exp(\lambda f(x)) dx \sim h(x_0) \exp(\lambda f(x_0)) \sqrt{\frac{2\pi}{\lambda |f''(x_0)|}}. \tag{2.1}$$

It can be easily seen from this that as $\lambda \rightarrow \infty$,

$$\int_0^{\infty} \exp(-\lambda(x + x^{-2})) dx \sim \exp(-3\lambda 2^{-2/3}) \sqrt{\frac{2^{4/3}\pi}{3\lambda}}. \tag{2.2}$$

Similarly, as $t \rightarrow \infty$,

$$\int_0^{\infty} \exp\left(-\frac{at}{u^2} - bu\right) du \sim \sqrt{\frac{\pi}{3}} 2^{2/3} a^{1/6} b^{-2/3} t^{1/6} \exp(-3a^{1/3} b^{2/3} 2^{-2/3} t^{1/3}). \tag{2.3}$$

This follows from equation (2.2) after making the change of variables $u = (atb^{-1})^{1/3}x$.

Finally, we obtain, as $t \rightarrow \infty$,

$$\int_0^{\infty} u \exp\left(-\frac{at}{u^2} - bu\right) du \sim 2\sqrt{\frac{\pi}{3}} a^{1/2} b^{-1} t^{1/2} \exp(-3a^{1/3} b^{2/3} 2^{-2/3} t^{1/3}). \tag{2.4}$$

Writing the power series for the cosine function we easily see that as $t \rightarrow \infty$,

$$\int_0^{\infty} x \cos(\pi K/x) \exp\left(-\frac{\pi^2 t}{2x^2} - \lambda_D x\right) dx \sim 2\sqrt{\frac{\pi}{3}} \left(\frac{\pi^2}{2}\right)^{1/2} \lambda_D^{-1} t^{1/2} \exp\left(-\frac{3}{2}\pi^{2/3} \lambda_D^{2/3} t^{1/3}\right). \tag{2.5}$$

We next state a version of de Bruijn’s Tauberian Theorem (Kasahara [17] Th. 3 and Bingham, Goldie and Teugels [6] p. 254).

Theorem 2.1 (de Bruijn’s Tauberian Theorem). *Let ξ be a positive random variable. Then, for $\alpha > 0$ and $\beta \in \mathbb{R}$*

$$\log P[\xi \leq \epsilon] \sim -C\epsilon^{-\alpha} |\log \epsilon|^\beta \quad \text{as } \epsilon \rightarrow 0^+$$

if and only if

$$\log E[\exp(-\lambda\xi)] \sim -(1 + \alpha)^{1-\beta/(1+\alpha)} \alpha^{-\alpha/(1+\alpha)} C^{1/(1+\alpha)} \lambda^{\alpha/(1+\alpha)} (\log \lambda)^{\beta/(1+\alpha)}$$

as $\lambda \rightarrow \infty$.

In the case $\alpha = 0$, if $x \rightarrow 0^+$

$$[|\log x|]^{-1} \log P[\xi \leq x] \sim -c/2.$$

Then as $\lambda \rightarrow \infty$

$$[\log \lambda]^{-1} \log E[\exp(-\lambda\xi)] \sim -c/2.$$

Next, we will recall some lemmas that will be used in Sections 3 and 4. The following lemma is proved in [12] Lemma A1 (it also follows from more general results on “intrinsic ultracontractivity”). We include it for completeness.

For $I \subset \mathbb{R}$ an open interval, we write

$$\eta_I = \eta(I) = \inf\{t \geq 0 : Y_t \notin I\}.$$

Lemma 2.1. *As $t \rightarrow \infty$,*

$$P_x[\eta_{(0,1)} > t] \sim \frac{4}{\pi} e^{-\frac{\pi^2 t}{2}} \sin \pi x, \quad \text{uniformly for } x \in (0, 1).$$

We recall a result from Nane [23] Lemma 6.2, that will be used for the process Z^1 .

Lemma 2.2. *Let $B = \{u > 0 : t/u^2 > M\}$ for M large. Then on B ,*

$$\frac{d}{du} P_0[\eta_{(-u,u)} > t] \sim \exp\left(-\frac{\pi^2 t}{8u^2}\right) \frac{\pi t}{u^3}. \tag{2.6}$$

3. ITERATED BROWNIAN MOTION IN BOUNDED DOMAINS

If $D \subset \mathbb{R}^n$ is an open set, write

$$\tau_D^\pm(z) = \inf\{t \geq 0 : X_t^\pm + z \notin D\},$$

and if $I \subset \mathbb{R}$ is an open interval, write

$$\eta_I = \eta(I) = \inf\{t \geq 0 : Y_t \notin I\}.$$

Recall that $\tau_D(Z)$ stands for the first exit time of iterated Brownian motion from D . As in DeBlassie [12, §3.], we have by the continuity of the paths for $Z_t = z + X(Y_t)$, if f is the probability density of $\tau_D^\pm(z)$

$$P_z[\tau_D(Z) > t] = \int_0^\infty \int_0^\infty P_0[\eta_{(-u,v)} > t] f(u) f(v) dv du. \tag{3.1}$$

The proof of Theorem 1.1. The following is well-known

$$P_0[\eta_{(-u,v)} > t] = \frac{4}{\pi} \sum_{n=0}^{\infty} \frac{1}{2n+1} \exp\left(-\frac{(2n+1)^2\pi^2}{2(u+v)^2}t\right) \sin \frac{(2n+1)\pi u}{u+v}, \quad (3.2)$$

(see Feller [16] pp. 340–342).

Let $\epsilon > 0$. From Lemma 2.1, choose $M > 0$ so large that

$$(1 - \epsilon) \frac{4}{\pi} e^{-\frac{\pi^2 t}{2}} \sin \pi x \leq P_x[\eta_{(0,1)} > t] \leq (1 + \epsilon) \frac{4}{\pi} e^{-\frac{\pi^2 t}{2}} \sin \pi x, \quad (3.3)$$

for $t \geq M$, uniformly $x \in (0, 1)$. For a bounded domain with regular boundary it is well-known (see [28] pp. 121–127) that there exists an increasing sequence of eigenvalues, $\lambda_1 < \lambda_2 \leq \lambda_3 \cdots$, and eigenfunctions ψ_k corresponding to λ_k such that,

$$P_z[\tau_D \leq t] = \sum_{k=1}^{\infty} \exp(-\lambda_k t) \psi_k(z) \int_D \psi_k(y) dy. \quad (3.4)$$

From the arguments in DeBlassie [12] Lemma A.4

$$f(t) = \frac{d}{dt} P_z[\tau_D \leq t] = \sum_{k=1}^{\infty} \lambda_k \exp(-\lambda_k t) \psi_k(z) \int_D \psi_k(y) dy. \quad (3.5)$$

Finally choose $K > 0$ so large that

$$A(z)(1 - \epsilon) \exp(-\lambda_D u) \leq f(u) \leq A(z)(1 + \epsilon) \exp(-\lambda_D u)$$

for all $u \geq K$, where

$$A(z) = \lambda_1 \psi_1(z) \int_D \psi_1(y) dy = \lambda_D \psi(z) \int_D \psi(y) dy.$$

We further assume that t is so large that $K < \frac{1}{2}\sqrt{t/M}$. Define A for $K > 0$ and $M > 0$ as

$$A = \left\{ (u, v) : K \leq u \leq \frac{1}{2}\sqrt{\frac{t}{M}}, u \leq v \leq \sqrt{\frac{t}{M}} - u \right\}.$$

By equation (3.3) and from equation (3.10) in [12],

$$\begin{aligned} P_z[\tau_D(Z) > t] &= 2 \int_0^\infty \int_u^\infty P_{\frac{u}{u+v}} \left[\eta_{(0,1)} > \frac{t}{(u+v)^2} \right] f(u) f(v) dv du \\ &\geq C^1 \int_K^{\frac{1}{2}\sqrt{t/M}} \int_u^{\sqrt{t/M}-u} \sin\left(\frac{\pi u}{u+v}\right) \exp\left(-\frac{\pi^2 t}{2(u+v)^2}\right) \exp(-\lambda_D(u+v)) dv du, \end{aligned}$$

where $C^1 = C^1(z) = 2(4/\pi)A(z)^2(1 - \epsilon)^3$. Changing the variables $x = u + v$, $z = u$ the integral is

$$= C^1 \int_K^{\frac{1}{2}\sqrt{t/M}} \int_{2z}^{\sqrt{t/M}} \sin\left(\frac{\pi z}{x}\right) \exp\left(-\frac{\pi^2 t}{2x^2}\right) \exp(-\lambda_D x) dx dz,$$

and reversing the order of integration

$$\begin{aligned} &= C^1 \int_{2K}^{\sqrt{t/M}} \int_K^{\frac{1}{2}x} \sin\left(\frac{\pi z}{x}\right) \exp\left(-\frac{\pi^2 t}{2x^2}\right) \exp(-\lambda_D x) dz dx \\ &= C^1/\pi \int_{2K}^{\sqrt{t/M}} x \cos\left(\frac{\pi K}{x}\right) \exp\left(-\frac{\pi^2 t}{2x^2}\right) \exp(-\lambda_D x) dx. \end{aligned}$$

From equation (2.5) as $t \rightarrow \infty$,

$$\int_0^\infty x \cos\left(\frac{\pi K}{x}\right) \exp\left(-\frac{\pi^2 t}{2x^2}\right) \exp(-\lambda_D x) dx \sim 2\sqrt{\frac{\pi}{3}} \left(\frac{\pi^2}{2}\right)^{1/2} \lambda_D^{-1} t^{1/2} \exp\left(-\frac{3}{2}\pi^{2/3} \lambda_D^{2/3} t^{1/3}\right). \quad (3.6)$$

Now for some $c_1 > 0$,

$$\int_0^{2K} x \exp\left(-\frac{\pi^2 t}{2x^2} - \lambda_D x\right) dx \leq e^{-\pi^2 t/2K^2} \int_0^{2K} x \exp(-\lambda_D x) dx \lesssim e^{-c_1 t}, \quad (3.7)$$

and

$$\int_{\sqrt{t/M}}^\infty x \exp\left(-\frac{\pi^2 t}{2x^2}\right) \exp(-\lambda_D x) dx \leq \int_{\sqrt{t/M}}^\infty x \exp(-\lambda_D x) dx = (\sqrt{t/M} \lambda_D^{-1} + \lambda_D^{-2}) \exp(-\lambda_D \sqrt{t/M}). \quad (3.8)$$

Now from equations (3.6)–(3.8) we get

$$\liminf_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3} \lambda_D^{2/3} t^{1/3}\right) P_z[\tau_D(Z) > t] \geq (C^1/\pi) 2\sqrt{\frac{\pi}{3}} \left(\frac{\pi^2}{2}\right)^{1/2} \lambda_D^{-1}. \quad (3.9)$$

For the upper bound for $P[\tau_D(Z) > t]$ from equation (3.10) in [12],

$$P_z[\tau_D(Z) > t] = 2 \int_0^\infty \int_u^\infty P_{\frac{u}{u+v}}[\eta_{(0,1)} > \frac{t}{(u+v)^2}] f(u) f(v) dv du. \quad (3.10)$$

We define the following sets that make up the domain of integration,

$$\begin{aligned} A_1 &= \{(u, v) : v \geq u \geq 0, u + v \geq \sqrt{t/M}\}, \\ A_2 &= \{(u, v) : u \geq 0, v \geq K, u \leq v, u + v \leq \sqrt{t/M}\}, \\ A_3 &= \{(u, v) : 0 \leq u \leq v \leq K\}. \end{aligned}$$

Over the set A_1 we have for some $c > 0$,

$$\int \int_{A_1} P_{\frac{u}{u+v}}[\eta_{(0,1)} > \frac{t}{(u+v)^2}] f(u) f(v) dv du \leq \int \int_{A_1} f(u) f(v) dv du \leq \exp(-c\sqrt{t/M}). \quad (3.11)$$

The equation (3.11) follows from the distribution of τ_D from equation (3.5).

Since on A_3 , $t/(u+v)^2 \geq M$,

$$\begin{aligned} & \int \int_{A_3} P_{\frac{u}{u+v}} \left[\eta_{(0,1)} > \frac{t}{(u+v)^2} \right] f(u)f(v)dvdu \\ & \lesssim \int_0^K \int_0^K \exp\left(-\frac{\pi^2 t}{2(u+v)^2}\right) f(u)f(v)dvdu \\ & \leq \exp\left(-\frac{\pi^2 t}{8K^2}\right) \int_0^K \int_0^K f(u)f(v)dvdu \leq \exp\left(-\frac{\pi^2 t}{8K^2}\right). \end{aligned} \quad (3.12)$$

Let $C_1 = C_1(z) = 2(4/\pi)A(z)^2(1+\epsilon)^3$. For the integral over A_2 we get,

$$\begin{aligned} & \int \int_{A_2} P_{\frac{u}{u+v}} \left[\eta_{(0,1)} > \frac{t}{(u+v)^2} \right] f(u)f(v)dvdu \\ & \leq C_{11} \int_0^K \int_K^{\sqrt{t/M}-u} f(u) \exp\left(-\frac{\pi^2 t}{2(u+v)^2} - \lambda_D v\right) dvdu \\ & + C_1 \int_K^{1/2\sqrt{t/M}} \int_u^{\sqrt{t/M}-u} \sin\left(\frac{\pi u}{u+v}\right) \exp\left(-\frac{\pi^2 t}{2(u+v)^2} - \lambda_D(u+v)\right) dvdu \\ & = I + II. \end{aligned} \quad (3.13)$$

Changing variables $u+v=z$, $u=w$

$$\begin{aligned} I & \approx \int_0^K \int_K^{\sqrt{t/M}-u} \exp\left(-\frac{\pi^2 t}{2(u+v)^2}\right) f(u) \exp(-\lambda_D v) dvdu \\ & \leq \int_0^K \int_{w+K}^{\sqrt{t/M}} \exp\left(-\frac{\pi^2 t}{2z^2}\right) f(w) \exp(-\lambda_D z) \exp(\lambda_D w) dz dw \\ & \leq \exp(\lambda_D K) \int_0^K f(w) dw \int_0^\infty \exp\left(-\frac{\pi^2 t}{2z^2}\right) \exp(-\lambda_D z) dz \\ & \lesssim t^{1/6} \exp\left(-\frac{3}{2}\pi^{2/3}\lambda_D^{2/3}t^{1/3}\right). \end{aligned} \quad (3.14)$$

Equation (3.14) follows from equation (2.3), with $a = \pi^2/2$, $b = \lambda_D$.

Changing variables $u+v=z$, $u=w$

$$\begin{aligned} II & \leq C_1 \int_K^{1/2\sqrt{t/M}} \int_{2w}^{\sqrt{t/M}} \sin\left(\frac{\pi w}{z}\right) \exp\left(-\frac{\pi^2 t}{2z^2} - \lambda_D z\right) dz dw \\ & = C_1 \int_{2K}^{\sqrt{t/M}} \int_K^{z/2} \sin\left(\frac{\pi w}{z}\right) \exp\left(-\frac{\pi^2 t}{2z^2} - \lambda_D z\right) dw dz \end{aligned} \quad (3.15)$$

$$\begin{aligned} & \leq C_1/\pi \int_{2K}^{\sqrt{t/M}} z \cos\left(\frac{\pi K}{z}\right) \exp\left(-\frac{\pi^2 t}{2z^2} - \lambda_D z\right) dz \\ & \leq (1+\epsilon)(C_1/\pi)2\sqrt{\frac{\pi}{3}}\left(\frac{\pi^2}{2}\right)^{1/2} \lambda_D^{-1} t^{1/2} \exp\left(-\frac{3}{2}\pi^{2/3}\lambda_D^{2/3}t^{1/3}\right). \end{aligned} \quad (3.16)$$

Equation (3.15) follows by changing the order of the integration. Equation (3.16) follows from equation (2.5).

Now from equations (3.11), (3.12), (3.14) and (3.16) we obtain

$$\limsup_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3}\lambda_D^{2/3}t^{1/3}\right) P_z[\tau_D(Z) > t] \leq (1 + \epsilon) \left(\frac{C_1}{\pi}\right) 2\sqrt{\frac{\pi}{3}} \left(\frac{\pi^2}{2}\right)^{1/2} \lambda_D^{-1}. \quad (3.17)$$

Finally, from equations (3.9) and (3.17) and letting $\epsilon \rightarrow 0$,

$$\begin{aligned} C(z) &\leq \liminf_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3}\lambda_D^{2/3}t^{1/3}\right) P_z[\tau_D(Z) > t] \\ &\leq \limsup_{t \rightarrow \infty} t^{-1/2} \exp\left(\frac{3}{2}\pi^{2/3}\lambda_D^{2/3}t^{1/3}\right) P_z[\tau_D(Z) > t] \leq C(z), \end{aligned}$$

where $C(z) = (\lambda_D 2^{7/2}) / \sqrt{3\pi} (\psi(z) \int_D \psi(y) dy)^2$. □

4. BROWNIAN-TIME BROWNIAN MOTION IN UNBOUNDED DOMAINS

In this section we study Brownian-time Brownian motion (BTBM), Z_t^1 started at $z \in \mathbb{R}$, in several unbounded domains.

If $D \subset \mathbb{R}^n$ is an open set, write

$$\tau_D(z) = \inf\{t \geq 0 : X_t + z \notin D\},$$

and if $I \subset \mathbb{R}$ is an open interval, we write

$$\eta_I = \inf\{t \geq 0 : Y_t \notin I\}.$$

Let $\tau_D(Z^1)$ stand for the first exit time of BTBM from D . We have by the continuity of paths

$$P_z[\tau_D(Z^1) > t] = P[\eta(-\tau_D(z), \tau_D(z)) > t]. \quad (4.1)$$

Proof of Theorem 1.2. Let $\epsilon > 0$. From Lemma 2.2, choose $M > 0$ so large that

$$(1 - \epsilon) \exp\left(-\frac{\pi^2 t}{8u^2}\right) \frac{\pi t}{u^3} \leq \frac{d}{du} P_0[\eta_{(-u, u)} > t] \leq (1 + \epsilon) \exp\left(-\frac{\pi^2 t}{8u^2}\right) \frac{\pi t}{u^3}$$

for all $u \leq \sqrt{t/M}$.

Let $C = C(\gamma)$. From the hypothesis choose $K > 0$ so large that

$$u^{-C(1+\epsilon)} \leq P(\tau_D(z) > u) \leq u^{-C(1-\epsilon)} \quad \text{for } u \geq K. \quad (4.2)$$

We further assume that t is so large that $K < \sqrt{t/M}$. Now by (4.1), after integration by parts,

$$\begin{aligned} P_z[\tau_D(Z^1) > t] &= \int_0^\infty \left(\frac{d}{du} P_0[\eta_{(-u, u)} > t]\right) P[\tau_D(z) > u] du \\ &\gtrsim t \int_K^{\sqrt{t/M}} \exp\left(-\frac{\pi^2 t}{8u^2}\right) u^{-(C(1+\epsilon)+3)} du \end{aligned} \quad (4.3)$$

and changing variables $u^{-2} = x$, $du = -1/2x^{-3/2}dx$ the integral is

$$\gtrsim t \int_{M/t}^{K^{-2}} \exp\left(-\frac{\pi^2 t x}{8}\right) x^{C(1+\epsilon)/2} dx. \quad (4.4)$$

Changing variables, $z = \pi^2 tx/8$, the integral is

$$\gtrsim t^{-C(1+\epsilon)/2} \int_{\pi^2 M/8}^{K^{-2}\pi^2 t/8} e^{-z} z^{C(1+\epsilon)/2} dz. \quad (4.5)$$

Now since for some $c_0 > 0$,

$$\begin{aligned} \int_{K^{-2}\pi^2 t/8}^{\infty} e^{-z} z^{C(1+\epsilon)/2} dz &\leq e^{-c_0 t}, \\ \int_0^{\pi^2 M/8} e^{-z} z^{C(1+\epsilon)/2} dz &< \infty, \end{aligned}$$

and

$$\int_0^{\infty} e^{-z} z^{C(1+\epsilon)/2} dz = \Gamma(1 + C(1 + \epsilon)/2).$$

We have

$$P_z[\tau_D(Z^1) > t] \gtrsim t^{-C(1+\epsilon)/2}. \quad (4.6)$$

We now give an upper bound.

$$\begin{aligned} P_z[\tau_D(Z^1) > t] &= \int_0^{\infty} P_0(\eta_{(-u,u)} > t) f(u) du \\ &\lesssim \int_0^{\sqrt{t/M}} e^{-\frac{\pi^2 t}{8u^2}} f(u) du + \int_{\sqrt{t/M}}^{\infty} f(u) du \\ &\lesssim E \left[\exp \left(-\frac{\pi^2 t}{8(\tau_D(z))^2} \right) \right] + (\sqrt{t/M})^{-C(1-\epsilon)} \\ &\lesssim t^{-C(1-\epsilon)/2}. \end{aligned} \quad (4.7)$$

Equation (4.7) follows from Theorem 2.1 and the asymptotics of $\tau_D(z)$.

Now from Equations (4.6) and (4.7) we have

$$t^{-C(1+\epsilon)/2} \lesssim P_z[\tau_D(Z^1) > t] \lesssim t^{-C(1-\epsilon)/2}.$$

Now taking logarithm of the above inequalities, dividing by $\log t$, letting $\epsilon \rightarrow 0$, we obtain the desired result. \square

Proof of Theorem 1.3. Let $\epsilon > 0$. From Lemma 2.2, choose $M > 0$ so large that

$$(1 - \epsilon) \exp \left(-\frac{\pi^2 t}{8u^2} \right) \frac{\pi t}{u^3} \leq \frac{d}{du} P_0[\eta_{(-u,u)} > t] \leq (1 + \epsilon) \exp \left(-\frac{\pi^2 t}{8u^2} \right) \frac{\pi t}{u^3} \quad (4.8)$$

for all $u \leq \sqrt{t/M}$.

Let $C = j_{\nu}^2$. Given $\epsilon > 0$, choose $K > 0$ so large that

$$\exp(-C(1 + \epsilon)u(\log u)^{-2/p}) \leq P(\tau_{P_f}(z) > u) \leq \exp(-C(1 - \epsilon)u(\log u)^{-2/p}) \quad (4.9)$$

for $u \geq K$. We further assume that t is so large that $K < \sqrt{t/M}$.

Then, by equations (4.8) and (4.9)

$$P_z[\tau_{P_f}(Z^1) > t] \gtrsim t \int_K^{\sqrt{t/M}} u^{-3} \exp \left(-\frac{\pi^2 t}{8u^2} \right) \exp(-C(1 + \epsilon)u(\log u)^{-2/p}) du \quad (4.10)$$

changing variables $u^{-2} = x$, $du = -1/2x^{-3/2}dx$ the integral is

$$\gtrsim t \int_{M/t}^{K^{-2}} \exp\left(-\frac{\pi^2 tx}{8}\right) \exp(-C(1+\epsilon)x^{-1/2}(\log x^{-1/2})^{-2/p}) dx.$$

Now we set

$$v(x) = C(1+\epsilon)x^{-1/2}(\log x^{-1/2})^{-2/p}$$

which gives

$$dv = C(1+\epsilon)x^{-3/2}(\log x^{-1/2})^{-2/p}[-1/2 + 1/p(\log x^{-1/2})^{-1}]dx = -Vdx.$$

Then the integral is

$$\gtrsim t \int_{M/t}^{K^{-2}} VV^{-1} \exp\left(-\frac{\pi^2 tx}{8}\right) \exp(-C(1+\epsilon)x^{-1/2}(\log x^{-1/2})^{-2/p}) dx.$$

Now for $x \geq M/t$

$$V^{-1} \gtrsim t^{-3/2}[1/2 - 1/p(\log \sqrt{t/M})^{-1}]^{-1} \gtrsim t^{-3/2}.$$

Hence the integral is

$$\gtrsim t^{-1/2} \int_{M/t}^{K^{-2}} \exp\left(-\frac{\pi^2 tx}{8}\right) \exp(-v(x))V(x)dx. \quad (4.11)$$

Now from de Bruijn's Tauberian Theorem 2.1 we have

$$\log \int_0^\infty \exp\left(-\frac{\pi^2 tx}{8}\right) \exp(-v(x))V(x)dx \sim -C(p, \epsilon)t^{1/3}(\log t)^{-4/(3p)}$$

where $C(p, \epsilon) = (3/2)^{(4+3p)/3p}2^{1/3}((1+\epsilon)j_\nu^2 2^{2/p})^{2/3}(\pi^2/8)^{1/3}$.

For some $c_1, c_2 > 0$ we have the following bounds:

$$\int_0^{M/t} \exp\left(-\frac{\pi^2 tx}{8}\right) \exp(-v(x))V(x)dx \leq e^{-c_1 t^{1/2}(\log t)^{-2/p}},$$

and

$$\int_{K^{-2}}^\infty \exp\left(-\frac{\pi^2 tx}{8}\right) \exp(-v(x))V(x)dx \leq e^{-c_2 t},$$

from which we get

$$P_z[\tau_{P_f}(Z^1) > t] \gtrsim \exp\left(-(1+\epsilon)^2 C(p, \epsilon)t^{1/3}(\log t)^{-4/3p}\right). \quad (4.12)$$

We next give the upper bound

$$\begin{aligned} P_z[\tau_{P_f}(Z^1) > t] &= \int_0^\infty P_0(\eta_{(-u,u)} > t)f(u)du \\ &\lesssim \int_0^{\sqrt{t/M}} e^{-\frac{\pi^2 t}{8u^2}} f(u)du + \int_{\sqrt{t/M}}^\infty f(u)du \\ &\lesssim E\left[\exp\left(-\frac{\pi^2 t}{8(\tau_{P_f}(z))^2}\right)\right] + \exp(-C(1-\epsilon)t^{1/2}(\log t)^{-2/p}). \end{aligned}$$

The upper bound follows from de Bruijn's Tauberian Theorem by observing from (1.5) that as $x \rightarrow 0^+$

$$\log P[1/(\tau_{P_f}(z))^2 \leq x] = \log P[\tau_{P_f}(z) \geq x^{-1/2}] \sim -Cx^{-1/2}(|\log x^{-1/2}|)^{-2/p}.$$

Hence

$$P_z[\tau_{P_f}(Z^1) > t] \lesssim \exp(-(1-\epsilon)D(p, \epsilon)t^{1/3}(\log t)^{-4/3p}) \quad (4.13)$$

where $D(p, \epsilon) = (3/2)^{(4+3p)/3p} 2^{1/3} ((1-\epsilon)j_\nu^2 2^{2/p})^{2/3} (\pi^2/8)^{1/3}$.

Therefore from equations (4.12) and (4.13), we obtain

$$\begin{aligned} & \exp(-(1+\epsilon)^2 C(p, \epsilon)t^{1/3}(\log t)^{-4/3p}) \\ & \lesssim P_z[\tau_{P_f}(Z^1) > t] \\ & \lesssim \exp(-(1-\epsilon)^2 D(p, \epsilon)t^{1/3}(\log t)^{-4/3p}). \end{aligned}$$

Now taking logarithms, multiplying by $t^{-1/3}(\log t)^{4/3p}$, letting $t \rightarrow \infty$ and then letting $\epsilon \rightarrow 0$, we get the desired result. \square

Proof of Theorem 1.4. The proof follows the same steps of the proof of Theorem 1.3, so we omit the details. \square

Acknowledgements. I would like to thank Professor Rodrigo Bañuelos, my academic advisor, for suggesting this problem to me and for his guidance on this paper.

REFERENCES

- [1] H. Allouba, Brownian-time processes: The pde connection and the corresponding Feynman-Kac formula. *Trans. Amer. Math. Soc.* **354** (2002) 4627–4637.
- [2] H. Allouba and W. Zheng, Brownian-time processes: The pde connection and the half-derivative generator. *Ann. Probab.* **29** (2001) 1780–1795.
- [3] R. Bañuelos and R.D. DeBlassie, The exit distribution for iterated Brownian motion in cones. *Stochastic Processes Appl.* **116** (2006) 36–69.
- [4] R. Bañuelos, R.D. DeBlassie and R. Smits, The first exit time of planar Brownian motion from the interior of a parabola. *Ann. Probab.* **29** (2001) 882–901.
- [5] R. Bañuelos, R. Smits, Brownian motion in cones. *Probab. Theory Relat. Fields* **108** (1997) 299–319.
- [6] N.H. Bingham, C.M. Goldie and J.L. Teugels, *Regular Variation*. Cambridge University Press, Cambridge (1987).
- [7] K. Burdzy, Some path properties of iterated Brownian motion, in *Seminar on Stochastic Processes*, E. Çinlar, K.L. Chung and M.J. Sharpe, Eds., Birkhäuser, Boston (1993) 67–87.
- [8] K. Burdzy, Variation of iterated Brownian motion, in *Workshops and Conference on Measure-valued Processes, Stochastic Partial Differential Equations and Interacting Particle Systems*, D.A. Dawson, Ed., Amer. Math. Soc. Providence, RI (1994) 35–53.
- [9] K. Burdzy and D. Khoshnevisan, Brownian motion in a Brownian crack. *Ann. Appl. Probabl.* **8** (1998) 708–748.
- [10] E. Csàki, M. Csörgő, A. Földes and P. Révész, The local time of iterated Brownian motion. *J. Theoret. Probab.* **9** (1996) 717–743.
- [11] R.D. DeBlassie, Exit times from cones in \mathbb{R}^n of Brownian motion. *Prob. Th. Rel. Fields* **74** (1987) 1–29.
- [12] R.D. DeBlassie, Iterated Brownian motion in an open set. *Ann. Appl. Probab.* **14** (2004) 1529–1558.
- [13] R.D. DeBlassie and R. Smits, Brownian motion in twisted domains. *Trans. Amer. Math. Soc.* **357** (2005) 1245–1274.
- [14] N.G. De Bruijn, *Asymptotic methods in analysis*. North-Holland Publishing Co., Amsterdam (1957).
- [15] N. Eisenbaum and Z. Shi, Uniform oscillations of the local time of iterated Brownian motion. *Bernoulli* **5** (1999) 49–65.
- [16] W. Feller, *An Introduction to Probability Theory and its Applications*. Wiley, New York (1971).
- [17] Y. Kasahara, Tauberian theorems of exponential type. *J. Math. Kyoto Univ.* **12** (1978) 209–219.
- [18] D. Khoshnevisan and T.M. Lewis, Stochastic calculus for Brownian motion in a Brownian fracture. *Ann. Applied Probabl.* **9** (1999) 629–667.
- [19] D. Khoshnevisan and T.M. Lewis, Chung's law of the iterated logarithm for iterated Brownian motion. *Ann. Inst. H. Poincaré Probab. Statist.* **32** (1996) 349–359.
- [20] O. Laporte, Absorption coefficients for thermal neutrons. *Phys. Rev.* **52** (1937) 72–74.
- [21] W. Li, The first exit time of a Brownian motion from an unbounded convex domain. *Ann. Probab.* **31** (2003) 1078–1096.
- [22] M. Lifshits and Z. Shi, The first exit time of Brownian motion from a parabolic domain. *Bernoulli* **8** (2002) 745–765.

- [23] E. Nane, Iterated Brownian motion in parabola-shaped domains. *Potential Analysis* **24** (2006) 105–123.
- [24] E. Nane, Iterated Brownian motion in bounded domains in \mathbb{R}^n . *Stochastic Processes Appl.* **116** (2006) 905–916.
- [25] E. Nane, Higher order PDE's and iterated processes. Accepted *Trans. Amer. Math. Soc.* math.PR/0508262.
- [26] E. Nane, Laws of the iterated logarithm for α -time Brownian motion. *Electron. J. Probab.* **11** (2006) 34–459 (electronic).
- [27] E. Nane, Isoperimetric-type inequalities for iterated Brownian motion in \mathbb{R}^n . Submitted, math.PR/0602188.
- [28] S.C. Port and C.J. Stone, *Brownian motion and Classical potential theory*. Academic, New York (1978).
- [29] Y. Xiao, Local times and related properties of multidimensional iterated Brownian motion. *J. Theoret. Probab.* **11** (1998) 383–408.