

## LIFTINGS OF SOME TYPES OF TENSOR FIELDS AND CONNECTIONS TO TANGENT BUNDLES OF $p^r$ -VELOCITIES

AKIHIKO MORIMOTO

### § Introduction.

In the previous paper [6], we studied the liftings of tensor fields to tangent bundles of higher order. The purpose of the present paper is to generalize the results of [6] to the tangent bundles  ${}^{r,p}TM$  of  $p^r$ -velocities in a manifold  $M$ — notions due to C. Ehresmann [1] (see also [2]). In §1, we explain the  $p^r$ -velocities in a manifold and define the  $(\lambda)$ -lifting of differentiable functions for any multi-index  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_p)$  of non-negative integers  $\lambda_i$  satisfying  $\sum \lambda_i \leq r$ . In §2, we construct  $\langle \lambda \rangle$ -lifts of any vector fields and  $(\lambda)$ -lifts of 1-forms. The  $\langle \lambda \rangle$ -lift is a little bit different from the  $(\lambda)$ -lift of vector fields in [6].

In §3, we construct  $(\lambda)$ -lifting of  $(0, q)$ -tensor fields and then  $(\lambda)$ -lifting of  $(1, q)$ -tensor fields to  ${}^{r,p}TM$  for  $q \geq 1$ . Unfortunately, the author could not construct a natural lifting of  $(s, q)$ -tensor fields to  ${}^{r,p}TM$  for  $s \geq 2$ .

Nevertheless, our  $(\lambda)$ -liftings of  $(s, q)$ -tensor fields for  $s = 0$  or  $1$  are quite sufficient for the geometric applications, because the important tensor fields with which we encounter so far in differential geometry seem to be, fortunately, only of type  $(s, q)$  with  $s = 0$  or  $1$ .

As an application, we shall consider in §4, the prolongations of almost complex structures and prove that if  $M$  is a (homogeneous) complex manifold, then  ${}^{r,p}TM$  is also a (homogeneous) complex manifold.

In §5, we consider the liftings of affine connections to  ${}^{r,p}TM$  and prove that if  $M$  is locally affine symmetric then  ${}^{r,p}TM$  is also locally affine symmetric with respect to the lifted affine connection.

In §6, we shall give a proof for the fact that if  $M$  is an affine symmetric space then  ${}^{r,p}TM$  is also an affine symmetric space.

---

Received June 17, 1969.

In this paper, all manifolds and mappings (functions) are assumed to be differentiable of class  $C^\infty$ , unless otherwise stated.

We shall fix two positive integers  $r$  and  $p$  throughout the paper.

**§1. Tangent bundles of  $p^r$ -velocities.**

Consider the algebra  $C^\infty(R^p)$  of all  $C^\infty$ -functions on the  $p$ -dimensional euclidean space  $R^p$  with natural coordinates  $(t_1, t_2, \dots, t_p)$ . For any  $p$ -tuple  $\nu = (\nu_1, \nu_2, \dots, \nu_p)$  of non-negative integers  $\nu_j$  we denote as usual by  $(\partial/\partial t)^\nu$  the following partial differentiation

$$(1.1) \quad \left(\frac{\partial}{\partial t}\right)^\nu f = \frac{\partial^{\nu_1+\dots+\nu_p} f}{\partial t_1^{\nu_1} \dots \partial t_p^{\nu_p}}$$

for  $f \in C^\infty(R^p)$ . We define  $|\nu|$  and  $\nu!$  as follows:

$$|\nu| = \nu_1 + \dots + \nu_p, \quad \nu! = \nu_1! \nu_2! \dots \nu_p!$$

We denote by  $N(p, r)$  the set of all  $p$ -tuples  $\nu = (\nu_1, \dots, \nu_p)$  of non-negative integers  $\nu_i$  such that  $|\nu| \leq r$ . The set  $N(p, r)$  is a subset of the additive group  $Z^p$  of all  $p$ -tuples of integers.

Take two elements  $f, g \in C^\infty(R^p)$ . We say  $f$  is  $r$ -equivalent to  $g$  if  $(\partial/\partial t)^\nu f = (\partial/\partial t)^\nu g$  at  $t = (t_1, \dots, t_p) = 0$  for every  $\nu \in N(p, r)$  and denote it by  $f \underset{r}{\sim} g$ . Clearly  $\underset{r}{\sim}$  is an equivalence relation in  $C^\infty(R^p)$ .

Now, let  $M$  be an  $n$ -dimensional manifold. Consider the set  $S_p(M)$  of all maps  $\varphi: R^p \rightarrow M$ . Take two elements  $\varphi, \psi \in S_p(M)$ . We say that  $\varphi$  is  $r$ -equivalent to  $\psi$  if  $f \circ \varphi \underset{r}{\sim} f \circ \psi$  for every  $f \in C^\infty(M)$  and denote it by  $\varphi \underset{r}{\sim} \psi$ . The relation  $\underset{r}{\sim}$  is also an equivalence relation in  $S_p(M)$ . We denote by  $\overset{r,p}{TM}$  the set of all equivalence classes in  $S_p(M)$  with respect to the equivalence relation  $\underset{r}{\sim}$ . We denote by  $[\varphi]_r$  the equivalence class containing  $\varphi \in S_p(M)$ , and we shall call it a  $p^r$ -velocity in  $M$  at  $\varphi(0)$ . To introduce the manifold structure in  $\overset{r,p}{TM}$ , we define local coordinate system on  $\overset{r,p}{TM}$  as follows: Take a coordinate neighborhood  $U$  in  $M$  with coordinate system  $\{x_1, x_2, \dots, x_n\}$ . Define the coordinate functions  $\{x_i^{(\nu)}\}$   $i = 1, \dots, n; \nu \in N(p, r)$  on  $\overset{r,p}{TU}$  by

$$(1.2) \quad x_i^{(\nu)}([\varphi]_r) = \frac{1}{\nu!} \left[ \left(\frac{\partial}{\partial t}\right)^\nu (x_i \circ \varphi) \right]_{t=0}$$

for  $[\varphi]_r \in \overset{r,p}{TU}$  (cf. (1.1)). It is straightforward to see that  $\overset{r,p}{TM}$  becomes a manifold by the above coordinate systems  $\{x_i^{(\nu)}\}$ . The projection  $\overset{r,p}{\pi}$  defined

by  $\pi^{r,q}([\varphi]_r) = \varphi(0)$  for  $\varphi \in S_p(M)$  is clearly a differentiable map of  $T^r M$  onto  $M$ .

**DEFINITION 1.1.** The differentiable manifold  $T^r M$  with projection  $\pi^{r,p}$  will be called *the tangent bundle of  $p^r$ -velocities in  $M$* .

**DEFINITION 1.2.** For any  $f \in C^\infty(M)$ , we define *the  $(\lambda)$ -lift  $f^{(\lambda)}$  of  $f$* , for every  $\lambda \in N(p, r)$ , as follows:

$$(1.3) \quad f^{(\lambda)}([\varphi]_r) = \frac{1}{\lambda!} \left[ \left( \frac{\partial}{\partial t} \right)^\lambda (f \circ \varphi) \right]_{t=0}$$

for  $[\varphi]_r \in T^r M$ . Clearly,  $f^{(\lambda)}$  is a well-defined differentiable function on  $T^r M$ . We note also that  $(x_i)^{(\nu)} = x_i^{(\nu)}$  holds on  $TU$  for the above coordinate system  $\{x_1, \dots, x_n\}$ .

For the sake of convenience we define  $f^{(\lambda)} = 0$  for any  $\lambda \in Z^p$  such that  $\lambda \notin N(p, r)$ .

**LEMMA 1.3.** The  $(\lambda)$ -lifting  $f \rightarrow f^{(\lambda)}$  is a linear map of  $C^\infty(M)$  into  $C^\infty(T^r M)$  and satisfies the following equality

$$(1.4) \quad (f \cdot g)^{(\lambda)} = \sum_{\mu \in Z^p} f^{(\mu)} \cdot g^{(\lambda-\mu)}$$

for every  $f, g \in C^\infty(M)$  and  $\lambda \in N(p, r)$ .

*Proof.* Straightforward verification similar to the one of Lemma 1.2 [6].

**§ 2. Liftings of vector fields and 1-forms.**

Let  $\mathcal{S}(M) = \sum \mathcal{S}_i^s(M)$  be, as in [6], the tensor algebra of all tensor fields on  $M$ .

**LEMMA 2.1.** For any  $X \in \mathcal{S}_i^s(M)$  and any  $\lambda \in N(p, r)$  there exists one and only one  $X^{<\lambda>} \in \mathcal{S}_i^s(T^r M)$  satisfying the following equality

$$(2.1) \quad X^{<\lambda>} f^{(\mu)} = (Xf)^{(\mu-\lambda)}$$

for every  $f \in C^\infty(M)$  and  $\mu \in N(p, r)$ .

*Proof.* Take a coordinate neighborhood  $U$  in  $M$  with coordinate system  $\{x_1, \dots, x_n\}$  and let  $X = \sum a_i \cdot \partial / \partial x_i$  ( $a_i \in C^\infty(U)$ ) be the local expression of  $X$  in  $U$ . Consider the vector field  $\tilde{X} = \tilde{X}_U$  on  $(\pi^{r,p})^{-1}(U)$  defined by

$$(2.2) \quad \tilde{X} = \sum_{\mu \in N(p,r)} \sum_{j=1}^n a_j^{(\mu-\lambda)} \frac{\partial}{\partial x_j^{(\mu)}}.$$

We see that  $\tilde{X}(x_j^{(\mu)}) = a_j^{(\mu-\lambda)} = (Xx_j)^{(\mu-\lambda)}$  for  $j = 1, 2, \dots, n; \mu \in N(p,r)$ . Now, making use of same arguments as in the proof of Lemma 1.4 [6], we can prove that  $\tilde{X}(f^{(\mu)}) = (Xf)^{(\mu-\lambda)}$  for every  $f \in C^\infty(U)$  and  $\mu \in N(p,r)$ . We can also prove that if  $U'$  is a coordinate neighborhood in  $M$  such that  $U \cap U' = U'' \neq \emptyset$ , then  $\tilde{X}_U|_{U''} = \tilde{X}_{U'}|_{U''}$  holds. Thus we obtain a vector field  $X^{<\lambda>}$  on  $\overset{r,p}{TM}$  such that  $X^{<\lambda>}|_{(\pi^{r,p})^{-1}(U)} = \tilde{X}_U$  for every coordinate neighborhood  $U$  in  $M$ . This vector field  $X^{<\lambda>}$  clearly satisfies the condition (2.1) for every  $f \in C^\infty(M)$  and  $\mu \in N(p,r)$ . The uniqueness of  $X^{<\lambda>}$  is also easily verified. Q.E.D.

**COROLLARY 2.2.** *Let  $\{x_1, \dots, x_n\}$  be a local coordinate system on a neighborhood  $U$  in  $M$ . Then, we have*

$$(2.3) \quad \left(\frac{\partial}{\partial x_i}\right)^{<\lambda>} = \frac{\partial}{\partial x_i^{(\lambda)}}$$

for every  $i = 1, \dots, n$  and  $\lambda \in N(p,r)$ .

*Proof.* Clear from the expression (2.2) of  $X^{<\lambda>}$  in  $(\pi^{r,p})^{-1}(U)$ .

**COROLLARY 2.3.** *Notations being as in Corollary 2.2, we have*

$$(2.4) \quad \left(\frac{\partial f}{\partial x_i}\right)^{(\lambda-\mu)} = \frac{\partial f^{(\lambda)}}{\partial x_i^{(\mu)}}$$

for every  $i = 1, \dots, n$  and  $\lambda, \mu \in N(p,r)$ .

*Proof.* By Corollary 2.2, we have

$$\frac{\partial f^{(\lambda)}}{\partial x_i^{(\mu)}} = \left(\frac{\partial}{\partial x_i}\right)^{<\mu>} f^{(\lambda)} = \left(\frac{\partial f}{\partial x_i}\right)^{(\lambda-\mu)}. \quad \text{Q.E.D.}$$

**DEFINITION 2.4.** The vector field  $X^{<\lambda>}$  in Lemma 2.1 will be called the  $\langle \lambda \rangle$ -lift of  $X$  to  $\overset{r,p}{TM}$  for  $\lambda \in N(p,r)$ . For the sake of convenience, we define  $X^{<\lambda>} = 0$  for every  $\lambda \in Z^p$  such that  $\lambda \notin N(p,r)$ . The  $\langle \lambda \rangle$ -lifting  $X \rightarrow X^{<\lambda>}$  is a linear map of  $\mathcal{S}_0^1(M)$  into  $\mathcal{S}_0^1(\overset{r,p}{TM})$  for every  $\lambda \in Z^p$ .

**LEMMA 2.5.** *For  $X, Y \in \mathcal{S}_0^1(M)$ , we have*

$$(2.5) \quad [X^{<\lambda>}, Y^{<\mu>}] = [X, Y]^{<\lambda+\mu>}$$

for every  $\lambda, \mu \in N(p,r)$ .

*Proof.* Assume  $\lambda + \mu \in N(p, r)$ . Then, for any  $g \in C^\infty(M)$  and  $\nu \in N(p, r)$  we have

$$\begin{aligned} [X^{<\lambda>}, Y^{<\mu>}]g^{(\nu)} &= X^{<\lambda>}Y^{<\mu>}g^{(\nu)} - Y^{<\mu>}X^{<\lambda>}g^{(\nu)} \\ &= X^{<\lambda>}(Yg)^{(\nu-\mu)} - Y^{<\mu>}(Xg)^{(\nu-\mu)} \\ &= X^{<\lambda>}(Yg)^{(\nu-\mu)} - Y^{<\mu>}(Xg)^{(\nu-\lambda)} \\ &= (XYg - YXg)^{(\nu-\lambda-\mu)} = ([X, Y]g)^{(\nu-\lambda-\mu)} \\ &= [X, Y]^{<\lambda+\mu>}g^{(\nu)}. \end{aligned}$$

Since  $g \in C^\infty(M)$  and  $\nu \in N(p, r)$  are arbitrary we get (2.5) if  $\lambda + \mu \in N(p, r)$ .

Assume  $\lambda + \mu \notin N(p, r)$ , then by our convention, we have  $[X, Y]^{<\lambda+\mu>} = 0$ . On the other hand, for any  $g \in C^\infty(M)$  and  $\nu \in N(p, r)$  we have, by the same calculation as above,

$$[X^{<\lambda>}, Y^{<\mu>}]g^{(\nu)} = (XYg)^{(\nu-\mu-\lambda)} - (YXg)^{(\nu-\lambda-\mu)} = 0, \text{ since } \nu - \mu - \lambda \notin N(p, r).$$

Thus (2.5) is verified in any case. Q.E.D.

LEMMA 2.6. For  $X \in \mathcal{S}^0(M)$  and  $f \in \mathcal{S}^0(M)$ , we have

$$(2.6) \quad (f \cdot X)^{<\lambda>} = \sum_{\nu \in N(p, r)} f^{(\nu)} \cdot X^{<\lambda+\nu>}$$

for every  $\lambda \in N(p, r)$ .

*Proof.* For any  $g \in \mathcal{S}^0(M)$  and  $\mu \in N(p, r)$ , we have

$$\begin{aligned} (f \cdot X)^{<\lambda>}g^{(\mu)} &= (fX \cdot g)^{(\mu-\lambda)} = (f \cdot Xg)^{(\mu-\lambda)} \\ &= \sum_{\nu \in \mathbb{Z}^p} f^{(\nu)} \cdot (Xg)^{(\mu-\lambda-\nu)} = \sum f^{(\nu)} \cdot X^{<\lambda+\nu>}g^{(\mu)} \\ &= (\sum f^{(\nu)} X^{<\lambda+\nu>})g^{(\mu)}. \end{aligned}$$

Since  $g$  and  $\mu$  are arbitrary, we get (2.6) for every  $\lambda \in N(p, r)$ . Q.E.D.

Remark 2.7. By our convention (cf. Def. 1.2) we can write (2.6) as follows:

$$(2.7) \quad (f \cdot X)^{<\lambda>} = \sum_{\nu \in \mathbb{Z}^p} f^{(\nu)} X^{<\lambda+\nu>}.$$

LEMMA 2.8. Let  $f_i, g_i \in C^\infty(M)$  ( $i = 1, \dots, k$ ) be such that  $\sum g_i df_i = 0$  on  $M$ . Then the following equality

$$(2.8) \quad \sum_{i=1}^k \sum_{\mu \in \mathbb{Z}^p} g_i^{(\mu)} df_i^{<\lambda-\mu>} = 0$$

holds on  $\overset{r,p}{TM}$  for every  $\lambda \in N(p, r)$ .

*Proof.* Similar to the proof of Lemma 2.1 [6].

Q.E.D.

LEMMA 2.9. *There is one and only one lifting  $L_\lambda: \mathcal{S}_1^0(M) \rightarrow \mathcal{S}_1^0(\overset{r,p}{TM})$  for every  $\lambda \in N(p, r)$  satisfying the following condition:*

$$(2.9) \quad L_\lambda(f \cdot dg) = \sum_{\lambda \in \mathbb{Z}^p} f^{(\mu)} dg^{(\lambda-\mu)}$$

for every  $f, g \in \mathcal{S}_1^0(M)$ .

*Proof.* Similar to the proof of Lemma 2.2 [6].

LEMMA 2.10. *For  $f \in \mathcal{S}_1^0(M)$  and  $\theta \in \mathcal{S}_1^0(M)$ , we have*

$$(2.10) \quad (f \cdot \theta)^{(\lambda)} = \sum_{\mu \in \mathbb{Z}^p} f^{(\mu)} \cdot \theta^{(\lambda-\mu)}$$

for every  $\lambda \in N(p, r)$ .

*Proof.* Similar to the proof of Corollary 2.4 [6].

LEMMA 2.11. *For  $\theta \in \mathcal{S}_1^0(M)$  and  $X \in \mathcal{S}_1^0(M)$ , we have*

$$(2.11) \quad \theta^{(\lambda)}(X^{<\mu>}) = (\theta(X))^{(\lambda-\mu)}$$

for every  $\lambda, \mu \in N(p, r)$ .

*Proof.* Let  $\theta = \sum f_i dx_i$  be the local expression of  $\theta$ . Making use of Lemma 2.1, we calculate as follows:

$$\begin{aligned} \theta^{(\lambda)}(X^{<\mu>}) &= (\sum f_i dx_i)^{(\lambda)}(X^{<\mu>}) \\ &= \sum_i \sum_{\nu \in \mathbb{Z}^p} f_i^{(\nu)} dx_i^{(\lambda-\nu)}(X^{<\mu>}) \\ &= \sum_i \sum_{\nu} f_i^{(\nu)}(X^{<\mu>} x_i^{(\lambda-\nu)}) = \sum_i \sum_{\nu} f_i^{(\nu)}(X x_i)^{(\lambda-\nu-\mu)} \\ &= \sum_i \sum_{\nu} f_i^{(\nu)}(dx_i(X))^{(\lambda-\nu-\mu)} \\ &= \sum (f_i \cdot dx_i(X))^{(\lambda-\mu)} = (\theta(X))^{(\lambda-\mu)}. \end{aligned}$$

Q.E.D.

### § 3. Lifting of $(1, q)$ -tensor fields.

Let  $\mathcal{S}_*(M)$  be the subalgebra of  $\mathcal{S}(M)$  consisting of all covariant tensor fields on  $M$ . We denote by  $\overset{r,p}{\mathcal{S}}_*M$  the  $m(r, p)$  times direct sum of  $\overset{r,p}{\mathcal{S}}_*(TM)$ , where  $m(r, p)$  denotes the number of elements in  $N(p, r)$ . i.e.

$$\mathcal{F}_*^{r,p}(M) = \sum_{q=0}^{\infty} \sum_{\lambda \in N(p,r)} (\mathcal{F}_q^{r,p}(TM))_{\lambda},$$

where  $(\mathcal{F}_q^{r,p}(TM))_{\lambda} = \mathcal{F}_q^{r,p}(TM)$  for all  $\lambda \in N(p,r)$ .

Take two elements  $\theta = (\theta^{\lambda})$  and  $\eta = (\eta^{\lambda})$  in  $\mathcal{F}_*^{r,p}(M)$ . We define the multiplication  $\theta \otimes \eta$  of  $\theta$  and  $\eta$  by the following:

$$(3.1) \quad (\theta \otimes \eta)^{\lambda} = \sum_{\mu, \lambda - \mu \in N(p,r)} \theta^{\mu} \otimes \eta^{\lambda - \mu}$$

for  $\lambda \in N(p,r)$ . We can readily see that  $\mathcal{F}_*^{r,p}(M)$  is an associative graded algebra over  $C^{\infty}(TM)$  by this multiplication  $\otimes$ .

We have defined, in Lemma 2.9, the lifting  $L_{\lambda}$  of  $\mathcal{F}_1^0(M)$  into  $\mathcal{F}_1^{r,p}(TM)$  for  $\lambda \in N(p,r)$ . Define  $L: \mathcal{F}_1^0(M) \rightarrow \mathcal{F}_*^{r,p}(TM)$  by  $L(\theta) = (L_{\lambda}(\theta))_{\lambda \in N(p,r)}$  for  $\theta \in T_1^0(M)$ .

**LEMMA 3.1.** *There exists one and only one homomorphism  $\tilde{L}: \mathcal{F}_*^0(M) \rightarrow \mathcal{F}_*^{r,p}(M)$  such that  $\tilde{L}|_{\mathcal{F}_1^0(M)} = L$ .*

*Proof.* Define  $L^q: (\mathcal{F}_1^0(M))^q \rightarrow \mathcal{F}_*^{r,p}(M)$  by

$$L^q(\theta_1, \dots, \theta_q) = L(\theta_1) \otimes \dots \otimes L(\theta_q)$$

for  $\theta_i \in \mathcal{F}_1^0(M)$   $i = 1, 2, \dots, q$ . Then,  $L$  is a multilinear map satisfying the following condition:

$$L^q(f_1\theta_1, \dots, f_q\theta_q) = L(f_1 \dots f_q) \otimes L^q(\theta_1, \dots, \theta_q)$$

for  $\theta_i \in \mathcal{F}_1^0(M)$  and  $f_i \in \mathcal{F}_0^0(M)$   $i = 1, \dots, q$ , from which we conclude that there is a linear map  $\tilde{L}^q$  of  $\mathcal{F}_q^{r,p}(M)$  into  $\mathcal{F}_*^0(M)$  such that

$$\tilde{L}^q(\theta_1 \otimes \dots \otimes \theta_q) = L(\theta_1) \otimes \dots \otimes L(\theta_q)$$

for  $\theta_i \in \mathcal{F}_1^0(M)$ ,  $i = 1, \dots, q$ . Thus  $\tilde{L}^q (q \geq 0)$  define a homomorphism  $\tilde{L}: \mathcal{F}_*^{r,p}(M) \rightarrow \mathcal{F}_*^0(M)$  such that  $\tilde{L}(\theta) = L(\theta)$  for  $\theta \in \mathcal{F}_1^0(M)$ . Q.E.D.

**DEFINITION 3.2.** For  $K \in \mathcal{F}_q^0(M)$  we denote by  $K^{(\lambda)}$  the  $\lambda$ -component of  $\tilde{L}(K)$  for  $\lambda \in N(p,r)$ , i.e.

$$\tilde{L}(K) = (K^{(\lambda)}).$$

We shall call  $K^{(\lambda)}$  the  $(\lambda)$ -lift of  $K$ . For the sake of convenience we put  $K^{(\lambda)} = 0$  for  $\lambda \in Z^p$  such that  $\lambda \notin N(p,r)$ .

LEMMA 3.3. *The notation  $\alpha_X^k$  being as in Lemma 3.7 [6], for any  $K \in \mathcal{S}_q^0(M)$  and  $X \in \mathcal{S}_0^1(M)$ , we have*

$$(3.2) \quad \alpha_X^{k < \lambda >} K^{(\mu)} = (\alpha_X^k K)^{(\mu - \lambda)}$$

for  $\lambda, \mu \in N(p, r)$ .

*Proof.* Using Lemma 2.11, we can prove the lemma in the same way as the one of Lemma 3.7 [6].

COROLLARY 3.4. *For  $K \in \mathcal{S}_q^0(M)$  and  $X_i \in \mathcal{S}_0^1(M)$   $i = 1, \dots, q$ , we have*

$$K^{(\lambda)}(X_1^{<\mu_1>}, \dots, X_q^{<\mu_q>}) = (K(X_1, \dots, X_q))^{(\lambda - \sum \mu_i)}$$

for every  $\lambda, \mu_i \in N(p, r)$ ,  $i = 1, \dots, q$ .

*Proof.* We use Lemma 3.3  $q$ -times. Q.E.D.

LEMMA 3.5. *For any  $K \in \mathcal{S}_q^1(M)$  and  $\nu \in N(p, r)$ , there is a unique  $\tilde{K} = K^{(\nu)} \in \mathcal{S}_q^{r,p}(\tilde{T}M)$  such that*

$$(3.3) \quad \tilde{K}(X_1^{<\lambda_1>}, \dots, X_q^{<\lambda_q>}) = (K(X_1, \dots, X_q))^{<\lambda + \nu>}$$

for every  $X_i \in \mathcal{S}_0^1(M)$  and  $\lambda_i \in N(p, r)$ , where  $\lambda = \sum_i \lambda_i$ .

*Proof.* Define  $\tilde{L}_\nu : \mathcal{S}_0^1(M) \times \mathcal{S}_q^0(M) \rightarrow \mathcal{S}_q^{r,p}(\tilde{T}M)$  by the following

$$(3.4) \quad \tilde{L}_\nu(X, T) = \sum_{\mu \in \mathbb{Z}^p} X^{<\mu + \nu>} \otimes T^{(\mu)}$$

for  $X \in \mathcal{S}_0^1(M)$  and  $T \in \mathcal{S}_q^0(M)$ . It is clear that  $\tilde{L}$  is a bilinear map over  $R$ . We now assert that the following

$$(3.5) \quad \tilde{L}_\nu(fX, T) = \tilde{L}_\nu(X, fT)$$

holds for every  $X \in \mathcal{S}_0^1(M)$ ,  $T \in \mathcal{S}_q^0(M)$  and  $f \in \mathcal{S}_q^0(M)$ . For, making use of Remark 2.7 and Lemma 3.1, we calculate as follows:

$$\begin{aligned} \tilde{L}_\nu(fX, T) &= \sum_{\mu} (fX)^{<\mu + \nu>} \otimes T^{(\mu)} \\ &= \sum_{\mu} \sum_{\lambda} f^{(\lambda)} X^{<\lambda + \mu + \nu>} \otimes T^{(\mu)} \\ &= \sum_{\mu} \sum_{\lambda'} f^{(\lambda' - \mu - \nu)} X^{<\lambda'>} \otimes T^{(\mu)} \\ &= \sum_{\lambda'} \sum_{\mu} X^{<\lambda'>} \otimes f^{(\lambda' - \mu - \nu)} T^{(\mu)} \end{aligned}$$



$$\begin{aligned} &= \sum_{\lambda'} X^{<\lambda'>} \otimes (fT)^{(\lambda'-\nu)} \\ &= \sum_{\lambda} X^{<\lambda+\nu>} \otimes (fT)^{(\lambda)} = \tilde{L}_\nu(X, fT), \end{aligned}$$

which proves our assertion. Thus, we obtain a linear map  $L_\nu$  of  $\mathcal{S}^{\frac{1}{q}}(M)$  into  $\mathcal{S}^{\frac{r,p}{q}}(TM)$  such that

$$L_\nu(X \otimes T) = \sum_{\mu \in \mathbb{Z}^p} X^{<\mu+\nu>} \otimes T^{(\mu)}$$

for  $X \in \mathcal{S}^{\frac{1}{q}}(M)$  and  $T \in \mathcal{S}^{\frac{0}{q}}(M)$ . Put  $\tilde{K} = L_\nu(K)$ . It is now sufficient to prove (3.3) for  $K = X \otimes T$  with  $X \in \mathcal{S}^{\frac{0}{q}}(M)$  and  $T \in \mathcal{S}^{\frac{0}{q}}(M)$ . Using Corollary 3.4 and Lemma 2.6, we can calculate as follows:

$$\begin{aligned} \tilde{K}(X_1^{<\lambda_1>}, \dots, X_q^{<\lambda_q>}) &= \sum_{\mu} T^{(\mu)}(X_1^{<\lambda_1>}, \dots, X_q^{<\lambda_q>}) X^{<\mu+\nu>} \\ &= \sum_{\mu} (T(X_1, \dots, X_q))^{(\mu-\lambda)} X^{<\mu+\nu>} \\ &= \sum_{\mu'} (T(X_1, \dots, X_q))^{(\mu')} X^{<\mu'+\nu+\lambda>} \\ &= (T(X_1, \dots, X_q) \cdot X)^{<\nu+\lambda>} = (K(X_1, \dots, X_q))^{<\nu+\lambda>}. \end{aligned}$$

The uniqueness of  $\tilde{K}$  is clear, since (3.3) holds for every  $X_i \in \mathcal{S}^{\frac{1}{q}}(M)$  and  $\lambda_i \in N(p, r)$ . Q.E.D.

**DEFINITION 3.6.** For  $K \in \mathcal{S}^{\frac{1}{q}}(M)$  and  $\nu \in N(p, r)$ , we denote  $\tilde{K}$  in Lemma 3.5 by  $\tilde{K} = K^{(\nu)}$  and call it *the  $(\nu)$ -lift of  $K$* , i.e.

$$(3.6) \quad K^{(\nu)}(X_1^{<\lambda_1>}, \dots, X_q^{<\lambda_q>}) = (K(X_1, \dots, X_q))^{<\lambda+\nu>}$$

for  $X_i \in \mathcal{S}^{\frac{1}{q}}(M)$ ,  $\lambda_i \in N(p, r)$ , where  $\lambda = \sum \lambda_i$ . We call  $K^{(0)}$  the complete lift of  $K$  to  $\mathcal{T}M$ .

**LEMMA 3.7.** For  $K \in \mathcal{S}^{\frac{1}{q}}(M)$  ( $q \geq 1$ ) and  $X \in \mathcal{S}^{\frac{1}{q}}(M)$ , we have

$$(3.7) \quad \alpha_X^k \langle \lambda \rangle K^{(\nu)} = (\alpha_X^k K)^{(\nu+\lambda)}$$

for  $k \leq q$  and  $\lambda, \mu \in N(p, r)$ .

*Proof.* It suffices to prove (3.7) for  $K = Y \otimes T$  with  $Y \in \mathcal{S}^{\frac{1}{q}}(M)$ ,  $T \in \mathcal{S}^{\frac{0}{q}}(M)$ . Using Lemma 3.3, we calculate as follows:

$$\begin{aligned} \alpha_X^k \langle \lambda \rangle K^{(\nu)} &= \alpha_X^k \langle \lambda \rangle \sum_{\nu} Y^{<\nu+\mu>} \otimes T^{(\nu)} \\ &= \sum_{\nu} Y^{<\nu+\mu>} \otimes \alpha_X^k \langle \lambda \rangle T^{(\nu)} \end{aligned}$$

$$\begin{aligned}
 &= \sum_{\nu} Y^{<\nu+\mu>} \otimes (\alpha_X^k T)^{(\nu-\lambda)} \\
 &= \sum Y^{<\nu'+\lambda+\mu>} \otimes (\alpha_X^k T)^{(\nu')} \\
 &= (Y \otimes \alpha_X^k T)^{(\lambda+\mu)} = (\alpha_X^k K)^{(\lambda+\mu)}
 \end{aligned}
 \tag{Q.E.D.}$$

COROLLARY 3.8. *We have*

$$\alpha_X^{k<0>} K^{(\mu)} = (\alpha_X^k K)^{(\mu)}$$

for every  $X \in \mathcal{F}_0^1(M)$ ,  $K \in \mathcal{F}_q^1(M)$  and  $\mu \in N(p, r)$ .

§4. Prolongations of almost complex structures.

LEMMA 4.1. *For any  $A, B \in \mathcal{F}_1^1(M)$ , we have*

$$(A \circ B)^{(0)} = A^{(0)} \circ B^{(0)}. \tag{4.1}$$

Let  $I_M \in \mathcal{F}_1^1(M)$  be the (1,1)-tensor field of identity transformations of tangent spaces to  $M$ . Then, we have

$$(I_M)^{(0)} = I_{p, \tau}^{TM} \tag{4.2}$$

*Proof.* Making use of (3.6), we have, for any  $X \in \mathcal{F}_0^1(M)$ ,

$$\begin{aligned}
 A^{(0)} \circ B^{(0)}(X^{<\lambda>}) &= A^{(0)}(B^{(0)}(X^{<\lambda>})) \\
 &= A^{(0)}((B(X))^{<\lambda>}) = (ABX)^{<\lambda>} \\
 &= ((A \circ B)X)^{<\lambda>} = (A \circ B)^{(0)}(X^{<\lambda>})
 \end{aligned}$$

for every  $\lambda \in N(p, r)$ . Therefore we get (4.1).

To prove (4.2), let  $I_M = \sum(\partial/\partial x_i) \otimes dx_i$  be the local expression of  $I_M$ , where  $\{x_i, \dots, x_n\}$  is a local coordinate system. Then, we have

$$\begin{aligned}
 (I_M)^{(0)} &= \sum_{i, \mu} \left(\frac{\partial}{\partial x_i}\right)^{<\mu>} \otimes (dx_i)^{(\mu)} \\
 &= \sum_{i, \mu} \frac{\partial}{\partial x_i^{(\mu)}} \otimes dx_i^{(\mu)} = I_{p, \tau}^{TM},
 \end{aligned}$$

which proves (4.2).

COROLLARY 4.2. *For any polynomial  $P(x)$  of one variable  $x$  with real coefficients and for any  $A \in \mathcal{F}_1^1(M)$ , we have*

$$(P(A))^{(0)} = P(A^{(0)}). \tag{4.3}$$

*Proof.* Use (4.1) and (4.2) repeatedly. Q.E.D.

**THEOREM 4.3.** *Let  $J$  be an almost complex structure on  $M$  with its Nijenhuis tensor  $N_J$ . Then, the bundle  ${}^{r,p}TM$  of  $p^r$ -velocities in  $M$  has an almost complex structure  $J^{(0)}$  with its Nijenhuis tensor  $(N_J)^{(0)}$ .*

**THEOREM 4.4.** *If a manifold  $M$  is a complex manifold with almost complex structure  $J$ , so is the bundle  ${}^{r,p}TM$  of  $p^r$ -velocities in  $M$  with almost complex structure  $J^{(0)}$ .*

**§ 5. Lifting of affine connections.**

Let  $\nabla$  be the covariant differentiation defined by an affine connection of  $M$ .

**THEOREM 5.1.** *There exists one and only one affine connection of  ${}^{r,p}TM$  whose covariant differentiation  $\tilde{\nabla}$  satisfies the following condition:*

$$(5.1) \quad \tilde{\nabla}_{X^{<\lambda>}} Y^{<\mu>} = (\nabla_X Y)^{<\lambda+\mu>}$$

for every  $X, Y \in \mathcal{S}_0^1(M)$  and  $\lambda, \mu \in N(p, r)$ .

*Proof.* Take a coordinate neighborhood  $U$  with coordinate system  $\{x_1, \dots, x_n\}$  and let  $\Gamma_{ij}^k$  be the connection components of  $\nabla$  with respect to  $\{x_1, \dots, x_n\}$ , i.e.

$$(5.2) \quad \nabla_{\frac{\partial}{\partial x_i}} \frac{\partial}{\partial x_j} = \sum_k \Gamma_{ij}^k \frac{\partial}{\partial x_k}$$

for  $i, j = 1, \dots, n$ . Let  $\Gamma'_{ij}^k$  be the connection components of  $\nabla$  with respect to another coordinate system  $\{y_1, \dots, y_n\}$  on  $U$ . Then, we have the following equalities:

$$(5.3) \quad \Gamma'_{ij}^k = \sum_{a,b,c} \frac{\partial x_b}{\partial y_i} \frac{\partial x_c}{\partial y_j} \frac{\partial y_k}{\partial x_a} \Gamma_{bc}^a + \sum \frac{\partial^2 x_a}{\partial y_i \partial y_j} \frac{\partial y_k}{\partial x_a}$$

for  $i, j, k = 1, 2, \dots, n$ . (cf. for instance [3] p. 27). Let  $\{x_i^{(\nu)} | i = 1, \dots, n; \nu \in N(p, r)\}$  (resp.  $\{y_i^{(\nu)}\}$ ) be the induced coordinate system on  $(\pi)^{-1}(U)$ . Define

$$(5.4) \quad \tilde{\Gamma}_{(i,\nu)(j,\mu)}^{(k,\lambda)} = (\Gamma_{ij}^k)^{(\lambda-\nu-\mu)}$$

for  $i, j, k = 1, 2, \dots, n; \lambda, \mu, \nu \in N(p, r)$ . We can now prove that there exists a connection  $\tilde{\nabla}$  whose connection components with respect to  $\{x_i^{(\nu)}\}$  are given

by (5.4). For, we can verify (5.5) [6] for  $\lambda, \mu, \nu \in N(p, r)$  in the same way as the proof of (5.5) [6], since we can use the equalities

$$\frac{\partial f^{(\lambda)}}{\partial x_i^{(\mu)}} = \left( \frac{\partial f}{\partial x_i} \right)^{(\lambda-\mu)}$$

for every  $\lambda, \mu \in N(p, r)$  and  $f \in C^\infty(U)$  (cf. Cor. 2.3).

Next, we shall verify the following

$$(5.5) \quad \tilde{\nabla}_{X_i^{<\lambda>} X_j^{<\mu>}} = (\nabla_{X_i} X_j)^{<\lambda+\mu>}$$

for every  $i, j = 1, \dots, n$  and  $\lambda, \mu \in N(p, r)$ , where we have put  $X_i = \frac{\partial}{\partial x_i}$ .

Making use of Lemma 2.6 we calculate as follows:

$$\begin{aligned} \tilde{\nabla}_{X_i^{<\lambda>} X_j^{<\mu>}} &= \tilde{\nabla}_{\frac{\partial}{\partial x_i^{<\lambda>}}} \left( \frac{\partial}{\partial x_j^{<\mu>}} \right) = \sum_{\nu, k} \tilde{\Gamma}_{(i, \lambda), (j, \mu)}^{(k, \nu)} \frac{\partial}{\partial x_k^{<\nu>}} \\ &= \sum_{\nu, k} (\Gamma_{ij}^k)^{(\nu-\lambda-\mu)} \frac{\partial}{\partial x_k^{<\nu>}} = \sum_{\nu, k} (\Gamma_{ij}^k)^{<\nu>} \left( \frac{\partial}{\partial x_k} \right)^{<\lambda+\mu+\nu>} \\ &= \left( \sum_k \Gamma_{ij}^k \frac{\partial}{\partial x_k} \right)^{<\lambda+\mu>} = (\nabla_{X_i} X_j)^{<\lambda+\mu>}. \end{aligned}$$

Now, we shall verify

$$(5.6) \quad \tilde{\nabla}_{(f \cdot X_i)^{<\lambda>} X_j^{<\mu>}} = (\nabla_{f X_i} X_j)^{<\lambda+\mu>}$$

for  $f \in C^\infty(U)$ ,  $i, j = 1, \dots, n$  and  $\lambda, \mu \in N(p, r)$ .

For, the left hand side of (5.6) is equal to

$$\begin{aligned} \tilde{\nabla}_{(f \cdot X_i)^{<\lambda+\nu>} X_j^{<\mu>}} &= \sum f^{<\nu>} \tilde{\nabla}_{X_i^{<\lambda+\nu>} X_j^{<\mu>}} \\ &= \sum f^{<\nu>} (\nabla_{X_i} X_j)^{<\lambda+\nu+\mu>} = (f \cdot \nabla_{X_i} X_j)^{<\lambda+\mu>} = (\nabla_{f X_i} X_j)^{<\lambda+\mu>}, \end{aligned}$$

which proves (5.6). Thus (5.1) is proved for  $Y = \frac{\partial}{\partial x_j}$  and for every  $X \in \mathcal{F}_0^1(M)$ .

Finally, we shall verify (5.1) for  $Y = \sum f_i X_j \in \mathcal{F}_0^1(M)$  as follows:

$$\begin{aligned} \tilde{\nabla}_{X^{<\lambda>} (\sum f_i X_j)^{<\mu>}} &= \tilde{\nabla}_{X^{<\lambda>}} \sum_{j, \nu} f_j^{<\nu>} X_j^{<\nu+\mu>} \\ &= \sum_{j, \nu} \{ f_j^{<\nu>} \tilde{\nabla}_{X^{<\lambda>}} X_j^{<\nu+\mu>} + X^{<\lambda>} f_j^{<\nu>} \cdot X_j^{<\nu+\mu>} \} \\ &= \sum_{j, \nu} \{ f_j^{<\nu>} (\nabla_X X_j)^{<\nu+\lambda+\mu>} + (X f_j)^{<\nu-\lambda>} X_j^{<\nu+\mu>} \} \\ &= \sum_j \{ (f_j \cdot \nabla_X X_j)^{<\lambda+\mu>} + (X f_j \cdot X_j)^{<\lambda+\mu>} \} \\ &= (\nabla_X (\sum f_j X_j))^{<\lambda+\mu>} \end{aligned}$$

The uniqueness of  $\tilde{\nabla}$  is clear, since (5.1) holds for every  $X, Y \in \mathcal{S}_0^1(M)$  and  $\lambda, \mu \in N(p, r)$ . Q.E.D.

DEFINITION 5.2. We denote  $\tilde{\nabla}$  in Theorem 5.1 by  $\tilde{\nabla} = \tilde{\nabla}^{r,p}$  and call it the complete lift of  $\nabla$  to  $\tilde{T}M$ .

PROPOSITION 5.3. Let  $\tilde{T}, \tilde{R}$  be the torsion and the curvature tensor field of  $\tilde{\nabla} = \tilde{\nabla}^{r,p}$ . Then we have

$$(5.7) \quad \tilde{T} = T^{(0)} \text{ and } R = \tilde{R},$$

where  $T^{(0)}$  and  $R^{(0)}$  are the complete lift of  $T$  and  $R$  (cf. Def. 3.6).

Proof. Using the relation (3.6), we calculate as follows:

$$\begin{aligned} T^{(0)}(X^{<\lambda>}, Y^{<\mu>}) &= (T(X, Y))^{<\lambda+\mu>} \\ &= (\nabla_X Y - \nabla_Y X - [X, Y])^{<\lambda+\mu>} \\ &= \tilde{\nabla}_{X^{<\lambda>}} Y^{<\mu>} - \tilde{\nabla}_{Y^{<\mu>}} X^{<\lambda>} - [X^{<\lambda>}, Y^{<\mu>}] = \tilde{T}(X^{<\lambda>}, Y^{<\mu>}) \end{aligned}$$

for every  $X, Y \in \mathcal{S}_0^1(M)$  and  $\lambda, \mu \in N(p, r)$ , which proves  $T^{(0)} = \tilde{T}$ .

Similarly, we have:

$$\begin{aligned} R^{(0)}(X^{<\lambda>}, Y^{<\mu>})Z^{<\nu>} &= (R(X, Y)Z)^{<\lambda+\mu+\nu>} \\ &= (\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X, Y]} Z)^{<\lambda+\mu+\nu>} \\ &= \tilde{\nabla}_{X^{<\lambda>}} \tilde{\nabla}_{Y^{<\mu>}} Z^{<\nu>} - \tilde{\nabla}_{Y^{<\mu>}} \tilde{\nabla}_{X^{<\lambda>}} Z^{<\nu>} - \tilde{\nabla}_{[X^{<\lambda>}, Y^{<\mu>}]} Z^{<\nu>} \\ &= \tilde{R}(X^{<\lambda>}, Y^{<\mu>})Z^{<\nu>} \end{aligned}$$

for every  $X, Y, Z \in \mathcal{S}_0^1(M)$  and  $\lambda, \mu, \nu \in N(p, r)$ , which proves  $R^{(0)} = \tilde{R}$ .

Q.E.D.

PROPOSITION 5.4. For any  $K \in \mathcal{S}_q^s(M)$  ( $s = 0$  or  $1$ ) and  $X \in \mathcal{S}_0^1(M)$ , we have

$$(5.8) \quad \tilde{\nabla}_{X^{<0>}} K^{(\mu)} = (\nabla_X K)^{(\mu)},$$

$$(5.9) \quad \tilde{\nabla} K^{(\mu)} = (\nabla K)^{(\mu)}$$

for every  $\mu \in N(p, r)$ .

Proof. It is sufficient to prove (5.8) for  $K = Y \otimes T$ , where  $Y \in \mathcal{S}_0^1(M)$ ,  $T \in \mathcal{S}_q^0(M)$ . Now, since  $K^{(\mu)} = \sum Y^{<\nu+\mu>} \otimes T^{(\nu)}$ , and since  $\tilde{\nabla}_{X^{<0>}}$  is a derivation of  $\tilde{\mathcal{S}}^{r,p}(TM)$ , it suffices to verify (5.8) in the special cases, where

$K = f \in \mathcal{S}^0(M)$  and  $K = Y \in \mathcal{S}^1_0(M)$  and  $K = \theta \in \mathcal{S}^0_1(M)$ .

If  $K = f$ , then we have

$$\check{\nabla}_{X^{<0>}} f^{(\mu)} = X^{<0>} f^{(\mu)} = (Xf)^{(\mu)} = (\nabla_X f)^{(\mu)}.$$

If  $K = Y$ , then we have

$$\check{\nabla}_{X^{<0>}} Y^{<\mu>} = (\nabla_X Y)^{<\mu>} = (\nabla_X Y)^{(\mu)}.$$

If  $K = \theta$ , then we have, for  $\mu, \nu \in N(p, r)$  and  $Y \in \mathcal{S}^1_0(M)$

$$\begin{aligned} (\check{\nabla}_{X^{<0>}} \theta^{(\mu)}) Y^{<\nu>} &= \check{\nabla}_{X^{<0>}} (\theta^{(\mu)} Y^{<\nu>}) - \theta^{(\mu)} (\check{\nabla}_{X^{<0>}} Y^{<\nu>}) \\ &= \check{\nabla}_{X^{<0>}} (\theta(Y))^{(\mu-\nu)} - \theta^{(\mu)} ((\nabla_X Y))^{<\nu>} \\ &= (\nabla_X \theta(Y))^{(\mu-\nu)} - (\theta(\nabla_X Y))^{(\mu-\nu)} \\ &= ((\nabla_X \theta) Y)^{(\mu-\nu)} = (\nabla_X \theta)^{(\mu)} (Y^{<\nu>}), \end{aligned}$$

and hence we get  $\check{\nabla}_{X^{<0>}} \theta^{(\mu)} = (\nabla_X \theta)^{(\mu)}$ .

To prove (5.9), using Corollary 3.8, we calculate as follows

$$\alpha_{X^{<0>}} \check{\nabla} K^{(\mu)} = \check{\nabla}_{X^{<0>}} K^{(\mu)} = (\nabla_X K)^{(\mu)} = (\alpha_X(\nabla K))^{(\mu)} = \alpha_{X^{<0>}} (\nabla K)^{(\mu)}.$$

Since  $(X^{<0>})_{[\varphi],r} (X \in \mathcal{S}^1_0(M))$  spans the tangent space to  $\overset{r,p}{T}M$  at  $[\varphi]_r \in \overset{r,p}{T}M$ , we conclude that (5.9) holds. Q.E.D.

Combining Proposition 5.3 and 5.4 we have proved the following

**THEOREM 5.5.** *Let  $T$  and  $R$  be the torsion and the curvature tensor field of an affine connection  $\nabla$  of  $M$ . According as  $T = 0$ ,  $R = 0$ ,  $\nabla R = 0$  or  $\nabla T = 0$ , we have  $T^{(0)} = 0$ ,  $\check{\nabla} T^{(0)} = 0$ ,  $R^{(0)} = 0$  or  $\check{\nabla} R^{(0)} = 0$ . In particular, if  $M$  is affine locally symmetric with respect to  $\nabla$ , so is  $\overset{r,p}{T}M$  with respect to  $\overset{r,p}{\nabla}$ .*

**§ 6. Affine symmetric spaces.**

Let  $\Phi : M \rightarrow N$  be a map of a manifold  $M$  into another manifold  $N$ . Then, the map  $\Phi$  induces a map  $\overset{r,p}{T}\Phi$  of  $\overset{r,p}{T}M$  into  $\overset{r,p}{T}N$  as follows:

$$(6.1) \quad (\overset{r,p}{T}\Phi)([\varphi]_r) = [\Phi \circ \varphi]_r$$

for  $[\varphi]_r \in \overset{r,p}{T}M$ . The map  $\overset{r,p}{T}\Phi$  is a well-defined differentiable map, which will be called the  $(r, p)$ -tangent to  $\Phi$ . It is clear that if  $\Phi$  is a diffeomorphism then  $\overset{r,p}{T}\Phi$  is also a diffeomorphism.

LEMMA 6.1. For any  $f \in C^\infty(N)$ , we have

$$(6.2) \quad f^{(\mu)} \circ \overset{r,p}{T}\Phi = (f \circ \Phi)^{(\mu)}$$

for every  $\mu \in N(p, r)$ .

*Proof.* Take a point  $[\varphi]_r \in \overset{r,p}{T}M$ . Then we have

$$\begin{aligned} (f^{(\mu)} \circ \overset{r,p}{T}\Phi)([\varphi]_r) &= f^{(\mu)}([\Phi \circ \varphi]_r) = \frac{1}{\mu!} \left[ \left( \frac{\partial}{\partial t} \right)^\mu (f \circ ([\Phi \circ \varphi])) \right]_{t=0} \\ &= \frac{1}{\mu!} \left[ \left( \frac{\partial}{\partial t} \right)^\mu ((f \circ \Phi) \circ \varphi) \right]_{t=0} = (f \circ \Phi)^{(\mu)}([\varphi]_r). \end{aligned} \quad \text{Q.E.D.}$$

LEMMA 6.2. Let  $\Phi : M \rightarrow N$  be a diffeomorphism of  $M$  onto  $N$ . Then for any  $X \in \mathcal{S}_0^1(M)$  we have

$$(6.3) \quad \overset{r,p}{T}\overset{r,p}{T}\Phi(X^{<\lambda>}) = (T\Phi X)^{<\lambda>}$$

for every  $\lambda \in N(p, r)$ .

*Proof.* Take a function  $f \in C^\infty(N)$ . Then, by making use of Lemma 6.1 and 2.1, we have, for any  $\mu \in N(p, r)$ :

$$\begin{aligned} \overset{r,p}{T}\overset{r,p}{T}\Phi(X^{<\lambda>})f^{(\mu)} &= X^{<\lambda>}(f^{(\mu)} \circ \overset{r,p}{T}\Phi) = X^{<\lambda>}(f \circ \Phi)^{(\mu)} \\ &= (X(f \circ \Phi))^{(\mu-\lambda)} = ((T\Phi X)f)^{(\mu-\lambda)} = (T\Phi X)^{<\lambda>}f^{(\mu)}. \end{aligned}$$

Since  $f \in C^\infty(N)$  and  $\mu \in N(p, r)$  are arbitrary, we get (6.3). Q.E.D.

LEMMA 6.3. Let  $\nabla$  (resp.  $\nabla'$ ) be an affine connection on  $M$  (resp.  $N$ ) and let  $\Phi : M \rightarrow N$  be a diffeomorphism transforming  $\nabla$  onto  $\nabla'$ , i.e. we have

$$T\Phi(\nabla_X Y) = \nabla'_{T\Phi X} T\Phi Y$$

for  $X, Y \in \mathcal{S}_0^1(M)$ . Then the map  $\overset{r,p}{T}\Phi$  transforms  $\overset{r,p}{\nabla}$  onto  $\overset{r,p}{\nabla}'$ .

*Proof.* Put  $\tilde{\Phi} = \overset{r,p}{T}\overset{r,p}{T}\Phi$ . It suffices to verify

$$(6.4) \quad \tilde{\Phi}\nabla_{X^{<\lambda>}} Y^{<\mu>} = \tilde{\nabla}_{\tilde{\Phi}X^{<\lambda>}} \tilde{\Phi}Y^{<\mu>}$$

for every  $X, Y \in \mathcal{S}_0^1(M)$  and  $\lambda, \mu \in N(p, r)$ . Now, by making use of Theorem 5.1 and Lemma 6.2, we see the left hand side of (6.7) is equal to

$$\begin{aligned} T\overset{r,p}{T}\Phi(\nabla_X Y)^{\langle\lambda+\mu\rangle} &= (T\Phi(\nabla_X Y))^{\langle\lambda+\mu\rangle} \\ &= (\nabla'_{T\phi_X} T\Phi Y)^{\langle\lambda+\mu\rangle} = \overset{r,p}{\nabla}'_{(T\phi_X)^{\langle\lambda\rangle}} (T\Phi Y)^{\langle\mu\rangle} = \overset{r,p}{\nabla}'_{\tilde{\phi}_X^{\langle\lambda\rangle}} \tilde{\Phi} Y^{\langle\mu\rangle}. \end{aligned}$$

Q.E.D.

LEMMA 6.4. Take a point  $x_0 \in M$  and let  $\Phi$  be a diffeomorphism of  $M$  onto itself such that  $\Phi(x_0) = x_0$  and that  $T_{x_0}\Phi = -1_{T_{x_0}M}$ . Consider the constant map  $\gamma_{x_0}$  of  $R^p$  into  $M$  defined by  $\gamma_{x_0}(u) = x_0$  for  $u \in R^p$ . Put  $\tilde{x}_0 = [\gamma_{x_0}]_r$ . Then, we have  $\overset{r,p}{T}\Phi(\tilde{x}_0) = \tilde{x}_0$  and that

$$(6.5) \quad T_{\tilde{x}_0} \overset{r,p}{T}\Phi = -1_{T_{\tilde{x}_0}(\overset{r,p}{T}M)}.$$

*Proof.* Take an element  $[\varphi]_1 \in T_{\tilde{x}_0}^{\overset{r,d}{T}}(\overset{r,p}{T}M)$ , where  $\varphi : R \rightarrow \overset{r,p}{T}M$  with  $\varphi(0) = \tilde{x}_0$ . Making use of the same arguments as in the proof of Lemma 1.1 [5], we can find a differentiable map  $\psi : R^{p+1} \rightarrow M$  such that  $\varphi(t) = [\psi_t]_r$  for small  $t$ , where we have put  $\psi_t(u) = \psi(t, u)$  for  $t \in R$  and  $u \in R^p$ . Put  $\psi^u(t) = \psi(t, u)$ . Then, since  $\varphi(0) = [\psi_0]_r = \tilde{x}_0 = [\gamma_{x_0}]_r$ , we can assume that  $\psi(0, u) = x_0$  for small  $u \in R^p$  (cf. the expression of  $(\tilde{\varphi})$  in the proof of Lemma 1.1 [5]). Take a coordinate neighborhood  $U$  of  $x_0$  with coordinate system  $\{x_1, \dots, x_n\}$ . Put  $x_{i,\nu} = x_i^{(\nu)}$  for  $i = 1, \dots, n$  and  $\nu \in N(p, r)$ . Then  $\{x_{i,\nu}\}$  is a coordinate system around  $\tilde{x}_0$ . We have to prove  $T\overset{r,p}{T}\Phi([\varphi]_1) = -[\varphi]_1$ , i.e. to prove  $[\overset{r,p}{T}\Phi \circ \varphi]_1 = -[\varphi]_1$ . To prove this, it suffices to prove the following

$$(6.6) \quad (x_{i,\nu})^{(1)}([\overset{r,p}{T}\Phi \circ \varphi]_1) = -(x_{i,\nu})^{(1)}([\varphi]_1)$$

for  $i = 1, 2, \dots, n$  and  $\nu \in N(p, r)$ .

Since  $(\overset{r,p}{T}\Phi \circ \varphi)(t) = \overset{r,p}{T}\Phi(\varphi(t)) = \overset{r,p}{T}\Phi([\psi_t]_r) = [\Phi \circ \psi_t]_r$ , we calculate as follows:

$$\begin{aligned} (x_{i,\nu})^{(1)}([\overset{r,p}{T}\Phi \circ \varphi]_1) &= \left[ \frac{\partial}{\partial t} (x_{i,\nu} \circ \overset{r,p}{T}\Phi \circ \varphi) \right]_{t=0} = \left[ \frac{\partial}{\partial t} (x_{i,\nu}([\Phi \circ \psi_t]_r)) \right]_{t=0} \\ &= \frac{1}{\nu!} \left[ \frac{\partial}{\partial t} \left( \left( \frac{\partial}{\partial u} \right)^\nu (x_i \circ \Phi \circ \psi_t) \right) \right]_{t=0} \\ &= \frac{1}{\nu!} \left[ \frac{\partial}{\partial t} \left( \left( \frac{\partial}{\partial u} \right)^\nu x_i(\Phi(\psi(t, u))) \right) \right]_{t=0} \\ &= \frac{1}{\nu!} \left[ \left( \frac{\partial}{\partial u} \right)^\nu \left( \left[ \frac{\partial}{\partial t} x_i(\Phi(\psi^u(t))) \right]_{t=0} \right) \right]_{u=0} \end{aligned}$$

Now, making use of our assumption  $T_{x_0}\Phi = -1_{T_{x_0}M}$  and the fact that  $\psi^u(0) =$



$\phi(0, u) = x_0$  for small  $u \in R^p$ , we have

$$\begin{aligned} \left[ \frac{\partial}{\partial t} x_i(\phi(\phi^u(t))) \right]_{t=0} &= x_i^{(1)}([\phi \circ \phi^u]_1) = x_i^{(1)}(T\phi[\phi^u]_1) \\ &= -x_i^{(1)}([\phi^u]_1) = -\left[ \frac{\partial}{\partial t} x_i(\phi^u(t)) \right]_{t=0}. \end{aligned}$$

Therefore, we can continue the above calculation as follows:

$$\begin{aligned} (x_{i,v})^{(1)}([\overset{r,p}{T}\phi \circ \varphi]_1) &= -\frac{1}{\nu!} \left[ \left( \frac{\partial}{\partial u} \right)^\nu \left( \left[ \frac{\partial}{\partial t} x_i \phi^u(t) \right]_{t=0} \right) \right]_{u=0} \\ &= -\frac{1}{\nu!} \left[ \frac{\partial}{\partial t} \left( \left( \frac{\partial}{\partial u} \right)^\nu (x_i \circ \phi_t) \right) \right]_{u=0} = -(x_{i,v})^{(1)}([\varphi]_1), \end{aligned}$$

which proves (6.6).

Q.E.D.

**COROLLARY 6.5.** *Let  $M$  be an affine symmetric space with affine connection  $\nabla$ . Let  $\phi : M \rightarrow M$  be the affine symmetry at a point  $x_0 \in M$ . Then the  $(r, p)$ -tangent  $\overset{r,p}{T}\phi$  to  $\phi$  is also the affine symmetry of  $\overset{r,p}{TM}$  with affine connection  $\overset{r,p}{\nabla}$  at the point  $\tilde{x}_0$ .*

*Proof.* Since  $\phi$  leaves  $\nabla$  invariant,  $\overset{r,p}{T}\phi$  also leaves  $\overset{r,p}{\nabla}$  invariant by Lemma 6.3. Next, since  $\phi$  is an affine symmetry we see that  $T_{x_0}\phi = -1_{T_{x_0}M}$ . Thus, by Lemma 6.4, we get (6.5), which means that  $\overset{r,p}{T}\phi$  is the affine symmetry at  $\tilde{x}_0$ . Q.E.D.

**LEMMA 6.6.** *Let  $\nabla$  be an affine connection on a manifold  $M$ , and let  $X \in \mathcal{S}_0^1(M)$  be an infinitesimal affine transformation of  $\nabla$ . Then, the  $\langle \lambda \rangle$ -lift  $X^{\langle \lambda \rangle}$  of  $X$  is also an infinitesimal affine transformation of  $\overset{r,p}{\nabla} = \overset{r,p}{\nabla}$  on  $\overset{r,p}{TM}$  for every  $\lambda \in N(p, r)$ .*

*Proof.* A necessary and sufficient condition for  $X$  to be an infinitesimal affine transformation of  $M$  is that

$$\mathcal{L}_X \circ \nabla_X - \nabla_X \circ \mathcal{L}_X = \nabla_{[X, Y]}$$

for every  $Y \in \mathcal{S}_0^1(M)$ , where  $\mathcal{L}_X$  denotes the Lie derivation with respect to  $X$ . Therefore, we have to prove the following

$$(6.7) \quad \mathcal{L}_{X^{\langle \lambda \rangle}} \circ \overset{r,p}{\nabla}_{\tilde{Y}} K - \overset{r,p}{\nabla}_{\tilde{Y}} \circ \mathcal{L}_{X^{\langle \lambda \rangle}} K = \overset{r,p}{\nabla}_{[X^{\langle \lambda \rangle}, \tilde{Y}]} K$$

for every  $K \in \overset{r,p}{\mathcal{S}}(TM)$  and  $\tilde{Y} \in \overset{r,p}{\mathcal{S}}_0^1(TM)$ . To prove (6.7) it suffices to prove (6.7) for the special cases, where  $\tilde{Y} = Y^{\langle \mu \rangle}$  with  $Y \in \mathcal{S}_0^1(M)$ ,  $\mu \in N(p, r)$  and  $K = Z^{\langle \nu \rangle}$  or  $\theta^{(\nu)}$  with  $Z \in \mathcal{S}_0^1(M)$ ,  $\theta \in \mathcal{S}_1^0(M)$  and  $\nu \in N(p, r)$ . Moreover, to prove (6.7) for the case  $K = \theta^{(\nu)}$  with  $\theta \in \mathcal{S}_1^0(M)$ , it suffices to prove (6.7)

for  $\theta = df$  with  $f \in \mathcal{F}_0^{\circ}(M)$ .

If  $K = Z^{<\nu>}$ , then we calculate as follows:

$$\begin{aligned} & \mathcal{L}_X^{<\lambda>} \tilde{\nabla}_Y^{<\mu>} Z^{<\nu>} - \tilde{\nabla}_Y^{<\mu>} \mathcal{L}_X^{<\lambda>} Z^{<\nu>} \\ &= [X^{<\lambda>}, (\nabla_Y Z)^{<\mu+\nu>}] - \tilde{\nabla}_Y^{<\mu>} [X^{<\lambda>}, Z^{<\nu>}] \\ &= [X, \nabla_Y Z]^{<\lambda+\mu+\nu>} - (\nabla_Y [X, Z])^{<\lambda+\mu+\nu>} \\ &= ([X, \nabla_Y Z] - \nabla_Y [X, Z])^{<\lambda+\mu+\nu>} = (\mathcal{L}_X \circ \nabla_Y) Z - (\nabla_Y \circ \mathcal{L}_X) Z^{<\lambda+\mu+\nu>} \\ &= (\nabla_{[X, Y]} Z)^{<\lambda+\mu+\nu>} = \nabla_{[X^{<\lambda>}, Y^{<\mu>}]} Z^{<\nu>}, \end{aligned}$$

which proves (6.7) for  $K = Z^{<\nu>}$ .

To prove (6.7) for the case  $K = df^{(\nu)}$  with  $f \in \mathcal{F}_0^{\circ}(M)$ , we first note that the following equalities hold:

$$(6.8) \quad (\mathcal{L}_X \theta)(Y) = X(\theta(Y)) - \theta([X, Y])$$

$$(6.9) \quad (\nabla_X(df))(Y) = XYf - (\nabla_X Y)f$$

for  $X, Y \in \mathcal{F}_0^{\circ}(M)$ ,  $f \in \mathcal{F}_0^{\circ}(M)$  and  $\theta \in \mathcal{F}_1^{\circ}(M)$ .

Take a vector field  $Z \in \mathcal{F}_0^{\circ}(M)$  and  $\rho \in N(p, r)$ . Making use of (6.8), (6.9), Lemma 2.5 and (5.1), we calculate as follows:

$$\begin{aligned} & \{ \mathcal{L}_X^{<\lambda>} (\tilde{\nabla}_Y^{<\mu>} (df^{(\nu)})) - \tilde{\nabla}_Y^{<\mu>} \mathcal{L}_X^{<\lambda>} (df^{(\nu)}) \} (Z^{<\rho>}) \\ &= X^{<\lambda>} ((\tilde{\nabla}_Y^{<\mu>} (df^{(\nu)})) (Z^{<\rho>})) - \tilde{\nabla}_Y^{<\mu>} df^{(\nu)} ([X^{<\lambda>}, Z^{<\rho>}]) \\ &\quad - (\tilde{\nabla}_Y^{<\mu>} d(X^{<\lambda>} f^{(\nu)})) (Z^{<\rho>}) \\ &= X^{<\lambda>} (Y^{<\mu>} Z^{<\rho>} f^{(\nu)} - (\tilde{\nabla}_Y^{<\mu>} Z^{<\rho>}) f^{(\nu)}) \\ &\quad - \{ Y^{<\mu>} [X^{<\lambda>}, Z^{<\rho>}] f^{(\nu)} - (\tilde{\nabla}_Y^{<\mu>} [X^{<\lambda>}, Z^{<\rho>}]) f^{(\nu)} \} \\ &\quad - \{ Y^{<\mu>} Z^{<\rho>} X^{<\lambda>} f^{(\nu)} - (\tilde{\nabla}_Y^{<\mu>} Z^{<\rho>}) X^{<\lambda>} f^{(\nu)} \} \\ &= [X\{YZf - (\nabla_Y Z)f\} - \{Y[X, Z]f - (\nabla_Y [X, Z])f - (\nabla_Y [X, Z])f\} \\ &\quad - \{YZXf - (\nabla_Y Z)Xf\}]^{(\nu-\mu-\rho-\lambda)} \\ &= [\{\mathcal{L}_X(\nabla_Y df) - \nabla_Y \mathcal{L}_X df\} (Z)]^{(\nu-\mu-\rho-\lambda)} \\ &= ((\nabla_{[X, Y]}(df))(Z))^{(\nu-\mu-\rho-\lambda)} = ([X, Y]Zf - (\nabla_{[X, Y]}Z)f)^{(\nu-\mu-\rho-\lambda)} \\ &= [X^{<\lambda>}, Y^{<\mu>}] Z^{<\rho>} f^{(\nu)} - (\nabla_{[X^{<\lambda>}, Y^{<\mu>}]} Z^{<\rho>}) f^{(\nu)} \\ &= (\nabla_{[X^{<\lambda>}, Y^{<\mu>}]} df^{(\nu)})(Z^{<\rho>}), \end{aligned}$$

which proves (6.7) for  $K = df^{(\nu)}$ , since  $Z \in \mathcal{F}_0^{\circ}(M)$  and  $\rho \in N(p, r)$  are arbitrary. Thus (6.7) holds for any  $K$  and  $\tilde{Y}$ . Q.E.D.

From Lemma 6.6 we obtain

PROPOSITION 6.7. *If the group of affine transformations of  $M$  with  $\nabla$  is transitive on  $M$ , then the group of affine transformations of  ${}^{r,p}TM$  with respect to  ${}^{r,p}\nabla$  is transitive on  ${}^{r,p}TM$ .*

From Proposition 6.7 and Corollary 6.5 we obtain the following

THEOREM 6.8. *If  $M$  is an affine symmetric space with connection  $\nabla$ , then  ${}^{r,p}TM$  is also an affine symmetric space with connection  ${}^{r,p}\nabla$ .*

§ 7. **Remarks.**

Let  $P(M, \pi, G)$  be a principal fibre bundle with base  $M$ , projection  $\pi$  and structure group  $G$ . We shall be able to prove that  ${}^{r,p}TP$  ( ${}^{r,p}TM, {}^{r,p}T\pi, {}^{r,p}TG$ ) becomes canonically a principal fibre bundle with structure group  ${}^{r,p}TG$ , which is a Lie group by the natural group multiplication. Let  $\omega$  be a connection form on  $P$ . Then by the same methods as in [5], we can construct the prolongation  $\omega^{(r,p)}$  of  $\omega$  to  ${}^{r,p}TP$ . If  $P = F(M)$  is the frame bundle of  $M$  then a linear connection on  $M$  will induce a linear connection on  ${}^{r,p}TM$  by the above procedure. We shall investigate the relationships between this procedure and the liftings of affine connections in § 5 in a forthcoming paper, where we shall also study the prolongations of  $G$ -structures to the tangent bundles of  $p^r$ -velocities, which will generalize the results in [4].

REFERENCES

[ 1 ] C. Ehresmann, Les prolongements d'une variété différentiable I. Calcul des jets, prolongement principal, C.R. Acad. Sci. Paris, **233**(1951), 598-600.  
 [ 2 ] —————, Les prolongements d'une variété différentiable II, L'espace des jets d'ordre  $r$  de  $V_n$  dans  $V_m$ , *ibid.*, 777-779.  
 [ 3 ] S. Helgason, Differential geometry and symmetric spaces, Acad. Press, 1962.  
 [ 4 ] A. Morimoto, Prolongations of  $G$ -structures to tangent bundles of higher order, Nagoya Math. J. **38**(1970), 153-179.  
 [ 5 ] —————, Prolongations of connections to tangential fibre bundles of higher order, to appear.  
 [ 6 ] —————, Liftings of tensor fields and connections to tangent bundles of higher order, to appear.

*Nagoya University*