



Published in final edited form as:

Nature. 2010 December 16; 468(7326): 921–926. doi:10.1038/nature09576.

## Light-avoidance-mediating photoreceptors tile the *Drosophila* larval body wall

Yang Xiang<sup>1</sup>, Quan Yuan<sup>1</sup>, Nina Vogt<sup>2</sup>, Loren L. Looger<sup>3</sup>, Lily Yeh Jan<sup>1</sup>, and Yuh Nung Jan<sup>1</sup>

<sup>1</sup> Howard Hughes Medical Institute, Departments of Physiology, Biochemistry, and Biophysics, University of California San Francisco, San Francisco, California 94158, USA

<sup>2</sup> Center for Developmental Genetics, New York University, New York, New York 10003, USA

<sup>3</sup> Howard Hughes Medical Institute, Janelia Farm Research Campus, Ashburn, Virginia 20147, USA

### Abstract

Photoreceptors for visual perception, phototaxis or light avoidance are typically clustered in eyes or related structures such as the Bolwig organ of *Drosophila* larvae. Unexpectedly, we found that the class IV dendritic arborization neurons of *Drosophila melanogaster* larvae respond to ultraviolet, violet and blue light, and are major mediators of light avoidance, particularly at high intensities. These class IV dendritic arborization neurons, which are present in every body segment, have dendrites tiling the larval body wall nearly completely without redundancy. Dendritic illumination activates class IV dendritic arborization neurons. These novel photoreceptors use phototransduction machinery distinct from other photoreceptors in *Drosophila* and enable larvae to sense light exposure over their entire bodies and move out of danger.

Light sensing is critical for animal life. Whereas image-forming visual perception allows animals to identify and track mates, predators and prey, non-image-forming functions regulate pupil reflex, phototaxis and circadian entrainment<sup>1,2</sup>. In addition to eyes<sup>1,2</sup>, extra-ocular photoreceptors exist<sup>1–5</sup>. For example, many eyeless or blinded animals can sense illumination of their body surfaces<sup>3–5</sup>. Birds possess deep-brain photoreceptors in their hypothalamus<sup>6</sup>, and extra-ocular photoreceptors are required for magnetic orientation of amphibians<sup>7</sup>. Recent studies demonstrate that eyeless animals such as *Caenorhabditis elegans* nonetheless have photoreceptors controlling light avoidance<sup>8–10</sup>.

*Drosophila* larvae spend most of the time feeding by digging into food. Light avoidance is a crucial behaviour to minimize body exposure. When tested in groups in a dark/light choice assay, *Drosophila* larvae prefer darkness<sup>11,12</sup>. This behaviour requires the pair of Bolwig organs on the larval head<sup>12</sup>; that is, primitive eye structures each comprised of 12

Correspondence and requests for materials should be addressed to Y.N.J. (yuhnung.jan@ucsf.edu).

Supplementary Information is linked to the online version of the paper at [www.nature.com/nature](http://www.nature.com/nature).

**Author Contributions** Y.X. designed and carried out the experiments and analysed the data; Q.Y. characterized molecular information of Gr28b, rhodopsin and cryptochrome. L.L.L. created GCaMP3 and did the bioinformatic analyses of Gr28b; N.V. cleaned up the *Rh3*<sup>1</sup> and *Rh4*<sup>1</sup> mutants; Y.N.J. helped to design the experiments and supervised the work; Y.X., L.L.L., L.Y.J. and Y.N.J. wrote the manuscript.

**Author Information** Reprints and permissions information is available at [www.nature.com/reprints](http://www.nature.com/reprints). The authors declare no competing financial interests. Readers are welcome to comment on the online version of this article at [www.nature.com/nature](http://www.nature.com/nature).

photoreceptors expressing Rh5 or Rh6, rhodopsins sensing blue and green light, respectively<sup>13</sup>.

## Cells besides Bolwig organs contribute to photoavoidance

We designed a photoavoidance assay for a single larva with sunlight-level intensities ( $-1 \text{ mW mm}^{-2}$  in San Francisco on a clear day in June, consistent with previous reports<sup>14</sup>). Wild-type *Drosophila* larvae showed avoidance of a white light spot of  $0.57 \text{ mW mm}^{-2}$  (Fig. 1a and Supplementary Movie 1). Surprisingly, similar avoidance (Fig. 1b and Supplementary Movie 2) was exhibited by larvae with their Bolwig organs ablated by the pro-apoptotic gene *Head involution defective* (*Hid*; also called *Wrinkled* (*W*))<sup>15</sup> expressed via the Bolwig-organ-specific promoter *Glass Multimer Reporter* (*GMR*)<sup>16</sup> (Supplementary Fig. 1a). Lower light intensities elicited less photoavoidance of wild-type animals, and even less of Bolwig-organ-ablated animals (Fig. 1c). However, at light intensities of  $0.57 \text{ mW mm}^{-2}$  or higher, *GMR-Hid* larvae showed avoidance comparable to wild-type animals ( $P > 0.05$ ). Thus, although the Bolwig organs are responsible for dim light avoidance and dark congregation<sup>12</sup>, *Drosophila* larvae must contain extra-ocular photoreceptors.

Testing the wavelength dependence of photoavoidance using bandpass filters letting through ultraviolet ( $-360 \text{ nm}$ ; Fig. 1d), violet ( $-402 \text{ nm}$ ; Fig. 1e), blue ( $-470 \text{ nm}$ ; Fig. 1f), green ( $-525 \text{ nm}$ ; Fig. 1g) or red light ( $-620 \text{ nm}$ ; Fig. 1h), we found that wild-type animals showed increased photoavoidance with higher light intensity (Fig. 1d–h), and were most sensitive to blue, violet and ultraviolet, and largely unresponsive to green and red light. Bolwig-organ-ablated animals showed less photoavoidance at low light intensity, but exhibited nearly normal avoidance response to high-intensity, short-wavelength light (Fig. 1d–f), demonstrating the existence of light-sensitive cells in addition to Bolwig organs. Because there was no detectable temperature increase associated with  $0.11 \text{ mW mm}^{-2}$  violet light (Supplementary Fig. 2)—which triggered avoidance in nearly 80% of the wild-type and *GMR-Hid* animals—and animals showed little response to high-intensity green or red light but strongly avoided low-intensity short-wavelength light (Fig. 1d–h), light avoidance probably involves wavelength-dependent photoreceptors but not local heating.

## Class IV neurons tiling larval body wall sense light

Given the report of diffusely distributed dermal photoreceptors triggering shadow reaction<sup>3–5</sup>, we tested whether sensory neurons in the larval body wall could be candidate photoreceptors. Using GCaMP3, a genetically encoded calcium indicator<sup>17–19</sup>, we found that blue light delivered for 5 s to the dorsal cluster (Fig. 2a, see Fig. 3c for whole larva image) generated a marked fluorescence increase specifically in the soma, axon and dendrites of ddaC, a class IV dendritic arborization neuron (Fig. 2b, e, f), but not in nearby sensory neurons (Fig. 2b, e, f). ddaC also responded to ultraviolet light (which also caused photo-bleaching), but not to green light (Fig. 2c–f). There were also  $\text{Ca}^{2+}$  increases specifically in class IV dendritic arborization neurons of the ventral and lateral cluster (V 'ada and VdaB, respectively) in response to ultraviolet and blue light, but not green light (Supplementary Figs 3 and 4). Similar GCaMP3 fluorescence responses were seen in class IV dendritic arborization neurons in body segments from head to tail (Supplementary Fig. 5).

Extracellular recording further revealed a progressive increase in action potential frequency when the dorsal class IV dendritic arborization neuron ddaC was illuminated with increasing intensity of blue light (Fig. 2g), ultraviolet light and violet light, but not red light (Supplementary Fig. 6). Responses were:  $340 \text{ nm} > 380 \text{ nm} > 402 \text{ nm} > 470 \text{ nm} \gg 525 \text{ nm}$  or  $620 \text{ nm}$  light (Fig. 2h).

The wavelength dependence of ddaC firing rate increase was similar to that observed with GCaMP3 imaging and the light avoidance behavioural assay. The latency between the onset of light stimulation and action potential burst firing decreased with higher light intensity, and was as short as 1 s with bright illumination (Supplementary Fig. 6). When illuminated with  $1.4 \text{ mW mm}^{-2}$  of white light (approximating sunlight), ddaC neurons in the dorsal cluster showed a significant firing increase (Fig. 2i). Similar robust activation of ventral (VdaB) and lateral (V'ada) class IV dendritic arborization neurons was induced by  $52.8 \text{ mW mm}^{-2}$  blue light (Supplementary Fig. 7a, b). The response of class IV dendritic arborization neurons was similar regardless of their location along the body axis (data not shown), as in the case of GCaMP3 imaging (Supplementary Fig. 5).

We did not observe any significant effects of light on firing rate of class I or III dendritic arborization neurons (Supplementary Fig. 7c, d,  $P > 0.05$ ). Because class I dendritic arborization neurons progressively increased their firing rate as the temperature was raised above  $30 \text{ }^{\circ}\text{C}$ , whereas class IV dendritic arborization neurons showed an abrupt increase of firing rate only above  $40 \text{ }^{\circ}\text{C}$  (Supplementary Fig. 8), thermal responses cannot account for the light-induced increase of firing in class IV but not class I dendritic arborization neurons. Moreover, application of  $10 \text{ }\mu\text{M H}_2\text{O}_2$ , which elevates the reactive oxygen species (ROS) level in *Drosophila* larvae<sup>20</sup>, had no effect on the firing rate of class IV dendritic arborization neurons (Supplementary Fig. 9). These studies demonstrate that ultraviolet, violet and blue light activate class IV dendritic arborization neurons in an intensity-dependent manner. Responses occur at sunlight-level intensities, are not induced by heat or ROS, correlate with behaviour, and are confined to this specific class of sensory neurons throughout the animal.

### Light activates class IV neurons and dendrites in isolation

The dendritic arborization neurons have dendrites in contact with epithelial cells whereas their somas and axons are wrapped by glia<sup>21</sup>. To test whether class IV dendritic arborization neurons can sense light by themselves, we prepared primary neuronal cultures<sup>22,23</sup> from embryos expressing GCaMP3 and RFP specifically in class IV dendritic arborization neurons by means of *pickpocket-GAL4 (ppk-GAL4)*<sup>24</sup>. Ultraviolet and blue light illumination of isolated class IV dendritic arborization neurons generated a robust increase of GCaMP3 signals (Fig. 3a and Supplementary Fig. 10). In contrast, cultured class III dendritic arborization neurons expressing GCaMP3 and RFP via *19-12-GAL4* yielded no light response (Fig. 3b). Thus, class IV dendritic arborization neurons have the intrinsic ability to detect light.

Dendrites of class IV dendritic arborization neurons tile the larval body wall with non-overlapping but complete coverage of the dendritic field<sup>25,26</sup> (Fig. 3c). Illumination of only the dendrites of class IV dendritic arborization neurons (Fig. 3d) with ultraviolet, violet and blue light, but not green or red light, activated the neurons (Fig. 3e). The activation spectrum is similar to that for illumination of the entire class IV dendritic arborization neurons (Fig. 2h), indicating the presence of phototransduction machinery in the dendrites.

### Gr28b is critical for light transduction in class IV neurons

No defects in light response of class IV dendritic arborization neurons were found in available mutants of rhodopsins<sup>13,27</sup> and cryptochrome (*cry*)<sup>28</sup>, as well as a mutant in *no receptor potential A (norpA)*, which encodes phospholipase C (PLC), downstream of rhodopsins<sup>29</sup> (Fig. 4a). We then tested the *Drosophila* homologue of Lite-1, a *C. elegans* light sensor<sup>8–10</sup>. The closest homologue of *lite-1* in *Drosophila* is gustatory receptor 28b (*Gr28b*), annotated as encoding a gustatory G-protein-coupled receptor. Several *Gr28b-GAL4* lines carrying different promoter regions revealed consistent expression in all class IV

dendritic arborization neurons, two sensory neurons in the lateral body wall, plus several neurons in the ventral nerve cord (Supplementary Fig. 11), as reported previously<sup>30</sup>. To test for the functional role of *Gr28b* in the light-induced electrophysiological responses, we recorded from class IV dendritic arborization neurons in *Dmel\Mi{ET1}Gr28b<sup>MB03888</sup>* (*MiET1*) and *Dmel\PBac{PB}Gr28b<sup>c01884</sup>* (*PBac*) larvae with P-element insertion into the *Gr28b* coding and intronic regions, respectively (<http://flybase.org/reports/FBgn0045495.html>). Whereas these P-element insertions did not alter the basal firing rate (data not shown), they caused a significant reduction in light-induced responses of class IV dendritic arborization neurons (Fig. 4b), as in hemizygous larvae carrying one *MiET1* allele and one deletion encompassing *Gr28b* (*Df(2L)Exel7031*; <http://flybase.org/reports/FBab0037910.html>) (Fig. 4c). The *MiET1* P-element inserts into the coding sequence common to all reported transcripts, and its mobilization for excision restored the light-induced response in class IV dendritic arborization neurons (Fig. 4d). Moreover, knockdown of *Gr28b* expression with *UAS-RNAi* driven by *ppk-GAL4* caused an overall reduction of light response of class IV dendritic arborization neurons (Fig. 4e). Taken together, our data indicate that *Gr28b* is expressed in class IV dendritic arborization neurons, and is required for proper light responses. Whether *Gr28b* is the direct photosensing molecule awaits further experimentation.

Sequence analysis revealed that *Gr28b* has a rhodopsin-like structure plus one extra transmembrane segment (Supplementary Fig. 12), raising the question of whether the *Gr28b*-dependent light response involves G-protein signalling. To test whether G-protein signalling is required in class IV dendritic arborization neurons, we applied the myristoylated  $\beta\gamma$ -binding peptide mSIRK, and found that the light response in class IV dendritic arborization neurons was significantly reduced (Supplementary Fig. 13). Thus, G-protein signalling is probably involved in the light response of class IV dendritic arborization neurons, similar to findings in *C. elegans*<sup>10</sup>. We further tested cyclic nucleotide-gated (CNG) channels, which are known to act downstream of Lite-1 and G proteins in *C. elegans*<sup>10</sup>. Unlike in *C. elegans*, blocking CNG channels with *L-cis*-diltiazem in class IV dendritic arborization neurons had no effect on their light responses (Supplementary Fig. 14).

## TrpA1 is required in light transduction in class IV neurons

Transient receptor potential (TRP) channels were first identified and characterized in the *Drosophila* compound eye<sup>29,31</sup>, with TRP and TRP-like (*trpl*) having key roles in phototransduction<sup>29</sup>. However, our electrophysiological studies revealed no defects in the light response of class IV dendritic arborization neurons in *trpl* or *painless* mutant larvae (Supplementary Fig. 15).

*TrpA1*, a *Drosophila* homologue of mammalian *TrpA*, may function as a thermosensor in larvae and adults<sup>32–34</sup>, and a receptor for reactive electrophiles such as allyl isothiocyanate (AITC)<sup>35</sup>. A *TrpA1* mutant exhibited normal basal firing in class IV dendritic arborization neurons (data not shown), but no light-induced firing increase (Fig. 4f). As reported previously<sup>32</sup>, we detected strong *TrpA1* immunoreactivity in several neurons in the larval brain but not in peripheral neurons (data not shown). We then performed MARCM (mosaic analysis with a repressible cell marker)<sup>36</sup>, and found no light response in class IV dendritic arborization neurons lacking *TrpA1* (Fig. 4g), and a significant reduction of light-induced firing of the heterozygous class IV dendritic arborization neurons (Supplementary Fig. 16), indicating that *TrpA1* is present in levels below immunodetection, but nonetheless of functional importance. In support of this notion, AITC caused strong activation of class IV dendritic arborization neurons, and this activation was abolished in the *TrpA1* mutant (Supplementary Fig. 17a). Moreover, *TrpA1* RNAi expression specifically in class IV

dendritic arborization neurons eliminated the light-induced firing change (Supplementary Fig. 18). Taken together, our observations suggest that TrpA1 is required cell-autonomously for light transduction in class IV dendritic arborization neurons.

Given the lack of AITC activation of class I or class III dendritic arborization neurons (Supplementary Fig. 17b), we expressed TrpA1 in class I dendritic arborization neurons and found that it conferred AITC sensitivity but not light response (Supplementary Fig. 17c, d), indicating that TrpA1 is not sufficient for light sensing. Because *trans*-heterozygotes carrying one mutant allele of *TrpA1* and one copy of the *MiET1* P-element insertion in the *Gr28b* gene showed reduced light response (Supplementary Fig. 19), it is likely that Gr28b and TrpA1 function in the same phototransduction pathway.

## Class IV neurons mediate light avoidance behaviour

To test whether class IV dendritic arborization neurons are involved in light avoidance, we genetically ablated class IV dendritic arborization neurons of third instar larvae by expressing the pro-apoptotic genes *Hid* (ref. <sup>15</sup>) and *reaper* (*rpr*) (ref. <sup>37</sup>) via *ppk-GAL4* (*ppk-GAL4; UAS-Hid, rpr*) (Supplementary Fig. 1). We also constructed a line lacking Bolwig organs as well as class IV dendritic arborization neurons (*UAS-Hid, rpr; GMR-Hid; ppk-GAL4*). Notably, both lines showed markedly decreased white-light-avoidance behaviour compared to wild-type and *GMR-Hid* (Bolwig-organ-ablated) larvae (Fig. 5a, b). Class IV dendritic-arborization-neurons-ablated animals showed a significant decrease of avoidance versus wild type, for all white light intensities tested (Fig. 5c–g). Ablation of class IV dendritic arborization neurons in animals lacking Bolwig organs produced a further decrease in white light avoidance (Fig. 5d–g). Avoidance of high-intensity ( $>0.57$  mW mm<sup>-2</sup>) white light was normal when Bolwig organs were ablated in wild type (Fig. 5e–g), and in control strains with either *GAL4* or *UAS* (Fig. 5f). Taken together with similar findings with ultraviolet, violet and blue light (Supplementary Figs 20–22), these results demonstrate that class IV dendritic arborization neurons are necessary to elicit photoavoidance at high intensities. It thus seems that the Bolwig organs and class IV dendritic arborization neurons operate in different light intensity regimes: Bolwig organs are tuned to low light, whereas class IV dendritic arborization neurons, required in low light, are the primary sensors at high intensities.

Careful examination of *ppk-GAL4* revealed additional expression in four mouth hook neurons, but not in the central nervous system (Supplementary Fig. 23a–d). Laser ablation of these four neurons in the *GMR-Hid* background had no effect on light avoidance behaviour (Supplementary Fig. 23e). Therefore, the class IV dendritic arborization neurons in the body wall are the ones important for the light avoidance behaviour.

*Pickpocket*, a Degenerin/Epithelial sodium Channel (DEG/ENaC) family member specifically expressed in class IV dendritic arborization neurons<sup>24,38</sup> (Fig. 3c), has been implicated in locomotion control<sup>39–41</sup>. However, nose-touch experiments<sup>42</sup> revealed that larvae lacking class IV dendritic arborization neurons responded normally to gentle touch by retracting or turning away their heads (Supplementary Fig. 24). Moreover, direct recording of class IV dendritic arborization neurons in *ppk* mutant larvae revealed no defect in light response (Supplementary Fig. 15b). These results demonstrate that reduced light avoidance in class IV dendritic-arborization-ablated larvae is not due to non-specific effects.

To probe sufficiency, we expressed channelrhodopsin-2 (ChR2), a retinal-dependent cation channel gated by light from ultraviolet to green<sup>43–45</sup>, specifically in class IV dendritic arborization neurons. ChR2 conferred green light sensitivity to dendritic arborization neurons from larvae fed with retinal (Supplementary Fig. 25), as well as robust avoidance of

green light of retinal-fed larvae without Bolwig organs (Fig. 5h). Thus, activation of class IV dendritic arborization neurons is sufficient to induce avoidance.

With or without Bolwig organs, *TrpA1* mutant larvae showed deficient avoidance of 1 mW mm<sup>-2</sup> white light (Fig. 5i). Moreover, reducing *TrpA1* expression in class IV dendritic arborization neurons by RNAi was sufficient to abolish the light avoidance behaviour in animals without Bolwig organs (Supplementary Fig. 26). Together, our physiological and behavioural studies indicate that a light transduction pathway involving *TrpA1* and *Gr28b* in class IV dendritic arborization neurons is necessary for light avoidance.

## Discussion

Extra-ocular photoreceptors, previously found in reptiles, birds, amphibians and fish, provide a good measure of ambient light luminance and serve mainly non-image-forming functions such as phototaxis, circadian photo-entrainment, pupal reflex, shadow reaction and magnetic orientation<sup>1-7</sup>. Usually, these extra-ocular photoreceptors have much lower light sensitivity and slower kinetics than ocular photoreceptors<sup>3</sup>.

*Drosophila* larvae have primitive eye structures, the Bolwig organs, which control avoidance of dim light<sup>12</sup>. Here we report that the class IV dendritic arborization neurons, previously implicated in mechanosensory response and motion control<sup>38-41,46</sup>, are surprisingly also photoreceptors. Our behavioural analysis suggests that Bolwig organs and class IV dendritic arborization neurons have different regimes of light sensing in acute photoavoidance. Bolwig organs, packed with photopigments<sup>47</sup>, are preferentially required for avoidance of low light. Class IV dendritic arborization neurons, which also contribute to low light avoidance, are the primary sensors at sunlight-level intensities. This organization ensures that larvae can detect the full range of ambient light intensities, from dim to strong.

Class IV dendritic arborization neurons have the intrinsic ability to sense light, even after isolation in culture, and their dendrites are capable of sensing light (Fig. 3 and Supplementary Fig. 10). Importantly, the dendrites of class IV dendritic arborization neurons have complete and non-redundant coverage of the body wall (Fig. 3c), allowing animals to perceive illumination of any body part, and initiate an appropriate behavioural response. Larvae spend much of the time with their heads digging into food, making their Bolwig organs on the head less likely to be exposed to light. Thus, the ability to sense light with sensory neurons tiling the body wall is critical for detection of exposure.

Class IV dendritic arborization neurons use a novel light transduction pathway. Like in *C. elegans*, a putative chemosensory G-protein-coupled receptor, *Gr28b*, is involved for phototransduction in class IV dendritic arborization neurons (Fig. 4b-e). *TrpA1* also is essential (Fig. 4f, g and Supplementary Fig. 18). *Drosophila* larval class IV dendritic arborization neurons may function as nociceptors<sup>46,48,49</sup>. They are required for thermal and mechanical nociception, and activation of class IV dendritic arborization neurons is sufficient to induce a behaviour pattern similar to nocifension<sup>46,48,49</sup>. Given that class IV dendritic arborization neurons are required for larvae to avoid harmful light stimuli, these neurons seem to be poised to alert the animal to a variety of adversities.

Our study has uncovered unexpected light-sensing machinery, which could be critical for foraging larvae to avoid harmful sunlight, desiccation and predation. By providing precedence for photoreceptors strategically placed away from the eyes, our finding of an array of class IV dendritic arborization neurons with elaborate dendrites tiling the entire body wall, and acting as light-sensing antennae, raises the question of whether other animals with eyes might also possess extra-ocular photoreceptors for more thorough light detection and behavioural response.

## METHODS SUMMARY

### Light avoidance assay

Light avoidance was scored if the third instar larva reversed in direction or turned its head completely away from the 1.7-mm light spot on its head during the 5-s illumination. Two-tailed Fisher exact test (20–40 larvae per condition), \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .

### Electrophysiology

Action potentials were monitored via extracellular recordings from a third instar larval fillet with muscles removed, using an Axon 700B amplifier and pCLAMP 10 software.

### GCaMP3 imaging

Third instar larval fillets were imaged on a Zeiss LS510 META confocal microscope with an Olympus  $\times 40/0.8$  NA water immersion objective, with a 488-nm laser. GCaMP3 cDNA is available from AddGene.

### Cell culture

Embryos homozygous for Canton S (*Cs*); *UAS-GCaMP3*; *ppk-GAL4*, *UAS-RFP* (for class IV dendritic arborization neuron culture) or *Cs*; *UAS-GCaMP3*; *19-12-GAL4*, *UAS-RFP* (for class III dendritic arborization neuron culture) were used for culture<sup>22,23</sup>.

### MARCM analysis

We recorded from class IV dendritic arborization neuron clones marked with GFP for lacking *TrpA1* (ref. <sup>36</sup>).

Full Methods and any associated references are available in the online version of the paper at [www.nature.com/nature](http://www.nature.com/nature).

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank P. Garrity, C. Desplan, J. Blau, C. Montell, J. Hall, H. Amrein, L. Tian, S. Younger and S. Zhu for fly stocks and reagents; T. Jin for technical support; C. Han for generating whole larval images; H. H. Lee for collaboration to identify the *19-12-GAL4* and *21-7-GAL4* lines; R. Yang, H. H. Lee, B. Ye, J. Parrish, P. Soba, B. Schroeder and J. Bagley for discussions and advice; B. Ye, R. Yang, W. Ge and J. Berg for critical reading of the manuscript; and Jan laboratory members for discussions. Y.X. was a recipient of a Long-term Fellowship from the Human Frontier Science Program, N.V. is supported by Deutsche Forschungsgemeinschaft. This work is supported by a NIH grant (2R37NS040929) to Y.N.J. Y.X. and Q.Y. are associates, L.Y.J. and Y.N.J. are investigators of the Howard Hughes Medical Institute. L.L.L. is supported by the Howard Hughes Medical Institute, Janelia Farm Campus.


## References

1. Fu Y, Liao HW, Do MT, Yau KW. Non-image-forming ocular photoreception in vertebrates. *Curr Opin Neurobiol* 2005;15:415–422. [PubMed: 16023851]
2. Yau KW, Hardie RC. Phototransduction motifs and variations. *Cell* 2009;139:246–264. [PubMed: 19837030]
3. Steven DM. The dermal light sense. *Biol Rev Camb Philos Soc* 1963;38:204–240. [PubMed: 4385905]
4. Millott N. The dermal light sense. *Symp Zool Soc Lond* 1968;23:1–36.

5. Yoshida, M. Handbook of Sensory Physiology. Vol. 7/6A. Springer; 1979. Extraocular photoreception; p. 581-640.
6. Halford S, et al. VA opsin-based photoreceptors in the hypothalamus of birds. *Curr Biol* 2009;19:1396–1402. [PubMed: 19664923]
7. Phillips JB, Deutschlander ME, Freake MJ, Borland SC. The role of extraocular photoreceptors in newt magnetic compass orientation: parallels between light-dependent magnetoreception and polarized light detection in vertebrates. *J Exp Biol* 2001;204:2543–2552. [PubMed: 11511670]
8. Ward A, Liu J, Feng Z, Xu XZ. Light-sensitive neurons and channels mediate phototaxis in *C. elegans*. *Nature Neurosci* 2008;11:916–922. [PubMed: 18604203]
9. Edwards SL, et al. A novel molecular solution for ultraviolet light detection in *Caenorhabditis elegans*. *PLoS Biol* 2008;6:e198. [PubMed: 18687026]
10. Liu J, et al. *C. elegans* phototransduction requires a G protein-dependent cGMP pathway and a taste receptor homolog. *Nature Neurosci* 2010;13:715–722. [PubMed: 20436480]
11. Sawin-McCormack EP, Sokolowski MB, Campos AR. Characterization and genetic analysis of *Drosophila melanogaster* photobehavior during larval development. *J Neurogenet* 1995;10:119–135. [PubMed: 8592272]
12. Mazzoni EO, Desplan C, Blau J. Circadian pacemaker neurons transmit and modulate visual information to control a rapid behavioral response. *Neuron* 2005;45:293–300. [PubMed: 15664180]
13. Sprecher SG, Desplan C. Switch of rhodopsin expression in terminally differentiated *Drosophila* sensory neurons. *Nature* 2008;454:533–537. [PubMed: 18594514]
14. Willson RC, Gulkis S, Janssen M, Hudson HS, Chapman GA. Observations of solar irradiance variability. *Science* 1981;211:700–702. [PubMed: 17776650]
15. Grether ME, Abrams JM, Agapite J, White K, Steller H. The head involution defective gene of *Drosophila melanogaster* functions in programmed cell death. *Genes Dev* 1995;9:1694–1708. [PubMed: 7622034]
16. Hay BA, Maile R, Rubin GM. P element insertion-dependent geneactivation in the *Drosophila* eye. *Proc Natl Acad Sci USA* 1997;94:5195–5200. [PubMed: 9144214]
17. Nakai J, Ohkura M, Imoto K. A high signal-to-noise Ca<sup>2+</sup> probe composed of a single green fluorescent protein. *Nature Biotechnol* 2001;19:137–141. [PubMed: 11175727]
18. Wang JW, Wong AM, Flores J, Vosshall LB, Axel R. Two-photon calcium imaging reveals an odor-evoked map of activity in the fly brain. *Cell* 2003;112:271–282. [PubMed: 12553914]
19. Tian L, et al. Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators. *Nature Methods* 2009;6:875–881. [PubMed: 19898485]
20. Ueda A, Wu CF. Effects of *hyperkinetic*, a  $\beta$  subunit of *Shaker* voltage-dependent K<sup>+</sup> channels, on the oxidation state of presynaptic nerve terminals. *J Neurogenet* 2008;22:103–115.
21. Auld VJ, Fetter RD, Broadie K, Goodman CS. Gliotactin, a novel transmembrane protein on peripheral glia, is required to form the blood-nerve barrier in *Drosophila*. *Cell* 1995;81:757–767. [PubMed: 7539719]
22. Saito M, Wu CF. Expression of ion channels and mutational effects in giant *Drosophila* neurons differentiated from cell division-arrested embryonic neuroblasts. *J Neurosci* 1991;11:2135–2150. [PubMed: 1712379]
23. Bai J, Sepp KJ, Perrimon N. Culture of *Drosophila* primary cells dissociated from gastrula embryos and their use in RNAi screening. *Nature Protocols* 2009;4:1502–1512.
24. Grueber WB, et al. Projections of *Drosophila* multidendritic neurons in the central nervous system: links with peripheral dendrite morphology. *Development* 2007;134:55–64. [PubMed: 17164414]
25. Grueber WB, Jan LY, Jan YN. Tiling of the *Drosophila* epidermis by multidendritic sensory neurons. *Development* 2002;129:2867–2878. [PubMed: 12050135]
26. Grueber WB, Ye B, Moore AW, Jan LY, Jan YN. Dendrites of distinct classes of *Drosophila* sensory neurons show different capacities for homotypic repulsion. *Curr Biol* 2003;13:618–626. [PubMed: 12699617]
27. O'Tousa JE, et al. The *Drosophila ninaE* gene encodes an opsin. *Cell* 1985;40:839–850. [PubMed: 2985266]




28. Dolezelova E, Dolezel D, Hall JC. Rhythm defects caused by newly engineered null mutations in *Drosophila's* *cryptochrome* gene. *Genetics* 2007;177:329–345. [PubMed: 17720919]
29. Montell C. Visual transduction in *Drosophila*. *Annu Rev Cell Dev Biol* 1999;15:231–268. [PubMed: 10611962]
30. Thorne N, Amrein H. Atypical expression of *Drosophila* gustatory receptor genes in sensory and central neurons. *J Comp Neurol* 2008;506:548–568. [PubMed: 18067151]
31. Montell C, Jones K, Hafen E, Rubin G. Rescue of the *Drosophila* phototransduction mutation *trp* by germline transformation. *Science* 1985;230:1040–1043. [PubMed: 3933112]
32. Rosenzweig M, et al. The *Drosophila* ortholog of vertebrate TRPA1 regulates thermotaxis. *Genes Dev* 2005;19:419–424. [PubMed: 15681611]
33. Hamada FN, et al. An internal thermal sensor controlling temperature preference in *Drosophila*. *Nature* 2008;454:217–220. [PubMed: 18548007]
34. Kwon Y, Shim HS, Wang X, Montell C. Control of thermotactic behavior via coupling of a TRP channel to a phospholipase C signaling cascade. *Nature Neurosci* 2008;11:871–873. [PubMed: 18660806]
35. Kang K, et al. Analysis of *Drosophila* TRPA1 reveals an ancient origin for human chemical nociception. *Nature* 2010;464:597–600. [PubMed: 20237474]
36. Lee T, Luo L. Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. *Neuron* 1999;22:451–461. [PubMed: 10197526]
37. White K, et al. Genetic control of programmed cell death in *Drosophila*. *Science* 1994;264:677–683. [PubMed: 8171319]
38. Adams CM, et al. Ripped pocket and pickpocket, novel *Drosophila* DEG/ENaC subunits expressed in early development and in mechanosensory neurons. *J Cell Biol* 1998;140:143–152. [PubMed: 9425162]
39. Ainsley JA, Kim MJ, Wegman LJ, Pettus JM, Johnson WA. Sensory mechanisms controlling the timing of larval developmental and behavioral transitions require the *Drosophila* DEG/ENaC subunit, Pickpocket1. *Dev Biol* 2008;322:46–55. [PubMed: 18674528]
40. Ainsley JA, et al. Enhanced locomotion caused by loss of the *Drosophila* DEG/ENaC protein Pickpocket1. *Curr Biol* 2003;13:1557–1563. [PubMed: 12956960]
41. Xu K, et al. The *fragile X-related* gene affects the crawling behavior of *Drosophila* larvae by regulating the mRNA level of the DEG/ENaC protein Pickpocket1. *Curr Biol* 2004;14:1025–1034. [PubMed: 15202995]
42. Kernan M, Cowan D, Zuker C. Genetic dissection of mechanosensory transduction: mechanoreception-defective mutations of *Drosophila*. *Neuron* 1994;12:1195–1206. [PubMed: 8011334]
43. Nagel G, et al. Channelrhodopsin-2, a directly light-gated cation-selective membrane channel. *Proc Natl Acad Sci USA* 2003;100:13940–13945. [PubMed: 14615590]
44. Schroll C, et al. Light-induced activation of distinct modulatory neurons triggers appetitive or aversive learning in *Drosophila* larvae. *Curr Biol* 2006;16:1741–1747. [PubMed: 16950113]
45. Zhang F, Aravanis AM, Adamantidis A, de Lecea L, Deisseroth K. Circuit-breakers: optical technologies for probing neural signals and systems. *Nature Rev Neurosci* 2007;8:577–581. [PubMed: 17643087]
46. Zhong L, Hwang RY, Tracey WD. Pickpocket is a DEG/ENaC protein required for mechanical nociception in *Drosophila* larvae. *Curr Biol* 2010;20:429–434. [PubMed: 20171104]
47. Pollock JA, Benzer S. Transcript localization of four opsin genes in the three visual organs of *Drosophila*; RH2 is ocellus specific. *Nature* 1988;333:779–782. [PubMed: 2968518]
48. Tracey WD Jr, Wilson RI, Laurent G, Benzer S. *painless*, a *Drosophila* gene essential for nociception. *Cell* 2003;113:261–273. [PubMed: 12705873]
49. Hwang RY, et al. Nociceptive neurons protect *Drosophila* larvae from parasitoid wasps. *Curr Biol* 2007;17:2105–2116. [PubMed: 18060782]




**Figure 1. Photoreceptors in addition to Bolwig organs contribute to photoavoidance**

**a, b**, Examples of light avoidance of wild-type (**a**) and *GMR-Hid* (**b**) larvae exposed to white light (0.57 mW mm<sup>-2</sup>) applied from 0 to 5 s. The light spot is indicated by the dotted circle. The arrow indicates the direction of larval locomotion; arrowheads at 2 s (**a**) and 5 s (**b**) indicate larval head turning. **c–h**, Percentage of animals avoiding white light (**c**), light of 360 nm (ultraviolet; **d**), 402 nm (violet; **e**), 470 nm (blue; **f**), 525 nm (green; **g**) and 620 nm (red; **h**) at different intensities. \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001, two-tailed Fisher exact test. Twenty to forty larvae were tested for each condition. Scale bar: 1 mm (**a, b**), shown at –2 s.




### Figure 2. Light activates class IV dendritic arborization neurons

**a**, Pre-stimulation image showing larval dorsal cluster sensory neurons (dbd, bipolar dendrite neuron; ddaD and ddaE, class I dendritic arborization neurons; ddaB, class II dendritic arborization neurons; ddaA and ddaF, class III dendritic arborization neurons; ddaC, class IV dendritic arborization neurons; ES, external sensory organ). Up is dorsal; right is anterior. For an atlas of the larval peripheral nervous system, see ref. <sup>25</sup>. **b**, Responses of the dorsal cluster neurons in **a** to 5 s blue light (470 nm) illumination. The boxed area in the left panel and insets in all three panels show the somas of ddaC, ddaF and ddaD dendritic arborization neurons. Left, pre-stimulation; middle, post-stimulation; right, GCaMP3 intensity difference (middle panel minus left panel), with ddaC dendrites (arrow) and axon (arrowhead) marked. **c**, **d**, Similar experiments with 5 s green (546 nm; **c**) and ultraviolet light (365 nm; **d**) revealed ddaC activation by ultraviolet, but not green, light. Scale bar in **a–d**, 20  $\mu\text{m}$ ; colour scale in right panels of **b–d** shows dynamic range (0–4,095). **e**, Time course of somatic GCaMP3 signals of dorsal cluster neurons shown in **a–d**. Time frames are indicated. **f**, Summary of somatic fluorescence changes ( $\Delta F/F$ ) of dorsal cluster neurons in response to 5 s light stimulation,  $n = 7–16$ . **g**, Example firing traces of ddaC in response to 5 s 470 nm blue light. **h**, Summary of firing frequency changes (average frequency of 5 s before light exposure subtracted from average frequency during 5 s of light exposure) of ddaC induced by white, 340, 380, 402, 470, 525 and 620 nm light. For clarity, significance is only shown for the 340 nm curve. Light intensity is reported as the log of ( $I$  normalized to  $I_0 = 1 \text{ mW mm}^{-2}$ ). Green (525 nm) or red (620 nm) light has no effect ( $P > 0.05$ ).  $n = 5–9$ . **i**, Effect of  $1.4 \text{ mW mm}^{-2}$  white light on ddaC, average frequencies of 5 s before (control) and during the 5 s of light exposure (light) are plotted.  $n = 6$ . \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ; two-tailed paired  $t$ -test. All error bars indicate s.e.m.




**Figure 3. Cell-autonomous activation of class IV dendritic arborization neurons by light**  
**a, b**, Quantification of somatic fluorescence changes ( $\Delta F/F$ ) in response to 5 s light and 100  $\mu$ M allyl isothiocyanate (AITC) stimulation of cultured class IV (**a**) and III (**b**) dendritic arborization neurons; RFP signals serve as control.  $n = 10-13$  (light) and  $n = 4$  (AITC) in **a**,  $n = 9$  in **b**. **c**, Larva with class IV dendritic arborization neurons labelled with GFP by *ppk-GAL4*. Dendrites tile the body wall. Boxed area shows an abdominal hemi-segment; three dotted circles mark soma positions of D (dorsal, *ddaC*), L (lateral, *V'ada*) and V (ventral, *VdaB*) class IV dendritic arborization neurons, respectively. Up, dorsal; left, anterior. Scale bar, 200  $\mu$ m. **d**, Illumination of dendrites within the dotted circle of GFP-labelled *ddaC* dendrites. Up, dorsal. Scale bar, 50  $\mu$ m. **e**, Responses of *ddaC* with dendritic illumination.  $n = 5$ . \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ; two-tailed paired *t*-test. All error bars indicate s.e.m.



**Figure 4. Gr28b and TrpA1 are essential for class IV dendritic arborization neuron light responses**

**a.** No significant defects were detected between wild-type and mutants of known phototransduction molecules with 340, 380, 402, 470, or 620 nm light.  $n=5-10$ . **b.** Reduced light response of class IV dendritic arborization neurons in *MiET1* and *PBac* larvae.  $n=8-29$ . **c.** Reduced light response of class IV dendritic arborization neurons in *MiET1*/deficiency larvae.  $n=5-12$ . **d.** Precise excision of *MiET1* P-element insertion restores light response in class IV dendritic arborization neurons.  $n=6-9$ . **e.** Reduced light responses of class IV dendritic arborization neurons with *Gr28b* RNAi knockdown.  $n=5-8$ . **f.** Abolished light responses of class IV dendritic arborization neurons in *TrpA1*<sup>-/-</sup> mutants.  $n=8-13$ . **g.** MARCM analysis of *TrpA1*<sup>+/-</sup> and *TrpA1*<sup>-/-</sup> class IV dendritic arborization neurons' response to light.  $n=5-8$ . For **a-g**, Light intensities ( $\text{mW mm}^{-2}$ ) are: 1.15 (340 nm), 5.79 (380 nm), 11.4 (402 nm), 52.8 (470 nm), 43.4 (525 nm), 29.6 (620 nm) and 94.7 (white). For **a, b, c, e**, \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ; one-way ANOVA followed by a Bonferroni post test; for **d, f, g**, \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ; two-tailed unpaired *t*-test. All error bars indicate s.e.m.



**Figure 5. Class IV dendritic arborization neurons are the extra-ocular photoreceptors that contribute to light avoidance**

**a, b**, Examples of larvae with either class IV dendritic arborization neurons ablated (**a**) or both Bolwig organs and class IV dendritic arborization neurons ablated (**b**) that failed to respond to white light ( $0.57 \text{ mW mm}^{-2}$ ) applied from 0 to 5 s (dotted circle). Arrow indicates locomotion direction. Scale bar, 1 mm (**a, b**), shown at  $-2 \text{ s}$ . **c–g**, Percentage of animals avoiding white light of different intensities (in  $\text{mW mm}^{-2}$ : **c**, 0.088; **d**, 0.24; **e**, 0.57; **f**, 1.0; **g**, 1.67). Wild-type larvae, Bolwig-organ-ablated larvae (*GMR-Hid*), larvae with class IV dendritic arborization neurons ablated (*ppk-GAL4; UAS-Hid, rpr*) and larvae with both ablated (*UAS-Hid, rpr; GMR-Hid; ppk-GAL4*) were examined. **h**, Percentage of Bolwig-organ-ablated animals avoiding  $0.25 \text{ mW mm}^{-2}$  525 nm green light when class IV dendritic arborization neurons express ChR2 with or without dietary retinal. **i**, Percentage of animals avoiding white light at  $1 \text{ mW mm}^{-2}$ . For **c–i**, controls are black bars. Twenty to forty animals were tested for each condition; \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ; two-tailed Fisher exact test. ChR2, channelrhodopsin-2; *rpr*, *reaper*; NS, not significant.