

## Limitations of the use of the mtp40 fragment as a marker of differentiation between *Mycobacterium tuberculosis* and *M. bovis*

CRISTINA VIANA-NIERO, SYLVIA CARDOSO LEÃO

The bacilli that cause tuberculosis (TB) belong to the *Mycobacterium tuberculosis* complex, which is composed of *M. tuberculosis*, *M. bovis* subsp. *bovis*, *M. africanum* and *M. microti*, as well as the *M. bovis* BCG strain used for vaccination. It has been proposed that newly discovered species should also be included in this complex. These new species include *M. canettii*, a variant of *M. tuberculosis* found in the Somalia region<sup>(1)</sup>, *M. bovis* subsp. *caprae*, the etiologic agent of TB in goats<sup>(2)</sup> and *M. pinnipedii*, which causes TB in sea lions and may also infect humans<sup>(3)</sup>.

Studies involving DNA-DNA hybridization and sequence analysis of the 16S rDNA sequence, the 16S-23S intergenic spacer sequence and the gene encoding the hsp65 heat shock protein<sup>(4)</sup> have shown that this complex, in fact, constitutes a single species. For reasons essentially related to the medical and veterinary significance of this group of bacteria, as well as to its pathogenic power and the wide spectrum of hosts receptive to each species, the nomenclature remains unchanged.

Within this complex of bacteria, *M. tuberculosis* is the principal pathogen in humans. However, cases of human TB resulting from infection with *M. africanum* and *M. canettii* have been reported, mainly in Africa<sup>(1)</sup>. In addition, *M. bovis*, the etiologic agent of bovine TB, may also infect humans and other animals. Studies conducted in Argentina and England showed that *M. bovis* is responsible for 0.4% to 1% of human TB cases<sup>(5,6)</sup>. According to the Pan American Health Organization<sup>(7)</sup>, 7000 new cases of TB appear annually in South America. There are no data available regarding human cases of TB resulting from *M. bovis* infection in Brazil. This species is naturally resistant to pyrazinamide (PZA), a drug used in the treatment of TB in humans, which, in certain cases, makes the differentiation between species relevant. Such cases include those in which

epidemiological evidence suggests *M. bovis* involvement and those in which the patient fails to respond to treatment regimes that include PZA.

A diagnosis of TB caused by *M. tuberculosis* or by *M. bovis* can be made through analysis of clinical data and radiological evidence. However, bacteriological diagnosis is necessary in order to confirm the diagnosis and identify the species involved. *M. tuberculosis* and *M. bovis* can be differentiated using phenotyping techniques such as tests for niacin production and nitrate reductase, as well as cultures to assess bacterial growth in the presence of thiophene-2-carboxylic acid hydrazide and PZA<sup>(4)</sup>. The polymerase chain reaction (PCR) method has been incorporated into the routine of many laboratories as a diagnostic alternative due to its greater speed, sensitivity and specificity. This technique allows the distinction between *M. tuberculosis* and *M. bovis* to be made through differential amplification of the *pncA* and *oxyR* gene sequences<sup>(8)</sup>, amplification and analysis of enzymatic restriction within the *gyrB* sequence<sup>(9)</sup>, multiple amplifications (multiplex PCR) of the DR fragment regions, the insertion sequence 6110 (IS6110) and the hsp65 gene<sup>(10)</sup>, as well as amplification of the mtp40 fragment, which is exclusive to *M. tuberculosis* and therefore absent from *M. bovis*<sup>(11)</sup>.

The mtp40 fragment, contained within the *plcA* gene sequence, which encodes the phospholipase C enzyme of *M. tuberculosis*, has been widely used for specific diagnosis of *M. tuberculosis* in (uncultured) clinical samples<sup>(11)</sup> and in cultures of isolated strains<sup>(12)</sup>. The absence of mtp40 from *M. bovis* has been verified by various authors, although the claim that mtp40 is present in all clinically isolated *M. tuberculosis* strains has been challenged<sup>(13)</sup>.

In our laboratory, we analyzed the results from the amplification of the mtp40 fragment in 790 *M. tuberculosis* strains isolated at the *Rijksinstituut voor Volksgezondheid en Milieu* (RIVM, National Institute

for Public Health and the Environment) reference laboratory (Bilthoven, the Netherlands). The primers PT1 (CAACGCGCCGTCGGTGG) and PT2 (CCCCCACGGCACCGC) were employed. We also evaluated the results of such amplification in 105 *M. africanum* strains and 10 *M. canettii* strains isolated at the *Centre National de Référence des Méningocoques, Institut Pasteur* (National Reference Center for Meningococci, Pasteur Institute, Paris France). Positive amplification results were obtained in 94.6%, 54.6% and 70%, respectively, of the isolated strains of each species (data from the RIVM)<sup>(14)</sup>.

We studied, in detail, a set of 32 strains belonging to these two collections. These strains were chosen because they presented varying PCR results in relation to the mtp40 fragment. In 17 strains, a fragment of the expected size (396 bp) was amplified, no amplification occurred in 13, and a fragment larger than expected (1700 bp) was amplified in 2 of the strains (both *M. tuberculosis*). The presence of genetic polymorphisms in the phospholipase C gene, which would explain the absence of the mtp40 fragment from the isolates, was evaluated. The results show that, in most cases, the lack of mtp40 fragment amplification results from complete deletion of the *plcA* gene, as well as of the adjacent genes. In addition, insertion of a copy of the *IS6110* element into the mtp40 fragment was observed in 2 of the isolates, thereby impeding amplification of the fragment of the correct size (Figure 1)<sup>(15)</sup>.

There are no available data regarding the relevance of using this marker in Brazilian strains. However, the existence of *M. tuberculosis*, *M. africanum* and *M. canettii* isolates presenting genetic polymorphisms serves as an indicator that we should be choosing molecular markers that are capable of definitively identifying *M. tuberculosis* and differentiating it from other members of the complex. Currently, it is advisable to use a combination of phenotypic and genotypic markers in the differential diagnosis between *M. tuberculosis* and *M. bovis*.

Cristina Viana-Niero, Sylvia Cardoso Leão.  
Department of Microbiology, Immunology and  
Parasitology, UNIFESP-EPM  
e-mail: cviana@ecb.epm.br

FINANCIAL SUPPORT: FAPESP (grant no. 00/02525-3);  
CABBIO-CNPq (grant no. 480382/01-8);

Rede Brasileira de Pesquisa em TB (Rede-TB, Brazilian  
Tuberculosis Research Network)/grant no. 62.0055/01-4-  
PACDT-Milenio




Figure 1. Location of the mtp40 fragment in the *Mycobacterium tuberculosis* genome: (A) *M. tuberculosis* H37Rv; (B) insertion of a copy of the *IS6110* element into the mtp40 fragment; (C) deletion of the 8.6-kilobase fragment including the mtp40 fragment. The black square within the *plcA* gene represents the mtp40 fragment and the hashed rectangle represents the inserted sequence *6110*. This region does not exist within the *M. bovis* genome.

## REFERENCES

1. van Soolingen D, Hoogenboezem T, de Haas PEW, Hermans PWM, Koedam MA, Teppema KS, et al. A novel pathogenic taxon of the *Mycobacterium tuberculosis* complex, *Canettii*: Characterization of an exceptional isolate from Africa. *Int J Syst Bacteriol* 1997; 47(4):1236-45.
2. Aranaz A, Cousins D, Mateos A, Domínguez L. Elevation of *Mycobacterium tuberculosis* subsp. *caprae* Aranaz et al. 1999 to species rank as *Mycobacterium caprae* comb. nov., sp. nov. *Int J Syst Evol Microbiol* 2003; 53:1785-9.
3. Cousins D, Bastida R, Cataldi A, Quse V, Redrobe S, Dow S, et al. A Tuberculosis in seals caused by a novel member of the *Mycobacterium tuberculosis* complex: *Mycobacterium pinnipedii* sp. nov. *Int J Syst Evol Microbiol* 2003; 53:1305-14.
4. Euzéby JP. List of Bacterial names with Standing in Nomenclature – Société de Bactériologie Systématique et Vétérinaire – France. Disponível em (URL: <http://www.bacterio.cict.fr>). Acesso em 20 out 2003.
5. Barrera L, De Kantor IN. Nontuberculous mycobacteria and *Mycobacterium bovis* as a cause of human disease in Argentina. *Trop Geogr Méd* 1987; 39:222-7.
6. Yates MD, Grange JM. Incidence and nature of human tuberculosis due to bovine tubercle bacilli in South-East England: 1977-1987. *Epidemiol Infect* 1988; 101:225-9.
7. Pan American Health Organization. 1991. Health conditions in the Americas, vol I. Scientific publication n°524. Pan American Health Organization, Washington, DC.
8. Espinosa de los Monteros LE, Galan JC, Gutierrez M,

- Samper S, Garcia Marin JF, Martin C, Dominguez L, et al. Allele-specific PCR method based on *pncA* and *oxyR* sequences for distinguishing *Mycobacterium bovis* from *Mycobacterium tuberculosis*: Intraspecific *M. bovis pncA* sequence polymorphism. *J Clin Microbiol* 1998; 36:239-42.
9. Chimara E, Ferrazoli L, Leão SC. *Mycobacterium tuberculosis* complex differentiation using *gyrB*-restriction fragment length polymorphism (*gyrB*-RFLP) analysis. Submetido a Memórias do Instituto Oswaldo Cruz. 2004.
  10. Yeboah-Manu D, Yates MD, Wilson SM. Application of a Simple Multiplex PCR To Aid in Routine Work of the Mycobacterium Reference Laboratory. *J Clin Microbiol* 2001; 39(11): 4166-8.
  11. Del Portillo P, Murillo LA, Patarroyo ME. Amplification of a species-specific DNA fragment *Mycobacterium tuberculosis* and its possible use in diagnosis. *J Clin Microbiol* 1991; 29(10):2163-8.
  12. Liébana E, Aranaz A, Francis B, Cousins D. Assessment of genetic markers for species differentiation within the *Mycobacterium tuberculosis* complex. *J Clin Microbiol* 1996; 34(4):933-8.
  13. Vera-Cabrera L, Hoard ST, Laszlo A, Johnson WM. Analysis of genetic polymorphism in the phospholipase region of *Mycobacterium tuberculosis*. *J Clin Microbiol* 1997; 35(5):1190-5.
  14. Viana-Niero C, Vincent V. 1999. Étude moléculaire des bacilles de la tuberculose d'origine africaine: *Mycobacterium africanum* et souches "canetti". Tese de mestrado apresentada à Université Paris V et Université Paris XI.
  15. Viana-Niero C, de Haas PE, van Soolingen D, Leão S C. Analysis of genetic polymorphisms affecting the four phospholipase C (*plc*) genes in *Mycobacterium tuberculosis* complex clinical isolates. *Microbiol* 2004; 150:967-78.