

LINEAR ALGEBRAIC GROUPS AND COUNTABLE BOREL EQUIVALENCE RELATIONS

SCOT ADAMS AND ALEXANDER S. KECHRIS

This paper is a contribution to the study of Borel equivalence relations on standard Borel spaces (*i.e.*, Polish spaces equipped with their Borel structure). In mathematics one often deals with problems of classification of objects up to some notion of equivalence by invariants. Frequently these objects can be viewed as elements of a standard Borel space X and the equivalence turns out to be a Borel equivalence relation E on X . A *complete classification* of X up to E consists of finding a set of invariants I and a map $c : X \rightarrow I$ such that $xEy \Leftrightarrow c(x) = c(y)$. For this to be of any interest both I and c must be explicit or definable and as simple and concrete as possible. The theory of Borel equivalence relations studies the set-theoretic nature of possible invariants and develops a mathematical framework for measuring the complexity of such classification problems.

In organizing this study, the following concept of reducibility is fundamental. Let E, F be equivalence relations on standard Borel spaces X, Y , resp. We say that E is *Borel reducible* to F , in symbols,

$$E \leq_B F,$$

if there is a Borel map $f : X \rightarrow Y$ such that

$$xEy \Leftrightarrow f(x)Ff(y).$$

This simply means that any complete invariants for F work as well for E (after composing with f) and therefore, in some sense, E has a classification problem at most as complicated as that of F . We also let

$$E \sim_B F \Leftrightarrow E \leq_B F \ \& \ F \leq_B E,$$

and this means that E, F have classification problems of equal complexity. Finally, we use

$$E <_B F \Leftrightarrow E \leq_B F \ \& \ F \not\leq_B E$$

to signify that E has a (strictly) simpler classification problem than that of F .

Another way to look at this notion is to notice that $E \leq_B F$ simply means that there is an injection from the quotient space X/E into Y/F which is “Borel”, in the sense that it has a Borel lifting. Thus $E \leq_B F$ can be interpreted as saying that X/E has “Borel cardinality” less than or equal to that of Y/F and similarly

Received by the editors March 27, 1999 and, in revised form, April 21, 2000.

2000 *Mathematics Subject Classification*. Primary 03E15; Secondary 37A20.

Key words and phrases. Borel equivalence relations, Borel reducibility, cocycles, superrigidity.

The first author’s research was partially supported by NSF Grant DMS 9703480.

The second author’s research was partially supported by NSF Grant DMS 9619880 and a Visiting Miller Research Professorship at U.C. Berkeley.

$E \sim_B F$ means that $X/E, Y/F$ have the same “Borel cardinality”, while $E <_B F$ means that X/E has strictly smaller “Borel cardinality” than that of Y/F . Thus the study of the (pre-)order \leq_B on Borel equivalence relations can also be viewed as the study of the “Borel cardinalities” of their quotient spaces.

In the beginning the structure of \leq_B is quite simple. Denoting by X also the equality relation on a standard Borel space X , and letting n be any such space of finite cardinality n , we have that the following is an initial segment of \leq_B :

$$1 <_B 2 <_B 3 <_B \cdots <_B \mathbb{N},$$

and $\mathbb{N} <_B E$ for any Borel equivalence E not in this sequence. The first non-trivial theorem concerning the structure of Borel equivalence relations under \leq_B is the *Silver Dichotomy* (see Silver [32]) which asserts that any Borel (in fact even coanalytic) equivalence relation on a Polish space has either countably many or else perfectly many equivalence classes, in the sense that there is a perfect set any two distinct elements of which belong to different equivalence classes. This shows that a strong form of the Continuum Hypothesis is valid for quotient spaces of Borel equivalence relations. It implies that

$$1 <_B 2 <_B 3 <_B \cdots <_B \mathbb{N} <_B \mathbb{R}$$

is an initial segment of \leq_B and $\mathbb{R} <_B E$ for any Borel equivalence relation E not in this list.

For the next step, notice that the equivalence relations $E \leq_B \mathbb{R}$ are exactly those for which there is a Borel map $f : X \rightarrow Y$, where X is the space of E and Y some standard Borel space, such that $xEy \Leftrightarrow f(x) = f(y)$. Such equivalence relations are called *tame* (or *smooth* or *concretely classifiable* as they can be completely classified by invariants which are members of some standard Borel space). The canonical example of a non-tame equivalence relation is the *Vitali equivalence relation* E_0 on \mathbb{R} :

$$xE_0y \Leftrightarrow x - y \in \mathbb{Q}.$$

Thus $\mathbb{R} <_B E_0$. The second dichotomy theorem for Borel equivalence relations, called the *General Glimm-Effros Dichotomy* (see Harrington-Kechris-Louveau [14]), asserts that for any Borel equivalence relation E , either E is tame or else contains a (Borel) copy of E_0 , and so $E_0 \leq_B E$. (This generalizes earlier results of Glimm [12] and Effros [6].) Thus we have that

$$1 <_B 2 <_B 3 <_B \cdots <_B \mathbb{N} <_B \mathbb{R} <_B E_0$$

is an initial segment of \leq_B and $E_0 <_B E$ for any Borel equivalence relation E not in this list.

The linearity of the order \leq_B breaks down beyond E_0 . This has been known for some time but it has been amplified more recently by results of Woodin (unpublished), who showed that there are uncountably many pairwise \leq_B -incomparable Borel equivalence relations, and then by Louveau-Velickovic [25], who showed that the partial order of subsets of the integers under inclusion modulo finite sets embeds into \leq_B on Borel equivalence relations. This non-linearity is in fact a crucial point in which Borel cardinality theory differs from the classical Cantor cardinality theory.

We now turn our attention to a particular class of Borel equivalence relations, the so-called countable ones. A Borel equivalence relation E on X is called *countable* if every equivalence class $[x]_E$ is countable. A typical example is the equivalence

relation induced by a Borel action of a countable group on a standard Borel space, e.g., the one induced by the orbits of a single Borel automorphism. Conversely, a theorem of Feldman-Moore [8] asserts that every countable Borel equivalence relation is induced by a Borel action of a countable group. Such equivalence relations have long been studied in ergodic theory and the theory of operator algebras (see, e.g., the surveys [27] and [30]). It also turns out that many equivalence relations, although not necessarily countable, nevertheless are up to \sim_B countable, so they fall, for our purposes, in the domain of countable Borel equivalence relations. These include, for example, all the equivalence relations induced by Borel actions of second countable locally compact groups (e.g., Lie groups); see Feldman-Hahn-Moore [7] and Kechris [20]. Also the isomorphism relation on various classes of countable models, which in some sense have “finite type”, e.g., finitely generated groups, locally finite connected graphs, finite rank torsion-free abelian groups, turn out to be \sim_B to countable Borel equivalence relations (see Hjorth-Kechris [16]). There is thus a great variety of interesting examples of (up to \sim_B) countable Borel equivalence relations occurring in many areas of mathematics. See, for example, the papers [19], [22], [23], [24], [5], [18] and the references contained therein, for further information concerning the descriptive study of countable Borel equivalence relations. The facts we are about to state in the next paragraph come from the last paper.

To get some feeling about the general structure of \leq_B on countable Borel equivalence relations, we first note that there is a largest one, naturally dubbed *universal*, and denoted by E_∞ . Thus E_∞ is countable Borel and $E \leq_B E_\infty$ for every countable Borel E . Such an E_∞ is uniquely determined up to \sim_B . Thus, excluding the tame ones (which are those in the list $1, 2, 3, \dots, \mathbb{N}, \mathbb{R}$), all countable Borel equivalence relations fall in the interval

$$E_0 \leq_B E \leq_B E_\infty.$$

(This should be contrasted to the situation concerning general Borel equivalence relations, where, by a result of Friedman-Stanley [9], for every Borel equivalence relation E there is a Borel equivalence relation F with $E <_B F$.) It also turns out that there are intermediate ones:

$$E_0 <_B E <_B E_\infty.$$

The first examples were found by applying results of Adams [1], [2], and thus made heavy use of methods and results of ergodic theory. However, rather remarkably, and despite the great plethora of examples of countable Borel equivalence relations, until recently, only two examples of distinct up to \sim_B intermediate relations had been found, say E, F , and they satisfied $E_0 <_B E <_B F <_B E_\infty$. In particular, it has been a long-standing problem in this theory whether there are incomparable under \leq_B countable Borel equivalence relations (see, e.g., Kechris [19]).

The first main result of this paper provides an affirmative answer to this problem and in fact shows that the partial order \leq_B on countable Borel equivalence relations is quite complex.

Theorem 1. *The partial ordering of Borel sets under inclusion can be embedded in the partial (pre-)order of countable Borel equivalence relations under Borel reducibility. More precisely, there is a map $A \mapsto E_A$ assigning to each Borel subset of $2^{\mathbb{N}}$, A , a countable Borel equivalence relation, E_A , so that*

$$A_1 \subseteq A_2 \Leftrightarrow E_{A_1} \leq_B E_{A_2}.$$

In particular, this shows that every Borel partial ordering can be embedded into \leq_B on countable Borel equivalence relations.

Our second main result computes the exact descriptive complexity of the relations \leq_B, \sim_B on countable Borel equivalence relations. We first fix a parametrization of binary Borel relations on a standard Borel space, which for definiteness we take to be the Baire space $\mathbb{N}^{\mathbb{N}}$. This consists of a Π_1^1 (i.e., co-analytic) set $D \subseteq 2^{\mathbb{N}}$ and a map $d \in D \mapsto D_d$, assigning to each $d \in D$ a binary Borel relation on $\mathbb{N}^{\mathbb{N}}$ so that every such relation is of the form D_d for some d . Moreover, $(D, d \mapsto D_d)$ satisfies some natural definability properties spelled out in Section 5 below. Using this, we let

$$\mathcal{E}_{\leq} = \{(d, e) \in D^2 : D_d, D_e \text{ are countable Borel equivalence relations and } D_d \leq_B D_e\}$$

and we similarly define \mathcal{E}_{\sim} . A straightforward calculation shows that $\mathcal{E}_{\leq}, \mathcal{E}_{\sim}$ are Σ_2^1 (= projections of co-analytic sets). The following computes their precise complexities.

Theorem 2. *The sets $\mathcal{E}_{\leq}, \mathcal{E}_{\sim}$ are Σ_2^1 -complete. In other words, the notions of Borel reducibility and bi-reducibility of countable Borel equivalence relations are Σ_2^1 -complete.*

(Recall here that a set A in a standard Borel space X is Σ_2^1 -complete if it is Σ_2^1 and for any Σ_2^1 set $B \subseteq Y$, Y a standard Borel space, there is a Borel function $f : Y \rightarrow X$ with $B = f^{-1}(A)$.)

The proofs of these results depend heavily on the Superrigidity Theorems of Zimmer [35], 5.2.5 and 10.1.6, which deal with reduction of cocycles of measure-preserving actions of linear algebraic groups. In [35] (and other papers referred to in this book) Zimmer also shows, loosely speaking, that, under certain circumstances, the equivalence relation induced by a group action together with (the measure class of) an associated invariant measure “encodes” or “remembers” quite a lot of information concerning the group and the action from which it came. In some sense, the main point in the proofs of the results here is that, as an application of the Zimmer Superrigidity Theory, there is also a “rigidity” phenomenon occurring in the purely descriptive context: Under certain circumstances, quite a bit of information concerning a group G acting on a space X is “encoded” or “remembered” simply by the “Borel cardinality” of the quotient space X/G of the orbits. This “set theoretic rigidity” is undoubtedly an interesting phenomenon that needs to be further explored.

A similar set of ideas is used by Zimmer in studying orbit equivalence. (See Theorem 5.2.1, p. 95 of Zimmer [35].) However, in that application of superrigidity, the cocycles that arise are readily seen, via the Borel Density Theorem, to be Zariski dense. In the situation presented in this paper, we construct cocycles whose algebraic hulls must be analyzed. However, we compensate for this complication by being very careful to arrange things so that the domain groups of our cocycles are, in some sense, incompatible with the target groups.

The organization of this paper is as follows. After an introductory Section 1, we discuss in Section 2 the fundamental notion of cocycle of a group action, and collect together various known facts and techniques concerning this concept that we will need later on. These are scattered in [35] and other publications or they are folklore. It seemed though that it would be helpful to collect them together

in one place and present them in as detailed and self-contained way as possible, for the convenience of the reader who is not familiar with these kinds of methods. In Section 3, we use the Superrigidity Theorems of Zimmer [35] to derive some cocycle reduction results that we will need in the proofs of our main theorems. Although the presentation here is much more detailed than is customary in a paper written for experts in the ergodic theory of linear algebraic groups, it is still far from self-contained and depends heavily on results from this theory (as well as from the theory of linear algebraic groups that are needed here). At least every attempt has been made to state explicitly the results that are being quoted, and give precise references to standard books, whenever possible. In Section 4, the results in the preceding section are put together to give the proof of Theorem 1 and in Section 5 of Theorem 2.

Section 6 contains an application of these ideas to another problem that has received attention recently, namely the issue of the complexity of the isomorphism problem for torsion-free abelian groups of finite rank. See Hjorth-Kechris [16] and Hjorth [15] for some background on this problem. We denote by \cong_n the isomorphism relation on torsion-free abelian groups of rank $\leq n$. These are, up to isomorphism, simply the subgroups of $(\mathbb{Q}^n, +)$. We also denote by \cong_n^* the restriction of \cong_n to the *rigid groups*, *i.e.*, those whose only automorphisms are $x \mapsto \pm x$. There is a classical result of Baer which classifies torsion-free abelian groups of rank 1. However, it has been a long-standing problem in abelian group theory whether a satisfactory classification can be found for rank 2 or higher groups (see Fuchs [10]). As it is not hard to see that \cong_n is \sim_B to a countable Borel equivalence relation, one can place all this in the context of the theory of countable Borel equivalence relations and try to compute the complexity of the classification problem for \cong_n and \cong_n^* . Baer's result implies that $(\cong_1) \sim_B (\cong_1^*) \sim_B E_0$ and clearly $(\cong_n^*) \leq_B (\cong_{n+1}^*)$, so $E_0 \leq_B (\cong_n^*) \leq_B (\cong_{n+1}^*) \leq_B (\cong_{n+1}) \leq_B E_\infty$ for any $n \geq 1$. It has been conjectured in Hjorth-Kechris [16] that for $n \geq 2$, \cong_n is universal, *i.e.*, $(\cong_n) \sim_B E_\infty$. This is still open but if correct it would quantify the exact complexity of the classification problem of rank 2 or higher torsion-free abelian groups, and would show that in some sense no simple classification is possible. Recently Hjorth [15] has taken a first step in that direction by showing that $E_0 <_B (\cong_n)$ for $n \geq 2$, and Thomas [33] extended this by showing that $E_0 <_B (\cong_n^*)$ for $n \geq 2$. Thus the classification problem for rigid rank 2 torsion-free abelian groups is, in some precise sense, definitely more complex than that of the rank 1 case. We prove in Section 6 the following result which shows that for rigid groups the complexity increases with the rank.

Theorem 3. *For any $n \geq 1$, $(\cong_n^*) <_B (\cong_{n+1}^*)$.*

In particular, this shows that \cong_n^* is not universal for any n .

Finally, in Section 7 we discuss some additional facts that can be proved by the methods used in this paper and discuss some open problems.

1. PRELIMINARIES

We will use more or less standard notation and terminology concerning group actions and descriptive set theory (see Zimmer [35] and Kechris [21], resp.).

Let G be a locally compact second countable (lcsc) group. By a *standard Borel G -space* we mean a standard Borel space X together with a Borel action $(g, x) \mapsto g \cdot x$

of G on X . We denote by E_G^X the corresponding equivalence relation

$$xE_G^X y \Leftrightarrow \exists g(g \cdot x = y),$$

by $G \cdot x$ the *orbit* of x , and by $G_x = \{g : g \cdot x = x\}$ the *stabilizer* of x .

For a standard Borel space X (i.e., a Polish space equipped with its Borel structure) by a *measure* on X we will always mean a *probability Borel measure* on X . Such a measure is *non-atomic* if $\mu(\{x\}) = 0, \forall x \in X$. The measures on X also form a standard Borel space denoted by $P(X)$. If $f : X \rightarrow Y$ is Borel and $\mu \in P(X)$, then $f_*(\mu) \in P(Y)$ is defined by $f_*(\mu)(A) = \mu(f^{-1}(A))$. If X is a G -space, $\mu \in P(X)$ is *invariant* if $g \cdot \mu = \mu, \forall g \in G$, and *ergodic* if every G -invariant Borel set $A \subseteq X$ has μ -measure 0 or 1. This is equivalent to saying that every Borel function $f : X \rightarrow Y, Y$ a standard Borel space, which is G -invariant, is constant μ -a.e.

An equivalence relation E on a standard Borel space is called *tame* (or *smooth* in the terminology of Zimmer [35]) if there is a Borel function $S : X \rightarrow Y, Y$ a standard Borel space, such that $xEy \Leftrightarrow S(x) = S(y)$. It is of course enough to take $Y = [0, 1]$ here. If $E = E_G^X$, for a standard Borel G -space X , then this is equivalent to saying that E admits a Borel *transversal*, i.e., a Borel set meeting every orbit in exactly one point. (This result is even true for G a Polish group (Burgess); see Kechris [21], 18.20 (iii).) Recall that if G is a Polish group and H is a closed subgroup, then the equivalence relation $xE_H^G y \Leftrightarrow y \in xH$, whose quotient space is the set G/H of left-cosets hH of H , is tame (see Kechris [21], 12.17). We call a standard Borel G -space X *tame* if the equivalence relation E_G^X is tame. In particular, every standard Borel G -space X is tame, when G is compact (second countable); see Zimmer [35], 2.1.21.

A *Borel equivalence* relation on a standard Borel space X is an equivalence relation $E \subseteq X^2$ which is a Borel set in X^2 . It is *countable* if every equivalence class $[x]_E, x \in X$, is countable. Every countable Borel equivalence relation E on X is of the form $E = E_G^X$, for a Borel action of a countable group G on X (Feldman-Moore [8]; see also Kechris [21], 18.16). A measure μ on X is called *E -invariant* if it is G -invariant for any countable G with $E = E_G^X$ (this is easily seen to be independent of G and the action). It is *E -ergodic* if every E -invariant Borel set $A \subseteq X$ has μ -measure 0 or 1.

Finally, given equivalence relations E, F on standard Borel spaces X, Y , resp., we say that E is *Borel reducible* to F , in symbols,

$$E \leq_B F,$$

if there is a Borel function $f : X \rightarrow Y$ such that

$$xEy \Leftrightarrow f(x)Ff(y).$$

We call such an f a *reduction* of E into F . We also let

$$E \sim_B F \Leftrightarrow E \leq_B F \text{ and } F \leq_B E$$

and

$$E <_B F \Leftrightarrow E \leq_B F \text{ and } F \not\leq_B E.$$

2. BASIC FACTS ABOUT COCYCLES

Let G be a lsc group, and X a standard Borel G -space with invariant probability Borel measure μ . Let H be a lsc group. A *cocycle* of this G -space into H is a Borel map $\alpha : G \times X \rightarrow H$ such that for all $g, h \in G$

$$\alpha(hg, x) = \alpha(h, g \cdot x)\alpha(g, x) \quad \mu\text{-a.e.}(x).$$

If this equation holds for *all* x we say that α is a *strict cocycle*. If $\beta : G \times X \rightarrow H$ is also a cocycle, we say that α is *equivalent or cohomologous* to β , in symbols $\alpha \sim \beta$, if there is a Borel map $A : X \rightarrow H$ such that for all $g \in G$,

$$\alpha(g, x) = A(g \cdot x)\beta(g, x)A(x)^{-1} \quad \mu\text{-a.e.}(x).$$

This is clearly an equivalence relation.

For example, suppose that $E = E_G^X$ and $F = E_H^Z$, where H acts freely on Z . Let $f : X \rightarrow Z$ be a function such that xEy implies $f(x)Ff(y)$. Then the function $\alpha : G \times X \rightarrow H$ defined by $f(g \cdot x) = \alpha(g, x) \cdot f(x)$ is a cocycle. Moreover, if $A : X \rightarrow H$ is any Borel function, if we define $f_0 : X \rightarrow Z$ by $f_0(x) = A(x) \cdot f(x)$ and if we define $\alpha_0 : G \times X \rightarrow H$ by $f_0(g \cdot x) = \alpha_0(g, x) \cdot f_0(x)$, then the cocycle α_0 is equivalent to α ; in fact, we have $\alpha_0(g, x) = A(g \cdot x)\alpha(g, x)(A(x))^{-1}$.

The main results in this paper are based on various theorems that show that cocycles α as above, under certain circumstances, reduce, *i.e.*, are equivalent to, cocycles β , whose range $\beta(G \times X)$ is contained in a “small” subgroup of H . The following elementary reformulation is useful for this purpose. We need the following terminology, using the above notation: If H acts on some set Y , then a function $f : X \rightarrow Y$ will be called *α -invariant* if for all $g \in G$,

$$\alpha(g, x) \cdot f(x) = f(g \cdot x) \quad \mu\text{-a.e.}(x).$$

Also for any closed subgroup $H_0 \subseteq H$ we denote by H/H_0 the space of left-cosets hH_0 of H_0 and we view H as acting on H/H_0 by $h_1 \cdot hH = h_1hH$.

We now have:

Proposition 2.1 (see Zimmer [35], 4.2.18(b)). *Let $H_0 \subseteq H$ be a closed subgroup of H . Then a cocycle $\alpha : G \times X \rightarrow H$ is equivalent to a cocycle $\beta : G \times X \rightarrow H$ with $\beta(G \times X) \subseteq H_0$ iff there exists a Borel α -invariant function $f : X \rightarrow H/H_0$.*

Proof. Say $\alpha(g, x) = A(g \cdot x)\beta(g, x)A(x)^{-1} \quad \mu\text{-a.e.}(x)$, where $A : X \rightarrow H$ is Borel and $\beta(G \times X) \subseteq H_0$. Then if $f(x) = A(x)H_0, f : X \rightarrow H/H_0$ is Borel and for each $g, \alpha(g, x) \cdot f(x) = f(g \cdot x) \quad \mu\text{-a.e.}(x)$, as $\beta(g, x) \in H_0$. Conversely, given a Borel α -invariant f , and letting $i : H/H_0 \rightarrow H$ be a Borel injection such that $i(hH) \in hH$, we let $A(x) = i(f(x))$. Then as $\alpha(g, x) \cdot f(x) = f(g \cdot x) \quad \mu\text{-a.e.}(x)$, we have that $\alpha(g, x)A(x) \in A(g \cdot x)H_0 \quad \mu\text{-a.e.}(x)$, so if $\beta(g, x) = A(g \cdot x)^{-1}\alpha(g, x)A(x)$, then $\beta(g, x) \in H_0 \quad \mu\text{-a.e.}(x)$ and $\alpha \sim \beta$. Now we change $\beta(g, x)$ on a Borel set in $G \times X$, whose G -sections are μ -null to make sure that $\beta(g, x) \in H_0$ for all $g, x, i.e., \beta(G \times X) \subseteq H_0.$ ◻

The next proposition, called the *Cocycle Reduction Lemma*, gives a convenient tool for showing that cocycles reduce to other ones with small range. Recall that the measure μ on the standard G -space X is called *ergodic* if every G -invariant Borel set has μ -measure 0 or 1. Also a Borel action of a lsc group H on a standard Borel space Y is called *tame* if the corresponding equivalence relation E_Y^H is *tame, i.e.*,

there is Borel $S : Y \rightarrow Z$, Z some Polish space, such that $y_1 E_Y^H y_2 \Leftrightarrow S(y_1) = S(y_2)$. We can of course always take Z to be $[0,1]$ here.

Proposition 2.2 (see Zimmer [35], 5.2.11). *Suppose X is a standard Borel G -space with ergodic, invariant measure μ and $\alpha : G \times X \rightarrow H$ is a cocycle. Assume also that Y is a tame standard Borel H -space. If there is an α -invariant Borel function $f : X \rightarrow Y$, then there is $y_0 \in Y$ such that $\alpha \sim \beta$, for a cocycle $\beta : G \times X \rightarrow H$ with $\beta(G \times X) \subseteq H_{y_0} =$ the stabilizer of y_0 . Moreover $f(x) \in H \cdot y_0$, μ -a.e.(x).*

Proof. Let $S : Y \rightarrow [0, 1]$ be Borel with $y_1 E_Y^H y_2 \Leftrightarrow S(y_1) = S(y_2)$. Let $\hat{f}(x) = S(f(x))$. Then, as $\alpha(g, x) \cdot f(x) = f(g \cdot x)$ μ -a.e.(x), clearly $\hat{f}(g \cdot x) = \hat{f}(x)$ μ -a.e.(x). So $\forall g \forall_\mu^* x (\hat{f}(g \cdot x) = \hat{f}(x))$ (where $\forall_\mu^* x P(x) \Leftrightarrow \{x : P(x)\}$ is μ -conull), thus, by Fubini, $\forall_\mu^* x \forall_\lambda^* g (\hat{f}(g \cdot x) = \hat{f}(x))$, where λ is the Haar measure on G . Now the set

$$X_0 = \{x : g \mapsto \hat{f}(g \cdot x) \text{ is } \lambda\text{-a.e. constant}\} = \{x : \forall_\lambda^* g \forall_\lambda^* h (\hat{f}(g \cdot x) = \hat{f}(h \cdot x))\}$$

is Borel (see, e.g., Kechris [21], 17.25) and, by the translation invariance of λ , G -invariant. So if we define $\hat{f}_0(x) =$ the value of $\hat{f}(g \cdot x)$, λ -a.e.(g), if $x \in X_0$, $\hat{f}_0(x) = 0$, if $x \notin X_0$, then clearly \hat{f}_0 is G -invariant and $\hat{f}_0 = \hat{f}$, μ -a.e. By ergodicity, it follows that \hat{f}_0 , and thus \hat{f} , is constant μ -a.e., and this means that f takes values in a single H -orbit, say $H \cdot y_0$, μ -a.e., and we can assume that f actually takes values in $H \cdot y_0$ everywhere, by changing f on a set of measure 0. Now the H -action on $H \cdot y_0$ can be clearly identified with the H -action on H/H_{y_0} and thus, since f is an α -invariant function, by 2.1, we have that $\alpha \sim \beta$ with $\beta(G \times X) \subseteq H_{y_0}$. \dashv

We will next discuss the concepts of *induced actions and cocycles*.

Suppose $\Gamma \subseteq G$ is a closed subgroup of G and fix a Borel transversal T for the left-cosets of Γ with $1 \in T$ (so T meets every such coset in exactly one point). We can clearly identify T with G/Γ , identifying t with $t\Gamma$. So the action of G on G/Γ induces a Borel action of G on T given by

$$g \cdot t = \text{the unique element of } T \text{ in the coset } gt\Gamma.$$

Also let

$$\begin{aligned} \rho(g, t) &= \text{the unique element } \gamma \in \Gamma \\ &\text{such that } (g \cdot t)\gamma = gt \\ &= (g \cdot t)^{-1}gt. \end{aligned}$$

Then it is easy to check that ρ is a strict cocycle, $\rho : G \times T \rightarrow \Gamma$. We call this the *cocycle associated* to Γ (and the choice of T).

Now assume Γ acts in a Borel way on a standard Borel space X with invariant measure μ . Assume also that Γ is a *lattice* in G , i.e., Γ is discrete and the action of G on G/Γ , and thus equivalently on T , admits an invariant probability Borel measure, say ν . We define the *induced action* of G as follows: Let $Y = X \times T$ and define the action of G on Y by

$$g \cdot (x, t) = (\rho(g, t) \cdot x, g \cdot t).$$

It is easy to check, using Fubini, that $\mu \times \nu$ is an invariant measure for this action: If $p : X \times T \rightarrow \mathbb{R}$ is bounded Borel, then

$$\begin{aligned} \int \int (g \cdot p) d(\mu \times \nu) &= \int \left(\int p(g \cdot (x, t)) d\mu(x) \right) d\nu(t) \\ &= \int \left(\int p(\rho(g, t) \cdot x, g \cdot t) d\mu(x) \right) d\nu(t) \\ &= \int \left(\int p(x, g \cdot t) d\mu(x) \right) d\nu(t) \\ &= \int \left(\int p(x, g \cdot t) d\nu(t) \right) d\mu(x), \end{aligned}$$

so

$$\begin{aligned} \int \int (g \cdot p) d(\mu \times \nu) &= \int \left(\int p(g \cdot (x, t)) d\mu(x) \right) d\nu(t) \\ &= \int \left(\int p(x, t) d\nu(t) \right) d\mu(x) \\ &= \int \int p d(\mu \times \nu). \end{aligned}$$

Next we check that if μ is ergodic, so is $\mu \times \nu$. Indeed, let $A \subseteq X \times T$ be Borel G -invariant, and put $A^t = \{x : (x, t) \in A\}$. Then $\rho(g, t) \cdot A^t = A^{g \cdot t}$ and A^1 is Γ -invariant, so $\mu(A^1) = 0$ or 1 , thus, since $\rho(g, 1) \cdot A^1 = A^{g \cdot 1}$, all A^t have the same μ -measure as A^1 , so $(\mu \times \nu)(A) = 0$ or 1 .

Now given any strict cocycle $\alpha : \Gamma \times X \rightarrow H$ we define the strict cocycle $\beta : G \times Y \rightarrow H$ by

$$\beta(g, (x, t)) = \alpha(\rho(g, t), x),$$

and call this the *induced cocycle*.

For example, suppose that $E = E_\Gamma^X$ and $F = E_H^Z$, where H acts freely on Z . Let $f : X \rightarrow Z$ be a function such that xEx' implies $f(x)Ff(x')$. Let $\alpha : \Gamma \times X \rightarrow H$ be the cocycle defined by $f(\gamma \cdot x) = \alpha(\gamma, x) \cdot f(x)$. Let G act on $Y = X \times T$ by $g \cdot (x, t) = (\rho(g, t) \cdot x, g \cdot t)$. Define $f_0 : Y \rightarrow Z$ by $f_0(x, t) = f(x)$. Let $E_0 = E_G^Y$. Then yE_0y' implies $f_0(y)Ff_0(y')$. Let $\alpha_0 : G \times Y \rightarrow H$ be the cocycle defined by $f_0(g \cdot y) = \alpha_0(g, y) \cdot f_0(y)$. Then α_0 is induced from α ; that is, we have $\alpha_0(g, (x, t)) = \alpha(\rho(g, t), x)$.

We have the following fact.

Proposition 2.3. *Suppose that Γ is a lattice in G , that X is a standard Borel Γ -space with invariant measure μ , that $\alpha : \Gamma \times X \rightarrow H$ is a strict cocycle and that $\beta : G \times Y \rightarrow H$ is the induced cocycle for the induced G -space Y . Assume that H_0 is a closed subgroup of H and that there is $\beta_1 : G \times Y \rightarrow H$ with $\beta_1 \sim \beta$ and with $\beta_1(G \times Y) \subseteq H_0$. Then there is $\alpha_1 : \Gamma \times X \rightarrow H$ with $\alpha_1 \sim \alpha$ and $\alpha_1(\Gamma \times X) \subseteq H_0$.*

Proof. By 2.1, there is a β -invariant Borel function $f : Y = (X \times T) \rightarrow H/H_0$, i.e., for all $g \in G$,

$$\beta(g, (x, t)) \cdot f(x, t) = f(g \cdot (x, t)) \quad (\mu \times \nu)\text{-a.e.}(x, t).$$

Let U be the space of all Borel functions from X into H/H_0 , two functions being identified if they agree μ -a.e. Consider then the map $\Phi : T \rightarrow U$ given by $\Phi(t)(x) =$

$f(x, t)$. Let Γ act on U by

$$\gamma \cdot p(x) = \alpha(\gamma, \gamma^{-1} \cdot x) \cdot p(\gamma^{-1} \cdot x).$$

We then claim that Φ is ρ -invariant, *i.e.*, for any $g \in G$,

$$\rho(g, t) \cdot \Phi(t) = \Phi(g \cdot t) \quad \nu\text{-a.e.}(t).$$

Indeed,

$$\begin{aligned} (\rho(g, t) \cdot \Phi(t))(x) &= \alpha(\rho(g, t), \rho(g, t)^{-1} \cdot x) \cdot \Phi(t)(\rho(g, t)^{-1} \cdot x) \\ &= \alpha(\rho(g, t), \rho(g, t)^{-1} \cdot x) \cdot f(\rho(g, t)^{-1} \cdot x, t). \end{aligned}$$

Now

$$\begin{aligned} \beta(g, (x, t)) \cdot f(x, t) &= f(g \cdot (x, t)) \\ &= f(\rho(g, t) \cdot x, g \cdot t) \quad (\mu \times \nu)\text{-a.e.}(x, t), \end{aligned}$$

so

$$\alpha(\rho(g, t), x) \cdot f(x, t) = f(\rho(g, t) \cdot x, g \cdot t) \quad (\mu \times \nu)\text{-a.e.}(x, t).$$

Thus for $\nu\text{-a.e.}(t)$ and using the Γ -invariance of μ to substitute $\rho(g, t)^{-1} \cdot x$ for x , we have

$$\begin{aligned} f(x, g \cdot t) &= \alpha(\rho(g, t), \rho(g, t)^{-1} \cdot x) \cdot f(\rho(g, t)^{-1} \cdot x, t) \\ &= (\rho(g, t) \cdot \Phi(t))(x) \quad \mu\text{-a.e.}(x), \end{aligned}$$

and, since

$$f(x, g \cdot t) = \Phi(g \cdot t)(x),$$

this shows that

$$\rho(g, t) \cdot \Phi(t) = \Phi(g \cdot t) \quad \nu\text{-a.e.}(t).$$

We now claim that this implies that there is a fixed point $p \in U$ for the Γ -action, *i.e.*, $\gamma \cdot p = p$ for all $\gamma \in \Gamma$. Granting this we have for any $\gamma \in \Gamma$,

$$\begin{aligned} \gamma \cdot p(x) &= \alpha(\gamma, \gamma^{-1} \cdot x) \cdot p(\gamma^{-1} \cdot x) \\ &= p(x) \quad \mu\text{-a.e.}(x), \end{aligned}$$

so putting $\gamma \cdot x$ for x and using again the Γ -invariance of μ , we have

$$p(\gamma \cdot x) = \alpha(\gamma, x) \cdot p(x) \quad \mu\text{-a.e.}(x),$$

i.e., p is an α -invariant function from X into H/H_0 , so by 2.1, $\alpha \sim \alpha_1$, for some cocycle $\alpha_1 : \Gamma \times X \rightarrow H$ with $\alpha_1(\Gamma \times X) \subseteq H_0$.

To prove the existence of such a fixed point p we argue as follows (see Zimmer [35], 4.2.19): Let $\sigma : G \rightarrow \Gamma$ be defined by $\sigma(g) = \rho(g, 1)$. Then notice that for any $g \in G$, $t \in T$, $\sigma(gt) = \rho(gt, 1) = \rho(g, t \cdot 1)\rho(t, 1) = \rho(g, t)\sigma(t)$, so $\rho(g, t) = \sigma(gt)\sigma(t)^{-1}$. It follows that for each $g \in G$,

$$\sigma(gt)^{-1} \cdot \Phi(g \cdot t) = \sigma(t)^{-1} \cdot \Phi(t) \quad \nu\text{-a.e.}(t).$$

So, by Fubini, there is $t_0 \in T$ such that for almost all $g \in G$ (with respect to the Haar measure), $\sigma(gt_0)^{-1} \cdot \Phi(g \cdot t_0) = \sigma(t_0)^{-1} \cdot \Phi(t_0)$. Put $p = \sigma(t_0)^{-1} \cdot \Phi(t_0)$. Then viewing now Φ as defined on G/Γ by identifying $t \in T$ with $t\Gamma \in G/\Gamma$, and using the translation invariance of the Haar measure, we conclude that for almost all $g \in G$, $\sigma(g)^{-1} \cdot \Phi(g\Gamma) = p$. Now notice that for $\gamma \in \Gamma$, $\sigma(g\gamma) = \sigma(g)\gamma$, so $\gamma^{-1} \cdot p = \gamma^{-1}\sigma(g)^{-1} \cdot \Phi(g\Gamma) = \sigma(g\gamma)^{-1} \cdot \Phi(g\Gamma) = \sigma(g\gamma)^{-1} \cdot \Phi(g\gamma\Gamma)$, for almost all $g \in G$, so, again by translation invariance, $\gamma^{-1} \cdot p = p$ and we are done. \dashv

Proposition 2.4. *Suppose H is lcsc and $H_1 \subseteq H$ is a closed subgroup. Suppose G acts on X with invariant, ergodic measure μ and $\alpha : G \times X \rightarrow H_1$ is a Borel cocycle. Assume that viewing α as a cocycle into H , $\alpha : G \times X \rightarrow H$, we have a cocycle $\beta : G \times X \rightarrow H$ with $\beta \sim \alpha$ and $\beta(G \times X) \subseteq H_0$, where H_0 is a closed subgroup of H . If the action of H_0 on H/H_1 is tame, then there is $h \in H$ and a cocycle $\alpha_1 : G \times X \rightarrow H_1$ such that $\alpha \sim \alpha_1$ and $\alpha_1(G \times X) \subseteq H_1 \cap hH_0h^{-1}$.*

In particular, this conclusion holds if H_0 is compact or if H_0 is arbitrary (closed) and H_1 has finite index in H .

Proof. By assumption and 2.1 there is an α -invariant Borel function $f : X \rightarrow H/H_0$. Now we claim that the action of H_1 on H/H_0 is tame. Granting this, we have, by 2.2, that there is $hH_0 \in H/H_0$ such that α is equivalent (as a cocycle into H_1) to a cocycle taking values in the stabilizer of hH_0 in the action by H_1 . This stabilizer is the set $\{h_1 \in H_1 : h_1 \cdot hH_0 = hH_0\} = H_1 \cap hH_0h^{-1}$.

To prove the claim, notice that there is a canonical bijection between the double cosets $H_1hH_0, h \in H$, of (H_1, H_0) and the orbits of the H_1 action on H/H_0 , namely $H_1hH_0 \mapsto H_1 \cdot (hH_0)$. Thus the action of H_1 on H/H_0 is tame iff the equivalence relation induced by the double cosets is tame. Clearly the equivalence relation induced by the double cosets H_1hH_0 is tame iff the equivalence relation induced by the double cosets H_0hH_1 is tame (using $x \mapsto x^{-1}$) iff the action of H_0 into H/H_1 is tame, and this last statement is assumed in our hypothesis.

Finally, notice that if H_0 is compact or H_1 has finite index in H , clearly H_0 acts tamely on H/H_1 . ◻

Suppose again that G acts on X preserving a measure μ . By a *finite extension* of this action we mean a standard Borel G -space \hat{X} with an invariant measure $\hat{\mu}$, together with a Borel map $\pi : \hat{X} \rightarrow X$ such that

- (i) for all $g \in G$ and for all $x \in \hat{X}$, we have $\pi(g \cdot x) = g \cdot \pi(x)$,
- (ii) π is finite-to-1,
- (iii) $\pi_*(\hat{\mu}) = \mu$.

If $\alpha : G \times X \rightarrow H$ is a cocycle, then its *lift* to \hat{X} , $\hat{\alpha} : G \times \hat{X} \rightarrow H$, is defined by $\hat{\alpha}(g, \hat{x}) = \alpha(g, \pi(\hat{x}))$.

The argument in the next proposition is similar to that of the proof of [35], 9.2.6.

Proposition 2.5. *Suppose that X is a standard Borel G -space with ergodic, invariant measure μ . Let $\alpha : G \times X \rightarrow H$ be a Borel cocycle. Suppose $H_0 \subseteq H$ is a closed subgroup of H with finite index. Then there exists a finite extension \hat{X} of X which is also ergodic such that the lift $\hat{\alpha}$ of α to \hat{X} , $\hat{\alpha} : G \times \hat{X} \rightarrow H$, is equivalent to a cocycle taking values in H_0 .*

Proof. Let H_1 be a minimal closed subgroup of H satisfying $H_0 \subseteq H_1 \subseteq H$ such that α is equivalent to a cocycle β taking values into H_1 . Then β has the property that no cocycle equivalent to it takes values into a proper closed subgroup of H_1 containing H_0 . So by replacing α by β if necessary, we can assume that α is not equivalent to a cocycle taking values into a proper closed subgroup of H containing H_0 , and show how to find an ergodic finite extension \hat{X} of X so that the lift $\hat{\alpha}$ of α is equivalent to a cocycle taking values in H_0 .

Let $\hat{X} = X \times (H/H_0)$ and define the action of G on \hat{X} by

$$g \cdot (x, hH_0) = (g \cdot x, \alpha(g, x)hH_0)$$

and let $\pi(x, hH_0) = x$. We give \hat{X} the product measure $\mu \times \nu$, where ν is the counting measure on H/H_0 . Clearly this is a finite extension of X . Consider the second projection map $\rho(x, hH_0) = hH_0, \rho : \hat{X} \rightarrow H/H_0$. It is clearly $\hat{\alpha}$ -invariant, so by 2.1, $\hat{\alpha}$ is equivalent to a cocycle taking values into H_0 .

So it only remains to show that $\mu \times \nu$ is ergodic. Assume $A \subseteq \hat{X}$ is a G -invariant Borel set and $(\mu \times \nu)(A) > 0$, in order to show that $(\mu \times \nu)(A) = 1$. Then $\mu(\pi(A)) > 0$, so since $\pi(A)$ is a G -invariant Borel subset of X , we have, by ergodicity in X , that $\mu(\pi(A)) = 1$. Put $B = \pi(A)$, and letting $F =$ power set of H/H_0 , define $f : X \rightarrow F$ by

$$f(x) = \begin{cases} \rho(\pi^{-1}(x) \cap A), & \text{if } x \in B, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Then considering the obvious action of H on F (induced by its action on H/H_0) we see that f is α -invariant; so by 2.2 there is $s_0 \in F$ such that $f(x) \in H \cdot s_0$ μ -a.e.(x), and there is β with $\alpha \sim \beta$ and β taking values in the stabilizer (in H) of s_0 , say H_1 . If $s_0 \neq H/H_0$, then clearly H_1 is a closed proper subgroup of H with $H_0 \subseteq H_1$, a contradiction. So $s_0 = H/H_0$ and $f(x) = H/H_0$ μ -a.e.(x), thus $\nu(A_x) = 1$ μ -a.e.(x), so $(\mu \times \nu)(A) = 1$. -1

Proposition 2.6. *Suppose that X is a standard Borel G -space with ergodic, invariant measure μ . Let $\alpha : G \times X \rightarrow H$ be a cocycle. Let \hat{X} be a finite extension of X and $\hat{\alpha} : G \times \hat{X} \rightarrow H$ the lift of α . If $\hat{\alpha}$ is equivalent to a cocycle taking values in a compact subgroup of H , then α is equivalent to a cocycle taking values in a compact subgroup of H .*

Proof. Say $\hat{\alpha} \sim \hat{\beta}$ with $\hat{\beta}(G \times \hat{X}) \subseteq K$, K a compact subgroup of H . Then, by 2.1 there is a Borel map $\hat{f} : \hat{X} \rightarrow H/K$ which is $\hat{\alpha}$ -invariant. Let F be the standard Borel space of non-empty finite subsets of H/K and let $f : X \rightarrow F$ be defined by $f(x) = \hat{f}(\pi^{-1}(x))$. Then it is easy to check that f is α -invariant. By considering the obvious action of H on F (induced by the action of H on H/K), we conclude, using 2.2, that, if this action is tame, then α is equivalent to a cocycle taking values into the stabilizer of some point for the action of H on F , which is clearly compact.

To see that this action is tame, it is enough to check, for each n , that the action of H on the subsets of H/K of cardinality n is tame, and this is the same as checking that the action of $H \times S_n$ ($S_n =$ the symmetric group in n elements) on the space of distinct n -tuples $\langle a_1, \dots, a_n \rangle$ from H/K is tame, where for $(h, \rho) \in H \times S_n$, $(h, \rho) \cdot \langle a_1, \dots, a_n \rangle = \langle h \cdot a_{\rho(1)}, \dots, h \cdot a_{\rho(n)} \rangle$. Since H has finite index in $H \times S_n$, it is enough to check that the action of H on the space of distinct n -tuples is tame. Call E the corresponding equivalence relation. Now the action of K on the space of distinct n -tuples is tame, as K is compact. Denote by E' its corresponding equivalence relation. Let $\vartheta : H/K \rightarrow H$ be Borel such that $\vartheta(hK) \cdot hK = K$. Then notice that $\langle a_1, \dots, a_n \rangle E \langle b_1, \dots, b_n \rangle$ iff $\vartheta(a_1) \cdot \langle a_1, \dots, a_n \rangle E \vartheta(b_1) \cdot \langle b_1, \dots, b_n \rangle$ iff $\vartheta(a_1) \cdot \langle a_1, \dots, a_n \rangle E' \vartheta(b_1) \cdot \langle b_1, \dots, b_n \rangle$, so E is tame as well. -1

Here is the final fact that we will need.

Proposition 2.7. *Suppose X is a standard Borel G -space with ergodic, invariant measure and $\alpha : G \times X \rightarrow H$ is a cocycle. Let $\pi : G' \rightarrow G$ be a continuous homomorphism such that $\pi(G')$ has finite index in G . Let G' act on X by $g' \cdot x = \pi(g') \cdot x$ and let $\alpha' : G' \times X \rightarrow H$ be given by $\alpha'(g', x) = \alpha(\pi(g'), x)$. Assume that*

α' is equivalent to a cocycle taking values in a compact subgroup of H . Then α is equivalent to a cocycle taking values in a compact subgroup of H .

Proof. Suppose $\alpha' \sim \beta'$ with $\beta'(G' \times X) \subseteq K$, K a compact subgroup of H . Then, by 2.1, there is an α' -invariant map $f : X \rightarrow H/K$. As in the proof of 2.3, let U be the space of all Borel maps from X into H/K , two functions being identified if they agree a.e. Let G act on U by

$$g \cdot p(x) = \alpha(g, g^{-1} \cdot x) \cdot p(g^{-1} \cdot x).$$

Then f is $\pi(G')$ -invariant, so the orbit of f under G is finite, say equal to the set $\{f_0, f_1, \dots, f_{n-1}\}$. Then let f^* be the Borel map from X into the space F of non-empty finite subsets of H/K , defined by $f^*(x) = \{f_0(x), \dots, f_{n-1}(x)\}$. Then if we let for $g \in G$, $g \cdot f^*(x) = \{g \cdot f_0(x), \dots, g \cdot f_{n-1}(x)\}$, $g \cdot f^* = f^*$ a.e., that is, $g \cdot f^*(x) = \alpha(g, g^{-1} \cdot x) \cdot f^*(g^{-1} \cdot x) = f^*(x)$, a.e.(x), thus $\alpha(g, x) \cdot f^*(x) = f^*(g \cdot x)$, a.e.(x), which means that f^* is α -invariant, where we view H as acting on F by the action induced from the action of H on H/K . Now the action of H on F is tame, by the proof in 2.6, and has compact stabilizers, so, by 2.2, α is equivalent to a cocycle taking values in a compact subgroup of H . \dashv

3. COCYCLE REDUCTION RESULTS

Recall that a lcsc group G is *amenable* if there is a (left)-invariant *mean* on $L^\infty(G)$ ($= L^\infty(G, \lambda)$, where λ is the Haar measure on G), i.e., a normalized ($\Lambda(1) = 1$) positive linear functional $\Lambda \in (L^\infty(G))^*$, such that $\Lambda(f) = \Lambda(f_g)$, where $f_g(h) = f(gh)$.

Also a lcsc group G is called a *Kazhdan group* (or a group with *property (T)*) if for any unitary representation $\pi : G \rightarrow U(H)$, where $U(H)$ is the unitary group of a separable Hilbert space H , if π almost has invariant vectors (i.e., for every $\epsilon > 0$ and every compact $K \subseteq G$ there is a unit vector $v \in H$ such that $\|\pi(g)(v) - v\| < \epsilon, \forall g \in K$), then π has a non-0 invariant vector.

Theorem 3.1 (Zimmer [35], 9.1.1). *Let G be a Kazhdan group and X a standard Borel G -space with an invariant, ergodic measure. Let H be an amenable group. Then for every cocycle $\alpha : G \times X \rightarrow H$ there is an equivalent cocycle $\beta : G \times X \rightarrow H$ and a compact subgroup $K \subseteq H$ such that $\beta(G \times X) \subseteq K$.*

Theorem 3.2 (Zimmer [36], 2.2). *Let G be a Kazhdan group and X a standard Borel G -space with an ergodic, invariant measure. Let Γ be a countable group. Then for every cocycle $\alpha : G \times X \rightarrow \Gamma$ there is an equivalent cocycle $\beta : G \times X \rightarrow \Gamma$ and a finitely generated subgroup $\Gamma_0 \subseteq \Gamma$ with $\beta(G \times X) \subseteq \Gamma_0$.*

Next we will review some cocycle reduction results for actions of algebraic groups. We will adapt the terminology and (with some minor exceptions) the notation of [35], Chapter 3. In particular, we take Ω to be an algebraically closed field of characteristic 0 containing \mathbb{R} and all p -adic fields \mathbb{Q}_p, p a prime. Then an algebraic group G is a subgroup of some $GL_n(\Omega)$ ($=$ the group of invertible $n \times n$ matrices over Ω), which is Zariski closed in $GL_n(\Omega)$, i.e., consists of all matrices M in $GL_n(\Omega)$ which satisfy a set of equations $f_1(M) = 0, \dots, f_k(M) = 0$, where each f_i is a polynomial in $\Omega[x_1, \dots, x_{n^2}]$ (we thus view M as a member of Ω^{n^2}). If the equations defining G have coefficients in a subfield $k \subseteq \Omega$, then we call G a

k-group. For any subring $A \subseteq \Omega$, we let $\mathrm{GL}_n(A) = \{(a_{ij}) \in \mathrm{GL}_n(\Omega) : a_{ij} \in A \text{ and } (\det(a_{ij}))^{-1} \in A\}$, and define for any algebraic group $G \subseteq \mathrm{GL}_n(\Omega)$,

$$G(A) = G \cap \mathrm{GL}_n(A).$$

In particular, if G is a k -group and $A = k$, $G(k)$ is the group of all matrices in G with coefficients in k .

The topology of an algebraic group G is always the Zariski topology on G . When G is a k -group and k is one of the fields $\mathbb{R}, \mathbb{C}, \mathbb{Q}_p$, then $G(k)$, as a group of matrices over these fields, is a lsc group with the topology it inherits from k^{n^2} . So any topological notions about the group $G(k)$ refer to this topology (sometimes called the *Hausdorff topology* to distinguish it from the Zariski topology).

The proofs of our main results are based on the *superrigidity theorems* of Zimmer [35]. We only state the particular instances of these results that we need.

The first result is a special case of [35], 10.1.6, and we use the context of Chapter 10 in that book.

Theorem 3.3 (Zimmer [35], 10.1.6). *Suppose p, q are distinct rational primes and let G be a connected semisimple \mathbb{Q}_p -group which is moreover algebraically simply connected, almost \mathbb{Q}_p -simple and satisfies $\mathbb{Q}_p\text{-rank}(G) \geq 2$. Let H be a connected \mathbb{Q}_q -simple, \mathbb{Q}_q -group. Assume that X is a standard Borel $G(\mathbb{Q}_p)$ -space, with invariant, ergodic measure, and that $\alpha : G(\mathbb{Q}_p) \times X \rightarrow H(\mathbb{Q}_q)$ is a cocycle. For every proper \mathbb{Q}_q -subgroup L of H , assume that α is not equivalent to any cocycle with range contained in $L(\mathbb{Q}_q)$. Then α is equivalent to a cocycle with range contained in a compact subgroup of $H(\mathbb{Q}_q)$.*

We will briefly comment on why this is a special case of Zimmer [35], 10.1.6. We take $S = \{p\}$ in that theorem, so that our G is G_p in his notation, and $\mathrm{rank}(G) = \mathbb{Q}_p\text{-rank}(G)$ (see top of page 189 in [35]—note however that what he calls G is our $G(\mathbb{Q}_p)$). Then by the first paragraph of Chapter 10, page 187 of this book, the hypothesis that $\mathbb{Q}_p\text{-rank}(G) \geq 2$ implies that $G(\mathbb{Q}_p)$ is not compact and thus has no compact factors, *i.e.*, there is no proper normal \mathbb{Q}_p -subgroup $H \subseteq G$ with $G(\mathbb{Q}_p)/H(\mathbb{Q}_p)$ compact, since the hypothesis of almost \mathbb{Q}_p -simplicity means exactly that every such H is finite. Finally, we have to check that X is an irreducible $G(\mathbb{Q}_p)$ -space, which according to the definition used in this context in page 188, line 5 in [35] (with the word “proper” omitted in this line, being a typo), this reduces in our case to the fact that the action is ergodic, since almost \mathbb{Q}_p -simplicity means exactly that there are no infinite normal \mathbb{Q}_p -subgroups of G . Thus all the hypotheses of 10.1.6 are satisfied. Then since, in the notation of 10.1.6, $k = \mathbb{Q}_q$ in our case with $q \neq p$, (b), (c) of this theorem do not apply, thus by (a), α is equivalent to a cocycle with values in a compact subgroup of $H(\mathbb{Q}_q)$.

The second and final superrigidity result that we will need is the following, which is a special case of [35], 5.2.5.

Theorem 3.4 (Zimmer [35], 5.2.5). *Let G be a connected, semisimple \mathbb{R} -group and assume that $G(\mathbb{R})$ is connected (in the Hausdorff topology), that every normal subgroup of $G(\mathbb{R})$ is contained in its center which is finite, and that $\mathbb{R}\text{-rank}(G) \geq 2$. Let X be a standard Borel G -space with invariant, ergodic measure. Let H be a connected k -simple, k -group, where $k = \mathbb{Q}_p$ or \mathbb{R} . If $\alpha : G(\mathbb{R}) \times X \rightarrow H(k)$ is a cocycle such that α is not equivalent to a cocycle with range contained in a subgroup*

of $H(k)$ of the form $L(k)$, for a proper k -subgroup L of H , then either α is equivalent to a cocycle taking values into a compact subgroup of $H(k)$ or else $k = \mathbb{R}$ and there is a non-trivial \mathbb{R} -morphism from G into H .

We will again briefly comment on why this is a special case of [35], 5.2.5. Since $G(\mathbb{R})$ is connected, we have that $G_{\mathbb{R}}^0$ in Zimmer’s notation, which is the connected component of the identity in $G(\mathbb{R})$ with respect to the Hausdorff topology (see page 35, paragraph 4), is the same as $G(\mathbb{R})$. The irreducibility condition (see [35], 2.2.11) becomes just ergodicity, since every normal subgroup of $G(\mathbb{R})$ is assumed to be contained in the center. So we must have (i) or (ii) in 5.2.5. If $k = \mathbb{R}$ and $H(k)$ is compact, there is nothing to prove. Otherwise, since H is assumed to be k -simple we have a k -morphism $\pi : G \rightarrow H$, such that α is equivalent to the cocycle $\alpha_0(g, x) = \pi(g)$. If π is non-trivial we are done; otherwise α_0 obviously takes values in $\{1\}$, which is compact. If $k = \mathbb{Q}_p$, then (iii) applies and we are also done.

We are now ready to derive a cocycle reduction result that we will need in Section 4.

Below let F be the quadratic form defined by

$$F(x_1, x_2, \dots, x_7) = x_1^2 + \dots + x_7^2$$

and let SO_F be the \mathbb{Q} -group (thus \mathbb{Q}_p -group) of all 7×7 matrices preserving F and having determinant 1, that is, all 7×7 matrices A with $\det(A) = 1, AA^t = I$.

Note that $SO_F(\mathbb{R})$ is compact. Consequently, for any set S of primes, the group $\Gamma_S = SO_F(\mathbb{Z}[S^{-1}])$ may be thought of as a lattice in $\prod_{p \in S} SO_F(\mathbb{Q}_p)$. In the sequel, we will be using induction of cocycles together with 10.1.6 of [35] to show that, if S and T are sets of primes and if $S \neq T$, then a finite measure preserving action of Γ_S cannot be reducible to a free finite measure preserving action of Γ_T .

Fix an integer $n \geq 3$. Note, for all primes p , that $\Gamma'_p = SL_n(\mathbb{Z}[1/p])$ is a lattice in $SL_n(\mathbb{R}) \times SL_n(\mathbb{Q}_p)$; moreover, if we project Γ'_p into $SL_n(\mathbb{Q}_p)$, it becomes a dense subgroup, and is not a lattice. So, if p and q are distinct primes, then it will be difficult to use induction of cocycles and 10.1.6 of [35] to show that a finite measure preserving action of Γ'_p cannot be reducible to a free finite measure preserving action of Γ'_q . The problem is that the ambient groups are $SL_n(\mathbb{R}) \times SL_n(\mathbb{Q}_p)$ and $SL_n(\mathbb{R}) \times SL_n(\mathbb{Q}_q)$; these two groups have a common factor, making 10.1.6 of [35] difficult to apply. It may be possible to surmount this difficulty, but we avoid it entirely by using SO_F in place of SL_n .

We have the following cocycle reduction result.

Theorem 3.5. *Let p and q be distinct rational primes. Let X be a standard Borel $SO_F(\mathbb{Q}_p)$ -space with an invariant, ergodic probability measure. Let $\alpha : SO_F(\mathbb{Q}_p) \times X \rightarrow SO_F(\mathbb{Q}_q)$ be a cocycle. Then α is equivalent to a cocycle taking values in a compact subgroup of $SO_F(\mathbb{Q}_q)$.*

Proof. We will appropriately apply 3.3. We will first need a few lemmas.

The following result is standard. The proof given is intended for non-experts in the theory of algebraic groups and its first part is addressed to those with some knowledge of model theory.

Lemma 3.6. *SO_F is connected and simple (as an abstract group), thus also \mathbb{Q}_p -simple.*

Proof. For the first assertion notice that SO_F over Ω is connected iff SO_F over \mathbb{C} is connected (as Ω and \mathbb{C} are elementarily equivalent) and by [35], page 35,

3rd paragraph, SO_F over \mathbb{C} is connected iff $\mathrm{SO}_F(\mathbb{C})$ is connected in the Hausdorff topology, which is a standard fact (see Gorbatsevich et al. [13], p. 23).

For the second assertion, we quote some results from Dieudonné [4]. Our group SO_F is $O_7^+(\Omega, F)$ in Dieudonné's notation. Again in his notation, F has index $\nu \geq 1$ (see Section 9, page 17), since there is some non-zero $(x_1, \dots, x_7) \in \Omega^7$ with $x_1^2 + \dots + x_7^2 = 0$. So by Prop. 12, page 28, $O_7^+(\Omega, F) = C_7(\Omega, F)$, since every element of Ω is a square, where $C_7(\Omega, F)$ is the commutator group of $O_7^+(\Omega, F)$ (see Section 12, pp. 23-24). Then by Section 14, Th. 2, page 29, $PC_7(\Omega, F)$ is simple, where $PC_7(\Omega, F)$ is the quotient of $C_7(\Omega, F)$ by the center of $O_7^+(\Omega, F)$ (see section 13, page 25). But this center is trivial (again see Section 13), so $\mathrm{SO}_F = O_7^+(\Omega, F) = C_7(\Omega, F) = PC_7(\Omega, F)$ is simple. \dashv

Lemma 3.7. \mathbb{Q}_p -rank(SO_F) ≥ 2 .

Proof. According to the definition in [35], page 85, the \mathbb{Q}_p -rank of SO_F is the maximal dimension of a \mathbb{Q}_p -subgroup of SO_F , which can be diagonalized over \mathbb{Q}_p . It will then be enough to find a \mathbb{Q}_p -subgroup H of the diagonal matrices in GL_7 and a matrix M in $\mathrm{GL}_7(\mathbb{Q}_p)$, such that H has dimension ≥ 2 and $MHM^{-1} \subseteq \mathrm{SO}_F$.

This follows from some standard facts on quadratic forms over \mathbb{Q}_p . We use Serre [31], Ch. IV, as a reference. By Serre [31], Theorem 6 in page 36 and Definitions 7, 4' and Proposition 3' in pp. 32-33, the form $F = x_1^2 + \dots + x_7^2$ over \mathbb{Q}_p is equivalent to $F' = x_1x_2 + x_3x_4 + g(x_5, x_6, x_7)$ over \mathbb{Q}_p for some form g . Then if the matrix $M \in \mathrm{GL}_7(\mathbb{Q}_p)$ implements the equivalence and $\mathrm{SO}_{F'}$ is the \mathbb{Q} -group of matrices in GL_7 preserving F' , then $M[\mathrm{SO}_{F'}(\mathbb{Q}_p)]M^{-1} \subseteq \mathrm{SO}_F(\mathbb{Q}_p)$. Let H be the group of all diagonal 7×7 matrices $(\lambda_1, \dots, \lambda_7)$ such that $\lambda_1\lambda_2 = 1$, $\lambda_3\lambda_4 = 1$ and $\lambda_5 = \lambda_6 = \lambda_7 = 1$. Then H is a \mathbb{Q} -group, thus a \mathbb{Q}_p -group, and has dimension 2. Moreover, $H(\mathbb{Q}_p) \subseteq \mathrm{SO}_{F'}(\mathbb{Q}_p)$, so $M[H(\mathbb{Q}_p)]M^{-1} \subseteq \mathrm{SO}_F(\mathbb{Q}_p)$ and thus, by Zariski density of $H(\mathbb{Q}_p)$ in H , we see that $MHM^{-1} \subseteq \mathrm{SO}_F$. \dashv

Note also that SO_F is semisimple, *i.e.*, the (solvable) radical of SO_F , which is defined to be the maximal normal connected solvable subgroup of SO_F , is trivial, since SO_F is simple.

To be in a position to apply 3.3, we now consider the algebraic universal cover of SO_F , say G (see the paragraph following 3.1.11 in Zimmer [35] or Margulis [26], 1.4, particularly 1.4.12). This maneuver is necessary to ensure the hypothesis that the group is algebraically simply connected.

Then G is a connected semisimple \mathbb{Q}_p -group, which is also algebraically simply connected and almost \mathbb{Q}_p -simple. (Indeed, assume N is a proper normal \mathbb{Q}_p -subgroup of G , and $\pi : G \rightarrow \mathrm{SO}_F$ is a \mathbb{Q}_p -isogeny, *i.e.*, a \mathbb{Q}_p -epimorphism with finite kernel. Since SO_F is simple as an abstract group, either $\pi(N) = 1$ or $\pi(N) = \mathrm{SO}_F$. In the former case, N is a finite subgroup of G . In the latter case, N has finite index in G , which contradicts the fact that G is connected.)

Moreover, by 3.7 and Margulis [26], 1.4.6(a), \mathbb{Q}_p -rank(G) ≥ 2 (in applying 1.4.6(a) recall that semisimple \Rightarrow reductive).

Let $\pi : G \rightarrow \mathrm{SO}_F$ as above be a \mathbb{Q}_p -isogeny. Then we can use π to lift the action of $\mathrm{SO}_F(\mathbb{Q}_p)$ on X to an action of $G(\mathbb{Q}_p)$ on X defined by $g \cdot x = \pi(g) \cdot x$ (note that $\pi(G(\mathbb{Q}_p)) \subseteq \mathrm{SO}_F(\mathbb{Q}_p)$). We can also lift the cocycle $\alpha : \mathrm{SO}_F(\mathbb{Q}_p) \times X \rightarrow \mathrm{SO}_F(\mathbb{Q}_q)$ to the cocycle $\alpha_1 : G(\mathbb{Q}_p) \times X \rightarrow \mathrm{SO}_F(\mathbb{Q}_q)$ given by $\alpha_1(g, x) = \alpha(\pi(g), x)$.

We claim that it suffices to show that α_1 is equivalent to a cocycle taking values in a compact subgroup, say K_1 , of $\mathrm{SO}_F(\mathbb{Q}_q)$. To see this argue as follows: In

general, by Zimmer [35], page 34, paragraph 6, when H_1, H_2 are k -groups and $\varphi : H_1 \rightarrow H_2$ a k -morphism, then $\varphi(H_1)$ is a k -group and $\varphi(H_1(k))$ is a subgroup of finite index in $(\varphi(H_1))(k)$. Thus $\pi(G(\mathbb{Q}_p))$ has finite index in $\text{SO}_F(\mathbb{Q}_p)$ (actually in this particular case it turns out that $\pi(G(\mathbb{Q}_p)) = \text{SO}_F(\mathbb{Q}_p)$ – see Margulis [26], page 42, line 7). So by 2.7, α is equivalent to a cocycle taking values in a compact subgroup of $\text{SO}_F(\mathbb{Q}_p)$.

Now G satisfies all the hypotheses of 3.3, but unfortunately we cannot take $H = \text{SO}_F(\mathbb{Q}_q)$ and $\alpha = \alpha_1$ in 3.3. One problem is that α_1 might be equivalent to a cocycle with range contained in a proper subgroup of the form $L(\mathbb{Q}_q)$ for a \mathbb{Q}_q -subgroup $L \subseteq H$. So we proceed as follows:

Call a cocycle $\beta : G(\mathbb{Q}_p) \times Y \rightarrow \text{SO}_F(\mathbb{Q}_q)$ close to $\alpha : G(\mathbb{Q}_p) \times X \rightarrow \text{SO}_F(\mathbb{Q}_q)$ if Y is a finite extension of X and if β is equivalent to the lift of α to Y . (Note that this notion is not symmetric.)

Now let \mathcal{C} denote the collection of all \mathbb{Q}_q -subgroups L of SO_F such that there is a cocycle $\beta : G(\mathbb{Q}_p) \times Y \rightarrow \text{SO}_F(\mathbb{Q}_q)$, β close to α_1 , with $\beta(G(\mathbb{Q}_p) \times Y) \subseteq L(\mathbb{Q}_q)$.

By the descending chain condition for algebraic groups, there is a minimal element H in \mathcal{C} . Then choose a cocycle $\beta : G(\mathbb{Q}_p) \times Y \rightarrow \text{SO}_F(\mathbb{Q}_q)$ which is close to α_1 such that $\beta(G(\mathbb{Q}_p) \times Y) \subseteq H(\mathbb{Q}_q)$. By 2.6, in order to show that α_1 is equivalent to a cocycle taking values in a compact subgroup of $\text{SO}_F(\mathbb{Q}_q)$, it is enough to show that the same holds for β .

First we will use the minimality of H to deduce that H must be connected, *i.e.*, that $H = H^0$, where H^0 is the Zariski connected component of the identity of H . Now H^0 is a normal \mathbb{Q}_q -subgroup of H of finite index, so $H^0(\mathbb{Q}_q)$ is a closed normal subgroup of finite index in $H(\mathbb{Q}_q)$. Then, by 2.5, there exists a finite extension Z of Y such that if $\hat{\beta}$ is the lift of β to Z , then there exists a cocycle $\gamma \sim \hat{\beta}$ such that γ takes values in $H^0(\mathbb{Q}_q)$. But γ is close to α_1 ; and so, if H^0 is a proper subgroup of H , then this violates the minimality of H .

Now H might not be semisimple, so let R be the (solvable) radical of H and $S = H/R, \tau : H \rightarrow S$ the canonical epimorphism. Then S is a connected semisimple \mathbb{Q}_q -group and τ is a \mathbb{Q}_q -morphism. By Margulis [26], first two paragraphs of 0.24 in pages 20, 21, there are normal connected almost \mathbb{Q}_q -simple, \mathbb{Q}_q -subgroups of S , say G_1, \dots, G_k , so that the multiplication operation from $G_1 \times \dots \times G_k$ into S is an isogeny (*i.e.*, an epimorphism with finite kernel), say ρ . Let Z_i be the center of G_i and let $S_i = G_i/Z_i$. Then since G_i is connected and almost \mathbb{Q}_q -simple, every normal \mathbb{Q}_q -subgroup is finite, so contained in the center (see Margulis [26], 0.18), so S_i is a connected \mathbb{Q}_q -simple, \mathbb{Q}_q -group, and since the kernel of ρ is finite it must be contained in the center of $G_1 \times \dots \times G_k$, since this group is connected. So the kernel of ρ is contained in the product $Z_1 \times \dots \times Z_k$ and there is a canonical \mathbb{Q}_p -epimorphism $\tau_* : S \rightarrow S_1 \times \dots \times S_k$. Denote by τ_i the composition of τ_* with the projection of $S_1 \times \dots \times S_k$ onto S_i .

Let $\sigma_i = \tau_i \circ \tau$ and $\beta_i = \sigma_i \circ \beta : G(\mathbb{Q}_p) \times Y \rightarrow S_i(\mathbb{Q}_q)$. Now all the hypotheses of 3.3 are clearly satisfied for $H = S_i, \alpha = \beta_i$. Recall that $\sigma_i(H(\mathbb{Q}_q))$ has finite index in $S_i(\mathbb{Q}_q)$, as σ_i maps H onto S_i . View $\beta_i : G(\mathbb{Q}_p) \times Y \rightarrow \sigma_i(H(\mathbb{Q}_q))$ as a cocycle into $\sigma_i(H(\mathbb{Q}_q))$. Suppose that, viewed as a cocycle into $S_i(\mathbb{Q}_q)$, β_i is equivalent to a cocycle taking values in $L(\mathbb{Q}_q)$, where L is a proper \mathbb{Q}_q -subgroup of S_i . Apply 2.4 to get that, as a cocycle into $\sigma_i(H(\mathbb{Q}_q))$, β_i is equivalent to a cocycle with values in $L_i = \sigma_i(H(\mathbb{Q}_q)) \cap hL(\mathbb{Q}_q)h^{-1}$, for some $h \in S_i(\mathbb{Q}_q)$. But then, by 2.1, there is a β_i -invariant Borel function $f : X \rightarrow \sigma_i(H(\mathbb{Q}_q))/L_i$, so clearly there

is a β -invariant Borel function $f' : X \rightarrow H(\mathbb{Q}_q)/\sigma_i^{-1}(L_i)$, and so β is equivalent to a cocycle taking values in $\sigma_i^{-1}(L_i) = \sigma_i^{-1}(hL(\mathbb{Q}_q)h^{-1})$. But $L' = hLh^{-1}$ is a proper k -subgroup of S_i , and so $\sigma_i^{-1}(L')$ is a proper k -subgroup of H , while $\sigma_i^{-1}(hL(\mathbb{Q}_p)h^{-1}) = \sigma_i^{-1}(L'(\mathbb{Q}_p)) = \sigma_i^{-1}(L')(\mathbb{Q}_p)$, contradicting the minimality of H .

We conclude, for all i , that β_i is equivalent to a cocycle with values in a compact subgroup K_i of $S_i(\mathbb{Q}_p)$. Thus $\tau_* \circ \tau \circ \beta$ is equivalent to a cocycle taking values in $K_1 \times \dots \times K_n$. Then, as in the preceding argument, an application of 2.4 shows that β is equivalent to cocycle γ taking values into $A = \tau^{-1}(\tau_*(K))$, where K is a compact subgroup of $S_1(\mathbb{Q}_q) \times \dots \times S_n(\mathbb{Q}_q)$. In fact, K is a conjugate of $K_1 \times \dots \times K_n$. Now $\tau_*^{-1}(K)$ is an extension of a compact group by a finite group, so it is amenable and $\tau^{-1}(\tau_*(K))$ is an extension of an amenable group by a solvable group (*i.e.*, R), so A is amenable (see [35], 4.6 and 4.1.2).

By Margulis [26], III 5.3 in page 130, $G(\mathbb{Q}_p)$ is Kazhdan. So, by 3.1, the cocycle γ is equivalent to a cocycle taking values in a compact subgroup of A , and so β is equivalent to a cocycle taking values in a compact subgroup of $H(\mathbb{Q}_q)$ and the proof is complete. -

The following cocycle reduction result will be needed in Section 6.

Theorem 3.8. *Let $n \geq 3$ and let X be a standard Borel $SL_n(\mathbb{R})$ -space with an invariant, ergodic probability Borel measure. Let $m < n$ and let $H^m = SL_m^{\pm 1} / \{I_m, -I_m\}$ be the group of all $m \times m$ matrices with determinant ± 1 , modulo the subgroup $\{I_m, -I_m\}$, where $I_m =$ the identity $m \times m$ matrix. Let p be a prime or ∞ and let $\mathbb{Q}_\infty = \mathbb{R}$. If $\alpha : SL_n(\mathbb{R}) \times X \rightarrow H^m(\mathbb{Q}_p)$ is a cocycle, then α is equivalent to a cocycle taking values in a compact subgroup of $H^m(\mathbb{Q}_p)$.*

Proof. The argument is similar to that of 3.5, using 3.4 instead of 3.3, so we will only indicate the changes.

First since $SL_n(\mathbb{C})$ is connected (see, e.g., Gorbatsevich et al. [13], page 22, Prop. 4.4), so is SL_n . Also $SL_n(\mathbb{R})$ is connected (same reference). Every normal subgroup of $SL_n(K)$, K any field of characteristic 0, is contained in its center which is finite (see Rotman [29], 8.9 and 9.4.6). In particular, SL_n is semisimple. Finally, \mathbb{R} -rank($SL_n(\mathbb{R})$) = $n - 1$ (see Zimmer [35], page 85, first paragraph), so since $n \geq 3$ all the hypotheses about $G = SL_n(\mathbb{R})$ in 3.4 are satisfied.

Then, by repeating the argument in 3.5, we end up with a cocycle of the action of $SL_n(\mathbb{R})$ on X into a group of the form $S_i(k)$, where $k = \mathbb{Q}_p$ or \mathbb{R} , and S_i is a connected k -simple, k -group for which there is a k -epimorphism from H onto S_i , where H is a connected k -subgroup of H^m . We have to conclude that this cocycle is equivalent to one with values in some compact subgroup of $S_i(k)$. Applying 3.4, this is the case unless $k = \mathbb{R}$ and there is a non-trivial \mathbb{R} -morphism ρ from SL_n into S_i . But this is impossible, since the dimension of S_i is less than or equal to that of H^m , which is $m^2 - 1$, and thus strictly smaller to that of SL_n , which is $n^2 - 1$, and the kernel of ρ is finite (see Humphreys [17], 7.4).

Finally, we need that $SL_n(\mathbb{R})$, for $n \geq 3$, is Kazhdan (see de la Harpe and Valette [3], 2.a.4, page 21). -

We conclude this section with the proof of a well-known fact that we will need in the next section.

Proposition 3.9. *Let $\mathbb{Z}[\frac{1}{p}]$ be the ring of rationals whose denominators (in reduced form) are powers of p . Then $\text{SO}_F(\mathbb{Z}[\frac{1}{p}])$ is a Kazhdan group.*

Proof. We will need the following standard result:

Lemma 3.10. *Let G, H be lsc groups, $\Gamma \subseteq G$ a lattice in G and $f : G \rightarrow H$ a continuous epimorphism with compact kernel. Then $f(\Gamma)$ is a lattice in H .*

Proof. Let $K = \text{kernel of } f$. Then $K\Gamma = \Gamma K$ is a closed subgroup of G and let ϕ be the canonical map from G/Γ onto $G/K\Gamma$ given by $\phi(h\Gamma) = hK\Gamma$. Then for the canonical actions of G on G/Γ and $G/K\Gamma$ we clearly have $\phi(g \cdot x) = g \cdot \phi(x)$, so if μ is a G -invariant measure on G/Γ , then $\nu = \phi_*(\mu)$ is a G -invariant probability Borel measure on $G/K\Gamma$. Now let $\psi : G/K\Gamma \rightarrow H/f(\Gamma)$ be the canonical bijection given by

$$\psi(hK\Gamma) = f(h)f(\Gamma).$$

Then, as $\psi(g \cdot x) = f(g) \cdot \psi(x)$, it follows that $\psi_*(\nu)$ is H -invariant. Finally it is easy to check that $f(\Gamma)$ is discrete in H . ⊣

Now take $\Gamma = \text{SO}_F(\mathbb{Z}[\frac{1}{p}])$ and let $K = \text{SO}_F(\mathbb{R}), H = \text{SO}_F(\mathbb{Q}_p)$. Then K is compact. Define $\delta : \Gamma \rightarrow K \times H$ by $\delta(x) = (x, x)$. Then by Zimmer [35], 10.1.1 (with $G = \text{SO}_F, S = \{p\}$ in his notation), $\delta(\Gamma)$ is a lattice in $K \times H$, so if $\rho : K \times H \rightarrow H$ is the projection, by the previous lemma, $\rho(\delta(\Gamma)) = \Gamma$ is a lattice in $\text{SO}_F(\mathbb{Q}_p)$. By Margulis [26], III 5.3 in page 130, $\text{SO}_F(\mathbb{Q}_p)$ is a Kazhdan group and by Zimmer [35], 7.4.3, every lattice in a Kazhdan group is also a Kazhdan group, so $\text{SO}_F(\mathbb{Z}[\frac{1}{p}])$ is a Kazhdan group. ⊣

4. EMBEDDING BOREL SETS UNDER INCLUSION

Our goal in this section is to prove the first main theorem of this paper.

Theorem 4.1. *The partial ordering of Borel sets under inclusion can be embedded in the partial (pre)order of countable Borel equivalence relations under Borel reducibility. More precisely, there is a map $A \mapsto E_A$ assigning to each Borel subset of $2^{\mathbb{N}}$, A , a countable Borel equivalence relation, E_A , so that*

$$A_1 \subseteq A_2 \Leftrightarrow E_{A_1} \leq_B E_{A_2}.$$

The proof of 4.1 is based on the following result.

Theorem 4.2. *There is a map $x \mapsto E_x$ assigning to each $x \in 2^{\mathbb{N}}$ a countable Borel equivalence relation E_x on $2^{\mathbb{N}}$ such that:*

- (i) E_x admits an ergodic, non-atomic, invariant measure μ_x .
- (ii) If $x_0 \neq x_1$ and $f : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ is a Borel function such that $yE_{x_0}z \Rightarrow f(y)E_{x_1}f(z)$, then there is a Borel E_{x_0} -invariant set M of μ_{x_0} -measure 1 such that $f(M)$ is contained in a single E_{x_1} -class. In particular, for any Borel set N of μ_{x_0} -measure 1, $E_{x_0}|N \not\leq_B E_{x_1}$.
- (iii) The assignment $x \mapsto E_x$ is “Borel”, in the sense that $E \subseteq (2^{\mathbb{N}})^3$ below is Borel:

$$(x, y, z) \in E \Leftrightarrow yE_xz.$$

Moreover $x \mapsto \mu_x$ is Borel.

Granting 4.2 we can prove 4.1 as follows:

To each Borel set $A \subseteq 2^{\mathbb{N}}$ assign the following equivalence relation:

$$E_A = \bigsqcup_{x \in A} E_x,$$

that is, the direct sum of the E_x with $x \in A$. This is precisely defined as follows: E_A is an equivalence relation on $A \times 2^{\mathbb{N}}$, and

$$(x, y)E_A(x', z) \Leftrightarrow x = x' \ \& \ yE_x z.$$

Then E_A is Borel and, since there is a Borel bijection of $A \times 2^{\mathbb{N}}$ with $2^{\mathbb{N}}$, we may, if we wish, think of E_A as being an equivalence relation on $2^{\mathbb{N}}$. It is clear that if $A_1 \subseteq A_2$, then $E_{A_1} \leq_B E_{A_2}$ via the identity function. Now assume that $A_1 \not\subseteq A_2$ and let $x \in A_1 \setminus A_2$. Then clearly $E_x \leq_B E_{A_1}$ (via $y \mapsto (x, y)$). On the other hand, we claim that $E_x \not\leq_B E_{A_2}$, so $E_{A_1} \not\leq_B E_{A_2}$. Because, otherwise, there is a Borel function $f : 2^{\mathbb{N}} \rightarrow A_2 \times 2^{\mathbb{N}}$ with $yE_x z \Leftrightarrow f(y)E_{A_2} f(z)$, and so, in particular, $yE_x z \Rightarrow \pi(f(y)) = \pi(f(z))$, where $\pi(a, b) = a$. By ergodicity, there is a Borel set M of μ_x -measure 1 with $\pi(f(y))$ constant for $y \in M$, say with value $\pi(f(y)) = x_0$. Then $y \mapsto \rho(f(y))$, where $\rho(a, b) = b$, is a Borel reduction of $E_x|_M$ into E_{x_0} , which contradicts 4.2(ii), as $x \neq x_0$.

We will devote the rest of this section to the proof of Theorem 4.2.

For any infinite countable group Γ , consider the compact Polish space 2^Γ of all functions from Γ into $2 = \{0, 1\}$ with the product topology (so that it is homeomorphic to $2^{\mathbb{N}}$). Γ acts on 2^Γ by (left) shift: $g \cdot x(h) = x(g^{-1}h)$. We denote by $E(\Gamma, 2)$ the associated equivalence relation

$$xE(\Gamma, 2)y \Leftrightarrow \exists g(g \cdot x = y).$$

We also let $F(\Gamma, 2)$ be the restriction of $E(\Gamma, 2)$ to the free part of the action, *i.e.*,

$$F(\Gamma, 2) = E(\Gamma, 2)|_{\text{Fr}_\Gamma},$$

where $\text{Fr}_\Gamma = \{x \in 2^\Gamma : \forall g \neq 1(g \cdot x \neq x)\}$. We also denote by μ_Γ the product measure on 2^Γ (where each bit in $\{0, 1\}$ has measure $1/2$). It is clearly invariant under the shift action and it is non-atomic. The next lemma is quite standard.

Lemma 4.3. $\mu_\Gamma(\text{Fr}_\Gamma) = 1$.

Proof. It is clearly enough to show that for each $g \neq 1$, $X_g = \{x \in 2^\Gamma : \forall h(x(gh) = x(h))\}$ has μ_Γ -measure 0. It is clear that if $x \in X_g$, then x is constant on each right coset $\langle g \rangle h$, $h \in \Gamma$, of the subgroup $\langle g \rangle$ generated by g . Since either $\langle g \rangle$ is infinite or there are infinitely many cosets, this implies that $\mu_\Gamma(X_g) = 0$. \dashv

If Δ is a subgroup of Γ , then also Δ acts on 2^Γ by left shift and we denote by $E(\Delta, \Gamma, 2)$ the corresponding equivalence relation, so that

$$xE(\Delta, \Gamma, 2)y \Leftrightarrow \exists g \in \Delta(g \cdot x = y).$$

The following lemma is also standard. In fact, the action of Γ on 2^Γ turns out to be mixing, a property which is inherited by infinite subgroups of Γ , and which implies ergodicity. For the sake of completeness, we include a proof which does not depend on understanding the mixing property.

Lemma 4.4. *For any infinite subgroup Δ of Γ , μ_Γ is $E(\Delta, \Gamma, 2)$ -ergodic.*

Proof. Suppose $X \subseteq 2^\Gamma$ is Borel and $E(\Delta, \Gamma, 2)$ -invariant. Enumerate $\Gamma = \{\gamma_0, \gamma_1, \dots, \gamma_n, \dots\}$. Then for each n , choose $\delta_n \in \Delta$ such that δ_n is not one of $\gamma_i \gamma_j^{-1}$, for $i, j < n$. Then $\delta_n \{\gamma_0, \dots, \gamma_{n-1}\} \cap \{\gamma_0, \dots, \gamma_{n-1}\} = \emptyset$. Consider the Boolean algebra \mathcal{A} of clopen subsets of 2^Γ . Denote by MALG_{μ_Γ} the measure algebra of μ_Γ , i.e., the Boolean algebra of Borel subsets of 2^Γ modulo sets of μ_Γ -measure 0. Then \mathcal{A} is dense in MALG_{μ_Γ} (see, e.g., Kechris [21], 17.43), so for each n there is $A_n \in \mathcal{A}$ with $\mu_\Gamma(X + A_n) \rightarrow 0$, where $+$ denotes here the symmetric difference of two sets. Now each A_n is a finite union of sets of the form

$$N_s = \{x \in 2^\Gamma : x|F = s\},$$

where $s : F \rightarrow 2$, for some finite subset $F \subseteq \Gamma$. Call F the *support* of N_s and if $A_n = N_{s_1} \cup \dots \cup N_{s_k}$, with the support of N_{s_i} equal to F_i , call $F_1 \cup \dots \cup F_k$ the support of A_n . Say it is included in $\{\gamma_0, \dots, \gamma_{k_n-1}\}$. Then it is clear that if $\epsilon_n = \delta_{k_n}^{-1}$, A_n and $\epsilon_n \cdot A_n$ are independent for μ_Γ , so $\mu_\Gamma(A_n \cap \epsilon_n \cdot A_n) = \mu_\Gamma(A_n)\mu_\Gamma(\epsilon_n \cdot A_n) = \mu_\Gamma(A_n)^2$ by the invariance of μ_Γ . Also $\mu_\Gamma(A_n + X) = \mu_\Gamma(\epsilon_n \cdot (A_n + X)) = \mu_\Gamma(\epsilon_n \cdot A_n + X)$, as $\epsilon_n \cdot X = X$. So $A_n \rightarrow X$ and $\epsilon_n \cdot A_n \rightarrow X$ in the measure algebra, so, as intersection is continuous (see Kechris [21], 17.43), $A_n \cap \epsilon_n \cdot A_n \rightarrow X$ and thus $\mu_\Gamma(A_n \cap \epsilon_n \cdot A_n) = \mu_\Gamma(A_n)^2 \rightarrow \mu_\Gamma(X)$. But also $\mu_\Gamma(A_n) \rightarrow \mu_\Gamma(X)$, so $\mu_\Gamma(X)^2 = \mu_\Gamma(X)$, i.e., $\mu_\Gamma(X) = 0$ or 1. ◻

Recall that SO_F is the \mathbb{Q} -group of all 7×7 matrices A with $\det(A) = 1, AA^t = I$. For each non-empty subset S of primes, let $\mathbb{Z}[S^{-1}]$ be the ring of rationals whose denominators (in reduced form) have prime factors in S , and let

$$\Gamma_S = \text{SO}_F(\mathbb{Z}[S^{-1}])$$

be the set of all 7×7 matrices in SO_F with coefficients in $\mathbb{Z}[S^{-1}]$. Let $E_S = F(\Gamma_S, 2)$ and $X_S = \text{Fr}_{\Gamma_S}, \mu_S = \mu_{\Gamma_S}$. Then we have the following lemma:

Lemma 4.5. *If $S \not\subseteq T$, and $f : X_S \rightarrow X_T$ is Borel such that $x E_S y \Rightarrow f(x) E_T f(y)$, then there is a Borel E_S -invariant set M with $\mu_S(M) = 1$ such that f maps M into a single E_T -class.*

Granting this, we can prove 4.2 as follows: Let π be a Borel function from $2^\mathbb{N}$ into the power set of the primes (which we identify with $2^P, P = \text{the set of primes}$) so that each $\pi(x)$ is infinite and

$$x \neq y \Rightarrow \pi(x) \cap \pi(y) \text{ is finite,}$$

i.e., $\{\pi(x)\}_{x \in 2^\mathbb{N}}$ is an almost disjoint family of subsets of primes. (The existence of such a Borel function π is a standard fact and here is a quick proof: It is clearly enough to replace P by \mathbb{N} . Fix a bijection $\langle \cdot \rangle : 2^{<\mathbb{N}} \rightarrow \mathbb{N}$ of the set of finite sequences $\langle s_0, \dots, s_{n-1} \rangle$ from \mathbb{N} with \mathbb{N} , and for $x \in 2^\mathbb{N}$ let $\pi(x) = \{\langle x(0), \dots, x(n-1) \rangle : n \in \mathbb{N}\} \subseteq \mathbb{N}$.) For $x \in 2^\mathbb{N}$, put

$$E_x = E_{\pi(x)}$$

and $\mu_x = \mu_{\pi(x)}$. Then (i)-(iii) of 4.2 are clear. (Literally E_x is not an equivalence relation on $2^\mathbb{N}$ but it can be considered as such by fixing Borel bijections of each $X_{\pi(x)}$ with $2^\mathbb{N}$, uniformly in x .)

Proof of Lemma 4.5. Fix $p \in S \setminus T$, and consider $\Gamma_p = \Gamma_{\{p\}} \subseteq \Gamma_S$, i.e., $\Gamma_p = \text{SO}_F(\mathbb{Z}[\frac{1}{p}])$. By 4.4 the measure μ_S is ergodic and invariant for the action of Γ_p on X_S . We can now define the following strict Borel cocycle $\alpha : \Gamma_p \times X_S \rightarrow \Gamma_T$:

$$\begin{aligned} \alpha(g, x) &= \text{the unique } h \in \Gamma_T \text{ such that} \\ h \cdot f(x) &= f(g \cdot x), \end{aligned}$$

which exists since $x E_S y \Rightarrow f(x) E_T f(y)$, and the action of Γ_T is free.

Since, by 3.9, Γ_p is a Kazhdan group, it follows from 3.2 that $\alpha \sim \beta$, where $\beta(\Gamma_p \times X_S) \subseteq \Lambda_T$, with $\Lambda_T \subseteq \Gamma_T$ a finitely generated subgroup of Γ_T . So there is a finite $T_0 \subseteq T$ such that $\Lambda_T \subseteq \Gamma_{T_0}$. Say $T_0 = \{p_1, \dots, p_n\}$. By Zimmer [35], B. 9 (on page 200), we can also assume that β is a strict cocycle.

Now $\Gamma_{T_0} = \text{SO}_F(\mathbb{Z}[T_0^{-1}])$ is a discrete subgroup of $\text{SO}_F(\mathbb{R}) \times \text{SO}_F(\mathbb{Q}_{p_1}) \times \dots \times \text{SO}_F(\mathbb{Q}_{p_n})$ (as $\mathbb{Z}[T_0^{-1}]$ is a discrete subring of $\mathbb{R} \times \mathbb{Q}_{p_1} \times \dots \times \mathbb{Q}_{p_n}$) and thus of $\text{SO}_F(\mathbb{Q}_{p_1}) \times \dots \times \text{SO}_F(\mathbb{Q}_{p_n})$, as $\text{SO}_F(\mathbb{R})$ is compact. (Here an element $g \in \Gamma_{T_0}$ is identified with (g, \dots, g) in this product and similarly for elements of $\mathbb{Z}[T_0^{-1}]$.) Also $\Gamma_p = \text{SO}_F(\mathbb{Z}[\frac{1}{p}])$ is a lattice in $\text{SO}_F(\mathbb{R}) \times \text{SO}_F(\mathbb{Q}_p)$, by Zimmer [35], 10.1.1 (since SO_F is a connected semisimple \mathbb{Q} -group), and so by 3.10, it is a lattice in $\text{SO}_F(\mathbb{Q}_p)$.

Consider the action of Γ_p on (X_S, μ_S) and the strict cocycle $\beta : \Gamma_p \times X_S \rightarrow \Gamma_{T_0}$. Using Section 2 (paragraph following 2.2), consider the induced Borel action of $\text{SO}_F(\mathbb{Q}_p)$ on Y (which also has an invariant, ergodic measure) and the induced strict cocycle $\hat{\beta} : \text{SO}_F(\mathbb{Q}_p) \times Y \rightarrow \Gamma_{T_0}$. Viewing $\hat{\beta}$ as a cocycle $\hat{\beta} : \text{SO}_F(\mathbb{Q}_p) \times Y \rightarrow \text{SO}_F(\mathbb{Q}_{p_1}) \times \dots \times \text{SO}_F(\mathbb{Q}_{p_n})$ and letting π_i be the projection of $\text{SO}_F(\mathbb{Q}_{p_1}) \times \dots \times \text{SO}_F(\mathbb{Q}_{p_n})$ to $\text{SO}_F(\mathbb{Q}_{p_i}), i = 1, \dots, n$, consider the cocycle $\hat{\beta}_i = \pi_i \circ \hat{\beta} : \text{SO}_F(\mathbb{Q}_p) \times Y \rightarrow \text{SO}_F(\mathbb{Q}_{p_i})$. By 3.5, $\hat{\beta}_i \sim \gamma_i$, where γ_i is a Borel cocycle with $\gamma_i(\text{SO}_F(\mathbb{Q}_p) \times Y) \subseteq K_i, K_i$ a compact subgroup of $\text{SO}_F(\mathbb{Q}_{p_i})$. Set $K = K_1 \times \dots \times K_n$. Then $\hat{\beta} \sim \gamma$ where $\gamma : \text{SO}_F(\mathbb{Q}_p) \times Y \rightarrow \text{SO}_F(\mathbb{Q}_{p_1}) \times \dots \times \text{SO}_F(\mathbb{Q}_{p_n})$ is a Borel cocycle with $\gamma(\text{SO}_F(\mathbb{Q}_p) \times Y) \subseteq K$. Then by 2.4, now viewing $\hat{\beta}$ as a cocycle into Γ_{T_0} , we have that $\hat{\beta} \sim \beta'$ where $\beta' : \text{SO}_F(\mathbb{Q}_p) \times Y \rightarrow \Gamma_{T_0}$ is a Borel cocycle with $\beta'(\text{SO}_F(\mathbb{Q}_p) \times Y) \subseteq \Delta_0, \Delta_0$ a finite subgroup of Γ_{T_0} . Now using 2.3, this shows that $\beta \sim \beta_0$, where $\beta_0 : \Gamma_p \times X_S \rightarrow \Gamma_{T_0}$ is a Borel cocycle with $\beta_0(\Gamma_p \times X_S) \subseteq \Delta_0$ and so $\alpha \sim \beta_0$. Thus there is a Borel function $A : X_S \rightarrow \Gamma_T$ such that for all $g \in \Gamma_p, \alpha(g, x) = A(g \cdot x)\beta_0(g, x)A(x)^{-1}, \mu_S$ -a.e.(x). Since Γ_T is countable, find some Borel set $N_0 \subseteq X_S$ of positive μ_S -measure, such that A is constant, say with value A_0 , on N_0 , and, since Γ_p is countable, we can assume that $\alpha(g, x) = A(g \cdot x)\beta_0(g, x)A(x)^{-1}$, for all $g \in \Gamma_p, x \in N_0$. Then if $x, g \cdot x \in N_0$ for some $g \in \Gamma_p, \alpha(g, x) = A_0\beta_0(g, x)A_0^{-1} \in A_0\Delta_0A_0^{-1}$. Since $\alpha(g, x) \cdot f(x) = f(g \cdot x)$ it follows that as g varies over Γ_p so that $g \cdot x \in N_0, f(g \cdot x)$ takes only finitely many values. Let $M_0 = \Gamma_p \cdot N_0$ be the Γ_p -saturation of N_0 . Then M_0 has μ_S -measure 1 by ergodicity. Moreover, the map on M_0 defined by $F(x) = f((\Gamma_p \cdot x) \cap N_0)$ is Γ_p -invariant and assigns in a Borel way to x a finite subset of X_T , so, by ergodicity again, it is fixed on a Borel set $M_1 \subseteq M_0$ of μ_S -measure 1. Note that if $x_0 \in M_0$, then $F(x)$ is a finite subset of the E_T -class which f assigns to the E_S -class containing $\Gamma_p \cdot x$. Hence f maps M_1 into a single E_T -class. Finally, let $M = \Gamma_S \cdot M_1$. Then clearly M is a Borel E_S -invariant subset of μ_S -measure 1 which is mapped into a single E_T -class. -1

5. THE DESCRIPTIVE COMPLEXITY OF BOREL REDUCIBILITY

Let us first fix a parametrization of Borel relations on $\mathbb{N}^{\mathbb{N}}$. This consists of a set $D \subseteq 2^{\mathbb{N}}$ and two sets $P, S \subseteq (\mathbb{N}^{\mathbb{N}})^3$ such that

- (i) D is Π_1^1 , P is Π_1^1 and S is Σ_1^1 .
- (ii) For $d \in D$, $P_d = S_d$ (where $P_d = \{(x, y) : (d, x, y) \in P\}$). Denote by D_d the set $P_d (= S_d)$, for $d \in D$.
- (iii) $\{D_d : d \in 2^{\mathbb{N}}\} = \{B \subseteq (\mathbb{N}^{\mathbb{N}})^2 : B \text{ is Borel}\}$.
- (iv) If $A \subseteq X \times (\mathbb{N}^{\mathbb{N}})^2$, X a Polish space, is Borel, there is a Borel function $p : X \rightarrow 2^{\mathbb{N}}$, such that $A_x = D_{p(x)}$, for all $x \in X$ (see Kechris [21], 35.5).

It is straightforward, using Kechris [21], 29.19, that

$$\mathcal{E} = \{d \in D : D_d \text{ is a countable equivalence relation}\}$$

is Π_1^1 and it is also easy to check that

$$\mathcal{E}_{\leq} = \{(d, e) \in D^2 : d, e \in \mathcal{E} \ \& \ D_d \leq_B D_e\},$$

$$\mathcal{E}_{\sim} = \{(d, e) \in D^2 : d, e \in \mathcal{E} \ \& \ D_d \sim_B D_e\}$$

are Σ_2^1 .

Recall that a set A in a standard Borel space X is called Σ_2^1 -complete if it is Σ_2^1 and for any Σ_2^1 set $B \subseteq Y$, Y a standard Borel space, there is a Borel function $f : Y \rightarrow X$ such that $B = f^{-1}(A)$. The following therefore computes the exact complexity of $\mathcal{E}_{\leq}, \mathcal{E}_{\sim}$.

Theorem 5.1. *The sets $\mathcal{E}_{\leq}, \mathcal{E}_{\sim}$ are Σ_2^1 -complete. In other words, the notions of Borel reducibility and bi-reducibility of countable Borel equivalence relations are Σ_2^1 -complete.*

Proof. We will prove in fact the stronger statement that there is a fixed countable Borel equivalence relation E^* such that the countable Borel equivalence relations to which E^* Borel reduces, resp. which are Borel bi-reducible with E^* , are both Σ_2^1 -complete or more precisely that

$$\mathcal{E}_{\leq}^* = \{d \in D : d \in \mathcal{E} \ \& \ E^* \leq_B D_d\},$$

$$\mathcal{E}_{\sim}^* = \{d \in D : d \in \mathcal{E} \ \& \ E^* \sim_B D_d\}$$

are Σ_2^1 -complete. The relation E^* is simply defined as follows: For any countable Borel equivalence relation E on X let cE be the sum of continuum many copies of E . We view cE as defined on $\mathbb{N}^{\mathbb{N}} \times X$ by

$$(a, x)cE(b, y) \Leftrightarrow a = b \ \& \ xEy.$$

By trivially modifying 4.2, fix a map $x \mapsto E_x$ from $\mathbb{N}^{\mathbb{N}}$ into Borel equivalence relations on $\mathbb{N}^{\mathbb{N}}$, satisfying (i)-(iii) of 4.2 (with $\mathbb{N}^{\mathbb{N}}$ replacing $2^{\mathbb{N}}$), and put

$$E_x^* = cE_x$$

and

$$E^* = \bigsqcup_x E_x^*$$

(where we can of course view all of these as living on $\mathbb{N}^{\mathbb{N}}$).

Consider a tree T on $\mathbb{N} \times \mathbb{N}$ (see Kechris [21], 2.1). Each such T defines a closed subset $[T] \subseteq (\mathbb{N}^{\mathbb{N}})^2$ given by

$$[T] = \{(x, y) \in (\mathbb{N}^{\mathbb{N}})^2 : \forall n(x|n, y|n) \in T\}.$$

Associate with T the following countable Borel equivalence relation, where Δ is the equality relation on $\mathbb{N}^{\mathbb{N}}$:

$$E_T = \bigsqcup_{(x,y) \in [T]} E_x^* \sqcup \bigsqcup_{(x,y) \notin [T]} \Delta.$$

Let us say that a closed set $F \subseteq (\mathbb{N}^{\mathbb{N}})^2$ admits a *full Borel uniformization* if there is a Borel function $f : \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that for every x , $(x, f(x)) \in F$. The main observation that we will use in the calculation of the complexity of \mathcal{E}_{\leq}^* , \mathcal{E}_{\sim}^* is now the following:

Lemma 5.2. *For each tree T we have*

$$\begin{aligned} E^* \leq_B E_T &\Leftrightarrow E^* \sim_B E_T \\ &\Leftrightarrow [T] \text{ admits a full Borel uniformization.} \end{aligned}$$

Proof. Since $E_T \leq_B \bigsqcup_{(x,y) \in (\mathbb{N}^{\mathbb{N}})^2} E_x^* \leq_B \bigsqcup_x (cE_x^*) \cong_B \bigsqcup_x c(cE_x) \cong_B \bigsqcup_x cE_x = E^*$, where \cong_B denotes Borel isomorphism, clearly

$$E^* \leq_B E_T \Leftrightarrow E^* \sim_B E_T.$$

Now assume that $[T]$ admits a full Borel uniformization, say f . View E_x^* as being defined on $\mathbb{N}^{\mathbb{N}}$ and E^* as defined on $\mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$ by

$$(x, y)E^*(x', z) \Leftrightarrow x = x' \ \& \ yE_x^*z,$$

and let $g : \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \rightarrow (\mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}) \times \mathbb{N}^{\mathbb{N}}$ be defined by

$$g(x, y) = (x, f(x), y).$$

Then g is a Borel reduction of E^* into E_T .

Conversely, let the Borel function $g : \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \rightarrow (\mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}) \times \mathbb{N}^{\mathbb{N}}$ reduce E^* into E_T . Then the function $g_x(y) = g(x, y)$ is a Borel reduction of E_x^* into E_T . We clearly have a Borel function ρ reducing each E_x into E_x^* (ρ is defined independently of x), so $g_x \circ \rho$ is a Borel reduction of E_x into E_T , thus in particular

$$yE_xz \Rightarrow \pi_1((g_x \circ \rho)(y)) = \pi_1((g_x \circ \rho)(z))$$

where $\pi_1(u, v, w) = (u, v)$. Since μ_x (as in 4.2(i)) is ergodic, this implies that there is a set M_x of μ_x -measure 1, so that $\pi_1 \circ g_x \circ \rho$ is constant on M_x , say with value $(u(x), f(x))$. But then, if $\pi_2(u, v, w) = w$, $\pi_2 \circ g_x \circ \rho$ is a Borel reduction of $E_x|_{M_x}$ into $E_{u(x)}^*$, if $(u(x), f(x)) \in [T]$, or into Δ , if $(u(x), f(x)) \notin [T]$. By applying ergodicity one more time, we conclude that $(u(x), f(x)) \in [T]$ and there is a Borel set of μ_x -measure 1, N_x , so that $E_x|_{N_x} \leq_B E_{u(x)}$, so by 4.2(ii), $u(x) = x$. So clearly f is a full uniformization of $[T]$ and it only remains to check that it is Borel. To see this, notice that the graph of f is given by

$$\begin{aligned} (x, y) \in G_f &\Leftrightarrow y = f(x) \\ &\Leftrightarrow \mu_x(\{z : (\pi_1 \circ g_x \circ \rho)(z) = (x, y)\}) = 1, \end{aligned}$$

so G_f is Borel, by Kechris [21], 17.25, thus f is Borel. -1

Denote by FBU the set of all trees T for which $[T]$ has a full Borel uniformization:

$$\text{FBU} = \{T : T \text{ is a tree on } \mathbb{N} \times \mathbb{N} \ \& \ [T] \text{ has a full Borel uniformization}\}.$$

We can identify here a tree with its characteristic function, a member of $2^{\mathbb{N}^{<\mathbb{N}} \times \mathbb{N}^{<\mathbb{N}}}$, which we can simply identify with $2^{\mathbb{N}}$. Clearly the set of trees is Borel. Since the set $E \subseteq 2^{\mathbb{N}} \times (\mathbb{N}^{\mathbb{N}})^2$ given by

$$(T, x, y) \in E \Leftrightarrow T \text{ is a tree and } (x, y) \in E_T$$

is clearly Borel, it follows that there is a Borel function p such that for each tree T , $p(T) \in D$ and

$$E_T = D_{p(T)}.$$

So by 5.2, $p^{-1}(\mathcal{E}_{\leq}^*) = p^{-1}(\mathcal{E}_{\sim}^*) = \text{FBU}$; thus in order to show that $\mathcal{E}_{\leq}^*, \mathcal{E}_{\sim}^*$ are Σ_2^1 -complete it is enough to show that FBU is Σ_2^1 -complete.

Following up a conversation with one of the authors, John Steel first found a proof of this using non-standard models. We give below a different proof based on effective descriptive set theory. We assume that the reader is familiar with this theory as exposed for example in Moschovakis [28].

Lemma 5.3. *The set FBU is Σ_2^1 -complete.*

Proof. It is enough to show that every Σ_2^1 subset of $\mathbb{N}^{\mathbb{N}}$ is of the form $f^{-1}(\text{FBU})$ for some Borel f .

We will first need some simple facts about trees (see Kechris [21], Section 2). Let S, T be trees on sets A, B , resp. We let

$$S \preceq T \Leftrightarrow \text{there is a strictly monotone map}$$

$$\varphi : S \rightarrow T.$$

Also let $S * T$ be the tree on $A \times B$ given by

$$S * T = \{(s, t) \in A^{<\mathbb{N}} \times B^{<\mathbb{N}} : \text{length}(s) = \text{length}(t) \ \& \ s \in S \ \& \ t \in T\}.$$

Then, recalling that a tree is called *illfounded* if it has an infinite branch, we clearly have that $S * T$ is illfounded iff both S, T are illfounded, so $S * T$ is wellfounded (*i.e.*, not illfounded) iff at least one of S, T is wellfounded. Moreover, if S is wellfounded but T is illfounded, then clearly $S \preceq S * T$. To see this, let $x \in [T]$ be an infinite branch of T and define $\varphi(s) = (s, x|\text{length}(s))$. Finally, recall that the set WF for all wellfounded trees is Π_1^1 -complete and that the map

$$\rho : \text{WF} \rightarrow \omega_1$$

(ω_1 = the first uncountable ordinal) given by

$$\rho(S) = \text{the rank of } S$$

is a Π_1^1 -rank (also called Π_1^1 -norm) on WF. Note also that for $T \in \text{WF}$, $S \preceq T \Leftrightarrow S \in \text{WF} \ \& \ \rho(S) \leq \rho(T)$.

Now fix a Σ_2^1 set $A \subseteq \mathbb{N}^{\mathbb{N}}$ and let $B \subseteq (\mathbb{N}^{\mathbb{N}})^2$ be Π_1^1 so that $u \in A \Leftrightarrow \exists v(u, v) \in B$ and let $g : (\mathbb{N}^{\mathbb{N}})^2 \rightarrow \mathbb{N}^{\mathbb{N}}$ be recursive so that $g(u, v)$ is a tree on \mathbb{N} , for each u, v , and $(u, v) \in B \Leftrightarrow g(u, v) \in \text{WF}$. Thus

$$\begin{aligned} u \in A &\Leftrightarrow \exists v(g(u, v) \in \text{WF}) \\ &\Leftrightarrow \exists T(T \in \text{WF} \ \& \ \exists \varphi : g(u, v) \rightarrow T, \varphi \text{ strictly monotone}). \end{aligned}$$

Since $\{(x, y) : y \in \Delta_1^1(x)\}$ is Π_1^1 , let f also be a recursive function such that $f(x, y)$ is a tree on \mathbb{N} for all x, y and

$$y \in \Delta_1^1(x) \Leftrightarrow f(x, y) \in \text{WF}.$$

Now define $R \subseteq (\mathbb{N}^{\mathbb{N}})^3$ by

$$(u, x, \langle y, T, v, \varphi \rangle) \in R \Leftrightarrow \varphi : g(u, v) \rightarrow f(x, y) * T \text{ is strictly monotone,}$$

where $\langle \rangle$ is an appropriate recursive coding of the objects indicated by a member of $\mathbb{N}^{\mathbb{N}}$. It is clear that R is closed in $(\mathbb{N}^{\mathbb{N}})^3$, so it is of the form $R = [U] = \{(x, y, z) \in (\mathbb{N}^{\mathbb{N}})^3 : \forall n(x|n, y|n, z|n) \in U\}$, for some tree U on $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$. Then if $f(u) = U(u) = \{(s, t) : \text{length}(s) = \text{length}(t) \ \& \ (u| \text{length}(s), s, t) \in U\}$, f is a Borel map and $R_u = [f(u)]$. We show that

$$\begin{aligned} u \in A &\Leftrightarrow R_u \text{ has a full Borel uniformization} \\ &\Leftrightarrow f(u) \in \text{FBU}, \end{aligned}$$

which shows that $A = f^{-1}(\text{FBU})$.

Recall that a closed set $C \subseteq (\mathbb{N}^{\mathbb{N}})^2$ admits a full Borel uniformization iff there is a parameter $p \in \mathbb{N}^{\mathbb{N}}$ such that for every x there is $y \in \Delta_1^1(x, p)$ with $(x, y) \in C$ (Moschovakis [28], 4D.4). So it is enough to show that:

$$u \in A \Leftrightarrow \exists p \forall x \exists \langle y, T, v, \varphi \rangle \in \Delta_1^1(x, p) R(u, x, \langle y, T, v, \varphi \rangle).$$

\Leftarrow : Fix such a p . Then take $x = p$ and find $y, T, v, \varphi \in \Delta_1^1(x, p) = \Delta_1^1(x)$ such that $\varphi : g(u, v) \rightarrow f(x, y) * T$ is strictly monotone. As $y \in \Delta_1^1(x), f(x, y) \in \text{WF}$, so $f(x, y) * T \in \text{WF}$, so $g(u, v) \in \text{WF}$ and thus $u \in A$.

\Rightarrow : Let T_0 be a fixed recursive illfounded tree, e.g., $T_0 = \mathbb{N}^{<\mathbb{N}}$. Let $u_0 \in A$. Pick v_0 so that $g(u_0, v_0) = S_0 \in \text{WF}$. Put $p = \langle u_0, v_0 \rangle$ (so also S_0 is recursive in p). Given any x we now have to find $\langle y, T, v, \varphi \rangle \in \Delta_1^1(x, p)$ satisfying $R(u_0, \langle y, T, v, \varphi \rangle)$. We take $v = v_0 \in \Delta_1^1(x, p)$. To find y, T, φ we consider two cases:

(i) $\omega_1^x > \rho(S_0)$ (where ω_1^x is the first ordinal not recursive in x).

Then let S_0^* be a wellfounded tree recursive in x with $\rho(S_0) = \rho(S_0^*)$. It follows that there must be some $y \in \Delta_1^1(x) \subseteq \Delta_1^1(x, p)$ such that $\rho(S_0) = \rho(S_0^*) \leq \rho(f(x, y))$, since otherwise

$$\forall y (y \in \Delta_1^1(x) \Leftrightarrow \rho(f(x, y)) < \rho(S_0^*)),$$

so that $\{y : y \in \Delta_1^1(x)\}$ is $\Delta_1^1(x)$, a contradiction (Moschovakis [28], 4D.16). Therefore there is also some $\varphi_1 : S_0 \rightarrow f(x, y)$ strictly monotone such that $\varphi_1 \in \Delta_1^1(x, y, S_0) \subseteq \Delta_1^1(x, p)$ and this immediately gives some $\varphi : g(u_0, v_0)(= S_0) \rightarrow f(x, y) * T_0, \varphi \in \Delta_1^1(x, p)$, which is strictly monotone. Thus if we choose $T = T_0, \langle y, T, v, \varphi \rangle$ works.

(ii) $\omega_1^x \leq \rho(S_0)$.

Then we claim that we can find $y \notin \Delta_1^1(x)$ such that $\mathcal{O}^{x, y} \in \Delta_1^1(x, p)$, where \mathcal{O}^a is the Kleene \mathcal{O} relative to a , i.e., the complete $\Pi_1^1(a)$ subset of \mathbb{N} .

Granting this claim, $f(x, y)$ is illfounded and we pick a branch $b \in [f(x, y)]$, which is recursive in $\mathcal{O}^{x, y}$, thus $b \in \Delta_1^1(x, p)$. Then let $\varphi(s) = (b| \text{length}(s), s)$ for $s \in g(u_0, v_0)$. Clearly $\varphi \in \Delta_1^1(x, p)$ and $\varphi : g(u_0, v_0)(= S_0) \rightarrow f(x, y) * S_0$ is strictly monotone. Thus if we now choose $T = S_0, \langle y, T, v, \varphi \rangle$ works.

It remains to prove the claim: Look at the set

$$\{y : \omega_1^{x, y} = \omega_1^x \ \& \ y \notin \Delta_1^1(x)\}.$$

Since $\omega_1^x \leq \rho(S_0) < \omega_1^{S_0,x}$ we have that $\mathcal{O}^x \in \Delta_1^1(S_0, x)$, so this set is $\Delta_1^1(S_0, x)$. It is also comeager, so it has an element $y \in \Delta_1^1(S_0, x)$ (Moschovakis [28], 4F.20). Then as $\omega_1^{x,y} < \omega_1^{S_0,x} = \omega_1^{S_0,x,y}$, we have that $\mathcal{O}^{x,y} \in \Delta_1^1(S_0, x, y) \subseteq \Delta_1^1(S_0, x) \subseteq \Delta_1^1(x, p)$, so we are done. \dashv

Now let us say that a closed set $F \subseteq (\mathbb{N}^{\mathbb{N}})^2$ has *full projection* if for every x there is some y with $(x, y) \in F$. By the Jankov, von Neumann Uniformization Theorem (see Kechris [21], 18.1), this is equivalent to saying the F admits a full $\sigma(\Sigma_1^1)$ -measurable uniformization, where $\sigma(\Sigma_1^1)$ is the smallest σ -algebra generated by the analytic sets.

Then by inspecting the argument in 5.2, we see the following, where $E \leq_{\sigma(\Sigma_1^1)} F$ means that E can be reduced to F via a $\sigma(\Sigma_1^1)$ -measurable function, and similarly for $\sim_{\sigma(\Sigma_1^1)}$.

Lemma 5.4. *For each tree T we have*

$$\begin{aligned} E^* \leq_{\sigma(\Sigma_1^1)} E_T &\Leftrightarrow E^* \sim_{\sigma(\Sigma_1^1)} E_T \\ &\Leftrightarrow [T] \text{ has full projection.} \end{aligned}$$

Let

$$\text{FP} = \{T : T \text{ is a tree on } \mathbb{N} \times \mathbb{N} \ \& \ [T] \text{ has full projection}\}.$$

Then it is quite standard that FP is a Π_2^1 -complete set: If $A \subseteq \mathbb{N}^{\mathbb{N}}$ is Π_2^1 , then $x \in A \Leftrightarrow \forall y(x, y) \in B$, where B is Σ_1^1 , so $(x, y) \in B \Leftrightarrow \exists z(x, y, z) \in F$, where F is closed in $(\mathbb{N}^{\mathbb{N}})^3$, thus $F = [U]$, for some tree U on $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$. For each $x \in \mathbb{N}^{\mathbb{N}}$, let

$$U(x) = \{(s, t) \mid \text{length}(s) = \text{length}(t) \text{ and } (x \upharpoonright \text{length}(s), s, t) \in U\}.$$

Then $x \in A \Leftrightarrow U(x) \in \text{FP}$, and clearly $x \mapsto U(x)$ is Borel.

We have $\text{FBU} \subseteq \text{FP}$ and since the first set is Σ_2^1 -complete while the second is Π_2^1 -complete, they are clearly not equal, *i.e.*, $\text{FBU} \not\subseteq \text{FP}$. (This is a well-known fact; see, e.g., Kechris [21], 18.17, but the argument used here seems to be new.) In particular, it follows trivially that there is a tree T such that $E^* \leq_{\sigma(\Sigma_1^1)} E_T$ but $E^* \not\leq_B E_T$. So we have:

Theorem 5.5. *There is a pair of countable Borel equivalence relations E, F such that $E \leq_{\sigma(\Sigma_1^1)} F$ but $E \not\leq_B F$ (and we moreover have that $F \leq_B E$).*

6. FINITE RANK TORSION-FREE ABELIAN GROUPS

Let \cong_n be the isomorphism relation on torsion-free abelian groups of rank $\leq n$. These are exactly the groups isomorphic to subgroups of $(\mathbb{Q}^n, +)$. Let us also denote by \cong_n^* the restriction of \cong_n to the *rigid groups*, where we call an abelian group rigid if its only automorphisms are $x \mapsto x, x \mapsto -x$.

It follows from a result of Baer (see, e.g., Fuchs [10]) that $(\cong_1) \sim_B (\cong_1^*) \sim_B E_0$, where E_0 is the equivalence relation on $2^{\mathbb{N}}$ defined by: $x E_0 y \Leftrightarrow \exists n \forall m \geq n (x_m = y_m)$. Hjorth [15] has shown that $E_0 <_B (\cong_n)$ for all $n \geq 2$ and Thomas [33] extended this to show that $E_0 <_B (\cong_n^*)$ for $n \geq 2$. Also Hjorth [15] showed that for $n \geq 3$, \cong_n is not essentially treeable, *i.e.*, there is no treeable countable Borel equivalence relation E with $(\cong_n) \sim_B E$, and Thomas [33] extended this to \cong_n^* , for $n \geq 3$. Recently Kechris showed that \cong_n is not essentially treeable for all $n \geq 2$. This is still open for \cong_2^* . (Recall that a countable Borel equivalence relation E on

X is *treeable* if there is a Borel acyclic graph on X whose connected components are the E -classes.) We show here the following, which in particular implies that \cong_n^* is universal.

Theorem 6.1. *For any $n \geq 1$, $(\cong_n^*) <_B (\cong_{n+1}^*)$.*

Proof. Let $S(\mathbb{Q}^n)$ be the space of all subgroups of \mathbb{Q}^n (a closed subset of the space of all subsets of \mathbb{Q}^n). Since every isomorphism between subgroups of \mathbb{Q}^n is induced by a matrix in $GL_n(\mathbb{Q})$, we can view, up to \sim_B , \cong_n as being the equivalence relation on $S(\mathbb{Q}^n)$ induced by the canonical action of $GL_n(\mathbb{Q})$.

Let $Z_n \subseteq S(\mathbb{Q}^n)$ consist of all subgroups containing \mathbb{Z}^n . Then $SL_n(\mathbb{Z})$ acts on Z_n and Hjorth [15] shows that there is a probability measure ν_n on Z_n invariant under this action with the following properties:

- (i) The equivalence relation on Z_n induced by this action is not tame, even on any set of ν_n -measure 1.
- (ii) (Thomas [33]) There is a Borel set of ν_n -measure 1, Y_n , such that every element of Y_n is rigid.

Now consider the ergodic decomposition of the action of $SL_n(\mathbb{Z})$ on Z_n , as in Varadarajan [34]. More precisely, let \mathcal{E} be the standard Borel space of ergodic, invariant measures of this action. Then there is a Borel surjection $\pi : Z_n \rightarrow \mathcal{E}$, which is $SL_n(\mathbb{Z})$ -invariant, such that if $X_e = \{x \in Z_n : \pi(x) = e\}$, then $e(X_e) = 1, \forall e \in \mathcal{E}$, and for any Borel set $A \subseteq Z_n$, and any $SL_n(\mathbb{Z})$ -invariant probability measure, ν , we have $\nu(A) = \int e(A) d\nu_*(e)$, where $\nu_* = \pi_*\nu$ is the measure on \mathcal{E} defined by $\nu_*(B) = \nu(\pi^{-1}(B))$.

Since $\nu_n(Y_n) = 1$, it follows that there is a Borel set $\mathcal{E}_1 \subseteq \mathcal{E}$ such that $(\nu_n)_*(\mathcal{E}_1) = 1$ and for $e \in \mathcal{E}_1, e(Y_n) = 1$. Then there must be some $e_1 \in \mathcal{E}_1$ which is non-atomic, since otherwise for each $e \in \mathcal{E}_1$, there is a unique orbit of $SL_n(\mathbb{Z})$, say θ_e , with $e(\theta_e) = 1$, and then the set $W = \{x \in Z_n : \pi(x) \in \mathcal{E}_1 \ \& \ x \in \theta_{\pi(x)}\} = \{x \in Z_n : \pi(x) \in \mathcal{E}_1 \ \& \ \pi(x)(SL_n(\mathbb{Z}) \cdot x) = 1\}$ is Borel, $SL_n(\mathbb{Z})$ -invariant, and the action of $SL_n(\mathbb{Z})$ on W is tame, as for $x, y \in W, y \in SL_n(\mathbb{Z}) \cdot x \Leftrightarrow \pi(y) = \pi(x)$. But clearly $\nu_n(W) = 1$, which is a contradiction.

So we can assume that there is an ergodic, non-atomic, $SL_n(\mathbb{Z})$ -invariant measure $\mu_n (= e_1)$ on Z_n and there is a Borel $SL_n(\mathbb{Z})$ -invariant set $X_n (= X_{e_1} \cap Y_n) \subseteq Z_n$ with $\mu_n(X_n) = 1$, such that every element of X_n is rigid.

It is clear that $(\cong_n^*) \leq_B (\cong_{n+1}^*)$ and Thomas' result shows that $(\cong_1^*) <_B (\cong_2^*)$. So assume $n \geq 2$ and $(\cong_{n+1}^*) \leq_B (\cong_n^*)$, towards a contradiction. Say f is a Borel function reducing \cong_{n+1}^* to \cong_n^* . In particular, for $x, y \in X_{n+1}$ we have

$$xE_{SL_{n+1}(\mathbb{Z})}^{X_{n+1}}y \Rightarrow f(x) \cong_n^* f(y).$$

By ergodicity, there is $m \leq n$ and a set of μ_{n+1} -measure 1, so that for x in that set $f(x)$ actually has rank exactly m , so that we can actually assume (by shrinking X_{n+1} a bit if necessary) that for $x \in X_{n+1}, f(x) \in S(\mathbb{Q}^m)$, $f(x)$ has rank m , and

$$xE_{SL_{n+1}(\mathbb{Z})}^{X_{n+1}}y \Rightarrow f(x) \cong_m^* f(y).$$

It follows that there is a strict Borel cocycle $\alpha : SL_{n+1}(\mathbb{Z}) \times X_{n+1} \rightarrow A^m(\mathbb{Q})$, where $A^m = GL_m / \{I_m, -I_m\}$, with $I_m =$ the identity $m \times m$ matrix, such that

$$\alpha(g, x) = [\text{the unique } h \in A^m(\mathbb{Q}) \text{ such that } h \cdot f(x) = f(g \cdot x)].$$

Now $|\det(\alpha)|$ is also a strict Borel cocycle, $|\det(\alpha)| : SL_{n+1}(\mathbb{Z}) \times X_{n+1} \rightarrow \mathbb{Q}_+^*$ ($=$ the multiplicative group of the positive rationals), so since \mathbb{Q}_+^* is amenable and

$SL_{n+1}(\mathbb{Z})$ is Kazhdan (see Zimmer [35], 4.1.2, and de la Harpe and Valette [3], 3.a.5, page 34, resp.), it follows by 3.1 that there is a Borel cocycle $\delta : SL_{n+1}(\mathbb{Z}) \times X_{n+1} \rightarrow \mathbb{Q}_+^*$ taking values in a finite subgroup of \mathbb{Q}_+^* , thus into $\{1\}$, such that $|\det(\alpha)| \sim \delta$. So there is Borel $F : X_{n+1} \rightarrow \mathbb{Q}_+^*$ with $|\det(\alpha(g, x))| = F(g \cdot x)F(x)^{-1}$ μ_{n+1} -a.e.

Choose for each $a \in \mathbb{Q}_+^*$ an element $\bar{a} \in A^m(\mathbb{Q})$ with $|\det(\bar{a})| = a$ and let $\bar{F} : X_{n+1} \rightarrow A^m(\mathbb{Q})$ be defined by $\bar{F}(x) = \overline{F(x)}$. Then if

$$\bar{\alpha}(g, x) = (\bar{F}(g \cdot x))^{-1} \alpha(g, x) \bar{F}(x),$$

$\bar{\alpha}$ is a Borel cocycle from $SL_{n+1}(\mathbb{Z}) \times X_{n+1}$ into $A^m(\mathbb{Q})$, $\alpha \sim \bar{\alpha}$, and as

$$\begin{aligned} |\det(\bar{\alpha}(g, x))| &= |\det(\bar{F}(g \cdot x))^{-1}| \cdot |\det(\alpha(g, x))| \cdot |\det(\bar{F}(x))| \\ &= F(g \cdot x)^{-1} \cdot |\det(\alpha(g, x))| \cdot F(x) = 1 \quad \mu_{n+1}\text{-a.e.}, \end{aligned}$$

we conclude that $\bar{\alpha}(g, x) \in H^m(\mathbb{Q})$, μ_{n+1} -a.e., with $H^m = SL_m^\pm / \{I_m, -I_m\} \subseteq A^m$, where SL_m^\pm consists of all $m \times m$ matrices with determinant ± 1 . Now $SL_{n+1}(\mathbb{Z})$ is a Kazhdan group, so by 3.2 we have that $\bar{\alpha} \sim \beta$, where $\beta(SL_{n+1}(\mathbb{Z}) \times X_{n+1}) \subseteq \Lambda$, a finitely generated subgroup of $H^m(\mathbb{Q})$, and we can assume that β is a strict cocycle. So there is a finite set of primes p_1, \dots, p_n such that $\Lambda \subseteq H^m(\mathbb{Z}[1/p_1, \dots, 1/p_n])$.

Now $H^m(\mathbb{Z}[1/p_1, \dots, 1/p_n])$ is a discrete subgroup of $H^m(\mathbb{R}) \times H^m(\mathbb{Q}_{p_1}) \times \dots \times H^m(\mathbb{Q}_{p_n})$ (see the corresponding argument in the proof of 4.5) and $SL_{n+1}(\mathbb{Z})$ is a lattice in $SL_{n+1}(\mathbb{R})$ (see Zimmer [35], page 1, Example (c)).

Consider the action of $SL_{n+1}(\mathbb{Z})$ on X_{n+1} and the cocycle $\beta : SL_{n+1}(\mathbb{Z}) \times X_{n+1} \rightarrow H^m(\mathbb{Z}[1/p_1, \dots, 1/p_n])$, and using Section 2 (paragraph following 2.2) consider the induced Borel action of $SL_{n+1}(\mathbb{R})$ on Y (which also has an invariant, ergodic measure) and the induced cocycle $\hat{\beta} : SL_{n+1}(\mathbb{R}) \times Y \rightarrow H^m(\mathbb{Z}[1/p_1, \dots, 1/p_n])$. Viewing $\hat{\beta}$ as a cocycle into $H^m(\mathbb{R}) \times H^m(\mathbb{Q}_{p_1}) \times \dots \times H^m(\mathbb{Q}_{p_n})$ and letting π be the projection of this product to one of its factors, consider the cocycle $\pi \circ \hat{\beta}$. By 3.8, $\pi \circ \hat{\beta}$ is equivalent to a cocycle with values in a compact subgroup, so this shows that $\hat{\beta} \sim \gamma_1$, where γ_1 is a cocycle with values into a compact subgroup, so, by 2.4, $\hat{\beta}$ is equivalent to a cocycle with values in a finite subgroup of $H^m(\mathbb{Z}[1/p_1, \dots, 1/p_n])$ and thus, by 2.3, β is equivalent to a cocycle γ with values in a finite subgroup of $H^m(\mathbb{Z}[1/p_1, \dots, 1/p_n]) \subseteq A^m(\mathbb{Q})$. Since $\alpha \sim \bar{\alpha} \sim \beta \sim \gamma$, we have $\alpha \sim \gamma$ as well. Say $\alpha(g, x) = B(g \cdot x)\gamma(g, x)B(x)^{-1}$, μ_{n+1} -a.e., $B : X_{n+1} \rightarrow A^m(\mathbb{Q})$ a Borel function. Then let $M \subseteq X_{n+1}$ be a Borel set of positive μ_{n+1} -measure on which B is constant. It follows that $\phi(x) = \{f(y) : y \in M \text{ \& } yE_{SL_{n+1}(\mathbb{Z})}^{X_{n+1}}x\}$ is finite, for all $x \in M$. By ergodicity, $\phi(x)$ is fixed on a Borel set $M_0 \subseteq M$ of positive μ_{n+1} -measure, and so if $M_1 = SL_{n+1}(\mathbb{Z}) \cdot M_0$, it follows that $\mu_{n+1}(M_1) = 1$ and $f(M_1)$ is contained in a single \cong_m^* -class. But since f is a reduction, this means that M_1 is contained in a single \cong_{n+1}^* -class, *i.e.*, in a single $GL_{n+1}(\mathbb{Q})$ orbit, so it is countable, violating the non-atomicity of μ_{n+1} . -1

7. MISCELLANEA

We mention here some other results that can be proved by the methods used in this paper.

(i) Consider the canonical action of $GL_n(\mathbb{Z})$ on $\mathbb{T}^n = (\mathbb{R}/\mathbb{Z})^n$ and denote by R_n the corresponding equivalence relation. Then we show (Theorem 7.1) that $m < n \Rightarrow R_n \not\leq_B R_m$. (In fact, in Theorem 7.1, we will prove a more general result concerning “essential” Borel reducibility, defined below.) We also show that $m \leq n \Rightarrow R_m \leq_B R_n$ (Theorem 7.2), therefore, $m < n$ implies that $R_m <_B R_n$.

On the other hand, consider the canonical action of $SL_n(\mathbb{Z})$ on $\mathbb{T}^n = (\mathbb{R}/\mathbb{Z})^n$ and denote by R'_n the corresponding equivalence relation. It should be pointed out here that we do not know whether, in general, $m < n \Rightarrow R'_m <_B R'_n$.

If R is a countable Borel equivalence relation on a standard Borel space X , if μ is a measure on X , and if S is a countable equivalence relation on a standard Borel space Y , then we say that R is *essentially reducible* to S if there is a conull Borel subset $X_0 \subseteq X$ such that $R|X_0 \leq_B S$. This will be denoted $R \leq_B S$ (a.e.). The notation $R \not\leq_B S$ (a.e.) will mean that R is not essentially reducible to S , i.e., that, for any conull Borel subset $X_0 \subseteq X$, we have $R|X_0 \not\leq_B S$.

For all integers $v \geq 1$, let $\mathbb{Z}_v := (1/v)\mathbb{Z} = \{i/v : i \in \mathbb{Z}\} \subseteq \mathbb{Q}$. For all integers $l, v \geq 1$, let R_{lv} be the equivalence relation defined by the orbits of the action of $GL_l(\mathbb{Z})$ on $(\mathbb{R}/\mathbb{Z}_v)^l$.

Theorem 7.1. *For all integers $m, n, s, t \geq 1$: $m < n \Rightarrow R_{nt} \not\leq_B R_{ms}$ (a.e.).*

Proof. The proof is by induction on m , and we may assume $m \geq 2$. Fix integers $n, s, t \geq 1$ and assume that $R_{nt} \leq_B R_{ms}$ (a.e.). We aim for a contradiction.

Let $G := GL_m(\mathbb{Z})$, $R := R_{ms}$ and $S := R_{nt}$. Then $S \leq_B R$ (a.e.). Let $A := (\mathbb{R}/\mathbb{Z}_s)^m$ and let $B := (\mathbb{R}/\mathbb{Z}_t)^n$. We denote the action of G on A by $(g, a) \mapsto ga : G \times A \rightarrow A$. Let $A' := \{a \in A : \exists g \in G \setminus \{1_G\} (ga = a)\}$ and let μ be the canonical measure for S . By the argument in 6.1, we see that $S \not\leq_B R|(A \setminus A')$ (a.e.). Ergodicity of μ then implies that $S \leq_B R|A'$ (a.e.).

For any integer $d \geq 1$, for any $a = (a_1, \dots, a_d) \in \mathbb{R}^d$, we will say that a is *totally irrational* if $1, a_1, \dots, a_d$ are linearly independent over \mathbb{Q} . For any integer $d \geq 1$, for any $a = (a_1, \dots, a_d) \in \mathbb{R}^d$, we will say that a is in *canonical position* if there exists $i \in \{0, \dots, d\}$ such that $(a_1, \dots, a_i) \in \mathbb{Q}^i$ and such that (a_{i+1}, \dots, a_d) is totally irrational. We call i the *index* of a and we call (a_1, \dots, a_i) the *rational part* of a . (Note: If $a \in \mathbb{Q}^d$, then a is in canonical position, with index d and rational part a . If a is totally irrational, then a is in canonical position, with index 0 and with no rational part.)

The conditions that a be totally irrational or in canonical position are both invariant under translation by elements of \mathbb{Q}^d . It follows that, if Γ is an additive subgroup of \mathbb{Q}^d and if $a \in \mathbb{R}^d/\Gamma$ and if a preimage of a in \mathbb{R}^d is totally irrational (resp. in canonical position), then all preimages of a in \mathbb{R}^d are totally irrational (resp. in canonical position). In this case, we will say that a is *totally irrational* (resp. in *canonical position*).

We claim, for all additive subgroups $\Gamma \subseteq \mathbb{Q}^d$, for all $a \in \mathbb{R}^d/\Gamma$, that there exists $g \in GL_d(\mathbb{Z})$ such that ga is in canonical position. *Sketch of proof:* It suffices to prove this when $\Gamma = \{0\}$. By induction on d , it suffices to show that if a is not totally irrational, then, for some $g \in GL_d(\mathbb{Z})$, the first coordinate of ga is rational. However, if a is not totally irrational, then there exists $k \in \mathbb{Z}^d \setminus \{0\}$ such that $k \cdot a \in \mathbb{Q}$, where \cdot denotes the standard dot product. Dividing k by the gcd of its entries, we may assume that k is primitive (i.e., that the entries of k are relatively prime). Then k is the first vector in some base of the finitely generated abelian group \mathbb{Z}^d . So, making a change of base, we may replace a by some element of $(GL_d(\mathbb{Z}))a$ and k by $(1, 0, \dots, 0)$. The first coordinate of a is then $k \cdot a$, which is rational, as desired. *The claim is proved.*

For all $i \in \{1, \dots, m\}$, let $Q_i := (\mathbb{Q}/\mathbb{Z}_s)^i$. For all $i \in \{1, \dots, m\}$, for all $q \in Q_i$, let A'_{iq} denote the collection of all $a \in A$ such that a is in canonical position with

index i and with rational part q . Note, for all $q \in Q_m$, that we have $A'_{mq} = \{q\}$, so $|A'_{mq}| = 1$. Let A'_0 be the union of all A'_{iq} over $i \in \{1, \dots, m\}$ and $q \in Q_i$.

Let $a \in A'$. Then there is some $g \in G \setminus \{1_G\}$ such that $(g - 1_G)a = 0$. Then if k is a non-zero row in $g - 1_G$, we have $k \cdot a = 0$, where \cdot denotes the standard dot product. This proves that a is not totally irrational. Choosing $g_0 \in G$ such that g_0a is in canonical position, we find that g_0a is not totally irrational, so the index i of g_0a is not 0. If q is the rational part of g_0a , then we have $g_0a \in A'_{iq}$, so $a \in g_0^{-1}A'_{iq} \subseteq GA'_0$. This argument proves that $A' \subseteq GA'_0$. Since A' is G -invariant and since $A'_0 \subseteq A'$, it follows that $GA'_0 \subseteq A'$. Then $A' = GA'_0$.

Then $R|A' \leq_B R|A'_0$, so, by ergodicity of S , fix $i \in \{1, \dots, m\}$ and $q \in Q_i$ such that $S \leq_B R|A'_{iq}$ (a.e.). Since μ is not atomic, we see that $|A'_{iq}| \neq 1$, which implies that $i \neq m$, so $i \in \{1, \dots, m - 1\}$. Choose $q_1, \dots, q_i \in \mathbb{Q}/\mathbb{Z}_s$ such that $q = (q_1, \dots, q_i)$.

Let $U := \{s, 2s, 3s, \dots\}$. For all $u \in U$, let $p_u : \mathbb{R}/\mathbb{Z}_s \rightarrow \mathbb{R}/\mathbb{Z}_u$ be the canonical homomorphism. Choose $r \in U$ such that $\mathbb{Z}q_1 + \dots + \mathbb{Z}q_i = \ker p_r$. Let $j := m - i$. Then $1 \leq j \leq m - 1$.

Define $p : (\mathbb{R}/\mathbb{Z}_s)^j \rightarrow (\mathbb{R}/\mathbb{Z}_r)^j$ by $p(c_1, \dots, c_j) = (p_r(c_1), \dots, p_r(c_j))$. Define $f : A'_{iq} \rightarrow (\mathbb{R}/\mathbb{Z}_r)^j$ by $f(a_1, \dots, a_m) = p(a_{i+1}, \dots, a_m)$. It remains to prove that f is a Borel reduction of $R_{ms}|A'_{iq}$ to R_{jr} , whereupon we will have $S \leq_B R_{jr}$ (a.e.), contradicting the induction assumption.

So fix $a = (a_1, \dots, a_m) \in A'_{iq}$ and $b = (b_1, \dots, b_m) \in A'_{iq}$. Let $a_0 := f(a)$ and let $b_0 := f(b)$ so $a_0, b_0 \in (\mathbb{R}/\mathbb{Z}_r)^j$. We wish to show:

$$\exists g \in G (ga = b) \quad \text{iff} \quad \exists g_0 \in \text{GL}_j(\mathbb{Z}) (g_0a_0 = b_0).$$

Set $a' = (a_1, \dots, a_i)$ and $b' := (b_1, \dots, b_i)$. Set $a'' = (a_{i+1}, \dots, a_m)$ and $b'' := (b_{i+1}, \dots, b_m)$. Then $a' = q = b'$. Moreover, a'' and b'' are totally irrational. Also, $a_0 = p(a'')$ and $b_0 = p(b'')$.

Proof of "if". By definition of r , we have $\ker p = \mathbb{Z}^{j \times i}q$. So, since we have $p(b'' - g_0a'') = b_0 - g_0a_0 = 0$, choose $Z \in \mathbb{Z}^{j \times i}$ such that $b'' - g_0a'' = Za'$. Recall that $a' = q = b'$.

Let I denote the $i \times i$ identity matrix and let 0 denote the $i \times j$ matrix with all entries equal to 0. Let $g := \begin{pmatrix} I & 0 \\ Z & g_0 \end{pmatrix}$. Then $ga = b$. *This concludes the proof of "if".*

Proof of "only if". Choose $X \in \mathbb{Z}^{i \times i}$, $Y \in \mathbb{Z}^{i \times j}$, $Z \in \mathbb{Z}^{j \times i}$ and $g_0 \in \mathbb{Z}^{j \times j}$ such that

$$g = \begin{pmatrix} X & Y \\ Z & g_0 \end{pmatrix}.$$

Then $ga = b$ implies that $Xa' + Ya'' = b'$ and $Za' + g_0a'' = b''$. We wish to show that $\det g_0 \in \{\pm 1\}$ and that $g_0a_0 = b_0$.

We have $Ya'' = b' - Xa' \in Q_i + \mathbb{Z}^{i \times i}Q_i \subseteq Q_i$. On the other hand, a'' is totally irrational. It follows that $Y = 0$. Then $\det g = (\det X)(\det g_0)$ and $\det X, \det g_0 \in \mathbb{Z}$ and $\det g \in \{\pm 1\}$. So $\det g_0 \in \{\pm 1\}$.

Define $\pi : (\mathbb{R}/\mathbb{Z}_s)^i \rightarrow (\mathbb{R}/\mathbb{Z}_r)^i$ by $\pi(c_1, \dots, c_i) = (p_r(c_1), \dots, p_r(c_i))$. Then $b_0 = p(b'') = p(Za' + g_0a'') = Z(\pi(a')) + g_0(p(a''))$. Since $a' = q$, it follows from the definition of r that $\pi(a') = 0$. Moreover, $p(a'') = a_0$. Then $b_0 = g_0a_0$, as desired. *This concludes the proof of "only if".* \dashv

Theorem 7.2. *For all integers $m, n \geq 1 : m \leq n \Rightarrow R_m \leq_B R_n$.*

Proof. We will use the notation and terminology established in the proof of 7.1. It is of course enough to show that for $m \geq 1$, $R_m \leq_B R_{m+1}$.

We start by noticing that if $d \geq 1$, then $a \in (\mathbb{R}/\mathbb{Z})^d$ is in canonical position with index $i \in \{0, \dots, d\}$ and $b \in (\mathbb{R}/\mathbb{Z})^d$ is in canonical position with index $j \in \{0, \dots, d\}$, and if $i \neq j$, then a, b are not in the same $\text{GL}_d(\mathbb{Z})$ -orbit.

Next we verify that if $(a_1, \dots, a_d) \in (\mathbb{R}/\mathbb{Z})^d$ is in canonical position and has index $i \geq 1$, then there exists $g \in \text{GL}_d(\mathbb{Z})$ with

$$ga = (0, \dots, 0, q, a_{i+1}, \dots, a_d),$$

for some $q \in \mathbb{Q}/\mathbb{Z}$. For that it is enough to check that if $i \geq 1$ and $(q_1, \dots, q_i) \in (\mathbb{Q}/\mathbb{Z})^i$, then there is $q \in \mathbb{Q}/\mathbb{Z}$ and $g \in \text{GL}_i(\mathbb{Z})$ with

$$g(0, \dots, 0, q) = (q_1, \dots, q_i).$$

To see this, notice that we can write $q_\ell = m_\ell r \pmod{\mathbb{Z}}$, where $(m_1, \dots, m_i) \in \mathbb{Z}^i$ are relatively prime and $r \in \mathbb{Q}$, so there is $g \in \text{GL}_i(\mathbb{Z})$ with last column equal to (m_1, \dots, m_i) (viewed as a column vector). If $q = r \pmod{\mathbb{Z}}$, then $g(0, \dots, 0, q) = (q_1, \dots, q_i)$.

So for $0 \leq i \leq m$, let

$$A_m^i = \{a \in (\mathbb{R}/\mathbb{Z})^m : a = (0, \dots, q, a_{i+1}, \dots, a_m), \\ \text{for some } q \in \mathbb{Q}/\mathbb{Z} \text{ and } (a_{i+1}, \dots, a_m) \\ \text{totally irrational}\}.$$

Then it is enough to show that

$$R_m|A_m^i \leq_B R_{m+1}|A_{m+1}^{i+1}.$$

We consider cases on $i \in \{0, \dots, m\}$.

Case 1: $i = 0$. Then $(a_1, \dots, a_m) \in A_m^0 \mapsto (0, a_1, \dots, a_m) \in A_{m+1}^1$ is a reduction.

Case 2: $i > 1$. Then again we claim that

$$(a_1, \dots, a_m) \in A_m^1 \mapsto (0, a_1, \dots, a_m) \in A_{m+1}^{i+1}$$

is a reduction. The main point here is to observe that if $(0, 0, \dots, 0, q), (0, 0, \dots, 0, r) \in (\mathbb{Q}/\mathbb{Z})^{i+1}$ are in the same $\text{GL}_{i+1}(\mathbb{Z})$ -orbit and $i > 1$, then q, r generate the same subgroup of \mathbb{R}/\mathbb{Z} , so $(0, \dots, 0, q), (0, \dots, 0, r) \in (\mathbb{Q}/\mathbb{Z})^i$ are in the same $\text{GL}_i(\mathbb{Z})$ -orbit.

Case 3: $i = 1$. Fix a set of representatives $T = \{t_j : j = 0, 1, 2, \dots\}$ for the following equivalence relation on $\mathbb{Q}/\mathbb{Z} : q \sim q'$ iff $q = q'$ or $q = -q'$. We agree that $t_0 = 0 \pmod{\mathbb{Z}}$. Every $(q, a_2, \dots, a_m) \in A_m^1$ can be sent to some $(t_i, a_2, \dots, a_m) \in A_m^1$ by an element of $\text{GL}_m(\mathbb{Z})$, so if

$$B_m^1 = \{(q, a_2, \dots, a_m) \in A_m^1 : q \in T\},$$

it is enough to show that

$$R_m|B_m^1 \leq_B R_{m+1}|A_{m+1}^2.$$

For each $j \geq 1$, choose $m_j, n_j \in \mathbb{Z}, n_j \neq 0$, relatively prime so that $t_j = \frac{m_j}{n_j} \pmod{\mathbb{Z}}$. Then notice that by induction on $j \geq 1$, we can find $0 \neq k_j \in \mathbb{Z}$ such that

k_j, m_j are relatively prime and such that, if we put

$$s_j = \frac{1}{k_j n_j} \pmod{\mathbb{Z}} \quad \text{and} \quad u_j = \frac{m_j}{k_j n_j} \pmod{\mathbb{Z}},$$

then for $\ell < j$, $(s_j, u_j), (s_\ell, u_\ell)$ are in distinct $\text{GL}_2(\mathbb{Z})$ -orbits. The main point here is that for $0 \neq k \in \mathbb{Z}$, $(\frac{1}{kn_j} \pmod{\mathbb{Z}}, 0)$ is in the $\text{GL}_2(\mathbb{Z})$ -orbit of $(\frac{1}{kn_j} \pmod{\mathbb{Z}}, t_j)$, and for $k \neq k'$, $(\frac{1}{kn_j} \pmod{\mathbb{Z}}, 0)$ and $(\frac{1}{k'n_j} \pmod{\mathbb{Z}}, 0)$ are in distinct $\text{GL}_2(\mathbb{Z})$ -orbits.

For each $k = 1, 2, \dots$, fix a Borel map $x \mapsto x/k : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}/\mathbb{Z}$ such that $k(x/k) = x$. We are now ready to define our reduction: First map $(0, a_2, \dots, a_m) \in B_m^1$ to $(0, 0, a_2, \dots, a_m)$. Then, for each $j \geq 1$, map $(t_j, a_2, \dots, a_m) \in B_m^1$ to $(s_j, u_j, a_2/k_j, \dots, a_m/k_j) \in A_{m+1}^2$. \dashv

(ii) Recall from the introduction that there is a universal countable Borel equivalence relation E_∞ . This equivalence relation has many manifestations, *i.e.*, there are various examples of countable Borel equivalence relations E with $E \sim_B E_\infty$. For example, the translation action of the free group with two generators F_2 acting on its subsets gives rise to a universal equivalence relation. It has been observed that there is no known *free* Borel action of a countable group which gives rise to a universal equivalence relation. So this led to the following open problem: Is there a countable group Γ and a free Borel action of Γ so that the corresponding equivalence relation is universal? We only remark that one can show by the methods used here that no Γ which is (up to isomorphism) a discrete subgroup of some $\text{GL}_n(\mathbb{C})$ or $\text{GL}_n(\mathbb{Q}_p)$ can possibly work.

(iii) Finally, although in Section 5 we have computed the exact descriptive complexity of the equivalence relation \mathcal{E}_\sim (of bi-reducibility of countable Borel equivalence relations), there are other questions that can be raised concerning the possible complexity of \mathcal{E}_\sim . For example, Theorem 4.1 implies that if R is the equivalence relation on the set of codes D of Borel sets, say of $2^{\mathbb{N}}$ (defined in a similar fashion as in Section 5), and $d \in D \mapsto D_d$ is the associated map (so that the Borel sets are exactly those of the form D_d , for $d \in D$), then the equivalence relation

$$R(d, e) \Leftrightarrow D_d = D_e$$

is Borel reducible to \mathcal{E}_\sim . From this it immediately follows that every Borel equivalence relation is Borel reducible to \mathcal{E}_\sim and Gao [11] has extended this to show that every Σ_1^1 equivalence relation is Borel reducible to \mathcal{E}_\sim . One can further ask here: Is every Π_1^1 or even Σ_2^1 equivalence relation Borel reducible to \mathcal{E}_\sim ?

ACKNOWLEDGMENT

We would like to thank G. Hjorth for asking a question (related to Section 6) that triggered our investigations in this paper. We would also like to thank an anonymous referee for many detailed comments that helped improve the presentation.

REFERENCES

1. S. Adams, Indecomposability of treed equivalence relations, *Israel J. Math.*, **86 (3)**, 362-380, 1988. MR **90m**:28022
2. S. Adams, Trees and amenable equivalence relations, *Erg. Th. and Dyn. Syst.*, **10**, 1-14, 1990. MR **91d**:28041

3. P. de la Harpe and A. Valette, *La Propriété (T) de Kazhdan pour les Groupes Localement Compacts*, Astérisque, **175**, Soc. Math. de France, 1989. MR **90m**:22001
4. J. Dieudonné, *Sur les Groupes Classiques*, Hermann, Paris, 1973. MR **49**:9094
5. R. Dougherty, S. Jackson, and A.S. Kechris, The structure of hyperfinite Borel equivalence relations, *Trans. Amer. Math. Soc.*, **341** (1), 193-225, 1994. MR **94c**:03066
6. E.G. Effros, Transformation groups and C^* -algebras, *Ann. of Math.*, **81**, 38-55, 1965. MR **30**:5175
7. J. Feldman, P. Hahn, and C.C. Moore, Orbit structure and countable sections for actions of continuous groups, *Adv. in Math.*, **26**, 186-230, 1979. MR **58**:1121
8. J. Feldman and C.C. Moore, Ergodic equivalence relations and von Neumann algebras, I, *Trans. Amer. Math. Soc.*, **234**, 289-324, 1977. MR **58**:28261a
9. H. Friedman and L. Stanley, A Borel reducibility theory for classes of countable structures, *J. Symb. Logic*, **54** (3), 894-914, 1989. MR **91f**:03062
10. L. Fuchs, *Infinite Abelian Groups*, Academic Press, 1970. MR **41**:333
11. S. Gao, Some applications of the Adams-Kechris technique, preprint, 2000.
12. J. Glimm, Locally compact transformation groups, *Trans. Amer. Math. Soc.*, **101**, 124-138, 1961. MR **25**:146
13. V.V. Gorbatsevich, A.L. Onishchik, and E.B. Vinberg, *Foundations of Lie Theory and Lie Transformation Groups*, Springer, 1997. MR **99c**:22009
14. L. Harrington, A.S. Kechris, and A. Louveau, A Glimm-Effros dichotomy for Borel equivalence relations, *J. Amer. Math. Soc.*, **3** (4), 903-928, 1990. MR **91h**:28023
15. G. Hjorth, Around nonclassifiability for countable torsion-free abelian groups, preprint, 1998.
16. G. Hjorth and A.S. Kechris, Borel equivalence relations and classifications of countable models, *Ann. Pure and Appl. Logic*, **82**, 221-272, 1996. MR **99m**:03073
17. J.E. Humphreys, *Linear Algebraic Groups*, Grad. Texts in Math., **21**, Springer-Verlag, 1975. MR **53**:633
18. S. Jackson, A.S. Kechris, and A. Louveau, Countable Borel equivalence relations, preprint, 2000.
19. A.S. Kechris, The structure of Borel equivalence relations in Polish spaces, *Set Theory of the Continuum*, H. Judah, W. Just, and W.H. Woodin (eds.), MSRI Publications, Springer-Verlag, **26**, 89-102, 1992. MR **94h**:03093
20. A.S. Kechris, Countable sections for locally compact group actions, *Erg. Th. and Dyn. Syst.*, **12**, 283-295, 1992. MR **94b**:22003
21. A.S. Kechris, *Classical Descriptive Set Theory*, Grad. Texts in Math., **156**, Springer-Verlag, 1995. MR **96e**:03057
22. A.S. Kechris, Actions of Polish groups and classification problems, preprint, 1998.
23. A.S. Kechris, New directions in descriptive set theory, *Bull. Symb. Logic*, **5** (2), 161-174, 1999.
24. A.S. Kechris, Descriptive dynamics, *Descriptive Set Theory and Dynamical Systems*, Ed. by M. Foreman, A.S. Kechris, A. Louveau, and B. Weiss, London Math. Society Lecture Note Series, **277**, 231-258, Cambridge Univ. Press, 2000.
25. A. Louveau and B. Velickovic, A note on Borel equivalence relations, *Proc. Amer. Math. Soc.*, **120**, 255-259, 1994. MR **94f**:54076
26. G.A. Margulis, *Discrete Subgroups of Semisimple Lie Groups*, Erg. der Math. und Ihrer. Grenz., **17**, Springer-Verlag, 1991. MR **92h**:22021
27. C.C. Moore, Ergodic theory and von Neumann algebras, *Proc. Symp. Pure Math.*, Amer. Math. Soc., **38**, 179-226, 1982. MR **84f**:22013
28. Y.N. Moschovakis, *Descriptive Set Theory*, North Holland, 1980. MR **82e**:03002
29. J.J. Rotman, *An Introduction to the Theory of Groups*, Grad. Texts in Math., **148**, Springer-Verlag, 1995. MR **95m**:20001
30. K. Schmidt, *Algebraic Ideas in Ergodic Theory*, CBMS Regional Conference Series in Math., **76**, Amer. Math Soc., 1990. MR **92k**:28029
31. J.-P. Serre, *A Course in Arithmetic*, Grad. Texts. in Math., **7**, Springer-Verlag, 1973. MR **49**:8956
32. J. Silver, Counting the number of equivalence classes of Borel and co-analytic equivalence relations, *Ann. Math. Logic*, **18**, 1-28, 1980. MR **81d**:03051
33. S. Thomas, Notes on TFA₂ groups, preprint, 1998.

34. V.S. Varadarajan, Groups of automorphisms of Borel spaces, *Trans. Amer. Math. Soc.*, **109**, 191-220, 1963. MR **28**:3139
35. R. Zimmer, *Ergodic Theory and Semisimple Groups*, Birkhäuser, 1984. MR **86j**:22014
36. R. Zimmer, Groups generating transversals to semisimple Lie group actions, *Israel J. Math.*, **73 (2)**, 151-159, 1991. MR **93e**:22012

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA, MINNEAPOLIS, MINNESOTA 55455
E-mail address: `adams@math.umn.edu`

DEPARTMENT OF MATHEMATICS, CALTECH, PASADENA, CALIFORNIA 91125
E-mail address: `kechris@caltech.edu`