Linearized alternating direction method with parallel splitting and adaptive penalty for separable convex programs in machine learning

Zhouchen Lin · Risheng Liu · Huan Li

Received: 14 January 2014 / Accepted: 3 October 2014 / Published online: 20 November 2014 © The Author(s) 2014

Abstract Many problems in machine learning and other fields can be (re)formulated as linearly constrained separable convex programs. In most of the cases, there are multiple blocks of variables. However, the traditional alternating direction method (ADM) and its linearized version (LADM, obtained by linearizing the quadratic penalty term) are for the two-block case and cannot be naively generalized to solve the multi-block case. So there is great demand on extending the ADM based methods for the multi-block case. In this paper, we propose LADM with parallel splitting and adaptive penalty (LADMPSAP) to solve multi-block separable convex programs efficiently. When all the component objective functions have bounded subgradients, we obtain convergence results that are stronger than those of ADM and LADM, e.g., allowing the penalty parameter to be *unbounded* and proving the sufficient and necessary conditions for global convergence. We further propose a simple optimality measure and reveal the convergence rate of LADMPSAP in an ergodic sense. For programs with extra convex set constraints, with refined parameter estimation we devise a practical version of LADMPSAP for faster convergence. Finally, we generalize LADMPSAP to handle programs with more difficult objective functions by linearizing part of the objective function as well. LADMPSAP is particularly suitable for sparse representation and low-rank recovery problems because its subproblems have closed form solutions and the sparsity and low-rankness of the iterates can be preserved during the iteration. It is also highly parallelizable and hence fits for parallel or

Z. Lin

R. Liu (⊠) School of Software Technology, Dalian University of Technology, Dalian, China e-mail: rsliu@dlut.edu.cn

H. Li School of Software and Microelectronics, Peking University, Beijing, China e-mail: lihuan_ss@pku.edu.cn

Editors: Cheng Soon Ong, Tu Bao Ho, Wray Buntine, Bob Williamson, and Masashi Sugiyama.

Key Laboratory of Machine Perception (MOE), School of EECS, Peking University, Beijing, China e-mail: zlin@pku.edu.cn

distributed computing. Numerical experiments testify to the advantages of LADMPSAP in speed and numerical accuracy.

Keywords Convex programs · Alternating direction method · Linearized alternating direction method · Proximal alternating direction method · Parallel splitting · Adaptive renalty

1 Introduction

In recent years, convex programs have become increasingly popular for solving a wide range of problems in machine learning and other fields, ranging from theoretical modeling, e.g., latent variable graphical model selection (Chandrasekaran et al. 2012), low-rank feature extraction [e.g., matrix decomposition (Candès et al. 2011) and matrix completion (Candès and Recht 2009)], subspace clustering (Liu et al. 2012), and kernel discriminant analysis (Ye et al. 2008), to real-world applications, e.g., face recognition (Wright et al. 2009), saliency detection (Shen and Wu 2012), and video denoising (Ji et al. 2010). Most of the problems can be (re)formulated as the following linearly constrained separable convex program¹:

$$\min_{\mathbf{x}_1,\dots,\mathbf{x}_n} \sum_{i=1}^n f_i(\mathbf{x}_i), \quad s.t. \quad \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i) = \mathbf{b},\tag{1}$$

where \mathbf{x}_i and \mathbf{b} could be either vectors or matrices,² f_i is a closed proper convex function, and $\mathcal{A}_i : \mathbb{R}^{d_i} \to \mathbb{R}^m$ is a linear mapping. Without loss of generality, we may assume that none of the \mathcal{A}_i 's is a zero mapping, the solution to $\sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i) = \mathbf{b}$ is non-unique, and the mapping $\mathcal{A}(\mathbf{x}_1, \dots, \mathbf{x}_n) \equiv \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i)$ is onto³.

1.1 Exemplar problems in machine learning

In this subsection, we present some examples of machine learning problems that can be formulated as the model problem (1).

1.1.1 Latent low-rank representation

Low-rank representation (LRR) (Liu et al. 2010, 2012) is a recently proposed technique for robust subspace clustering and has been applied to many machine learning and computer vision problems. However, LRR works well only when the number of samples is more than the dimension of the samples, which may not be satisfied when the data dimension is high.

¹ If the objective function is not separable or there are extra convex set constraints, $\mathbf{x}_i \in X_i$, i = 1, ..., n, where X_i 's are convex sets, the program can be transformed into (1) by introducing auxiliary variables, c.f. (26)–(28).

² In this paper we call each \mathbf{x}_i a "block" of variables because it may consist of multiple scalar variables. We will use bold capital letters if a block is known to be a matrix.

³ The last two assumptions are equivalent to that the matrix $\mathbf{A} \equiv (\mathbf{A}_1 \dots \mathbf{A}_n)$ is not full column rank but full row rank, where \mathbf{A}_i is the matrix representation of \mathcal{A}_i .

So Liu and Yan (2011) proposed latent LRR to overcome this difficulty. The mathematical model of latent LRR is as follows:

$$\min_{\mathbf{Z},\mathbf{L},\mathbf{E}} \|\mathbf{Z}\|_* + \|\mathbf{L}\|_* + \mu \|\mathbf{E}\|_1, \quad s.t. \quad \mathbf{X} = \mathbf{X}\mathbf{Z} + \mathbf{L}\mathbf{X} + \mathbf{E},$$
(2)

where **X** is the data matrix, each column being a sample vector, $\|\cdot\|_*$ is the nuclear norm (Fazel 2002), i.e., the sum of singular values, and $\|\cdot\|_1$ is the ℓ_1 norm (Candès et al. 2011), i.e., the sum of absolute values of all entries. Latent LRR is to decompose data into principal feature **XZ** and salient feature **LX**, up to sparse noise **E**.

1.1.2 Nonnegative matrix completion

Nonnegative matrix completion (NMC) (Xu et al. 2011) is a novel technique for dimensionality reduction, text mining, collaborative filtering, and clustering, etc. It can be formulated as:

$$\min_{\mathbf{X},\mathbf{e}} \|\mathbf{X}\|_* + \frac{1}{2\mu} \|\mathbf{e}\|^2, \quad s.t. \quad \mathbf{b} = \mathcal{P}_{\Omega}(\mathbf{X}) + \mathbf{e}, \ \mathbf{X} \ge 0,$$
(3)

where **b** is the observed data in the matrix **X** contaminated by noise **e**, Ω is an index set, \mathcal{P}_{Ω} is a linear mapping that selects those elements whose indices are in Ω , and $\|\cdot\|$ is the Frobenius norm. NMC is to recover the nonnegative low-rank matrix **X** from the observed noisy data **b**.

To see that the NMC problem can be reformulated as (1), we introduce an auxiliary variable **Y** and rewrite (3) as

$$\min_{\mathbf{X},\mathbf{Y},\mathbf{e}} \|\mathbf{X}\|_* + \chi_{\geq 0}(\mathbf{Y}) + \frac{1}{2\mu} \|\mathbf{e}\|^2, \quad s.t. \quad \begin{pmatrix} \mathcal{P}_{\Omega}(\mathbf{X}) \\ \mathbf{X} \end{pmatrix} - \begin{pmatrix} \mathbf{0} \\ \mathbf{Y} \end{pmatrix} + \begin{pmatrix} \mathbf{e} \\ \mathbf{0} \end{pmatrix} = \begin{pmatrix} \mathbf{b} \\ \mathbf{0} \end{pmatrix}, \quad (4)$$

where

$$\chi_{\geq 0}(\mathbf{Y}) = \begin{cases} 0, & \text{if } \mathbf{Y} \geq 0, \\ +\infty, & \text{otherwise,} \end{cases}$$

is the characteristic function of the set of nonnegative matrices.

1.1.3 Group sparse logistic regression with overlap

Besides unsupervised learning models shown above, many supervised machine learning problems can also be written in the form of (1). For example, using logistic function as the loss function in the group LASSO with overlap (Jacob et al. 2009; Deng et al. 2011), one obtains the following model:

$$\min_{\mathbf{w},b} \frac{1}{s} \sum_{i=1}^{s} \log\left(1 + \exp\left(-y_i(\mathbf{w}^T \mathbf{x}_i + b)\right)\right) + \mu \sum_{j=1}^{t} \|\mathbf{S}_j \mathbf{w}\|,\tag{5}$$

where \mathbf{x}_i and y_i , i = 1, ..., s, are the training data and labels, respectively, and \mathbf{w} and b parameterize the linear classifier. \mathbf{S}_j , j = 1, ..., t, are the selection matrices, with only one 1 at each row and the rest entries are all zeros. The groups of entries, $\mathbf{S}_j \mathbf{w}$, j = 1, ..., t, may overlap each other. This model can also be considered as an extension of the group sparse logistic regression problem (Meier et al. 2008) to the case of overlapped groups.

Introducing $\mathbf{\bar{w}} = (\mathbf{w}^T, b)^T, \mathbf{\bar{x}}_i = (\mathbf{x}_i^T, 1)^T, \mathbf{z} = (\mathbf{z}_1^T, \mathbf{z}_2^T, \dots, \mathbf{z}_t^T)^T$, and $\mathbf{\bar{S}} = (\mathbf{S}, \mathbf{0})$, where $\mathbf{S} = (\mathbf{S}_1^T, \dots, \mathbf{S}_t^T)^T$, (5) can be rewritten as

$$\min_{\bar{\mathbf{w}},\mathbf{z}} \frac{1}{s} \sum_{i=1}^{s} \log\left(1 + \exp\left(-y_i(\bar{\mathbf{w}}^T \bar{\mathbf{x}}_i)\right)\right) + \mu \sum_{j=1}^{t} \|\mathbf{z}_j\|, \quad s.t. \quad \mathbf{z} = \bar{\mathbf{S}}\bar{\mathbf{w}}, \tag{6}$$

which is a special case of (1).

1.2 Related work

Although general theories on convex programs are fairly complete nowadays, e.g., most of them can be solved by the interior point method (Boyd and Vandenberghe 2004), when faced with large scale problems, which are typical in machine learning, the general theory may not lead to efficient algorithms. For example, when using CVX,⁴ an interior point based toolbox, to solve nuclear norm minimization problems [i.e., one of the f_i 's is the nuclear norm of a matrix, e.g., (2) and (3)], such as matrix completion (Candès and Recht 2009), robust principal component analysis (Candès et al. 2011), and low-rank representation (Liu et al. 2010, 2012), the complexity of each iteration is $O(q^6)$, where $q \times q$ is the matrix size. Such a complexity is unbearable for large scale computing.

To address the scalability issue, first order methods are often preferred. The accelerated proximal gradient (APG) algorithm (Beck and Teboulle 2009; Toh and Yun 2010) is popular due to its guaranteed $O(K^{-2})$ convergence rate, where *K* is the iteration number. However, APG is basically for unconstrained optimization. For constrained optimization, the constraints have to be added to the objective function as penalties, resulting in approximated solutions only. The alternating direction method (ADM)⁵ (Fortin and Glowinski 1983; Boyd et al. 2011; Lin et al. 2009a) has regained a lot of attention recently and is also widely used. It is especially suitable for separable convex programs like (1) because it fully utilizes the separable structure of the objective function. Unlike APG, ADM can solve (1) exactly. Another first order method is the split Bregman method (Goldstein and Osher 2008; Zhang et al. 2011), which is closely related to ADM (Esser 2009) and is influential in image processing.

An important reason that first order methods are popular for solving large scale convex programs in machine learning is that the convex functions f_i 's are often matrix or vector norms or characteristic functions of convex sets, which enables the following subproblems [called the proximal operation of f_i (Rockafellar 1970)]

$$\operatorname{prox}_{f_i,\sigma}(\mathbf{w}) = \operatorname{argmin}_{\mathbf{x}_i} f_i(\mathbf{x}_i) + \frac{\sigma}{2} \|\mathbf{x}_i - \mathbf{w}\|^2$$
(7)

to have closed form solutions. For example, when f_i is the ℓ_1 norm, $\operatorname{prox}_{f_i,\sigma}(\mathbf{w}) = \mathcal{T}_{\sigma^{-1}}(\mathbf{w})$, where $\mathcal{T}_{\varepsilon}(x) = \operatorname{sgn}(x) \max(|x| - \varepsilon, 0)$ is the soft-thresholding operator (Goldstein and Osher 2008); when f_i is the nuclear norm, the optimal solution is: $\operatorname{prox}_{f_i,\sigma}(\mathbf{W}) = \mathbf{U}\mathcal{T}_{\sigma^{-1}}(\Sigma)\mathbf{V}^T$, where $\mathbf{U}\Sigma\mathbf{V}^T$ is the singular value decomposition (SVD) of \mathbf{W} (Cai et al. 2010); and when f_i is the characteristic function of the nonnegative cone, the optimal solution is $\operatorname{prox}_{f_i,\sigma}(\mathbf{w}) = \max(\mathbf{w}, 0)$. Since subproblems like (7) have to be solved in each iteration when using first order methods to solve separable convex programs, that they have closed form solutions greatly facilitates the optimization.

⁴ Available at http://stanford.edu/~boyd/cvx.

⁵ Also called the alternating direction method of multipliers (ADMM) in some literatures, e.g., (Boyd et al. 2011; Zhang et al. 2011; Deng and Yin 2012).

or full-rankness on some linear operators.

However, when applying ADM to solve (1) with non-unitary linear mappings (i.e., $A_i^{\dagger} A_i$ is not the identity mapping, where A_i^{\dagger} is the adjoint operator of A_i), the resulting subproblems may not have closed form solutions,⁶ hence need to be solved iteratively, making the optimization process awkward. Some work (Yang and Yuan 2013; Lin et al. 2011) has considered this issue by linearizing the quadratic term $||A_i(\mathbf{x}_i) - \mathbf{w}||^2$ in the subproblems, hence such a variant of ADM is called the linearized ADM (LADM). Deng and Yin (2012) further propose the generalized ADM that makes both ADM and LADM as its special cases and prove its globally linear convergence by imposing strong convexity on the objective function

Nonetheless, most of the existing theories on ADM and LADM are for the *two-block* case, i.e., n = 2 in (1) (Fortin and Glowinski 1983; Boyd et al. 2011; Lin et al. 2011; Deng and Yin 2012). The number of blocks is restricted to two because the proofs of convergence for the two-block case are not applicable for the multi-block case, i.e., n > 2 in (1). Actually, a naive generalization of ADM or LADM to the multi-block case may diverge [see (15) and Chen et al. 2013]. Unfortunately, in practice multi-block convex programs often occur, e.g., robust principal component analysis with dense noise (Candès et al. 2011), latent low-rank representation (Liu and Yan 2011) [see (2)], and when there are extra convex set constraints [see (3) and (26)–(27)]. So it is desirable to design practical algorithms for the multi-block case.

Recently He and Yuan (2013) and Tao (2014) considered the multi-block LADM and ADM, respectively. To safeguard convergence, He and Yuan (2013) proposed LADM with Gaussian back substitution (LADMGB), which destroys the sparsity or low-rankness of the iterates during iterations when dealing with sparse representation and low-rank recovery problems, while Tao (2014) proposed ADM with parallel splitting, whose subproblems may not be easily solvable. Moreover, they all developed their theories with the penalty parameter being fixed, resulting in difficulty of tuning an optimal penalty parameter that fits for different data and data sizes. This has been identified as an important issue (Deng and Yin 2012).

1.3 Contributions and differences from prior work

To propose an algorithm that is more suitable for convex programs in machine learning, in this paper we aim at combining the advantages of He and Yuan (2013), Tao (2014), and Lin et al. (2011), i.e., combining LADM, parallel splitting, and adaptive penalty. Hence we call our method LADM with parallel splitting and adaptive penalty (LADMPSAP). With LADM, the subproblems will have forms like (7) and hence can be easily solved. With parallel splitting, the sparsity and low-rankness of iterates can be preserved during iterations when dealing with sparse representation and low-rank recovery problems, saving both the storage and the computation load. With adaptive penalty, the convergence can be faster and it is unnecessary to tune an optimal penalty parameter. Parallel splitting also makes the algorithm highly parallelizable, making LADMPSAP suitable for parallel or distributed computing, which is important for large scale machine learning. When all the component objective functions have bounded subgradients, we prove convergence results that are stronger than the existing theories on ADM and LADM. For example, the penalty parameter can be unbounded and the sufficient and necessary conditions of the global convergence of LADMPSAP can be obtained as well. We also propose a simple optimality measure and prove the convergence rate of LADMPSAP in an ergodic sense under this measure. Our proof is simpler than those in He and Yuan (2012) and Tao (2014) which relied on a complex optimality measure. When

⁶ Because $\|\mathbf{x}_i - \mathbf{w}\|^2$ in (7) becomes $\|\mathcal{A}_i(\mathbf{x}_i) - \mathbf{w}\|^2$, which cannot be reduced to $\|\mathbf{x}_i - \tilde{\mathbf{w}}\|^2$.

a convex program has extra convex set constraints, we further devise a practical version of LADMPSAP that converges faster thanks to better parameter analysis. Finally, we generalize LADMPSAP to cope with more difficult f_i 's, whose proximal operation (7) is not easily solvable, by further linearizing the smooth components of f_i 's. Experiments testify to the advantage of LADMPSAP in speed and numerical accuracy.

Note that Goldfarb and Ma (2012) also proposed a multiple splitting algorithm for convex optimization. However, they only considered a special case of our model problem (1), i.e., all the linear mappings A_i 's are identity mappings.⁷ With their simpler model problem, linearization is unnecessary and a faster convergence rate, $O(K^{-2})$, can be achieved. In contrast, in this paper we aim at proposing a practical algorithm for efficiently solving more general problems like (1).

We also note that Hong and Luo (2012) used the same linearization technique for the smooth components of f_i 's as well, but they only considered a special class of f_i 's. Namely, the non-smooth component of f_i is a sum of ℓ_1 and ℓ_2 norms or its epigraph is polyhedral. Moreover, for parallel splitting (Jacobi update) Hong and Luo (2012) has to incorporate a postprocessing to guarantee convergence, by interpolating between an intermediate iterate and the previous iterate. Third, Hong and Luo (2012) still focused on a fixed penalty parameter. Again, our method can handle more general f_i 's, does not require postprocessing, and allows for an adaptive penalty parameter.

A more general splitting/linearization technique can be founded in Zhang et al. (2011). However, the authors only proved that any accumulation point of the iteration is a Kuhn– Karush–Tucker (KKT) point and did not investigate the convergence rate. There was no evidence that the iteration could converge to a unique point. Moreover, the authors only studied the case of fixed penalty parameter.

Although dual ascent with dual decomposition (Boyd et al. 2011) can also solve (1) in a parallel way, it may break down when some f_i 's are not strictly convex (Boyd et al. 2011), which typically happens in sparse or low-rank recovery problems where ℓ_1 norm or nuclear norm are used. Even if it works, since f_i is not strictly convex, dual ascent becomes dual *subgradient* ascent (Boyd et al. 2011), which is known to converge at a rate of $O(K^{-1/2})$ —slower than our $O(K^{-1})$ rate. Moreover, dual ascent requires choosing a good step size for each iteration, which is less convenient than ADM based methods.

1.4 Organization

The remainder of this paper is organized as follows. We first review LADM with adaptive penalty (LADMAP) for the two-block case in Sect. 2. Then we present LADMPSAP for the multi-block case in Sect. 3. Next, we propose a practical version of LADMPSAP for separable convex programs with convex set constraints in Sect. 4. We further extend LADMPSAP to proximal LADMPSAP for programs with more difficult objective functions in Sect. 5. We compare the advantage of LADMPSAP in speed and numerical accuracy with other first order methods in Sect. 6. Finally, we conclude the paper in Sect. 7.

This paper is an extension of our prior work Lin et al. (2011) and Liu et al. (2013).

2 Review of LADMAP for the two-block case

We first review LADMAP (Lin et al. 2011) for the two-block case of (1). It consists of four steps:

⁷ The multi-block problems introduced in Boyd et al. (2011) also fall within this category.

1. Update \mathbf{x}_1 :

$$\mathbf{x}_{1}^{k+1} = \underset{\mathbf{x}_{1}}{\operatorname{argmin}} f_{1}(\mathbf{x}_{1}) + \frac{\sigma_{1}^{(k)}}{2} \|\mathbf{x}_{1} - \mathbf{x}_{1}^{k} + \mathcal{A}_{1}^{\dagger}\left(\tilde{\boldsymbol{\lambda}}_{1}^{k}\right) / \sigma_{1}^{(k)} \|^{2},$$
(8)

2. Update \mathbf{x}_2 :

$$\mathbf{x}_{2}^{k+1} = \underset{\mathbf{x}_{2}}{\operatorname{argmin}} \ f_{2}(\mathbf{x}_{2}) + \frac{\sigma_{2}^{(k)}}{2} \left\| \mathbf{x}_{2} - \mathbf{x}_{2}^{k} + \mathcal{A}_{2}^{\dagger} \left(\tilde{\boldsymbol{\lambda}}_{2}^{k} \right) / \sigma_{2}^{(k)} \right\|^{2}, \tag{9}$$

3. Update λ :

$$\boldsymbol{\lambda}^{k+1} = \boldsymbol{\lambda}^k + \beta_k \left(\sum_{i=1}^2 \mathcal{A}_i \left(\mathbf{x}_i^{k+1} \right) - \mathbf{b} \right), \tag{10}$$

4. Update β :

$$\beta_{k+1} = \min(\beta_{\max}, \rho\beta_k), \tag{11}$$

where λ is the Lagrange multiplier, β_k is the penalty parameter, $\sigma_i^{(k)} = \eta_i \beta_k$ with $\eta_i > ||\mathcal{A}_i||^2$ ($||\mathcal{A}_i||$ is the operator norm of \mathcal{A}_i),

$$\tilde{\boldsymbol{\lambda}}_{1}^{k} = \boldsymbol{\lambda}^{k} + \beta_{k} \left(\mathcal{A}_{1} \left(\mathbf{x}_{1}^{k} \right) + \mathcal{A}_{2} \left(\mathbf{x}_{2}^{k} \right) - \mathbf{b} \right),$$
(12)

$$\tilde{\boldsymbol{\lambda}}_{2}^{k} = \boldsymbol{\lambda}^{k} + \beta_{k} \left(\mathcal{A}_{1} \left(\mathbf{x}_{1}^{k+1} \right) + \mathcal{A}_{2} \left(\mathbf{x}_{2}^{k} \right) - \mathbf{b} \right), \tag{13}$$

and ρ is an adaptively updated parameter [see (20)]. Please refer to (Lin et al. 2011) for details. Note that the latest \mathbf{x}_1^{k+1} is immediately used to compute \mathbf{x}_2^{k+1} [see (13)]. So \mathbf{x}_1 and \mathbf{x}_2 have to be updated alternately, hence the name alternating direction method.

3 LADMPSAP for the multi-block case

In this section, we extend LADMAP for multi-block separable convex programs (1). We also provide the *sufficient and necessary conditions* for global convergence when subgradients of the objective functions are all bounded. We further prove the convergence rate in an ergodic sense.

3.1 LADM with parallel splitting and adaptive penalty

Contrary to our intuition, the multi-block case is actually fundamentally different from the two-block one. For the multi-block case, it is very natural to generalize LADMAP for the two-block case in a straightforward way, with

$$\tilde{\boldsymbol{\lambda}}_{i}^{k} = \boldsymbol{\lambda}^{k} + \beta_{k} \left(\sum_{j=1}^{i-1} \mathcal{A}_{j} \left(\mathbf{x}_{j}^{k+1} \right) + \sum_{j=i}^{n} \mathcal{A}_{j} \left(\mathbf{x}_{j}^{k} \right) - \mathbf{b} \right), \quad i = 1, \dots, n.$$
(14)

Unfortunately, we were unable to prove the convergence of such a naive LADMAP using the same proof for the two-block case. This is because their Fejér monotone inequalities (see Remark 4) cannot be the same. That is why He et al. has to introduce an extra Gaussian back substitution (He et al. 2012; He and Yuan 2013) for correcting the iterates. Actually,

the above naive generalization of LADMAP may be divergent (which is even worse than converging to a wrong solution), e.g., when applied to the following problem:

$$\min_{\mathbf{x}_{1},...,\mathbf{x}_{n}} \sum_{i=1}^{n} \|\mathbf{x}_{i}\|_{1}, \quad s.t. \quad \sum_{i=1}^{n} \mathbf{A}_{i} \mathbf{x}_{i} = \mathbf{b},$$
(15)

where $n \ge 5$ and \mathbf{A}_i and \mathbf{b} are Gaussian random matrix and vector, respectively, whose entries fulfil the standard Gaussian distribution independently. Chen et al. (2013) also analyzed the naively generalized ADM for the multi-block case and showed that even for three blocks the iteration could still be divergent. They also provided sufficient conditions, which basically require that the linear mappings \mathcal{A}_i should be orthogonal to each other $(\mathcal{A}_i^{\dagger} \mathcal{A}_j = 0, i \neq j)$, to ensure the convergence of naive ADM.

Fortunately, by modifying $\tilde{\lambda}_i^k$ slightly we are able to prove the convergence of the corresponding algorithm. More specifically, our algorithm for solving (1) consists of the following steps:

1. Update \mathbf{x}_i 's in parallel:

$$\mathbf{x}_{i}^{k+1} = \underset{\mathbf{x}_{i}}{\operatorname{argmin}} f_{i}(\mathbf{x}_{i}) + \frac{\sigma_{i}^{(k)}}{2} \left\| \mathbf{x}_{i} - \mathbf{x}_{i}^{k} + \mathcal{A}_{i}^{\dagger}\left(\hat{\boldsymbol{\lambda}}^{k}\right) / \sigma_{i}^{(k)} \right\|^{2}, \quad i = 1, \dots, n, \quad (16)$$

2. Update λ :

$$\boldsymbol{\lambda}^{k+1} = \boldsymbol{\lambda}^k + \beta_k \left(\sum_{i=1}^n \mathcal{A}_i \left(\mathbf{x}_i^{k+1} \right) - \mathbf{b} \right), \tag{17}$$

3. Update β :

$$\beta_{k+1} = \min(\beta_{\max}, \rho\beta_k), \tag{18}$$

where $\sigma_i^{(k)} = \eta_i \beta_k$,

$$\hat{\boldsymbol{\lambda}}^{k} = \boldsymbol{\lambda}^{k} + \beta_{k} \left(\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k} \right) - \mathbf{b} \right), \tag{19}$$

and

$$\rho = \begin{cases} \rho_0, & \text{if } \beta_k \max\left(\left\{\sqrt{\eta_i} \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\|, i = 1, \dots, n\right\}\right) / \|\mathbf{b}\| < \varepsilon_2, \\ 1, & \text{otherwise,} \end{cases}$$
(20)

with $\rho_0 > 1$ being a constant and $0 < \varepsilon_2 \ll 1$ being a threshold. Indeed, we replace $\tilde{\lambda}_i^{\kappa}$ with $\hat{\lambda}^k$ as (19), which is independent of *i*, and the rest procedures of the algorithm, including the scheme (18) and (20) to update the penalty parameter, are all inherited from Lin et al. (2011), except that η_i 's have to be made larger (see Theorem 1). As now \mathbf{x}_i 's are updated in parallel and β_k changes adaptively, we call the new algorithm LADM with *parallel splitting* and *adaptive penalty* (LADMPSAP).

Algorithm 1 LADMPSAP for Solving (1)

Initialize: Set $\rho_0 > 1$, $\varepsilon_1 > 0$, $\varepsilon_2 > 0$, $\beta_{\max} \gg 1 \gg \beta_0 > 0$, λ^0 , $\eta_i > n \|\mathcal{A}_i\|^2$, \mathbf{x}_i^0 , i = 1, ..., n. **while** (21) or (22) is not satisfied **do Step 1:** Compute $\hat{\boldsymbol{\lambda}}^k$ as (19). **Step 2:** Update \mathbf{x}_i 's in parallel by solving

$$\mathbf{x}_{i}^{k+1} = \underset{\mathbf{x}_{i}}{\operatorname{argmin}} f_{i}(\mathbf{x}_{i}) + \frac{\eta_{i}\beta_{k}}{2} \left\| \mathbf{x}_{i} - \mathbf{x}_{i}^{k} + \mathcal{A}_{i}^{\dagger}(\hat{\mathbf{\lambda}}^{k}) / (\eta_{i}\beta_{k}) \right\|^{2}, \ i = 1, \dots, n.$$
(23)

Step 3: Update λ by (17) and β by (18) and (20). end while

3.2 Stopping criteria

Some existing work (e.g., Liu et al. 2010; Favaro et al. 2011) proposed stopping criteria out of intuition only, which may not guarantee that the correct solution is approached. Recently, Lin et al. (2009a) and Boyd et al. (2011) suggested that the stopping criteria can be derived from the KKT conditions of a problem. Here we also adopt such a strategy. Specifically, the iteration terminates when the following two conditions are met:

$$\left\|\sum_{i=1}^{n} \mathcal{A}_{i}\left(\mathbf{x}_{i}^{k+1}\right) - \mathbf{b}\right\| / \|\mathbf{b}\| < \varepsilon_{1},$$
(21)

$$\beta_k \max\left(\left\{\sqrt{\eta_i} \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\|, i = 1, \dots, n\right\}\right) / \|\mathbf{b}\| < \varepsilon_2.$$
(22)

The first condition measures the feasibility error. The second condition is derived by comparing the KKT conditions of problem (1) and the optimality condition of subproblem (23). The rules (18) and (20) for updating β are actually hinted by the above stopping criteria such that the two errors are well balanced.

For better reference, we summarize the proposed LADMPSAP algorithm in Algorithm 1. For fast convergence, we suggest that $\beta_0 = \alpha m \varepsilon_2$ and $\alpha > 0$ and $\rho_0 > 1$ should be chosen such that β_k increases steadily along with iterations.

3.3 Global convergence

In the following, we always use $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*, \boldsymbol{\lambda}^*)$ to denote the KKT point of problem (1). For the global convergence of LADMPSAP, we have the following theorem, where we denote $\{\mathbf{x}_i^k\} = \{\mathbf{x}_1^k, \dots, \mathbf{x}_n^k\}$ for simplicity.

Theorem 1 (Convergence of LADMPSAP)⁸ If $\{\beta_k\}$ is non-decreasing and upper bounded, $\eta_i > n \|\mathcal{A}_i\|^2$, i = 1, ..., n, then $\{(\{\mathbf{x}_i^k\}, \boldsymbol{\lambda}^k)\}$ generated by LADMPSAP converge to a KKT point of problem (1).

3.4 Enhanced convergence results

Theorem 1 is a convergence result for general convex programs (1), where f_i 's are general convex functions and hence $\{\beta_k\}$ needs to be bounded. Actually, almost all the existing theories on ADM and LADM even assumed a fixed β . For adaptive β_k , it will be more convenient if a user needs not to specify an upper bound on $\{\beta_k\}$ because imposing a large

⁸ Please see "Appendix" for all the proofs of our theoretical results hereafter.

upper bound essentially equals to allowing $\{\beta_k\}$ to be unbounded. Since many machine learning problems choose f_i 's as matrix/vector norms, which result in bounded subgradients, we find that the boundedness assumption can be removed. Moreover, we can further prove the *sufficient and necessary* condition for global convergence.

Theorem 2 (Sufficient condition for global convergence) If $\{\beta_k\}$ is non-decreasing and $\sum_{k=1}^{+\infty} \beta_k^{-1} = +\infty, \eta_i > n \|\mathcal{A}_i\|^2, \partial f_i(\mathbf{x})$ is bounded, i = 1, ..., n, then the sequence $\{\mathbf{x}_i^k\}$ generated by LADMPSAP converges to an optimal solution to (1).

Remark 1 Theorem 2 does not claim that $\{\lambda^k\}$ converges to a point λ^{∞} . However, as we are more interested in $\{\mathbf{x}_i^k\}$, such a weakening is harmless.

We also have the following result on the necessity of $\sum_{k=1}^{+\infty} \beta_k^{-1} = +\infty$.

Theorem 3 (Necessary condition for global convergence) If $\{\beta_k\}$ is non-decreasing, $\eta_i > n \|\mathcal{A}_i\|^2$, $\partial f_i(\mathbf{x})$ is bounded, i = 1, ..., n, then $\sum_{k=1}^{+\infty} \beta_k^{-1} = +\infty$ is also a necessary condition for the global convergence of $\{\mathbf{x}_i^k\}$ generated by LADMPSAP to an optimal solution to (1).

With the above analysis, when all the subgradients of the component objective functions are bounded we can remove β_{max} in Algorithm 1.

3.5 Convergence rate

The convergence rate of ADM and LADM in the traditional sense is an open problem (Goldfarb and Ma 2012). Although Hong and Luo (2012) claimed that they proved the linear convergence rate of ADM, their assumptions are actually quite strong. They assumed that the non-smooth part of f_i is a sum of ℓ_1 and ℓ_2 norms or its epigraph is polyhedral. Moreover, the convex constraint sets should all be polyhedral and bounded. So although their results are encouraging, for general convex programs the convergence rate is still a mystery. Recently, He and Yuan (2012) and Tao (2014) proved an O(1/K) convergence rate of ADM and ADM with parallel splitting in an ergodic sense, respectively. Namely $\frac{1}{K} \sum_{k=1}^{K} \mathbf{x}_i$ violates an optimality measure in O(1/K). Their proof is lengthy and is for fixed penalty parameter only.

In this subsection, based on a simple optimality measure we give a simple proof for the convergence rate of LADMPSAP. For simplicity, we denote $\mathbf{x} = (\mathbf{x}_1^T, \dots, \mathbf{x}_n^T)^T, \mathbf{x}^* = ((\mathbf{x}_1^*)^T, \dots, (\mathbf{x}_2^*)^T)^T$, and $f(\mathbf{x}) = \sum_{i=1}^n f_i(\mathbf{x}_i)$. We first have the following proposition.

Proposition 1 $\tilde{\mathbf{x}}$ is an optimal solution to (1) if and only if there exists $\alpha > 0$, such that

$$f(\tilde{\mathbf{x}}) - f(\mathbf{x}^*) + \sum_{i=1}^n \left\langle \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*), \tilde{\mathbf{x}}_i - \mathbf{x}_i^* \right\rangle + \alpha \left\| \sum_{i=1}^n \mathcal{A}_i(\tilde{\mathbf{x}}_i) - \mathbf{b} \right\|^2 = 0.$$
(24)

Since the left hand side of (24) is always nonnegative and it becomes zero only when $\tilde{\mathbf{x}}$ is an optimal solution, we may use its magnitude to measure how far a point $\tilde{\mathbf{x}}$ is from an optimal solution. Note that in the unconstrained case, as in APG (Beck and Teboulle 2009), one may simply use $f(\tilde{\mathbf{x}}) - f(\mathbf{x}^*)$ to measure the optimality. But here we have to deal with the constraints. Our criterion is simpler than that in (He and Yuan 2012; Tao 2014), which has to compare $(\{\mathbf{x}_i^k\}, \lambda^k)$ with all $(\mathbf{x}_1, \dots, \mathbf{x}_n, \boldsymbol{\lambda}) \in \mathbb{R}^{d_1} \times \ldots \times \mathbb{R}^{d_n} \times \mathbb{R}^m$.

Then we have the following convergence rate theorem for LADMPSAP in an ergodic sense.

Theorem 4 (Convergence rate of LADMPSAP) Define $\bar{\mathbf{x}}^K = \sum_{k=0}^K \gamma_k \mathbf{x}^{k+1}$, where $\gamma_k = \beta_k^{-1} / \sum_{j=0}^K \beta_j^{-1}$. Then the following inequality holds for $\bar{\mathbf{x}}^K$:

$$f(\bar{\mathbf{x}}^{K}) - f(\mathbf{x}^{*}) + \sum_{i=1}^{n} \left\langle \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}), \bar{\mathbf{x}}_{i}^{K} - \mathbf{x}_{i}^{*} \right\rangle + \frac{\alpha\beta_{0}}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i}\left(\bar{\mathbf{x}}_{i}^{K}\right) - \mathbf{b} \right\|^{2}$$

$$\leq C_{0} / \left(2 \sum_{k=0}^{K} \beta_{k}^{-1} \right), \qquad (25)$$

where

$$\alpha^{-1} = (n+1) \max\left(1, \left\{\frac{\|\mathcal{A}_i\|^2}{\eta_i - n\|\mathcal{A}_i\|^2}, i = 1, \dots, n\right\}\right)$$

and

$$C_{0} = \sum_{i=1}^{n} \eta_{i} \|\mathbf{x}_{i}^{0} - \mathbf{x}_{i}^{*}\|^{2} + \beta_{0}^{-2} \|\boldsymbol{\lambda}^{0} - \boldsymbol{\lambda}^{*}\|^{2}.$$

Theorem 4 means that $\bar{\mathbf{x}}^K$ is by $O\left(1/\sum_{k=0}^K \beta_k^{-1}\right)$ from being an optimal solution. This theorem holds for both bounded and unbounded $\{\beta_k\}$. In the bounded case, $O\left(1/\sum_{k=0}^K \beta_k^{-1}\right)$ is simply O(1/K). Theorem 4 also hints that $\sum_{k=0}^K \beta_k^{-1}$ should approach infinity to guarantee the convergence of LADMPSAP, which is consistent with Theorem 3.

4 Practical LADMPSAP for convex programs with convex set constraints

In real applications, we are often faced with convex programs with convex set constraints:

$$\min_{\mathbf{x}_{1},...,\mathbf{x}_{n}} \sum_{i=1}^{n} f_{i}(\mathbf{x}_{i}), \ s.t. \ \sum_{i=1}^{n} \mathcal{A}_{i}(\mathbf{x}_{i}) = \mathbf{b}, \quad \mathbf{x}_{i} \in X_{i}, \ i = 1, ..., n,$$
(26)

where $X_i \subseteq \mathbb{R}^{d_i}$ is a closed convex set. In this section, we consider to extend LADMPSAP to solve the more complex convex set constraint model (26). We assume that the projections onto X_i 's are all easily computable. For many convex sets used in machine learning, such an assumption is valid, e.g., when X_i 's are nonnegative cones or positive semi-definite cones. In the following, we discuss how to solve (26) efficiently. For simplicity, we assume $X_i \neq \mathbb{R}^{d_i}$, $\forall i$. Finally, we assume that **b** is an interior point of $\sum_{i=1}^n \mathcal{A}_i(X_i)$.

We introduce auxiliary variables \mathbf{x}_{n+i} to convert $\mathbf{x}_i \in X_i$ into $\mathbf{x}_i = \mathbf{x}_{n+i}$ and $\mathbf{x}_{n+i} \in X_i$, i = 1, ..., n. Then (26) can be reformulated as:

$$\min_{\mathbf{x}_1,...,\mathbf{x}_{2n}} \sum_{i=1}^{2n} f_i(\mathbf{x}_i), \quad s.t. \quad \sum_{i=1}^{2n} \hat{\mathcal{A}}_i(\mathbf{x}_i) = \hat{\mathbf{b}},$$
(27)

where

$$f_{n+i}(\mathbf{x}) \equiv \chi_{X_i}(\mathbf{x}) = \begin{cases} 0, & \text{if } \mathbf{x} \in X_i, \\ +\infty, & \text{otherwise,} \end{cases}$$

Springer

is the characteristic function of X_i ,

$$\hat{\mathcal{A}}_{i}(\mathbf{x}_{i}) = \begin{pmatrix} \mathcal{A}_{i}(\mathbf{x}_{i}) \\ 0 \\ \vdots \\ \mathbf{x}_{i} \\ \vdots \\ 0 \end{pmatrix}, \quad \hat{\mathcal{A}}_{n+i}(\mathbf{x}_{n+i}) = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ -\mathbf{x}_{n+i} \\ \vdots \\ 0 \end{pmatrix}, \quad \text{and} \quad \hat{\mathbf{b}} = \begin{pmatrix} \mathbf{b} \\ 0 \\ \vdots \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \quad (28)$$

where i = 1, ..., n.

The adjoint operator \hat{A}_{i}^{\dagger} is

$$\hat{\mathcal{A}}_{i}^{\dagger}(\mathbf{y}) = \mathcal{A}_{i}^{\dagger}(\mathbf{y}_{1}) + \mathbf{y}_{i+1}, \quad \hat{\mathcal{A}}_{n+i}^{\dagger}(\mathbf{y}) = -\mathbf{y}_{i+1}, \quad i = 1, \dots, n,$$
(29)

where \mathbf{y}_i is the *i*th sub-vector of \mathbf{y} , partitioned according to the sizes of \mathbf{b} and \mathbf{x}_i , $i = 1, \dots, n$.

Then LADMPSAP can be applied to solve problem (27). The Lagrange multiplier λ and the auxiliary multiplier $\hat{\lambda}$ are respectively updated as

$$\boldsymbol{\lambda}_{1}^{k+1} = \boldsymbol{\lambda}_{1}^{k} + \beta_{k} \left(\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} \right) - \mathbf{b} \right), \quad \boldsymbol{\lambda}_{i+1}^{k+1} = \boldsymbol{\lambda}_{i+1}^{k} + \beta_{k} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{n+i}^{k+1} \right), \quad (30)$$

$$\hat{\boldsymbol{\lambda}}_{1}^{k} = \boldsymbol{\lambda}_{1}^{k} + \beta_{k} \left(\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k} \right) - \mathbf{b} \right), \quad \hat{\boldsymbol{\lambda}}_{i+1}^{k} = \boldsymbol{\lambda}_{i+1}^{k} + \beta_{k} \left(\mathbf{x}_{i}^{k} - \mathbf{x}_{n+i}^{k} \right), \tag{31}$$

and \mathbf{x}_i is updated as (see 16)

$$\mathbf{x}_{i}^{k+1} = \underset{\mathbf{x}}{\operatorname{argmin}} \quad f_{i}(\mathbf{x}) + \frac{\eta_{i}\beta_{k}}{2} \left\| \mathbf{x} - \mathbf{x}_{i}^{k} + \left[\mathcal{A}_{i}^{\dagger} \left(\hat{\boldsymbol{\lambda}}_{1}^{k} \right) + \hat{\boldsymbol{\lambda}}_{i+1}^{k} \right] / (\eta_{i}\beta_{k}) \right\|^{2}, \quad (32)$$

$$\mathbf{x}_{n+i}^{k+1} = \underset{\mathbf{x}\in X_i}{\operatorname{argmin}} \frac{\eta_{n+i}\beta_k}{2} \left\| \mathbf{x} - \mathbf{x}_{n+i}^k - \hat{\boldsymbol{\lambda}}_{i+1}^k / (\eta_{n+i}\beta_k) \right\|^2$$
$$= \pi_{X_i} \left(\mathbf{x}_{n+i}^k + \hat{\boldsymbol{\lambda}}_{i+1}^k / (\eta_{n+i}\beta_k) \right), \tag{33}$$

where π_{X_i} is the projection onto X_i and i = 1, ..., n.

As for the choice of η_i 's, although we can simply apply Theorem 1 to assign their values as $\eta_i > 2n(||A_i||^2 + 1)$ and $\eta_{n+i} > 2n$, i = 1, ..., n, such choices are too pessimistic. As η_i 's are related to the magnitudes of the differences in \mathbf{x}_i^{k+1} from \mathbf{x}_i^k , we had better provide tighter estimate on η_i 's in order to achieve faster convergence. Actually, we have the following better result.

Theorem 5 For problem (27), if $\{\beta_k\}$ is non-decreasing and upper bounded and η_i 's are chosen as $\eta_i > n \|\mathcal{A}_i\|^2 + 2$ and $\eta_{n+i} > 2, i = 1, ..., n$, then the sequence $\{(\{\mathbf{x}_i^k\}, \boldsymbol{\lambda}^k)\}$ generated by LADMPSAP converge to a KKT point of problem (27).

Finally, we summarize LADMPSAP for problem (27) in Algorithm 2, which is a practical algorithm for solving (26).

Remark 2 Analogs of Theorems 2 and 3 are also true for Algorithm 2 although ∂f_{n+i} 's are unbounded, thanks to our assumptions that all ∂f_i , i = 1, ..., n, are bounded and **b** is an interior point of $\sum_{i=1}^{n} A_i(X_i)$, which result in an analog of Proposition 4. Consequently, β_{max} can also be removed if all ∂f_i , i = 1, ..., n, are bounded.

Algorithm 2 LADMPSAP for (27), also a Practical Algorithm for (26).

Initialize: Set $\rho_0 > 1$, $\varepsilon_1 > 0$, $\varepsilon_2 > 0$, $\beta_{\max} \gg 1 \gg \beta_0 > 0$, $\lambda^0 = ((\lambda_1^0)^T, \dots, (\lambda_{n+1}^0)^T)^T$, $\eta_i > n \|\mathcal{A}_i\|^2 + 2$, $\eta_{n+i} > 2$, $\mathbf{x}_i^0, \mathbf{x}_{n+i}^0 = \mathbf{x}_i^0$, $i = 1, \dots, n$. **while** (21) or (22) is not satisfied **do Step 1:** Compute $\hat{\lambda}^k$ as (31). **Step 2:** Update \mathbf{x}_i , $i = 1, \dots, 2n$, in parallel as (32)-(33). **Step 3:** Update λ by (30) and β by (18) and (20). **end while** (Note that in (20), (21), and (22), n and \mathcal{A}_i should be replaced by 2n and $\hat{\mathcal{A}}_i$, respectively.)

Remark 3 Since Algorithm 2 is an application of Algorithm 1 to problem (27), only with refined parameter estimation, its convergence rate in an ergodic sense is also $O\left(1/\sum_{k=0}^{K} \beta_k^{-1}\right)$, where *K* is the number of iterations.

5 Proximal LADMPSAP for even more general convex programs

In LADMPSAP we have assumed that the subproblems (16) are easily solvable. In many machine learning problems, the functions f_i 's are often matrix or vector norms or characteristic functions of convex sets. So this assumption often holds. Nonetheless, this assumption is not always true, e.g., when f_i is the logistic loss function [see (6)]. So in this section we aim at generalizing LADMPSAP to solve even more general convex programs (1).

We are interested in the case that f_i can be decomposed into two components:

$$f_i(\mathbf{x}_i) = g_i(\mathbf{x}_i) + h_i(\mathbf{x}_i), \tag{34}$$

where both g_i and h_i are convex, g_i is $C^{1,1}$:

$$\|\nabla g_i(\mathbf{x}) - \nabla g_i(\mathbf{y})\| \le L_i \|\mathbf{x} - \mathbf{y}\|, \quad \forall x, y \in \mathbb{R}^{d_i},$$
(35)

and h_i may not be differentiable but its proximal operation is easily solvable. For brevity, we call L_i the Lipschitz constant of ∇g_i .

Recall that in each iteration of LADMPSAP, we have to solve subproblem (16). Since now we do not assume that the proximal operation of f_i (7) is easily solvable, we may have difficulty in solving subproblem (16). By (34), we write down (16) as

$$\mathbf{x}_{i}^{k+1} = \underset{\mathbf{x}_{i}}{\operatorname{argmin}} h_{i}(\mathbf{x}_{i}) + g_{i}(\mathbf{x}_{i}) + \frac{\sigma_{i}^{(k)}}{2} \left\| \mathbf{x}_{i} - \mathbf{x}_{i}^{k} + \mathcal{A}_{i}^{\dagger}\left(\hat{\boldsymbol{\lambda}}^{k}\right) / \sigma_{i}^{(k)} \right\|^{2}, \quad i = 1, \dots, n,$$
(36)

Since $g_i(\mathbf{x}_i) + \frac{\sigma_i^{(k)}}{2} \|\mathbf{x}_i - \mathbf{x}_i^k + \mathcal{A}_i^{\dagger}(\hat{\mathbf{\lambda}}^k) / \sigma_i^{(k)}\|^2$ is $C^{1,1}$, we may also linearize it at \mathbf{x}_i^k and add a proximal term. Such an idea leads to the following updating scheme of \mathbf{x}_i :

$$\mathbf{x}_{i}^{k+1} = \operatorname*{argmin}_{\mathbf{x}_{i}} h_{i}(\mathbf{x}_{i}) + g_{i}\left(\mathbf{x}_{i}^{k}\right) + \frac{\sigma_{i}^{(k)}}{2} \left\|\mathcal{A}_{i}^{\dagger}\left(\hat{\boldsymbol{\lambda}}^{k}\right) / \sigma_{i}^{(k)}\right\|^{2} \\ + \left\langle\nabla g_{i}\left(\mathbf{x}_{i}^{k}\right) + \mathcal{A}_{i}^{\dagger}\left(\hat{\boldsymbol{\lambda}}^{k}\right), \mathbf{x}_{i} - \mathbf{x}_{i}^{k}\right\rangle + \frac{\tau_{i}^{(k)}}{2} \left\|\mathbf{x}_{i} - \mathbf{x}_{i}^{k}\right\|^{2} \\ = \operatorname*{argmin}_{\mathbf{x}_{i}} h_{i}(\mathbf{x}_{i}) + \frac{\tau_{i}^{(k)}}{2} \left\|\mathbf{x}_{i} - \mathbf{x}_{i}^{k} + \frac{1}{\tau_{i}^{(k)}} \left[\mathcal{A}_{i}^{\dagger}\left(\hat{\boldsymbol{\lambda}}^{k}\right) + \nabla g_{i}\left(\mathbf{x}_{i}^{k}\right)\right]\right\|^{2}, \quad (37)$$

Algorithm 3 Proximal LADMPSAP for Solving (1) with f_i Satisfying (34).

Initialize: Set $\rho_0 > 1$, $\beta_0 > 0$, λ^0 , $T_i \ge L_i$, $\eta_i > n ||\mathcal{A}_i||^2$, \mathbf{x}_i^0 , i = 1, ..., n. **while** (39) or (40) is not satisfied **do Step 1:** Compute $\hat{\boldsymbol{\lambda}}^k$ as (19). **Step 2:** Update \mathbf{x}_i 's in parallel by solving

$$\mathbf{x}_{i}^{k+1} = \underset{\mathbf{x}_{i}}{\operatorname{argmin}} h_{i}(\mathbf{x}_{i}) + \frac{\tau_{i}^{(k)}}{2} \left\| \mathbf{x}_{i} - \mathbf{x}_{i}^{k} + \frac{1}{\tau_{i}^{(k)}} \left[\mathcal{A}_{i}^{\dagger}(\hat{\boldsymbol{\lambda}}^{k}) + \nabla g_{i}\left(\mathbf{x}_{i}^{k}\right) \right] \right\|^{2}, \ i = 1, \dots, n,$$
(41)

where $\tau_i^{(k)} = T_i + \beta_k \eta_i$. **Step 3:** Update λ by (17) and β by (18) with ρ defined in (38). **end while**

where i = 1, ..., n. The choice of $\tau_i^{(k)}$ is presented in Theorem 6, i.e. $\tau_i^{(k)} = T_i + \beta_k \eta_i$, where $T_i \ge L_i$ and $\eta_i > n \|\mathcal{A}_i\|^2$ are both positive constants.

By our assumption on h_i , the above subproblems are easily solvable. The update of Lagrange multiplier λ and β are still respectively goes as (17) and (18) but with

$$\rho = \begin{cases}
\rho_0, & \text{if max}\left(\left\{\|\mathcal{A}_i\|^{-1} \left\|\nabla g_i\left(\mathbf{x}_i^{k+1}\right) - \nabla g_i\left(\mathbf{x}_i^{k}\right) - \tau_i^{(k)}\left(\mathbf{x}_i^{k+1} - \mathbf{x}_i^{k}\right)\right\|, \\
i = 1, \dots, n\right\}\right) / \|\mathbf{b}\| < \varepsilon_2, \\
1, & \text{otherwise.}
\end{cases}$$
(38)

The iteration terminates when the following two conditions are met:

$$\left\|\sum_{i=1}^{n} \mathcal{A}_{i}(\mathbf{x}_{i}^{k+1}) - \mathbf{b}\right\| / \|\mathbf{b}\| < \varepsilon_{1},$$

$$\max\left(\left\{\|\mathcal{A}_{i}\|^{-1} \left\|\nabla g_{i}(\mathbf{x}_{i}^{k+1}) - \nabla g_{i}(\mathbf{x}_{i}^{k}) - \tau_{i}^{(k)}(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k})\right\|,$$

$$i = 1, \dots, n\right\}\right) / \|\mathbf{b}\| < \varepsilon_{2}.$$

$$(40)$$

These two conditions are also deduced from the KKT conditions.

We call the above algorithm as proximal LADMPSAP and summarize it in Algorithm 3. As for the convergence of proximal LADMPSAP, we have the following theorem.

Theorem 6 (Convergence of proximal LADMPSAP) If β_k is non-decreasing and upper bounded, $\tau_i^{(k)} = T_i + \beta_k \eta_i$, where $T_i \ge L_i$ and $\eta_i > n \|\mathcal{A}_i\|^2$ are both positive constants, i = 1, ..., n, then {($\{\mathbf{x}_i^k\}, \boldsymbol{\lambda}^k$)} generated by proximal LADMPSAP converge to a KKT point of problem (1).

We further have the following convergence rate theorem for proximal LADMPSAP in an ergodic sense.

Theorem 7 (Convergence rate of proximal LADMPSAP) Define $\bar{\mathbf{x}}_i^K = \sum_{k=0}^K \gamma_k \mathbf{x}_i^{k+1}$, where $\gamma_k = \beta_k^{-1} / \sum_{j=0}^K \beta_j^{-1}$. Then the following inequality holds for $\bar{\mathbf{x}}_i^K$:

$$\sum_{i=1}^{n} \left(f_i\left(\bar{\mathbf{x}}_i^K\right) - f_i\left(\mathbf{x}_i^*\right) + \left\langle \mathcal{A}_i^{\dagger}(\lambda^*), \bar{\mathbf{x}}_i^K - \mathbf{x}_i^* \right\rangle \right) + \frac{\alpha\beta_0}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_i\left(\bar{\mathbf{x}}_i^K\right) - \mathbf{b} \right\|^2$$
$$\leq C_0 / \sum_{k=0}^{K} 2\beta_k^{-1}, \tag{42}$$

🖉 Springer

where

$$\alpha^{-1} = (n+1) \max\left(1, \left\{\frac{\|\mathcal{A}_i\|^2}{\eta_i - n\|\mathcal{A}_i\|^2}, i = 1, \dots, n\right\}\right)$$

and

$$C_0 = \sum_{i=1}^n \beta_0^{-1} \tau_i^{(0)} \| \mathbf{x}_i^0 - \mathbf{x}_i^* \|^2 + \beta_0^{-2} \| \lambda^0 - \lambda^* \|^2.$$

When there are extra convex set constraints, $\mathbf{x}_i \in X_i$, i = 1, ..., n, we can also introduce auxiliary variables as in Sect. 4 and have an analogy of Theorems 5 and 4.

Theorem 8 For problem (27), where f_i is described at the beginning of Section 5, if β_k is non-decreasing and upper bounded and $\tau_i^{(k)} = T_i + \eta_i \beta_k$, where $T_i \ge L_i$, $T_{n+i} = 0$, $\eta_i > n \|\mathcal{A}_i\|^2 + 2$, and $\eta_{n+i} > 2$, i = 1, ..., n, then $\{(\{\mathbf{x}_i^k\}, \mathbf{\lambda}^k)\}$ generated by proximal LADMPSAP converge to a KKT point of problem (27). The convergence rate in an ergodic sense is also $O\left(1/\sum_{k=0}^{K} \beta_k^{-1}\right)$, where K is the number of iterations.

6 Numerical results

In this section, we test the performance of LADMPSAP on three specific examples of problem (1), i.e., Latent low-rank representation [see (2)], nonnegative matrix completion [see (3)], and group sparse logistic regression with overlap [see (6)].

6.1 Solving latent low-rank representation

We first solve the latent LRR problem (Liu and Yan 2011) (2). In order to test LADMPSAP and related algorithms with data whose characteristics are controllable, we follow (Liu et al. 2010) to generate synthetic data, which are parameterized as (s, p, d, \tilde{r}) , where s, p, d, and \tilde{r} are the number of independent subspaces, points in each subspace, and ambient and intrinsic dimensions, respectively. The number of scale variables and constraints is $(sp) \times d$.

As first order methods are popular for solving convex programs in machine learning (Boyd et al. 2011), here we compare LADMPSAP with several conceivable first order algorithms, including APG (Beck and Teboulle 2009), naive ADM, naive LADM, LADMGB, and LADMPS. Naive ADM and naive LADM are generalizations of ADM and LADM, respectively, which are straightforwardly generalized from two variables to multiple variables, as discussed in Sect. 3.1. Naive ADM is applied to solve (2) after rewriting the constraint of (2) as $\mathbf{X} = \mathbf{XP} + \mathbf{QX} + \mathbf{E}$, $\mathbf{P} = \mathbf{Z}$, $\mathbf{Q} = \mathbf{L}$. For LADMPS, β_k is fixed in order to show the effectiveness of adaptive penalty. The parameters of APG and ADM are the same as those in (Lin et al. 2009b) and (Liu and Yan 2011), respectively. For LADM, we follow the suggestions in (Yang and Yuan 2013) to fix its penalty parameter β at 2.5/min(d, sp), where $d \times sp$ is the size of \mathbf{X} . For LADMGB, as there is no suggestion in He and Yuan (2013) on how to choose a fixed β , we simply set it the same as that in LADM. The rest of the parameters are the same as those suggested in He et al. (2012). We fix $\beta = \sigma_{max}(\mathbf{X}) \min(d, sp)\varepsilon_2$ in LADMPS and set $\beta_0 = \sigma_{max}(\mathbf{X}) \min(d, sp)\varepsilon_2$ and $\rho_0 = 10$ in LADMPSAP. For LADMPSAP, we also set $\eta_Z = \eta_L = 1.02 \times 3\sigma_{max}^2(\mathbf{X})$, where η_Z and η_L are the parameters η_i 's in Algorithm 1 for \mathbf{Z} and \mathbf{L} , respectively. For the stopping criteria, $\|\mathbf{XZ}^k + \mathbf{L}^k\mathbf{X} + \mathbf{E}^k - \mathbf{X}\|/\|\mathbf{X}\| \le \varepsilon_1$ and $\max(\|\mathbf{Z}^k - \mathbf{Z}^{k-1}\|, \|\mathbf{L}^k - \mathbf{L}^{k-1}\|, \|\mathbf{E}^k - \mathbf{E}^{k-1}\|)/\|\mathbf{X}\| \le \varepsilon_2$, with $\varepsilon_1 = 10^{-3}$ and $\varepsilon_2 = 10^{-4}$

(s, p, d, \tilde{r})	Method	Time	No. of iteration	$\frac{\ \hat{\mathbf{Z}}-\mathbf{Z}^*\ }{\ \mathbf{Z}^*\ }$	$\frac{\ \hat{\mathbf{L}}-\mathbf{L}^*\ }{\ \mathbf{L}^*\ }$	$\frac{\ \hat{\mathbf{E}}-\mathbf{E}^*\ }{\ \mathbf{E}^*\ }$	Acc.
(5, 50, 250, 5)	APG	18.20	236	0.3389	0.3167	0.4500	95.6
	nADM	16.32	172	0.3993	0.3928	0.5592	95.6
	nLADM	21.34	288	0.4553	0.4408	0.5607	95.6
	LADMGB	24.10	290	0.4520	0.4355	0.5610	95.6
	LADMPS	17.15	232	0.0163	0.0139	0.0446	95.6
	LADMPSAP	8.04	109	0.0089	0.0083	0.0464	95.6
(10, 50, 500, 5)	APG	85.03	234	0.1020	0.0844	0.7161	95.8
	nADM	78.27	170	0.0928	0.1026	0.6636	95.8
	nLADM	181.42	550	0.2077	0.2056	0.6623	95.8
	LADMGB	214.94	550	0.1877	0.1848	0.6621	95.8
	LADMPS	64.65	200	0.0167	0.0089	0.1059	95.8
	LADMPSAP	37.85	117	0.0122	0.0055	0.0780	95.8
(20, 50, 1000, 5)	APG	544.13	233	0.0319	0.0152	0.2126	95.2
	nADM	466.78	166	0.0501	0.0433	0.2676	95.2
	nLADM	1,888.44	897	0.1783	0.1746	0.2433	95.2
	LADMGB	2,201.37	897	0.1774	0.1736	0.2434	95.2
	LADMPS	367.68	177	0.0151	0.0105	0.0872	95.2
	LADMPSAP	260.22	125	0.0106	0.0041	0.0671	95.2

 Table 1
 Comparisons of APG, naive ADM (nADM), naive LADM (nLADM), LADMGB, LADMPS, and LADMPSAP on the latent LRR problem (2)

The quantities include computing time (in seconds), number of iterations, relative errors, and clustering accuracy (in percentage). They are averaged over 10 runs

Best results are shown in bold

are used for all the algorithms. For the parameter μ in (2), we empirically set it as $\mu = 0.01$. To measure the relative errors in the solutions we run LADMPSAP 2,000 iterations with $\rho_0 = 1.01$ to obtain the estimated ground truth solution ($\mathbb{Z}^*, \mathbb{L}^*, \mathbb{E}^*$). The experiments are run and timed on a notebook computer with an Intel Core i7 2.00 GHz CPU and 6 GB memory, running Windows 7 and Matlab 7.13.

Table 1 shows the results of related algorithms. We can see that LADMPS and LADMP-SAP are faster and more numerically accurate than LADMGB, and LADMPSAP is even faster than LADMPS thanks to the adaptive penalty. Moreover, naive ADM and naive LADM have relatively poorer numerical accuracy, possibly due to converging to wrong solutions. The numerical accuracy of APG is also worse than those of LADMPS and LADMPSAP because it only solves an approximate problem by adding the constraint to the objective function as penalty. Note that although we do not require { β_k } to be bounded, this does not imply that β_k will grow infinitely. As a matter of fact, when LADMPSAP terminates the final values of β_k are 21.1567, 42.2655, and 81.4227 for the three data settings, respectively.

We then test the performance of the above six algorithms on the Hopkins155 database (Tron and Vidal 2007), which consists of 156 sequences, each having 39–550 data vectors drawn from two or three motions. For computational efficiency, we preprocess the data by projecting them to be 5-dimensional using PCA. We test all algorithms with $\mu = 2.4$, which is the best parameter for LRR on this database (Liu et al. 2010). Table 2 shows the results on the Hopkins155 database. We can also see that LADMPSAP is faster than other methods in comparison. In particular, LADMPSAP is faster than LADMPS, which uses a fixed β . This testify to the advantage of using an adaptive penalty.

Method	Time (s)	No. of iteration	Error (%)
APG	10.37	67	8.33
nADM	24.76	144	8.33
nLADM	15.50	112	8.33
LADMGB	16.05	113	8.36
LADMPS	15.58	113	8.33
LADMPSAP	3.80	26	8.33

 Table 2
 Comparisons of APG, naive ADM (nADM), naive LADM (nLADM), LADMGB, LADMPS, and LADMPSAP on the Hopkins155 database

The quantities include average computing time, average number of iterations, and average classification errors on all 156 sequences

Best results are shown in bold

Table 3 Comparisons on the NMC problem (3) with synthetic data, averaged on 10 runs

X			LADM	LADM				LADMPSAP		
n	q (%)	t/d_r	No. of iteration	Time (s)	RelErr	FA	No. of iteration	Time (s)	RelErr	FA
1,000	20	10.05	375	177.92	1.35E-5	6.21E-4	58	24.94	9.67E-6	0
	10	5.03	1,000	459.70	4.60E-5	6.50E-4	109	42.68	1.72E-5	0
5,000	20	50.05	229	1,613.68	1.08E-5	1.93E-4	49	369.96	9.05E-6	0
	10	25.03	539	2,028.14	1.20E-5	7.70E-5	89	365.26	9.76E-6	0
10,000	10	50.03	463	6,679.59	1.11E-5	4.18E-5	89	1584.39	1.03E-5	0

q, t, and d_r denote, respectively, the sample ratio, the number of measurements t = q(mn), and the "degree of freedom" defined by $d_r = r(m + n - r)$ for an $m \times n$ matrix with rank r and q. Here we set m = n and fix r = 10 in all the tests

Best results are shown in bold

6.2 Solving nonnegative matrix completion

This subsection evaluates the performance of the practical LADMPSAP proposed in Sect. 4 for solving nonnegative matrix completion (Xu et al. 2011) (3).

We first evaluate the numerical performance on synthetic data to demonstrate the superiority of practical LADMPSAP over the conventional LADM⁹ (Yang and Yuan 2013). The nonnegative low-rank matrix X_0 is generated by truncating the singular values of a randomly generated matrix. As LADM cannot handle the nonnegativity constraint, it actually solve the standard matrix completion problem, i.e., (3) without the nonnegativity constraint. For LADMPSAP, we follow the conditions in Theorem 5 to set η_i 's and set the rest of the parameters the same as those in Sect. 6.1. The stopping tolerances are set as $\varepsilon_1 = \varepsilon_2 = 10^{-5}$. The numerical comparison is shown in Table 3, where the relative nonnegative feasibility (FA) is defined as (Xu et al. 2011):

FA: =
$$\|\min(\hat{\mathbf{X}}, 0)\| / \|\mathbf{X}_0\|,$$

in which \mathbf{X}_0 is the ground truth and $\hat{\mathbf{X}}$ is the computed solution. It can be seen that the numerical performance of LADMPSAP is much better than that of LADM, thus again verifies

⁹ Code available at http://math.nju.edu.cn/~jfyang/IADM_NNLS/index.html.



Fig. 1 Image inpainting by FPCA, LADM and LADMPSAP

 Table 4
 Comparisons on the image inpainting problem. "PSNR" stands for "Peak Signal to Noise Ratio" measured in decibel (dB)

Method	No. of iteration	Time (s)	PSNR (dB)	FA
FPCA	179	228.99	27.77	9.41E-4
LADM	228	207.95	26.98	2.92E-3
LADMPSAP	143	134.89	31.39	0

Best results are shown in bold

the efficiency of our proposed parallel splitting and adaptive penalty scheme for enhancing ADM/LADM type algorithms.

We then consider the image inpainting problem, which is to fill in the missing pixel values of a corrupted image. As the pixel values are nonnegative, the image inpainting problem can be formulated as the NMC problem. To prepare a low-rank image, we also truncate the singular values of a $1,024 \times 1,024$ grayscale image "man"¹⁰ to obtain an image of rank 40, shown in Fig. 1a, b. The corrupted image is generated from the original image (all pixels have been normalized in the range of [0, 1]) by sampling 20% of the pixels uniformly at random and adding Gaussian noise with mean zero and standard deviation 0.1.

Besides LADM, here we also consider another recently proposed fixed point continuation with approximate SVD [FPCA (Ma et al. 2011)] on this problem. Similar to LADM, the code of FPCA¹¹ can only solve the standard matrix completion problem without the nonnegativity constraint. This time we set $\varepsilon_1 = 10^{-3}$ and $\varepsilon_2 = 10^{-1}$ as the thresholds for stopping criteria. The recovered images are shown in Fig. 1c–e and the quantitative results are in Table 4. One can see that on our test image both the qualitative and the quantitative results of LADMPSAP are better than those of FPCA and LADM. Note that LADMPSAP is faster than FPCA and LADM even though they do not handle the nonnegativity constraint.

6.3 Solving group sparse logistic regression with overlap

In this subsection, we apply proximal LADMPSAP to solve the problem of group sparse logistic regression with overlap (5).

The Lipschitz constant of the gradient of logistic function with respect to $\bar{\mathbf{w}}$ can be proven to be $L_{\bar{w}} \leq \frac{1}{4s} \|\bar{\mathbf{X}}\|_2^2$, where $\bar{\mathbf{X}} = (\bar{\mathbf{x}}_1, \bar{\mathbf{x}}_2, \dots, \bar{\mathbf{x}}_s)$. Thus (5) can be directly solved by Algorithm 3.

¹⁰ Available at http://sipi.usc.edu/database/.

¹¹ Code available at http://www1.se.cuhk.edu.hk/~sqma/softwares.html.

6.3.1 Synthetic data

To assess the performance of proximal LADMPSAP, we simulate data with p = 9t + 1 variables, covered by *t* groups of ten variables with overlap of one variable between two successive groups: $\{1, ..., 10\}, \{10, ..., 19\}, ..., \{p - 9, ..., p\}$. We randomly choose *q* groups to be the support of **w**. If the chosen groups have overlapping variables with the unchosen groups, the overlapping variables are removed from the support of **w**. So the support of **w** may be less than 10q. $\mathbf{y} = (y_1, ..., y_s)^T$ is chosen as $(1, -1, 1, -1, ...)^T$. $\mathbf{X} \in \mathbb{R}^{p \times s}$ is generated as follows. For $\mathbf{X}_{i,j}$, if *i* is in the support of **w** and $\mathbf{y}_j = -1$, then $\mathbf{X}_{i,j}$ is generated uniformly on [-1.5, -0.5]; if *i* is not in the support of **w** are statistically different from the remaining rows in **X**, hence can be considered as informative rows. We use model (6) to select the informative rows for classification, where $\mu = 0.1$. If the ground truth support of **w** is recovered, then the two groups of data are linearly separable by considering only the coordinates in the support of **w**.

We compare proximal LADMPSAP with a series of ADM based methods, including ADM, LADM, LADMPS, and LADMPSAP, where the subproblems for **w** and **b** have to be solved iteratively, e.g., by APG (Beck and Teboulle 2009). We terminate the inner loop by APG when the norm of gradient of the objective function of the subproblem is less than 10^{-6} . As for the outer loop, we choose $\varepsilon_1 = 2 \times 10^{-4}$ and $\varepsilon_2 = 2 \times 10^{-3}$ as the thresholds to terminate the iterations.

For ADM, LADM, and LADMPS, which use a fixed penalty β , as we do not find any suggestion on its choice in the literature (the choice suggested in Yang and Yuan (2013) is for nuclear norm regularized least square problem only) we try multiple choices of β and choose the one that results in the fastest convergence. For LADMPSAP, we set $\beta_0 = 0.2$ and $\rho_0 = 5$. For proximal LADMPSAP we set $T_1 = \frac{1}{4s} \|\bar{\mathbf{X}}\|_2^2$, $\eta_1 = 2.01 \|\bar{\mathbf{S}}\|_2^2$, $T_2 = 0$, $\eta_2 = 2.01$, $\beta_0 = 1$, and $\rho_0 = 5$. To measure the relative errors in the solutions we iterate proximal LADMPSAP for 2,000 times and regard its output as the ground truth solution ($\bar{\mathbf{w}}^*, \mathbf{z}^*$).

Table 5 shows the comparison among related algorithms. The ground truth support of **w** is recovered by all the compared algorithms. We can see that ADM, LADM, LADMPS, and LADMPSAP are much slower than proximal LADMPSAP because of the time-consuming subproblem computation, although they have much smaller number of outer iterations. Their numerical accuracies are also inferior to that of proximal LADMPSAP. We can also see that LADMPSAP is faster and more numerically accurate than ADM, LADM, and LADMPS. This again testifies to the effectiveness of using adaptive penalty.

6.3.2 Pathway analysis on breast cancer data

Then we consider the pathway analysis problem using the breast cancer gene expression data set (Vijver and He 2002), which consists of 8,141 genes in 295 breast cancer tumors (78 metastatic and 217 non-metastatic). We follow Jacob et al. (2009) and use the canonical pathways from MSigDB (Subramanian et al. 2005) to generate the overlapping gene sets, which contains 639 groups of genes, 637 of which involve genes from our study. The statistics of the 637 gene groups are summarized as follows: the average number of genes in each group is 23.7, the largest gene group has 213 genes, and 3,510 genes appear in these 637 groups with an average appearance frequency of about four. We follow Jacob et al. (2009) to restrict the analysis to the 3,510 genes and balance the data set by using three replicates of each

(s, p, t, q)	Method	Time	No. of iteration	$rac{\ \hat{oldsymbol{w}}-oldsymbol{ar{w}}^*\ }{\ oldsymbol{ar{w}}^*\ }$	$\tfrac{\ \hat{\mathbf{z}} - \mathbf{z}^*\ }{\ \mathbf{z}^*\ }$
(300, 901, 100, 10)	ADM	294.15	43	0.4800	0.4790
	LADM	229.03	43	0.5331	0.5320
	LADMPS	105.50	47	0.2088	0.2094
	LADMPSAP	57.46	39	0.0371	0.0368
	pLADMPSAP	1.97	141	0.0112	0.0112
(450, 1351, 150, 15)	ADM	450.96	33	0.4337	0.4343
	LADM	437.12	36	0.5126	0.5133
	LADMPS	201.30	39	0.1938	0.1937
	LADMPSAP	136.64	37	0.0321	0.0306
	pLADMPSAP	4.16	150	0.0131	0.0131
(600, 1801, 200, 20)	ADM	1,617.09	62	1.4299	1.4365
	LADM	1,486.23	63	1.5200	1.5279
	LADMPS	494.52	46	0.4915	0.4936
	LADMPSAP	216.45	32	0.0787	0.0783
	pLADMPSAP	5.77	127	0.0276	0.0277

Table 5 Comparisons among ADM, LADM, LADMPS, LADMPSAP, and proximal LADMPSAP (pLADMPSAP) on the group sparse logistic regression with overlap problem. The quantities include the computing time (in seconds), number of outer iterations, and relative errors

Best results are shown in bold

metastasis patient in the training set. We use model (6) to select genes, where $\mu = 0.08$. We want to predict whether a tumor is metastatic ($y_i = 1$) or non-metastatic ($y_i = -1$).

We compare proximal LADMPSAP with the active set method, which was adopted in (Jacob et al. 2009),¹² LADM, and LADMPSAP. In LADMPSAP and proximal LADMP-SAP, we both set $\beta_0 = 0.8$ and $\rho_0 = 1.1$. For LADM, we try multiple choices of β and choose the one that results in the fastest convergence. In LADM and LADMPSAP, we terminate the inner loop by APG when the norm of gradient of the objective function of the subproblem is less than 10^{-6} . The thresholds for terminating the outer loop are all chosen as $\varepsilon_1 = 10^{-3}$ and $\varepsilon_2 = 6 \times 10^{-3}$. For the three LADM based methods, we first solve (6) to select genes. Then we use the selected genes to re-train a traditional logistic regression model and use the model to predict the test samples. As in Jacob et al. (2009) we partition the whole data set into three subsets to do the experiment three times. Each time we select one subset as the test set and the other two as the training set (i.e., there are $(78 + 217) \times 2/3 = 197$ samples for training). It is worth mentioning that Jacob et al. (2009) only kept the 300 genes that are the most correlated with the output in the pre-processing step. In contrast, we use all the 3,510 genes in the training phase.

Table 6 shows that proximal LADMPSAP is more than ten times faster than the active set method used in Jacob et al. (2009), although it computes with a more than ten times larger training set. Proximal LADMPSAP is also much faster than LADM and LADMPSAP due to the lack of inner loop to solve subproblems. The prediction error and the sparseness at the pathway level by proximal LADMPSAP is also competitive with those of other methods in comparison.

¹² Code available at http://cbio.ensmp.fr/~ljacob/documents/overlasso-package.tgz.

Method	Time	Error	No. of pathway		
Active set	2,179	0.36 ± 0.03	6, 5, 78		
LADM	2,433	0.315 ± 0.049	7, 9, 10		
LADMPSAP	1,593	0.329 ± 0.011	7, 9, 9		
pLADMPSAP	179	0.312 ± 0.026	4, 6, 6		

 Table 6
 Comparisons among the Active Set method (Jacob et al. 2009), LADM, LADMPSAP, and proximal LADMPSAP (pLADMPSAP) on the pathway analysis

We present the CPU time (in seconds), classification error rate, and number of pathways. Results are estimated by three-fold cross validation. No. of Pathway gives the number of pathways that the selected genes belong to in each of the cross validation

Best results are shown in bold

7 Conclusions

In this paper, we propose linearized alternating direction method with parallel splitting and adaptive penalty (LADMPSAP) for efficiently solving linearly constrained multi-block separable convex programs, which are abundant in machine learning. LADMPSAP fully utilizes the properties that the proximal operations of the component objective functions and the projections onto convex sets are easily solvable, which are usually satisfied by machine learning problems, making each of its iterations cheap. It is also highly parallel, making it appealing for parallel or distributed computing. Numerical experiments testify to the advantages of LADMPSAP over other possible first order methods.

Although LADMPSAP is inherently parallel, when solving the proximal operations of component objective functions we will still face basic numerical algebraic computations. So for particular large scale machine learning problems, it will be interesting to integrate the existing distributed computing techniques [e.g., parallel incomplete Cholesky factorization (Chang et al. 2007; Chang 2011) and caching factorization techniques (Boyd et al. 2011)] with our LADMPSAP in order to effectively address the scalability issues.

Acknowledgments Z. Lin is supported by 973 Program of China (No. 2015CB3525), NSFC (Nos. 61272341 and 61231002), and Microsoft Research Asia Collaborative Research Program. R. Liu is supported by NSFC (No. 61300086), the China Postdoctoral Science Foundation (Nos. 2013M530917 and 2014T70249), the Fundamental Research Funds for the Central Universities (No. DUT12RC(3)67) and the Open Project Program of the State Key Lab of CAD&CG (No. A1404), Zhejiang University. Z. Lin also thanks Xiaoming Yuan, Wotao Yin, and Edward Chang for valuable discussions and HTC for financial support.

Appendix 1: Proof of Theorem 1

To prove this theorem, we first have the following lemmas and propositions.

Lemma 1 (KKT Condition) *The Kuhn–Karush–Tucker (KKT) condition of problem (1) is that there exists* $(\mathbf{x}_1^*, \ldots, \mathbf{x}_n^*, \lambda^*)$ *, such that*

$$\sum_{i=1}^{n} \mathcal{A}_i \left(\mathbf{x}_i^* \right) = \mathbf{b},\tag{43}$$

$$-\mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \in \partial f_{i}\left(\mathbf{x}_{i}^{*}\right), \quad i = 1, \dots, n,$$

$$\tag{44}$$

where ∂f_i is the subgradient of f_i .

The first is the feasibility condition and the second is the duality condition. Such $(\mathbf{x}_1^*, \ldots, \mathbf{x}_n^*, \lambda^*)$ is called a KKT point of problem (1).

Lemma 2 For $\{(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k, \lambda^k)\}$ generated by Algorithm 1, we have that

$$-\sigma_{i}^{(k)}\left(\mathbf{x}_{i}^{k+1}-\mathbf{u}_{i}^{k}\right)\in\partial f_{i}\left(\mathbf{x}_{i}^{k+1}\right),\quad i=1,\ldots,n,$$
(45)

where $\mathbf{u}_i^k = \mathbf{x}_i^k - \mathcal{A}_i^{\dagger}(\hat{\boldsymbol{\lambda}}^k) / \sigma_i^{(k)}$.

This can be easily proved by checking the optimality conditions of (16).

Lemma 3 For $\{(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k, \lambda^k)\}$ generated by Algorithm 1 and a KKT point $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*, \lambda^*)$ of problem (1), the following inequality holds:

$$\left\langle -\sigma_i^{(k)} \left(\mathbf{x}_i^{k+1} - \mathbf{u}_i^k \right) + \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*), \, \mathbf{x}_i^{k+1} - \mathbf{x}_i^* \right\rangle \ge 0, \quad i = 1, \dots, n.$$
(46)

This can be deduced by the monotonicity of subgradient mapping (Rockafellar 1970).

Lemma 4 For $\{(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k, \lambda^k)\}$ generated by Algorithm 1 and a KKT point $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*, \lambda^*)$ of problem (1), we have that

$$\beta_{k} \sum_{i=1}^{n} \sigma_{i}^{(k)} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} + \left\| \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{*} \right\|^{2}$$
(47)

$$=\beta_{k}\sum_{i=1}^{n}\sigma_{i}^{(k)}\left\|\mathbf{x}_{i}^{k}-\mathbf{x}_{i}^{*}\right\|^{2}+\left\|\boldsymbol{\lambda}^{k}-\boldsymbol{\lambda}^{*}\right\|^{2}$$
(48)

$$-2\beta_k \sum_{i=1}^n \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, -\sigma_i^{(k)} \left(\mathbf{x}_i^{k+1} - \mathbf{u}_i^k \right) + \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*) \right\rangle$$
(49)

$$-\beta_{k}\sum_{i=1}^{n}\sigma_{i}^{(k)}\left\|\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\right\|^{2}-\left\|\boldsymbol{\lambda}^{k+1}-\boldsymbol{\lambda}^{k}\right\|^{2}$$
(50)

$$-2\beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathbf{x}_i^k - \mathbf{u}_i^k \right\rangle$$
(51)

$$+2\left(\boldsymbol{\lambda}^{k+1}-\boldsymbol{\lambda}^{k},\boldsymbol{\lambda}^{k+1}\right).$$
(52)

Proof This can be easily checked. First, we add (49) and (51) to have

$$-2\beta_k \sum_{i=1}^n \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, -\sigma_i^{(k)} \left(\mathbf{x}_i^{k+1} - \mathbf{u}_i^k \right) + \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*) \right\rangle$$
(53)

$$-2\beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathbf{x}_i^k - \mathbf{u}_i^k \right\rangle$$
(54)

$$= -2\beta_k \sum_{i=1}^n \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathcal{A}_i^{\dagger}(\lambda^*) \right\rangle + 2\beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\rangle$$
(55)

$$= -2\beta_k \sum_{i=1}^n \left\langle \mathcal{A}_i \left(\mathbf{x}_i^{k+1} - \mathbf{x}_i^* \right), \mathbf{\lambda}^* \right\rangle + 2\beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\rangle$$
(56)

$$= -2\left\langle\beta_{k}\sum_{i=1}^{n}\mathcal{A}_{i}\left(\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{*}\right),\boldsymbol{\lambda}^{*}\right\rangle + 2\beta_{k}\sum_{i=1}^{n}\sigma_{i}^{(k)}\left\langle\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{*},\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\right\rangle$$
(57)

$$= -2\left\langle \beta_k \left(\sum_{i=1}^n \mathcal{A}_i \left(\mathbf{x}_i^{k+1} \right) - \mathbf{b} \right), \mathbf{\lambda}^* \right\rangle$$
(58)

$$+2\beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\rangle$$
(59)

$$= -2\left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}, \boldsymbol{\lambda}^{*} \right\rangle + 2\beta_{k} \sum_{i=1}^{n} \sigma_{i}^{(k)} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*}, \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\rangle,$$
(60)

where we have used (43) in (57). Then we apply the identity

$$2\langle \mathbf{a}_{k+1} - \mathbf{a}^*, \mathbf{a}_{k+1} - \mathbf{a}_k \rangle = \|\mathbf{a}_{k+1} - \mathbf{a}^*\|^2 - \|\mathbf{a}_k - \mathbf{a}^*\|^2 + \|\mathbf{a}_{k+1} - \mathbf{a}_k\|^2$$
(61)
that (47)–(52) holds.

to see that (47)–(52) holds.

Proposition 2 For $\{(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k, \lambda^k)\}$ generated by Algorithm 1 and a KKT point $(\mathbf{x}_1^*, \ldots, \mathbf{x}_n^*, \lambda^*)$ of problem (1), the following inequality holds:

$$\beta_{k} \sum_{i=1}^{n} \sigma_{i}^{(k)} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} + \left\| \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{*} \right\|^{2}$$
(62)

$$\leq \beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\| \mathbf{x}_i^k - \mathbf{x}_i^* \right\|^2 + \left\| \boldsymbol{\lambda}^k - \boldsymbol{\lambda}^* \right\|^2 \tag{63}$$

$$-2\beta_k \sum_{i=1}^n \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, -\sigma_i^{(k)} \left(\mathbf{x}_i^{k+1} - \mathbf{u}_i^k \right) + \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*) \right\rangle$$
(64)

$$-\beta_{k}\sum_{i=1}^{n}\left(\sigma_{i}^{(k)}-n\beta_{k}\|\mathcal{A}_{i}\|^{2}\right)\left\|\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\right\|^{2}-\left\|\boldsymbol{\lambda}^{k}-\hat{\boldsymbol{\lambda}}^{k}\right\|^{2}.$$
(65)

Proof We continue from (51)–(52). As $\sigma_i^{(k)}(\mathbf{x}_i^k - \mathbf{u}_i^k) = \mathcal{A}_i^{\dagger}(\hat{\boldsymbol{\lambda}}^k)$, we have

$$-2\beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathbf{x}_i^k - \mathbf{u}_i^k \right\rangle + 2\left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^k, \boldsymbol{\lambda}^{k+1} \right\rangle$$
(66)

$$= -2\beta_k \sum_{i=1}^n \left\langle \mathcal{A}_i \left(\mathbf{x}_i^{k+1} - \mathbf{x}_i^* \right), \hat{\boldsymbol{\lambda}}^k \right\rangle + 2 \left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^k, \boldsymbol{\lambda}^{k+1} \right\rangle$$
(67)

$$= -2\beta_k \left\langle \sum_{i=1}^n \mathcal{A}_i \left(\mathbf{x}_i^{k+1} \right) - \sum_{i=1}^n \mathcal{A}_i \left(\mathbf{x}_i^* \right), \hat{\boldsymbol{\lambda}}^k \right\rangle + 2\left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^k, \boldsymbol{\lambda}^{k+1} \right\rangle$$
(68)

$$= -2\left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}, \, \hat{\boldsymbol{\lambda}}^{k} \right\rangle + 2\left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}, \, \boldsymbol{\lambda}^{k+1} \right\rangle \tag{69}$$

$$= 2\left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}, \, \boldsymbol{\lambda}^{k+1} - \hat{\boldsymbol{\lambda}}^{k} \right\rangle \tag{70}$$

$$= \|\boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}\|^{2} + \|\boldsymbol{\lambda}^{k+1} - \hat{\boldsymbol{\lambda}}^{k}\|^{2} - \|\boldsymbol{\lambda}^{k} - \hat{\boldsymbol{\lambda}}^{k}\|^{2}$$
(71)

$$= \|\boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}\|^{2} + \beta_{k}^{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right) \right\|^{2} - \|\boldsymbol{\lambda}^{k} - \hat{\boldsymbol{\lambda}}^{k}\|^{2}$$
(72)

$$\leq \|\boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}\|^{2} + \beta_{k}^{2} \left(\sum_{i=1}^{n} \|\mathcal{A}_{i}\| \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\| \right)^{2} - \|\boldsymbol{\lambda}^{k} - \hat{\boldsymbol{\lambda}}^{k}\|^{2}$$
(73)

$$\leq \|\boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}\|^{2} + n\beta_{k}^{2}\sum_{i=1}^{n}\|\mathcal{A}_{i}\|^{2}\left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2} - \|\boldsymbol{\lambda}^{k} - \hat{\boldsymbol{\lambda}}^{k}\|^{2}$$
(74)

Plugging the above into (51)–(52), we have (62)–(65).

Remark 4 Proposition 2 shows that the sequence $\{(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k, \lambda^k)\}$ is Fejér monotone. Proposition 2 is different from Lemma 1 in Supplementary Material of Lin et al. (2011) because for n > 2 we cannot obtain an (in)equality that is similar to Lemma 1 in Supplementary Material of Lin et al. (2011) such that each term with minus sign could be made non-positive. Such Fejér monotone (in)equalities are the corner stones for proving the convergence of Lagrange multiplier based optimization algorithms. As a result, we cannot prove the convergence of the naively generalized LADM for the multi-block case.

Then we have the following proposition.

Proposition 3 Let $\sigma_i^{(k)} = \eta_i \beta_k$, i = 1, ..., n. If $\{\beta_k\}$ is non-decreasing, $\eta_i > n \|\mathcal{A}_i\|^2$, i = 1, ..., n, $\{(\mathbf{x}_1^k, ..., \mathbf{x}_n^k, \lambda^k)\}$ is generated by Algorithm 1, and $(\mathbf{x}_1^*, ..., \mathbf{x}_n^*, \lambda^*)$ is any KKT point of problem (1), then

(1)
$$\left\{\sum_{i=1}^{n} \eta_{i} \|\mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*}\|^{2} + \beta_{k}^{-2} \|\boldsymbol{\lambda}^{k} - \boldsymbol{\lambda}^{*}\|^{2}\right\}$$
 is nonnegative and non-increasing.

(2)
$$\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\| \to 0, i = 1, \dots, n, \text{ and } \beta_{k}^{-1} \|\boldsymbol{\lambda}^{k} - \boldsymbol{\hat{\lambda}}^{k}\| \to 0.$$

(2) $\sum_{i=1}^{i} \sum_{j=1}^{i} \rho^{-1} |\mathbf{x}_{i}^{k+1} - \mathbf{x}_{j}^{k} - \rho^{(k)} (\mathbf{x}_{i}^{k+1} - \mathbf{x}_{j}^{k})| \to 0.$

(3)
$$\sum_{k=1}^{+\infty} \beta_k^{-1} \left(\mathbf{x}_i^{k+1} - \mathbf{x}_i^*, -\sigma_i^{(\kappa)} (\mathbf{x}_i^{k+1} - \mathbf{u}_i^k) + \mathcal{A}_i^{\dagger} (\boldsymbol{\lambda}^*) \right) < +\infty, \quad i = 1, \dots, n.$$

Proof We divide both sides of (62)–(65) by β_k^2 to have

$$\sum_{i=1}^{n} \eta_{i} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} + \beta_{k}^{-2} \left\| \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{*} \right\|^{2}$$
(75)

$$\leq \sum_{i=1}^{n} \eta_{i} \|\mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*}\|^{2} + \beta_{k}^{-2} \|\boldsymbol{\lambda}^{k} - \boldsymbol{\lambda}^{*}\|^{2}$$
(76)

$$-2\beta_{k}^{-1}\sum_{i=1}^{n}\left\langle \mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{*},-\sigma_{i}^{(k)}\left(\mathbf{x}_{i}^{k+1}-\mathbf{u}_{i}^{k}\right)+\mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*})\right\rangle$$
(77)

$$-\sum_{i=1}^{n} \left(\eta_{i} - n \|\mathcal{A}_{i}\|^{2}\right) \left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2}$$
(78)

$$-\beta_k^{-2} \|\boldsymbol{\lambda}^k - \hat{\boldsymbol{\lambda}}^k\|^2.$$
(79)

Then by (46), $\eta_i > n \|\mathcal{A}_i\|^2$ and the non-decrement of $\{\beta_k\}$, we can easily obtain (1). Second, we sum both sides of (75)–(79) over *k* to have

$$2\sum_{k=0}^{+\infty}\beta_k^{-1}\sum_{i=1}^n \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}^*, -\sigma_i^{(k)}\left(\mathbf{x}_i^{k+1} - \mathbf{u}_i^k\right) + \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*) \right\rangle$$
(80)

+
$$\sum_{i=1}^{n} (\eta_i - n \|\mathcal{A}_i\|^2) \sum_{k=0}^{+\infty} \|\mathbf{x}_i^{k+1} - \mathbf{x}_i^k\|^2$$
 (81)

$$+\sum_{k=0}^{+\infty}\beta_{k}^{-2}\|\mathbf{\lambda}^{k}-\hat{\mathbf{\lambda}}^{k}\|^{2}$$
(82)

$$\leq \sum_{i=1}^{n} \eta_{i} \left\| \mathbf{x}_{i}^{0} - \mathbf{x}^{*} \right\|^{2} + \beta_{0}^{-2} \| \boldsymbol{\lambda}^{0} - \boldsymbol{\lambda}^{*} \|^{2}.$$
(83)

Then (2) and (3) can be easily deduced.

Now we are ready to prove Theorem 1. The proof resembles that in (Lin et al. 2011).

Proof of Theorem 1 By Proposition 3-(1) and the boundedness of $\{\beta_k\}, \{(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k, \boldsymbol{\lambda}^k)\}$ is bounded, hence has an accumulation point, say $(\mathbf{x}_1^{k_j}, \dots, \mathbf{x}_n^{k_j}, \boldsymbol{\lambda}^{k_j}) \to (\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty}, \boldsymbol{\lambda}^{\infty}).$ We accomplish the proof in two steps.

1. We first prove that $(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty}, \boldsymbol{\lambda}^{\infty})$ is a KKT point of problem (1). By Proposition 3-(2),

$$\sum_{i=1}^{n} \mathcal{A}_{i}\left(\mathbf{x}_{i}^{k}\right) - \mathbf{b} = \beta_{k}^{-1}\left(\hat{\boldsymbol{\lambda}}^{k} - \boldsymbol{\lambda}^{k}\right) \to 0.$$

So any accumulation point of $\{(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k)\}$ is a feasible solution. Since $-\sigma_i^{(k_j-1)}(\mathbf{x}_i^{k_j} - \mathbf{u}_i^{k_j-1}) \in \partial f_i(\mathbf{x}_i^{k_j})$, we have

$$\sum_{i=1}^{n} f_{i}\left(\mathbf{x}_{i}^{k_{j}}\right) \leq \sum_{i=1}^{n} f_{i}\left(\mathbf{x}_{i}^{*}\right) + \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{k_{j}} - \mathbf{x}_{i}^{*}, -\sigma_{i}^{(k_{j}-1)}\left(\mathbf{x}_{i}^{k_{j}} - \mathbf{u}_{i}^{k_{j}-1}\right)\right\rangle$$
$$= \sum_{i=1}^{n} f_{i}\left(\mathbf{x}_{i}^{*}\right) + \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{k_{j}} - \mathbf{x}_{i}^{*}, -\eta_{i}\beta_{k_{j}-1}\left(\mathbf{x}_{i}^{k_{j}} - \mathbf{x}_{i}^{k_{j}-1}\right) - \mathcal{A}_{i}^{\dagger}\left(\hat{\boldsymbol{\lambda}}^{k_{j}-1}\right)\right\rangle$$

Let $j \to +\infty$. By observing Proposition 3-(2) and the boundedness of $\{\beta_k\}$, we have

$$\sum_{i=1}^{n} f_i \left(\mathbf{x}_i^{\infty} \right) \leq \sum_{i=1}^{n} f_i \left(\mathbf{x}_i^{*} \right) + \sum_{i=1}^{n} \left\langle \mathbf{x}_i^{\infty} - \mathbf{x}_i^{*}, -\mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^{\infty}) \right\rangle$$
$$= \sum_{i=1}^{n} f_i \left(\mathbf{x}_i^{*} \right) - \sum_{i=1}^{n} \left\langle \mathcal{A} \left(\mathbf{x}_i^{\infty} - \mathbf{x}_i^{*} \right), \boldsymbol{\lambda}^{\infty} \right\rangle$$
$$= \sum_{i=1}^{n} f_i \left(\mathbf{x}_i^{*} \right) - \left\langle \sum_{i=1}^{n} \mathcal{A} \left(\mathbf{x}_i^{\infty} \right) - \mathbf{b}, \boldsymbol{\lambda}^{\infty} \right\rangle$$
$$= \sum_{i=1}^{n} f_i \left(\mathbf{x}_i^{*} \right).$$

So we conclude that $(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty})$ is an optimal solution to (1). Again by $-\sigma_i^{(k_j-1)}(\mathbf{x}_i^{k_j} - \mathbf{u}_i^{k_j-1}) \in \partial f_i(\mathbf{x}_i^{k_j})$ we have

$$f_{i}(\mathbf{x}) \geq f_{i}\left(\mathbf{x}_{i}^{k_{j}}\right) + \left\langle \mathbf{x} - \mathbf{x}_{i}^{k_{j}}, -\sigma_{i}^{(k_{j}-1)}(\mathbf{x}_{i}^{k_{j}} - \mathbf{u}_{i}^{k_{j}-1})\right\rangle$$

= $f_{i}\left(\mathbf{x}_{i}^{k_{j}}\right) + \left\langle \mathbf{x} - \mathbf{x}_{i}^{k_{j}}, -\eta_{i}\beta_{k_{j}-1}\left(\mathbf{x}_{i}^{k_{j}} - \mathbf{x}_{i}^{k_{j}-1}\right) - \mathcal{A}_{i}^{\dagger}\left(\hat{\boldsymbol{\lambda}}^{k_{j}-1}\right)\right\rangle.$

Fixing **x** and letting $j \to +\infty$, we see that

$$f_i(\mathbf{x}) \ge f_i(\mathbf{x}_i^{\infty}) + \left\langle \mathbf{x} - \mathbf{x}_i^{\infty}, -\mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^{\infty}) \right\rangle, \quad \forall \mathbf{x}.$$

Springer

So $-\mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^{\infty}) \in \partial f_i(\mathbf{x}_i^{\infty}), i = 1, ..., n$. Thus $(\mathbf{x}_1^{\infty}, ..., \mathbf{x}_n^{\infty}, \boldsymbol{\lambda}^{\infty})$ is a KKT point of problem (1).

2. We next prove that the whole sequence $\{(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k, \lambda^k)\}$ converges to $(\mathbf{x}_1^\infty, \ldots, \mathbf{x}_n^\infty)$ **λ**[∞]).

By choosing $(\mathbf{x}_1^*, \ldots, \mathbf{x}_n^*, \boldsymbol{\lambda}^*) = (\mathbf{x}_1^{\infty}, \ldots, \mathbf{x}_n^{\infty}, \boldsymbol{\lambda}^{\infty})$ in Proposition 3, we have

$$\sum_{i=1}^{n} \eta_i \left\| \mathbf{x}_i^{k_j} - \mathbf{x}_i^{\infty} \right\|^2 + \beta_{k_j}^{-2} \left\| \boldsymbol{\lambda}^{k_j} - \boldsymbol{\lambda}^{\infty} \right\|^2 \to 0.$$

By Proposition 3-(1), we readily have

$$\sum_{i=1}^{n} \eta_i \left\| \mathbf{x}_i^k - \mathbf{x}_i^\infty \right\|^2 + \beta_k^{-2} \left\| \boldsymbol{\lambda}^k - \boldsymbol{\lambda}^\infty \right\|^2 \to 0.$$

So $(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k, \boldsymbol{\lambda}^k) \to (\mathbf{x}_1^\infty, \dots, \mathbf{x}_n^\infty, \boldsymbol{\lambda}^\infty)$. As $(\mathbf{x}_1^\infty, \dots, \mathbf{x}_n^\infty, \boldsymbol{\lambda}^\infty)$ can be an arbitrary accumulation point of $\{(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k, \boldsymbol{\lambda}^k)\}$, we conclude that $\{(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k, \boldsymbol{\lambda}^k)\}$ converge to a KKT point of problem (1).

Appendix 2: Proof of Theorem 2

We first have the following proposition.

Proposition 4 If $\{\beta_k\}$ is non-decreasing and unbounded, $\eta_i > n \|A_i\|^2$ and $\partial f_i(\mathbf{x})$ is bounded for i = 1, ..., n, then Proposition 3 holds and

$$\beta_k^{-1} \boldsymbol{\lambda}^k \to 0. \tag{84}$$

Proof As the conditions here are stricter than those in Proposition 3, Proposition 3 holds. Then we have that $\{\beta_k^{-1} \| \boldsymbol{\lambda}^k - \boldsymbol{\lambda}^* \|\}$ is bounded due to Proposition 3-(1). So $\{\beta_k^{-1} \boldsymbol{\lambda}^k\}$ is bounded due to $\beta_k^{-1} \| \hat{\boldsymbol{\lambda}}^k \| \le \beta_k^{-1} \| \boldsymbol{\lambda}^k - \boldsymbol{\lambda}^* \| + \beta_k^{-1} \| \boldsymbol{\lambda}^* \|$. $\{\beta_k^{-1} \hat{\boldsymbol{\lambda}}^k\}$ is also bounded thanks to Proposition 3-(2).

We rewrite Lemma 2 as

$$-\eta_i \left(\mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right) - \mathcal{A}_i^{\dagger} \left(\beta_k^{-1} \hat{\boldsymbol{\lambda}}^k \right) \in \beta_k^{-1} \partial f_i \left(\mathbf{x}_i^{k+1} \right), \quad i = 1, \dots, n.$$
(85)

Then by the boundedness of $\partial f_i(x)$, the unboundedness of $\{\beta_k\}$ and Proposition 3-(2), letting $k \to +\infty$, we have that

$$\mathcal{A}_{i}^{\dagger}(\check{\boldsymbol{\lambda}}^{\infty}) = 0, \quad i = 1, \dots, n.$$
(86)

where $\check{\boldsymbol{\lambda}}^{\infty}$ is any accumulation point of $\{\beta_k^{-1}\hat{\boldsymbol{\lambda}}^k\}$, which is the same as that of $\{\beta_k^{-1}\boldsymbol{\lambda}^k\}$ due to Proposition 3-(2).

Recall that we have assumed that the mapping $\mathcal{A}(\mathbf{x}_1, \dots, \mathbf{x}_n) \equiv \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i)$ is onto. So $\bigcap_{i=1}^{n} null(\mathcal{A}_{i}^{\dagger}) = 0.$ Therefore by (86), $\check{\lambda}^{\infty} = 0.$

Based on Proposition 4, we can prove Theorem 2 as follows.

Proof of Theorem 2 When $\{\beta_k\}$ is bounded, the convergence has been proven in Theorem 1. In the following, we only focus on the case that $\{\beta_k\}$ is unbounded.

By Proposition 3-(1), $\{(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k)\}$ is bounded, hence has at least one accumulation point $(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty})$. By Proposition 3-(2), $(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty})$ is a feasible solution.

Since $\sum_{k=1}^{+\infty} \beta_k^{-1} = +\infty$ and Proposition 3-(3), there exists a subsequence $\{(\mathbf{x}_1^{k_j}, \dots, \mathbf{x}_n^{k_j})\}$ such that

$$\left|\mathbf{x}_{i}^{k_{j}}-\mathbf{x}_{i}^{*},-\sigma_{i}^{(k_{j}-1)}\left(\mathbf{x}_{i}^{k_{j}}-\mathbf{u}_{i}^{k_{j}-1}\right)+\mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*})\right|\rightarrow0,\quad i=1,\ldots,n.$$
(87)

As $\mathbf{p}_i^{k_j} \equiv -\sigma_i^{(k_j-1)}(\mathbf{x}_i^{k_j} - \mathbf{u}_i^{k_j-1}) \in \partial f_i(\mathbf{x}_i^{k_j})$ and ∂f_i is bounded, we may assume that

$$\mathbf{x}_i^{k_j} \to \mathbf{x}_i^{\infty}$$
 and $\mathbf{p}_i^{k_j} \to \mathbf{p}_i^{\infty}$.

It can be easily proven that

$$\mathbf{p}_i^{\infty} \in \partial f_i(\mathbf{x}_i^{\infty}).$$

Then letting $j \to \infty$ in (87), we have

$$\left\langle \mathbf{x}_{i}^{\infty} - \mathbf{x}_{i}^{*}, \mathbf{p}_{i}^{\infty} + \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle = 0, \quad i = 1, \dots, n.$$
 (88)

Then by $\mathbf{p}_i^{k_j} \in \partial f_i(\mathbf{x}_i^{k_j})$,

$$\sum_{i=1}^{n} f_i\left(\mathbf{x}_i^{k_j}\right) \le \sum_{i=1}^{n} f_i\left(\mathbf{x}_i^*\right) + \sum_{i=1}^{n} \left\langle \mathbf{x}_i^{k_j} - \mathbf{x}_i^*, \mathbf{p}_i^{k_j} \right\rangle.$$
(89)

Letting $j \to \infty$ and making use of (88), we have

$$\sum_{i=1}^{n} f_{i} \left(\mathbf{x}_{i}^{\infty} \right) \leq \sum_{i=1}^{n} f_{i} \left(\mathbf{x}_{i}^{*} \right) + \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{\infty} - \mathbf{x}_{i}^{*}, \mathbf{p}_{i}^{\infty} \right\rangle$$
$$= \sum_{i=1}^{n} f_{i} \left(\mathbf{x}_{i}^{*} \right) - \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{\infty} - \mathbf{x}_{i}^{*}, \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle$$
$$= \sum_{i=1}^{n} f_{i} \left(\mathbf{x}_{i}^{*} \right) - \sum_{i=1}^{n} \left\langle \mathcal{A}_{i} \left(\mathbf{x}_{i}^{\infty} - \mathbf{x}_{i}^{*} \right), \boldsymbol{\lambda}^{*} \right\rangle$$
$$= \sum_{i=1}^{n} f_{i} \left(\mathbf{x}_{i}^{*} \right). \tag{90}$$

So together with the feasibility of $\{(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty})\}$ we have that $\{(\mathbf{x}_1^{k_j}, \dots, \mathbf{x}_n^{k_j})\}$ converges to an optimal solution $\{(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty})\}$ to (1). Finally, we set $\mathbf{x}_i^* = \mathbf{x}_i^{\infty}$ and λ^* be the corresponding Lagrange multiplier λ^{∞} in Propo-

sition 3. By Proposition 4, we have that

$$\sum_{i=1}^{n} \eta_i \left\| \mathbf{x}_i^{k_j} - \mathbf{x}_i^{\infty} \right\|^2 + \beta_{k_j}^{-2} \left\| \boldsymbol{\lambda}^{k_j} - \boldsymbol{\lambda}^{\infty} \right\|^2 \to 0.$$

By Proposition 3-(1), we readily have

$$\sum_{i=1}^n \eta_i \|\mathbf{x}_i^k - \mathbf{x}_i^\infty\|^2 + \beta_k^{-2} \|\boldsymbol{\lambda}^k - \boldsymbol{\lambda}^\infty\|^2 \to 0.$$

So $(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k) \to (\mathbf{x}_1^\infty, \ldots, \mathbf{x}_n^\infty)$.

Springer

Appendix 3: Proof of Theorem 3

Proof of Theorem 3 We first prove that there exist linear mappings \mathcal{B}_i , i = 1, ..., n, such that \mathcal{B}_i 's are not all zeros and $\sum_{i=1}^n \mathcal{B}_i \mathcal{A}_i^{\dagger} = 0$. Indeed, $\sum_{i=1}^n \mathcal{B}_i \mathcal{A}_i^{\dagger} = 0$ is equivalent to

$$\sum_{i=1}^{n} \mathbf{B}_i \mathbf{A}_i^T = 0, \tag{91}$$

where \mathbf{A}_i and \mathbf{B}_i are the matrix representations of \mathcal{A}_i and \mathcal{B}_i , respectively. (91) can be further written as

$$(\mathbf{A}_1 \ \dots \ \mathbf{A}_n) \begin{pmatrix} \mathbf{B}_1^T \\ \vdots \\ \mathbf{B}_n^T \end{pmatrix} = 0.$$
(92)

Recall that we have assumed that the solution to $\sum_{i=1}^{n} A_i(\mathbf{x}_i) = \mathbf{b}$ is non-unique. So $(\mathbf{A}_1 \dots \mathbf{A}_n)$ is not full column rank hence (92) has nonzero solutions. Thus there exist \mathcal{B}_i 's such that they are not all zeros and $\sum_{i=1}^{n} \mathcal{B}_i A_i^{\dagger} = 0$.

By Lemma 2,

$$-\sigma_i^{(k)}\left(\mathbf{x}_i^{k+1} - \mathbf{u}_i^k\right) \in \partial f_i\left(\mathbf{x}_i^{k+1}\right), \quad i = 1, \dots, n.$$
(93)

As ∂f_i is bounded, $i = 1, \ldots, n$, so is

$$\sum_{i=1}^{n} \mathcal{B}_{i} \left(\sigma_{i}^{(k)} \left(\mathbf{x}_{i}^{k+1} - \mathbf{u}_{i}^{k} \right) \right) = \beta_{k} \left(\mathbf{v}^{k+1} - \mathbf{v}^{k} \right),$$
(94)

where $\mathbf{v}^k = \phi(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k)$ and

$$\phi(\mathbf{x}_1,\ldots,\mathbf{x}_n) = \sum_{i=1}^n \eta_i \mathcal{B}_i(\mathbf{x}_i).$$
(95)

In (94) we have utilized $\sum_{i=1}^{n} \mathcal{B}_i \mathcal{A}_i^{\dagger} = 0$ to cancel $\hat{\lambda}^k$, whose boundedness is uncertain.

Then we have that there exists a constant C > 0 such that

$$\|\mathbf{v}^{k+1} - \mathbf{v}^k\| \le C\beta_k^{-1}.$$
(96)

If $\sum_{k=1}^{+\infty} \beta_k^{-1} < +\infty$, then $\{\mathbf{v}^k\}$ is a Cauchy sequence, hence has a limit \mathbf{v}^{∞} . Define $\mathbf{v}^* = \phi(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*)$, where $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*)$ is any optimal solution. Then

$$\|\mathbf{v}^{\infty} - \mathbf{v}^*\| = \left\|\mathbf{v}^0 + \sum_{k=0}^{\infty} (\mathbf{v}^{k+1} - \mathbf{v}^k) - \mathbf{v}^*\right\|$$
(97)

$$\geq \|\mathbf{v}^{0} - \mathbf{v}^{*}\| - \sum_{k=0}^{\infty} \|\mathbf{v}^{k+1} - \mathbf{v}^{k}\|$$
(98)

$$\geq \|\mathbf{v}^{0} - \mathbf{v}^{*}\| - C \sum_{k=0}^{\infty} \beta_{k}^{-1}.$$
(99)

So if $(\mathbf{x}_1^0, \dots, \mathbf{x}_n^0)$ is initialized badly such that

$$\|\mathbf{v}^{0} - \mathbf{v}^{*}\| > C \sum_{k=0}^{\infty} \beta_{k}^{-1},$$
(100)

then $\|\mathbf{v}^{\infty} - \mathbf{v}^*\| > 0$, which implies that $(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k)$ cannot converge to $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*)$. Note that (100) is possible because ϕ is not a zero mapping given the conditions on \mathcal{B}_i .

Appendix 4: Proofs of Proposition 1 and Theorem 4

Proof of Proposition 1 If $\tilde{\mathbf{x}}$ is optimal, it is easy to check that (24) holds. Since $-\mathcal{A}_i^{\dagger}(\lambda^*) \in \partial f_i(\mathbf{x}_i^*)$, we have

$$f(\tilde{\mathbf{x}}) - f(\mathbf{x}^*) + \sum_{i=1}^n \left\langle \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*), \, \tilde{\mathbf{x}}_i - \mathbf{x}_i^* \right\rangle \ge 0.$$

So if (24) holds, we have

$$f(\tilde{\mathbf{x}}) - f(\mathbf{x}^*) + \sum_{i=1}^n \left\langle \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*), \, \tilde{\mathbf{x}}_i - \mathbf{x}_i^* \right\rangle = 0, \tag{101}$$

$$\sum_{i=1}^{n} \mathcal{A}_i(\tilde{\mathbf{x}}_i) - \mathbf{b} = 0.$$
(102)

With (102), we have

$$\sum_{i=1}^{n} \left\langle \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}), \, \tilde{\mathbf{x}}_{i} - \mathbf{x}_{i}^{*} \right\rangle = \sum_{i=1}^{n} \left\langle \boldsymbol{\lambda}^{*}, \, \mathcal{A}_{i}\left(\tilde{\mathbf{x}}_{i} - \mathbf{x}_{i}^{*} \right) \right\rangle = \left\langle \boldsymbol{\lambda}^{*}, \, \sum_{i=1}^{n} \mathcal{A}_{i}\left(\tilde{\mathbf{x}}_{i} - \mathbf{x}_{i}^{*} \right) \right\rangle = 0.$$
(103)

So (101) reduces to $f(\tilde{\mathbf{x}}) = f(\mathbf{x}^*)$. As $\tilde{\mathbf{x}}$ satisfies the feasibility condition, it is an optimal solution to (1).

Proof of Theorem 4 We first deduce

$$\begin{split} & \left\|\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1}\right) - \mathbf{b}\right\|^{2} \\ &= \left\|\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k}\right) - \mathbf{b} + \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right)\right\|^{2} \\ &\leq \left(\left\|\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k}\right) - \mathbf{b}\right\| + \sum_{i=1}^{n} \left\|\mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right)\right\|\right)^{2} \\ &\leq (n+1) \left(\left\|\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k}\right) - \mathbf{b}\right\|^{2} + \sum_{i=1}^{n} \left\|\mathcal{A}_{i}\right\|^{2} \left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2}\right) \\ &\leq (n+1) \left(\left\|\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k}\right) - \mathbf{b}\right\|^{2} + \max\left\{\frac{\left\|\mathcal{A}_{i}\right\|^{2}}{\eta_{i} - n \left\|\mathcal{A}_{i}\right\|^{2}}\right\} \sum_{i=1}^{n} \left(\eta_{i} - n \left\|\mathcal{A}_{i}\right\|^{2}\right) \left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2}\right) \end{split}$$

$$\leq (n+1) \max\left\{1, \left\{\frac{\|\mathcal{A}_{i}\|^{2}}{\eta_{i}-n\|\mathcal{A}_{i}\|^{2}}\right\}\right\} \left(\beta_{k}^{-2} \|\boldsymbol{\lambda}_{k}-\hat{\boldsymbol{\lambda}}_{k}\|^{2} + \sum_{i=1}^{n} \left(\eta_{i}-n\|\mathcal{A}_{i}\|^{2}\right) \|\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\|^{2}\right)$$
$$= \alpha^{-1} \left(\beta_{k}^{-2} \|\boldsymbol{\lambda}_{k}-\hat{\boldsymbol{\lambda}}_{k}\|^{2} + \sum_{i=1}^{n} \left(\eta_{i}-n\|\mathcal{A}_{i}\|^{2}\right) \|\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\|^{2}\right).$$
(104)

By Proposition 2, we have

$$\beta_{k}^{-1} \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*}, -\sigma_{i}^{(k)} \left(\mathbf{x}_{i}^{k+1} - \mathbf{u}_{i}^{k} \right) + \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle \\ + \frac{1}{2} \sum_{i=1}^{n} \left(\eta_{i} - n \|\mathcal{A}_{i}\|^{2} \right) \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} + \frac{1}{2} \beta_{k}^{-2} \|\boldsymbol{\lambda}^{k} - \hat{\boldsymbol{\lambda}}^{k}\|^{2} \\ \leq \frac{1}{2} \left(\sum_{i=1}^{n} \eta_{i} \| \mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*} \|^{2} + \beta_{k}^{-2} \| \boldsymbol{\lambda}^{k} - \boldsymbol{\lambda}^{*} \|^{2} \right) \\ - \frac{1}{2} \left(\sum_{i=1}^{n} \eta_{i} \| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \|^{2} + \beta_{k+1}^{-2} \| \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{*} \|^{2} \right).$$
(105)

So by Lemma 2 and combining the above inequalities, we have

$$\beta_{k}^{-1} \left(f(\mathbf{x}^{k+1}) - f(\mathbf{x}^{*}) + \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*}, \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle + \frac{\alpha \beta_{0}}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} \right) - \mathbf{b} \right\|^{2} \right)$$

$$\leq \beta_{k}^{-1} \left(\sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*}, -\sigma_{i}^{(k)} \left(\mathbf{x}_{i}^{k+1} - \mathbf{u}_{i}^{k} \right) \right\rangle + \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*}, \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle \right)$$

$$+ \frac{\alpha}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} \right) - \mathbf{b} \right\|^{2}$$

$$\leq \beta_{k}^{-1} \sum_{i=1}^{n} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*}, -\sigma_{i}^{(k)} \left(\mathbf{x}_{i}^{k+1} - \mathbf{u}_{i}^{k} \right) + \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle$$

$$+ \frac{1}{2} \beta_{k}^{-2} \left\| \boldsymbol{\lambda}_{k} - \hat{\boldsymbol{\lambda}}_{k} \right\|^{2} + \frac{1}{2} \sum_{i=1}^{n} \left(\eta_{i} - n \left\| \mathcal{A}_{i} \right\|^{2} \right) \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2}$$

$$\leq \frac{1}{2} \left(\sum_{i=1}^{n} \eta_{i} \left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*} \right\|^{2} + \beta_{k}^{-2} \left\| \boldsymbol{\lambda}^{k} - \boldsymbol{\lambda}^{*} \right\|^{2} \right)$$

$$- \frac{1}{2} \left(\sum_{i=1}^{n} \eta_{i} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} + \beta_{k-1}^{-2} \left\| \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{*} \right\|^{2} \right).$$
(106)

Here we use the fact that $\beta_k \ge \beta_0$, which is guaranteed by (18) and (20). Summing the above inequalities from k = 0 to K, and dividing both sides with $\sum_{k=0}^{K} \beta_k^{-1}$, we have

$$\sum_{k=0}^{K} \gamma_k f(\mathbf{x}^{k+1}) - f(\mathbf{x}^*) + \sum_{i=1}^{n} \left\langle \sum_{k=0}^{K} \gamma_k \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*) \right\rangle$$
(107)

D Springer

$$+\frac{\alpha\beta_0}{2}\sum_{k=0}^{K}\gamma_k \left\|\sum_{i=1}^{n}\mathcal{A}_i\left(\mathbf{x}_i^{k+1}\right) - \mathbf{b}\right\|^2$$
(108)

$$\leq \frac{1}{2\sum_{k=0}^{K}\beta_{k}^{-1}} \left(\sum_{i=1}^{n}\eta_{i} \|\mathbf{x}_{i}^{0}-\mathbf{x}_{i}^{*}\|^{2} + \beta_{0}^{-2} \|\boldsymbol{\lambda}^{0}-\boldsymbol{\lambda}^{*}\|^{2} \right).$$
(109)

Next, by the convexity of f and the squared Frobenius norm $\|\cdot\|^2$, we have

$$f(\bar{\mathbf{x}}^{K}) - f(\mathbf{x}^{*}) + \sum_{i=1}^{n} \left\langle \bar{\mathbf{x}}_{i}^{K} - \mathbf{x}_{i}^{*}, \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle + \frac{\alpha \beta_{0}}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i}(\bar{\mathbf{x}}_{i}^{K}) - \mathbf{b} \right\|^{2}$$
(110)

$$\leq \sum_{k=0}^{K} \gamma_k f(\mathbf{x}^{k+1}) - f(\mathbf{x}^*) + \sum_{i=1}^{n} \left\langle \sum_{k=0}^{K} \gamma_k \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathcal{A}_i^{\dagger}(\boldsymbol{\lambda}^*) \right\rangle$$
(111)

$$+ \frac{\alpha\beta_0}{2} \sum_{k=0}^{K} \gamma_k \left\| \sum_{i=1}^{n} \mathcal{A}_i\left(\mathbf{x}_i^{k+1}\right) - \mathbf{b} \right\|^2.$$
(112)

Combining (107)–(109) and (110)–(112), we have

$$f(\bar{\mathbf{x}}^{K}) - f(\mathbf{x}^{*}) + \sum_{i=1}^{n} \left\langle \bar{\mathbf{x}}_{i}^{K} - \mathbf{x}_{i}^{*}, \mathcal{A}_{i}^{\dagger}(\boldsymbol{\lambda}^{*}) \right\rangle + \frac{\alpha\beta_{0}}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i}\left(\bar{\mathbf{x}}_{i}^{K} \right) - \mathbf{b} \right\|^{2}$$
(113)

$$\leq \frac{1}{2\sum_{k=0}^{K}\beta_{k}^{-1}} \left(\sum_{i=1}^{n}\eta_{i} \left\| \mathbf{x}_{i}^{0} - \mathbf{x}_{i}^{*} \right\|^{2} + \beta_{0}^{-2} \left\| \boldsymbol{\lambda}^{0} - \boldsymbol{\lambda}^{*} \right\|^{2} \right).$$
(114)

Appendix 5: Proof of Theorem 5

We only need to prove the following proposition. Then by the same technique for proving Theorem 1, we can prove Theorem 5.

Proposition 5 For $\{(\mathbf{x}_1^k, \ldots, \mathbf{x}_{2n}^k, \lambda^k)\}$ generated by Algorithm 2 and a KKT point $(\mathbf{x}_1^*, \ldots, \mathbf{x}_{2n}^*, \lambda^*)$ of problem (27), we have that

$$\beta_k \sum_{i=1}^{2n} \sigma_i^{(k)} \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i^* \right\|^2 + \| \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^* \|^2$$
(115)

$$\leq \beta_k \sum_{i=1}^n \sigma_i^{(k)} \left\| \mathbf{x}_i^k - \mathbf{x}_i^* \right\|^2 + \left\| \boldsymbol{\lambda}^k - \boldsymbol{\lambda}^* \right\|^2 \tag{116}$$

$$-2\beta_k \sum_{i=1}^{2n} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, -\sigma_i^{(k)} \left(\mathbf{x}_i^{k+1} - \mathbf{u}_i^k \right) + \hat{\mathcal{A}}_i^{\dagger}(\boldsymbol{\lambda}^*) \right\rangle$$
(117)

$$-\beta_k \sum_{i=1}^n \left(\sigma_i^{(k)} - \beta_k (n \|\mathcal{A}_i\|^2 + 2) \right) \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\|^2$$
(118)

$$-\beta_k \sum_{i=n+1}^{2n} \left(\sigma_i^{(k)} - 2\beta_k\right) \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\|^2$$
(119)

$$-\|\boldsymbol{\lambda}^k - \hat{\boldsymbol{\lambda}}^k\|^2.$$
(120)

Proof We continue from (72):

$$-2\beta_k \sum_{i=1}^{2n} \sigma_i^{(k)} \left\langle \mathbf{x}_i^{k+1} - \mathbf{x}_i^*, \mathbf{x}_i^k - \mathbf{u}_i^k \right\rangle + 2\left\langle \boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^k, \boldsymbol{\lambda}^{k+1} \right\rangle$$
(121)

$$= \|\boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}\|^{2} + \beta_{k}^{2} \left\| \sum_{i=1}^{2n} \hat{\mathcal{A}}_{i} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right) \right\|^{2} - \|\boldsymbol{\lambda}^{k} - \hat{\boldsymbol{\lambda}}^{k}\|^{2}$$
(122)

$$= \left\|\boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}\right\|^{2} + \beta_{k}^{2} \left\|\sum_{i=1}^{n} \mathcal{A}_{i}\left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right)\right\|^{2}$$
(123)

$$+\beta_{k}^{2}\sum_{i=1}^{n}\left\|\left(\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\right)-\left(\mathbf{x}_{n+i}^{k+1}-\mathbf{x}_{n+i}^{k}\right)\right\|^{2}-\|\boldsymbol{\lambda}^{k}-\hat{\boldsymbol{\lambda}}^{k}\|^{2}$$
(124)

$$\leq \|\boldsymbol{\lambda}^{k+1} - \boldsymbol{\lambda}^{k}\|^{2} + n\beta_{k}^{2} \sum_{i=1}^{n} \|\mathcal{A}_{i}\|^{2} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2}$$
(125)

$$+2\beta_{k}^{2}\sum_{i=1}^{n}\left(\left\|\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\right\|^{2}+\left\|\mathbf{x}_{n+i}^{k+1}-\mathbf{x}_{n+i}^{k}\right\|^{2}\right)-\left\|\boldsymbol{\lambda}^{k}-\hat{\boldsymbol{\lambda}}^{k}\right\|^{2}.$$
 (126)

Then we can have (115)-(120).

Appendix 6: Proof of Theorem 6

To prove Theorem 6, we need the following proposition:

Proposition 6 For $\{(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k, \lambda^k)\}$ generated by Algorithm 3 and a KKT point $(\mathbf{x}_1^*, \ldots, \mathbf{x}_n^*, \lambda^*)$ of problem (1) with f_i described in Sect. 5, we have that

$$\sum_{i=1}^{n} \left(f_i \left(\mathbf{x}_i^{k+1} \right) - f_i \left(\mathbf{x}_i^* \right) + \left\langle \mathcal{A}_i^{\dagger}(\lambda^*), \mathbf{x}_i^{k+1} - \mathbf{x}_i^* \right\rangle \right)$$

$$\leq \frac{1}{2} \sum_{i=1}^{n} \tau_i^{(k)} \left(\left\| \mathbf{x}_i^k - \mathbf{x}_i^* \right\|^2 - \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i^* \right\|^2 \right)$$
(127)

$$+ \frac{1}{2\beta_k} \left(\|\lambda^k - \lambda^*\|^2 - \|\lambda^{k+1} - \lambda^*\|^2 \right)$$
(128)

$$-\frac{1}{2}\sum_{i=1}^{n} \left(\tau_{i}^{(k)} - L_{i} - n\beta_{k} \|\mathcal{A}_{i}\|^{2}\right) \left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2} - \frac{1}{2\beta_{k}} \|\hat{\lambda}^{k} - \lambda^{k}\|^{2}$$
(129)

Proof It can be observed that

$$0 \in \partial h_i\left(\mathbf{x}_i^{k+1}\right) + \nabla g_i\left(\mathbf{x}_i^k\right) + \mathcal{A}_i^{\dagger}(\hat{\lambda}^k) + \tau_i^{(k)}\left(\mathbf{x}_i^{k+1} - \mathbf{x}_i^k\right).$$

So we have

$$h_{i}(\mathbf{x}_{i}) - h_{i}\left(\mathbf{x}_{i}^{k+1}\right) \geq \left\langle -\nabla g_{i}\left(\mathbf{x}_{i}^{k}\right) - \mathcal{A}_{i}^{\dagger}(\hat{\lambda}^{k}) - \tau_{i}^{(k)}\left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right), \mathbf{x}_{i} - \mathbf{x}_{i}^{k+1}\right\rangle, \quad \forall \mathbf{x}_{i},$$

and

$$\sum_{i=1}^{n} f_{i}\left(\mathbf{x}_{i}^{k+1}\right) = \sum_{i=1}^{n} \left(h_{i}\left(\mathbf{x}_{i}^{k+1}\right) + g_{i}\left(\mathbf{x}_{i}^{k+1}\right)\right)$$

$$\leq \sum_{i=1}^{n} \left(h_{i}\left(\mathbf{x}_{i}^{k+1}\right) + g_{i}\left(\mathbf{x}_{i}^{k}\right) + \left\langle\nabla g_{i}\left(\mathbf{x}_{i}^{k}\right), \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\rangle + \frac{L_{i}}{2} \left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2}\right)$$

$$= \sum_{i=1}^{n} \left(h_{i}\left(\mathbf{x}_{i}^{k+1}\right) + g_{i}\left(\mathbf{x}_{i}^{k}\right) + \left\langle\nabla g_{i}\left(\mathbf{x}_{i}^{k}\right), \mathbf{x}_{i} - \mathbf{x}_{i}^{k}\right\rangle + \left\langle\nabla g_{i}\left(\mathbf{x}_{i}^{k}\right), \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}\right\rangle$$

$$+ \frac{L_{i}}{2} \left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2}\right)$$

$$\leq \sum_{i=1}^{n} \left(g_{i}(\mathbf{x}_{i}) + h_{i}(\mathbf{x}_{i}) + \left\langle\mathcal{A}_{i}^{\dagger}(\hat{\lambda}^{k}) + \tau_{i}^{(k)}\left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right), \mathbf{x}_{i} - \mathbf{x}_{i}^{k+1}\right) + \frac{L_{i}}{2} \left\|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\right\|^{2}\right).$$

On the one hand,

$$\begin{split} \sum_{i=1}^{n} \left(f_{i} \left(\mathbf{x}_{i}^{k+1} \right) - f_{i} (\mathbf{x}_{i}) + \left(\mathcal{A}_{i}^{\dagger} (\hat{\lambda}^{k}), \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i} \right) \right) - \left\langle \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} \right) - \mathbf{b}, \hat{\lambda}^{k} - \lambda \right\rangle \end{split} \tag{130} \\ &\leq \sum_{i=1}^{n} \left(-\tau_{i}^{(k)} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}, \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i} \right\rangle + \frac{L_{i}}{2} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right) - \left\langle \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} \right) - \mathbf{b}, \hat{\lambda}^{k} - \lambda \right\rangle \\ &= \sum_{i=1}^{n} \left(-\tau_{i}^{(k)} \left\langle \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}, \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i} \right\rangle + \frac{L_{i}}{2} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right) - \left| \frac{1}{\beta_{k}} \left\langle \lambda^{k+1} - \lambda^{k}, \hat{\lambda}^{k} - \lambda \right\rangle \\ &= \sum_{i=1}^{n} \left[\frac{\tau_{i}^{(k)}}{2} \left(\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right) + \frac{L_{i}}{2} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right] \\ &- \frac{1}{2\beta_{k}} \left(\left\| \lambda^{k+1} - \lambda \right\|^{2} - \left\| \lambda^{k} - \lambda \right\|^{2} + \left\| \hat{\lambda}^{k} - \lambda^{k} \right\|^{2} - \left\| \lambda^{k+1} - \lambda^{k} \right\|^{2} \right) + \frac{L_{i}}{2} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right] \\ &= \sum_{i=1}^{n} \left[\frac{\tau_{i}^{(k)}}{2} \left(\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right) + \frac{L_{i}}{2} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right] \\ &= \sum_{i=1}^{n} \left[\frac{\tau_{i}^{(k)}}{2} \left(\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right) + \frac{L_{i}}{2} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \right] \\ &= \frac{1}{2\beta_{k}} \left(\left\| \lambda^{k+1} - \lambda \right\|^{2} - \left\| \lambda^{k} - \lambda \right\|^{2} + \left\| \hat{\lambda}^{k} - \lambda^{k} \right\|^{2} - \beta_{k}^{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right) \right\|^{2} \right) \\ &\leq \frac{1}{2} \sum_{i=1}^{n} \tau_{i}^{(k)} \left(\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i} \right\|^{2} \right) - \frac{1}{2} \sum_{i=1}^{n} \left(\tau_{i}^{(k)} - L_{i} - n\beta_{k} \left\| \mathcal{A}_{i} \right\|^{2} \right) \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} \\ &+ \frac{1}{2\beta_{k}} \left(\left\| \lambda^{k} - \lambda^{k} \right\|^{2} - \left\| \lambda^{k+1} - \lambda^{k} \right\|^{2} - \left\| \lambda^{k} - \lambda^{k} \right\|^{2} \right) \right\}$$

On the other hand,

$$\sum_{i=1}^{n} \left(f_i\left(\mathbf{x}_i^{k+1}\right) - f_i(\mathbf{x}_i) + \left\langle \mathcal{A}_i^{\dagger}(\lambda), \mathbf{x}_i^{k+1} - \mathbf{x}_i \right\rangle \right) - \left\langle \sum_{i=1}^{n} \mathcal{A}_i(\mathbf{x}_i) - \mathbf{b}, \hat{\lambda}^k - \lambda \right\rangle$$
$$= \sum_{i=1}^{n} \left(f_i\left(\mathbf{x}_i^{k+1}\right) - f_i(\mathbf{x}_i) + \left\langle \mathcal{A}_i^{\dagger}(\hat{\lambda}^k), \mathbf{x}_i^{k+1} - \mathbf{x}_i \right\rangle \right) - \left\langle \sum_{i=1}^{n} \mathcal{A}_i\left(\mathbf{x}_i^{k+1}\right) - \mathbf{b}, \hat{\lambda}^k - \lambda \right\rangle.$$

D Springer

So we have

$$\begin{split} &\sum_{i=1}^{n} \left(f_i \left(\mathbf{x}_i^{k+1} \right) - f_i (\mathbf{x}_i) + \left\langle \mathcal{A}_i^{\dagger}(\lambda), \mathbf{x}_i^{k+1} - \mathbf{x}_i \right\rangle \right) - \left\langle \sum_{i=1}^{n} \mathcal{A}_i (\mathbf{x}_i) - \mathbf{b}, \hat{\lambda}^k - \lambda \right\rangle \\ &\leq \frac{1}{2} \sum_{i=1}^{n} \tau_i^{(k)} \left(\left\| \mathbf{x}_i^k - \mathbf{x}_i \right\|^2 - \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i \right\|^2 \right) \\ &- \frac{1}{2} \sum_{i=1}^{n} \left(\tau_i^{(k)} - L_i - n\beta_k \|\mathcal{A}_i\|^2 \right) \left\| \mathbf{x}_i^{k+1} - \mathbf{x}_i^k \right\|^2 \\ &+ \frac{1}{2\beta_k} \left(\| \lambda^k - \lambda \|^2 - \| \lambda^{k+1} - \lambda \|^2 - \| \hat{\lambda}^k - \lambda^k \|^2 \right). \end{split}$$

Let $\mathbf{x}_i = \mathbf{x}_i^*$ and $\lambda = \lambda^*$, we have

$$\sum_{i=1}^{n} \left(f_{i} \left(\mathbf{x}_{i}^{k+1} \right) - f_{i} \left(\mathbf{x}_{i}^{*} \right) + \left\langle \mathcal{A}_{i}^{\dagger}(\lambda^{*}), \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\rangle \right)$$

$$\leq \frac{1}{2} \sum_{i=1}^{n} \tau_{i}^{(k)} \left[\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} \right] + \frac{1}{2\beta_{k}} \left(\left\| \lambda^{k} - \lambda^{*} \right\|^{2} - \left\| \lambda^{k+1} - \lambda^{*} \right\|^{2} \right)$$

$$- \frac{1}{2} \sum_{i=1}^{n} \left(\tau_{i}^{(k)} - L_{i} - n\beta_{k} \left\| \mathcal{A}_{i} \right\|^{2} \right) \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} - \frac{1}{2\beta_{k}} \left\| \hat{\lambda}^{k} - \lambda^{k} \right\|^{2}.$$

Proof of Theorem 6 As \mathbf{x}^* minimizes $\sum_{i=1}^n f(\mathbf{x}_i) + \langle \lambda^*, \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i) - \mathbf{b} \rangle$, we have

$$0 \leq \sum_{i=1}^{n} \left(f_i\left(\mathbf{x}_i^{k+1}\right) - f_i\left(\mathbf{x}_i^*\right) + \left\langle \mathcal{A}_i^{\dagger}(\lambda^*), \mathbf{x}_i^{k+1} - \mathbf{x}_i^* \right\rangle \right).$$

By Proposition 6, we have

$$\begin{split} &\sum_{i=1}^{n} \frac{1}{2} \left(\tau_{i}^{(k)} - L_{i} - n\beta_{k} \|\mathcal{A}_{i}\|^{2} \right) \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right\|^{2} + \frac{1}{2\beta_{k}} \|\hat{\lambda}^{k} - \lambda^{k}\|^{2} \\ &\leq \frac{1}{2} \sum_{i=1}^{n} \tau_{i}^{(k)} \left(\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} \right) + \frac{1}{2\beta_{k}} \left(\|\lambda^{k} - \lambda^{*}\|^{2} - \|\lambda^{k+1} - \lambda^{*}\|^{2} \right). \end{split}$$

Dividing both sides by β_k and using $\tau_i^{(k)} - L_i - n\beta_k \|\mathcal{A}_i\|^2 \ge \beta_k (\eta_i - n\|\mathcal{A}_i\|^2)$, the nondecrement of β_k and the non-increment of $\beta_k^{-1} \tau_i^{(k)}$, we have

$$\frac{1}{2} \sum_{i=1}^{n} (\eta_{i} - n \|\mathcal{A}_{i}\|^{2}) \|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\|^{2} + \frac{1}{2\beta_{k}^{2}} \|\hat{\lambda}^{k} - \lambda^{k}\|^{2}$$

$$\leq \frac{1}{2} \sum_{i=1}^{n} \left(\beta_{k}^{-1} \tau_{i}^{(k)} \|\mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*}\|^{2} - \beta_{k+1}^{-1} \tau_{i}^{(k+1)} \|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*}\|^{2}\right)$$

$$+ \left(\frac{1}{2\beta_{k}^{2}} \|\lambda^{k} - \lambda^{*}\|^{2} - \frac{1}{2\beta_{k+1}^{2}} \|\lambda^{k+1} - \lambda^{*}\|^{2}\right).$$
(132)

It can be easily seen that $(\mathbf{x}_1^k, \dots, \mathbf{x}_n^k, \lambda^k)$ is bounded, hence has an accumulation point, say $(\mathbf{x}_1^{k_j}, \dots, \mathbf{x}_n^{k_j}, \lambda^{k_j}) \to (\mathbf{x}_1^\infty, \dots, \mathbf{x}_n^\infty, \lambda^\infty)$. Summing (132)–(133) over $k = 0, \dots, \infty$, we have

$$\begin{split} &\frac{1}{2}\sum_{i=1}^{n}\left(\eta_{i}-n\|\mathcal{A}_{i}\|^{2}\right)\sum_{k=0}^{\infty}\left\|\mathbf{x}_{i}^{k+1}-\mathbf{x}_{i}^{k}\right\|^{2}+\sum_{k=0}^{\infty}\frac{1}{2\beta_{k}^{2}}\|\hat{\lambda}^{k}-\lambda^{k}\|^{2}\\ &\leq\frac{1}{2}\sum_{i=1}^{n}\beta_{0}^{-1}\tau_{i}^{(0)}\left\|\mathbf{x}_{i}^{0}-\mathbf{x}_{i}^{*}\right\|^{2}+\frac{1}{2\beta_{0}^{2}}\|\lambda^{0}-\lambda^{*}\|^{2}. \end{split}$$

So $\|\mathbf{x}_i^{k+1} - \mathbf{x}_i^k\| \to 0$ and $\beta_k^{-2} \|\hat{\lambda}^k - \lambda^k\| \to 0$ as $k \to \infty$. Hence $\|\sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i^k) - \mathbf{b}\| \to 0$, which means that $\mathbf{x}_1^{\infty}, \ldots, \mathbf{x}_n^{\infty}$ is a feasible solution.

From (130)–(131), we have

$$\begin{split} &\sum_{i=1}^{n} \left(f_{i} \left(\mathbf{x}_{i}^{k_{j}+1} \right) - f_{i}(\mathbf{x}_{i}) + \left\langle \mathcal{A}_{i}^{\dagger}(\hat{\lambda}^{k_{j}}), \mathbf{x}_{i}^{k_{j}+1} - \mathbf{x}_{i} \right\rangle \right) - \left\langle \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k_{j}+1} \right) - \mathbf{b}, \hat{\lambda}^{k_{j}} - \lambda \right\rangle \\ &\leq \frac{1}{2} \sum_{i=1}^{n} \tau_{i}^{(k_{j})} \left(\left\| \mathbf{x}_{i}^{k_{j}} - \mathbf{x}_{i} \right\|^{2} - \left\| \mathbf{x}_{i}^{k_{j}+1} - \mathbf{x}_{i} \right\|^{2} \right) \\ &- \frac{1}{2} \sum_{i=1}^{n} \left(\tau_{i}^{(k_{j})} - L_{i} - n\beta_{k_{j}} \|\mathcal{A}_{i}\|^{2} \right) \left\| \mathbf{x}_{i}^{k_{j}+1} - \mathbf{x}_{i}^{k_{j}} \right\|^{2} \\ &+ \frac{1}{2\beta_{k_{j}}} \left(\| \lambda^{k_{j}} - \lambda \|^{2} - \| \lambda^{k_{j}+1} - \lambda \|^{2} - \| \hat{\lambda}^{k_{j}} - \lambda^{k_{j}} \|^{2} \right). \end{split}$$

Let $j \to \infty$. By the boundedness of $\tau_i^{(k_j)}$ we have

$$\sum_{i=1}^{n} \left(f_i(\mathbf{x}_i^{\infty}) - f_i(\mathbf{x}_i) + \left(\mathcal{A}_i^{\dagger}(\lambda^{\infty}), \mathbf{x}_i^{\infty} - \mathbf{x}_i \right) \right) \le 0, \quad \forall \mathbf{x}_i.$$

Together with the feasibility of $(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty})$, we can see that $(\mathbf{x}_1^{\infty}, \dots, \mathbf{x}_n^{\infty}, \lambda^{\infty})$ is a KKT point.

By choosing $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*, \boldsymbol{\lambda}^*) = (\mathbf{x}_1^\infty, \dots, \mathbf{x}_n^\infty, \boldsymbol{\lambda}^\infty)$ we have

$$\sum_{i=1}^{n} \eta_i \left\| \mathbf{x}_i^{k_j} - \mathbf{x}_i^{\infty} \right\|^2 + \frac{1}{\beta_{k_j}^2} \left\| \boldsymbol{\lambda}^{k_j} - \boldsymbol{\lambda}^{\infty} \right\|^2 \to 0.$$

Using (132)–(133), we have

$$\sum_{i=1}^{n} \eta_i \left\| \mathbf{x}_i^k - \mathbf{x}_i^{\infty} \right\|^2 + \frac{1}{\beta_k^2} \left\| \boldsymbol{\lambda}^k - \boldsymbol{\lambda}^{\infty} \right\|^2 \to 0.$$

So $(\mathbf{x}_1^k, \ldots, \mathbf{x}_n^k, \boldsymbol{\lambda}^k) \to (\mathbf{x}_1^\infty, \ldots, \mathbf{x}_n^\infty, \boldsymbol{\lambda}^\infty).$

🖄 Springer

Appendix 7: Proof of Theorem 7

Proof of Theorem 7 By the definition of α and $\tau_i^{(k)}$,

$$\frac{1}{2} \left[\sum_{i=1}^{n} \left(\tau_{i}^{(k)} - L_{i} - n\beta_{k} \|\mathcal{A}_{i}\|^{2} \right) \|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\|^{2} + \frac{1}{\beta_{k}} \|\hat{\lambda}^{k} - \lambda^{k}\|^{2} \right]$$

$$\geq \frac{\beta_{k}}{2} \left[\sum_{i=1}^{n} \left(\eta_{i} - n \|\mathcal{A}_{i}\|^{2} \right) \|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\|^{2} + \frac{1}{\beta_{k}^{2}} \|\hat{\lambda}^{k} - \lambda^{k}\|^{2} \right]$$

$$\geq \frac{\alpha\beta_{k}}{2} (n+1) \left(\sum_{i=1}^{n} \|\mathcal{A}_{i}\|^{2} \|\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k}\|^{2} + \frac{1}{\beta_{k}^{2}} \|\hat{\lambda}^{k} - \lambda^{k}\|^{2} \right)$$

$$\geq \frac{\alpha\beta_{k}}{2} (n+1) \left(\sum_{i=1}^{n} \|\mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right) \|^{2} + \frac{1}{\beta_{k}^{2}} \|\hat{\lambda}^{k} - \lambda^{k}\|^{2} \right)$$

$$= \frac{\alpha\beta_{k}}{2} (n+1) \left(\sum_{i=1}^{n} \|\mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{k} \right) \|^{2} + \|\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k} \right) - \mathbf{b} \|^{2} \right)$$

$$\geq \frac{\alpha\beta_{k}}{2} \|\sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} \right) - \mathbf{b} \|^{2}.$$
(134)

So by (127)–(129) and the non-decrement of β_k , we have

$$\sum_{i=1}^{n} \left(f_{i}\left(\mathbf{x}_{i}^{k+1}\right) - f_{i}\left(\mathbf{x}_{i}^{*}\right) + \left\langle \mathcal{A}_{i}^{\dagger}(\lambda^{*}), \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\rangle \right) + \frac{\alpha\beta_{0}}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i}\left(\mathbf{x}_{i}^{k+1}\right) - \mathbf{b} \right\|^{2}$$
(136)
$$\leq \sum_{i=1}^{n} \left(f_{i}\left(\mathbf{x}_{i}^{k+1}\right) - f_{i}\left(\mathbf{x}_{i}^{*}\right) + \left\langle \mathcal{A}_{i}^{\dagger}(\lambda^{*}), \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\rangle \right) + \frac{\alpha\beta_{0}}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i}\left(\mathbf{x}_{i}^{k+1}\right) - \mathbf{b} \right\|^{2}$$
(137)
$$\leq \frac{1}{2} \sum_{i=1}^{n} \tau_{i}^{(k)} \left(\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} \right) + \frac{1}{2\beta_{k}} \left(\left\| \lambda^{k} - \lambda^{*} \right\|^{2} - \left\| \lambda^{k+1} - \lambda^{*} \right\|^{2} \right).$$
(138)

Dividing both sides by β_k and using the non-decrement of β_k and the non-increment of $\beta_k^{-1} \tau_i^{(k)}$, we have

$$\frac{1}{\beta_{k}} \left[\sum_{i=1}^{n} \left(f_{i} \left(\mathbf{x}_{i}^{k+1} \right) - f_{i} \left(\mathbf{x}_{i}^{*} \right) + \left\langle \mathcal{A}_{i}^{\dagger}(\lambda^{*}), \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\rangle \right) + \frac{\alpha\beta_{0}}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_{i} \left(\mathbf{x}_{i}^{k+1} \right) - \mathbf{b} \right\|^{2} \right]$$
(139)
$$\leq \frac{1}{2} \sum_{i=1}^{n} \beta_{k}^{-1} \tau_{i}^{(k)} \left(\left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*} \right\|^{2} - \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} \right) + \frac{1}{2\beta_{k}^{2}} \left(\left\| \lambda^{k} - \lambda^{*} \right\|^{2} - \left\| \lambda^{k+1} - \lambda^{*} \right\|^{2} \right)$$

$$\leq \frac{1}{2} \sum_{i=1}^{n} \left(\beta_{k}^{-1} \tau_{i}^{(k)} \left\| \mathbf{x}_{i}^{k} - \mathbf{x}_{i}^{*} \right\|^{2} - \beta_{k+1}^{-1} \tau_{i}^{(k+1)} \left\| \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\|^{2} \right)$$

D Springer

$$+\left(\frac{1}{2\beta_{k}^{2}}\left\|\lambda^{k}-\lambda^{*}\right\|^{2}-\frac{1}{2\beta_{k+1}^{2}}\left\|\lambda^{k+1}-\lambda^{*}\right\|^{2}\right).$$
(140)

Summing over k = 0, ..., K and dividing both sides by $\sum_{k=0}^{K} \beta_k^{-1}$, we have

$$\sum_{i=1}^{n} \left(\sum_{k=0}^{K} \gamma^{k} f_{i} \left(\mathbf{x}_{i}^{k+1} \right) - f_{i} \left(\mathbf{x}_{i}^{*} \right) + \left\langle \mathcal{A}_{i}^{\dagger}(\lambda^{*}), \sum_{k=0}^{K} \gamma^{k} \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\rangle \right)$$
(141)

$$+\frac{\alpha\beta_0}{2}\sum_{k=0}^{K}\gamma^k \left\|\sum_{i=1}^{n}\mathcal{A}_i\left(\mathbf{x}_i^{k+1}\right) - \mathbf{b}\right\|^2$$
(142)

$$\leq \left(\sum_{i=1}^{n} \beta_{0}^{-1} \tau_{i}^{(0)} \|\mathbf{x}_{i}^{0} - \mathbf{x}_{i}^{*}\|^{2} + \beta_{0}^{-2} \|\lambda^{0} - \lambda^{*}\|^{2}\right) / \sum_{k=0}^{K} 2\beta_{k}^{-1}.$$
 (143)

Using the convexity of f_i and $\|\cdot\|^2$, we have

$$\sum_{i=1}^{n} \left(f_i \left(\bar{\mathbf{x}}_i^K \right) - f_i \left(\mathbf{x}_i^* \right) + \left\langle \mathcal{A}_i^{\dagger}(\lambda^*), \bar{\mathbf{x}}_i^K - \mathbf{x}_i^* \right\rangle \right) + \frac{\alpha \beta_0}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_i \left(\bar{\mathbf{x}}_i^K \right) - \mathbf{b} \right\|^2$$
(144)

$$\leq \sum_{i=1}^{n} \left(\sum_{k=0}^{K} \gamma^{k} f_{i} \left(\mathbf{x}_{i}^{k+1} \right) - f_{i} \left(\mathbf{x}_{i}^{*} \right) + \left\langle \mathcal{A}_{i}^{\dagger}(\lambda^{*}), \sum_{k=0}^{K} \gamma^{k} \mathbf{x}_{i}^{k+1} - \mathbf{x}_{i}^{*} \right\rangle \right)$$
(145)

$$+ \frac{\alpha\beta_0}{2} \sum_{k=0}^{K} \gamma^k \left\| \sum_{i=1}^{n} \mathcal{A}_i\left(\mathbf{x}_i^{k+1}\right) - \mathbf{b} \right\|^2.$$
(146)

So we have

$$\sum_{i=1}^{n} \left(f_i\left(\bar{\mathbf{x}}_i^K\right) - f_i\left(\mathbf{x}_i^*\right) + \left\langle \mathcal{A}_i^{\dagger}(\lambda^*), \bar{\mathbf{x}}_i^K - \mathbf{x}_i^* \right\rangle \right) + \frac{\alpha\beta_0}{2} \left\| \sum_{i=1}^{n} \mathcal{A}_i\left(\bar{\mathbf{x}}_i^K\right) - \mathbf{b} \right\|^2$$
(147)

$$\leq \left(\sum_{i=1}^{n} \beta_{0}^{-1} \tau_{i}^{(0)} \|\mathbf{x}_{i}^{0} - \mathbf{x}_{i}^{*}\|^{2} + \beta_{0}^{-2} \|\lambda^{0} - \lambda^{*}\|^{2}\right) / \sum_{k=0}^{K} 2\beta_{k}^{-1}.$$
(148)

- Beck, A., & Teboulle, M. (2009). A fast iterative shrinkage-thresholding algorithm for linear inverse problems. SIAM Journal of Imaging Sciences, 2(1), 183–202.
- Boyd, S., Parikh, N., Chu, E., Peleato, B., & Eckstein, J. (2011) Distributed optimization and statistical learning via the alternating direction method of multipliers. In M. Jordan (Ed.), *Foundations and trends in machine learning*. Hanover, MA: Now Publishers Inc.
- Boyd, S., & Vandenberghe, L. (2004). Convex optimization. Cambridge: Cambridge University Press.
- Cai, J., Candès, E., & Shen, Z. (2010). A singular value thresholding algorithm for matrix completion. SIAM Journal of Optimization, 20(4), 1956–1982.
- Candès, E., & Recht, B. (2009). Exact matrix completion via convex optimization. Foundations of Computational Mathematics, 9(6), 717–772.
- Candès, E., Li, X., Ma, Y., & Wright, J. (2011). Robust principal component analysis? JACM, 58(3), 11.
- Chandrasekaran, V., Parrilo, P., & Willsky, A. (2012). Latent variable graphical model selection via convex optimization. *The Annals of Statistics*, 40(4), 1935–1967.

- Chang, E., Zhu, K., Wang, H., Bai, H., Li, J., Qiu, Z., & Cui, H. (2007). Psvm: Parallelizing support vector machines on distributed computers. In NIPS.
- Chang, E. (2011). Foundations of large-scale multimedia information management and retrieval: Mathematics of perception. Berlin: Springer.
- Chen, C., He, B., Ye, Y., & Yuan, X. (2013). The direct extension of ADMM for multi-block convex minimization problems is not necessarily convergent. Preprint.
- Deng, W., & Yin, W. (2012). On the global and linear convergence of the generalized alternating direction method of multipliers. DTIC Document: Tech. rep.
- Deng, W., Yin, W., & Zhang, Y. (2011). Group sparse optimization by alternating direction method. TR11-06, Department of Computational and Applied Mathematics, Rice University.
- Esser, E. (2009). Applications of Lagrangian-based alternating direction methods and connections to split Bregman. CAM Report 09-31, UCLA.
- Favaro, P., Vidal, R., Ravichandran, A. (2011). A closed form solution to robust subspace estimation and clustering. In CVPR.
- Fazel, M. (2002). Matrix rank minimization with applications. Ph.D. thesis.
- Fortin, M., & Glowinski, R. (1983). Augmented Lagrangian methods. Amsterdam: North-Holland.
- Goldfarb, D., & Ma, S. (2012). Fast multiple splitting algorithms for convex optimization. SIAM Journal of Optimization, 22(2), 533–556.
- Goldstein, T., & Osher, S. (2008). The split Bregman method for l₁ regularized problems. SIAM Journal of Imaging Sciences, 2(2), 323–343.
- He, B., & Yuan, X. (2012). On the O(1/n) convergence rate of the Douglas–Rachford alternating direction method. SIAM Journal of Numerical Analysus, 50(2), 700–709.
- He, B., Tao, M., & Yuan, X. (2012). Alternating direction method with Gaussian back substitution for separable convex programming. SIAM Journal of Optimization, 22(2), 313–340.
- He, B., & Yuan, X. (2013). Linearized alternating direction method with Gaussian back substitution for separable convex programming. *Numerical Algebra, Control and Optimization*, 3(2), 247–260.
- Hong, M., Luo, Z. Q. (2012). On the linear convergence of the alternating direction method of multipliers. Preprint, arXiv:12083922.
- Jacob, L., Obozinski, G., & Vert, J. (2009). Group Lasso with overlap and graph Lasso. In ICML.
- Ji, H., Liu, C., Shen, Z., & Xu, Y. (2010). Robust video denoising using low rank matrix completion. In CVPR.
- Lin, Z., Chen, M., & Ma, Y. (2009a). The augmented Lagrange multiplier method for exact recovery of corrupted low-rank matrices. UIUC Technical Report UILU-ENG-09-2215.
- Lin, Z., Ganesh, A., Wright, J., Wu, L., Chen, M., & Ma, Y. (2009b). Fast convex optimization algorithms for exact recovery of a corrupted low-rank matrix. UIUC Technical Report UILU-ENG-09-2214.
- Lin, Z., Liu, R., & Su, Z. (2011). Linearized alternating direction method with adaptive penalty for low-rank representation. In NIPS.
- Liu, G., & Yan, S. (2011). Latent low-rank representation for subspace segmentation and feature extraction. In ICCV.
- Liu, R., Lin, Z., & Su, Z. (2013). Linearized alternating direction method with parallel splitting and adaptive penalty for separable convex programs in machine learning. In ACML.
- Liu, G., Lin, Z., & Yu, Y. (2010). Robust subspace segmentation by low-rank representation. In ICML.
- Liu, G., Lin, Z., Yan, S., Sun, J., Yu, Y., & Ma, Y. (2012). Robust recovery of subspace structures by low-rank representation. *IEEE Transactions on PAMI*, 35(1), 171–184.
- Ma, S., Goldfarb, D., & Chen, L. (2011). Fixed point and bregman iterative methods for matrix rank minimization. *Mathematical Programming*, 128(1–2), 321–359.
- Meier, L., Geer, S. V. D., & Bühlmann, P. (2008). The group Lasso for logistic regression. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 70(1), 53–71.
- Rockafellar, R. (1970). Convex analysis. Princeton: Princeton University Press.
- Shen, X., & Wu, Y. (2012). A unified approach to salient object detection via low rank matrix recovery. In CVPR.
- Subramanian, A., Tamayo, P., Mootha, V., Mukherjee, S., et al. (2005). Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences*, 102(43), 267–288.
- Tao, M. (2014). Some parallel splitting methods for separable convex programming with O(1/t) convergence rate. *Pacific Journal of Optimization*, 10(2), 359–384.
- Toh, K., & Yun, S. (2010). An accelerated proximal gradient algorithm for nuclear norm regularized least squares problems. *Pacific Journal of Optimization*, 6(15), 615–640.
- Tron, R., & Vidal, R. (2007). A benchmark for the comparison of 3D montion segmentation algorithms. In CVPR.

- van de Vijver, M., He, Y., van't Veer, L., Dai, H., et al. (2002). A gene-expression signature as a predictor of survival in breast cancer. *The New England Journal of Medicine*, 347(25), 1999–2009.
- Wright, J., Yang, A., Ganesh, A., Sastry, S., & Ma, Y. (2009). Robust face recognition via sparse representation. IEEE Transactions on PAMI, 31(2), 210–227.
- Xu, Y., Yin, W., & Wen, Z. (2011). An alternating direction algorithm for matrix completion with nonnegative factors. CAAM Technical Report TR11-03.
- Yang, J., & Yuan, X. (2013). Linearized augmented Lagrangian and alternating direction methods for nuclear norm minimization. *Mathematics of Computation*, 82(281), 301–329.
- Ye, J., Ji, S., & Chen, J. (2008). Multi-class discriminant kernel learning via convex programming. JMLR, 9, 719–758.
- Zhang, X., Burger, M., & Osher, S. (2011). A unified primal-dual algorithm framework based on Bregman iteration. *Journal of Scientific Computing*, 46(1), 20–46.