

2003

Linking Landscape And Water Quality In The Mississippi River Basin For 200 Years

R. Eugene Turner
eturne@lsu.edu

Nancy N. Rabalais
nrabalais@lumcon.edu

Follow this and additional works at: https://digitalcommons.lsu.edu/oceanography_coastal_pubs

Recommended Citation

Turner, R., & Rabalais, N. N. (2003). Linking Landscape And Water Quality In The Mississippi River Basin For 200 Years. *Bioscience*, 53 (6), 563-572. [https://doi.org/10.1641/0006-3568\(2003\)053\[0563:LLAWQI\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2003)053[0563:LLAWQI]2.0.CO;2)

This Article is brought to you for free and open access by the Department of Oceanography & Coastal Sciences at LSU Digital Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of LSU Digital Commons. For more information, please contact ir@lsu.edu.

Linking Landscape and Water Quality in the Mississippi River Basin for 200 Years

R. EUGENE TURNER AND NANCY N. RABALAIS

Two centuries of land use in the Mississippi River watershed are reflected in the water quality of its streams and in the continental shelf ecosystem receiving its discharge. The most recent influence on nutrient loading—intense and widespread farming and especially fertilizer use—has had a more significant effect on water quality than has land drainage or the conversion of native vegetation to cropland and grazing pastures. The 200-year record of nutrient loading to offshore water is reflected in the paleoreconstructed record of plankton in dated sediments. This record illustrates that the development of fair, sustained management of inland ecosystems is linked to the management of offshore systems. Land use in this fully occupied watershed is under the strong influence of national policies affecting all aspects of the human ecosphere. These policies can be modified for better or worse, but water quality will probably change only gradually because of the strong buffering capacity of the soil ecosystem.

Keywords: Mississippi River, water quality, agriculture, sustainability, environmental history

The local linkages between land use and water quality have cumulative effects within a region, its watershed, and the receiving coastal waters. The effects of these linkages vary as the cultural and ecological landscape varies with population growth, changes in land use, and climatic events. These changes have been particularly evident in North America over the past four centuries, as European culture was inserted into the North American continent, as the Native American Indians died from newly introduced diseases and were subdued through military and political means, and as the mostly European population grew and became urbanized. A high-intensity agricultural–economic system has turned the American Midwest into what is now known as the nation’s “breadbasket,” where 65% of the land in the 14 states of the Mississippi River Basin (MRB; figure 1) is farmland and 25% is harvestable cropland.

As the landscape changed in response to these developments, so too did the Mississippi River. Its main channel has been shortened and dredged, its banks stabilized for navigation, and flood protection levees built, which extend continuously south of Vicksburg, Mississippi. These levees have isolated the alluvial soils from the main channel, promoting agriculture development on alluvial soils (Abernethy and Turner 1987). The volume of river water extracted and returned by industrial plants in Louisiana is now greater than that discharged into the Gulf of Mexico (EARI 1975).

Today’s significant water quality problems in the MRB are related to these landscape and industrial developments. Forty-four percent of the surveyed rivers in 15 MRB states were “impaired” in 2000 (EPA 2002). Nitrate concentration in the Des Moines River is sometimes greater than 10 milligrams per liter, which is the statutory maximum limit for drinkable water supplies (Hallberg 1987), and there were 1557 fish consumption advisories in the MRB in September 1996 alone (EPA 1998). Water-quality changes in the lower Mississippi River in this century can be linked in a plausible cause-and-effect relationship to the formation of a 20,000-square-kilometer (km²) low-oxygen zone in coastal waters (Rabalais et al. 2002). The food web in this zone is poised to change from a productive food web of diatoms, zooplankton, and fish—which supports 25% of US fish landings—to one with diminished pelagic and demersal fisheries (Turner et al. 1998).

In this article we discuss two centuries of landscape changes that affect water quality in the MRB and in continental shelf

R. Eugene Turner (e-mail: eturne@lsu.edu) is a professor in the Department of Oceanography and Coastal Sciences and the Coastal Ecology Institute at Louisiana State University, Baton Rouge, LA 70803. Nancy N. Rabalais (e-mail: nrabalais@lumcon.edu) is a professor at the Louisiana Universities Marine Consortium in Chauvin, LA 70344. © 2003 American Institute of Biological Sciences.




Figure 1. The Mississippi River Basin and the major watersheds discussed in this article. Key: 1, upper Mississippi River; 2, Ohio River; 3, Missouri River; 4, Arkansas River; 5, lower Mississippi River; 6, Red River. The total combined area is 41% of the contiguous states.

ecosystems. First, we reconstruct the timing of population growth and soil disturbance resulting from deforestation, cultivation, and drainage. Next, we draw connections between the subsequently higher sediment yields and the loss of nutrients from the newly disturbed landscape. To support the conclusion that the release of nutrients stored in the pre-European period was large, we cite experiments on small watersheds in which variables can be controlled, as well as anecdotal accounts from before the 1930s. Various analyses of the variability in water quality among watersheds support the conclusion that land use and population density are directly related to higher nitrogen yields and that these yields change important nutrient ratios. A paleoreconstruction of continental shelf sediments confirms the significance of these water quality changes to coastal food webs and suggests that the most recent influence on nutrient loading—intense and widespread farming—has affected water quality more than all previous landscape changes have.

Soil disturbance leading to soil loss

The sediment yield of a watershed is, in general, inversely related to the amount of vegetation present and to the accumulation of soil organic matter. The vegetation cover in the MRB was reduced as populations expanded during the 1800s. The population density of Native American Indians in the early 1600s was relatively low (no more than 106,000, or less than 0.1 person per km²; Ubelaker 1992), in part because of diseases introduced by the Europeans. Population growth in the midwestern states began earlier than in the Missouri

basin, reaching 1 to 10 people per km² by the 1850s (figure 2a), when the population center of the United States crossed the Appalachian Mountains and headed into the MRB in a west–southwest trajectory.

The area of land brought under cultivation rose with population growth (figure 2b). Cultivation was preceded by tree cutting, often by girdling the trunk, or burning. Trees were not routinely sold, so we inferred changes in forest area from the records of land use. Forest loss during the 1800s was quite rapid. The area of forest in Ohio, for example, went from 54% in 1853 to 18% in 1883 (Leue 1886). Greeley (1925) documented that the virgin forests of 1850 in the United States were largely remnants by 1920. Humphreys and Abbot (1876) estimated that 15% of the MRB was under cultivation or had “improvements” by 1860.

Did this agricultural expansion result in a significant increase in the sediment yield of the MRB? To find out, we assembled the suspended sediment records collected at the Carrollton water treatment plant in New Orleans (Louisiana) since 1903, when the existing unified treatment system was under construction. Additional data were reported by Quinn (1894), who sampled the Mississippi River for suspended sediments from 1879 to 1899. Data are also reported in Humphreys and Abbot (1876) for 96 weekly measurements from February 1851 to February 1853, for 7 additional measurements later in 1853, and for 36 data points for May through August 1846. These data (figure 2c) indicate that the suspended sediment concentration at New Orleans was highest in the late 1800s, when the area of new agricultural land

brought into production each year was at its peak. Sediment concentration declined after 1910, especially after the period of extensive dam construction that began in the early 1950s on the Missouri River. Meade and colleagues (1990) documented that these dams trapped large amounts of sediments and dramatically altered the transport patterns of suspended sediments downstream in the basin all the way to New Orleans. Improved soil conservation practices may also have contributed to this decline (Trimble 1999).

Was there a long-term climate change in river discharge that caused these long-term variations in sediment yield? If discharge varies, then both the nutrient and sediment loads from a watershed are affected (load = concentration \times discharge). Thus it is important to document the variability in water supply for the basin as a whole. A portion of the Mississippi River is diverted westward through control structures to join with the Red River to form the Atchafalaya River. This diversion, which was controlled in earnest after the record-setting 1927 flood, now constitutes about 30% of the main stem flow. The longest-kept discharge records are from upstream of this diversion at Vicksburg, Mississippi, the location of the US Army Corps of Engineers Waterways Experiment Station. These discharge records started in 1817, although some of the earlier records, which were based on empirical stage–discharge relationships calibrated without the benefit of today's technical conveniences, may be suspect. The discharge volume since 1817 varies around a long-term annual mean of 17,000 cubic meters per second (figure 3). The 170-year record of river discharge demonstrates no permanent rise or fall in the annual discharge volume. The influences of variable climate changes, therefore, are not considered an important cause of variations in sediment yields on the scale of decades.

The suspended sediment yield (mass per unit area) from the landscape certainly increased after European colonization of what was once an American Indian aboriginal province of hunter–gatherers with sparse and casual crop cultivation (Cronon 1983). The new colonists practiced row farming at an intensity previously unknown in these lands, and the switch to higher-intensity farming happened quickly. Row crop cultivation meant that existing vegetation was removed and the soil surface severely disrupted and compacted. Livestock roamed the woods for forage and further disrupted the soil surface. A visitor in the 1800s noted, “There is no portion of the globe that is being exhausted of its fertility by injudicious cultivation, so rapidly as the Mississippi Valley” (Bateham 1849, quoted in Whitney 1994, p. 226).

The effects of land use on soil erosion can be inferred from well-documented examples inside and outside the MRB. For example, Wolman (1967) discussed the changes in sediment yield from 1800 to 1960 for a




Figure 2. A summary interpretation of the relationships among population growth, land conversion to agriculture, and fertilizer use in the Mississippi River Basin (MRB) and coastal diatom production. (a) Population density in two regional groupings of states. The Midwest group consists of Ohio, Indiana, Illinois, Iowa, Michigan, Wisconsin, and Minnesota. The Missouri group consists of Montana, Wyoming, North Dakota, South Dakota, Nebraska, and Missouri. Population data are from the US Census Bureau and exclude Native Americans and slaves in most cases before 1850 (Anonymous 1992). (b) The new area of harvested cropland and land under drainage added each year in the MRB. Cropland data for before 1860 are from Humphreys and Abbot (1876), and subsequent census estimates are from the US Department of Agriculture (USDA). The rise in harvested cropland after a decade of decline is probably taking place on previously farmed land. The new land under drainage is from USDA census estimates (irregular intervals) (USCB 1961, 1973). (c) The annual average suspended sediment concentration (milligrams per liter) at New Orleans, Louisiana. Data are from the annual reports of the New Orleans Water and Sewerage Board, Quinn (1894), and Humphreys and Abbot (1876). (d) The percentage of biogenic silica (BSi) in sediments from dated sediment cores collected near the mouth of the Mississippi River (Turner and Rabalais 1994). Percentage BSi is an indicator of biogenic silica found in diatom remains (dry weight basis). Also included is the annual flux of nitrogen in the Mississippi River from 1920 to 1987.




Figure 3. Mississippi River discharge at Vicksburg, Mississippi, for individual years and a 10-year moving average.

Maryland Piedmont stream: Erosion increased when the forest cover was removed for crops, declined when farms were abandoned, rose again with urban growth in the 1960s, and declined again as concrete and roads held soil in place while construction activities dropped off. In the MRB, Brune (1948) documented a 50-fold increase in sediment yield as cropland area increased (the amount of sediment was also affected by the drainage area). Brune also showed similar relations of sediment yield to drainage area and cropland for the Ohio River Basin. Meade and colleagues (1990) noted that conversion of forest to agricultural fields “caused orders-of-magnitude increases in soil erosion and corresponding increases in the sediment yields of rivers.” In a letter to Jared Eliot in the mid-18th century, the naturalist John Bartram wrote,

About 20 years past when the woods was not pastured and full of high weeds and the ground light(,) then the rain sunk much more into the earth and did not wash and tear up the surface (as now). The rivers and brooks in floods would be black with mud but now the rain runs most of it off on the surface (and) is collected [sic] into the hollows which it wears to the sand and clay which it bears away with the swift current down to brooks and rivers whose banks it overflows. (Cronon 1983, p. 147)

Trimble’s provocative review of the history of soil management in the United States begins with these conclusions:

Much of the American soil has been poorly treated since European settlement. Early travelers’ accounts and a few systematic studies make this clear. Many early settlers brought poor agricultural practices with them: the Scots, for example, were known for their crude agriculture as late as 1780. On this continent, the cheap and unlimited land promoted a widespread attitude that land could be used, exhausted, or destroyed as the case may be, and then abandoned for new land. Such a system existed well into the twentieth century, at least in much of the southeast, and the depredations on the landscape are still visible today. (Trimble 1985, p. 162)

These effects of agricultural expansion on soil erosion (figure 4) were widely acknowledged by the 1930s as having

major consequences for farm management. The amounts of soil loss under cultivation could be staggering and quick. A 1928 US Department of Agriculture publication on soil erosion described how “certain piedmont areas whose records are known have, within a period of 30 years, lost their topsoil entirely, 10 inches or more of loam and clay loam having washed off down to the clay subsoil” (Bennett and Chapline 1928). The authors of that report described how an apple orchard near Lookout Mountain in northeastern Kansas had the trunks of trees completely buried by overwash of silt from neighboring lands, so that the level of the ground was at the branches, and they described gullies that were 300 feet wide in places and three-quarters of a mile long (Bennett and Chapline 1928). They concluded that “it seems scarcely necessary to state the perfectly obvious fact that a very large part of this impoverishment and wastage has taken place since the clearing of the forests, the breaking of the prairie sod, and the overgrazing of pasture lands. A little is being done here and there to check this loss—an infinitesimal part of what should be done” (p. 23).

A 1935 Iowa State Planning Commission document (cited in Prince 1997) noted that disturbance of the state’s prairie had caused the loss of 192,643 metric tons per km² of soil and that 40% of Iowa had lost 50% to 75% of its surface soil. Data for farmland at an Oklahoma agriculture experiment station showed that the water runoff from land cultivated continuously in cotton was 11 times greater, and the soil losses 760 times greater, than from the same kind of land covered with ungrazed Bermuda grass (6-year average, 1930–1935; USDA 1938). Soil losses from 1894 to 1935 for land continuously planted with corn at the Ohio Agricultural Experiment Station were 63% of the organic matter and 4.05 centimeters (cm) of soil; the losses for land planted with oats over the same period were 36% of organic matter and 2.64 cm of soil (USDA 1938). Such high sediment losses are not unusual. Erosion rates of 100, or even 1000, times higher after land clearing are common (Novotny 1999). Within the MRB, the sediment yields are highest with cultivation of row crops and lowest with dense plant cover. For example, the suspended sediment in runoff from modern-day forests (which are not mature forests) is one-third of that from land planted with corn and soybeans (table 1).

As a result of these massive changes, sediments, including soil organic matter, were washed off the land and into small creeks, rivers, and, at least to some extent, the coastal zone. Recognition of the seriousness of soil erosion led to the formation of the Soil Erosion Service (renamed the Soil Conservation Service in 1935, now the Natural Resources Conservation Service) and to the classic 1934 Reconnaissance Erosion Survey.

Although preimmigration suspended sediment yields in the MRB were not recorded, there are proxy measures. These proxy measures are in sediment cores from the Mississippi River Delta that have been dated and analyzed for indicators

Table 1. Suspended sediment, total phosphorus, and nitrate yields in runoff by dominant land use in the United States for 1980–1989.

Land use	Runoff (kilograms per square kilometer per year)		
	Suspended sediment	Total phosphorus	Nitrate
Wheat	3503	3.5	11.2
Urban	8056	41.7	192
Forest	10,858	22.1	89.3
Rangeland	11,559	6.0	10.9
Mixed crops	27,671	23.1	107
Corn and soybeans	35,026	57.1	326

Source: Smith et al. (1996).

of phytoplankton production (Turner and Rabalais 1994). Diatoms, in particular, leave a record of their abundance in the siliceous frustules deposited on the bottom. Quantifying these frustules in sediments yields a surrogate or proxy measurement of algal production in the surface waters. The accumulation of this biologically bound silica (BSi) showed a distinctive rise at the beginning of the 1800s (figure 2d). It later declined to a low around the 1900s, underwent another rise

and fall, and then rose again over the last 30 to 40 years. The diatom production rate is thought to be limited by the supply of nitrogen, much as an agricultural crop is nutrient limited, but in a different physical setting. The annual BSi accumulation has risen in proportion to the documented loading of nitrogen in the river from 1950 to 1990 (Turner and Rabalais 1994). Is the source of the nitrogen that caused the rise in BSi offshore in the 1800s and early 1900s related to an increase in nutrient yields accompanying soil erosion from the MRB?

Nutrient release from soils

When soils are disturbed enough during cultivation, the ecological processes that keep nutrients bound up in the soil and organic matter are subdued, and the stored nitrogen is released until the soil is, as an agriculturist might say, “exhausted”—meaning that the natural fertility of the soil is diminished to the point that crop growth is compromised. This is the MRB equivalent of the slash-and-burn agriculture found in the tropics, a farming practice in which crops are planted amid a shifting mosaic of soils that are newly exploited, in decline, or abandoned to natural rehabilitation. The US presidents Washington and Jefferson wrote about soil exhaustion. Later, so did naturalist John Muir:




Figure 4. Photographs of the effects of soil erosion, 1910 to 1939. Top left: erosion in Chilton County, Alabama, circa 1935; cotton was grown on this field in 1910. Top right: line fence crossing a tributary in Winon County, Minnesota, in 1939. The posts and wire show the form of the stream and floodplain at the time the fence was rebuilt 5 years earlier; the photograph shows the channel trenching and widening since then. The man in the photograph is 6 feet tall. Bottom left: Two mules on a hillside plowing corn, circa 1935. Bottom right: buried machinery in a barn lot in Dallas, South Dakota, in 1936. Photographs: National Archives (NLR-PHOCO-53227 [600], WR 10 26, RG 083 G 36711, and 114 SC 5089).

At first, wheat, corn, and potatoes were the principal crops we raised; wheat especially. But in four or five years the soil was so exhausted that only five or six bushels an acre, even in the better fields, were obtained, although when first plowed twenty and twenty-five were the ordinary yield. More attention was then paid to corn, but without fertilizers the corn crop also became very meagre. At last it was discovered that English clover would grow on even the exhausted fields, and that when plowed under and planted with corn, or even wheat, wonderful crops were raised. (Muir 1965, p. 164)

Gray's (1933) review of southern agriculture up to 1860 characterized soil exhaustion as an expected consequence of a way of farming: "Planters bought land as they might buy a wagon—with the expectation of wearing it out...[as the] wave of migration passed like a devastating scrooge [sic]. Especially in the rolling piedmont lands the planting of corn and cotton in hill and drill hastened erosion, leaving the hill-sides gullied and bare" (p. 446).

Soil nitrogen is one of the exhausted nutrients, and the nitrate ion is particularly mobile. When a forest was clear-cut in Gale River, New Hampshire, the nitrogen losses went from an annual *retention* of 200 kilograms (kg) nitrogen as nitrate (nitrate-N) per km² as an undisturbed forest to a loss of 3800 and 5700 kg nitrate-N for the first and second years, respectively, after clear-cutting (Pierce et al. 1972). These loss rates are 10 to 20 times higher than the present total nitrogen (TN) yields for the whole MRB (489 kg TN per km² per year; Goolsby et al. 1999). To put these loss rates in perspective, consider that the present carbon, nitrogen, and phosphorus yields from the MRB are less than 0.1% of the carbon, nitrogen, and phosphorus in a layer of soil 1 meter thick with a 2% organic content. A small change in the element inventory on land can thus significantly change water quality in stream channels. These disturbances can have long-term consequences, too. Aber and colleagues (1998) studied the nitrogen dynamics of a northeastern forest and came to the conclusion that land-use history going back as far as 200 years had a stronger influence on nitrate losses than did modern-day nitrogen inventories or depositions.

The application of technological inventions in the 1800s introduced changes to farming practices. The wide-scale use of the iron plow, threshing machine, mower, and reaper began between 1825 and 1850. The first patent for a chemical fertilizer was

issued in 1849 to the Chappell brothers in Baltimore, and subsequent phosphate fertilizer production centered around Charleston, South Carolina, in the late 1800s. The first agricultural journals appeared—*The American Farmer* (Baltimore, 1819), *The Plow Boy* (Albany, New York, 1819), *The New England Farmer* (Boston, 1822), *The New York Farmer* (New York, 1827), and *The Southern Agriculturist* (Charleston, 1828)—as land management practices evolved toward less land clearing and more intensive use. The newer approaches to soil management also led to high nutrient losses relative to the losses from continuously vegetated land. A 2-year study from one of the new agricultural experiment stations in the early 1900s documented that the conversion of the bluegrass native prairie vegetation to continuous wheat or corn, or to plowed fields, resulted in the loss of up to 160 times more nitrogen and 350 times more phosphorus than lost from the native landscape (figure 5).

Ditching and tile drainage further stimulated nitrogen losses from soils. Soil drainage is intended to dry out soils by promoting belowground drainage. The majority of dissolved nitrogen lost in a drained field is in the form of nitrate, and the nitrate is lost by movement as shallow, subsurface flow or as deeper groundwater, not as overland flow (Lowrance 1992).




Figure 5. Average nitrogen (N) and phosphorus (P) losses (kilograms per square kilometer per year), 1 May 1926 to 1 May 1928, in runoff from experimental plots at the Missouri Agriculture Experiment Station (Miller and Krusekopf 1932). A belowground vertical profile of the roots in wheat (*Triticum aestivum*) and native big bluestem (*Andropogon gerardi*) prairie vegetation is shown.

Several studies have documented a strong and proportional relationship between nitrogen applications to surface soils and the nitrogen content in soil pore water and groundwater (Baker and Laflen 1983, Keeney 1986, Hallberg 1989). Nitrate accumulation in soil water of experimental plots in Virginia and Nebraska, for example, increased as nitrogen fertilizer application rates increased (Hahne et al. 1977, Schepers et al. 1991). Tile drainage promotes these leaching losses, reduces the amount of soil nitrogen, and increases nitrate in the groundwater, which eventually becomes surface water (Hallberg 1989).

John Johnston was the first person to lay drain tile in the United States, which he did on his farm in Ithaca, New York, in 1835 (USDA 1938). Drainage improved crop yields on marginal lands, but it took decades for the practice to spread to the Midwest (figure 2b), as marginal farmland came increasingly under cultivation after the passage of the Swamp Land Acts and the expansion of the railroads, which owned vast tracts of land. These drainage activities may not have been the only cause of greater losses of nitrogen in the soil, but they are indicative of the increased level of soil disturbance to the existing land and of expansion into formerly waterlogged soils.

The sequence of colonization, land clearing, agricultural expansion, soil erosion, increased nitrogen loading, and offshore diatom growth is summarized in figure 2. The BSi peaks and declines coincide with land-use changes resulting from land clearing, expansion of agriculture, and land drainage efforts within the MRB; they are modulated by the natural restoration of abandoned land. Note that the percentage of BSi in sediments accumulating offshore is higher now than it was during the peak in the mid-1800s. This recent rise in BSi concentration is undoubtedly related to increased nitrogen loading from the river, which occurred as a direct consequence of fertilizer application rising dramatically after World War II (figure 2d). Several analyses of whole watersheds have shown that the amount of fertilizer applied is sufficient to account for the postwar rise in nitrate in the Mississippi River (Peierls et al. 1991, Turner and Rabalais 1991, Howarth et al. 1996, Caraco and Cole 1999, Goolsby et al. 1999) and that urban and atmospheric nitrogen sources are relatively small (Howarth et al. 1996, Jordan and Weller 1996, Burkart and James 1999, Caraco and Cole 1999, Goolsby et al. 1999). However, the soil nitrogen pool is also significant, and it undoubtedly still contributes nitrate to the main channel of the Mississippi River.

Nutrient concentrations today

The effect of present land use on river water quality throughout the United States can be seen by examining the variability in water quality among watersheds. Jordan and colleagues (1997) showed that nitrate yields went up, and the ratios of dissolved silicate to nitrate went down, as the area of cropland increased in 27 watersheds of the Chesapeake Bay. Smart and colleagues (1985) studied watersheds in the Missouri Ozarks in the summer of 1979 and found that the silicate:

nitrate ratio and the nitrogen content in water went up as the land in pasture increased (there were apparently few row crops in that area); they concluded that the stream nutrient concentrations were more strongly related to land use than to bedrock geology. They developed a statistical model that used watershed size and the percentage of urban land to describe 43% of the variation in total phosphorus yields among watersheds. Eighty percent of the variation in total nitrogen was described using the percentage of land in pasture and urban area. Perkins and colleagues (1998) showed similar results for all four major types of Missouri watersheds, as did Jones and colleagues (1976) for 34 watersheds in northwestern Iowa (3-year data set). One interesting pattern in Jones and colleagues' (1976) data set is that the nitrogen yield dropped rapidly with a small increase in the percentage of land as marsh, and it continued to fall to less than 1 kg per hectare




Figure 6. Relationship between the percentage of a watershed in Iowa that is marsh and nitrogen from nitrate and ammonium (kilograms per hectare per year) for 17 watersheds over 100 hectares, 1971 to 1973. Adapted from data in Jones and colleagues (1976).

(ha) when the watershed was 15% marsh (figure 6). This finding suggests that strategic restoration of wetlands throughout the upper watershed may be a useful approach to restoring water quality (Mitsch et al. 2001), especially where the nitrate concentration is high (Turner 2003a).

Many studies have concluded that the application of fertilizers is a major source of the increased nutrient loading among large river watersheds in the last 50 years (NRC 1993). The major contributor to these changes in the Mississippi River watershed is nitrate (figure 7). These observations substantiate the conclusion that human intervention within the natural landscape has transformed water quality on the scale of the world's largest river basins and smaller coastal watersheds (e.g., Turner and Rabalais 1991, Howarth et al. 1996, Jordan et al. 1997, Vitousek et al. 1997, Caraco and Cole 1999, Turner et al. 2000, 2003, Seitzinger et al. 2002).

Changes in nutrient ratios, and not just nutrient concentrations, also have important consequences for diatoms in the receiving coastal waters, which are generally limited by silica deficiency when the ratio of silicate to dissolved nitrogen

drops below 1:1. Zooplankton graze on diatoms (which can be relatively large phytoplankton) and become part of the food web leading to fish production, among other things. The present-day concentrations of nitrate and silicate at New Orleans are very different from those in the early part of this century (figure 7), so much so that the atomic ratio of silicate to nitrate has fallen from about 4:1 to about 1:1. When the atomic ratio of silicate to nitrate falls below 1:1, the food web off the Mississippi River seems to switch from a diatom-based ecosystem to another ecosystem state that may be less desirable (Turner et al. 1998, Turner 2003b). The trends in many large rivers are in a similar direction (Turner et al. 2003). These changing nutrient ratios offshore of the Mississippi River affect an area from which 25% of the US commercial fisheries capture occurs. Many large rivers are changing in a similar way (Turner et al. 2003) and leading to eutrophication and hypoxia, which is having an unfavorable effect on the world's marine fisheries (Diaz and Rosenberg 1995). Thus, compromises to the quality and quantity of the diatoms in coastal waters could have unfavorable and significant consequences to food webs, commercial fisheries and recreational fishers, and coastal economies.




Figure 7. (a) Relationship between total nitrogen (y-axis) and combined concentration of nitrate and nitrite (x-axis) for 42 subwatersheds of the Mississippi River Basin (from data in Goolsby et al. 1999). (b) Relationship between the atomic ratio of dissolved silicon to dissolved inorganic nitrogen (y-axis) and the concentration of dissolved nitrate and nitrite (x-axis) for 42 subwatersheds of the Mississippi River Basin (from data in Goolsby et al. 1999) and for 529 individual sampling events at New Orleans.

The scale of water quality changes in the MRB over the last 200 years is substantial, and the social infrastructure supporting the humanmade landscape is nontrivial and important to many. This particular watershed has been anything but an equilibrium system for the last 200 years, suggesting that additional changes are forthcoming. Population growth alone will cause new changes in water quality. The landscape developments described herein have already resulted in nitrate levels in drinking water sources that exceed national standards.

These developments have also contributed to the formation of the largest coastal low-oxygen zone in the western North Atlantic (Rabalais and Turner 2001, Rabalais et al. 2002) and to the eutrophication of inland and coastal waters, which is a widely acknowledged social and environmental concern (e.g., Nixon 1995). Concerns about flood protection for farmland and homes are intermingled with concerns about diminishing natural resource quality and quantity and with national political agendas involving international trade, global climate change, and food supplies. It is our challenge as scientists, citizens, land managers, and agriculturists, among others, to work toward a mutually satisfying equilibrium of interests that is sustainable, ethical, and socially responsive. Our view of soil and land management can be much broader than the articulated and commonly held view of a popular soil textbook of 40 years ago: “After all, our primary aim in soil management is to seek the highest yields we can maintain consistent with greatest profit” (Thompson 1957, p. 363). This view of “profit” could be enlarged to include off-farm social costs and benefits, the soil’s health over decades, and the sustenance of the farming social structure, including the family-owned farm.

One thing seems certain: It took decades for the present system to develop, which suggests that it will take decades of working together for water quality rehabilitation to succeed. Three examples provide a perspective on just how patient we may have to be. First, the unfertilized fallow soil at the experimental soil plots in Rothamstead (United Kingdom) continues to leak significant amounts of nitrogen after 40 years (Addiscott 1988). A second example is from Sweden, where nitrate leaching from a grain field continued almost unabated 13 years after fertilizers were no longer added (Löfgren et al. 1999). Data on river water quality following the collapse of agriculture (circa 1990) in the former Soviet republics of Estonia, Latvia, and Lithuania show that, although fertilizer application fell to the level of the 1950s, the concentration of inorganic phosphate and nitrogen was the same in 1994 as in 1987 (Löfgren et al. 1999). The soil nitrogen mineralization of the huge soil nitrogen pool in these Baltic states is 50 to 200 kg nitrogen per ha compared with 64 to 93 kg nitrogen per ha of fertilizer applied. The strong buffering capacity of soils is becoming evident (Stålnacke et al. 1999). Understanding and managing the soil nutrient pools and turnover is a key factor in water quality management and will be inextricably commingled with the social structure and diverse incentives that have dominated soil cultivation practices for the last 200 years.

Acknowledgments

This analysis was prepared under a grant from the US Environmental Protection Agency, Gulf of Mexico Program, Project MX994721-95-0. Our research was funded by the National Oceanic and Atmospheric Administration’s Coastal Ocean Program under award NA06OP0528 to Louisiana Universities Marine Consortium and award NA06OP0529 to Louisiana State University. Erick M. Swenson assisted in the

data compilation; colleagues John Downing, Brian Fry, Dubravko Justic, and Dan J. Conley provided encouragement and support for the examination of changes in nutrient dynamics in this watershed. We thank Jerry Glover of the Land Institute for finding the photograph of the bluestem and wheat root system.

References cited

- Aber J, McDowell W, Nadelhoffer K, Magill A, Bernston G, Kamakea M, McNulty S, Currie W, Rustad L, Fernandez I. 1998. Nitrogen saturation in temperate forest ecosystems. *BioScience* 48: 921–934.
- Abernethy Y, Turner RE. 1987. US forested wetlands: 1940–1980. *BioScience* 37: 721–727.
- Addiscott TM. 1988. Long-term leakage of nitrate from bare unmanured soil. *Soil Use Management* 4: 91–95.
- Anonymous. 1992. *World Almanac and Book of Facts*. New York: Pharos Books.
- Baker J, Laflen JM. 1983. Water quality consequences of conservation tillage. *Journal of Soil and Water Conservation* 38: 186–193.
- Bateham MB. 1849. Exhaustion of the soil. *Ohio Cultivator* 5: 71.
- Bennett HH, Chapline WR. 1928. *Soil Erosion: A National Menace*. Washington (DC): US Government Printing Office. US Department of Agriculture Circular 33.
- Brune GM. 1948. *Rates of Sediment Production in Midwestern United States*. Washington (DC): US Soil Conservation Service. Technical report no. SCS-TP 65.
- Burkart MR, James DE. 1999. Agricultural-nitrogen contribution to hypoxia in the Gulf of Mexico. *Journal of Environmental Quality* 28: 850–859.
- Caraco NF, Cole JJ. 1999. Human impact on nitrate export: An analysis using major world rivers. *Ambio* 28: 167–170.
- Cronon W. 1983. *Changes in the Land: Indians, Colonists, and the Ecology of New England*. New York: Farrar, Straus and Giroux.
- Diaz RJ, Rosenberg R. 1995. Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna. *Oceanography and Marine Biology Annual Reviews* 33: 245–303.
- [EARI] Engineering Agency for Resource Inventories. 1975. *Inventory of Basic Environmental Data: New Orleans–Baton Rouge Metropolitan Area*. Prepared for the U.S. Army Corps of Engineers, New Orleans District. Washington (DC): EARI, US Army Engineers Topographic Laboratory.
- [EPA] US Environmental Protection Agency. 1998. *National Water Quality Inventory: 1996 Report to Congress*. Washington (DC): EPA Office of Water. Report no. EPA841-F-97-003.
- . 2002. *National Water Quality Inventory: 2000 Report*. Washington (DC): EPA Office of Water. Report no. EPA841-R-02-001. (24 April 2003; www.epa.gov/305b/2000report)
- Goolsby DA, Battaglin WA, Lawrence GB, Artz RS, Aulenbach BT, Hooper RP, Keeney DR, Stensland GJ. 1999. Flux and sources of nutrients in the Mississippi-Atchafalaya River Basin: Topic 3 report for the integrated assessment of hypoxia in the Gulf of Mexico. NOAA Coastal Ocean Program, Decision Analysis Series No. 17. (24 April 2003; www.nos.noaa.gov/Products/hypox_t3final.pdf)
- Gray LC. 1933. *History of Agriculture in the Southern United States to 1860*. Washington (DC): Carnegie Institute of Washington.
- Greeley WB. 1925. The relation of geography to timber supply. *Economic Geography* 1: 1–14.
- Hahne HCH, Kroontje W, Lutz JA Jr. 1977. Nitrogen fertilization, I: Nitrate accumulation and losses under continuous corn cropping. *Journal of the Soil Science Society of America* 41: 562–567.
- Hallberg GR. 1987. Nitrates in groundwater in Iowa. Pages 23–68 in D'Itri RM, Wolson LG, eds. *Rural Groundwater Contamination*. Chelsea (MI): Lewis Publishing.
- . 1989. Nitrate in ground water in the United States. Pages 35–74 in Follet RF, ed. *Nitrogen Management and Ground Water Protection*. Developments in Agricultural and Managed-Forest Ecology, vol. 21. Amsterdam (Netherlands): Elsevier.
- Howarth RE, et al. 1996. Regional nitrogen budgets and riverine N & P fluxes for the drainages to the North Atlantic Ocean: Natural and human influences. *Biogeochemistry* 35: 75–139.
- Humphreys AA, Abbot HL. 1876. Report upon the Physics and Hydraulics of the Mississippi River; upon the Protection of the Alluvial Region against Overflow; and upon the Deepening of the Mouths. Submitted to the Bureau of Topographical Engineers, War Department, 1861. Reprinted with additions. Washington (DC): US Government Printing Office.
- Jones JR, Borofka BP, Bachmann RW. 1976. Factors affecting nutrient loads in some Iowa streams. *Water Research* 10: 117–122.
- Jordan TE, Weller DE. 1996. Human contributions to terrestrial nitrogen flux. *BioScience* 46: 655–664.
- Jordan TE, Correll DL, Weller DE. 1997. Relating nutrient discharges from watersheds to land use and streamflow variability. *Water Resources Research* 33: 2579–2590.
- Keeney DR. 1986. Sources of nitrate to ground water. *Critical Reviews in Environmental Control* 16: 257–304.
- Leue A. 1886. *The Forestal Relations of Ohio*. First Annual Report of the Ohio State Forestry Bureau. Columbus (OH): Westbote Co., State Printers.
- Löfgren, S, Gustafson A, Steineck S, Ståhlacke P. 1999. Agricultural development and nutrient flows in the Baltic states and Sweden after 1988. *Ambio* 28: 320–327.
- Lowrance RR. 1992. Nitrogen outputs from a field-size agricultural watershed. *Journal of Environmental Quality* 21: 602–607.
- Meade RH, Yuzik TR, Day TJ. 1990. Movement and storage of sediment in rivers of the United States and Canada. Pages 255–280 in Wolman MG, Riggs HC, eds. *Surface Water Hydrology. The Geology of North America*, vol. O-1. Boulder (CO): Geological Society of America.
- Miller MF, Krusekopf HH. 1932. *The Influence of Systems of Cropping and Methods of Culture on Surface Runoff and Soil Erosion*. Columbia (MO): Missouri Agricultural Experiment Station. Research Bulletin 177.
- Mitsch WJ, Day JW Jr, Gilliam JW, Groffman PM, Hey DL, Randall GW, Wang N. 2001. Reducing nitrogen loading to the Gulf of Mexico from the Mississippi River Basin: Strategies to counter a persistent ecological problem. *BioScience* 51: 373–388.
- Muir J. 1965. *The Story of My Boyhood and Youth*. Madison: University of Wisconsin Press.
- Nixon SW. 1995. Coastal marine eutrophication: A definition, social causes, and future concerns. *Ophelia* 41: 199–219.
- Novotny V. 1999. Diffuse pollution from agriculture—a worldwide outlook. *Water Science and Technology* 39: 1–13.
- [NRC] National Research Council. 1993. *Soil and Water Quality: An Agenda for Agriculture*. Washington (DC): National Academy Press.
- Peierls B, Caraco N, Pace M, Cole J. 1991. Human influence on river nitrogen. *Nature* 350: 386–387.
- Perkins BD, Lohman K, Van Nieuwenhuysse E, Jones JR. 1998. An examination of land cover and stream water quality among physiographic provinces of Missouri, U.S.A. *Verhandlungen der Internationalen Vereinigung für Theoretische und Angewandte Limnologie* 26: 940–947.
- Pierce RS, Martin CW, Reeves CC, Likens GE, Bormann FH. 1972. Nutrient loss from clearcuttings in New Hampshire. Pages 285–295 in Csallany SC, McLaughlin TG, Striffler WD, eds. *Watersheds in Transition: Symposium of the American Water Resources Association*, Colorado State University, Fort Collins, CO. Urbana (IL): American Water Resources Association.
- Prince H. 1997. *Wetlands of the American Midwest: A Historical Geography of Changing Attitudes*. Chicago: University of Chicago Press. University of Chicago Research Paper no. 241.
- Quinn JB. 1894. Report of the Chief of Engineers. Pages 1345–1347 in House of Representatives Executive Documents, 53 Cong. 2 sess, vol. 2, pt. 3.
- Rabalais NN, Turner RE, eds. 2001. *Coastal Hypoxia: Consequences for Living Resources and Ecosystems*. Washington (DC): American Geophysical Union. *Coastal and Estuarine Studies* 58.

- Rabalais NN, Turner RE, Scavia D. 2002. Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi River. *BioScience* 52: 129–142.
- Schepers JS, Moravek MG, Alberts EE, Frank KD. 1991. Maize production impacts on groundwater quality. *Journal of Environmental Quality* 20: 12–16.
- Seitzinger SP, Kroeze C, Bouwman AF, Caraco N, Dentener F, Styles R. 2002. Global patterns of dissolved inorganic and particulate nitrogen inputs to coastal systems: Recent conditions and future projections. *Estuaries* 25: 640–655.
- Smart MM, Jones JR, Sebaugh JL. 1985. Stream-watershed relations in the Missouri Ozark Plateau Province. *Journal of Environmental Quality* 14: 77–82.
- Smith RA, Alexander RB, Lanfear KJ. 1996. Stream water quality in the conterminous United States—status and trends of selected indicators during the 1980s. US Geological Survey Water Supply Paper no. 2400. (23 April 2003; <http://water.usgs.gov/nwsum/sal/index.html>)
- Stålnacke P, Vagstad N, Tamminen T, Wassmann P, Jansons V, Loigo E. 1999. Nutrient runoff and transfer from land and rivers to the Gulf of Riga. *Hydrobiologia* 410: 103–110.
- Thompson LM. 1957. *Soils and Soil Fertility*. New York: McGraw-Hill.
- Trimble SW. 1985. Perspectives on the history of soil erosion control in the eastern United States. *Agricultural History* 59: 162–180.
- . 1999. Decreased rates of alluvial sediment storage in the Coon Creek Basin, Wisconsin, 1975–1993. *Science* 285: 1244–1246.
- Turner RE. 2003a. Nitrogen and phosphorus concentration and retention in water flowing over freshwater wetlands. In Fredrikson LH, King SL, Kaminski RM, eds. *Ecology and Management of Bottomland Hardwood Systems: The State of Our Understanding*. Puxico (MO): University of Missouri–Columbia, Gaylord Memorial Laboratory. Forthcoming.
- . 2003b. Element ratios in aquatic food webs. *Estuaries* 25: 694–703.
- Turner RE, Rabalais NN. 1991. Changes in the Mississippi River this century: Implications for coastal food webs. *BioScience* 41: 140–147.
- . 1994. Coastal eutrophication near the Mississippi River delta. *Nature* 368: 619–621.
- Turner RE, Qureshi N, Rabalais NN, Dortch Q, Justic' D, Shaw R, Cope J. 1998. Fluctuating silicate:nitrate ratios and coastal plankton food webs. *Proceedings of the National Academy of Sciences* 95: 13,048–13,051.
- Turner RE, Stanley D, Brock D, Pennock J, Rabalais NN. 2000. A comparison of independent N-loading estimates for U.S. estuaries. Pages 107–118 in Valigura RW, Alexander RB, Castro MS, Meyers TP, Paerl HW, Stacey PE, Turner RE, eds. *Nitrogen Loading in Coastal Water Bodies: An Atmospheric Perspective*. Washington (DC): American Geophysical Union. *Coastal and Estuarine Studies* 57.
- Turner RE, Rabalais NN, Justic' D, Dortch Q. 2003. Global patterns of dissolved silicate and nitrogen in large rivers. *Biogeochemistry*. Forthcoming.
- Ubelaker DH. 1992. The sources and methodology for Mooney's estimates of North American Indian populations. Pages 243–288 in Denevan WH, ed. *The Native Population of the Americas in 1492*. Madison (WI): Wisconsin Press.
- [USCB] US Census Bureau. 1961. *U.S. Census of Agriculture: 1959, vol. 4: Drainage of Agricultural Lands*. Washington (DC): US Department of Commerce.
- . 1973. *1969 Census of Agriculture, vol. 6: Drainage of Agricultural Lands*. Washington (DC): US Department of Commerce.
- [USDA] US Department of Agriculture. 1938. *Soils and Men. Yearbook of Agriculture, 1938*. Washington (DC): USDA.
- Vitousek PM, Aber JD, Howarth RW, Likens GE, Matson PA, Schindler DW, Schlesinger WH, Tilman DG. 1997. Human alteration of the global nitrogen cycle: Sources and consequences. *Ecological Applications* 7: 737–750.
- Whitney GG. 1994. *From Coastal Wilderness to Fruited Plain: A History of Environmental Change in Temperate North America from 1500 to the Present*. Cambridge (United Kingdom): Cambridge University Press.
- Wolman MG. 1967. A cycle of sedimentation and erosion in urban river channels. *Geografiska Annaler* 49A: 385–395.