

LIOUVILLE BROWNIAN MOTION

BY CHRISTOPHE GARBAN¹, RÉMI RHODES² AND VINCENT VARGAS²

Université Lyon 1, Université Paris-Est, Marne la Vallée and CNRS, ENS Paris

We construct a stochastic process, called the *Liouville Brownian motion*, which is the Brownian motion associated to the metric $e^{\gamma X(z)} dz^2$, $\gamma < \gamma_c = 2$ and X is a Gaussian Free Field. Such a process is conjectured to be related to the scaling limit of random walks on large planar maps eventually weighted by a model of statistical physics which are embedded in the Euclidean plane or in the sphere in a conformal manner. The construction amounts to changing the speed of a standard two-dimensional Brownian motion B_t depending on the local behavior of the Liouville measure “ $M_\gamma(dz) = e^{\gamma X(z)} dz$ ”. We prove that the associated Markov process is a Feller diffusion for all $\gamma < \gamma_c = 2$ and that for all $\gamma < \gamma_c$, the Liouville measure M_γ is invariant under P_t . This Liouville Brownian motion enables us to introduce a whole set of tools of stochastic analysis in Liouville quantum gravity, which will be hopefully useful in analyzing the geometry of Liouville quantum gravity.

1. Introduction. An important issue for applications in $2d$ -Liouville quantum gravity is to construct a random metric on a two-dimensional Riemann manifold D , say a domain of \mathbb{R}^2 (or the sphere) equipped with the Euclidean metric dz^2 , which takes on the form

$$(1.1) \quad e^{\gamma X(z)} dz^2,$$

where X is a Gaussian Free Field (GFF) on the manifold D and $\gamma \in [0, 2)$ is a coupling constant that can be expressed in terms of the central charge of the model coupled to gravity (see [14, 29] for further details and also [18, 24, 31] for insights in Liouville quantum gravity). The simplicity of such an expression hides many highly nontrivial mathematical difficulties. Indeed, the correlation function of a GFF presents a short scale logarithmically divergent behavior that makes relation (1.1) nonrigorous. One has to apply a cutoff procedure to smooth down the singularity of the GFF and the method to do this at a metric level remains unclear. However, many geometric quantities are related to this metric and for some of them, the cutoff procedure may be applied properly. For instance, the construction

Received May 2014; revised June 2015.

¹Supported in part by the ANR Grant MAC2 10-BLAN-0123.

²Supported in part by Grant ANR-11-JCJC CHAMU.

MSC2010 subject classifications. 60D05, 28A80.

Key words and phrases. Liouville quantum gravity, Liouville Brownian motion, Gaussian multiplicative chaos.

of the volume form, also called the Liouville measure, was carried out by Kahane within the framework of Gaussian multiplicative chaos [27] (see also [21, 36, 37] for more recent constructions based on convolution techniques). This allows us to give a rigorous meaning to the expression

$$(1.2) \quad M_\gamma(A) = \int_A e^{\gamma X(z) - (\gamma^2/2)\mathbb{E}[X(z)^2]} dz,$$

where dz stands for the volume form (Lebesgue measure) on D [to be exhaustive, one should integrate against $h(z) dz$ where h is a deterministic function involving the conformal radius at z but this term does not play an important role for our concerns]. This strategy made possible an interpretation in terms of measures of the Knizhnik–Polyakov–Zamolodchikov formula (KPZ for short, see [29]) relating the fractal dimensions of sets as seen by the Lebesgue measure or the Liouville measure. The KPZ formula is proved in [21] when considering the fractal notion of expected box counting dimension whereas the fractal notion of almost sure Hausdorff dimension is considered in [6, 34] (see also [9]). The reader may consult [7, 9, 14, 16, 18–20, 23, 24, 31, 35] for more references on this topic. Another important part of the theory which we do not review here is that it is conjectured to be the scaling limit of discrete quantum gravity: the reader may consult [21] for more on this topic as well as physics references therein.

Another powerful tool in describing a Riemann geometry is the Brownian motion. With it are attached several analytic objects serving to describe the geometry: a semigroup, a Laplace–Beltrami operator, a heat kernel, Dirichlet forms, etc. Therefore, a relevant way to have further insights into Liouville quantum gravity geometry is to define the *Liouville Brownian motion* (LBM for short). This is the purpose of this paper. It can be constructed on any background $2d$ -Riemann manifold equipped with a GFF X and can be seen as the Brownian motion associated to the metric $e^{\gamma X(x)} dx^2$ where $\gamma \in [0, 2[$ is a parameter and dx^2 stands for the metric on the manifold.

In this paper and for pedagogical purposes, we will mostly describe the situation when the underlying manifold is the whole plane \mathbb{R}^2 , in which case it is natural to consider a Massive Gaussian Free Field X on \mathbb{R}^2 (MFF for short). We will also explain how to adapt our framework to the cases of the sphere \mathbb{S}^2 , the torus \mathbb{T}^2 or planar bounded domains. More generally, it is also clear that our methodology may apply to any 2-dimensional Riemann manifold equipped with a log-correlated Gaussian field and yields similar results. Let us also mention that another work [10] appeared online simultaneously to ours and is concerned with the LBM starting from one point: the paper [10] proves that for fixed x , one can define almost surely the LBM starting from x (the work [10] also initiates a multi-fractal analysis of the LBM starting from one point). We will show that almost surely we can define the LBM starting from all x , hence obtaining the existence of a diffusion process associated to the tensor (1.1) and all the related stochastic analysis tools.

Finally, we point out that the notions of diffusion or heat kernel are at the core of the physics literature about Liouville quantum gravity (see [3, 4, 12, 13, 15, 16, 41], e.g., among a huge amount of other works). For instance, a heat kernel derivation of the KPZ formula is obtained in [16]. The fractal structure of quantum space–time is also investigated in [3, 4, 15, 41] via diffusions and heat kernel properties, obtaining relations about the fractal dimensions of quantum space–time.

2. Liouville Brownian motion on the plane. In this section, we construct Liouville Brownian motion on the whole plane. Regarding the physics literature, the natural free field to consider on the whole plane is the Massive Gaussian Free Field (MFF for short). So we first remind the reader of the construction of the MFF after introducing a few basic notation. Then we recall the construction of Gaussian multiplicative chaos associated to the MFF and state a few basic properties, which are used thereafter to construct the Liouville Brownian motion.

2.1. Basic notation and terminology.

Basics. In what follows, the Liouville Brownian motion that we are going to construct will be denoted by $(\mathcal{B}_t)_{t \geq 0}$. We distinguish the (quantum) time t along \mathcal{B}_t and the (classical) time t along a standard Brownian motion B_t . The open ball centered at x with radius R is denoted $B(x, R)$ and the closed ball $\bar{B}(x, R)$. \mathbb{S}^2 and \mathbb{T}^2 stand for the two-dimensional sphere or torus.

Functional spaces and analysis. The space of continuous functions with compact support in a domain D (resp., vanishing at infinity on \mathbb{R}^2 , resp., bounded functions on D , resp., continuous functions on \mathbb{R}_+ equipped with the sup-norm topology over compact sets) is denoted by $C_c(D)$ [resp., $C_0(\mathbb{R}^2)$, resp., $C_b(D)$, resp., $C(\mathbb{R}_+)$].

The standard Laplace–Beltrami operator on a manifold is denoted by Δ . We will say that a semigroup on $C_b(D)$ is Feller if the semigroup maps $C_b(D)$ into itself. We will say that a Markov process on D is Feller if its semigroup is.

Positive continuous additive functionals and Revuz measures. Let us consider a standard Brownian motion $(\Omega_B, (B_t)_{t \geq 0}, (\mathcal{F}_t)_{t \geq 0}, (\mathbb{P}_x^B)_{x \in D})$ in D (with $D = \mathbb{R}^2$, \mathbb{S}^2 or \mathbb{T}^2). It is reversible for the canonical volume form dx of D . We suppose that the space Ω_B is equipped with the standard shifts $(\theta_t)_{t \geq 0}$ on the trajectory. One may then consider the classical notion of capacity Cap associated to the Brownian motion (see [22]). The set K is said polar when $\text{Cap}(K) = 0$.

A Revuz measure μ is a Radon measure on D which does not charge the polar sets. A positive continuous additive functional (PCAF) $(A_t)_{t \geq 0}$ is a \mathcal{F}_t -adapted continuous functional with values in $[0, \infty]$ that satisfies for all $\omega \in \Lambda$:

$$A_{t+s}(\omega) = A_s(\omega) + A_t(\theta_s(\omega)), \quad s, t \geq 0,$$

where Λ is a subset of Ω_B such that $\forall x \in D, \mathbb{P}_x^B(\Lambda) = 1$ and $\theta_t(\Lambda) \subset \Lambda$ for all $t \geq 0$. In particular, we will always consider PCAFs defined for all starting points $x \in D$ (they are sometimes called *PCAF in the strict sense* in the literature, especially in [22]).

Massive Gaussian Free Field on the plane. We consider a whole plane Massive Gaussian Free Field (MFF) (see [25, 39] for an overview of the construction of the MFF and applications). Given a real number $m > 0$, it is a centered Gaussian random distribution (in the sense of Schwartz) with covariance function given by the Green function G_m of the operator $m^2 - \Delta$, that is,

$$(m^2 - \Delta)G_m(x, \cdot) = 2\pi\delta_x.$$

Notice that G_m is π times the Green function of the Brownian motion killed at rate $m^2/2$. It is a standard fact that the massive Green function can be written as an integral of the transition densities of the Brownian motion weighted by the exponential of the mass:

$$(2.1) \quad \forall x, y \in \mathbb{R}^2, \quad G_m(x, y) = \int_0^\infty e^{-(m^2/2)u - |x-y|^2/(2u)} \frac{du}{2u}.$$

Clearly, it is a kernel of σ -positive type in the sense of Kahane [27] since we integrate a continuous function of positive type with respect to a positive measure. One can also check that

$$(2.2) \quad G_m(x, y) = \ln_+ \frac{1}{|x - y|} + g_m(x, y),$$

for some continuous and bounded function g_m and $\ln_+ x = \max(\ln x, 0)$.

It is furthermore a star-scale invariant kernel (see [2, 33]): it can be rewritten as

$$(2.3) \quad G_m(x, y) = \int_1^{+\infty} \frac{k_m(u(x - y))}{u} du,$$

for some continuous covariance kernel $k_m(z) = \frac{1}{2} \int_0^\infty e^{-(m^2/(2v))|z|^2 - v/2} dv$. In particular, we will make intensive use of the following relation, valid for $\epsilon \in]0, 1]$:

$$(2.4) \quad G_m(x, y) \leq G_m\left(\frac{x}{\epsilon}, \frac{y}{\epsilon}\right) + \ln \frac{1}{\epsilon}.$$

Now we consider an unbounded strictly increasing sequence $(c_n)_{n \geq 1}$ such that $c_1 = 1$. For each $n \geq 1$, we consider a centered Gaussian process Y_n with covariance kernel given by

$$(2.5) \quad \mathbb{E}[Y_n(x)Y_n(y)] = \int_{c_{n-1}}^{c_n} \frac{k_m(u(x - y))}{u} du.$$

The reader may check that such a process is stationary and has smooth sample paths (to check this point, apply the Kolmogorov criterion to Y_n as well as its

derivatives in the standard manner). The MGFF is the Gaussian distribution defined by

$$X(x) = \sum_{n \geq 1} Y_n(x),$$

where the processes $(Y_n)_n$ are assumed to be independent. We define the n -regularized field by

$$(2.6) \quad X_n(x) = \sum_{k=1}^n Y_k(x).$$

Actually, based on Kahane’s theory of multiplicative chaos [27], the choice of the decomposition (2.6) will not play a part in the forthcoming results, except that it is important that the covariance kernel of X_n be smooth in order to associate to this field a Riemann geometry.

NOTATION 2.1. In what follows, we will consider a Brownian motion on \mathbb{R}^2 (or other two-dimensional manifolds) $(\Omega_B, \mathcal{F}_B, (B_t)_{t \geq 0}, (\mathcal{F}_t)_{t \geq 0}, (\mathbb{P}_x^B)_{x \in \mathbb{R}^2})$. We will also consider a MFF X (and all the corresponding $(Y_n)_n$) defined on a probability space $(\Omega_X, \mathcal{F}_X, \mathbb{P}^X)$. So we consider a measurable space $(\Omega, \mathcal{F}) = (\Omega_X \times \Omega_B, \mathcal{F}_X \otimes \mathcal{F}_B)$ on which are defined both the MFF X and the Brownian motion B . On this measurable space are defined the probability measures $\mathbb{P}_x = \mathbb{P}^X \otimes \mathbb{P}_x^B$ (with expectation \mathbb{E}_x) for all $x \in \mathbb{R}^2$. Notice that under \mathbb{P}_x , the MFF X and the Brownian motion are independent. We will also denote by \mathcal{F}_n the sigma-algebra generated by the fields $(Y_k)_{k \leq n}$, that is, $\mathcal{F}_n = \sigma\{Y_k(x); k \leq n, x \in \mathbb{R}^2\}$. Finally, we mention that we will sometimes consider other Brownian motions B, W : the convention of notation will be the same as for B .

2.2. *Gaussian multiplicative chaos.* Let us fix $\gamma \geq 0$. We consider the random measure for $n \geq 1$ (the constant c_n is defined in the previous subsection)

$$(2.7) \quad M_n(dx) = c_n^{-\gamma^2/2} e^{\gamma X_n(x)} dx,$$

defined on the Borel sets of \mathbb{R}^2 , which will be called n -regularized Liouville measure. Classical theory of Gaussian multiplicative chaos ([27] or [35], Theorem 2.5) ensures that, \mathbb{P}^X almost surely, the family $(M_n)_{n \geq 1}$ weakly converges as $n \rightarrow \infty$ toward a limiting Radon measure M , which is called the Liouville measure. The limiting measure is nontrivial if and only if $\gamma \in [0, 2)$. We will denote by ξ_M the power law spectrum of M (see [2, 5, 34], e.g.):

$$(2.8) \quad \forall p \geq 0, \quad \xi_M(p) = \left(2 + \frac{\gamma^2}{2}\right)p - \frac{\gamma^2}{2}p^2.$$

Recall (see [27] or [35], Theorems 2.11 and 2.12) that for all bounded Borel set A and $p < 4/\gamma^2$, we have $\mathbb{E}[M(A)^p] < +\infty$ and that

$$(2.9) \quad \sup_{r < 1} r^{-\xi_M(p)} \mathbb{E}[M(rA)^p] \leq C_p$$

for some constant C_p only depending on p .

Let us emphasize that Kahane’s theory of Gaussian multiplicative chaos ensures that the law of the measure M does not depend on the chosen regularization $(X_n)_n$ of X (see [27, 35, 37]). Furthermore, in the case of a GFF X on a planar domain, this result is reinforced in [21] for the Liouville measure: the authors prove that circle average approximations of X and projections of X along any H^1 basis yields almost surely the same Liouville measure. For more recent results on existence and uniqueness, see [38].

We state below a result about the local modulus of continuity of the measure M as well as its approximating sequence $(M_n)_n$.

THEOREM 2.2. *We set $\alpha = 2(1 - \frac{\gamma}{2})^2 > 0$. Let $\epsilon > 0$ and $R > 0$. \mathbb{P}^X -almost surely, there exists a random constant $C > 0$ such that:*

$$\sup_{r \in (0,1)} \sup_{x \in [-R,R]^2} \sup_{n \geq 1} r^{-\alpha+\epsilon} (M_n(B(x,r))) + M(B(x,r)) \leq C.$$

PROOF. It suffices to prove the result only for M because $(M_n)_n$ is a martingale converging a.s. toward M (use Doob’s inequalities to estimate the $\sup_n M_n$ in terms of M).

We take $R = \frac{1}{2}$ for simplicity. Now, we partition $[-\frac{1}{2}, \frac{1}{2}]^2$ into 2^{2n} dyadic squares $(I_n^j)_{1 \leq j \leq 2^{2n}}$ of equal size. If p belongs to $]0, \frac{4}{\gamma^2}[$, we get

$$\begin{aligned} \mathbb{P}^X \left(\sup_{1 \leq j \leq 2^{2n}} M(I_n^j) \geq \frac{1}{2^{(\alpha-\epsilon)n}} \right) &\leq 2^{p(\alpha-\epsilon)n} \mathbb{E}^X \left[\sum_{1 \leq j \leq 2^{2n}} M(I_n^j)^p \right] \\ &\leq \frac{C_p}{2^{(\xi_M(p)-2-(\alpha-\epsilon)p)n}}. \end{aligned}$$

By taking $p = \frac{2}{\gamma}$ in the above inequalities [i.e., $\xi_M(p) - 2 - (\alpha - \epsilon)p > 0$] and by using Borel–Cantelli’s lemma, we obtain that, \mathbb{P}^X almost surely, there exists a random constant C such that

$$\sup_{1 \leq j \leq 2^{2n}} M(I_n^j) \leq \frac{C}{2^{(\alpha-\epsilon)n}} \quad \forall n \geq 1.$$

We conclude by the fact that each ball $B(x, r)$ is contained in at most 4 dyadic squares $(I_n^j)_{1 \leq j \leq 2^{2n}}$ when we choose n such that $\frac{1}{2^{n+1}} < r \leq \frac{1}{2^n}$. \square

2.3. Potential of the measure M . For each $R > 0$, let us introduce the Green function G_R of the Laplacian on the ball $B(0, R)$ with Dirichlet boundary conditions, that is, $(\delta_x$ stands for the Dirac mass at x)

$$(2.10) \quad \Delta G_R(x, \cdot) = -2\delta_x(\cdot), \quad G_R(x, \cdot)|_{\partial B(0,R)} = 0.$$

Keep in mind the distinction between the massive Green function G_m defined by (2.1) on \mathbb{R}^2 and the Green function G_R on a ball $B(0, R)$. Despite the similar notation, this should bring no confusion as we will always refer to the massive

Green function when the subscript is m and the Green function on balls when the subscript is R .

We introduce the R -potential of a Borel measure μ on \mathbb{R}^2 by

$$\forall x \in B(0, R), \quad g_R(\mu)(x) := \int_{B(0,R)} G_R(x, y)\mu(dy).$$

Let us consider the set of measures

$$\mathcal{M} = \{M_n; n \geq 1\} \cup \{M\}.$$

Furthermore, for $x \in \mathbb{R}^2$, we denote by M^z the shifted measure $M^z(\cdot) = M(z + \cdot)$.

Now we use Theorem 2.2 to prove the following.

PROPOSITION 2.3. *For any $R > 0$, \mathbb{P}^X -almost surely, we have:*

1. $\sup_{\mu \in \mathcal{M}} \sup_{x \in B(0,R)} g_R(\mu)(x) < +\infty$,
2. for any $\mu \in \mathcal{M}$, the mapping $x \in \bar{B}(0, R) \mapsto g_R(\mu)(x)$ is continuous,
3. $\sup_{x \in B(0,R)} |g_R(M_n)(x) - g_R(M)(x)| \rightarrow 0$ as $n \rightarrow \infty$,
4. for any $z_0 \in \mathbb{R}^2$, $\lim_{z \rightarrow z_0} \sup_{x \in B(0,R)} |g_R(M^z)(x) - g_R(M^{z_0})(x)| = 0$.

PROOF. Recall that the Green function G_R satisfies for all $x, y \in \bar{B}(0, R)$

$$(2.11) \quad G_R(x, y) \leq \frac{1}{\pi} \ln \frac{1}{|x - y|} + C$$

for some constant C . Therefore, it suffices to prove that

$$\sup_{\mu \in \mathcal{M}} \sup_{x \in B(0,R)} \int_{B(0,R)} \ln \frac{1}{|x - y|} \mu(dy) < +\infty.$$

From Theorem 2.2, we can find a constant C and $\alpha > 0$ (depending on γ and R) such that for all $x \in \bar{B}(0, 2R)$ and all $r \in (0, R)$

$$\sup_{\mu \in \mathcal{M}} \mu(B(x, r)) \leq Cr^\alpha.$$

For $\mu \in \mathcal{M}$, we have

$$(2.12) \quad \begin{aligned} \int_{B(0,R)} \ln \frac{1}{|x - y|} \mu(dy) &\leq \sum_{n \geq 0} \int_{B(0,R) \cap \{2^{-n}R < |x-y| \leq 2^{-n+1}R\}} \ln \frac{1}{|x - y|} \mu(dy) \\ &\leq \sum_{n \geq 0} (n \ln 2 - \ln R) \mu(B(x, 2^{-n+1}R)) \\ &\leq C2^\alpha \sum_{n \geq 1} (n \ln 2 - \ln R) R^\alpha 2^{-\alpha n}. \end{aligned}$$

This latter quantity is finite and does not depend on x or $\mu \in \mathcal{M}$. This proves the first part of our statement.

For the second statement, consider a function $\theta : \mathbb{R} \rightarrow \mathbb{R}$ such that $0 \leq \theta \leq 1$, $\theta(x) = 1$ for $|x| \leq 1$ and $\theta(x) = 0$ for $|x| \geq 2$. For $\delta > 0$ set $\theta_\delta(x) = \theta(x/\delta)$ and $\bar{\theta}_\delta(x) = 1 - \theta_\delta(x)$. Choose $\mu \in \mathcal{M}$. Observe that for all $\delta > 0$

$$\begin{aligned}
 g_R(\mu)(x) &= \int_{B(0,R)} G_R(x,y)\theta_\delta(x-y)\mu(dy) \\
 (2.13) \qquad &+ \int_{B(0,R)} G_R(x,y)\bar{\theta}_\delta(x-y)\mu(dy) \\
 &=: A_{\mu,\delta}(x) + D_{\mu,\delta}(x).
 \end{aligned}$$

We will show that the mappings $x \mapsto D_{\mu,\delta}(x)$ converge uniformly toward $x \mapsto g_R(\mu)(x)$ as $\delta \rightarrow 0$. As it is obvious to check that the mapping $x \mapsto D_{\mu,\delta}(x)$ is continuous with the help of standard theorems of continuity for parameterized integrals, this will show that $x \mapsto g_R(\mu)(x)$ is continuous. So, let us show that the family of mappings $x \mapsto A_{\mu,\delta}(x)$ converges uniformly toward 0 on $\bar{B}(0, R)$ as $\delta \rightarrow 0$. From (2.11) again, it is enough to show that

$$\sup_{x \in \bar{B}(0,R)} \int_{B(0,R)} \ln \frac{1}{|x-y|} \theta_\delta(x-y)\mu(dy) \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

In fact, we will prove a stronger statement. With computations similar to (2.12), we get

$$\begin{aligned}
 &\sup_{\mu \in \mathcal{M}} \int_{B(0,R)} \ln \frac{1}{|x-y|} \theta_\delta(x-y)\mu(dy) \\
 (2.14) \quad &\leq \sup_{\mu \in \mathcal{M}} \sum_{n \geq \ln 2/(-\ln 4\delta)} \int_{B(0,R) \cap \{2^{-n-1} < |x-y| \leq 2^{-n}\}} \ln \frac{1}{|x-y|} \mu(dy) \\
 &\leq C \ln 2 \sum_{n \geq \ln 2/(-\ln 4\delta)} (n+1)2^{-an}.
 \end{aligned}$$

This latter series converges to 0 as $\delta \rightarrow 0$. The function $g_R(\mu)$ is thus continuous as a uniform limit of continuous functions.

We now prove the third statement. Sticking to the previous notation (2.13), we have

$$g_R(M_n)(x) =: A_{M_n,\delta}(x) + D_{M_n,\delta}(x).$$

From (2.14), we have $\sup_{\mu \in \mathcal{M}} \sup_{x \in \bar{B}(0,R)} |A_{\mu,\delta}(x)| \rightarrow 0$ as $\delta \rightarrow 0$. Therefore, it suffices to prove that for each fixed $\delta > 0$, the family $(D_{M_n,\delta})_n$ converges uniformly on $\bar{B}(0, R)$ toward $D_{M,\delta}$. Point-wise convergence is ensured by the weak convergence of the family of measure $(M_n)_n$ toward M as $n \rightarrow \infty$. We just have to show that the family $(D_{M_n,\delta})_n$ is relatively compact for the topology of uniform convergence on $\bar{B}(0, R)$. For each fixed $\delta > 0$, the mapping

$(x, y) \mapsto G_R(x, y)\bar{\theta}_\delta(x - y)$ is continuous on $\bar{B}(0, R)^2$ and, therefore, uniformly continuous. The quantity

$$\omega(\eta) = \sup_{|x-x'|\leq\eta, (x,x',y)\in\bar{B}(0,R)^3} |G_R(x, y)\bar{\theta}_\delta(x - y) - G_R(x', y)\bar{\theta}_\delta(x' - y)|$$

thus converges toward 0 as $\eta \rightarrow 0$. We have for $x, x' \in \bar{B}(0, R)^2$

$$|D_{M_n,\delta}(x) - D_{M_n,\delta}(x')| \leq \omega(|x - x'|)M_n(\bar{B}(0, R)).$$

We complete the proof with the Arzelà–Ascoli criterion and the relation $\sup_n M_n(\bar{B}(0, R)) < +\infty$ almost surely. The proof of item 4 can be handled the same way, so we let the reader check the details. \square

2.4. *Approximation and construction of the PCAF of M .* For $R > 0$, we further introduce the stopping time

$$T_R = \inf\{t > 0; B_t \notin B(0, R)\}.$$

Observe that Theorem 2.2 (or Proposition 2.3) implies that each $\mu \in \mathcal{M}$ does not charge any polar set. Following [11], Proposition 3.2, we deduce that \mathbb{P}^X a.s. we can associate to each $\mu \in \mathcal{M}$ a unique PCAF $(F_t^{\mu,R})_t$ such that the process

$$\forall t \geq 0, \quad g_R(\mu)(B_{t \wedge T_R}) - g_R(\mu)(B_0) + F_t^{\mu,R}$$

is a mean zero martingale under \mathbb{P}_x^B for all $x \in B(0, R)$. The Revuz measure of the PCAF $F^{\mu,R}$ is μ [restricted to the ball $B(0, R)$].

NOTATION 2.4. When $\mu = M_n$ for some $n \geq 1$, we write $F_R^n(t)$ instead of the heavy notation $F_t^{M_n,R}$. Similarly, we write $F_R(t)$ for $F_t^{M,R}$.

It may be worth mentioning that we have the explicit expression

$$(2.15) \quad F_R^n(t) = c_n^{-\gamma^2/2} \int_0^{t \wedge T_R} e^{\gamma X_n(B_r)} dr.$$

The purpose of what follows is now to establish the convergence of the family of PCAFs $(F_R^n)_n$ toward F_R . Following [8], we consider the following distance between two Radon measures μ, ν on $\bar{B}(0, R)$:

$$d_R(\mu, \nu) = \sup_{x \in \bar{B}(0,R)} |g_R(\mu)(x) - g_R(\nu)(x)|.$$

Now we state the following lemma; the proof of which is omitted as a straightforward adaptation of [8], Proposition 2.1.

LEMMA 2.5. For all $R, \eta > 0, x \in \bar{B}(0, R)$ and μ_1, μ_2 two Borel measures such that $d_R(\mu_1, \mu_2) \leq 1$, we have

$$\mathbb{P}_x^B \left(\sup_{t \geq 0} |F_t^{\mu_1, R} - F_t^{\mu_2, R}| \geq \eta \right) \leq c_R \exp \left(- \frac{\eta}{c_R \sqrt{d_R(\mu_1, \mu_2)}} \right)$$

for some constant c_R that only depends (increasingly) on

$$\sup_{i=1,2} \sup_{x \in B(0, R)} g_R(\mu_i)(x).$$

From Proposition 2.3 item 3, \mathbb{P}^X -almost surely and for all $R > 0$, we have $d_R(M_n, M) \rightarrow 0$ as $n \rightarrow \infty$. We deduce the following.

COROLLARY 2.6. \mathbb{P}^X -almost surely, for all $x \in \bar{B}(0, R)$, we have

$$\mathbb{P}_x^B \left(\sup_{t \geq 0} |F_R^n(t) - F_R(t)| \geq \eta \right) \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

THEOREM 2.7. \mathbb{P}^X -almost surely, there exists a unique PCAF denoted by F such that

$$F(t) = F_R(t), \quad \text{for } t < T_R.$$

Furthermore, \mathbb{P}^X -almost surely:

1. the Revuz measure of F is M ,
2. for all $x \in \mathbb{R}^2$ and $T > 0, \mathbb{P}_x^B(\sup_{t \leq T} |c_n^{-\gamma^2/2} \int_0^t e^{\gamma X_n(B_r)} dr - F(t)| \geq \eta) \rightarrow 0$ as $n \rightarrow \infty$,
3. for all $x \in \mathbb{R}^2, \mathbb{P}_x^B$ -a.s., F is strictly increasing,
4. for all $x \in \mathbb{R}^2, \mathbb{P}_x^B$ -a.s., $\lim_{t \rightarrow \infty} F(t) = +\infty$,
5. the law of the pair (B, F) under \mathbb{P}_x^B on the space of continuous functions on \mathbb{R}_+ equipped with the topology of uniform convergence over compact sets is a continuous function of x , meaning

$$\lim_{x \rightarrow x_0} \mathbb{E}_x^B [G(B, F)] = \mathbb{E}_{x_0}^B [G(B, F)]$$

for every bounded continuous function on $C([0, T], \mathbb{R}_+)$ (for $T > 0$) and $x_0 \in \mathbb{R}^2$.

PROOF. Existence and uniqueness of such a PCAF is a straightforward consequence of the previous results.

Item 2 results from Corollary 2.6 provided that one takes R large enough to make $\mathbb{P}_x^B(T_R < T)$ arbitrarily small.

Item 1 results from [22], Theorem 5.1.3 and Lemma 5.1.10, and [11], Proposition 3.2. Indeed, this shows that F coincides with the whole plane PCAF of the measure M (recall that it does not charge polar sets).

Now we focus on items 3 and 4. Obviously, \mathbb{P}^X -almost surely and for all $x \in \mathbb{R}^2$, the mapping $t \in \mathbb{R}_+ \mapsto F(t)$ is increasing \mathbb{P}_x^B -almost surely. This mapping thus defines a measure on \mathbb{R}_+ , which we still denote by F with a slight abuse of notation.

From now on, we will use a few auxiliary lemmas along the main argument: their proofs are postponed after that of Theorem 2.7. Recall the definition of $(\mathcal{F}_n)_n$ in Section 2.1.

LEMMA 2.8. *For each fixed $x \in \mathbb{R}^2$, \mathbb{P}_x^B -almost surely, for all $t \geq 0$ and $R > 0$, the family $(F_R^n(t))_n$ is a uniformly integrable martingale with respect to the filtration $(\mathcal{F}_n)_n$, which converges \mathbb{P}^X -almost surely toward $F_R(t)$. We have, \mathbb{P}_x^B -almost surely, $\mathbb{E}^X[F_R(t)] = t \wedge T_R$.*

We prove item 3. We fix $x \in \mathbb{R}^2$ and we first prove that \mathbb{P}_x -a.s. F is strictly increasing. It suffices to prove that it is strictly increasing on $[0, T_R[$ for all $R > 0$. We consider a nonempty interval $I = [s, t]$ with $t < T_R$. \mathbb{P}_x^B -a.s., the event $\{F_R(I) > 0\}$ is an event belonging to the asymptotic sigma-algebra generated by the random processes $(Y_n)_n$, that is,

$$\{F_R(I) > 0\} \in \bigcap_{N \geq 1} \sigma\{Y_k(x); x \in \mathbb{R}^2, k \geq N\}.$$

As the processes $(Y_k)_k$ are independent, we can use the Kolmogorov 0–1 law to deduce that, \mathbb{P}_x^B -almost surely, the event $\{F_R(I) > 0\}$ has \mathbb{P}^X -probability 0 or 1. From Lemma 2.8, we have \mathbb{P}_x^B -almost surely

$$\mathbb{E}^X[F_R(I)] = t - s > 0.$$

Therefore, \mathbb{P}_x^B -almost surely, on the event $\{t < T_R\}$, the event $\{F_R(I) > 0\}$ has \mathbb{P}^X -probability 1. Then we can consider a countable family $(I_p)_p$ of intervals generating the Borel sigma algebra on $[0, T_R[$. \mathbb{P}_x^B -almost surely, we have $F_R(I_p) > 0$ for all p (and all $R > 0$). This shows that \mathbb{P}^X -a.s., F has full support \mathbb{P}_x^B -a.s., which equivalently means that the random mapping $t \mapsto F(t)$ is strictly increasing.

So far, we have only proved that, for each $x \in \mathbb{R}^2$, there is a measurable set $S_x \subset \Omega$ such that $\mathbb{P}^X(S_x) = 1$ and on S_x , F is strictly increasing under \mathbb{P}_x^B . Now we want to show that there is a measurable set $S \subset \Omega$ such that $\mathbb{P}^X(S) = 1$ and for all $x \in \mathbb{R}^2$, on S , F is strictly increasing under \mathbb{P}_x^B . Clearly, we can find a set S such that $\mathbb{P}^X(S) = 1$ for all $x \in \mathbb{Q}^2$ and F is increasing on S under \mathbb{P}_x^B . Now we explain a coupling procedure that will serve to complete the proof of item 3 as well as 4 and 5. Let us consider another Brownian motion W independent of B , X (even if it means enlarging the space Ω , we may assume that W is defined on the same probability space than the MFF X and the Brownian motion B). We denote by $\mathbb{P}_{x,y}^{B,W}$ the probability measure $\mathbb{P}_x^B \otimes \mathbb{P}_y^W$. We state the following coupling lemma, the proof of which is rather elementary and thus left to the reader.

LEMMA 2.9. *Let us denote by τ_1 the first time at which the first components of B and W coincide and by τ_2 the first time at which the second components coincide after τ_1 :*

$$\tau_1 = \inf\{u > 0; B_u^1 = W_u^1\}, \quad \tau_2 = \inf\{u > \tau_1; B_u^2 = W_u^2\}.$$

Under $\mathbb{P}_{x,y}^{B,W}$, the random process \bar{B} defined by

$$\bar{B}_t = \begin{cases} (W_t^1, W_t^2), & \text{if } t \leq \tau_1, \\ (B_t^1, W_t^2), & \text{if } \tau_1 < t \leq \tau_2, \\ (B_t^1, B_t^2), & \text{if } \tau_2 < t, \end{cases}$$

is a Brownian motion on \mathbb{R}^2 starting from y , and coincides with W for all times $t > \tau_2$. Furthermore, we have

$$\forall \eta > 0, \quad \lim_{\delta \rightarrow 0} \sup_{x,y \in \mathbb{R}^2; |x-y| \leq \delta} \mathbb{P}_{x,y}^{B,X}(\tau_2 > \eta) \rightarrow 0 \quad \text{and} \quad \mathbb{P}_{x,y}^{B,X}(\tau_2 < \infty) = 1.$$

We can associate \mathbb{P}^X -a.s. to the Brownian motion \bar{B} a PCAF, denoted by $F(\bar{B}, t)$ to distinguish it from F , with Revuz measure M as prescribed in the beginning of the proof of Theorem 2.7. It is also plain to check that \mathbb{P}^X -a.s., for all $x, y \in \mathbb{R}^2$, under $\mathbb{P}_{x,y}^{B,W}$, the marginal laws of (B, F) and $(\bar{B}, F(\bar{B}, \cdot))$, respectively, coincide with the law of (B, F) under \mathbb{P}_x^B and \mathbb{P}_y^B .

Therefore, on S and for $y \in \mathbb{Q}^2$, $\mathbb{P}_{x,y}^{B,W}$ a.s., the PCAF $F(\bar{B}, \cdot)$ is strictly increasing on \mathbb{R}_+ . Furthermore, the coupling procedure (Lemma 2.9) also entails that the mappings $s \in [\tau_2, +\infty[\mapsto F(s) - F(\tau_2)$ and $s \in [\tau_2, +\infty[\mapsto F(\bar{B}, s) - F(\bar{B}, \tau_2)$ are equal $\mathbb{P}_{x,y}^{B,W}$ a.s. Therefore, for any $x \in \mathbb{R}^2$ and $y \in \mathbb{Q}^2$, the above discussion shows that $F(s)$ is strictly increasing for $s > \tau_2$. If y is chosen arbitrarily close to x , Lemma 2.9 shows that $\tau_2 \rightarrow 0$ in probability. We deduce that F is strictly increasing on \mathbb{R}_+ . This completes the proof of item 3 as we have shown that on S for all $x \in \mathbb{R}^2$, F is strictly increasing \mathbb{P}_x^B a.s.

We now prove item 4. Once again, the coupling procedure (Lemma 2.9) shows that it is enough to prove item 4 \mathbb{P}^X a.s. for only one $x \in \mathbb{R}^2$: because $\mathbb{P}_{x,y}^{B,W}(\tau_2 < \infty) = 1$ for all x, y , it is plain to deduce that $\mathbb{P}_x^B(F(t) \rightarrow \infty \text{ as } t \rightarrow \infty) = \mathbb{P}_y^B(F(t) \rightarrow \infty \text{ as } t \rightarrow \infty)$ for all $y \in \mathbb{R}^2$. So we work under \mathbb{P}_0^B .

We consider the following sequence of stopping times associated to the Brownian motion:

$$T_n = \inf\{t > 0, |B_t| = 2n\}, \quad \bar{T}_n = \inf\{t > T_n, |B_t - B_{T_n}| = \frac{1}{4}\}.$$

We also consider an increasing sequence of integers $(n_j)_{j \geq 1}$ such that the following property holds for all $l \leq k$:

$$(2.16) \quad \sum_{l \leq j < j' \leq k} \alpha_{j,j'} \leq k - l + 1,$$

with $\alpha_{j,j'} = \sup_{|x| \leq 2n_j + 1/4, |y| \geq 2n_{j'} - 1/4} G_m(x, y).$

Such a sequence exists because G_m defined by (2.1) satisfies $G_m(x, y) \leq ce^{-c|x-y|}$ for $|x - y| \geq 1$ and some constant $c > 0$. Recall that we identify F and its associated measure. From the Markov inequality and Fatou’s lemma, we obtain

$$\begin{aligned}
 & \mathbb{P}_0\left(\bigcap_{l \leq j \leq k} \{F(]T_{n_j}, \bar{T}_{n_j}] \leq c\}\right) \\
 & \leq c^{k-l+1} \mathbb{E}_0\left[\prod_{l \leq j \leq k} (F(]T_{n_j}, \bar{T}_{n_j}])^{-1}\right] \\
 (2.17) \quad & \leq \liminf_{n \rightarrow \infty} c^{k-l+1} \mathbb{E}_0\left[\left(\int_{\prod_{l \leq j \leq k}]T_{n_j}, \bar{T}_{n_j}] c_n^{-(k-l+1)(\gamma^2/2)} \right. \right. \\
 & \quad \left. \left. \times e^{\gamma(X_n(B_{s_l}) + \dots + X_n(B_{s_k}))} ds_l \dots ds_k\right)^{-1}\right].
 \end{aligned}$$

We want to get rid of the long range correlations of the MFF X . To this purpose, we introduce a log-correlated random distribution \bar{X} with covariance kernel $\mathbb{E}[\bar{X}(x)\bar{X}(y)] = \ln_+ \frac{1}{|y-x|}$. It is a kernel of σ -positive type [35], Proposition 2.15. We can find an approximation family “à la Kahane” $(\bar{X}_n)_n$ of \bar{X} such that $\mathbb{E}[\bar{X}_n(x)\bar{X}_n(y)] \leq \ln_+ \frac{1}{|y-x|}$ and

$$(2.18) \quad \mathbb{E}[\bar{X}_n(x)\bar{X}_n(y)] - D \leq \mathbb{E}[X_n(x)X_n(y)] \leq \mathbb{E}[\bar{X}_n(x)\bar{X}_n(y)] + D$$

for some constant D which does not depend on relevant quantities (in particular not on n , e.g., [35], proof of Proposition 2.15). By using this relation and (2.16), we get for all $(x_l, \dots, x_k), (y_l, \dots, y_k) \in \prod_{l \leq j \leq k}]T_{n_j}, \bar{T}_{n_j}]$

$$\begin{aligned}
 & \mathbb{E}^X\left[\left(\sum_{j=l}^k X_n(x_j)\right)\left(\sum_{j=l}^k X_n(y_j)\right)\right] \\
 & = \sum_{j=l}^k \mathbb{E}^X[X_n(x_j)X_n(y_j)] + \sum_{j, j'=l, j \neq j'}^k \mathbb{E}^X[X_n(x_j)X_n(y_{j'})] \\
 (2.19) \quad & \leq (k-l+1)D + \sum_{j=l}^k \mathbb{E}^X[\bar{X}_n(x_j)\bar{X}_n(y_j)] + 2 \sum_{l \leq j < j' \leq k} \alpha_{j, j'} \\
 & \leq (k-l+1)(D+2) + \mathbb{E}^{\bar{X}}\left[\left(\sum_{j=l}^k \bar{X}_n(x_j)\right)\left(\sum_{j=l}^k \bar{X}_n(y_j)\right)\right].
 \end{aligned}$$

In the last line, we have used the fact that $\mathbb{E}^{\bar{X}}[\bar{X}_n(x_j)\bar{X}_n(y_{j'})] = 0$ if $j \neq j'$.

Now we want to apply Lemma A.1 with $Y(x_l, \dots, x_k) = \sum_{j=l}^k X_n(x_j)$ (with kernel K) and $Y'(x_l, \dots, x_k) = \sum_{j=l}^k \bar{X}_n(x_j)$ (with kernel K'), $\nu(dx_l, \dots, dx_k) =$

$\mu_l(dx_l) \times \dots \times \mu_k(dx_k)$ where each μ_l stands for the occupation measure of the Brownian motion between the times T_{n_j} and \bar{T}_{n_j} and the convex function $x \mapsto 1/x$ (in fact, this function is discontinuous at 0 but this is not a problem: truncate it in order to have a continuous convex function, apply Kahane’s inequality and then remove the truncation). We have shown above that $K \leq K' + C$ with $C = (k - l + 1)(D + 2)$. The last point is that the exponential term in the expectation (2.17) is not renormalized by the variance. However, the above inequality shows that

$$\mathbb{E}^X \left[\left(\sum_{j=l}^k X_n(x_j) \right)^2 \right] \leq \sum_{j=l}^k \mathbb{E}^X [X_n(x_j)^2] + 2D(k - l + 1)$$

in such a way that $(k - l + 1) \ln c_n \leq \mathbb{E}^X [(\sum_{j=l}^k X_n(x_j))^2] \leq (k - l + 1) \ln c_n + 2D(k - l + 1)$.

Hence, even if it means multiplying the constant c by a deterministic constant that does not depend on k, l , we can replace the term $c_n^{-(k-l+1)(\gamma^2/2)}$ in (2.17) by $\exp(-\frac{\gamma^2}{2} \mathbb{E}^X [(\sum_{j=l}^k X_n(x_j))^2])$. We are then in position to apply Lemma A.1 item 2, which tells us that we can replace X_n in (2.19) by \bar{X}_n at the cost of replacing the constant c by another constant c' , which still does not depend on n (only on D, γ). The main advantage of this procedure is that we deal now with a field \bar{X}_n that possesses strong decorrelation properties: if, for a set $A \subset \mathbb{R}^2$, we denote by $\mathcal{F}_{n,A}$ the sigma algebra generated by the random variables $\{\bar{X}_n(x), x \in A\}$ then $\mathcal{F}_{n,A}$ is independent of $\mathcal{F}_{n,B}$ as soon as $\text{dist}(A, B) > 1$.

In what follows, we still stick to the notation \mathbb{E}_0 to denote expectation with respect to the probability measure $\mathbb{P}^{\bar{X}} \otimes \mathbb{P}_0^B$ and $\bar{c}_n = \mathbb{E}^{\bar{X}} [\bar{X}_n(x)^2]$, which does not depend on x by stationarity. We have by using the strong Markov property of the Brownian motion and the fact that \bar{X}_n is decorrelated at distance 1

$$\begin{aligned} & \mathbb{P}_0 \left(\bigcap_{l \leq j \leq k} \{F(]T_{n_j}, \bar{T}_{n_j}[) \leq c\} \right) \\ & \leq \liminf_{n \rightarrow \infty} (c')^{k-l+1} \mathbb{E}_0 \left[\left(\int_{\prod_{l \leq j \leq k}]T_{n_j}, \bar{T}_{n_j}[} \bar{c}_n^{-(k-l+1)(\gamma^2/2)} \right. \right. \\ & \quad \left. \left. \times e^{\gamma(\bar{X}_n(B_{s_l}) + \dots + \bar{X}_n(B_{s_k}))} ds_l \dots ds_k \right)^{-1} \right] \\ & = \liminf_{n \rightarrow \infty} (c')^{k-l+1} \mathbb{E}_0 \left[\left(\int_{]T_{n_1}, \bar{T}_{n_1}[} \bar{c}_n^{-\gamma^2/2} e^{\gamma \bar{X}_n(B_s)} ds \right)^{-1} \right]^{k-l+1}. \end{aligned}$$

Notice that $n \mapsto \int_{]T_{n_1}, \bar{T}_{n_1}[} \bar{c}_n^{-\gamma^2/2} e^{\gamma \bar{X}_n(B_s)} ds$ is a uniformly integrable martingale and converges toward a random variable denoted by

$$\int_{]T_{n_1}, \bar{T}_{n_1}[} e^{\gamma \bar{X}(B_s) - (\gamma^2/2) \mathbb{E}_0[\bar{X}^2(B_s)]} ds$$

(the proof is identical to Lemma 2.8 and the notation is due to the fact that the limit is a Gaussian multiplicative chaos). Therefore, the Jensen inequality leads to

$$\begin{aligned} & \mathbb{P}_0 \left(\bigcap_{l \leq j \leq k} \{F(I_{T_{n_j}}, \bar{T}_{n_j}) \leq c\} \right) \\ & \leq \left(c' \mathbb{E}_0 \left[\left(\int_{I_{T_{n_1}, \bar{T}_{n_1}}} e^{\gamma \bar{X}(B_s) - (\gamma^2/2) \mathbb{E}_0[\bar{X}^2(B_s)]} ds \right)^{-1} \right] \right)^{k-l+1}. \end{aligned}$$

Let us admit for a while that the above expectation in the right-hand side is finite (this will be proved below in Lemma 2.12). This inequality shows that we can choose c small enough such that

$$\mathbb{P}_0 \left(\bigcap_{l \leq j < \infty} \{F(I_{T_{n_j}}, \bar{T}_{n_j}) \leq c\} \right) = 0.$$

Thus, we get

$$\mathbb{P}_0 \left(\bigcap_{l \geq 1} \bigcup_{l \leq j < \infty} \{F(I_{T_{n_j}}, \bar{T}_{n_j}) > c\} \right) = 1.$$

Since $\lim_{t \rightarrow \infty} F(t) \geq c \sum_{j \geq 1} \mathbb{1}_{\{F(I_{T_{n_j}}, \bar{T}_{n_j}) > c\}}$, the proof of item 4 is complete.

It remains to prove item 5. For $y \in \mathbb{R}^2$, we denote by M^y the shifted measure $M^y(A) = M(A + y)$, and F^y its associated PCAF. We claim that it is enough to prove that, \mathbb{P}^x -a.s. for all $x \in \mathbb{R}^2$ and $\eta > 0$

$$(2.20) \quad \lim_{y \rightarrow x} \mathbb{P}_0^B \left(\sup_{t \leq T} |F^y(t) - F^x(t)| \geq \eta \right) = 0.$$

Indeed, if (2.20) is true, then for any uniformly continuous function G (with modulus m and bounded by K) on $C(\mathbb{R}_+; \mathbb{R}^2 \times \mathbb{R}_+)$, we have for all $\eta > 0$

$$\begin{aligned} & |\mathbb{E}_x^B[G(B, F)] - \mathbb{E}_y^B[G(B, F)]| \\ & = |\mathbb{E}_0^B[G(x + B, F^x)] - \mathbb{E}_0^B[G(y + B, F^y)]| \\ & \leq K \mathbb{1}_{\{|x-y| > \eta\}} + K \mathbb{P}_0^B \left(\sup_{t \leq T} |F^x(t) - F^y(t)| > \eta \right) + m(\eta). \end{aligned}$$

We can then pass to the limit as $y \rightarrow x$ and use (2.20) to prove that the first two terms go to 0 and then choose η arbitrarily small to conclude.

To establish (2.20), for all $R > 0$, we use Lemma 2.5 to get

$$\begin{aligned} & \mathbb{P}_0^B \left(\sup_{t \leq T} |F^y(t) - F^x(t)| \geq \eta \right) \\ & = \mathbb{P}_0^B \left(\sup_{t \leq 0} |F^y(t \wedge T_R) - F^x(t \wedge T_R)| \geq \eta \right) + \mathbb{P}_0^B(T_R \leq T) \\ & \leq c_R \exp \left(- \frac{\eta}{c_R \sqrt{d_R(M^y, M^x)}} \right) + \mathbb{1}_{\{d_R(M^y, M^x) \geq 1\}} + \mathbb{P}_0^B(T_R \leq T). \end{aligned}$$

Since $d_R(M^y, M^x) \rightarrow 0$ as $y \rightarrow x$ (cf. item 4 of Proposition 2.3), we deduce

$$\limsup_{y \rightarrow x} \mathbb{P}_0^B \left(\sup_{t \leq T} |F^y(t) - F^x(t)| \geq \eta \right) \leq \mathbb{P}_0^B(T_R \leq T).$$

We complete the proof by letting $R \rightarrow \infty$. \square

PROOF OF LEMMA 2.8. \mathbb{P}_x^B -a.s., the family $(F_R^n(t))_n$ is a nonnegative martingale w.r.t. the filtration $(\mathcal{F}_n)_n$ and, therefore, converges almost surely as $n \rightarrow \infty$.

Let us prove that it is uniformly integrable. Denote by ν the occupation measure of the Brownian motion B between 0 and $t \wedge T_R$. Observe that

$$\int_{\mathbb{R}^2} c_n^{-\gamma^2/2} e^{\gamma X_n(z)} \nu(dz) = F_R^n(t).$$

From [27] (see also [35]), we just have to prove that \mathbb{P}_x^B -a.s.

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{|z - z'|^\alpha} \nu(dz) \nu(dz') < +\infty$$

for some $\alpha < 2$. This statement is elementary (just compute the expectation), and thus left to the reader (much stronger statements are discussed in [17], Section 10, e.g.). \square

2.5. Study of the moments and power law spectrum. In this section, we investigate the finiteness of the moments of the PCAF. We will say that F possesses moments of order q if we have $\mathbb{E}_x[F(t)^q] < +\infty$ for all $t > 0$ and $x \in \mathbb{R}^2$.

THEOREM 2.10 (Positive moments and power law spectrum). (1) If $\gamma < 2$, the mapping F possesses moments of order q for $0 \leq q < 4/\gamma^2$. (2) If F admits moments of order $q \geq 1$ then, for all $s \in [0, 1]$ and $t \in [0, T]$:

$$\mathbb{E}_x[(F(t+s) - F(t))^q] \leq C_q s^{\xi(q)},$$

where

$$\xi(q) = \left(1 + \frac{\gamma^2}{4} \right) q - \frac{\gamma^2}{4} q^2$$

and $C_q > 0$ is some constant independent of x, T .

PROOF. There is here no exception to the rule in multiplicative chaos theory that studying finiteness of the moments is technically heavy. So, the entire Appendix B is devoted to the proof of item 1.

Now we assume that $\gamma < 2$ and that F possesses moments of order $q \geq 1$. We prove the estimate concerning the power law spectrum. By stationarity, we may

assume that $x = 0$. We first prove it when $t = 0$ and then we deduce the uniform estimate in t . Under \mathbb{P}_0 , we have

$$\begin{aligned}
 (2.21) \quad F(s) &= \int_0^s e^{\gamma X(B_r) - (\gamma^2/2)\mathbb{E}[X(B_r)^2]} dr \\
 &= s \int_0^1 e^{\gamma X(B_{us}) - (\gamma^2/2)\mathbb{E}[X(B_{us})^2]} du \\
 &\stackrel{\text{law}}{=} s \int_0^1 e^{\gamma X(\sqrt{s}B_u) - (\gamma^2/2)\mathbb{E}[X(\sqrt{s}B_u)^2]} du.
 \end{aligned}$$

Let us stress here that the above computations are of course only formal as all the quantities are understood as limits: yet, the final statement is correct as can be seen by applying the argument to the same regularized quantities and by passing to the limit to remove the cutoff. Then, from (2.4), we have $G_m(\sqrt{s}u, \sqrt{s}v) \leq \ln \frac{1}{\sqrt{s}} + G_m(u, v)$ for all $u, v \in \mathbb{R}^2$. Then, by taking the q th power and expectation in (2.21) and Kahane’s convexity inequalities (see Lemma A.1 item 2), we get for some Gaussian random variable Ω_s with mean 0 and variance $-\frac{1}{2} \ln s$ and independent of $\int_0^t e^{\gamma X(B_u) - (\gamma^2/2)\mathbb{E}[X(B_u)^2]} du$

$$\begin{aligned}
 \mathbb{E}_0[F(s)^q] &\leq s^q \mathbb{E}_0 \left[\left(e^{\gamma \Omega_s + (\gamma^2/4) \ln s} \int_0^1 e^{\gamma X(B_u) - (\gamma^2/2)\mathbb{E}[X(B_u)^2]} du \right)^q \right] \\
 &= s^q \mathbb{E}_0 [e^{q\gamma \Omega_s + q(\gamma^2/4) \ln s}] \mathbb{E}_0 \left[\left(\int_0^1 e^{\gamma X(B_u) - (\gamma^2/2)\mathbb{E}[X(B_u)^2]} du \right)^q \right] \\
 &= C_q s^{\xi(q)},
 \end{aligned}$$

where $C_q = \mathbb{E}_0[(\int_0^1 e^{\gamma X(B_u) - (\gamma^2/2)\mathbb{E}[X(B_u)^2]} du)^q]$ is independent of s, x .

Now we treat the general case $t \neq 0$. By using the Markov property of the Brownian motion and the stationarity of X , it is readily seen that

$$\mathbb{E}_x[(F(t + s) - F(t))^q] = \mathbb{E}_0[F(s)^q]. \quad \square$$

COROLLARY 2.11. *Set $\alpha = (1 - \frac{\gamma}{2})^2$. For each $T > 0$ and $\epsilon > 0$, there exists a random constant $C > 0$ such that \mathbb{P}_x a.s.*

$$\sup_{0 \leq s < t \leq T} |F(t) - F(s)| \leq C |t - s|^{\alpha - \epsilon}.$$

PROOF. Similar to Theorem 2.2 and thus left to the reader. \square

Now we investigate finiteness of moments of negative order. Denote by T_r^x the first exit time of the Brownian motion B out of the disk $B(x, r)$ for $r \in]0, 1]$.

PROPOSITION 2.12. *For all $q > 0$, there exists some constant $C_q > 0$ (depending on q) such that for all $x \in \mathbb{R}^2$ and $r \in [0, 1]$*

$$(2.22) \quad \sup_{n \geq 0} \mathbb{E}_x \left[\left(c_n^{-\gamma^2/2} \int_0^{T_r^x} e^{\gamma X_n(B_s)} ds \right)^{-q} \right] \leq C_q r^{2\xi(-q)}.$$

PROOF. Without loss of generality, we can take $x = 0$ by stationarity of the field X . Furthermore, from Kahane’s convexity inequalities Lemma A.1, it suffices to prove the result for one log-correlated Gaussian field with a kernel of σ -positive type. Let us choose the exact scale invariant field \bar{X} with covariance kernel given by

$$\mathbb{E}[\bar{X}(x)\bar{X}(y)] = \ln_+ \frac{2}{|x - y|}$$

with white noise decomposition $(\bar{X}_\epsilon)_{\epsilon \in]0,1]}$ of \bar{X} as constructed in [37]. More precisely, the correlation structure of $(\bar{X}_\epsilon)_{\epsilon \in]0,1]}$ is given for $\epsilon, \epsilon' \in]0, 1]$ by

$$\begin{aligned} & \mathbb{E}[\bar{X}_\epsilon(x)\bar{X}_{\epsilon'}(y)] \\ &= \begin{cases} 0, & \text{if } |x - y| > 2, \\ \ln \frac{2}{|x - y|}, & \text{if } \max(\epsilon, \epsilon') \leq |x - y| \leq 2, \\ \ln \frac{2}{\max(\epsilon, \epsilon')} + 2 \left(1 - \frac{|x - y|^{1/2}}{\max(\epsilon, \epsilon')^{1/2}} \right), & \text{if } |y - x| \leq \max(\epsilon, \epsilon'). \end{cases} \end{aligned}$$

This covariance structure entails some interesting properties that we detail now. The process $\epsilon \rightarrow \bar{X}_\epsilon$ has independent increments, meaning that for $\epsilon' > \epsilon$ the field $\bar{X}_\epsilon - \bar{X}_{\epsilon'}$ is independent of the sigma algebra $\sigma\{X_u(z); z \in \mathbb{R}^2, u \geq \epsilon'\}$. For $\epsilon' > \epsilon$, the field $\bar{X}_{\epsilon, \epsilon'} := \bar{X}_\epsilon - \bar{X}_{\epsilon'}$ has a correlation cutoff of length ϵ' , meaning that the fields $(\bar{X}_{\epsilon, \epsilon'}(x))_{x \in A}$ and $(\bar{X}_{\epsilon, \epsilon'}(x))_{x \in B}$ are independent whenever the Euclidean distance between the two sets A, B is greater than ϵ' . Finally, we have the following relation in law for $r, \epsilon \leq 1$:

$$(2.23) \quad (X_{r\epsilon}(rx))_{|x| \leq 1} \stackrel{\text{law}}{=} (\Omega_r + X_\epsilon(x))_{|x| \leq 1},$$

where Ω_r is a Gaussian distribution $\mathcal{N}(0, \ln \frac{1}{r})$ and is independent of $(\bar{X}_\epsilon)_\epsilon$. In what follows, we stick to the Notation 2.1 with \bar{X} instead of X . Finally, we just write T_r for T_r^0 .

So, we have to prove

$$(2.24) \quad \sup_{\epsilon \in]0,1]} \mathbb{E}_0 \left[\left(2^{-\gamma^2/2} \epsilon^{\gamma^2/2} \int_0^{T_r} e^{\gamma \bar{X}_\epsilon(B_s)} ds \right)^{-q} \right] \leq C_q r^{2\xi(-q)}.$$

Notice that the supremum is reached for $\epsilon \rightarrow 0$ by the martingale property and the Jensen inequality. Now, if $\tilde{T}_{1/4}$ is the first time the Brownian motion $(B_{t+T_{3/4}} -$

$B_{T_{3/4}})_{t \geq 0}$ hits the disk of radius $\frac{1}{4}$, we get for $\epsilon < 1/4$

$$\begin{aligned} & 2^{-\gamma^2/2} \epsilon^{\gamma^2/2} \int_0^{T_1} e^{\gamma \bar{X}_\epsilon(B_s)} ds \\ & \geq 2^{-\gamma^2/2} \epsilon^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_\epsilon(B_s)} ds + 2^{-\gamma^2/2} \epsilon^{\gamma^2/2} \int_0^{\tilde{T}_{1/4}} e^{\gamma \bar{X}_\epsilon(B_{s+T_{3/4}})} ds \\ & \geq 8^{-\gamma^2/2} e^{\gamma \inf_{|x| \leq 1} \bar{X}_{1/4}(x)} \left((4\epsilon)^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_s)} ds \right. \\ & \quad \left. + (4\epsilon)^{\gamma^2/2} \int_0^{\tilde{T}_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_{s+T_{3/4}})} ds \right). \end{aligned}$$

The main observation is that, under the annealed measure \mathbb{E}_0 , the above two integrals are independent identically distributed random variables. Indeed, by considering two bounded continuous functionals F, G , we get

$$\begin{aligned} P(F, G) & := \mathbb{E}_0 \left[F \left((4\epsilon)^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_s)} ds \right) \right. \\ & \quad \left. \times G \left((4\epsilon)^{\gamma^2/2} \int_0^{\tilde{T}_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_{s+T_{3/4}})} ds \right) \right] \\ & = \mathbb{E}_0^B \left[\mathbb{E}^{\bar{X}} \left[F \left((4\epsilon)^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_s)} ds \right) \right] \right. \\ & \quad \left. \times \mathbb{E}^{\bar{X}} \left[G \left((4\epsilon)^{\gamma^2/2} \int_0^{\tilde{T}_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_{s+T_{3/4}})} ds \right) \right] \right], \end{aligned}$$

where we have used the fact that $\bar{X}_{\epsilon,1/4}$ has a correlation cutoff of length $1/4$. If we use now the stationarity of the field $\bar{X}_{\epsilon,1/4}$ and the independence of the increments of the standard Brownian motion, we obtain

$$\begin{aligned} P(F, G) & = \mathbb{E}_0^B \left[\mathbb{E}^{\bar{X}} \left[F \left((4\epsilon)^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_s)} ds \right) \right] \right. \\ & \quad \left. \times \mathbb{E}^{\bar{X}} \left[G \left((4\epsilon)^{\gamma^2/2} \int_0^{\tilde{T}_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_{s+T_{3/4}-B_{T_{3/4}}})} ds \right) \right] \right] \\ & = \mathbb{E}_0 \left[F \left((4\epsilon)^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_s)} ds \right) \right] \\ & \quad \times \mathbb{E}_0 \left[G \left((4\epsilon)^{\gamma^2/2} \int_0^{\tilde{T}_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_{s+T_{3/4}-B_{T_{3/4}}})} ds \right) \right] \\ & = \mathbb{E}_0 \left[F \left((4\epsilon)^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_s)} ds \right) \right] \\ & \quad \times \mathbb{E}_0 \left[G \left((4\epsilon)^{\gamma^2/2} \int_0^{T_{1/4}} e^{\gamma \bar{X}_{\epsilon,1/4}(B_s)} ds \right) \right]. \end{aligned}$$

Furthermore, for all $r \in]0, 1]$, we have

$$\begin{aligned} (r\varepsilon/2)^{\gamma^2/2} \int_0^{T_r} e^{\gamma \bar{X}_{r\varepsilon}(B_s)} ds &= (r\varepsilon/2)^{\gamma^2/2} r^2 \int_0^{T_r/r^2} e^{\gamma \bar{X}_{r\varepsilon}(r(B_{r^2 s'/r}))} ds' \\ &= (r\varepsilon/2)^{\gamma^2/2} r^2 \int_0^{\tilde{T}_1} e^{\gamma \bar{X}_{r\varepsilon}(r\tilde{B}_{s'})} ds', \end{aligned}$$

where $\tilde{B}_{s'} = r^{-1} B_{r^2 s'}$ is a Brownian motion and $\tilde{T}_1 = \frac{T_r}{r^2}$ is the first time it hits the disk of radius 1. Therefore, from (2.23), we get the following scaling relation in distribution for all $r \in]0, 1]$ under the annealed measure \mathbb{P}_0 :

$$\begin{aligned} (2.25) \quad &(r\varepsilon/2)^{\gamma^2/2} \int_0^{T_r} e^{\gamma \bar{X}_{r\varepsilon}(B_s)} ds \\ &\stackrel{\text{law}}{=} r^2 e^{\gamma \Omega_r - \gamma^2/2 \ln(1/r)} (\varepsilon/2)^{\gamma^2/2} \int_0^{T_1} e^{\gamma \bar{X}_\varepsilon(B_s)} ds. \end{aligned}$$

From this scaling relation and the above considerations, we deduce that we can find some variable N with negative moments and such that we have the following stochastic domination:

$$Y \geq N(Y_1 + Y_2),$$

where (Y_1, Y_2) are i.i.d. of distribution Y , independent of N , $\mathbb{E}[N^{-q}] < +\infty$ (see [1], Theorem 2.1.1) and Y is distributed like $\lim_{\varepsilon \rightarrow 0} \varepsilon^{\gamma^2/2} \int_0^{T_1} e^{\gamma \bar{X}_\varepsilon(B_s)} ds$. We get (2.24) is true with $r = 1$ by adapting [30] (see also [20], Section B.4). Then one deduces inequality (2.24) from (2.25) for all r . \square

One can then deduce the following results.

COROLLARY 2.13. *For all $q > 0$, there exists some constant $C > 0$ (depending on q) such that for all $x \in \mathbb{R}^2$ and all $0 \leq s < t \leq 1$:*

$$(2.26) \quad \sup_{n \geq 0} \mathbb{E}_x \left[\left(c_n^{-\gamma^2/2} \int_s^t e^{\gamma X_n(B_r)} dr \right)^{-q} \right] \leq C(t-s)^\xi (-q).$$

PROOF. Without loss of generality, we can take $x = 0$ and $s = 0$ by stationarity of the field X and the strong Markov property of the Brownian motion. Then it suffices to show that the expectation in (2.26) is finite for $t = 1$. Indeed, one can then use the techniques in the proof of Proposition 2.10 in order to obtain the right-hand side of (2.26) for any $t \in [0, 1]$ ($x \mapsto x^{-q}$ is convex).

Recall that T_r denote the first exit times of the Brownian motion out of the ball $B(0, r)$. Recall that, by scaling and [40], Theorem 1.2 (page 93), we have the existence of some absolute constant $c > 0$ such that for all $t > 0$

$$(2.27) \quad \mathbb{P}_0^B(T_r \geq t) \leq \frac{ct}{r^2} e^{-(ct)/r^2}.$$

We have

$$F(1)^{-q} \leq \sum_{n=1}^{\infty} \mathbb{1}_{\{T_{1/2^n} < 1 \leq T_{1/2^{n-1}}\}} F(T_{1/2^n})^{-q}.$$

Therefore, we get by Proposition 2.12 and the bound (2.27)

$$\begin{aligned} \mathbb{E}_0[F(1)^{-q}] &\leq \sum_{n=1}^{\infty} \mathbb{E}_0[\mathbb{1}_{\{T_{1/2^n} < 1 \leq T_{1/2^{n-1}}\}} F(T_{1/2^n})^{-q}] \\ &\leq \sum_{n=1}^{\infty} \mathbb{P}_0^B(T_{1/2^{n-1}} \geq 1)^{1/2} \mathbb{E}_0[F(T_{1/2^n})^{-2q}]^{1/2} \\ &\leq C \sum_{n=1}^{\infty} 2^{2n} e^{-c2^{2n}} (2^n)^{2q + ((q(1+2q))/2)\gamma^2}. \end{aligned}$$

This latter series is obviously finite. \square

COROLLARY 2.14. *Set $\beta = (1 + \frac{\gamma}{2})^2$. For each $T > 0$ and $\epsilon > 0$, there exists a random constant $C > 0$ such that \mathbb{P}_x a.s.*

$$\forall 0 \leq s < t \leq T, \quad |F(t) - F(s)| \geq C|t - s|^{\beta + \epsilon}.$$

PROOF. We take $T = 1$ for simplicity. Now, we partition $[0, 1]$ into 2^n dyadic squares $(I_n^j)_{1 \leq j \leq 2^{2n}}$ of equal size. If $p > 0$, we get

$$\begin{aligned} \mathbb{P}_x \left(\inf_{1 \leq j \leq 2^n} F(I_n^j) \leq \frac{1}{2^{(\beta + \epsilon)n}} \right) &\leq 2^{-p(\beta + \epsilon)n} \mathbb{E}_x \left[\sum_{1 \leq j \leq 2^n} F(I_n^j)^{-p} \right] \\ &\leq C_p 2^{-p(\beta + \epsilon)n} 2^{(1 - \xi(-p))n}. \end{aligned}$$

We conclude as in the proof of Theorem 2.2 by choosing $p > 0$ such that $\xi(-p) - 1 + p(\beta + \epsilon) > 0$. \square

2.6. Definition of the Liouville Brownian motion. Before giving the explicit description of the Liouville Brownian motion, let us first motivate the forthcoming definitions. The reader may consult [26] for an introductory background on Brownian motion on manifolds. Recall that our purpose is to define the Brownian motion associated to the metric tensor formally written (using conventional notation in Riemann geometry) $e^{\gamma X(x)} dx^2$, where dx^2 stands for the standard Euclidean metric on \mathbb{R}^2 . Because of the obvious divergences of such a direct definition, it is natural to regularize the field X and to consider instead the Riemann metric tensor

$$(2.28) \quad g_n = c_n^{-\gamma^2/2} e^{\gamma X_n(x)} dx^2.$$

The renormalization sequence $(c_n^{-\gamma^2/2})_n$ appears here to regulate the divergences when $n \rightarrow \infty$.

The Riemann volume on the manifold (\mathbb{R}^2, g_n) is nothing but the measure M_n defined by (2.7). One can also associate to the Riemann manifold (\mathbb{R}^2, g_n) a Brownian motion \mathcal{B}^n . Following the standard construction, such a Brownian motion can be constructed as follows. Consider a standard Brownian motion Z on \mathbb{R}^2 and, for any $x \in \mathbb{R}^2$, consider the solution $\mathcal{Z}^{n,x}$ of the following stochastic differential equation:

$$(2.29) \quad \begin{cases} \mathcal{Z}_{\mathbf{t}=0}^{n,x} = x, \\ d\mathcal{Z}_{\mathbf{t}}^{n,x} = c_n^{\gamma^2/4} e^{-\gamma/2 X_n(\mathcal{Z}_{\mathbf{t}}^{n,x})} dZ_{\mathbf{t}}. \end{cases}$$

Notice that it is not clear for the time being that the solution of this SDE has an infinite lifetime. Actually, we will see that the solution of such a SDE does not converge in probability a $n \rightarrow \infty$. Yet, it is enough to investigate convergence in law in the space of continuous functions $C(\mathbb{R}_+, \mathbb{R}^2)$. By using the Dambis–Schwarz theorem, the solution of (2.29) has the same law as the following process.

DEFINITION 2.15. For any $n \geq 1$, we define

$$(2.30) \quad \mathcal{B}_{\mathbf{t}}^n = B_{\langle \mathcal{B}^n \rangle_{\mathbf{t}}},$$

where $(B_r)_{r \geq 0}$ is a planar Brownian motion, independent of the MFF X and where the quadratic variation $\langle \mathcal{B}^n \rangle$ of $\mathcal{B}^{n,x}$ is defined as follows:

$$(2.31) \quad \langle \mathcal{B}^n \rangle_{\mathbf{t}} := \inf \left\{ s \geq 0 : c_n^{-\gamma^2/2} \int_0^s e^{\gamma X_n(B_u)} du \geq \mathbf{t} \right\}.$$

We call this process *n-regularized Liouville Brownian motion* (*n-LBM* for short).

The infimum in (2.31) is necessarily finite. Indeed we have

$$c_n^{-\gamma^2/2} \int_0^s e^{\gamma X_n(B_u)} du \geq c_n^{-\gamma^2/2} e^{\min_{x \in B(0,1)} X_n(x)} \int_0^s \mathbb{1}_{\{B_r \in B(0,1)\}} dr$$

and the latter quantity tends to ∞ when $t \rightarrow \infty$ via recurrence arguments.

Observe that (2.31) amounts to saying that the increasing process $\langle \mathcal{B}^{n,x} \rangle : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is just the inverse function of the PCAF

$$(2.32) \quad F_n(t) = c_n^{-\gamma^2/2} \int_0^t e^{\gamma X_n(B_r)} dr$$

studied in Section 2.4. In particular, we can assert now that the solution of (2.29) has infinite lifetime because $F_n(t) \rightarrow \infty$ as $t \rightarrow \infty$ (adapt, e.g., the argument of item 4 in Theorem 2.7). Several standard facts can be deduced from the smoothness of X_n .

PROPOSITION 2.16. *Let $n \geq 1$ be fixed. \mathbb{P}^X -a.s., the n -regularized Liouville Brownian motion \mathcal{B}^n induces a Feller diffusion on \mathbb{R}^2 . Let us denote by $(P_t^{\gamma,n})_{t \geq 0}$ its semigroup. Also, \mathcal{B}^n is reversible with respect to the Riemann volume M_n , which is therefore invariant for \mathcal{B}^n .*

For a proof of this proposition, the reader may consult [22], for instance, though the Feller property is not discussed but this is easy (for the same reason as the Liouville Brownian motion is Feller; see item 1 of Proposition 2.19 below).

As we have seen that the PCAFs $(F_n)_n$ converge toward F , it is natural to introduce the following.

DEFINITION 2.17. Assume $\gamma < 2$. \mathbb{P}^X almost surely, we define the *Liouville Brownian motion* (LBM for short) as the time changed Brownian motion

$$\mathcal{B}_t = B_{F^{-1}(t)},$$

where F is the PCAF of Theorem 2.7.

Observe that the continuity of F makes sure that the LBM does not get stuck in some area of the state space \mathbb{R}^2 . Typically, this situation may happen over areas where the field X takes large values, therefore, having as consequence to slow down the LBM. Furthermore, strict monotonicity of F ensures that the LBM possesses no jumps and the fact that F tends to ∞ as $t \rightarrow \infty$ makes sure that the LBM has an infinite lifetime, it does not “reach ∞ ” in finite time.

2.7. *Main properties of the LBM.* In this subsection, we collect a few important properties of the LBM that can be deduced from our previous analysis. The reader is referred to [22] for the classical terminology on Dirichlet forms.

THEOREM 2.18. *For $\gamma \in [0, 2[$, \mathbb{P}^X -a.s., the LBM is a strong Markov process with continuous sample paths, that is a diffusion process. If we denote by $(P_t^{\gamma,X})_{t \geq 0}$ the associated semigroup on $L^2(\mathbb{R}^2, M)$ then:*

1. $(P_t^{\gamma,X})_{t \geq 0}$ is strongly continuous on $L^2(\mathbb{R}^2, M)$ and symmetric.
2. the associated Dirichlet form is strongly local, regular and possesses $C_c^\infty(\mathbb{R}^2)$ as special core.

Observe that the semigroup $(P_t^{\gamma,X})_{t \geq 0}$ is itself random as it depends on the randomness of the MFF X . Furthermore, the above theorem entails that we can classically extend the semigroup $(P_t^{\gamma,X})_{t \geq 0}$ to a strongly-continuous semi-group on $L^p(\mathbb{R}^2, M)$ for all $1 \leq p < \infty$.

PROOF OF THEOREM 2.18. From [22], Theorem 6.1.1, the LBM is a Hunt process, therefore strong Markov. Item 1 and 2 then result from [22], Theorem 6.2.1. Continuity of sample paths results from the fact that F is strictly increasing (Theorem 2.7 item 3), in which case F^{-1} is continuous. \square

PROPOSITION 2.19. For $\gamma \in [0, 2[, \mathbb{P}^X$ -a.s.:

1. the law in $C(\mathbb{R}_+)$ of the LBM \mathcal{B} under \mathbb{P}_x^B is continuous with respect to x .
2. for all $x \in \mathbb{R}^2$, under \mathbb{P}_x^B the n -LBM $(\mathcal{B}^n)_n$ converges in law in $C(\mathbb{R}_+)$ equipped with the topology of uniform convergence over compact sets toward \mathcal{B} ,
3. the semigroup $(P_t^{\gamma, X})_{t \geq 0}$ is the limit of the semigroups $(P_t^{\gamma, n})_n$ as $n \rightarrow \infty$ in the sense that

$$\forall f \in C_b(\mathbb{R}^2), \quad \lim_{n \rightarrow \infty} P_t^{\gamma, n} f(x) = P_t^{\gamma, X} f(x),$$

4. for all $x \in \mathbb{R}^2$ and for all $z \in \mathbb{R}^2, \mathbb{P}_x^B(\liminf_{t \rightarrow \infty} |\mathcal{B}_t - z| = 0) = 1,$
5. for all $x \in \mathbb{R}^2, \mathbb{P}_x^B(\limsup_{t \rightarrow \infty} |\mathcal{B}_t| = \infty) = 1.$

PROOF. Let us prove item 1. Observe that the mapping

$$(w, v) \in C(\mathbb{R}_+, \mathbb{R}^2) \times C(\mathbb{R}_+, \mathbb{R}_+) \mapsto w \circ v^{-1},$$

where $v^{-1}(t) = \inf\{s \geq 0; v(s) > t\}$ is continuous at all these pairs (w, v) such that v is strictly increasing and $\lim_{s \rightarrow \infty} v(s) = +\infty$. Therefore, item 1 results from Theorem 2.7 items 3 + 4 + 5.

The same argument and Theorem 2.7 items 2 + 3 + 4 prove item 2. Item 3 is a direct consequence of item 2. Items 4 and 5 results from the fact a standard Brownian motion satisfies items 4 + 5 and the fact that \mathcal{B} is a time changed Brownian motion, with a continuous time change that goes to ∞ as $t \rightarrow \infty$. \square

COROLLARY 2.20. For $\gamma \in [0, 2[, \mathbb{P}^X$ -a.s., the semigroup $(P_t^{\gamma, X})_{t \geq 0}$ is Feller.

For $\gamma < 2$, we define the Liouville Laplacian Δ_X as the generator of the Liouville Brownian motion times the usual extra factor $\sqrt{2}$. The Liouville Laplacian corresponds to an operator which can formally be written as

$$\Delta_X = e^{-\gamma X(x)} \Delta$$

and can be thought of as the Laplace–Beltrami operator of $2d$ -Liouville quantum gravity.

2.8. *Asymptotic independence of the Liouville Brownian motion and the Euclidean Brownian motion.* Recall the SDE (2.29) with solution $\mathcal{Z}^{n, x}$, which is measurable with respect to the planar Brownian motion Z . In this section, we prove that $(\mathcal{Z}^{n, x})_n$ does not converge in probability as $n \rightarrow \infty$. This will show that the time change representation of the LBM is more relevant.

THEOREM 2.21. If $\gamma < 2, \mathbb{P}^X$ -a.s. and for all $x \in \mathbb{R}^2$, the pair of processes $(Z, \mathcal{Z}^{n, x})_n$ converges in law toward a couple (Z, \mathcal{Z}^x) . The planar Brownian motion Z and the LBM \mathcal{Z}^x are independent.

The above theorem shows that some extra randomness is created by taking the limit $n \rightarrow \infty$. Indeed, the n -LBM is a measurable function of the planar Brownian motion Z . Yet, Liouville/Euclidean Brownian motions are independent at the limit, showing that convergence in probability cannot hold.

PROOF OF THEOREM 2.21. Before beginning the proof, let us first clarify a few points. The n -LBM (2.29) involves the planar Brownian motion Z . An equivalent definition in law of this n -LBM is given in Definition 2.15 by means of another Brownian motion B , constructed via the Dambis–Schwarz theorem. As such, it implicitly depends on n as well as Z . It is therefore relevant to write explicitly this dependence in this proof. So we will write B^n instead of B .

We begin with writing explicitly the dependence between Z and B^n . The Dambis–Schwarz theorem tells us that [recall (2.32)]

$$B_t^n = c_n^{\gamma^2/4} \int_0^{F_n(t)} e^{-\gamma/2 X_n(\mathcal{Z}_u^{n,x})} dZ_u = \mathcal{Z}_{F_n(t,x)}^{n,x}.$$

Now we prove the asymptotic independence of Z and B^n . Let us compute their predictable bracket by making the change of variables:

$$\begin{aligned} \langle B^n, Z \rangle_t &= c_n^{\gamma^2/4} \int_0^t e^{-(\gamma/2) X_n(\mathcal{Z}_r^{n,x})} dr \\ &= c_n^{\gamma^2/4} \int_0^t e^{-(\gamma/2) X_n(B_{F_n^{-1}(r)}^n)} dr \\ &= c_n^{-\gamma^2/4} \int_0^{F_n(t)} e^{(\gamma/2) X_n(B_u^n)} du \\ &= c_n^{-\gamma^2/8} \times c_n^{-\gamma^2/8} \int_0^{F_n(t)} e^{(\gamma/2) X_n(B_u)} du. \end{aligned}$$

From Theorem 2.7, the above expression corresponds to multiplying a tight family by a factor $c_n^{-\gamma^2/8}$ that goes to 0 as $n \rightarrow \infty$. Therefore, \mathbb{P}^X -a.s., the family $(\langle B^n, Z \rangle)_n$ (as random functions of t) converges \mathbb{P}^Z -a.s. in $C(\mathbb{R}_+)$ toward 0. The pair (B^n, Z) therefore converges in law in $C(\mathbb{R}_+)$ toward a pair (B, Z) of Brownian motions, the brackets of which vanish. Knight’s theorem [28], Theorem 4.13, implies that B and Z are independent (see also the Appendix in [32]). As a measurable function of B , the Liouville Brownian motion is independent of Z . \square

2.9. Liouville Brownian motion defined on other geometries: Torus, sphere and planar domains. So far, we constructed in detail the LBM for the (Massive) Free Field on \mathbb{R}^2 . Our method applies to other two-dimensional manifolds like the torus, sphere, planar domains, . . . equipped with a log-correlated Gaussian field (of special interest is the case of Gaussian Free Field), stationary or not. The main reason is that Kahane’s theory remains valid on C^1 -manifolds (see [27, 35]). Intuitively,

this is just because such manifolds are locally isometric to open domains of the Euclidean space.

There is at least one point in our proofs that must be changed in order to apply to the torus or the sphere, or any compact manifold without boundary: the fact that $\lim_{t \rightarrow \infty} F(x, t) = +\infty$. Indeed, our proof uses the “infinite volume” of the plane. In the case of the torus or sphere, the strategy is much simpler because of compactness arguments: the standard Brownian motion on \mathbb{S}^2 or \mathbb{T}^2 possesses an invariant probability measure, call it μ , which is nothing but the volume form of \mathbb{S}^2 or \mathbb{T}^2 . Apply the ergodic theorem to prove that $\mathbb{P}^X \otimes \mathbb{P}_\mu^B$ almost surely:

$$\lim_{t \rightarrow \infty} \frac{F(t)}{t} = G,$$

for some random variable G , which is shift-invariant. Since the Brownian motion on the sphere is ergodic, G is measurable with respect to the sigma algebra generated by $\sigma\{X_x; x \in \mathbb{T}^2 \text{ or } \mathbb{S}^2\}$. It is not clear that G is constant. Yet, the set $\{G > 0\}$ is measurable with respect to the asymptotic sigma-algebra of the $(Y_n)_n$. Therefore, \mathbb{P}^μ almost surely, the set $\{G > 0\}$ has \mathbb{P}^X -probability 0 or 1. Since G has expectation 1, this set has \mathbb{P}^X -probability 1. Therefore, \mathbb{P}^X almost surely, the change of times F goes to ∞ as $t \rightarrow \infty$ for μ almost every x . Then use the coupling trick to deduce that the property holds for all starting points.

APPENDIX A: KAHANE’S CONVEXITY INEQUALITY

For the classical terminology of Gaussian multiplicative chaos, the reader is referred to [27] (see also [35]).

We consider a locally compact separable metric space (D, d) , a Radon measure ν on the Borel subsets of (D, d) and two Gaussian random distributions Y, Y' (in the sense of Schwartz) with respective covariance kernels K, K' , which are of σ -positive type. We assume that the Gaussian multiplicative chaos associated to (Y, ν) and (Y', ν) are strongly nondegenerate (e.g., the kernels of Y_n [see (2.5)] or the kernel $\gamma^2 G_m$ of (2.1) for $0 \leq \gamma^2 < 4$). Here, we recall the following standard lemma that can be found in [27].

LEMMA A.1. *Let $F : \mathbb{R}_+ \rightarrow \mathbb{R}$ be some convex function such that*

$$\forall x \in \mathbb{R}_+, \quad |F(x)| \leq M(1 + |x|^\beta),$$

for some positive constants M, β .

(1) *Assume that $K(u, v) \leq K'(u, v)$ for all $u, v \in D$. Then*

$$\mathbb{E}\left[F\left(\int_D e^{Y_r - (1/2)\mathbb{E}[Y_r^2]} \nu(dr)\right)\right] \leq \mathbb{E}\left[F\left(\int_D e^{Y'_r - (1/2)\mathbb{E}[Y_r'^2]} \nu(dr)\right)\right].$$

(2) If $K(u, v) \leq K'(u, v) + C$ for some constant $C > 0$ for all $u, v \in D$ then

$$\mathbb{E} \left[F \left(\int_D e^{Y_r - (1/2)\mathbb{E}[Y_r^2]} \nu(dr) \right) \right] \leq \mathbb{E} \left[F \left(e^{\sqrt{C}Z - C/2} \int_D e^{Y_r' - (1/2)\mathbb{E}[Y_r'^2]} \nu(dr) \right) \right],$$

where Z is a standard Gaussian random variable independent of the other random quantities.

APPENDIX B: FINITENESS OF THE MOMENTS

In this section, our only goal is to prove that

$$\forall x \in \mathbb{R}^2, \quad \mathbb{E}_x[F(t)^p] < +\infty$$

for $p \in [0, 4/\gamma^2[$. We only treat the case when $2 \leq \gamma^2 < 4$ and, therefore, $1 < p < 4/\gamma^2$ (hence $p < 2$). This is mathematically the most complicated part and notationally the easiest part. The case $0 \leq \gamma^2 < 2$ is discussed in Remark B.6.

Furthermore, by stationarity of the field X , we may assume that $x = 0$. By using the concavity of the mapping $x \mapsto x^{p/2}$ and the Jensen inequality, we get

$$\begin{aligned} \mathbb{E}_0[F(t)^p] &\leq \mathbb{E}^X[\mathbb{E}_0^B[F(t)^2]^{p/2}] \\ \text{(B.1)} \quad &\leq \mathbb{E}^X \left[\left(\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} f(x, y) M(dx) M(dy) \right)^{p/2} \right], \end{aligned}$$

where we have set

$$\text{(B.2)} \quad f(x, y) = \int_0^t \int_s^t e^{-|x|^2/(2s) - |y-x|^2/(2|r-s|)} \frac{dr ds}{4\pi^2 s|r-s|}.$$

So we just have to prove that the expectation in the above right-hand side is finite. In what follows, ξ_M stands for the structure exponent of the measure M . Recall that, in dimension d , it reads

$$\text{(B.3)} \quad \xi_M(p) = \left(d + \frac{\gamma^2}{2} \right) p - \frac{\gamma^2}{2} p^2.$$

Of course, we can take here $d = 2$. But is worth recalling this fact since it will happen that some arguments below will be carried out in dimension 1. So the reader will take care of replacing d by 1 when reading a proof in dimension 1. The main idea of our proof is the following. First, we observe that the function f possesses singularities. They are logarithmic (see below) when x or $|x - y|$ is close to 0. We will also have to treat the behavior near infinity. So we split the space $\mathbb{R}^2 \times \mathbb{R}^2$ into 3 domains

$$\begin{aligned} \text{(B.4)} \quad \mathcal{D}_1 &= \{|x| \leq 1, |x - y| \leq 1\}, & \mathcal{D}_2 &= \{|x| \geq 1, |x - y| \leq 1\}, \\ \mathcal{D}_3 &= \{|x - y| \geq 1\} \end{aligned}$$

and, by subadditivity of the mapping $x \in \mathbb{R}_+ \mapsto x^{p/2}$, it suffices to evaluate the quantity in the right-hand side of (B.1) on each of these three domains.

Concerning the behavior of f , we claim the following.

LEMMA B.1. *We have:*

1. *for all $x, y \in \mathbb{R}^2$: $f(x, y) \leq D(1 + \ln_+ \frac{1}{|x-y|})(1 + \ln_+ \frac{1}{|x|})$,*
2. *for all $|x| \geq 1$ and $|x - y| \leq 1$: $f(x, y) \leq D(1 + \ln_+ \frac{1}{|x-y|}) \exp(-\frac{|x|^2}{4t})$, for some constant $D > 0$.*

PROOF. Recall (B.2). By making successive changes of variables, we obtain

$$\begin{aligned} f(x, y) &= \int_0^t \int_0^{(t-s)/|x-y|^2} e^{-|x|^2/(2s)-1/(2r)} \frac{dr ds}{4\pi^2 sr} \\ &= \int_0^{t/|x|^2} \int_0^{(t-s|x|^2)/|x-y|^2} e^{-1/(2s)-1/(2r)} \frac{dr ds}{4\pi^2 sr} \\ &\leq g\left(\frac{t}{|x|^2}\right)g\left(\frac{t}{|x-y|^2}\right), \end{aligned}$$

where we have set

$$g(t) = \int_0^t e^{-1/(2s)} \frac{ds}{2\pi s}.$$

It is obvious to check that, for some constant $D > 0$, we have $g(t) \leq D(1 + \ln_+ t)$, which completes the proof of item 1. The proof of item 2 is similar and left to the reader. \square

NOTATION B.2. Until the end of the proof, we will only deal with expectations with respect to the measure M . So there is no need to keep on using the superscript X of \mathbb{E}^X and we will just write \mathbb{E} instead of \mathbb{E}^X .

Domain $\{|x| \leq 1, |x - y| \leq 1\}$. The main purpose of this part is to show that

$$(B.5) \quad \mathbb{E} \left[\left(\int_{\max(|x|, |y|) \leq 1} \frac{1}{|x|^\delta |x - y|^\delta} M(dx) M(dy) \right)^{p/2} \right] < +\infty.$$

The first step is to prove the following.

LEMMA B.3. *For $\gamma^2 < 4$ and $p \in]1, \frac{4}{\gamma^2}[$, there exist $\delta > 0$ and $C > 0$ such that for all $n \geq 0$,*

$$\mathbb{E} \left[\left(\int_{\max(|x|, |y|) \leq 2^{-n}} \frac{1}{|x|^\delta} M(dx) M(dy) \right)^{p/2} \right] \leq C 2^{-n(\xi_M(p) - (\delta p)/2)}.$$

PROOF. We carry out the proof in dimension 1 since, apart from notational issues, the dimension 2 does not raise any further difficulty. We first have to prove

$$\mathbb{E} \left[\left(\int_{(x, y) \in [0, 1]^2} \frac{1}{|x|^\delta} M(dx) M(dy) \right)^{p/2} \right] < +\infty.$$

Furthermore, from Kahane’s convexity inequalities, it suffices to prove the above lemma for any 1d log-correlated Gaussian field. Let us use the kernel $K(x, y) =$

$\ln_+ \frac{1}{|x-y|}$ of [5]. In fact, whatever the covariance kernel, we just need to use the property (2.9), which is shared by all the reasonable $1d$ log-correlated Gaussian fields (see [35] for more on this).

We also remind the reader that the above integral is finite for $\delta = 0$ (see [27]). Therefore, by using subadditivity of the mapping $x \mapsto x^{p/2}$, we have

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{[0,1]^2} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ &= \mathbb{E} \left[\left(\sum_{n=0}^\infty \int_{[2^{-n-1}, 2^{-n}] \times [0,1]} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ &\leq \sum_{n=0}^\infty \mathbb{E} \left[\left(\int_{[2^{-n-1}, 2^{-n}] \times [0,1]} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ &\leq \sum_{n=0}^\infty 2^{\delta(n+1)p/2} \mathbb{E}[(M([2^{-n-1}, 2^{-n}])M([0, 1]))^{p/2}]. \end{aligned}$$

Now we use the standard inequality $ab \leq \epsilon a^2 + \frac{b^2}{\epsilon}$ for any $\epsilon > 0$ and subadditivity of the mapping $x \mapsto x^{p/2}$ to get $(ab)^{p/2} \leq \epsilon^{p/2} a^p + \epsilon^{-p/2} b^p$. Therefore, with $a = M([2^{-n-1}, 2^{-n}])$, $b = M([0, 1])$ and $\epsilon = 2^{(n+1)\xi_M(p)/p}$, we obtain

$$\begin{aligned} \mathbb{E}[(M([2^{-n-1}, 2^{-n}])M([0, 1]))^{p/2}] &\leq (2^{(n+1)\xi_M(p)/p})^{p/2} \mathbb{E}[M([2^{-n-1}, 2^{-n}])^p] \\ &\quad + (2^{(n+1)\xi_M(p)/p})^{-p/2} \mathbb{E}[M([0, 1])^p]. \end{aligned}$$

By using (2.9), we get

$$\mathbb{E}[M([2^{-n-1}, 2^{-n}])^p] \leq C_p 2^{-(n+1)\xi_M(p)}$$

and plugging this relation into the above expression yields:

$$\mathbb{E}[(M([2^{-n-1}, 2^{-n}])M([0, 1]))^{p/2}] \leq 2^{-(n+1)\xi_M(p)/2} (C_p + \mathbb{E}[M([0, 1])^p]).$$

To sum up, we have

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{(x,y) \in [0,1]^2} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ &\leq (C_p + \mathbb{E}[M([0, 1])^p]) \sum_{n=0}^\infty 2^{-(n+1)(\xi_M(p)/2 - \delta p/2)}. \end{aligned}$$

So, δ can clearly be chosen small enough to make the above series convergent.

Once the finiteness of the expectation is proved, the statement results from a scaling argument. For $\lambda < 1$, the measure M satisfies (see [5] but this is elementary) the following relation in law:

$$(B.6) \quad (M(\lambda A))_{A \subset [0,1]} = (\lambda^{1+\gamma^2/2} e^{\gamma \Omega_\lambda} M(A))_{A \subset [0,1]},$$

where Ω_λ is a centered Gaussian random variable with variance $-\ln \lambda$ independent of $(M(A))_{A \subset [0,1]}$. Thus, we have

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{[0,\lambda]^2} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ &= \lambda^{p(1+\gamma^2/2)-\delta p/2} \mathbb{E} [e^{p\gamma\Omega_\lambda}] \mathbb{E} \left[\left(\int_{[0,1]^2} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right]. \end{aligned}$$

The result follows by taking $\lambda = 2^{-n}$. \square

LEMMA B.4. *For any $\gamma^2 < 4$ and $p \in]1, \frac{4}{\gamma^2}[$, there exist $\delta > 0$ and a constant $C > 0$ such that for all n :*

$$\mathbb{E} \left[\left(\int_{\substack{\max(|x|,|y|) \leq 1 \\ 2^{-n-1} \leq |x-y| \leq 2^{-n}}} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \leq \frac{C}{1-\delta p/2} 2^{-n(\xi_M(p)-2)}.$$

PROOF. Once again and for the same reason as previously, we carry out the proof in dimension 1. In that case, we have to prove

$$\mathbb{E} \left[\left(\int_{\substack{(x,y) \in [0,1]^2 \\ 2^{-n-1} \leq |x-y| \leq 2^{-n}}} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \leq C 2^{-n(\xi_M(p)-1)}.$$

Once again Kahane’s convexity inequality shows that we can take the kernel $K(x, y) = \ln_+ \frac{1}{|x-y|}$ of [5]. We will use the following elementary geometric argument: for any $n \geq 1$, the set of points 2^{-n} -close to the diagonal

$$\{(x, y) \in [0, 1]^2; |x - y| \leq 2^{-n}\}$$

is entirely recovered by the union for $k = 0, \dots, 2^n - 2$ of the (overlapping) squares $[\frac{k}{2^n}, \frac{k+2}{2^n}]^2$. Therefore, by using subadditivity of the mapping $x \mapsto x^{p/2}$, we have

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\substack{x,y \in [0,1] \\ 2^{-n-1} \leq |x-y| \leq 2^{-n}}} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ & \leq \sum_{k=0, \dots, 2^n-2} \mathbb{E} \left[\left(\int_{x,y \in [k/2^n, (k+2)/2^n]^2} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ & \leq \mathbb{E} \left[\left(\int_{x,y \in [0, 2^{-n+1}]^2} \frac{1}{|x|^\delta} M(dx)M(dy) \right)^{p/2} \right] \\ & \quad + \sum_{k=1, \dots, 2^n-2} \frac{2^{n\delta p/2}}{k^{\delta p/2}} \mathbb{E} \left[M \left(\left[\frac{k}{2^n}, \frac{k+2}{2^n} \right] \right)^p \right]. \end{aligned}$$

By stationarity and scale invariance (B.6), we get

$$\begin{aligned} & \sum_{k=1, \dots, 2^{n-2}} \frac{2^{n\delta p/2}}{k^{\delta p/2}} \mathbb{E} \left[M \left(\left[\frac{k}{2^n}, \frac{k+2}{2^n} \right] \right)^p \right] \\ & \leq 2^{n\delta p/2} \sum_{k=1, \dots, 2^{n-2}} \frac{1}{k^{\delta p/2}} \mathbb{E} [M([0, 2^{-n+1}])^p] \\ & \leq 2^{n\delta p/2} 2^{-(n-1)\xi_M(p)} \mathbb{E} [M([0, 1])^p] \sum_{k=1, \dots, 2^{n-2}} \frac{1}{k^{\delta p/2}} \\ & \leq \frac{C}{1 - \delta p/2} 2^{-n(\xi_M(p)-1)}, \end{aligned}$$

where C only depends on $\mathbb{E} [M([0, 1])^p]$. We conclude with Lemma B.3 provided we impose $\delta p/2 < 1$. \square

Now we are equipped to prove (B.5). Choose another $\delta' > 0$ such that $0 < \delta' < \frac{2(\xi_M(p)-2)}{p}$. By using Lemma B.4 and by subadditivity, we have

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\max(|x|, |y|) \leq 1} \frac{1}{|x|^\delta |x-y|^{\delta'}} M(dx) M(dy) \right)^{p/2} \right] \\ & \leq \sum_{n=0}^{+\infty} \mathbb{E} \left[\left(\int_{\substack{\max(|x|, |y|) \leq 1 \\ 2^{-n-1} \leq |x-y| \leq 2^{-n}}} \frac{1}{|x|^\delta |x-y|^{\delta'}} M(dx) M(dy) \right)^{p/2} \right] \\ \text{(B.7)} \quad & \leq \sum_{n=0}^{+\infty} 2^{(n+1)(\delta' p)/2} \mathbb{E} \left[\left(\int_{\substack{\max(|x|, |y|) \leq 1 \\ 2^{-n-1} \leq |x-y| \leq 2^{-n}}} \frac{1}{|x|^\delta} M(dx) M(dy) \right)^{p/2} \right] \\ & \leq \sum_{n=0}^{+\infty} 2^{(n+1)(\delta' p)/2} C 2^{-n(\xi_M(p)-2)}. \end{aligned}$$

Since the latter series converges, the proof of (B.5) is complete.

By gathering (B.5) and Lemma B.1 item 1, we deduce

$$\text{(B.8)} \quad \mathbb{E} \left[\left(\int_{|x| \leq 1, |x-y| \leq 1} f(x, y) M(dx) M(dy) \right)^{p/2} \right] < +\infty.$$

Domain $\{|x - y| \geq 1\}$. Let us now investigate the situation when $|x - y| \geq 1$. This is the easy part because, in that case, the measures $M(dx)$ and $M(dy)$ are “almost” independent. Therefore, we can proceed more directly in the computations.

We use the Jensen inequality with the concave function $x \mapsto x^{p/2}$ to get

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{|x-y| \geq 1} f(x, y) M(dx) M(dy) \right)^{p/2} \right] \\ & \leq \left(\mathbb{E} \left[\int_{|x-y| \geq 1} f(x, y) M(dx) M(dy) \right] \right)^{p/2} \\ & \leq \left(\int_{|x-y| \geq 1} f(x, y) e^{\gamma^2 G_m(x, y)} dx dy \right)^{p/2}. \end{aligned}$$

Since $|x - y| \geq 1$, we have $G_m(x, y) \leq C$ for some fixed positive constant C . We deduce that the above integral is less than $e^{Cp/2} (\int_{\mathbb{R}^2 \times \mathbb{R}^2} f(x, y) dx dy)^{p/2}$, which is equal to $e^{Cp/2}$, hence finite.

Domain $\{|x| \geq 1, |x - y| \leq 1\}$. The final part of the proof consists in checking that

$$(B.9) \quad \mathbb{E} \left[\left(\int_{|x| \geq 1, |x-y| \leq 1} f(x, y) M(dx) M(dy) \right)^{p/2} \right] < +\infty.$$

Because of Lemma (B.1) item 2, the above relation just boils down to proving that there exist $\delta > 0$ such that

$$(B.10) \quad \mathbb{E} \left[\left(\int_{\substack{|x| \geq 1 \\ |x-y| \leq 1}} \frac{\exp(-|x|^2/(4t))}{|x-y|^\delta} M(dx) M(dy) \right)^{p/2} \right] < +\infty.$$

Once again, we first need to estimate the above expectation on stripes of the type $\{|x| \geq 1, 2^{-n-1} \leq |x - y| \leq 2^{-n}\}$. So we claim the following.

LEMMA B.5. Fix $t > 0$. For any $\gamma^2 < 4$ and $p \in]1, \frac{4}{\gamma^2}[$, there exists a constant $C > 0$ (only depending on $\mathbb{E}[M([0, 1])^p]$) such that for all n :

$$\mathbb{E} \left[\left(\int_{\substack{|x| \geq 1 \\ 2^{-n-1} \leq |x-y| \leq 2^{-n}}} \exp\left(-\frac{|x|^2}{4t}\right) M(dx) M(dy) \right)^{p/2} \right] \leq C 2^{-n(\xi_M(p)-2)}.$$

Let us admit for a while the above lemma to finish the proof of (B.10). If we choose δ such that $0 < \delta < \frac{2(\xi_M(p)-2)}{p}$, we can then use Lemma B.5 and sub-additivity to get to computations (B.7), similar

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\substack{|x| \geq 1 \\ |x-y| \leq 1}} \frac{\exp(-|x|^2/(4t))}{|x-y|^\delta} M(dx) M(dy) \right)^{p/2} \right] \\ & \leq \sum_{n=0}^{+\infty} 2^{(n+1)(\delta p)/2} C 2^{-n(\xi_M(p)-2)}, \end{aligned}$$

which is a converging series.

PROOF OF LEMMA B.5. We keep on carrying out the proof in dimension 1 with the kernel $K(x, y) = \ln_+ \frac{1}{|x-y|}$. It is also plain to check that the expectation is finite thanks to the exponential term. We will prove the result when integrating only over the domain $\{x \geq 1, 2^{-n-1} \leq |x - y| \leq 2^{-n}\}$. It will then be obvious to complete the proof (e.g., by using invariance of M in law under reflection). As previously, the reader may check that the stripe $\{x \geq 1, 2^{-n-1} \leq |x - y| \leq 2^{-n}\}$ may be covered by the squares $[\frac{k}{2^n}, \frac{k+2}{2^n}]^2$ for k running over the set $K_n = \mathbb{Z} \cap [2^n, +\infty[$. Therefore, by using subadditivity of the mapping $x \mapsto x^{p/2}$, we have

$$\begin{aligned} & \mathbb{E} \left[\left(\int_{\substack{x \geq 1 \\ 2^{-n-1} \leq |x-y| \leq 2^{-n}}} \exp\left(-\frac{|x|^2}{4t}\right) M(dx)M(dy) \right)^{p/2} \right] \\ & \leq \sum_{k \in K_n} \mathbb{E} \left[\left(\int_{[k/2^n, (k+2)/2^n]^2} \exp\left(-\frac{k^2}{t2^{2n+2}}\right) M(dx)M(dy) \right)^{p/2} \right] \\ & = \sum_{k \in K_n} \exp\left(-\frac{k^2 p}{t2^{2n+2}}\right) \mathbb{E} \left[M\left(\left[\frac{k}{2^n}, \frac{k+2}{2^n}\right]\right)^p \right]. \end{aligned}$$

By stationarity and scale invariance (B.6), we get

$$\begin{aligned} & \sum_{k \in K_n} \exp\left(-\frac{k^2 p}{t2^{2n+2}}\right) \mathbb{E} \left[M\left(\left[\frac{k}{2^n}, \frac{k+2}{2^n}\right]\right)^p \right] \\ & = \sum_{k \in K_n} \exp\left(-\frac{k^2 p}{t2^{2n+2}}\right) \mathbb{E} [M([0, 2^{n-1}])^p] \\ & = 2^{-(n-1)\xi_M(p)} \sum_{k \in K_n} \exp\left(-\frac{k^2 p}{t2^{2n+2}}\right) \mathbb{E} [M([0, 1])^p] \leq C 2^{-n(\xi_M(p)-1)}, \end{aligned}$$

where C only depends on $\mathbb{E}[M([0, 1])^p]$. The last line uses the standard trick of convergence of Riemann sums. \square

REMARK B.6. If $\gamma < 2$, it is expected in great generality that F possesses moments of order p for $p < \frac{4}{\gamma^2}$. We proved that this is true in the more complicated situation $\sqrt{2} \leq \gamma < 2$. If $0 < \gamma < \sqrt{2}$, we only gave the existence of moments for $p \leq 2$. However, our strategy could be easily adapted to treat the case $p < \frac{4}{\gamma^2}$. In that case, one has to choose an integer $n \geq 2$ such that $p/n < 1$ and apply the Jensen inequality to get an expression similar to (B.1) (replace 2 by n) excepted that we get an integral over $(\mathbb{R}^2)^n$ instead of $(\mathbb{R}^2)^2$. Then we can reproduce our strategy up to modifications that are obvious but notationally awful.

Acknowledgements. The authors wish to thank M. Bauer, F. David, J. Dubédat, M. Gubinelli, J. F. Le Gall for fruitful discussions and comments. We also wish to thank the anonymous referee for his/her careful reading.

REFERENCES

- [1] ADLER, R. J. and TAYLOR, J. E. (2007). *Random Fields and Geometry*. Springer, New York. [MR2319516](#)
- [2] ALLEZ, R., RHODES, R. and VARGAS, V. (2013). Lognormal \star -scale invariant random measures. *Probab. Theory Related Fields* **155** 751–788. [MR3034792](#)
- [3] AMBJØRN, J., ANAGNOSTOPOULOS, K. N., ICHIHARA, T., JENSEN, L. and WATABIKI, Y. (1998). Quantum geometry and diffusions. Unpublished manuscript.
- [4] AMBJØRN, J., BOULATOV, D., NIELSEN, J. L., ROLF, J. and WATABIKI, Y. (1998). The spectral dimension of 2D quantum gravity. *J. High Energy Phys.* Available at <http://iopscience.iop.org/article/10.1088/1126-6708/1998/02/010/meta>.
- [5] BACRY, E. and MUZY, J. F. (2003). Log-infinitely divisible multifractal processes. *Comm. Math. Phys.* **236** 449–475. [MR2021198](#)
- [6] BARRAL, J., JIN, X., RHODES, R. and VARGAS, V. (2013). Gaussian multiplicative chaos and KPZ duality. *Comm. Math. Phys.* **323** 451–485. [MR3096527](#)
- [7] BARRAL, J., KUPIAINEN, A., NIKULA, M., SAKSMAN, E. and WEBB, C. (2014). Critical Mandelbrot cascades. *Comm. Math. Phys.* **325** 685–711. [MR3148099](#)
- [8] BASS, R. F. and KHOSHNEVISAN, D. (1992). Local times on curves and uniform invariance principles. *Probab. Theory Related Fields* **92** 465–492. [MR1169015](#)
- [9] BENJAMINI, I. and SCHRAMM, O. (2009). KPZ in one dimensional random geometry of multiplicative cascades. *Comm. Math. Phys.* **289** 653–662. [MR2506765](#)
- [10] BERESTYCKI, N. (2015). Diffusion in planar Liouville quantum gravity. *Ann. Inst. Henri Poincaré Probab. Stat.* **51** 947–964. [MR3365969](#)
- [11] BROSAMLER, G. A. (1970). Quadratic variation of potentials and harmonic functions. *Trans. Amer. Math. Soc.* **149** 243–257. [MR0270442](#)
- [12] CALCAGNI, G. (2012). Diffusion in multiscale spacetimes. *Phys. Rev. E* **87** 012123.
- [13] CALCAGNI, G. (2012). Diffusion in quantum geometry. *Phys. Rev. D* **86** 044021.
- [14] DAVID, F. (1988). Conformal field theories coupled to 2-D gravity in the conformal gauge. *Modern Phys. Lett. A* **3** 1651–1656. [MR0981529](#)
- [15] DAVID, F. (1992). What is the intrinsic geometry of two-dimensional quantum gravity? *Nuclear Phys. B* **368** 671–700. [MR1146107](#)
- [16] DAVID, F. and BAUER, M. (2009). Another derivation of the geometrical KPZ relations. *J. Stat. Mech. Theory Exp.* **3** P03004, 9. [MR2495865](#)
- [17] DEMBO, A., PERES, Y., ROSEN, J. and ZEITOUNI, O. (2001). Thick points for planar Brownian motion and the Erdős–Taylor conjecture on random walk. *Acta Math.* **186** 239–270. [MR1846031](#)
- [18] DISTLER, J. and KAWAI, H. (1989). Conformal field theory and 2-D quantum gravity or who’s afraid of Joseph Liouville? *Nuclear Phys. B* **321** 509–517.
- [19] DUPLANTIER, B., RHODES, R., SHEFFIELD, S. and VARGAS, V. (2014). Critical Gaussian multiplicative chaos: Convergence of the derivative martingale. *Ann. Probab.* **42** 1769–1808. [MR3262492](#)
- [20] DUPLANTIER, B., RHODES, R., SHEFFIELD, S. and VARGAS, V. (2014). Renormalization of critical Gaussian multiplicative chaos and KPZ formula. *Comm. Math. Phys.* **330** 283–330.
- [21] DUPLANTIER, B. and SHEFFIELD, S. (2011). Liouville quantum gravity and KPZ. *Invent. Math.* **185** 333–393. [MR2819163](#)
- [22] FUKUSHIMA, M., OSHIMA, Y. and TAKEDA, M. (1994). *Dirichlet Forms and Symmetric Markov Processes*. *De Gruyter Studies in Mathematics* **19**. de Gruyter, Berlin. [MR1303354](#)
- [23] GARBAN, C. (2012). In quantum gravity and the KPZ formula, Séminaire Bourbaki, 64e Année 2011–2012, no 1052.

- [24] GINSPARG, P. and MOORE, G. (1993). Lectures on 2D gravity and 2D string theory. In *Recent Direction in Particle Theory, Proceedings of the 1992 TASI* (J. Harvey and J. Polchinski, eds.). World Scientific, Singapore.
- [25] GLIMM, J. and JAFFE, A. (1981). *Quantum Physics: A Functional Integral Point of View*. Springer, New York. [MR0628000](#)
- [26] HSU, P. (1988). Brownian motion and Riemannian geometry. In *Geometry of Random Motion (Ithaca, NY, 1987)*. *Contemp. Math.* (R. Durrett and M. Pinsky, eds.) **73** 95–104. Amer. Math. Soc., Providence, RI. [MR0954633](#)
- [27] KAHANE, J.-P. (1985). Sur le chaos multiplicatif. *Ann. Sci. Math. Québec* **9** 105–150. [MR0829798](#)
- [28] KARATZAS, I. and SHREVE, S. E. (1991). *Brownian Motion and Stochastic Calculus*, 2nd ed. *Graduate Texts in Mathematics* **113**. Springer, New York. [MR1121940](#)
- [29] KNIZHNIK, V. G., POLYAKOV, A. M. and ZAMOLODCHIKOV, A. B. (1988). Fractal structure of 2D-quantum gravity. *Modern Phys. Lett. A* **3** 819–826. [MR0947880](#)
- [30] MOLCHAN, G. M. (1996). Scaling exponents and multifractal dimensions for independent random cascades. *Comm. Math. Phys.* **179** 681–702. [MR1400758](#)
- [31] NAKAYAMA, Y. (2004). Liouville field theory: A decade after the revolution. *Internat. J. Modern Phys. A* **19** 2771–2930. [MR2073993](#)
- [32] PITMAN, J. and YOR, M. (1986). Asymptotic laws of planar Brownian motion. *Ann. Probab.* **14** 733–779. [MR0841582](#)
- [33] RHODES, R., SOHIER, J. and VARGAS, V. (2014). Levy multiplicative chaos and star scale invariant random measures. *Ann. Probab.* **42** 689–724. [MR3178471](#)
- [34] RHODES, R. and VARGAS, V. (2011). KPZ formula for log-infinitely divisible multifractal random measures. *ESAIM Probab. Stat.* **15** 358–371. [MR2870520](#)
- [35] RHODES, R. and VARGAS, V. (2014). Gaussian multiplicative chaos and applications: A review. *Probab. Surv.* **11** 315–392. [MR3274356](#)
- [36] ROBERT, R. and VARGAS, V. (2008). Hydrodynamic turbulence and intermittent random fields. *Comm. Math. Phys.* **284** 649–673. [MR2452591](#)
- [37] ROBERT, R. and VARGAS, V. (2010). Gaussian multiplicative chaos revisited. *Ann. Probab.* **38** 605–631. [MR2642887](#)
- [38] SHAMOV, A. (2014). On Gaussian multiplicative chaos. Preprint. Available at [arXiv:1407.4418](#).
- [39] SHEFFIELD, S. (2007). Gaussian free fields for mathematicians. *Probab. Theory Related Fields* **139** 521–541. [MR2322706](#)
- [40] SZNITMAN, A.-S. (1998). *Brownian Motion, Obstacles and Random Media*. Springer, Berlin. [MR1717054](#)
- [41] WATABIKI, Y. (1993). Analytic study of fractal structure of quantized surface in two-dimensional quantum gravity. *Progr. Theoret. Phys. Suppl.* **114** 1–17. [MR1268911](#)

C. GARBAN
 INSTITUT CAMILLE JORDAN
 UNIVERSITÉ LYON 1
 43 BD DU 11 NOVEMBRE 1918
 69622 VILLEURBANNE CEDEX
 FRANCE
 E-MAIL: garban@math.univ-lyon1.fr

R. RHODES
 LAMA
 UNIVERSITÉ PARIS-EST MARNE LA VALLÉE
 CHAMPS SUR MARNE
 FRANCE
 E-MAIL: remi.rhodes@u-pem.fr

V. VARGAS
 DMA
 ENS PARIS
 45 RUE D'ULM
 75005 PARIS
 FRANCE
 E-MAIL: Vincent.Vargas@ens.fr