

Lipid Rafts Act as Specialized Domains for Tetanus Toxin Binding and Internalization into Neurons

Judit Herreros,^{*†‡} Tony Ng,^{§||} and Giampietro Schiavo^{*}

^{*}Molecular Neuropathobiology and [§]Cell Biophysics Laboratories, Imperial Cancer Research Fund, WC2A 3PX London, United Kingdom; and ^{||}Richard Dimbleby Department of Cancer Research, St. Thomas' Hospital, SE1 7EH London, United Kingdom

Submitted March 1, 2001; Revised June 25, 2001; Accepted July 11, 2001
Monitoring Editor: Juan Bonifacino

Tetanus (TeNT) is a zinc protease that blocks neurotransmission by cleaving the synaptic protein vesicle-associated membrane protein/synaptobrevin. Although its intracellular catalytic activity is well established, the mechanism by which this neurotoxin interacts with the neuronal surface is not known. In this study, we characterize p15s, the first plasma membrane TeNT binding proteins and we show that they are glycosylphosphatidylinositol-anchored glycoproteins in nerve growth factor (NGF)-differentiated PC12 cells, spinal cord cells, and purified motor neurons. We identify p15 as neuronal Thy-1 in NGF-differentiated PC12 cells. Fluorescence lifetime imaging microscopy measurements confirm the close association of the binding domain of TeNT and Thy-1 at the plasma membrane. We find that TeNT is recruited to detergent-insoluble lipid microdomains on the surface of neuronal cells. Finally, we show that cholesterol depletion affects a raft subpool and blocks the internalization and intracellular activity of the toxin. Our results indicate that TeNT interacts with target cells by binding to lipid rafts and that cholesterol is required for TeNT internalization and/or trafficking in neurons.

INTRODUCTION

Tetanus (TeNT) and botulinum neurotoxins (BoNTs) block neurotransmitter release and are responsible for tetanus and botulism, respectively. These toxins share a common structure comprised of a heavy (H, 100 kDa) and a light (L, 50 kDa) chain linked by a disulfide bond. The H chain mediates binding and internalization in neurons, whereas the L chain is a metallo-protease that selectively cleaves synaptic proteins (Niemann *et al.*, 1994; Schiavo *et al.*, 2000). TeNT and

BoNTs bind to the neuromuscular junction, but their intracellular actions take place at different levels of the nervous system. TeNT undergoes retrograde transport to the cell body of spinal cord motor neurons (MNs), is transcytosed, and cleaves the synaptic vesicle protein vesicle-associated membrane protein (VAMP)/synaptobrevin in inhibitory synapses. Instead, BoNTs act at a peripheral level by blocking acetylcholine release at the motor nerve terminal. This differential sorting has been interpreted as a consequence of binding to different surface receptors (Habermann and Dreyer, 1986; Herreros *et al.*, 1999).

TeNT and BoNTs bind to polysialogangliosides of the G1b series (Halpern and Neale, 1995). However, the fact that their binding is sensitive to proteases (Lazarovici and Yavin, 1986; Pierce *et al.*, 1986; Yavin and Nathan, 1986) suggests the existence of specific protein receptors. Thus, a model in which TeNT and BoNTs interact with a complex constituted by both lipid and protein receptors has been proposed (Montecucco, 1986). Despite several efforts, these protein receptors have not been conclusively identified. Several BoNT serotypes interact with synaptotagmins (Nishiki *et al.*, 1994; Li and Singh, 1998), but the role of these proteins as physiological BoNT receptors remains controversial (Evans *et al.*, 1986; Bakry *et al.*, 1997). We have previously followed a cross-linking approach (Schiavo *et al.*, 1991) to demonstrate that the binding domain of TeNT (TeNT H_c) interacts with a glycoprotein of ~15 kDa (p15) in nerve growth factor (NGF)-differentiated PC12 cells and spinal cord MNs (Her-

[†] Present address: Department of Biological Sciences, Imperial College of Science, Technology, and Medicine, Exhibition Rd., SW7 2AY London, UK.

[‡] Corresponding authors. E-mail addresses: g.schiavo@icrf.icnet.uk and j.herreros@ic.ac.uk.

¹ $\text{Eff} = 1 - \tau_{\text{DA}}/\tau_{\text{D}}$, where τ_{DA} is the lifetime map of the donor in the presence of acceptor, and τ_{D} is the average lifetime of the donor in the absence of acceptor.

Abbreviations used: BoNT, botulinum neurotoxin; b-TeNT, biotinylated-TeNT; CT, cholera toxin; DIG, detergent-insoluble glycolipid-enriched membrane; FLIM, fluorescence lifetime imaging microscopy; FRET, fluorescence resonance energy transfer; GPI, glycosylphosphatidylinositol; H, heavy chain; H_c, binding domain; L, light chain; MCDX, methyl- β -cyclodextrin; MN, motor neuron; p15, TeNT binding protein of ~15 kDa; PI-PLC, phosphoinositol-phospholipase C; TeNT, tetanus neurotoxin; TX-100, Triton X-100; VSV-G, vesicular stomatitis virus protein G.

ros *et al.*, 2000a). In these neuronal cell types p15 showed the behavior of an integral membrane protein (Herreros *et al.*, 2000a).

Several toxins, including cholera toxin (CT) (Orlandi and Fishman, 1998; Wolf *et al.*, 1998; Shogomori and Futerman, 2001) and all the known pore-forming toxins (Fivaz *et al.*, 1999; Gordon *et al.*, 1999), bind to lipid raft components. Furthermore, some bacteria and viruses enter cells via lipid rafts (Dehio *et al.*, 1995; Baorto *et al.*, 1997; Parton and Lindsay, 1999; Shin *et al.*, 2000). Lipid rafts are microdomains of the plasma membrane enriched in sphingolipids (including gangliosides), cholesterol, and glycosylphosphatidylinositol (GPI)-anchored proteins (Brown and London, 2000; Simons and Toomre, 2000). They have been implicated in vesicular sorting, trafficking to the apical membrane, and signaling (reviewed in Simons and Ikonen, 1997; Simons and Toomre, 2000). Recently, several lines of evidence have supported the role of cholesterol in the control of intracellular membrane trafficking (Grimmer *et al.*, 2000; Hoekstra and van Ijzendoorn, 2000; Mukherjee and Maxfield, 2000; Simons and Gruenberg, 2000).

Here, we characterize the cross-linking products containing TeNT H_C and p15 to show that p15s are GPI-anchored proteins in different neuronal cell types. Immunoprecipitation experiments identify p15 as Thy-1, a GPI-anchored neuronal raft protein (Williams and Gagnon, 1982; Madore *et al.*, 1999) in NGF-differentiated PC12 cells. The association of TeNT H_C with Thy-1 at the plasma membrane is confirmed by fluorescence resonance energy transfer (FRET) measurements with the use of fluorescence lifetime imaging microscopy (FLIM). Cell-bound TeNT H_C and Thy-1 are found in detergent-insoluble glycolipid-enriched (DIGs) fractions, which represent *in vitro* isolated lipid rafts (Brown and London, 2000). Furthermore, we show that cholesterol depletion causes the displacement of TeNT H_C from a DIG subpool and protects neurons from the toxic activity of TeNT *in vivo*.

MATERIALS AND METHODS

Materials

Recombinant TeNT H_C fragments were produced and radiolabeled with [γ -³²P]ATP as previously described (Lalli *et al.*, 1999; Herreros *et al.*, 2000a). TeNT was purified and biotinylated as in Arribas *et al.* (1993). Aerolysin and anti-aerolysin antibodies were kindly provided by Dr. G. van der Goot (University of Geneva, Geneva, Switzerland). Monoclonal antibody against GAP-43 (clone NM4) was from Autogen Bioclear (Calne, Wilshire, United Kingdom) and anti-HPC-1/syntaxin-1 and anti-SNAP-25 antibodies were a kind gift from T.H. Söllner (Memorial Sloan-Kettering Cancer Center, New York, NY).

Binding, Cross-linking, and Immunoprecipitation

Rat pheochromocytoma (PC12) cells were differentiated with 75 ng/ml 7S NGF (Alomone, Jerusalem, Israel) for 6–7 d. Spinal cord MNs were purified from E14 rat embryos and mouse spinal cord cells were isolated from E13 mice embryos (Lalli *et al.*, 1999; Herreros *et al.*, 2000a).

For phosphoinositol-phospholipase C (PI-PLC) treatment, cells were washed in serum-free medium, pretreated (1 h, 37°C) with different amounts of PI-PLC (Sigma, Poole, Dorset, United Kingdom) and extensively washed. Cells were cooled on ice, washed with Hanks' buffer (Herreros *et al.*, 2000a), and incubated (1 h, 4°C) with 300 pM ³²P-

labeled TeNT H_C in Hanks' buffer containing 0.2% bovine serum albumin. After binding, cells were washed and cross-linked (10 min, 4°C) with 0.22 mM bis[2(succinimidyl)oxycarbonyloxyethyl]sulfone (Perbio-Science, Tatten Hall, Cheshire, United Kingdom) in Hanks' buffer. The reaction was stopped and cells were solubilized as previously described (Herreros *et al.*, 2000a). Proteins were analyzed in 6–12% acrylamide gradient gels followed by autoradiography.

In immunoprecipitation experiments, selected samples were incubated with 1 nM CT (1 h, 4°C) after ³²P-labeled TeNT H_C binding and cross-linking. Cells were scraped in Hanks' buffer containing 1 mM Pefabloc (Roche Molecular Biochemicals, Mannheim, Germany), 1 mM iodoacetamide, 1 mM benzamidine, 1 μ g/ml aprotinin, 1 μ g/ml leupeptin, and phosphatase inhibitors cocktail (all from Sigma). Samples were pelleted down and solubilized (30 min, 4°C) with 4% octyl- β -D-glucopyranoside, 0.5% Triton X-100 (TX-100) in Hanks' buffer plus inhibitors. After centrifugation (10,000 rpm, 10 min), supernatants were incubated (2 h, 4°C) with protein G or protein A beads precoupled with monoclonal anti-vesicular stomatitis virus protein G (VSV-G) (Herreros *et al.*, 2000a), anti-Thy-1 (clone OX7), anti-CT (Biogenesis, Pool, Dorset, United Kingdom), and mock antibodies or affinity-purified polyclonal anti-PrP antibodies (Affinity Bioreagents, Golden, CO), respectively. Beads were extensively washed with detergent-containing Hanks' buffer. In some cases, an additional wash with 1 M urea was performed after the detergent washes. Beads were resuspended in SDS-sample buffer and analyzed as described above.

Isolation of DIGs and Western Blot

PC12 cells (seeded at 110 cells/mm²) were treated with NGF for 6–7 d. A total of 3.9×10^6 mouse spinal cord neurons was cultured for 2 wk before the experiment. Cells were washed and in selected cases pretreated (1 h, 37°C) with 4.5 mM methyl- β -cyclodextrin (MCDX; Sigma) in serum-free medium. After extensive washing, 300 pM TeNT ³²P-H_C or 1 nM biotinylated-TeNT (b-TeNT) was bound as mentioned above. Cells were washed with Hanks' buffer and resuspended in 1 ml (PC12) or 0.25 ml (spinal cord cells) of 1% TX-100 in Hanks' buffer plus inhibitors. Cells were solubilized (30 min, 4°C), the cell lysate was adjusted to 41% sucrose in Hanks' buffer, and overlaid with 8.5 ml of 35% sucrose and 2.5 ml of 16% sucrose. DIGs were isolated by ultracentrifugation (35,000 rpm, 18 h, 4°C; Abrami *et al.*, 1998). Then 11–12 fractions of 1 ml were collected, precipitated with 6.5% trichloroacetic acid in the presence of 0.05% sodium deoxycholate (Sigma), and washed with 80% cold acetone. Samples were analyzed by SDS-PAGE followed by Western blot and autoradiography. b-TeNT was detected with the use of streptavidin-peroxidase (1.25 μ g/ml; Sigma). Western blots were developed with the use of the enhanced chemiluminescence method (Amersham Pharmacia Biotech UK, Little Chalfont, Buckinghamshire, United Kingdom). Alternatively, in the case of mouse spinal cord cells, gels were rinsed for 20 min with 50 mM Tris-HCl, pH 7.4, 20% glycerol and transferred to nitrocellulose (5 h, 150 mA) in 10 mM NaHCO₃, 3 mM Na₂CO₃, pH 9.8 (Abrami *et al.*, 1998). Blots were rinsed with binding buffer (50 mM NaH₂PO₄, pH 7.5, 0.3% Tween 20) and incubated with aerolysin (2 nM, 2 h), followed by washing and Western blot with the use of anti-aerolysin antibodies.

For cross-linking experiments, fractions 2–5 (DIGs) and 9–12 (soluble) were pooled and cross-linked with 0.5 mM bis[2(succinimidyl)oxycarbonyloxyethyl]sulfone (10 min, 4°C, under shaking). The reaction was stopped with 30 mM glycine and samples were analyzed as described above.

Immunofluorescence

Fab fragment from the OX7 clone was purified with the use of the ImmunoPure Fab Preparation kit, according to manufacturer's instructions (Perbio-Science). Direct conjugation of IgG and Fab to Cy3 and Cy5 fluorophores (Amersham Pharmacia Biotech UK) was

performed at pH 8.5 (IgG) or pH 9.0 (Fab) as described previously (Bastiaens and Jovin, 1998).

Cells were incubated with either TeNT H_C (80–120 nM at 4 or 37°C) or CT (2 nM, 1 h, 4°C) and fixed with 3.7% paraformaldehyde (10 min at room temperature). TeNT H_C was immunodetected with the use of purified monoclonal antibodies against the VSV-G epitope as previously described (Lalli *et al.*, 1999). CT in Figure 4 was detected by with the use of anti-CT polyclonal antibodies (Sigma). When permeabilization was needed, cells were incubated with 0.1% TX-100 for 5 min. Texas Red- (Amersham Pharmacia Biotech UK) or Alexa 488-coupled secondary antibodies (Molecular Probes, Eugene, OR) were used according to manufacturer's instructions. Cells were visualized through a Plan-APOCHROMAT, 63×/1.4 numerical aperture phase 3 oil objective with the use of a laser scanning confocal microscope (Zeiss LSM 510; Zeiss, Jena, Germany). Images correspond to selected optical sections (collected in the z-axis at intervals of 0.4 μm).

FRET Determination by FLIM

For FLIM measurements, immunocytochemical staining was performed as described above, without cell permeabilization, and included an additional fixation with 3.7% paraformaldehyde before mounting. A detailed description of the FLIM apparatus used for FRET determination can be found elsewhere (Squire and Bastiaens, 1999). The lifetime instrument performs phase- and modulation-based imaging fluorometry by microscopy. All images were taken with the use of a Zeiss Plan-APOCHROMAT 100×/1.4 numerical aperture phase 3 oil objective and the homodyne phase-sensitive images recorded at a modulation frequency of 80.224 MHz. Donors (Cy3-labeled monoclonal anti-VSV-G or anti-CT IgGs) were excited with the use of the 414 nm line of an argon/krypton laser and the resultant fluorescence separated with the use of a combination of dichroic beamsplitter (HQ 565 LP; Chroma, Brattleboro, VT) and emission filter (HQ 610/75; Chroma). Acceptor images (Cy5-anti Thy-1 Fab) were recorded with the use of a 100-W mercury arc lamp (Zeiss Attoarc) as a source of sample illumination combined with a high Q Cy3 filter set (exciter, HQ 620/60; dichroic, HQ 660 LP; emitter, HQ 700/75 LP; Chroma).

TeNT Intoxication and VAMP Cleavage

Mouse spinal cord cells (10–12 d in culture) were pretreated with 2 or 4.5 mM MCDX (1 h, 37°C) in serum-free medium and washed. The effect of adding cholesterol-MCDX complexes could not be assessed because this treatment was toxic in spinal cells. Treated and untreated cells were incubated with 200 pM TeNT (20 h, 37°C) in serum-free medium. Cells were scraped in Hanks' buffer plus Pefabloc and proteins recovered by trichloroacetic acid precipitation. Samples were analyzed by SDS-PAGE containing urea and Western blot with the use of anti-VAMP-2 monoclonal antibodies (Edelmann *et al.*, 1995). VAMP/synaptobrevin immunoreactivity was quantified with the use of NIH Image 1.61 and normalized to syntaxin-1 for equal loading.

Transferrin Uptake

MCDX-treated and untreated spinal cord cells were assessed for ¹²⁵I-transferrin uptake (human diferric form; 670 ng/ml; PerkinElmer Life Science Products, Boston, MA) in serum-free medium at 4°C or 37°C for different times. After two rounds of acid wash (0.2 M acetic acid, 0.5 M NaCl, pH 2.5) (Hopkins and Trowbridge, 1983) for 2 min on ice followed by a wash in medium, cell lysates were recovered in 1 M NaOH and counted in a gamma counter (Packard Instrument, Meriden, CT). Counts represent endocytosed transferrin. Competition was tested with the use of 200× excess of human holotransferrin (Sigma).

RESULTS

p15s Are GPI-anchored TeNT H_C Binding Proteins

We recently demonstrated that cross-linking products very similar in size are formed after binding of ³²P-labeled TeNT H_C (Figure 1A) to NGF-differentiated PC12 cells, rat MNs, and mouse spinal cord cells (Herrerros *et al.*, 2000a). These results indicated the interaction of TeNT H_C (48 kDa) with a protein of an apparent molecular weight of 15 kDa (p15) in different cell types. p15s are N-glycosylated and behave as integral membrane proteins in detergent-partitioning experiments (Herrerros *et al.*, 2000a). This last property of p15 could be mediated by the presence of transmembrane domain(s) or by various forms of covalent lipid modification.

GPI-anchored proteins are abundant components of the plasma membrane, behaving as integral membrane proteins. Their biochemical analysis is simplified by the use of PI-PLC, an enzyme that causes their selective release from the membrane. We therefore tested the possibility that p15s are GPI-anchored proteins. Pretreatment with PI-PLC before TeNT H_C binding and cross-linking inhibited the formation of the previously characterized ~65 kDa cross-linking product in NGF-differentiated PC12 cells in a dose-dependent manner (Figure 1B, filled arrowhead), whereas total binding was not significantly affected (Figure 1B, empty arrowhead). Formation of the corresponding cross-linking products in MNs and spinal cord cells (Figure 1B, filled arrowheads) was also inhibited by PI-PLC pretreatment. These results indicate that p15s, proteins interacting with TeNT H_C in our cross-linking assays, are GPI-anchored proteins in all three neuronal cell types.

p15 Is Thy-1 in NGF-differentiated PC12 Cells

In our attempts to identify p15s, we searched for known proteins with the properties described above. Among others, Thy-1 (Williams and Gagnon, 1982), a major cell-surface GPI-anchored glycoprotein of 25–29 kDa present in thymocytes and brain and expressed in PC12 cells (Jeng *et al.*, 1998), represented a likely candidate. To test whether p15 is Thy-1, immunoprecipitation experiments with a panel of different antibodies was performed in NGF-differentiated PC12 cells (Figure 2A). As expected, antibodies against the VSV-G tag of TeNT H_C pulled down both the unmodified binding fragment and the ~65-kDa cross-linking product (Figure 2A). The ~65-kDa cross-linking band was specifically recovered with monoclonal antibodies against Thy-1, but not with mock antibodies or with antibodies against PrP, a GPI-anchored protein similar in size to Thy-1 (Figure 2A). Antibodies against CT, which binds to GM1 and is used as a raft marker, did not immunoprecipitate the cross-linking product (Figure 2A), indicating that lipid rafts had been disrupted by solubilization with octyl-β-D-glucopyranoside under our experimental conditions (Arni *et al.*, 1998; Simons *et al.*, 1999; Simons and Toomre, 2000). Furthermore, incubation of cells with both CT and TeNT H_C followed by cross-linking did not result in the appearance of additional cross-linking products (our unpublished results). These findings suggest that the cross-linking of TeNT H_C to Thy-1 is not a consequence of high local concentration or proximity but rather due to their direct interaction. Interestingly, anti-




Figure 1. p15s are GPI-anchored proteins. (A) Schematic representation of TeNT, the binding fragment H_C and the carboxyl-terminal subdomain H_CC. (B) Binding and cross-linking of ³²P-labeled TeNT H_C after pretreatment with increasing doses of PI-PLC in NGF-differentiated PC12 cells, rat MNs, and mouse spinal cord cells. Empty arrowhead indicates cell-bound TeNT-H_C and filled arrowheads point to the corresponding cross-linking products, which do not form after PI-PLC treatment.

Thy-1 antibodies coimmunoprecipitated a fraction of TeNT H_C that could not be dissociated from the beads with 1 M urea (Figure 2A), suggesting a strong interaction between Thy-1 and TeNT H_C. Moreover, the decrease in Thy-1 immunoreactivity after PI-PLC treatment of NGF-differentiated PC12 cells strictly correlated with the inhibition of the formation of the ~65-kDa cross-linking product (our unpublished results). The apparent molecular weight of the cross-linking product is lower than the sum of the individual components (48 and 25 kDa). Anomalous migration of cross-linking adducts in SDS-PAGE has been reported (Tate and Khadse, 1987) and is dependent on the nature of the interacting proteins surfaces and the generation of conformational constraints stable in denaturing conditions.

We have previously demonstrated that the carboxyl-terminal subdomain of TeNT H_C (H_CC; Figure 1A) is necessary and sufficient for binding to p15 (Herreros *et al.*, 2000b). Consistently, anti-Thy-1 but not mock monoclonal antibodies immunoprecipitated the cross-linking product obtained after binding of ³²P-labeled H_CC to NGF-differentiated PC12 cells (Figure 2B).

The interaction of TeNT H_C with Thy-1 in its membrane environment was further analyzed by FLIM (Figure 3). With the use of FLIM, we determined the extent of FRET between a Cy3-labeled IgG directed against the VSV-G epitope of TeNT H_C (donor) and a Cy5-conjugated IgG Fab fragment recognizing Thy-1 (acceptor). FRET results in a shortening of the donor fluorescence lifetime, which is measured by two independent parameters, the phase shift (τ_p) and relative modulation depth (τ_m). FRET is only detected between two proteins that are closely associated or complexed in vivo, typically within 10 nm (Selvin, 2000). NGF-differentiated

PC12 cells were stained with a Cy3-VSV-G IgG to visualize cell-bound TeNT H_C after incubation with the toxin for 1 h at 4°C or 30 min at 37°C either alone, or together with a Cy5-labeled anti-Thy-1 Fab fragment. The fluorescence lifetime, $\langle\tau\rangle$, measured as the average phase shift and relative modulation depth [$(\tau_p + \tau_m)/2$] for Cy3-VSV-G antibodies was decreased at punctate structures (see below) of the plasma membrane (Figure 3, A and B, right). This was particularly evident when TeNT H_C had been incubated with the cells at 37°C. The presence of FRET was confirmed by photobleaching the Cy5 acceptor, resulting in a lengthening of the lifetime of the donor (our unpublished results). The FRET efficiency (Eff) pseudocolor plots also indicate that FRET occurs at the plasma membrane. Statistical analysis of cumulative results from all the cells analyzed demonstrates that TeNT H_C interacts with Thy-1 at 4°C and that their association is enhanced at 37°C (see two-dimensional τ_p vs. τ_m lifetime profiles and pixel counts vs. FRET efficiency plots in the lower part of Figure 3A). Permeabilized cells (allowing staining of putative intracellular structures) showed the same FRET efficiency (our unpublished results), indicating that FRET was due to the interaction of the TeNT H_C and Thy-1 at the plasma membrane. Parallel experiments with the use of a Cy3-labeled anti-CT IgG to visualize cell-bound CT either alone or together with a Cy5-labeled anti-Thy-1 Fab fragment showed no significant FRET, compared with the positive control cells (stained with Cy3-labeled anti-VSVG IgG against cell-bound TeNT H_C/Cy5-labeled anti-Thy-1 Fab fragment) (Figure 3B), indicating that localization to lipid rafts per se is not sufficient to produce FRET under our experimental conditions. Taken together, these results identify p15 as Thy-1 in NGF-differentiated PC12 cells and




Figure 2. Thy-1 is p15 in NGF-differentiated PC12 cells. Immunoprecipitation (IP) after binding and cross-linking (CL) of ^{32}P -labeled TeNT- H_C (A) or H_C (B) with the use of mock antibodies (anti-myc or anti-HA) or antibodies against VSV-G, Thy-1, CT, or PrP in NGF-differentiated PC12 cells. Filled arrowheads indicate the position of the corresponding cross-linking products.

indicate that TeNT H_C specifically associates with Thy-1 on the surface of these cells.

TeNT Interacts with Lipid Rafts

GPI-anchored proteins and gangliosides, together with cholesterol and other sphingolipids, concentrate in microdomains of the plasma membrane called lipid rafts (Jacobson and Dietrich, 1999; Simons and Toomre, 2000). The interaction of TeNT H_C with Thy-1, which is considered a neuronal raft marker (Aarts *et al.*, 1999; Madore *et al.*, 1999), therefore suggested that the binding of TeNT to the cell surface could be mediated by lipid rafts. To test this hypothesis, we first looked at the distribution of Thy-1 and TeNT H_C in NGF-differentiated PC12 cells. Bound TeNT H_C displays a punctate pattern on the plasma membrane (Lalli *et al.*, 1999; Herreros *et al.*, 2000a), which is reminiscent of lipid rafts (Mayor *et al.*, 1994; Harder *et al.*, 1998) and is shared by Thy-1 (Jeng *et al.*, 1998) (Figure 4, a–c). Furthermore, GM1, another raft marker (Orlandi and Fishman, 1998; Wolf *et al.*, 1998), showed a partial colocalization with TeNT H_C (Figure 4, d–f).

To investigate more directly whether TeNT H_C binds to lipid microdomains, we prepared DIG-enriched membranes. Isolation of DIGs is one of the most widely used

methods for studying lipid rafts (Brown and London, 2000). DIGs were purified from NGF-differentiated PC12 cells after binding of 300 pM ^{32}P -labeled TeNT H_C and radioactivity was followed along the gradient fractions. TeNT H_C concentrated in fractions 2–4 at the top of the gradient (Figure 5A), corresponding to DIGs. DIG fractions were defined by their enrichment in Thy-1 (Figure 5A) and other raft markers such as the ganglioside GM1 or PrP (our unpublished results). SNAP-25, a palmitoylated plasma membrane protein, remained concentrated in the bottom fractions of the gradient (Figure 5A), which contain the soluble material and the majority of proteins. Similarly, only a very small amount of VAMP/synaptobrevin, an integral membrane protein of synaptic vesicles, was found in DIGs (Figure 5A; Chamberlain *et al.*, 2001). The ability of TeNT H_C to reach the top of the gradient is dependent on the presence of cell lysate (our unpublished results), indicating that the flotation of TeNT H_C to the lighter fractions is not due to the presence of detergent and the centrifugation procedure.

To assess whether the interaction of TeNT H_C and Thy-1 is stable in the conditions used for DIG isolation, TeNT H_C was bound to NGF-differentiated PC12 cells and then DIGs or soluble fractions were isolated and used for cross-linking. DIGs showed the same pattern of cross-linking seen in intact





Figure 3. TeNT H_c and Thy-1 association in NGF-differentiated PC12 cells detected by FLIM. NGF-differentiated PC12 cells were incubated with TeNT H_c (1 h at 4°C or 30 min at 37°C) or CT (30 min at 37°C), washed, and then fixed in paraformaldehyde. Cells were stained with a Cy3-labeled anti-VSV-G IgG (donor) alone (-anti-Thy1-Cy5) or together with a Cy5-conjugated anti-Thy1 IgG Fab fragment (+anti-Thy1-Cy5) (A and right panels of B). In parallel experiments, cells were stained with Cy3-labeled anti-CT IgG (donor) alone (-anti-Thy1-Cy5) or together with a Cy5-conjugated anti-Thy-1 IgG Fab fragment (+anti-Thy1-Cy5) (B, left). The fluorescence images from the donor (TeNT H_c anti-VSV-G-Cy3 or anti-CT-Cy3) and acceptor (anti-Thy1-Cy5), the donor fluorescence lifetime $\langle \tau \rangle$ and its corresponding pseudocolor scales are shown. Eff is the pixel-by-pixel FRET efficiency represented on a pseudocolor scale (see details in the text) and the average lifetime of the donor in the absence of acceptor (τ_D) used to calculate Eff has been taken by averaging six anti-VSVG or anti-CT-Cy3 donor (alone without acceptor) measurements. The cumulative lifetimes of the donor alone (green) and those measured in the presence of the acceptor fluorophore (red) are plotted on the two-dimensional histograms ($n = 7$ in A and $n = 6$ in B). The pixel counts versus Eff profiles summarize all the FRET efficiency data observed. FRET between TeNT H_c and Thy-1 is greatly enhanced by incubation at 37°C and is not significant between CT and Thy-1.

Figure 4. TeNT H_C binding colocalizes with the raft markers Thy-1 and GM1. Binding of TeNT H_C (1 h, 4°C) was revealed with the use of Cy3 labeled-monoclonal anti-VSV-G antibodies (b) or Texas Red-coupled secondary antibodies (e) in nonpermeabilized NGF-differentiated PC12 cells. Thy-1 immunoreactivity was detected with the use of Cy5-labeled anti-Thy-1 Fab (a) (staining is shown in green pseudocolor). GM1 was detected by CT binding (d; 1 h, 4°C) followed by polyclonal anti-CT and Alexa 488-coupled secondary antibodies. Overlaps are shown in c and f (colocalization appears in yellow). Confocal images correspond to single z-sections taken at a similar distance from the substrate. Bar, 10 μm.



cells (Figure 5B). Both the ~65-kDa cross-linking product representing the interaction of TeNT H_C with Thy-1 and the ~85-kDa adduct, which is likely to reflect the formation of H_C homodimers (Herrerros *et al.*, 2000a), were obtained in purified DIGs (Figure 5B). In contrast, no cross-linking products were observed in the soluble fraction.

In parallel experiments, b-TeNT was bound to NGF-differentiated PC12 cells and DIGs were isolated. As shown for TeNT H_C, b-TeNT was found in DIG fractions, although in this case the association with DIGs was not complete and some of the bound toxin remained with the soluble material (Figure 5C, fractions 10–12). A possible explanation for this result would be that TeNT binding is more sensitive to TX-100 solubilization than that of H_C due to steric hindrance caused by the translocation domain of the holotoxin. The fact that both TeNT H_C and TeNT were found in DIGs is supported by their very similar punctate distribution on the plasma membrane (Herrerros *et al.*, 2000a).

These results indicate that the interaction of TeNT and its binding fragment H_C with the plasma membrane of NGF-differentiated PC12 cells is mediated by lipid microdomains. The two identified TeNT ligands, polysialogangliosides and Thy-1, are clustered in lipid rafts, which could therefore serve as concentration platforms for the toxin binding. However, Thy-1 appears to be a dispensable component of the machinery involved in TeNT binding to the neuronal surface. In fact, anti-Thy-1 antibodies were not able to immunoprecipitate the cross-linking product of TeNT H_C with a GPI-anchored protein present on the surface of MNs (our unpublished results), indicating that p15 is not Thy-1 in these cells (see DISCUSSION). TeNT interaction with Thy-1 and other GPI-anchored proteins could therefore be interpreted as an indication of the entry of the toxin in a raft environment and used as a probe to follow the recruitment of TeNT into lipid microdomains.

TeNT Binds to Lipid Rafts in Spinal Neurons

Our observations on the interaction of TeNT H_C with GPI-anchored proteins in all the different cell types tested prompted us to investigate TeNT recruitment to lipid microdomains in neurons. We prepared DIGs from mouse spinal cord cells, a mixed culture that has been extensively used for the study of TeNT binding and internalization (Lalli *et al.*, 1999; Williamson *et al.*, 1999). After binding of TeNT H_C, isolated DIGs contained ~60% of the total bound toxin and the majority of Thy-1.2 (the mouse allotype for Thy-1; Figure 6) as detected by overlay with the GPI-binding toxin aerolysin (Abrami *et al.*, 1998). A small subpool of TeNT H_C was found in fractions containing soluble proteins at the bottom of the gradient, possibly reflecting differences in the lipid composition of spinal cord and NGF-differentiated PC12 cells (Figure 6).

Cholesterol Depletion Protects Neurons from TeNT Intoxication

The use of cholesterol-sequestering drugs to disrupt membrane rafts is well established (reviewed in Simons and Toomre, 2000). Studies with these drugs led to the conclusion that cholesterol plays a key structural role in the lipid microdomain architecture (Schroeder *et al.*, 1998), although some raft components appear resistant to these drugs in nonneuronal cells (Abrami and van der Goot, 1999; Lipardi *et al.*, 2000).

Mouse spinal cord cells were pretreated with MCDX, a drug that extracts cholesterol from the membranes (Neufeld *et al.*, 1996), before binding of TeNT H_C and isolation of DIGs. Under these conditions GAP-43, a palmitoylated protein of the neuronal membrane, was completely extracted from DIGs (Laux *et al.*, 2000) and represents a positive control for the treatment (Figure 7; compare left and right, bottom panels). Interestingly, preincubation with MCDX in-




Figure 5. TeNT H_c and TeNT are found in DIGs isolated from NGF-differentiated PC12 cells. (A) Bound ^{32}P -labeled TeNT H_c (300 pM) concentrates in the lighter fractions (2–4; DIGs) of the gradient (see MATERIALS AND METHODS), which contain the majority of Thy-1. In contrast, SNAP-25 and VAMP are mainly found in soluble fractions (11 and 12). (B) TeNT H_c and Thy-1 interact in isolated DIGs. The cross-linking (CL) pattern observed in intact NGF-differentiated PC12 cells (total) is reproduced when DIGs (pooled fractions 2–5) but not the soluble fractions (9–12) were used for cross-linking. Filled arrowhead indicates the ~65-kDa cross-linking product representing the interaction of TeNT H_c with Thy-1 and the asterisk points to the ~83-kDa cross-linking band corresponding to TeNT H_c homodimers. (C) b-TeNT (1 nM) is found in the lighter fractions (2–5) of the gradient where it comigrates with Thy-1 and also in fractions 11–12. The band showed corresponds to the H chain of b-TeNT.

duced a moderate, but consistent, shift of TeNT H_c from DIGs to the soluble fractions of the gradient (Figure 7). The increase in the solubility of TeNT H_c correlates with an increase in Thy-1.2 in the same fractions. These results indicate that distinct raft components are differently affected by cholesterol depletion and suggest that in spinal cord cells TeNT H_c binds to distinct lipid rafts subpools, which display different sensitivities to MDCX.

After the effect of MCDX on the recruitment of TeNT H_c to DIGs, we investigated whether cholesterol plays a role in the internalization or intracellular sorting of TeNT. Pretreatment of spinal cord cells with MCDX inhibited the internalization of TeNT H_c as shown by immunofluorescence (Figure 8A). Endocytic structures labeled with TeNT H_c in control cells (which are internal in single confocal z-sections; Figure 8A) were greatly reduced in MCDX-treated neurons,




Figure 6. TeNT H_c fragment associate with DIGs in spinal cord cells. TeNT H_c (300 pM) concentrates in DIGs and only a small portion is found in the soluble fractions. Thy-1.2, the mouse Thy-1 allotype (which is not recognized by OX7 anti-Thy-1 antibodies), was detected with the use of the GPI-binding toxin aerolysin (Abrami *et al.*, 1998; see MATERIALS AND METHODS) and its identification among other GPI-anchored proteins was inferred from the comparison with samples prepared from Thy-1 knockout cells. Thy-1.2 comigrates with TeNT H_c in DIG fractions.




Figure 7. MCDX treatment partially extracts TeNT H_C from DIGs in spinal cord cells. TeNT H_C is partially solubilized from DIGs (lanes 2 and 3) and its recovery in soluble fractions (lanes 11 and 12) increases accordingly upon MCDX treatment (right). Thy-1.2 follows the same pattern of TeNT H_C before and after MCDX, whereas GAP-43 is completely extracted from DIGs (lanes 2 and 3) by MCDX (bottom right).

where TeNT H_C was concentrated on the plasma membrane (Figure 8A). The limited intracellular staining present in MCDX-treated cells closely resembled the background levels of control samples. These observations suggest that cholesterol depletion inhibits TeNT H_C internalization in neurons.

Because MCDX blocked TeNT internalization, we next asked whether MCDX-treated cells would be protected from the intracellular zinc-endopeptidase activity of TeNT. We followed the effect of MCDX preincubation on the proteolysis of VAMP/synaptobrevin, the intracellular target of TeNT (Schiavo *et al.*, 2000). In spinal cord neurons, ~50% of VAMP is cleaved upon overnight incubation with 200 pM TeNT, as detected by Western blot with an anti-VAMP antibody (Figure 8B) (Lalli *et al.*, 1999). This partial cleavage is due both to the resistance of VAMP to TeNT proteolysis when engaged in preformed soluble *N*-ethylmaleimide-sensitive fusion protein-attachment receptor complexes (SNARE) (Schiavo *et al.*, 2000) and to the contamination of the neuronal culture with VAMP-containing glial cells (Pappas *et al.*, 1995), which are not competent to TeNT intoxication. Cell pretreatment with 4.5 mM MCDX completely protected VAMP from TeNT cleavage (Figure 8B). The effect of MCDX is dose-dependent as demonstrated by the lack of protection seen with 2 mM MCDX (Figure 8B). We did not observe any significant change in the recovery of VAMP after treatment with the drug alone (our unpublished results). Transferrin uptake in untreated and MCDX-treated spinal cells was not significantly different (Figure 8C), indicating that under our experimental conditions, this treatment does not affect unspecifically cell functionality or viability (Figure 8C) (see DISCUSSION). Taken together, these results demonstrate that cholesterol plays a key role in the internalization of TeNT in spinal cord neurons.

DISCUSSION

In this study, we demonstrate that TeNT H_C interacts with GPI-anchored proteins, p15s, in PC12 cells, spinal cord neurons, and purified MNs. In NGF-differentiated PC12 cells, we identify p15 as Thy-1 and show by immunoprecipitation and FLIM experiments that Thy-1 interacts with TeNT H_C

and H_CC, the H_C subdomain sufficient for binding, in this neuronally differentiated cell line. Thy-1 is a major component of neurons and T lymphocytes, where it is expressed in several distinct glycoforms (Parekh *et al.*, 1987). Thy-1 has been implicated in multiple processes, including neurite outgrowth (Tiveron *et al.*, 1992), long-term potentiation (Nosten-Bertrand *et al.*, 1996), and T-cell receptor signaling (Hueber *et al.*, 1997). Despite the strong similarities observed between p15s in different neuronal cell types (Herrerros *et al.*, 2000a), several lines of evidence indicate that Thy-1 is not the neuronal receptor for TeNT. Spinal cord cells isolated from Thy-1 knockout mice (Nosten-Bertrand *et al.*, 1996) show TeNT H_C binding and internalization similar to those isolated from wild-type animals (our unpublished results). Together with the inability of an anti-Thy-1 antibody to immunoprecipitate the TeNT H_C cross-linking product in rat MNs (our unpublished results), our findings indicate that Thy-1 is not an essential component of the TeNT binding and internalization machinery and suggest that a still unidentified glycosylated GPI-anchored protein could act as cellular acceptor for TeNT in MNs.

The identification of GPI-anchored proteins (including Thy-1) as specific TeNT binding partners and the concentration of these proteins in lipid rafts led us to investigate the association of TeNT with lipid microdomains. Several experiments suggest that TeNT binds to lipid rafts. First, the punctate plasma membrane staining obtained after TeNT H_C binding to NGF-differentiated PC12 cells, spinal cord cells, and purified MNs is reminiscent of lipid rafts (Mayor *et al.*, 1994; Harder *et al.*, 1998). Second, TeNT as well as TeNT H_C associates with DIG fractions both in NGF-differentiated PC12 cells and spinal cord neurons. Third, the interaction of TeNT H_C with Thy-1 occurs in DIGs. Finally, binding of anti-TeNT H_C antibody induces clustering in unfixed cells (our unpublished results), a feature of lipid raft components (Mayor *et al.*, 1994). In this light, the molecular interaction between Thy-1 and TeNT H_C shown by cross-linking and FLIM in NGF-differentiated PC12 cells could be enhanced by the association of both proteins to lipid rafts (Friedrichson and Kurzchalia, 1998; Varma and Mayor, 1998). FRET efficiency was higher at 37°C than at 4°C, suggesting that physiological temperatures promote a closer




Figure 8. MCDX treatment blocks TeNT H_C internalization and the toxic activity of TeNT in spinal cord cells. (A) After internalization of TeNT H_C (100 nM, 1 h, 37°C) and permeabilization, untreated cells show a dotted intracellular staining, detected with the use of monoclonal anti-VSV-G antibodies. MCDX-treated (4.5 mM) cell bodies and neurites exhibit membrane staining but internalized TeNT H_C is not observed. Confocal images corresponding to single z-sections of 0.4 μm were taken at the same distance from the substrate. Bar, 10 μm. (B) TeNT-induced cleavage of VAMP (compare lane 2 vs. lane 1) is slightly (lane 3) or completely protected (lane 4) by pretreatment with 2 or 4.5 mM MCDX, respectively. Syntaxin-1 (isoforms 1a and 1b) is used as a loading control. Results from four independent experiments (±SD), normalized for syntaxin-1, are plotted in the panel below. (C) ¹²⁵I-Transferrin endocytosis at 37°C is similar in untreated (■) and MCDX-treated (4.5 mM, ▨) spinal cord cells. Results from two independent experiments (±SD; p > 0.05) are shown. ¹²⁵I-Transferrin binding (20 min, 4°C) and residual endocytosis upon competition with unlabeled transferrin accounts for 12 ± 3 and 25 ± 3% of maximal endocytosis, respectively (our unpublished results).

interaction between the two proteins directly or via changes in the lateral mobility of raft proteins. We did not however detect FRET between CT and Thy-1, suggesting that under our experimental conditions lipid raft clustering is not sufficient to account for the observed FRET between the TeNT H_C and Thy-1 (Kenworthy *et al.*, 2000).

Some BoNTs have also been found to bind to neuronal DIGs (Herreros and Schiavo, unpublished results), increasing the possibility that the interaction with lipid microdomains represents a general mechanism for the recruitment of TeNT and BoNTs to the neuronal membrane.

What is the physiological relevance of TeNT binding to lipid rafts? These lipid domains have been implicated in membrane trafficking and signaling (Simons and Ikonen, 1997; Simons and Toomre, 2000). Interestingly, lipid rafts have recently emerged as membrane domains essential for the binding and uptake of pathogens and virulence factors into cells (Dehio *et al.*, 1995; Baorto *et al.*, 1997; Orlandi and Fishman, 1998; Wolf *et al.*, 1998; Fivaz *et al.*, 1999; Gordon *et al.*, 1999; Parton and Lindsay, 1999; Ricci *et al.*, 2000; Shin *et al.*, 2000). A possible explanation for this common mechanism is that binding to lipid rafts promotes a local increase of the pathogen/toxin's concentration (Abrami and van der Goot, 1999), which is exploited by pore-forming and multivalent toxins (Fivaz *et al.*, 1999). TeNT binds to polysialogangliosides (Montecucco, 1986; Halpern and Neale, 1995), which are likely to be concentrated in lipid rafts as demonstrated for most sphingolipids (Prinetti *et al.*, 2000). PI-PLC treatment does not affect TeNT H_C total binding, suggesting a role of gangliosides as primary low-affinity TeNT acceptors (Williamson *et al.*, 1999) that trap TeNT on the neuronal surface. Thus lipid rafts, by clustering polysialogangliosides, GPI-anchored binding proteins, and possibly other proteins involved in TeNT binding (Lazarovici and Yavin, 1986; Pierce *et al.*, 1986) could act as concentrating platforms for TeNT at the plasma membrane. This multiple binding within lipid rafts could confer to TeNT the extreme high affinity and neurospecificity observed *in vitro* and *in vivo*

(Montecucco, 1986). Furthermore, binding to raft components could target CNTs to *hot spots* on the plasma membrane that would give access to the signaling cascades emerging from these microdomains. In agreement with this view, it has been reported that TeNT binding stimulates phosphorylation of trk A and mitogen-activated protein kinase activity (Gil *et al.*, 2000, 2001). Similar to the signaling cascades triggered by NGF binding (Huang *et al.*, 1999), these events are likely to involve lipid rafts.

Depletion of the cellular cholesterol by treatment with cholesterol-sequestering drugs (MDCX, filipin) or detergents (saponin) disrupts lipid rafts (Neufeld *et al.*, 1996; Simons and Toomre, 2000) and consequently reduces the recovery of typical raft components in DIGs (Abrami and van der Goot, 1999; Huang *et al.*, 1999; Martens *et al.*, 2000). MDCX treatment of spinal cord cells causes the displacement of a discrete fraction of the bound TeNT H_C and other raft markers from DIGs. These findings suggest the existence of heterogeneous raft pools on the neuronal membrane (Madore *et al.*, 1999) that could be differently affected by changes in the cholesterol content. This moderate increase in the solubility of TeNT H_C correlates with a blockade of its internalization in MDCX-treated spinal cord cells, suggesting that a specialized subpool of lipid rafts is responsible for the productive binding and internalization of TeNT. In this regard, specialized lipid microdomains have been implicated in the internalization of proteins and pathogens (Parton *et al.*, 1994; Baorto *et al.*, 1997) and endocytosis of CT is inhibited by MDCX in nonneuronal cells (Orlandi and Fishman, 1998).

Strikingly, MDCX treatment causes the complete protection of VAMP from TeNT proteolytic activity, indicating an essential role of cholesterol in the internalization and intracellular trafficking of the toxin. Although alternative explanations are possible (see below), these findings could now explain some unique physiological features of TeNT and BoNTs. One of these properties is the apparent irreversibility of the binding of these toxins to the neuronal surface (Schmitt *et al.*, 1981; Habermann and Dreyer, 1986). The association of TeNT with lipid rafts would be characterized by a very low dissociation constant due to the multivalent binding nature of both lipid rafts (containing polysialogangliosides and protein acceptors) and TeNT. Structural analysis of TeNT H_C reveals the presence of multiple binding sites for oligosaccharides at the extreme carboxy terminus (Emsley *et al.*, 2000), which is necessary and sufficient for the interaction with the neuronal surface (Herreros *et al.*, 2000b). Moreover, TeNT has been described to form homodimers (Ledoux *et al.*, 1994; Herreros *et al.*, 2000a), a process that would further strengthen the multimeric nature of these interactions. In addition, the existence of at least two subpopulations of TeNT H_C that are differently affected by MDCX treatment could explain the observation of a nonproductive and productive neurotoxin binding (Daniels-Holgate and Dolly, 1996). The fraction of TeNT interacting with MDCX-sensitive lipid rafts could thus represent the productive subpool of TeNT, which is internalized, cleaves VAMP, and leads to the inhibition of neurotransmitter release.

Recently, MDCX has been reported to inhibit clathrin-dependent internalization of transferrin in nonneuronal cells (Rodal *et al.*, 1999; Subtil *et al.*, 1999). The blockade of TeNT internalization by cholesterol depletion could therefore sug-

gest that TeNT is internalized by a clathrin-dependent pathway in spinal cord neurons (Parton *et al.*, 1987). MDCX-treated and control cells showed a similar transferrin uptake, suggesting that the endosomal recycling pathway is not affected in our experimental conditions. However, we cannot completely rule out a possible inhibition of clathrin-mediated endocytosis in a subset of cells in our mixed spinal culture. Cholesterol-sequestering drugs also inhibit endocytosis from lipid rafts/caveolae in nonneuronal cells (Schnitzer *et al.*, 1994; Deckert *et al.*, 1996; Orlandi and Fishman, 1998) and GPI-anchored proteins can be internalized by clathrin-dependent and -independent routes (Makiya *et al.*, 1992; Mayor *et al.*, 1994; Parton *et al.*, 1994; Skretting *et al.*, 1999). Thus, TeNT internalization in neurons could involve both coated and uncoated pathways (Schwab and Thoenen, 1978; Parton *et al.*, 1987).

In conclusion, we demonstrate that TeNT interacts with GPI-anchored proteins and binds to lipid rafts. Cholesterol depletion causes the displacement of a subpool of TeNT from DIGs and blocks the internalization and intracellular activity of TeNT. This finding highlights a key role of cholesterol in the trafficking of TeNT in neurons.

ACKNOWLEDGMENTS

We are indebted to Dr. R. Morris (Guy's Hospital, London, United Kingdom) for kindly providing Thy-1 knockout mice, Dr. G. van der Goot (University of Geneva, Geneva, Switzerland) for aerolysin reagents, G. Lalli for purified MNs and valuable help in preparing spinal cord cultures, and Dr. P. Bastiaens for the FLIM microscope setup. We thank G. van der Goot, T. Iglesias, H. McNeil, C. Montecucco, R. Morris, G. Stenbeck, and members of our laboratory for critical reading of the manuscript and useful discussion. This work was supported by the Imperial Cancer Research Fund and the Human Frontier Science Program (to J.H.).

REFERENCES

- Aarts, L.H.J., Verkade, P., van Dalen, J.J.W., van Rozen, A.J., Gispen, W.H., Schrama, L.H., and Schotman, P. (1999). B-50/GAP-43 potentiates cytoskeletal reorganization in raft domains. *Mol. Cell. Neurosci.* 14, 85–97.
- Abrami, L., Fivaz, M., Glauser, P.-E., Parton, R.G., and van der Goot, F.G. (1998). A pore-forming toxin interacts with a GPI-anchored protein and causes vacuolation of the endoplasmic reticulum. *J. Cell Biol.* 140, 525–540.
- Abrami, L., and van der Goot, F.G. (1999). Plasma membrane microdomains act as concentration platforms to facilitate intoxication by aerolysin. *J. Cell Biol.* 147, 175–184.
- Ami, S., Keilbaugh, S.A., Ostermeyer, A.G., and Brown, D.A. (1998). Association of GAP-43 with detergent-resistant membranes requires two palmitoylated cysteine residues. *J. Biol. Chem.* 273, 28478–28485.
- Arribas, M., Blasi, J., Egea, G., Fariñas, I., Solsona, C., and Marsal, J. (1993). High resolution labeling of cholinergic nerve terminals using a specific fully active biotinylated botulinum neurotoxin type A. *J. Neurosci. Res.* 36, 635–645.
- Bakry, N.M., Kamata, Y., and Simpson, L.L. (1997). Expression of botulinum toxin binding sites in *Xenopus* oocytes. *Infect. Immun.* 65, 2225–2232.

- Baorto, D.M., Gao, Z., Malaviya, R., Dustin, M.L., van der Merwe, A., Lublin, D.M., and Abraham, S.N. (1997). Survival of FimH-expressing enterobacteria in macrophages relies on glycolipid traffic. *Nature* 389, 636–639.
- Bastiaens, P.I.H., and Jovin, T.M. (1998) Fluorescence resonance energy transfer microscopy. In: *Cell Biology: A Laboratory Manual*, ed. J.E. Celis, San Diego, CA: Academic Press, 136–146.
- Brown, D.A., and London, E. (2000). Structure, and function of sphingolipid-, and cholesterol-rich membrane rafts. *J. Biol. Chem.* 275, 17221–17224.
- Chamberlain, L.H., Burgoyne, R.D., and Gould, G.W. (2001). SNARE proteins are highly enriched in lipid rafts in PC12 cells. Implications for the spatial control of exocytosis. *Proc. Natl. Acad. Sci. USA* 98, 5619–5624.
- Daniels-Holgate, P.U., and Dolly, J.O. (1996). Productive and non-productive binding of botulinum neurotoxin A to motor nerve endings are distinguished by its heavy chain. *J. Neurosci. Res.* 44, 263–271.
- Deckert, M., Ticchioni, M., and Bernard, A. (1996). Endocytosis of GPI-anchored proteins in human lymphocytes: role of glycolipid-based domains, actin cytoskeleton, and protein kinases. *J. Cell Biol.* 133, 791–799.
- Dehio, C., Prevost, M.C., and Sansonetti, P.J. (1995). Invasion of epithelial cells by *Shigella flexneri* induces tyrosine phosphorylation of cortactin by a pp60c-src-mediated signaling pathway. *EMBO J.* 14, 2471–2482.
- Edelmann, L., Hanson, P.I., Chapman, E.R., and Jahn, R. (1995). Synaptobrevin binding to synaptophysin: a potential mechanism for controlling the exocytotic fusion machine. *EMBO J.* 14, 224–231.
- Emsley, P., Fotinou, C., Black, I., Fairweather, N.F., Charles, I.G., Watts, C., Hewitt, E., and Isaacs, N.W. (2000). The structures of the H_C fragment of tetanus toxin with carbohydrate subunit complexes provide insight into ganglioside binding. *J. Biol. Chem.* 275, 8889–8894.
- Evans, D.M., Williams, R.S., Shone, C.C., Hambleton, P., Melling, J., and Dolly, J.O. (1986). Botulinum neurotoxin type B. Its purification, radioiodination and interaction with rat-brain synaptosomal membranes. *Eur. J. Biochem.* 154, 409–416.
- Fivaz, M., Abrami, L., and van der Goot, F.G. (1999). Landing on lipid rafts. *Trends Cell Biol.* 9, 212–213.
- Friedrichson, T., and Kurzchalia, T.V. (1998). Micromains of GPI-anchored proteins in living cells revealed by crosslinking. *Nature* 394, 802–805.
- Gil, C., Chaib-Oukadour, I., Blasi, J., and Aguilera, J. (2001). H_C fragment of tetanus toxin activates protein kinase C isoforms, and phosphoproteins involved in signal transduction. *Biochem. J.* 356, 97–103.
- Gil, C., Chaib-Oukadour, I., Pelliccioni, P., and Aguilera, J. (2000). Activation of signal transduction pathways involving trkA, PLCγ1, PKC isoforms, and ERK-1/2 by tetanus toxin. *FEBS Lett.* 481, 177–182.
- Gordon, V.M., Nelson, K.M., Buckley, J.T., Stevens, V.L., Tweten, R.K., Elwood, P.C., and Leppla, S.H. (1999). *Clostridium septicum* alpha toxin uses glycosylphosphatidylinositol-anchored protein receptors. *J. Biol. Chem.* 274, 27274–27280.
- Grimmer, S., Iversen, T.-G., van Deurs, B., and Sandvig, K. (2000) Endosome to Golgi transport of ricin is regulated by cholesterol. *Mol. Biol. Cell* 11, 4205–4216.
- Habermann, E., and Dreyer, F. (1986). Clostridial neurotoxins: handling and action at the cellular and molecular level. *Curr. Top. Microbiol. Immunol.* 129, 93–179.
- Halpern, J.L., and Neale, E.A. (1995). Neurospecific binding, internalization, and retrograde axonal transport. *Curr. Top. Microbiol. Immunol.* 195, 221–241.
- Harder, T., Scheiffele, P., Verkade, P., and Simons, K. (1998). Lipid domain-structure of the plasma-membrane revealed by patching of membrane-components. *J. Cell Biol.* 141, 929–942.
- Herreros, J., Lalli, G., Montecucco, C., and Schiavo, G. (1999). Pathophysiological properties of clostridial neurotoxins. In: *The Comprehensive Source Book of Bacterial Protein Toxin*, ed. J.H. Freer and J.E. Alouf, London, UK: Academic Press, London, 202–228.
- Herreros, J., Lalli, G., Montecucco, C., and Schiavo, G. (2000a). Tetanus toxin fragment C binds to a protein present in neuronal cell lines, and motor neurons. *J. Neurochem.* 74, 1941–1950.
- Herreros, J., Lalli, G., and Schiavo, G. (2000b). C-Terminal half of tetanus toxin fragment C is sufficient for neuronal binding, and interaction with a putative protein receptor. *Biochem. J.* 347, 199–204.
- Hoekstra, D., and van Ijzendoorn, S.C.D. (2000). Lipid trafficking, and sorting. how cholesterol is filling gaps. *Curr. Opin. Cell Biol.* 12, 496–502.
- Hopkins, C.R., and Trowbridge, T. (1983). Internalization and processing of transferrin and the transferrin receptor in human carcinoma A431 cells. *J. Cell Biol.* 97, 508–521.
- Huang, C.-S., Zhou, J., Feng, A.K., Lynch, C.C., Klumperman, J., DeArmond, S.J., and Mobley, W.C. (1999). Nerve growth factor signaling in caveolae-like domains at the plasma membrane. *J. Biol. Chem.* 274, 36707–36714.
- Hueber, A.-O., Bernard, A.-M., Langlet-El Battari, C., Marguet, D., Massol, P., Foa, C., Brun, N., Garcia, S., Stewart, C., Pierres, M., and He, H.T. (1997) Thymocytes in Thy^{-/-} mice show augmented TCR signaling and impaired differentiation. *Curr. Biol.* 7, 705–708.
- Jacobson, K., and Dietrich, C. (1999). Looking at lipid rafts? *Trends Cell Biol.* 9, 87–91.
- Jeng, C.-J., McCarroll, S.A., Martin, T.F.J., Floor, E., Adams, J., Krantz, D., Butz, S., Edwards, R., and Schweitzer, E.S. (1998). Thy-1 is a component common to multiple populations of synaptic vesicles. *J. Cell Biol.* 140, 685–698.
- Kenworthy, A.K., Petranova, N., and Edidin, M. (2000). High-resolution FRET microscopy of cholera toxin B-subunit, and GPI-anchored proteins in cell plasma membranes. *Mol. Biol. Cell.* 11, 1645–1655.
- Lalli, G., Herreros, J., Osborne, S.L., Montecucco, C., Rossetto, O., and Schiavo, G. (1999). Functional characterization of tetanus and botulinum neurotoxins binding domains. *J. Cell Sci.* 112, 2715–2714.
- Laux, T., Fukami, K., Thelen, M., Golub, T., Frey, D., and Caroni, P. (2000). GAP43, MARCKS, and CAP23 modulate PI(4,5)P₂ at plasmalemmal rafts, and regulate cell cortex actin dynamics through a common mechanism. *J. Cell Biol.* 149, 1455–1471.
- Lazarovici, P., and Yavin, E. (1986). Affinity-purified tetanus neurotoxin interaction with synaptic membranes: properties of a protease-sensitive receptor component. *Biochemistry* 25, 7047–7054.
- Ledoux, D.N., Be, X.H., and Singh, B.R. (1994). Quaternary structure of botulinum and tetanus neurotoxins as probed by chemical cross-linking and native gel electrophoresis. *Toxicon* 32, 1095–1104.
- Li, L., and Singh, B.R. (1998). Isolation of synaptotagmin as a receptor for type A and type E botulinum neurotoxin and analysis of their comparative binding using a new microtiter plate assay. *J. Nat. Toxins* 7, 215–226.
- Lipardi, C., Nitsch, L., and Zurzolo, C. (2000). Detergent-insoluble GPI-anchored proteins are apically sorted in Fisher rat thyroid cells, but interference with cholesterol or sphingolipids differentially af-

- fects detergent insolubility, and apical sorting. *Mol. Biol. Cell.* *11*, 531–542.
- Madore, N., Smith, K.L., Graham, C.H., Jen, A., Brady, K., Hall, S., and Morris, R. (1999). Functionally different GPI proteins are organized in different domains on the neuronal surface. *EMBO J.* *18*, 6917–6926.
- Makiya, R., Thornell, L.E., and Stigbrand, T. (1992). Placental alkaline phosphatase, a GPI-anchored protein, is clustered in clathrin-coated vesicles. *Biochem. Biophys. Res. Commun.* *183*, 803–808.
- Martens, J.R., NavarroPolanco, R., Coppock, E.A., Nishiyama, A., Parshley, L., Grobaski, T.D., and Tamkun, M.M. (2000). Differential targeting of shaker-like potassium channels to lipid rafts. *J. Biol. Chem.* *275*, 7443–7446.
- Mayor, S., Rothberg, K.G., and Maxfield, F.R. (1994). Sequestration of GPI-anchored proteins in caveolae triggered by cross-linking. *Science* *264*, 1948–1951.
- Montecucco, C. (1986). How do tetanus and botulinum toxins bind to neuronal membranes? *Trends Biochem. Sci.* *11*, 315–317.
- Mukherjee, S., and Maxfield, F.R. (2000). Role of membrane organization, and membrane domains in endocytic lipid trafficking. *Traffic* *1*, 203–211.
- Neufeld, E.B., Cooney, A.M., Pitha, J., Dawidowicz, E.A., Dwyer, N.K., Pentchev, P.G., and Blanchette-Mackie, E.J. (1996). Intracellular trafficking of cholesterol monitored with a cyclodextrin. *J. Biol. Chem.* *271*, 21604–21613.
- Niemann, H., Blasi, J., and Jahn, R. (1994). Clostridial neurotoxins: new tools for dissecting exocytosis. *Trends Cell Biol.* *4*, 179–185.
- Nishiki, T., Kamata, Y., Nemoto, Y., Omori, A., Ito, T., Takahashi, M., and Kozaki, S. (1994). Identification of protein receptor for *Clostridium botulinum* type B neurotoxin in rat brain synaptosomes. *J. Biol. Chem.* *269*, 10498–10503.
- Nosten-Bertrand, M., Errington, M.L., Murphy, K.P.S.J., Tokugawa, Y., Barboni, E., Kozlova, E., Michalovich, D., Morris, R.G., Silver, J., Stewart, C.L., Bliss, T.V., and Morris, R.J. (1996). Normal spatial learning despite regional inhibition of LTP in mice lacking Thy-1. *Nature* *379*, 826–829.
- Orlandi, P.A., and Fishman, P.H. (1998). Filipin-dependent inhibition of cholera-toxin: evidence for toxin internalization and activation through caveolae-like domains. *J. Cell Biol.* *141*, 905–915.
- Parekh, R.B., Tse, A.G.D., Dweek, R.A., Williams, A.F., and Rademacher, T.W. (1987). Tissue-specific N-glycosylation, site-specific oligosaccharide patterns and lentil lectin recognition of rat Thy-1. *EMBO J.* *6*, 1233–1244.
- Parpura, V., Fang, Y., Basarsky, T., Jahn, R., and Haydon, P.G. (1995). Expression of synaptobrevin II, cellubrevin and syntaxin but not SNAP-25 in cultured astrocytes. *FEBS Lett.* *377*, 489–492.
- Parton, R.G., and Lindsay, M. (1999). Exploitation of major histocompatibility complex class I molecules and caveolae by simian virus 40. *Immunol. Rev.* *168*, 23–31.
- Parton, R.G., Ockleford, C.D., and Critchley, D.R. (1987). A study of the mechanism of internalization of tetanus toxin by primary mouse spinal cord cultures. *J. Neurochem.* *49*, 1057–1068.
- Parton, R.G., Joggerst, B., and Simons, K. (1994). Regulated internalization of caveolae. *J. Cell Biol.* *127*, 1199–1215.
- Pierce, E.J., Davison, M.D., Parton, R.G., Habig, W.H., and Critchley, D.R. (1986). Characterization of tetanus toxin binding to rat brain membranes. Evidence for a high-affinity proteinase-sensitive receptor. *Biochem. J.* *236*, 845–852.
- Prinetti, A., Chigorno, V., Tettamanti, G., and Sonnino, S. (2000). Sphingolipid-enriched membrane domains from rat cerebellar granule cells differentiated in cultures. *J. Biol. Chem.* *275*, 11658–11665.
- Ricci, V., Galmiche, A., Doye, A., Necchi, V., Solcia, E., and Boquet, P. (2000). High cell sensitivity to *Helicobacter pylori* VacA toxin depends on a GPI-anchored protein, and is not blocked by inhibition of the clathrin-mediated pathway of endocytosis. *Mol. Biol. Cell* *11*, 3897–3909.
- Rodal, S.K., Skretting, G., Garred, O., Vilhardt, F., van Deurs, B., and Sandvig, K. (1999). Extraction of cholesterol with methyl-beta-cyclodextrin perturbs formation of clathrin-coated endocytic vesicles. *Mol. Biol. Cell* *10*, 961–974.
- Schiavo, G., Ferrari, G., Rossetto, O., and Montecucco, C. (1991). Specific cross-linking of tetanus toxin to a protein of NGF-differentiated PC12 cells. *FEBS Lett.* *290*, 227–230.
- Schiavo, G., Matteoli, M., and Montecucco, C. (2000). Neurotoxins affecting neuroexocytosis. *Physiol. Rev.* *80*, 717–766.
- Schmitt, A., Dreyer, F., and John, C. (1981). At least three sequential steps are involved in the tetanus toxin-induced block of neuromuscular transmission. *Naunyn-Schmiedeberg's Arch. Pharmacol.* *317*, 326–330.
- Schnitzer, J.E., Oh, P., Pinney, E., and Allard, J. (1994). Filipin-sensitive caveolae-mediated transport in endothelium: reduced transcytosis, scavenger endocytosis, and capillary permeability of select macromolecules. *J. Cell Biol.* *127*, 1217–1232.
- Schroeder, R.J., Ahmed, S.N., Zhu, Y., London, E., and Brown, D.A. (1998). Cholesterol and sphingolipid enhance the Triton X-100 insolubility of glycosylphosphatidylinositol-anchored proteins by promoting the formation of detergent-insoluble ordered membrane domains. *J. Biol. Chem.* *273*, 1150–1157.
- Schwab, M.E., and Thoenen, H. (1978). Selective binding, uptake, and retrograde transport of tetanus toxin by nerve terminals in the rat iris. An electron microscope study using colloidal gold as a tracer. *J. Cell Biol.* *77*, 1–13.
- Selvin, P.R. (2000). The renaissance of fluorescence resonance energy transfer. *Nat. Struct. Biol.* *7*, 730–734.
- Shin, J.S., Gao, Z., and Abraham, S.N. (2000). Involvement of cellular caveolae in bacterial entry into mast cells. *Science* *289*, 785–788.
- Shogomori, H., and Futerman, A.H. (2001). Cholera toxin is found in detergent-insoluble rafts/domains at the cell surface of hippocampal neurons but is internalized via a raft-independent mechanism. *J. Biol. Chem.* *276*, 9182–9188.
- Simons, M., Friedrichson, T., Schulz, J.B., Pitto, M., Masserini, M., and Kurzchalia, T.V. (1999). Exogenous administration of gangliosides displaces GPI-anchored proteins from lipid microdomains in living cells. *Mol. Biol. Cell* *10*, 3187–3196.
- Simons, K., and Gruenberg, J. (2000). Jamming the endosomal system. Lipid rafts and lysosomal storage diseases. *Trends Cell Biol.* *10*, 459–462.
- Simons, K., and Ikonen, E. (1997). Functional rafts in cell membranes. *Nature* *387*, 569–572.
- Simons, K., and Toomre, D. (2000). Lipid rafts, and signal transduction. *Nat. Rev. Mol. Cell Biol.* *1*, 31–39.
- Skretting, G., Torgersen, M.L., van Deurs, B., and Sandvig, K. (1999). Endocytic mechanisms responsible for uptake of GPI-linked diphtheria toxin receptor. *J. Cell Sci.* *112*, 3899–3909.
- Squire, A., and Bastiaens, P. (1999). Three dimensional image restoration in fluorescence lifetime imaging microscopy. *J. Microsc.* *193*, 36–49.
- Subtil, A., Gaidarov, I., Kobylarz, K., Lampson, M.A., Keen, J.H., and McGraw, T.E. (1999). Acute cholesterol depletion inhibits clathrin-coated pit budding. *Proc. Natl. Acad. Sci. USA* *96*, 6775–6780.

- Tate, S.S., and Khadse, V. (1987). Intramolecular crosslinking of gamma-glutamyl transpeptidase. *Arch. Biochem. Biophys.* 255, 304–308.
- Tiveron, M.-C., Barboni, E., Pliego Rivero, F.B., Gormley, A.M., Seeley, P.J., Grosveld, F., and Morris, R. (1992). Selective inhibition of neurite outgrowth on mature astrocytes by Thy-1 glycoprotein. *Nature* 355, 745–748.
- Varma, R., and Mayor, S. (1998). GPI-anchored proteins are organized in submicron domains at the cell surface. *Nature* 394, 798–801.
- Williams, A.F., and Gagnon, J. (1982). Neuronal cell Thy-1 glycoprotein: homology with immunoglobulin. *Science* 216, 696–703.
- Williamson, L.C., Bateman, K.E., Clifford, J.C.M., and Neale, E.A. (1999). Neuronal sensitivity to tetanus toxin requires gangliosides. *J. Biol. Chem.* 274, 25173–25180.
- Wolf, A.A., Jobling, M.G., Wimermackin, S., Fergusonmaltzman, M., Madara, J.L., Holmes, R.K., and Lencer, W.I. (1998). Ganglioside structure dictates signal-transduction by cholera-toxin and association with caveolae-like membrane domains in polarized epithelia. *J. Cell Biol.* 141, 917–927.
- Yavin, E., and Nathan, A. (1986). Tetanus toxin receptors on nerve cells contain a trypsin-sensitive component. *Eur. J. Biochem.* 154, 403–407.