

1-1-1979

Lithologic Discrimination by Fourier Analysis of Image Texture

Michael Daily

Susan Williams

William D. Stromberg

Follow this and additional works at: http://docs.lib.purdue.edu/lars_symp

Daily, Michael; Williams, Susan; and Stromberg, William D, "Lithologic Discrimination by Fourier Analysis of Image Texture" (1979).
LARS Symposia. Paper 310.
http://docs.lib.purdue.edu/lars_symp/310

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 27 - 29, 1979

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana 47907 USA

IEEE Catalog No.
79CH1430-8 MPRSD

Copyright © 1979 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

LITHOLOGIC DISCRIMINATION BY FOURIER ANALYSIS OF IMAGE TEXTURE

MICHAEL DAILY, SUSAN WILLIAMS AND
WILLIAM D. STROMBERG

Jet Propulsion Laboratory/California
Institute of Technology

Drainage density is influenced by the erosion resistance of the underlying rocks. Ideal circumstances for lithologic discrimination by measurement of drainage network parameters are:

- 1) Geomorphic maturity
- 2) Relatively homogeneous rock structure
- 3) Slope as the only source of gray-tone modulation (e.g., in jungle)

An example of such a terrain would be a crystalline shield area in the wet tropics.

Our reconnaissance study involves the use of digital topographic tapes and aircraft and spacecraft (Seasat-A) radar data. The digital topographic data are first processed by SHADY 2, an algorithm that creates a scene illuminated at a selectable geometry according to a cosine scattering law. SHADY 2 has been modified to offer radar-like scattering behavior and to generate shadows.

The Fourier transforms of the images are computed and filtered with ten annular bandpass filters. The inverse transforms of the bandpass-filtered data are generated to produce images containing spatial features having a restricted range of dimensions. Spatial regions having high activity in a given frequency band will have a large variance in these images, so we generate a variance picture whose DN's are a measure of the local variance of the input image. These variance pictures (~10) are used as inputs in a supervised classifier.

Frequency slice pictures of the Klamath Mountains of Northern California show:

- 1) Only weak dependence on illumination geometry

- 2) Distinct clumping of high-activity areas
- 3) Differences in the location of the high-activity areas as a function of frequency