

## Local human impacts disrupt relationships between benthic reef assemblages and environmental predictors

Ford, Amanda; Jouffray, Jean-Baptiste; Norstrom, Albert V.; Moore, Bradley; Nugues, Maggy; Williams, Gareth J.; Bejarano, Sonia; Magron, Frank; Wild, Christian; Ferse, Sebastian

**Frontiers in Marine Science**

DOI:

[10.3389/fmars.2020.571115](https://doi.org/10.3389/fmars.2020.571115)

Published: 21/10/2020

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*

Ford, A., Jouffray, J-B., Norstrom, A. V., Moore, B., Nugues, M., Williams, G. J., Bejarano, S., Magron, F., Wild, C., & Ferse, S. (2020). Local human impacts disrupt relationships between benthic reef assemblages and environmental predictors. *Frontiers in Marine Science*, 7, [571115]. <https://doi.org/10.3389/fmars.2020.571115>

### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Local human impacts disrupt relationships between benthic reef assemblages and environmental predictors

Amanda K Ford<sup>1,2,3,4\*</sup>, Jouffray J-B<sup>3,5</sup>, Norström AV<sup>3</sup>, Moore BR<sup>6,7</sup>, Nugues MM<sup>8,9</sup>, Williams GJ<sup>10</sup>, Bejarano S<sup>11</sup>, Magron F<sup>12</sup>, Wild C<sup>2</sup>, Ferse SCA<sup>1,2</sup>

<sup>1</sup>Ecology Department, Leibniz Centre for Tropical Marine Research (ZMT), Bremen, Germany; <sup>2</sup>Marine Ecology Department, Faculty of Biology and Chemistry (FB2), University of Bremen, Bremen, Germany; <sup>3</sup>Stockholm Resilience Centre, Stockholm, Sweden; <sup>4</sup>School of Marine Studies, University of the South Pacific, Suva, Fiji; <sup>5</sup>Global Economic Dynamics and the Biosphere Academy Programme, Royal Swedish Academy of Sciences, Stockholm, Sweden; <sup>6</sup>National Institute of Water and Atmospheric Research, Nelson, New Zealand; <sup>7</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia; <sup>8</sup>EPHE, PSL Research University, UPVD-CNRS, USR3278 CRILOBE, France; <sup>9</sup>Labex Corail, CRILOBE, 98729 Moorea, French Polynesia; <sup>10</sup>School of Ocean Sciences, Bangor University, Anglesey LL59 5AB, UK; <sup>11</sup>Reef Systems Research Group, Leibniz Centre for Tropical Marine Research (ZMT), Bremen, Germany, <sup>12</sup>Pacific Community, Noumea, New Caledonia.

## \*Correspondence:

Amanda K Ford  
amandakford@hotmail.com

**Keywords: climate change, reef degradation, ecological reorganisation, ecological homogenisation, generalised additive models, model selection**

Manuscript length = 6124 words

Number of figures = 4

Number of tables = 2

Article language style: British English

## Abstract

Human activities are changing ecosystems at an unprecedented rate, yet large-scale studies into how local human impacts alter natural systems and interact with other aspects of global change are still lacking. Here we provide empirical evidence that local human impacts fundamentally alter relationships between ecological communities and environmental drivers. Using tropical coral reefs as a study system, we investigated the influence of contrasting levels of local human impact using a spatially extensive dataset spanning 62 outer reefs around inhabited Pacific islands. We tested how local human impacts (low *versus* high determined using a threshold of 25 people km<sup>-2</sup> reef) affected benthic community (i) structure, and (ii) relationships with environmental predictors using pre-defined models and model selection tools. Data on reef depth, benthic assemblages, and herbivorous fish communities were collected from field surveys. Additional data on thermal stress, storm exposure, and market gravity (a function of human population size and reef accessibility) were extracted from public repositories. Findings revealed that reefs subject to high local human impact were characterised by relatively more turf algae (>10% higher mean absolute coverage) and lower live coral cover (9% less mean absolute coverage) than reefs subject to low local human impact, but had similar macroalgal cover and coral morphological composition. Models based on spatio-physical predictors were significantly more accurate in explaining the variation of benthic assemblages at sites with low (mean adjusted-R<sup>2</sup> = 0.35) rather than high local human impact, where relationships became much weaker (mean adjusted-R<sup>2</sup> = 0.10). Model selection procedures also identified a distinct shift in the relative importance of different herbivorous fish functional groups in explaining benthic communities depending on the local human impact level. These results demonstrate that local human impacts alter natural systems and indicate that projecting climate change impacts may be particularly challenging at reefs close to higher human populations, where dependency and pressure on ecosystem services is highest.

# 94 1 Introduction

95  
96 Humans have become a dominant force of planetary change (Steffen et al., 2007).  
97 Ecosystems worldwide are being fundamentally altered by climate change impacts  
98 against a diverse backdrop of local anthropogenic stressors. Our ability to reliably  
99 predict the future configuration of affected ecosystems requires a thorough  
100 understanding of interactions between these different stressor types (Williams et al.,  
101 2019). Increasing evidence indicates that ecosystems are being reorganised, or  
102 homogenised, into stress-tolerant or opportunistic communities, leading to novel  
103 systems that differ from their previous state in terms of their dominant constituents  
104 (Graham et al., 2014; Morse et al., 2014). Quantifying ecological reorganisation over  
105 broad scales remains challenging due to coarse taxonomic resolution inherent to  
106 large datasets that are necessary to address this topic. Nonetheless, this  
107 phenomenon may significantly alter a system's relationship with various  
108 environmental and climate change-related drivers (Côté and Darling, 2010; Williams  
109 et al., 2015a). Understanding what role local human impacts play in driving  
110 ecological reorganisation and changing a system's relationship with its biophysical  
111 environment is thus pivotal to improving predictive models and informing local  
112 management (Robinson et al., 2018; Jouffray et al., 2019; Henderson et al., 2020).

113  
114 Inherent high biodiversity and the presence of multiple stressors acting at local and  
115 global scales make tropical coral reefs a unique system to examine how local human  
116 impacts drive ecological states (Hoegh-Guldberg et al., 2007; Ban et al., 2014;  
117 Norström et al., 2016; Darling et al., 2019). Locally, rapidly expanding coastal  
118 development, sewage input, and agricultural practices are reducing water quality,  
119 whilst the modernisation of fishing gear and expedited market access are facilitating  
120 overexploitation of coastal fisheries resources (Fabricius et al., 2005; Brewer et al.,  
121 2012; Hamilton et al., 2012). Globally, the intensity and frequency of thermal  
122 anomalies, unusual weather patterns, and destructive storm events are increasing  
123 (Gattuso et al., 2015; van Hooijdonk et al., 2016; IPCC, 2019), and recovery windows  
124 between stress events are narrowing (Riegl et al., 2013; Hughes et al., 2018). The  
125 productivity and provision of ecosystem services (e.g. Woodhead et al., 2019) of  
126 coral reef benthic communities differ depending on their composition (Ferrario et al.,  
127 2014; Rogers et al., 2018), underlining the importance of quantifying how  
128 communities are affected by global change. The widespread negative effects of  
129 climate change-related stressors are relatively well-understood. However, despite  
130 more than half the world's coral reefs being located within 30 minutes travel time  
131 from human populations (Maire et al., 2016), the role of local stressors remains  
132 somewhat contested. These local stressors can range greatly in their intensity  
133 depending on inherent factors such as human population density and the level of  
134 exploitation, such as reliance on marine resources.

135  
136 Higher human population densities have been linked to reductions in reef fish  
137 biomass and coral cover, and to increases in fleshy (turf and macro-) algal cover  
138 (Sandin et al., 2008; Williams et al., 2015b; Heenan et al., 2016; Smith et al., 2016).  
139 However, coarse taxonomic resolution (inherent to large-scale datasets) can lead to  
140 the conclusion that there is no link between local stressors and reef condition (e.g.  
141 Bruno and Valdivia, 2016). This may be a particular issue in regions such as the  
142 tropical Pacific where species diversity is exceptionally high and fleshy macroalgae,

143 frequently stated as principal alternative organisms on degraded reefs, do not play  
144 such a dominant role in benthic dynamics compared, for instance, to the Caribbean  
145 region (Roff and Mumby, 2012). Furthermore, reefs are increasingly existing in a  
146 partially degraded condition between distinct regimes (Mumby, 2017), likely having  
147 undergone ecological homogenisation resulting from the non-random removal of  
148 species with particular traits in response to environmental factors (McWilliam et al.,  
149 2020). In this case, coral cover can remain moderate but comprises a less diverse  
150 community of stress-tolerant and opportunistic types (Côté and Darling, 2010; van  
151 Woesik et al., 2011; Riegl et al., 2013). Ecological homogenisation is visible across  
152 reef habitat types, with inshore reefs that are naturally exposed to a more  
153 challenging environment (e.g. in terms of light, temperature, and sediment input)  
154 favouring a smaller species pool of stress-resistant corals compared to nearby outer  
155 reef habitats (Rogers, 1990; Browne et al., 2013; Williams et al., 2013; Schoepf et  
156 al., 2015; Morgan et al., 2016).

157

158 A recent study demonstrated that whilst sea surface temperature (SST), chlorophyll  
159 concentration, and wave energy have strong power in predicting benthic  
160 assemblages at remote reefs, this predictive power is lost or the relationships  
161 fundamentally altered at reefs closer to human populations (Williams et al., 2015a).  
162 Considering the dominant role of humans in shaping ecosystems, factors associated  
163 with local anthropogenic impacts may have overtaken biophysical drivers in  
164 structuring these altered reefs. It has also been postulated that reorganisation  
165 towards a stress-tolerant coral community could increase resilience to climate  
166 change, assuming co-tolerance between local and climate change-related stressors  
167 (Côté and Darling, 2010), in line with the concept of 'intrinsic resistance' (Darling and  
168 Côté, 2018). Indeed, coral richness does not translate into higher resilience to  
169 disturbances (Zhang et al., 2014). Better understanding and accounting for the role  
170 that local human impacts play in shaping benthic communities and their relationships  
171 with environmental drivers is important for developing theories, designing  
172 experiments, setting baselines, informing management, as well as optimising large-  
173 scale spatial predictive models for coral reef futures.

174

175 Here, we investigate how local human impacts affect coral reef benthic communities  
176 in the tropical Pacific. We start by classifying 62 island sites into two levels of local  
177 human impact (low *versus* high) using a set threshold of human population density  
178 informed by previous work (i.e. D'agata et al., 2014). We then ask whether the level  
179 of local human impact influences benthic community structure or the relationship  
180 between benthic community structure and spatio-physical explanatory drivers. Lastly,  
181 we examine the relative roles of ten biophysical parameters as drivers of benthic  
182 community structure under low *versus* high local human impacts. We hypothesised  
183 that spatio-physical drivers would be more related to benthic community structure on  
184 reefs exposed to low local human impacts compared to those with high local human  
185 impacts, where we expect human-associated factors to have become more  
186 dominant. This approach allowed us to develop on the findings of Williams et al.  
187 (2015a) by testing whether decoupling between reefs and biophysical drivers in the  
188 presence of humans was also detectable between different levels of local human  
189 impact.

190

191

## 2 Material and methods

### 2.1 Study area and sampling design

This study intended to build on work from Williams et al. (2015a) by assessing in more detail how local population density, rather than human presence/absence, potentially decouples the relationship between reef benthos and larger-scale environmental drivers. To do so, we utilised a large-scale dataset that is unique in having both site-level ecological (fish and benthic) and socio-economic data (in particular human density per reef area at the site level), which is missing from other datasets that rely on global socioeconomic estimates for human population density, or in some cases district-level surveys. Fish and benthic communities at 62 reefs within 17 different Pacific Island countries and territories were surveyed once between 2003–2008 (see Supplementary Figure 1 for map) as part of the Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish) under the auspices of the Pacific Community. Importantly, all reefs were in close vicinity to, and used by, coastal human communities across a large range of intensities (e.g. relative human density ranged from 1.3–1705 humans km<sup>-2</sup> reef). For site disturbance history, sampling dates and detailed sampling methodology, refer to Supplementary Table 1 and Pinca et al. (2010). Though it is important to acknowledge that the sampling program was not originally designed in a way to address macroecological questions or aggregate beyond the state/territory level, we did our best to account for the shortcomings in the design by including additional information such as observer bias (see section 2.3 for details on environmental parameters included). Furthermore, while data may not be representative of the current-day scenario, it is the trends that are important for this study. Accordingly, we are confident that despite some inevitable compromising, this gave us the best possible dataset to look at the effects of site-specific human density.

### 2.2 Field surveys

Underwater surveys covered outer (fore-) reefs, with on average nine (n = 3–47) joint fish-benthic 50 m transects measured at each site. Transect data were pooled within each site. Reef fish communities were measured using the variable distance-sampling underwater visual census method along transects (described in Labrosse et al., 2002). Data on abundance and size were recorded to species-level for herbivorous fish. Counts were converted to biomass (g m<sup>-2</sup>) from established length-weight relationships (Kulbicki et al., 2005). Benthic cover data was obtained using the medium-scale approach described by Clua et al., (2006). This method is based on a semi-quantitative description of ten 25 m<sup>2</sup> (5 x 5 m) quadrats laid down on each side of the 50 m transect (i.e. 20 replicate quadrats / 500 m<sup>2</sup> per transect). Surveyors first recorded abiotic and live coral substrates, i.e. sand, rubble, rocky slab, boulders, and hard coral - live, bleached, and long dead, with live coral divided into broad morphologies (e.g. branching, encrusting, massive). Each component was quickly estimated using a semi-quantitative scale ranging from 0 to 100% per quadrat, in units of 5%. Secondly, benthic groups (e.g. macroalgae – inclusive of calcified and fleshy types, turf algae, crustose coralline algae [CCA], sponges, cyanobacteria) growing over abiotic substrate such as long dead coral were recorded using the same semi-quantitative scale (Table 1).

241 Fishing grounds were initially delineated from information given by local fishers and  
 242 quantified from satellite interpretations (similar to methods in Close and Hall, 2006;  
 243 Léopold et al., 2014). Total reef area (km<sup>2</sup>) within each fishing ground was then  
 244 derived from reef areas quantified by the Millennium Coral Reef Mapping Project  
 245 from satellite images (Andréfouët et al., 2006). Socioeconomic assessments  
 246 determined total population within communities with access to the fishing ground,  
 247 allowing subsequent calculation of human population relative to reef area (referred to  
 248 as 'relative human density'). Finfish landings for each site, determined from  
 249 interviews with fishers, were extrapolated to total finfish catch per year per reef area  
 250 ('relative fishing pressure') (see Pinca et al., 2010).

251  
 252 **Table 1** Benthic variables included in the models and their ranges.  
 253

Variable	Description	Range (%)
live hard coral	mean cover (%) of live hard coral	6.1–65.1
dead hard coral	mean cover (%) of dead hard coral; including rubble, boulders, and pavement. Dead hard coral may also include biotic groups (i.e. growing over the dead coral), which are further classified as additional categories below*	5.5–61.2
branching morphologies	proportion (%) of branching coral morphologies within live hard coral community	0.1–85.6
encrusting morphologies	proportion (%) of encrusting coral morphologies within live hard coral community	2.8–72.9
massive morphologies	proportion (%) of massive coral morphologies within live hard coral community	0.1–60.6
CCA	mean cover (%) of crustose coralline algae	0.2–43.5
macroalgae	mean cover (%) of macroalgae; inclusive of calcified and fleshy types	0.0–31.2
turf algae	mean cover (%) of turf algae	0.0–45.6

254  
 255 *\*not all categories are mutually exclusive – abiotic (e.g. dead coral, sand) and live*  
 256 *coral substrates were recorded up to 100%, and biotic cover (e.g. algal groups) was*  
 257 *recorded separately up to 100% (i.e. sum of substrate and biotic cover ≠100%).*  
 258

259  
 260 **2.3 Data analyses**

261  
 262 **2.3.1 Response variables**

263 As response variables, we selected the main benthic groups (i.e. those with a mean  
 264 composition >10% of benthic community): dead coral (incorporating long dead coral,

265 rubble, boulders, and pavement), live hard coral, and algal groups (differentiating turf  
266 algae, macroalgae, and CCA). We also assessed the proportional representation  
267 (within the live hard coral community) of the three most commonly observed  
268 morphological groups: branching, encrusting and massive morphologies. Pairwise  
269 relationship tests (*corvif* function – Zuur et al., 2009) established no collinearity  
270 among the response variables (correlation;  $R^2 < 0.5$ ). Table 1 provides an overview  
271 of the benthic variables and their ranges.

272

### 273 **2.3.2 Model predictors**

274 We had to make a feasible choice of how to determine low *versus* high impact sites.  
275 Rather than choosing a completely arbitrary threshold, we selected one that was  
276 informed by previous findings by D’agata et al., (2014) – using boosted regression  
277 trees for the same dataset, the authors identified 25 people km<sup>-2</sup> reef to be the  
278 threshold after which taxonomic diversity of parrotfishes significantly declined. We  
279 then ran a sensitivity analysis to demonstrate how robust our findings were (see  
280 section 2.3.4). To compare reefs exposed to different disturbance regimes, we thus  
281 categorised all sites into those subject to low (i.e. < 25 people km<sup>-2</sup> reef; n = 29) and  
282 high (i.e. > 25 people km<sup>-2</sup> reef; n = 33) local human impact. Relative human density  
283 correlates with relative exploitation – i.e. relative human density was collinear ( $R^2 =$   
284 0.8) with fishing pressure (tonnes fish km<sup>-2</sup> reef year<sup>-1</sup>; Table 2). This threshold was  
285 also a median point and allowed similar sample sizes in each impact level (see  
286 Supplementary Figure 2). Relative human density also showed weak positive  
287 correlation with market gravity – an index combining the population size of nearby  
288 human settlements and their accessibility to reefs (Cinner and Maire, 2018; Cinner et  
289 al., 2018). However, we decided to base our study on relative human density as we  
290 had unique site-level data and we see this metric to be more directly linked to  
291 benthic communities in terms of human density-dependent sewage and agricultural  
292 run-off in addition to subsistence and artisanal fishing.

293

294 The selected biophysical predictors included a variety of factors that were either  
295 collected during field surveys or extracted from public data repositories (Table 2).  
296 Due to inherent differences in coral reefs across latitudinal scales (e.g. Hughes et al.,  
297 1999; Harriott and Banks, 2002), latitude was represented by degree distance from  
298 the equator without differentiating between north and south (0–23.9°). Degree  
299 heating weeks (DHW) data were extracted from the NOAA Coral Reef Thermal  
300 Anomaly Database (CoRTAD version 4 - Casey et al., 2012)). Storm exposure was  
301 quantified from the NOAA IBTrACS-WMO data (Knapp et al., 2010a, 2010b) within  
302 ArcMAP 10.4 (ESRI, 2011), where the number of storms (categories 1–5 on the  
303 Saffir-Simpson Hurricane Scale) passing within a 50 km radius of each site  
304 (Behrmann projection) was extracted. Storm exposure and DHW data were confined  
305 to 12 years prior to each respective site’s survey date based on the premise that  
306 remote reefs can recover from acute disturbances within this timeframe (Sheppard et  
307 al., 2008; Gilmour et al., 2013). Reef depth, estimated during field surveys, was  
308 averaged over all transects at each site. Island relief refers to each site’s  
309 geomorphology, and was classified into three categories: atoll, low-lying island, and  
310 high island based on available information (see Supplementary Table 1 for  
311 references), and authors’ knowledge. Island relief was included as a predictor due to  
312 its known influence on coral reef benthic and fish communities (Donaldson, 2002;  
313 Houk et al., 2015). Herbivorous fish from selected families (e.g. excluding



314 herbivorous damselfish) encountered during visual surveys were classified into  
315 functional groups according to Heenan et al. (2016) (see Supplementary Table 2).  
316 Biomasses ( $\text{g m}^{-2}$ ) of the following functional groups were then incorporated as  
317 predictors: browsers, grazers, detritivores, scrapers and small excavators, and large  
318 excavators and bioeroders. Market gravity (Cinner and Maire, 2018) was extracted  
319 for each site in QGIS (QGIS Development Team, 2019) and was incorporated as a  
320 continuous predictor.

321

322 Prior to model fitting, paired plots were assessed for collinearity between model  
323 terms. Strong collinearity ( $R^2 > -0.9$ ) between latitude and DHW precluded their joint  
324 inclusion in subsequent models, and consequently latitude was selected because of  
325 its complete reef-specific dataset (DHW data limited to  $n = 55$  sites). Multi-collinearity  
326 was also then tested using the generalised variance inflation factor (*GVIF*) function  
327 (*car* package - Fox and Weisberg, 2019) where values  $>3$  suggest collinearity – as a  
328 result longitude was excluded from all models and the joint inclusion of browsers and  
329 scrapers was prohibited (i.e. individual best-fit models were constrained to contain  
330 only one or the other).

331

### 332 **2.3.3 Statistical models**

333 All statistical analyses were performed in R version 3.6.1 (R Development Core  
334 Team, 2019). Differences in benthic community structure between reefs exposed to  
335 low *versus* high local human impact level were tested using *t*-tests with appropriate  
336 variance structures depending on homogeneity of variance test outcomes. Due to  
337 surveyor discrepancies in recording turf algal cover, we created a random effect  
338 (*bias\_score*) to be incorporated within turf algae models (see Supplementary Figure  
339 3 for details). No surveyor-related discrepancies were evident for other benthic  
340 groups (see Supplementary Figure 3). To test how turf algal cover differed across  
341 the two local human impact levels we thus used a linear mixed effects model  
342 incorporating *bias\_score* as a random effect. All hereon described models were run  
343 separately for sites predetermined to be exposed to either low or high local human  
344 impacts to explicitly test for decoupling of abiotic and biotic predictors under different  
345 disturbance regimes.

346

347 To test whether the level of local human impact influences the relationship between  
348 benthic community structure and spatio-physical explanatory drivers, we developed  
349 a 'spatio-physical' model (i.e. focusing on spatial and physical predictors only) that  
350 included storm exposure, reef depth, latitude, and island relief. To account for non-  
351 linear relationships between response variables and predictors, we applied  
352 generalised additive mixed effects models (GAMM) using the (Wood and Scheipl,  
353 2014) and *lme4* packages. To account for possible spatial autocorrelation, ten  
354 unique island clusters (*cluster*) were identified and incorporated into models as a  
355 random effect (for details see Supplementary Figure 4). For turf algae models,  
356 *bias\_score* was additionally included as a random effect. To avoid overfitting, the  
357 number of knots within models was limited to four. We retained the adjusted- $R^2$  (Adj-  
358  $R^2$ ) values from the model output to quantify each model's explanatory power. Adj- $R^2$   
359 values for each response variable were then compared (by paired *t*-tests) to test  
360 overall differences in model performance in explaining benthic community structure  
361 under the two local human impact levels.

362

363 **Table 2** Predictor descriptions and ranges at outer reefs. <sup>a</sup> predictor incorporated in  
 364 'spatio-physical' models; <sup>b</sup> predictor used in model selection procedures. For sources  
 365 of data, see main text.  
 366

Predictor	Description	Range
reef depth <sup>a,b</sup>	mean depth (m) of transects	3.9–10.5
degree heating weeks (DHW)	measure of cumulative thermal stress – sum of previous 12 weeks where thermal stress anomaly $\geq 1^{\circ}\text{C}$ ; value averaged over 12 years preceding survey; negatively collinear ( $R^2 = -0.9$ ) with latitude; only available for $n = 55$ sites	0.6–3.5
latitude <sup>a,b</sup>	degrees ( $^{\circ}$ ) distance from equator (absolute value). Negatively collinear ( $R^2 = -0.9$ ) with DHW	0.0–23.9
longitude	degrees ( $^{\circ}$ ) longitude on continuous scale (i.e. -175 counted as 185), included to account for distance from the Coral Triangle biodiversity hotspot	134.3–214.2 (i.e. -145.8)
storm exposure <sup>a,b</sup>	total number of storms (cat. 1 to 5 on the Saffir-Simpson Hurricane Scale) passing within 50 km of site within previous 12 years	0–14
relief <sup>a,b</sup>	3 classifications: 1 = atoll; 2 = low-lying island; 3 = high island	
browser biomass <sup>b</sup>	biomass ( $\text{g m}^{-2}$ ) of browsers	0.1–58.5
detritivore biomass <sup>b</sup>	biomass ( $\text{g m}^{-2}$ ) of detritivores	0.5–62.8
excavator biomass <sup>b</sup>	biomass ( $\text{g m}^{-2}$ ) of large excavators and bioeroders	0.0–369.1
grazer biomass <sup>b</sup>	biomass ( $\text{g m}^{-2}$ ) of grazers	1.1–161.0
scraper biomass <sup>b</sup>	biomass ( $\text{g m}^{-2}$ ) of scrapers and small excavators	1.9–134.3
relative human density	number of people within communities of the primary/customary resource users (living adjacent to/accessing fishing grounds within the reef area) related to total reef area ( $\text{people km}^{-2}$ reef), positively collinear ( $R^2 = 0.7$ ) with relative fishing pressure. Used to determine local human impact level	1.3–1705
relative fishing pressure	annual reef finfish catch (tonnes) $\text{km}^{-2}$ reef year <sup>-1</sup> ; positively collinear ( $R^2 = 0.8$ ) with relative human density	0.1–78.2

market gravity<sup>b</sup>

index that combines human population  
size and reef accessibility

0–1140

367

368 Lastly, to examine the relative roles of ten biophysical parameters as predictors of  
369 benthic community structure under low versus high local human impacts, we applied  
370 model selection techniques using the *MuMIn* package (Barton, 2016). From an initial  
371 model containing ten biophysical predictors (i.e. those included in the ‘spatio-  
372 physical model’ as well as biomass of herbivorous fish functional groups, and market  
373 gravity – see Table 2), the dredge function was used to run all possible predictor  
374 combinations and rank models from best to worst based on Akaike weight. The  
375 function also returns a value between 0 and 1 for each predictor that reflects its  
376 relative importance (RI), representing the total Akaike weight of all models containing  
377 that predictor (i.e. higher values correspond to greater RI). Output models were  
378 restricted to comprising a maximum of four predictors. One sample (Niue) was  
379 removed from the model selection procedure due to a lack of data for market gravity  
380 (i.e.  $n = 61$ ). Model selection was run separately for sites with low and high levels of  
381 local human impact, and all models incorporated the *uGamm* wrapper function to  
382 allow the inclusion of random effects consistent with ‘spatio-physical’ model  
383 constructions. To assess incongruities between benthic communities exposed to  
384 different local human impact levels, we retained the ‘best-fit’ model structures (i.e. all  
385 predictors included in models with Akaike weight  $>0.05$ ) for each benthic response  
386 variable, as well as the RI of individual predictors.

387

#### 388 **2.3.4 Assumptions and sensitivity tests**

389 All response variables were logit-transformed (appropriate for percentage data -  
390 Warton and Hui, 2011) using the *car* package, following adjustment using each  
391 respective variable’s minimum value  $>0$ . All model predictors were standardised (z-  
392 scores) to allow comparisons between predictors with largely varying effect sizes  
393 and numeric values (Zuur et al., 2009). Model residuals were checked for the  
394 violation of model assumptions using the *gam.check* function (see Supplementary  
395 Figure 5). As part of a sensitivity analysis, ‘spatio-physical’ models were additionally  
396 run using  $\pm 5$  and  $\pm 10$  humans  $\text{km}^{-2}$  reef as a threshold from which to categorise  
397 ‘low’ and ‘high’ impact sites, with consistent outcomes observed (see Supplementary  
398 Table 3). Furthermore, we repeated the same for a range of thresholds based on  
399 human density relative to outer reef area (as opposed to full reef area), again with  
400 consistent outcomes observed (see Supplementary Table 4).

401


402

### 403 **3 Results**

#### 404 **3.1 Benthic assemblages under contrasting local human impacts**

405 Benthic communities subject to high local human impact were associated with lower  
406 live hard coral cover (-9.2% mean absolute coverage; *t*-test:  $p = 0.002$ ; Figure 1A).  
407 However, the relative contribution of different coral morphologies within the live hard  
408 coral community did not differ significantly with local human impact level (Figure 1B).  
409 The only algal group that differed significantly depending on the local impact level  
410 was turf algae, which was higher under high local human impact (+10.7% mean  
411 absolute coverage; linear mixed effects model:  $p = 0.015$ ; Figure 1C).

412




413  
 414 **Figure 1** Boxplots detailing (A) benthic cover (%) of substrate types (from left to  
 415 right; live hard coral, dead hard coral), (B) the relative proportion (%) of the three  
 416 most dominant morphologies in the live hard coral community (branching,  
 417 encrusting, massive), and (C) benthic cover (%) of different algal groups (CCA,  
 418 macroalgae, turf algae), at sites determined to have low or high local impacts.  
 419 Asterisk (\*) indicates significant differences ( $p < 0.05$ ) in cover of the respective  
 420 benthic variable between local impact levels according to two-sample  $t$ -tests, or for  
 421 turf algae according to linear mixed effect models incorporating *bias\_score* as a  
 422 random effect (see section 2.3.2).

423  
 424

### 425 3.2 Predictive strength of 'spatio-physical' model

426 The pre-defined 'spatio-physical' model exhibited relatively high power in explaining  
 427 benthic assemblage variance at sites with low local human impacts (mean Adj-R<sup>2</sup> ±  
 428 SE; 0.35 ± 0.09; Figure 2), but model performance was severely compromised when  
 429 local human impacts were high (0.10 ± 0.04; paired  $t$ -test:  $p = 0.01$ ). When  
 430 considering only the substrate types and dominant morphological groups (i.e. without  
 431 the algal groups) the mean adjusted-R<sup>2</sup> for sites with low local human impacts  
 432 increases to 0.44 ± 0.06 but stays unchanged at sites with high local human impacts.

433 When local human impacts were low, 'spatio-physical' predictors explained a high  
 434 proportion of the variance of live hard coral ( $\text{Adj-R}^2 = 0.52$ ) and macroalgae ( $\text{Adj-R}^2 =$   
 435  $0.59$ ) cover, and the relative contribution of branching ( $\text{Adj-R}^2 = 0.64$ ) and massive  
 436 ( $\text{Adj-R}^2 = 0.50$ ) coral morphologies. Conversely, these variables were consistently  
 437 poorly explained when local human impacts were high ( $\text{Adj-R}^2 = 0.00, 0.26, 0.17,$   
 438  $0.10,$  respectively). No variance in turf algae was explained by this model for reefs at  
 439 either local human impact level. Significant differences in the power of the 'spatio-  
 440 physical' model in explaining benthic assemblage variance between 'low' and 'high'  
 441 impact sites held constant when the threshold was moved  $\pm 5$  and  $10 \text{ humans km}^{-2}$   
 442 reef (i.e. paired  $t$ -test:  $p < 0.05$ ; Supplementary Table 3).  
 443




444 **Figure 2** Explanatory power ( $\text{Adj-R}^2$ ) of the 'spatio-physical' model (including the  
 445 predictors storm exposure, reef depth, latitude, and island relief) in predicting  
 446 individual benthic responses (live hard coral, dead hard coral, the proportion of  
 447 branching, encrusting, and massive morphologies, and cover of algal groups CCA,  
 448 macroalgae, and turf algae). Bars are absent where the  $\text{Adj-R}^2$  value = 0.  
 449


450  
 451 **3.3 Relative importance of predictors**

452 Best-fit' models tailored for each individual benthic variable comprised distinctly  
 453 different predictors depending on the level of local human impact (Figure 3A). Two of  
 454 the predictors where discrepancies were most apparent were storm exposure and

455 grazer biomass, which were only selected for 'best-fit' models at sites with low and  
 456 high local human impacts, respectively. Similarly, reef depth was selected as part of  
 457 'best-fit' models for more response variables (live hard coral, branching  
 458 morphologies, and macroalgae) at sites with low local human impacts. Further  
 459 discrepancies under the two levels of local human impact were revealed by  
 460 comparing the mean RI of each predictor (Figure 3B,C), as the RI trends for  
 461 predictors almost reversed between low *versus* high local human impact sites. At  
 462 sites with low local human impacts, the individual predictors with the highest RI in  
 463 explaining benthic communities were storm exposure, scraper biomass, island relief,  
 464 followed by reef depth (Figure 3B). Contrastingly, apart from island relief which was  
 465 on average the most important predictor, storm exposure, scraper biomass, and reef  
 466 depth were among the least important predictors when local human impact was high  
 467 (Figure 3C). For benthic communities exposed to high local human impact, biomass  
 468 of grazers, detritivores, and browsers, as well as market gravity, ranked as the most  
 469 important predictors of benthic community structure.  
 470



471  
 472  
 473 **Figure 3 (A)** Outcome of model selection procedures: coloured boxes indicate that  
 474 the predictor was selected for 'best-fit' models (i.e. those with Akaike weight > 0.05)  
 475 at sites subject to low and/or high levels of local impacts. The overall mean relative  
 476 importance of each predictor (RI - based on total Akaike weight of all models  
 477 including that predictor) in explaining the benthic response variables is shown  
 478 separately for sites subject to **(B)** low and **(C)** high local impacts. Note: though model  
 479 selection was limited to four predictors per model, all predictors included in models  
 480 with Akaike weight >0.05 are shown meaning there may be more than four predictors  
 481 shown within one impact level for a response variable. Model outputs for 'best-fit'  
 482 models are available in Supplementary Table 5.  
 483  
 484



485  
 486 **Figure 4** Smoother plots of normalised residuals from generalised additive models  
 487 with 95% confidence intervals (shaded areas) to exemplify discrepancies in  
 488 response-predictor relationships between local impact levels, using storm exposure  
 489 as an example. The plots represent the explanatory power of storm exposure in  
 490 predicting (A) live hard coral cover, and the proportion of (B) branching, (C)  
 491 encrusting, and (D) massive morphologies, separated for sites with low *versus* high  
 492 local impacts. Refer to Supplementary Figure 6 for all predictor-response plots from  
 493 'best-fit' models.

494  
 495 Individual relationships of 'best-fit' predictors for each response variable further  
 496 emphasised discrepancies between benthic communities and environmental  
 497 predictors driven by local human impacts (for all plots see Supplementary Figure 6).  
 498 A clear example of this discrepancy can be seen in coral compositional changes with

499 increasing storm exposure. Whilst live hard coral cover remained unrelated to storm  
500 exposure under both local human impact levels (Figure 4A), the morphological  
501 composition was closely correlated with storm exposure at sites where local human  
502 impacts were low (Figure 4B–D). Specifically, with increasing storm exposure, the  
503 relative proportion of branching morphologies decreased significantly (Figure 4B),  
504 whereas encrusting (Figure 4C) and massive (Figure 4D) morphologies increased.  
505 No morphological changes with storm exposure were observed at sites with high  
506 local human impacts. Though these relationships appear primarily driven by few  
507 points at the higher end of storm exposure, they remained consistent when all reefs  
508 exposed to >10 storms were removed from the analysis – i.e. significant at ‘low’  
509 impact sites (branching coral  $p = 0.02$ ; encrusting coral  $p = 0.02$ ; massive coral  $p =$   
510  $0.03$ ), insignificant at ‘high’ impact sites (all morphological growth forms  $p > 0.05$ ),  
511 and no relationship for either impact for total live hard coral cover ( $p > 0.05$ ).

512

513

## 514 **4 Discussion**

515

516 Under increasing climate change-associated stressors and local anthropogenic  
517 influence (Burke et al., 2011; Gattuso et al., 2015; IPCC, 2019), it is particularly  
518 important for researchers and planners to maximise the ability of models to predict  
519 ecosystem futures to allow appropriate mitigation strategies to be implemented. This  
520 study indicates that the role of local human impacts in changing coral reef ecological  
521 communities and their responses to environmental drivers should be accounted for.  
522 The results revealed that local human impacts influenced both benthic community  
523 structure and relationships with biophysical predictors. Specifically, models based on  
524 spatio-physical predictors (i.e. reef depth, latitude, storm exposure, and relief)  
525 exhibited high power at explaining benthic assemblages under low local human  
526 impacts but were strongly compromised where local human impacts were high.  
527 Importantly, these outcomes remained similar when our threshold for human  
528 impacts, informed by previous work, was reduced or increased in the frame of a  
529 sensitivity analysis. Increasing sewage input, agricultural run-off, and sedimentation  
530 are potential changes associated with increasing human densities that reduce water  
531 quality and affect benthic communities (Fabricius, 2005; Fabricius et al., 2005; Ford  
532 et al., 2017). Furthermore, as fishing removes biomass of functionally important fish  
533 species, important top-down control of some benthic organisms is lost (Bellwood et  
534 al., 2004). These localised human impacts may homogenise benthic communities by  
535 driving ecological reorganisation that favours tolerant taxa (Darling et al., 2019). Our  
536 results indicate that this homogenisation may be occurring at Pacific Island reefs that  
537 are exposed to local human impacts, leading to novel systems that react  
538 fundamentally differently and unpredictably to environmental predictors compared to  
539 reefs less influenced by humans (Williams et al., 2015a). In turn, we expect that local  
540 human impacts will influence responses of coral reefs to climate change-related  
541 stressors, and that reefs close to human populations will require context-specific  
542 management approaches to maximise their future sustainability and associated  
543 critical ecosystem services (Moberg and Folke, 1999).

544

545 The results emphasise the variation in benthic assemblages that exists among  
546 Pacific Island reefs exposed to different levels of local human impact. Benthic  
547 communities were not restricted to distinct regimes dominated by either hard corals



548 or macroalgae, supporting previous studies from the Pacific (Bruno et al., 2009;  
549 Albert et al., 2012; Jouffray et al., 2015, 2019; Smith et al., 2016). In fact,  
550 macroalgae was the least common of all the algal groups, with turf algae and CCA  
551 more prominent on these outer reefs. Reefs at sites with higher local human impacts  
552 comprised significantly more turf algae and less live hard coral. Abundant and  
553 diverse outer reef coral communities (e.g. Ellis et al., 2017) have likely retained  
554 sensitive species, facilitating measurable differences (i.e. reductions in live hard  
555 coral cover) under higher local human impacts. Higher turf algae coverage at sites  
556 with more local human impacts provides further evidence that turf algae may  
557 become the dominant benthic group on degraded Pacific Island reefs (Jouffray et al.,  
558 2015; Smith et al., 2016; Tebbett and Bellwood, 2019). This may contrast with coral  
559 reef systems in the Caribbean, where macroalgae naturally play a more dominant  
560 role (Roff and Mumby, 2012). Furthermore, a signal of local human impact may have  
561 been detected if the data had distinguished macroalgae into fleshy/frondose and  
562 calcified types (e.g. Smith et al., 2016; Cannon et al., 2019). A similar link between  
563 human population size, hard coral cover, and turf algae has also recently been  
564 reported from sites in the Indian Ocean (Brown et al., 2017).

565  
566 The 'spatio-physical' models were weak in explaining the variance of benthic  
567 communities exposed to relatively more local human impacts. At these sites, model  
568 selection identified mostly local biotic controls or ecological features such as fish  
569 biomass and market gravity to be of highest relative importance. These results  
570 suggest that reefs altered by chronic local human impacts become decoupled from  
571 spatio-physical factors (Williams et al., 2015a) and become more related to factors  
572 associated with human activities (e.g. market gravity) or local ecological features  
573 (e.g. biomass of functional groups of fish). Interestingly, this outcome contrasts to  
574 recent findings by Robinson et al. (2018) who did not detect decoupling at inhabited  
575 *versus* uninhabited reefs. Our different outcomes for decoupling could in part be  
576 explained by Robinson et al. (2018) combining (i) hard corals and CCA, and (ii)  
577 fleshy macroalgae and turf algae, each of which we found to exhibit different  
578 responses to local impacts (i.e. with higher local impacts hard coral cover reduced  
579 whereas CCA remained unchanged, turf algae increased whereas macroalgae  
580 remained unchanged). Model selection indicated that the specific spatio-physical  
581 predictors whose influence were most disrupted by local human impacts were storm  
582 exposure and reef depth. Storms can have mixed effects, benefitting reefs by  
583 alleviating thermal stress during warmer summer months but also causing physical  
584 destruction, particularly to delicate branching coral morphologies, leading to a higher  
585 proportion of more robust massive morphologies (Heron et al., 2005; Manzello et al.,  
586 2007). Accordingly, when local human impacts were low, coral communities  
587 comprised relatively less branching and more encrusting and massive morphologies  
588 at sites subject to more frequent storms. However, relationships between storm  
589 exposure and benthic assemblages were only observed at sites classified as having  
590 low local human impacts. Returning to the concept that local human impacts drive  
591 ecological reorganisation, we would expect reefs with less local human impacts to  
592 harbour a great diversity of species and morphologies, thus allowing for greater  
593 levels of ecological reorganisation in response to a stormier environment (i.e.  
594 favouring more robust morphologies). We anticipate that Pacific reefs would more  
595 likely display this phenomenon than Caribbean reefs due to a significantly larger  
596 initial species pool, with greater response diversity and functional redundancy

597 affording a higher level of ecological insurance (Elmqvist et al., 2003; Bellwood et al.,  
598 2004; Nyström, 2006), and a loss of structurally-complex coral species throughout  
599 most of the Caribbean over past decades (Alvarez-Filip et al., 2009). It would be  
600 interesting to test whether storm exposure results in more conspicuous negative  
601 impacts on coral cover (e.g. Gardner et al., 2005) in less diverse regions due to the  
602 limited capacity for ecological reorganisation to a tolerant community, even in sites  
603 with minimal local human impact. In terms of reef depth, benthic communities at sites  
604 with low local human impacts also exhibited a higher level of depth-structuring  
605 (particularly in terms of coral cover and composition, consistent with Huston, 1985)  
606 compared to those with high local human impacts. Island relief was also identified as  
607 being a strong predictor of benthic community structure (despite being found to be a  
608 weak predictor on central-western Pacific reefs - Robinson et al., 2018), and  
609 interestingly this role was maintained regardless of local human impact level.  
610 Importantly, when considering latitude – collinear with DHW/cumulative thermal  
611 stress – as an individual predictor, our results neither contradict nor confirm previous  
612 observations that local impacts exacerbate the sensitivity of coral communities to  
613 thermal stress (Wiedenmann et al., 2013; Ellis et al., 2019), instead highlighting a  
614 large variance among sites.

615  
616 Thermal stress and the intensity of storms are projected to increase under future  
617 climate change scenarios (IPCC, 2019), with profound implications for coral reefs  
618 and adjacent ecosystems. Our findings suggest that while the effects of factors  
619 associated with climate change (e.g. storm exposure) on Pacific reef benthic  
620 assemblages may be reasonably well-predicted where local impacts are low, system  
621 responses become less predictable as local human impacts increase. In both marine  
622 and terrestrial systems, structural changes caused by local impacts have profoundly  
623 changed how ecosystems respond to natural stressors: for example, local stressors  
624 have affected how parts of Australia's Great Barrier Reef have recovered from recent  
625 climate change impacts (MacNeil et al., 2019; Mellin et al., 2019) and habitat  
626 fragmentation and modification have exacerbated recent impacts of tropical and  
627 temperate forest fires (Brando et al., 2014; Alencar et al., 2015; Taylor et al., 2016).  
628 Because the effects of storms are strongly dependent not only on their intensity, but  
629 also the extent of the fetch, their frequency, and intrinsic reef properties such as  
630 topography (Lugo et al., 2000; Heron et al., 2005), this study incorporated all  
631 recorded storms (category 1–5 on the Saffir-Simpson Hurricane Scale) passing  
632 within 50 km of each site. We thus cannot deduce benthic community responses to  
633 increasing storm *intensity*, which is projected to occur (IPCC, 2019).

634  
635 Functional groups of herbivorous fishes also emerged to be of contrasting relative  
636 importance depending on the level of local human impact, shifting from scrapers and  
637 small excavators at less impacted reefs to grazers at more impacted reefs. Browsers  
638 were of similar importance at reefs exposed to both local human impact levels,  
639 perhaps linked to the fact that their food source (macroalgae) remained constant  
640 regardless of local human impact level. Additional reasons for this could be that  
641 browsers have remained more resilient to fishing pressure than other herbivores, or  
642 that visual survey data does not accurately represent browser populations (for  
643 example some browser species are known to be particularly wary of divers –  
644 Kulbicki, 1998). Fish that act on turf algae and/or on surfaces available for coral  
645 settlement seem to be more sensitive to local human impacts, likely in response to

646 benthic community shifts. Scrapers and small excavators clear substrate for  
647 calcifiers, justifying their higher relative importance in models focused on less locally  
648 impacted reefs where live hard coral cover was higher and conditions for settlement  
649 and growth of juvenile corals were likely better (e.g. less nutrients, lower  
650 sedimentation). Contrastingly, grazers crop and maintain algal turfs, explaining their  
651 importance under higher local human impacts where turf algae were more dominant.  
652 These results align with findings in the Hawaiian Archipelago, where biomass of  
653 grazers and scrapers were the most important predictors of turf/macroalgal, and  
654 calcified regimes, respectively (Jouffray et al., 2015; see also Robinson et al., 2018).  
655 Though collinearity tests ruled out significant, potentially confounding relationships  
656 between biomass of different functional groups with individual abiotic predictors, it is  
657 important to acknowledge that fish communities themselves can be affected by  
658 various physical predictors (Williams et al., 2015b; Samoily et al., 2019), which  
659 could in turn influence benthic structure.

660

661 Other factors not included in these analyses are known to structure benthic  
662 assemblages, including chlorophyll, SSTs, and wave exposure (Gove et al., 2013,  
663 2015; Williams et al., 2013; Robinson et al., 2018; Darling et al., 2019). Remotely  
664 sensed chlorophyll data captures offshore productivity, but we expected land-based  
665 input to dominate many of these reefs which are close to land, while the survey  
666 design (sometimes around the periphery of small islands/atolls) made wave  
667 exposure challenging to quantify. Additionally, wave exposure, mean SSTs, and  
668 climatological ranges can be relatively well-captured by latitude in the Pacific (Gove  
669 et al., 2013). Also, although this study goes into more detail than many similar large-  
670 scale analyses by evaluating coral growth forms, it is still limited in its ability to  
671 quantify ecological reorganisation, which would require higher resolution data (at  
672 least family or genus). This limitation can be overcome by broad-scale surveys  
673 refining the level at which hard coral communities are recorded, which will become  
674 easier with improvements in automated software tools. These results do however  
675 emphasise that even broad morphological groupings (i.e. branching, massive,  
676 encrusting morphologies) provide pertinent information on ecological changes and  
677 can improve model performance compared to when overall hard coral cover is  
678 considered (Gove et al., 2015).

679

680 Importantly, the threshold used in this study for determining low and high impact,  
681 while informed by a previous study, was set *a priori* and thus does not allow defining  
682 a 'carrying capacity' of human density – this would require a different survey design  
683 and analytical approach, and should be pursued in future studies. We would  
684 however suggest that although this metric is most applicable to areas where  
685 customary resource use is common practice, we anticipate that this threshold (25  
686 humans km<sup>-2</sup> reef) is highly relevant across the Pacific region (i.e. given that it is a  
687 median point in this dataset for which sites were selected due to being regionally  
688 representative of fished areas). Furthermore, the study design was not originally  
689 meant to address macroecological questions but was nonetheless the best available  
690 to study the questions we were interested in (with incorporation of appropriate  
691 secondary data on local environmental context). Again, future studies should  
692 address this by appropriate designs that allow for large-scale comparability and  
693 collect both ecological and socio-economic local data. Our findings indicate that  
694 island relief is a factor that should be considered in designing regional sampling

695 programs with comparable sites. Finally, we should aim to obtain context-specific  
696 information on factors associated with human population density (e.g. sewage  
697 treatment presence, farming and associated fertiliser-usage) that strengthens our  
698 ability to predict benthic communities under various levels of human population  
699 density and improve its use as a proxy of local human impact.

700

701 Our findings contribute to a better understanding of the role of local human impacts  
702 on diverse ecosystems such as tropical coral reefs. The results provide empirical  
703 evidence that local human impacts drive conspicuous changes in benthic community  
704 relationships with environmental predictors, with indications of ecological  
705 reorganisation. Even if decisive steps are taken to reduce fossil fuel emissions, most  
706 reefs will suffer long-term degradation from the effects of climate change by 2050,  
707 and >75% of reefs will experience annual severe bleaching (Frieler et al., 2013; van  
708 Hoodonk et al., 2016). Our results show that local human impacts can lead to  
709 increasingly unpredictable relationships between benthic communities and their  
710 physical environment, and that overlooking their role could pave way to significant  
711 errors in future projections, potentially compromising mitigation efforts.

712

713

## 714 **5 Conflict of Interest**

715

716 The authors declare that the research was conducted in the absence of any  
717 commercial or financial relationships that could be construed as a potential conflict of  
718 interest.

719

720

## 721 **6 Author Contributions**

722

723 AF developed the study idea with close support from JBJ, AN, BM and SF. AF led  
724 the statistical analyses together with input from JBJ, GW and SF. BM and FM  
725 provided expertise on the original data. All authors provided support throughout the  
726 interpretation of the results and development of the manuscript. All authors approve  
727 this final version.

728

729

## 730 **7 Funding**

731

732 The COFish and PROCFish-C Programmes were funded by the European Union.  
733 AF, SB, and SF were funded by the (German) Federal Ministry of Education and  
734 Research (BMBF) through the “Nachwuchsgruppen Globaler Wandel 4 + 1”  
735 (REPICORE, grant number 01LN1303A).

736

737

## 738 **8 Acknowledgements**

739

740 We would like to thank the Pacific Community survey team, in particular Samasoni  
741 Sauni, Pierre Boblin, Ribanataake Awira, Laurent Vigliola, and Silvia Pinca, as well  
742 as the fisheries officers across the PICTs who assisted with logistics and data

743 collection. Data collection was made possible by the support of in-country Fisheries  
744 Divisions, as well as the chiefs and people of all local communities at the surveyed  
745 sites. The research reported in this paper contributes to the Programme on  
746 Ecosystem Change and Society ([www.pecs-science.org](http://www.pecs-science.org)).

747

748

## 749 **9 References**

750

751 Albert, S., Dunbabin, M., Skinner, M., Moore, B., and Grinham, A. (2012). Benthic shift in a Solomon  
752 Islands' lagoon: corals to cyanobacteria. *Proceedings of the 12th International Coral Reef*  
753 *Symposium, Cairns, Australia, 9-13 July 2012*, 1–5.

754 Alencar, A. A., Brando, P. M., Asner, G. P., and Putz, F. E. (2015). Landscape fragmentation, severe  
755 drought, and the new Amazon forest fire regime. *Ecological Applications* 25, 1493–1505.  
756 doi:10.1890/14-1528.1.

757 Alvarez-Filip, L., Dulvy, N. K., Gill, J. A., Côté, I. M., and Watkinson, A. R. (2009). Flattening of  
758 Caribbean coral reefs: region-wide declines in architectural complexity. *Proceedings of the*  
759 *Royal Society B: Biological Sciences* 276, 3019–3025. doi:10.1098/rspb.2009.0339.

760 Andréfouët, S., Muller-Karger, F. E., Robinson, J. A., Kranenburg, C. J., Torres-Pulliza, D., Spraggins, S.  
761 A., et al. (2006). Global assessment of modern coral reef extent and diversity for regional  
762 science and management applications: a view from space. *Proceedings of 10th International*  
763 *Coral Reef Symposium*, 1732–1745.

764 Ban, S. S., Graham, N. A. J., and Connolly, S. R. (2014). Evidence for multiple stressor interactions and  
765 effects on coral reefs. *Global Change Biology* 20, 681–697. doi:10.1111/gcb.12453.

766 Barton, K. (2016). *MuMIn: Multi-Model Inference. R package version 1.15.6.* [https://CRAN.R-](https://CRAN.R-project.org/package=MuMIn)  
767 [project.org/package=MuMIn](https://CRAN.R-project.org/package=MuMIn). Available at: <https://CRAN.R-project.org/package=MuMIn>.

768 Bellwood, D. R., Hughes, T. P., Folke, C., and Nyström, M. (2004). Confronting the coral reef crisis.  
769 *Nature* 429, 827–833. doi:10.1038/nature02691.

770 Brando, P. M., Balch, J. K., Nepstad, D. C., Morton, D. C., Putz, F. E., Coe, M. T., et al. (2014). Abrupt  
771 increases in Amazonian tree mortality due to drought–fire interactions. *PNAS* 111, 6347–  
772 6352. doi:10.1073/pnas.1305499111.

773 Brewer, T. D., Cinner, J. E., Fisher, R., Green, A., and Wilson, S. K. (2012). Market access, population  
774 density, and socioeconomic development explain diversity and functional group biomass of  
775 coral reef fish assemblages. *Global Environmental Change* 22, 399–406.  
776 doi:10.1016/j.gloenvcha.2012.01.006.

777 Brown, K. T., Bender-Champ, D., Bryant, D. E. P., Dove, S., and Hoegh-Guldberg, O. (2017). Human  
778 activities influence benthic community structure and the composition of the coral-algal  
779 interactions in the central Maldives. *Journal of Experimental Marine Biology and Ecology*  
780 497, 33–40. doi:10.1016/j.jembe.2017.09.006.

781 Browne, N. K., Smithers, S. G., and Perry, C. T. (2013). Spatial and temporal variations in turbidity on  
782 two inshore turbid reefs on the Great Barrier Reef, Australia. *Coral Reefs* 32, 195–210.  
783 doi:10.1007/s00338-012-0965-1.

- 784 Bruno, J. F., Sweatman, H., Precht, W. F., Selig, E. R., and Schutte, V. G. W. (2009). Assessing  
785 evidence of phase shifts from coral to macroalgal dominance on coral reefs. *Ecology* 90,  
786 1478–1484. doi:10.1890/08-1781.1.
- 787 Bruno, J. F., and Valdivia, A. (2016). Coral reef degradation is not correlated with local human  
788 population density. *Scientific Reports* 6, 29778. doi:10.1038/srep29778.
- 789 Burke, L. M., Reyta, K., Spalding, M., and Perry, A. (2011). *Reefs at Risk Revisited*. Washington D.C.:  
790 World Resources Institute.
- 791 Cannon, S. E., Donner, S. D., Fenner, D., and Beger, M. (2019). The relationship between macroalgae  
792 taxa and human disturbance on central Pacific coral reefs. *Marine Pollution Bulletin* 145,  
793 161–173. doi:10.1016/j.marpolbul.2019.05.024.
- 794 Casey, K. S., Selig, E. R., and Foti, G. (2012). The Coral Reef Temperature Anomaly Database  
795 (CoRTAD) Version 4 - Global, 4 km Sea Surface Temperature and Related Thermal Stress  
796 Metrics for 1981-10-31 to 2010-12-31 (NODC Accession 0087989). Version 2.2. National  
797 Oceanographic Data Center, NOAA. Dataset.
- 798 Cinner, J. E., Maire, E., Huchery, C., MacNeil, M. A., Graham, N. A. J., Mora, C., et al. (2018). Gravity  
799 of human impacts mediates coral reef conservation gains. *PNAS* 115, E6116–E6125.  
800 doi:10.1073/pnas.1708001115.
- 801 Cinner, J., and Maire, E. (2018). Global gravity of coral reefs spatial layer. James Cook University.  
802 (dataset). <http://dx.doi.org/10.4225/28/5a0e7b1b3cc0e>.
- 803 Close, C. H., and Hall, G. B. (2006). A GIS-based protocol for the collection and use of local knowledge  
804 in fisheries management planning. *J. Environ. Manage.* 78, 341–352.  
805 doi:10.1016/j.jenvman.2005.04.027.
- 806 Clua, E., Legendre, P., Vigliola, L., Magron, F., Kulbicki, M., Sarramegna, S., et al. (2006). Medium  
807 scale approach (MSA) for improved assessment of coral reef fish habitat. *Journal of*  
808 *Experimental Marine Biology and Ecology* 333, 219–230. doi:10.1016/j.jembe.2005.12.010.
- 809 Côté, I. M., and Darling, E. S. (2010). Rethinking Ecosystem Resilience in the Face of Climate Change.  
810 *PLoS Biol* 8. doi:10.1371/journal.pbio.1000438.
- 811 D’agata, S., Mouillot, D., Kulbicki, M., Andréfouët, S., Bellwood, D. R., Cinner, J. E., et al. (2014).  
812 Human-Mediated Loss of Phylogenetic and Functional Diversity in Coral Reef Fishes. *Current*  
813 *Biology* 24, 555–560. doi:10.1016/j.cub.2014.01.049.
- 814 Darling, E. S., and Côté, I. M. (2018). Seeking resilience in marine ecosystems. *Science* 359, 986–987.  
815 doi:10.1126/science.aas9852.
- 816 Darling, E. S., McClanahan, T. R., Maina, J., Gurney, G. G., Graham, N. A. J., Januchowski-Hartley, F.,  
817 et al. (2019). Social–environmental drivers inform strategic management of coral reefs in the  
818 Anthropocene. *Nature Ecology & Evolution* 3, 1341–1350. doi:10.1038/s41559-019-0953-8.
- 819 Donaldson, T. J. (2002). High Islands Versus Low Islands: A Comparison of Fish Faunal Composition of  
820 the Palau Islands. *Environmental Biology of Fishes* 65, 241–248.  
821 doi:10.1023/A:1020067931910.

- 822 Ellis, J., Anlauf, H., Kürten, S., Lozano-Cortés, D., Alsaffar, Z., Cúrdia, J., et al. (2017). Cross shelf  
823 benthic biodiversity patterns in the Southern Red Sea. *Scientific Reports* 7, 437.  
824 doi:10.1038/s41598-017-00507-y.
- 825 Ellis, J. I., Jamil, T., Anlauf, H., Coker, D. J., Curdia, J., Hewitt, J., et al. (2019). Multiple stressor effects  
826 on coral reef ecosystems. *Global Change Biology* 25, 4131–4146. doi:10.1111/gcb.14819.
- 827 Elmqvist, T., Folke, C., Nyström, M., Peterson, G., Bengtsson, J., Walker, B., et al. (2003). Response  
828 diversity, ecosystem change, and resilience. *Frontiers in Ecology and the Environment* 1,  
829 488–494. doi:10.1890/1540-9295(2003)001[0488:RDECAR]2.0.CO;2.
- 830 ESRI (2011). *ArcGIS Desktop: Release 10*. Redlands, California: Environmental Systems Research  
831 Institute.
- 832 Fabricius, K., De'ath, G., McCook, L., Turak, E., and Williams, D. McB. (2005). Changes in algal, coral  
833 and fish assemblages along water quality gradients on the inshore Great Barrier Reef.  
834 *Marine Pollution Bulletin* 51, 384–398. doi:10.1016/j.marpolbul.2004.10.041.
- 835 Fabricius, K. E. (2005). Effects of terrestrial runoff on the ecology of corals and coral reefs: review  
836 and synthesis. *Marine Pollution Bulletin* 50, 125–146. doi:10.1016/j.marpolbul.2004.11.028.
- 837 Ferrario, F., Beck, M. W., Storlazzi, C. D., Micheli, F., Shepard, C. C., and Airoidi, L. (2014). The  
838 effectiveness of coral reefs for coastal hazard risk reduction and adaptation. *Nature*  
839 *Communications* 5, 3794. doi:10.1038/ncomms4794.
- 840 Ford, A. K., Hoytema, N. V., Moore, B. R., Pandihau, L., Wild, C., and Ferse, S. C. A. (2017). High  
841 sedimentary oxygen consumption indicates that sewage input from small islands drives  
842 benthic community shifts on overfished reefs. *Environmental Conservation* 44, 405–411.  
843 doi:10.1017/S0376892917000054.
- 844 Fox, J., and Weisberg, S. (2019). *An R Companion to Applied Regression, Third Edition*. Thousand  
845 Oaks CA: Sage Publications Available at:  
846 <https://socialsciences.mcmaster.ca/jfox/Books/Companion/>.
- 847 Frieler, K., Meinshausen, M., Golly, A., Mengel, M., Lebek, K., Donner, S. D., et al. (2013). Limiting  
848 global warming to 2 °C is unlikely to save most coral reefs. *Nature Climate Change* 3, 165–  
849 170. doi:10.1038/nclimate1674.
- 850 Gardner, T. A., Côté, I. M., Gill, J. A., Grant, A., and Watkinson, A. R. (2005). Hurricanes and  
851 Caribbean Coral Reefs: Impacts, Recovery Patterns, and Role in Long-Term Decline. *Ecology*  
852 86, 174–184. doi:10.1890/04-0141.
- 853 Gattuso, J.-P., Magnan, A., Billé, R., Cheung, W. W. L., Howes, E. L., Joos, F., et al. (2015). Contrasting  
854 futures for ocean and society from different anthropogenic CO<sub>2</sub> emissions scenarios. *Science*  
855 349. doi:10.1126/science.aac4722.
- 856 Gilmour, J. P., Smith, L. D., Heyward, A. J., Baird, A. H., and Pratchett, M. S. (2013). Recovery of an  
857 isolated coral reef system following severe disturbance. *Science* 340, 69–71.  
858 doi:10.1126/science.1232310.

- 859 Gove, J. M., Williams, G. J., McManus, M. A., Clark, S. J., Ehses, J. S., and Wedding, L. M. (2015). Coral  
860 reef benthic regimes exhibit non-linear threshold responses to natural physical drivers.  
861 *Marine Ecology Progress Series* 522, 33–48. doi:10.3354/meps11118.
- 862 Gove, J. M., Williams, G. J., McManus, M. A., Heron, S. F., Sandin, S. A., Vetter, O. J., et al. (2013).  
863 Quantifying Climatological Ranges and Anomalies for Pacific Coral Reef Ecosystems. *PLOS*  
864 *ONE* 8, e61974. doi:10.1371/journal.pone.0061974.
- 865 Graham, N. A., Cinner, J. E., Norström, A. V., and Nyström, M. (2014). Coral reefs as novel  
866 ecosystems: embracing new futures. *Current Opinion in Environmental Sustainability* 7, 9–  
867 14. doi:10.1016/j.cosust.2013.11.023.
- 868 Hamilton, R. J., Giningele, M., Aswani, S., and Ecochard, J. L. (2012). Fishing in the dark-local  
869 knowledge, night spearfishing and spawning aggregations in the Western Solomon Islands.  
870 *Biological Conservation* 145, 246–257. doi:10.1016/j.biocon.2011.11.020.
- 871 Harriott, V., and Banks, S. (2002). Latitudinal variation in coral communities in eastern Australia: a  
872 qualitative biophysical model of factors regulating coral reefs. *Coral Reefs* 21, 83–94.  
873 doi:10.1007/s00338-001-0201-x.
- 874 Heenan, A., Hoey, A. S., Williams, G. J., and Williams, I. D. (2016). Natural bounds on herbivorous  
875 coral reef fishes. *Proceedings of the Royal Society B: Biological Sciences* 283, 20161716.  
876 doi:10.1098/rspb.2016.1716.
- 877 Henderson, C. J., Gilby, B. L., Schlacher, T. A., Connolly, R. M., Sheaves, M., Maxwell, P. S., et al.  
878 (2020). Landscape transformation alters functional diversity in coastal seascapes. *Ecography*  
879 43, 138–148. doi:10.1111/ecog.04504.
- 880 Heron, S., Morgan, J., Eakin, M., and Skirving, W. (2005). “Hurricanes and their effects on coral  
881 reefs,” in *Wilkinson, C. and Souter, D. (eds.), Status of Caribbean Coral Reefs after Bleaching*  
882 *and Hurricanes in 2005*. (Global Coral Reef Monitoring Network and Reef and Rainforest  
883 Research Centre), 31–36.
- 884 Hoegh-Guldberg, O., Mumby, P. J., Hooten, A. J., Steneck, R. S., Greenfield, P., Gomez, E., et al.  
885 (2007). Coral Reefs Under Rapid Climate Change and Ocean Acidification. *Science* 318, 1737–  
886 1742. doi:10.1126/science.1152509.
- 887 Houk, P., Camacho, R., Johnson, S., McLean, M., Maxin, S., Anson, J., et al. (2015). The Micronesia  
888 Challenge: Assessing the Relative Contribution of Stressors on Coral Reefs to Facilitate  
889 Science-to-Management Feedback. *PLOS ONE* 10, e0130823.  
890 doi:10.1371/journal.pone.0130823.
- 891 Hughes, T. P., Anderson, K. D., Connolly, S. R., Heron, S. F., Kerry, J. T., Lough, J. M., et al. (2018).  
892 Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science* 359,  
893 80–83. doi:10.1126/science.aan8048.
- 894 Hughes, T. P., Baird, A. H., Dinsdale, E. A., Moltschanivskyj, N. A., Pratchett, M. S., Tanner, J. E., et al.  
895 (1999). Patterns of recruitment and abundance of corals along the Great Barrier Reef.  
896 *Nature* 397, 59–63. doi:10.1038/16237.
- 897 Huston, M. A. (1985). Patterns of Species Diversity on Coral Reefs. *Annu. Rev. Ecol. Syst.* 16, 149–  
898 177. doi:10.1146/annurev.es.16.110185.001053.




- 899 IPCC (2019). Summary for Policymakers. – In: H.-O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P.  
900 Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.  
901 Weyer (eds.), IPCC Special Report on the Ocean and Cryosphere in a Changing Climate.
- 902 Jouffray, J.-B., Nyström, M., Norström, A. V., Williams, I. D., Wedding, L. M., Kittinger, J. N., et al.  
903 (2015). Identifying multiple coral reef regimes and their drivers across the Hawaiian  
904 archipelago. *Philosophical Transactions of the Royal Society B: Biological Sciences* 370,  
905 20130268. doi:10.1098/rstb.2013.0268.
- 906 Jouffray, J.-B., Wedding, L. M., Norström, A. V., Donovan, M. K., Williams, G. J., Crowder, L. B., et al.  
907 (2019). Parsing human and biophysical drivers of coral reef regimes. *Proceedings of the*  
908 *Royal Society B: Biological Sciences* 286, 20182544. doi:10.1098/rspb.2018.2544.
- 909 Knapp, K. R., Applequist, S., Howard, J. D., Kossin, J. P., Kruk, M., and Schreck, C. (2010a). NCDC  
910 International Best Track Archive for Climate Stewardship (IBTrACS) Project, Version 3. NOAA  
911 National Centers for Environmental Information. DOI:10.7289/V5NK3BZP.
- 912 Knapp, K. R., Kruk, M. C., Levinson, D. H., Diamond, H. J., and Neumann, C. J. (2010b). The  
913 International Best Track Archive for Climate Stewardship (IBTrACS). *Bull. Amer. Meteor. Soc.*  
914 91, 363–376. doi:10.1175/2009BAMS2755.1.
- 915 Kulbicki, M. (1998). How the acquired behaviour of commercial reef fishes may influence the results  
916 obtained from visual censuses. *Journal of Experimental Marine Biology and Ecology* 222, 11–  
917 30. doi:10.1016/S0022-0981(97)00133-0.
- 918 Kulbicki, M., Guillemot, N., and Amand, M. (2005). A general approach to length-weight relationships  
919 for New Caledonian lagoon fishes. *Cybium* 29, 235–252.
- 920 Labrosse, P., Kulbicki, M., and Ferraris, J. (2002). Underwater visual fish census surveys: Proper use  
921 and implementation. Noumea, New Caledonia: Secretariat of the Pacific Community (SPC).
- 922 Léopold, M., Guillemot, N., Rocklin, D., and Chen, C. (2014). A framework for mapping small-scale  
923 coastal fisheries using fishers' knowledge. *ICES J Mar Sci* 71, 1781–1792.  
924 doi:10.1093/icesjms/fst204.
- 925 Lugo, A. E., Rogers, C. S., and Nixon, S. W. (2000). Hurricanes, Coral Reefs and Rainforests:  
926 Resistance, Ruin and Recovery in the Caribbean. *AMBIO: A Journal of the Human*  
927 *Environment* 29, 106–114.
- 928 MacNeil, M. A., Mellin, C., Matthews, S., Wolff, N. H., McClanahan, T. R., Devlin, M., et al. (2019).  
929 Water quality mediates resilience on the Great Barrier Reef. *Nature Ecology & Evolution* 3,  
930 620–627. doi:10.1038/s41559-019-0832-3.
- 931 Maire, E., Cinner, J., Velez, L., Huchery, C., Mora, C., D'agata, S., et al. (2016). How accessible are  
932 coral reefs to people? A global assessment based on travel time. *Ecology Letters* 19, 351–  
933 360. doi:10.1111/ele.12577.
- 934 Manzello, D. P., Brandt, M., Smith, T. B., Lirman, D., Hendee, J. C., and Nemeth, R. S. (2007).  
935 Hurricanes benefit bleached corals. *PNAS* 104, 12035–12039. doi:10.1073/pnas.0701194104.


- 936 McWilliam, M., Pratchett, M. S., Hoogenboom, M. O., and Hughes, T. P. (2020). Deficits in functional  
937 trait diversity following recovery on coral reefs. *Proceedings of the Royal Society B: Biological*  
938 *Sciences* 287, 20192628. doi:10.1098/rspb.2019.2628.
- 939 Mellin, C., Matthews, S., Anthony, K. R. N., Brown, S. C., Caley, M. J., Johns, K. A., et al. (2019). Spatial  
940 resilience of the Great Barrier Reef under cumulative disturbance impacts. *Global Change*  
941 *Biology* 25, 2431–2445. doi:10.1111/gcb.14625.
- 942 Moberg, F., and Folke, C. (1999). Ecological goods and services of coral reef ecosystems. *Ecological*  
943 *Economics* 29, 215–233. doi:10.1016/S0921-8009(99)00009-9.
- 944 Morgan, K. M., Perry, C. T., Smithers, S. G., Johnson, J. A., and Daniell, J. J. (2016). Evidence of  
945 extensive reef development and high coral cover in nearshore environments: implications  
946 for understanding coral adaptation in turbid settings. *Scientific Reports* 6, 29616.  
947 doi:10.1038/srep29616.
- 948 Morse, N., Pellissier, P., Cianciola, E., Brereton, R., Sullivan, M., Shonka, N., et al. (2014). Novel  
949 ecosystems in the Anthropocene: a revision of the novel ecosystem concept for pragmatic  
950 applications. *Ecology and Society* 19. doi:10.5751/ES-06192-190212.
- 951 Mumby, P. J. (2017). Embracing a world of subtlety and nuance on coral reefs. *Coral Reefs* 36, 1003–  
952 1011. doi:10.1007/s00338-017-1591-8.
- 953 Norström, A. V., Nyström, M., Jouffray, J.-B., Folke, C., Graham, N. A., Moberg, F., et al. (2016).  
954 Guiding coral reef futures in the Anthropocene. *Frontiers in Ecology and the Environment* 14,  
955 490–498. doi:10.1002/fee.1427.
- 956 Nyström, M. (2006). Redundancy and response diversity of functional groups: Implications for the  
957 resilience of coral reefs. *ambi* 35, 30–35. doi:10.1579/0044-7447-35.1.30.
- 958 Pinca, S., Kronen, M., Friedman, K., Magron, F., Chapman, L., Tardy, E., et al. (2010). Regional  
959 assessment report: profiles and results from survey work at 63 sites across 17 Pacific Island  
960 Countries and Territories. Noumea, New Caledonia: Pacific Regional Oceanic and Coastal  
961 Fisheries Development Programme (PROCFish/C/CoFish).
- 962 QGIS Development Team (2019). *QGIS Geographic Information System. Open Source Geospatial*  
963 *Foundation Project*. <http://qgis.osgeo.org>. Available at: <http://qgis.osgeo.org>.
- 964 R Development Core Team (2019). *R: A language and environment for statistical computing*. Vienna,  
965 Austria: R Foundation for Statistical Computing.
- 966 Riegl, B., Berumen, M., and Bruckner, A. (2013). Coral population trajectories, increased disturbance  
967 and management intervention: a sensitivity analysis. *Ecol Evol* 3, 1050–1064.  
968 doi:10.1002/ece3.519.
- 969 Robinson, J. P. W., Williams, I. D., Yeager, L. A., McPherson, J. M., Clark, J., Oliver, T. A., et al. (2018).  
970 Environmental conditions and herbivore biomass determine coral reef benthic community  
971 composition: implications for quantitative baselines. *Coral Reefs* 37, 1157–1168.  
972 doi:10.1007/s00338-018-01737-w.
- 973 Roff, G., and Mumby, P. J. (2012). Global disparity in the resilience of coral reefs. *Trends in Ecology &*  
974 *Evolution* 27, 404–413. doi:10.1016/j.tree.2012.04.007.

- 975 Rogers, A., Blanchard, J. L., and Mumby, P. J. (2018). Fisheries productivity under progressive coral  
976 reef degradation. *Journal of Applied Ecology* 55, 1041–1049. doi:10.1111/1365-2664.13051.
- 977 Rogers, C. (1990). Responses of coral reefs and reef organisms to sedimentation. *Mar. Ecol. Prog.*  
978 *Ser.* 62, 185–202. doi:10.3354/meps062185.
- 979 Samoily, M. A., Halford, A., and Osuka, K. (2019). Disentangling drivers of the abundance of coral  
980 reef fishes in the Western Indian Ocean. *Ecology and Evolution* 9, 4149–4167.  
981 doi:10.1002/ece3.5044.
- 982 Sandin, S. A., Smith, J. E., DeMartini, E. E., Dinsdale, E. A., Donner, S. D., Friedlander, A. M., et al.  
983 (2008). Baselines and Degradation of Coral Reefs in the Northern Line Islands. *PLOS ONE* 3,  
984 e1548. doi:10.1371/journal.pone.0001548.
- 985 Schoepf, V., Stat, M., Falter, J. L., and McCulloch, M. T. (2015). Limits to the thermal tolerance of  
986 corals adapted to a highly fluctuating, naturally extreme temperature environment. *Scientific*  
987 *Reports* 5, 17639. doi:10.1038/srep17639.
- 988 Sheppard, C., Harris, A., and Sheppard, A. L. S. (2008). Archipelago-wide coral recovery patterns since  
989 1998 in the Chagos Archipelago, central Indian Ocean. *Marine Ecology Progress Series* 362,  
990 109–117.
- 991 Smith, J. E., Brainard, R., Carter, A., Grillo, S., Edwards, C., Harris, J., et al. (2016). Re-evaluating the  
992 health of coral reef communities: baselines and evidence for human impacts across the  
993 central Pacific. *Proc. R. Soc. B* 283, 20151985. doi:10.1098/rspb.2015.1985.
- 994 Steffen, W., Crutzen, P. J., and McNeill, J. R. (2007). The Anthropocene: Are Humans Now  
995 Overwhelming the Great Forces of Nature? *Ambio* 36, 614–621. doi:10.2307/25547826.
- 996 Taylor, A. H., Trouet, V., Skinner, C. N., and Stephens, S. (2016). Socioecological transitions trigger  
997 fire regime shifts and modulate fire–climate interactions in the Sierra Nevada, USA, 1600–  
998 2015 CE. *PNAS* 113, 13684–13689. doi:10.1073/pnas.1609775113.
- 999 Tebbett, S. B., and Bellwood, D. R. (2019). Algal turf sediments on coral reefs: what’s known and  
1000 what’s next. *Marine Pollution Bulletin* 149, 110542. doi:10.1016/j.marpolbul.2019.110542.
- 1001 van Hooidonk, R., Maynard, J., Tamelander, J., Gove, J., Ahmadi, G., Raymundo, L., et al. (2016).  
1002 Local-scale projections of coral reef futures and implications of the Paris Agreement.  
1003 *Scientific Reports* 6, 39666. doi:10.1038/srep39666.
- 1004 van Woesik, R., Sakai, K., Ganase, A., and Loya, Y. (2011). Revisiting the winners and the losers a  
1005 decade after coral bleaching. *Marine Ecology Progress Series* 434, 67–76.  
1006 doi:10.3354/meps09203.
- 1007 Warton, D. I., and Hui, F. K. C. (2011). The arcsine is asinine: the analysis of proportions in ecology.  
1008 *Ecology* 92, 3–10. doi:10.1890/10-0340.1.
- 1009 Wiedenmann, J., D’Angelo, C., Smith, E. G., Hunt, A. N., Legiret, F.-E., Postle, A. D., et al. (2013).  
1010 Nutrient enrichment can increase the susceptibility of reef corals to bleaching. *Nature*  
1011 *Climate Change* 3, 160–164. doi:10.1038/nclimate1661.

- 1012 Williams, G. J., Gove, J. M., Eynaud, Y., Zgliczynski, B. J., and Sandin, S. A. (2015a). Local human  
1013 impacts decouple natural biophysical relationships on Pacific coral reefs. *Ecography* 38, 751–  
1014 761. doi:10.1111/ecog.01353.
- 1015 Williams, G. J., Graham, N. A. J., Jouffray, J.-B., Norström, A. V., Nyström, M., Gove, J. M., et al.  
1016 (2019). Coral reef ecology in the Anthropocene. *Functional Ecology* 33, 1014–1022.  
1017 doi:10.1111/1365-2435.13290.
- 1018 Williams, G. J., Smith, J. E., Conklin, E. J., Gove, J. M., Sala, E., and Sandin, S. A. (2013). Benthic  
1019 communities at two remote Pacific coral reefs: effects of reef habitat, depth, and wave  
1020 energy gradients on spatial patterns. *PeerJ* 1, e81. doi:10.7717/peerj.81.
- 1021 Williams, I. D., Baum, J. K., Heenan, A., Hanson, K. M., Nadon, M. O., and Brainard, R. E. (2015b).  
1022 Human, Oceanographic and Habitat Drivers of Central and Western Pacific Coral Reef Fish  
1023 Assemblages. *PLOS ONE* 10, e0120516. doi:10.1371/journal.pone.0120516.
- 1024 Wood, S., and Scheipl, F. (2014). *gamm4: Generalized additive mixed models using mgcv and lme4. R*  
1025 *package version 0.2–3*.
- 1026 Woodhead, A. J., Hicks, C. C., Norström, A. V., Williams, G. J., and Graham, N. A. J. (2019). Coral reef  
1027 ecosystem services in the Anthropocene. *Functional Ecology* 33, 1023–1034.  
1028 doi:10.1111/1365-2435.13331.
- 1029 Zhang, S. Y., Speare, K. E., Long, Z. T., McKeever, K. A., Gyoerkoe, M., Ramus, A. P., et al. (2014). Is  
1030 coral richness related to community resistance to and recovery from disturbance? *PeerJ* 2,  
1031 e308. doi:10.7717/peerj.308.
- 1032 Zuur, A., Ieno, E., Walker, N., Saveliev, A. A., and Smith, G. M. (2009). *Mixed effects models and*  
1033 *extensions in ecology with R*. New York, USA: Springer Science+Business Media.
- 1034




**Supplementary Figure 1** - Map of the 62 sites surveyed across 17 different Pacific Island countries and territories, coloured by local impact level.




**Supplementary Figure 2** - Number of sites by human density (humans km<sup>-2</sup> reef). Dashed line indicates threshold separating low impact from high impact sites.

(A)




(B)



**Supplementary Figure 3** - To account for possible spatial autocorrelation among sites, we included island cluster as a random effect within models (based on methodology of Williams et al. 2015\*). Island clusters were identified using (A) hierarchical clustering based on pairwise Euclidean distances between each of the 62 sites, by finding (B) an inflection point in the number of unique clusters and cluster distance which we identified to be at ~1300 km, resulting in ten unique island clusters.


\*Williams GJ, Gove JM, Eynaud Y, Zgliczynski BJ, Sandin SA. (2015) Local human impacts decouple natural biophysical relationships on Pacific coral reefs. *Ecography*, **38**, 751–761.




**Supplementary Figure 4** - Boxplots representing ranges of response variables recorded by the core survey diver team - the core survey team consisted of five divers, who recorded 94% of all transects. Additional surveyors were included in the remaining 6%. As surveyor bias is inherent to such datasets, we tested for surveyor effects on our response variables. These tests indicated that significantly higher turf algae coverage was reported by two of the core survey team (survey divers iii and v). It became apparent that even within one reef, there were discrepancies in the reported turf algae cover between these survey divers and survey divers i, ii, and iv. This was not an issue with other benthic groups nor morphological classifications of hard corals (i.e. massive, branching, and encrusting). To account for the differences in turf algae reporting, we scored the contribution of survey divers iii and v to total surveys at each reef on a continuous scale from 0 to 1 (i.e. 0 meaning survey divers iii and v were not present on any transects at the reef, and 1 meaning all transects at a reef were conducted by survey diver iii and/or v). This score (referred to as *bias\_score*) was included as a random effect in subsequent turf algae-focused models.




## Live hard coral




## Dead hard coral




## Branching morphologies




## Encrusting morphologies




## Massive morphologies




## CCA



## Fleshy algae





## Turf algae




**Supplementary Figure 5** - Residuals (`$gam.check`) of ‘spatio-physical’ models (for each response variable (in order from top to bottom: live hard coral, dead hard coral, branching morphologies, encrusting morphologies, massive morphologies, CCA, macroalgae, turf algae). Left set of graphs = models based on low local impact sites (< 25 humans km<sup>-2</sup> reef); right set of graphs = models based on high local impact reefs (> 25 humans km<sup>-2</sup> reef):

(A)





(B)



**Supplementary Figure 6** – Individual relationships between selected predictors for each respective response variables as per ‘best-fit’ model outcomes. (A) Smoother plots of normalised residuals from generalised additive mixed effect models (*gam4* package) with 95% confidence intervals (shaded areas). Plots are based on the \$gam output from these models. These graphs provide relationships between individual response variables and predictors that were selected for ‘best-fit’ models (those with Akaike weight > 0.05; see Figure 4; Supplementary Table 4) at outer reef sites with low and/or high local human impacts. Plots are provided in terms of reefs exposed to low (light grey/left panel) *versus* high (dark grey/right panel) local impact as determined using a threshold of 25 humans km<sup>-2</sup> reef. Large crosses (X) mean that the predictor was not selected for ‘best-fit’ models for reefs exposed to either impact level. (B) Boxplots conveying relationships between the predictor island relief and benthic response variables for which it was selected for ‘best-fit’ models (i.e. those with Akaike weight > 0.05; see Figure 4; Supplementary Table 4).

**Supplementary Table 1** - Table containing survey dates and timelines of disturbances (up to 12 years before surveys) with associated references provided.

*\*Additional survey information: Criteria for site selection: (i) having active reef fisheries, (ii) being representative of the country, (iii) being relatively closed systems, (iv) being appropriate in size, (v) possessing diverse habitats\*, (vi) presenting no major logistical problems, (vii) having been previously investigated and (viii) presenting particular interest for the respective Fisheries Agency. For full details see Pinca et al. (2010). Originals surveys grouped reefs into four geomorphological structures, but this study focused only on outer (fore-) reefs (primarily due to replication). Though 63 individual outer reefs were surveyed, one reef had less than three replicate transects and thus was removed, leaving 62 replicate reefs.*

<b>Country</b>	<b>Sites &amp; Survey Dates</b>	<b>History of Events</b>	<b>References</b>
<b>Cook Islands</b>	Aitutaki, Feb-2007 Mangaia Oct-2007 Palmerston Feb-2007 Rarotonga Oct-2007	1997: Several cyclones (including Cyclone Martin – category 3) 2000: Mass bleaching event (up to 80% <i>Acropora</i> bleached in Rarotonga) 2002/2003: Major cyclone 2003/2004: Major cyclone 2004/2005: Several destructive cyclones (six)	Lovell 2001, Cumming et al. 2002, Salvat 2002, de Scally 2008, Pinca et al. 2009a
<b>Fiji</b>	Dromuna Apr-2003 Lakeba Jun-2003 Mali Jun-2003 Muiavuso Apr-2003	2000: Mass bleaching event (40–80% coral mortality) 2001: Minor bleaching 2002: Bleaching in shallow areas	Goreau et al. 2000, Cumming et al. 2002, Lovell et al. 2004, Friedman et al. 2010
<b>French Polynesia</b>	Fakarava Mar-2004 Maatea May-2006 Mataiea May/Jun-2003 Raivavae Mar-2004 Tikehau Oct-2003	1994: Major bleaching but low mortality 1997: Cyclone Martin hit (southern) French Polynesia. Patchy bleaching and mortality 2002: Start of COTS outbreak (at Society Archipelago and Australes)	Salvat 2002, Vieux et al. 2004, Adjeroud et al. 2005, Kronen et al. 2008a, Trapon et al. 2011

<b>FSM</b>	Piis-Panewu Apr-2006 Riiken Apr-2006 Romanum Apr-2006 Yyin May-2006	2005: Concerning levels of COTs reported from rapid assessment	George et al. 2008, Kronen et al. 2009a
<b>Kiribati</b>	Abaiang Jun-2004 Abemama May-2004 Kiritimati Sep-2004 Kurua May-2004	2003: Fish kill in atolls of Gilbert Islands 2004: (post survey) first mass bleaching event internationally reported	Awira et al. 2008, Donner et al. 2010
<b>Marshall Islands</b>	Ailuk Aug-2007 Arno Sep-2007 Laura Sep-2007 Likiep Aug-2007	2001: Bleaching mortality observed (particularly shallow <i>Acropora</i> ) 2003: Bleaching event, high incidence of <i>Acropora</i> white disease 2004: Elevated COTS (>1000 km <sup>-2</sup> ) found in Majuro's SW lagoon, with associated coral mortality 2006: Storm caused large surf and storm surge (become Typhoon Soulik) – <i>Acropora</i> colonies damaged. Also bleaching event with up to 90% <i>Acropora</i> bleaching (20–50% mortality) in lagoons.	Beger et al. 2008 Pinca et al. 2009b
<b>Nauru</b>	Nauru Oct-2005	2000: Phosphate reserves are virtually exhausted (80% of the island's surface has been strip mined – mining started in 1908) 2002: High rainfall and some bleaching 2003: Coral bleaching and mass fish kills both linked to elevated SSTs, strong upwelling or drops in dissolved oxygen levels. 2004: Large number of	Sulu et al. 2002, Lovell et al. 2004, Sauni et al. 2007, Chin et al. 2011



		seabirds found dead without a known cause	
<b>New Caledonia</b>	Luengoni Jun-2004 Moindou May-2004 Ouasse Aug-2004 Oundjo Feb-2004 Thio Mar-2004	1998 and 2001: Extensive coral loss due to COTs, bleaching and disease 2003: Cyclone (Erica) led to 10–80% coral loss	Sulu et al. 2002, Lovell et al. 2004, Kronen et al. 2009b
<b>Niue</b>	All Niue May-2005	2004: Major cyclone (Cyclone Heta – category 5) caused 20–90% reefs on the western coast to be flattened, hence live coral in surveys sometimes <2%	Vieux et al. 2004, Kronen et al. 2008b, Chin et al. 2011
<b>Palau</b>	Airai Apr-2007 Koror May-2007 Ngarchelong Apr-2007 Ngatpang Apr-2007	1998: Mass bleaching from El Niño reduced coral cover from 50–70% to 14–23% (exposed barrier reefs suffered more than coastal reefs). High mortality of soft corals. <i>Area is not exposed to cyclones and COTs outbreaks remain local events</i>	Friedman et al. 2009a, Golbuu 2011
<b>Papua New Guinea</b>	Andra Aug-2006 Panapompom Nov-2006 Sideia Oct-2006 Tsoilaunung Aug-2006	1996: Mass bleaching event (>50% corals at some sites) 1998: Mass bleaching from El Niño (75% <i>Acropora</i> affected at Kimbe Bay) 1999: Bleaching event 2000/2001: Mass bleaching event (>50% corals at some sites)	Lovell 2001, Cumming et al. 2002, Foale 2006, Friedman et al. 2009b
<b>Samoa</b>	Manono-uta Jun-2005 Salelavalu Aug-2005 Vailoa Jun-2005 Vaisala Aug-2005	2004: Cyclone (Heta – category 5) damaged 13% of coral reefs	Sulu et al. 2002, Lovell et al. 2004, Vunisea et al. 2008, Chin et al. 2011

<b>Solomon Islands</b>	Chubikopi Dec-2006 Marau Jun-2006 Nggela Jun-2006 Rarumana Aug-2006	2000: Mass bleaching 2002: Major cyclone (Zoe – category 5)	Lovell 2001, Cumming et al. 2002, Sulu et al. 2002, Pinca et al. 2009c
<b>Tonga</b>	Ha’afutu2 Sep-2008 Koulo Oct-2008 Lofanga Oct-2008 Manuka Sep-2008	1997: Cyclone (Hina – category 3) 1999: Cyclone (Cora – category 3) 2000: Cyclone (Mona – category 3), mass bleaching (up to 90% of <i>Goniastrea</i> and <i>Platygyra</i> colonies bleached)	Lovell 2001, Salvat 2002, Friedman et al. 2009c
<b>Tuvalu</b>	Funafuti Nov-2004 Niutao Apr-2005 Nukufetau Oct/Nov-2004 Vaitipu Mar/Apr-2005	1997: Major cyclones (Oliwa and Keli) 2002: Mass bleaching (30–40% corals bleached)	Sulu et al. 2002, Lovell et al. 2004, Sauni et al. 2008
<b>Vanuatu</b>	Maskelynes Nov/Dec-2003 Moso Jul-2003 Paunangisu Jul-2003 Uri-Uripiv Nov/Dec-2003	2001: Bleaching event (documented in Moso but lack of monitoring data for other areas)	Sulu et al. 2002, Friedman et al. 2008
<b>Wallis and Futuna</b>	All Futuna Nov-2005 Halalo Sep-2005 Vailala Sep-2005		Kronen et al. 2008c

## References

Adjeroud M, Chancerelle Y, Schrimm M et al. (2005) Detecting the effects of natural disturbances on coral assemblages in French Polynesia: a decade survey at multiple scales. *Aquatic Living Resources*, **18**, 111–123.

Awira R, Friedman K, Sauni S et al. (2008) Kiribati country report: Profiles and results from survey work at Abaiang, Abemama, Kuria and Kirimati: (May to November 2004). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 313 pp.

Beger M, Jacobson D, Pinca S et al. (2008) The state of coral reef ecosystems of the Republic of the Marshall Islands. In: *Status of coral reefs of the world: 2008* (ed. Wilkinson CR) pp. 387–417. Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville, Australia, 296 pp.

Chin A, Lison de Loma T, Reytar K et al. (2011) Status of coral reefs of the Pacific and outlook: 2011. Global Coral Reef Monitoring Network, 260 pp.

Cumming RL, Toscano M, Lovell A et al. (2002) Mass coral bleaching in the Fiji Islands, 2000. In: *Proceedings of the Ninth International Coral Reef Symposium, Bali, 23–27 October 2000*, **2**, 1161–1167.

de Scally F (2008) Historical tropical cyclone activity and impacts in the Cook Islands. *Pacific Science*, **62**, 443–459

Donner SD, Kirata T, Vieux C (2010) Recovery from the 2004 coral bleaching event at the Gilbert Islands, Kiribati. *Atoll Research Bulletin*, **587**, 1–25.

Foale SJ (2006) The scale and epistemology of coral bleaching in Papua New Guinea. Bridging Scales and Epistemologies. Alexandria: Millennium Ecosystem Assessment.

Friedman K, Pakoa K, Kronen M et al. (2008) Vanuatu country report: profiles and results from survey work at Paunangisu Village, Moso Island, Uri and Uripiv Islands and the Maskelyne Archipelago (July to December 2003). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 357 pp.

Friedman K, Kronen M, Pinca S et al. (2009a) Palau country report: Profiles and results from survey work at Ngarchelong, Ngatpang, Airai and Koror (April to June 2007). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 411 pp.

Friedman K, Kronen M, Pinca S et al. (2009b) Papua New Guinea country report: profiles and results from survey work at Andra, Tsoilaunung, Sideia, and Panapompom (June to November 2006). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Noumea, New Caledonia: Secretariat of the Pacific Community (SPC). 471 pp.

Friedman K, Pakoa K, Kronen M et al. (2009c) Tonga country report: profiles and results from survey work at Ha'atafu, Manuka, Koulo and Lofanga (November and December 2001; March to June 2002; April to June, September and October 2008). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 369 pp.

Friedman K, Kronen M, Vunisea A et al. (2010) Fiji Islands country report: profiles and results from survey work at Dromuna, Muavuso, Mali and Lakeba (September to November 2002, April to June 2003, June and July 2007, and February 2009). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 467 pp.

George A, Luckymis M, Palik S et al. (2008) The state of coral reef ecosystems of the Federated States of Micronesia. In: *Status of coral reefs of the world: 2008* (ed. Wilkinson CR) pp. 419–436. Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville, Australia, 296 pp.

Golbuu, Y (2011) Responses of Palau's coral reefs to disturbances at multiple scales. PhD thesis, Southern Cross University, Lismore, NSW.

Goreau T, McClanahan T, Hayes R, Strong A (2000) Conservation of coral reefs after the 1998 Global Bleaching Event. *Conservation Biology*, **14**, 5–15.

Kronen M, Friedman K, Pinca S et al. (2008a) French Polynesia country report: Profiles and results from survey work at Fakarava, Maatea, Mataiea, Raivavae and Tikehau (September –

October 2003, January – March 2004, April – June 2006). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 383 pp.

Kronen M, Fisk D, Pinca S et al. (2008b) Niue country report: Profile and results from in-country survey work (May to June 2005). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 173 pp.

Kronen M, Tardy E, Boblin P et al. (2008c) Wallis and Futuna country report: profiles and results from survey work at Vailala, Halalo, Leava and Vele (August–December 2005 and March 2006). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 333 pp.

Kronen M, Friedman K, Boblin P et al. (2009a) Federated States of Micronesia country report: Profiles and results from survey work at Yyin and Riiken (YAP) and Piis-Panewu and Romanum (Chuuk) (April – May 2006). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 367 pp.

Kronen M, Boblin P, Friedman K et al. (2009b) New Caledonia country report: Profile and results from survey work at Ouassé, Thio, Luengoni, Oundjo and Moindou (March, April and November 2003; January, February, April, June, August and November 2004; April and May 2005; January to March 2006; and January and February 2007) . Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 409 pp.

Lovell ER (2001) Reef Check description of the 2000 mass coral bleaching event in Fiji with reference to the South Pacific. Unpubl report to University of Rhode Island. Available online at: [http://www.crc.uri.edu/download/fiji\\_mass\\_bleaching\\_report.pdf](http://www.crc.uri.edu/download/fiji_mass_bleaching_report.pdf)

Lovell ER, Sykes H, Debye M et al. (2004) Status of coral reefs in the South West Pacific: Fiji, Nauru, New Caledonia, Samoa, Solomon Islands, Tuvalu and Vanuatu. In: *Status of coral reefs of the world: 2004* (ed. Wilkinson CR), pp. 337–362. Australian Institute of Marine Science, Townsville, Australia.

Pinca S, Awira R, Kronen M et al. (2009a) Cook Islands country report: Profiles and results from survey work at Aitutaki, Palmerston, Mangaia and Rarotonga (February and October 2007). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 339 pp.

Pinca S, Tardy E, Awira R et al. (2009b) Marshall Islands country report: profiles and results from survey work at Likiep, Ailuk, Arno and Laura (August and September 2007). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 337 pp.

Pinca S, Vunise A, Lasi F et al. (2009c) Solomon Islands country report: profiles and results from survey work at Nggela, Marau, Rarumana and Chubikopi (June to September 2006 and December 2006). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 409 pp.

Salvat B (2002) Status of Southeast and Central Pacific coral reefs 'Polynesia Mana Node': Cook Islands, French Polynesia, Kiribati, Niue, Tokelau, Tonga, Wallis and Futuna. In: *Status of coral reefs of the world: 2002* (ed. Wilkinson CR), pp. 203–216. Australian Institute of Marine Science, Townsville, Australia.

Sauni S, Vunisea A, Kronen M et al. (2007) Nauru country report: profile and results from in-country survey work (October and November 2005). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 137 pp.

Sauni S, Kronen M, Pinca S et al. (2008) Tuvalu country report: profile and results from in-country survey work (October–November 2004 and March–April 2005). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 311 pp.

Sulu R, Cumming R, Wantiez L et al. (2002) Status of coral reefs in the South West Pacific: Fiji, Nauru, New Caledonia, Samoa, Solomon Islands, Tuvalu and Vanuatu. In: *Status of coral reefs of the world: 2002* (ed. Wilkinson CR), pp. 181–202. Australian Institute of Marine Science, Townsville, Australia.

Trapon ML, Pratchett MS, Penin L (2011) Comparative effects of different disturbances in coral reef habitats in Moorea, French Polynesia. *Journal of Marine Biology*, **2011**, doi:10.1155/2011/807625

Vieux C, Aubanel A, Axford JC et al. (2004) A century of change in coral reef status in Southeast and Central Pacific: Polynesia Mana Node, Cook Islands, French Polynesia, Kiribati, Niue, Tokelau, Tonga, Wallis and Futuna. In: *Status of coral reefs of the world: 2004* (ed. Wilkinson CR), pp. 363–380. Australian Institute of Marine Science, Townsville, Australia.

Vunisea A, Friedman K, Awira R et al. (2008) Samoa country report: profiles and results from survey work at Manono-Uta, Salelavalu, Vailoa and Vaisala (June 2005 and August/September 2005). Pacific Regional Oceanic and Coastal Fisheries Development Programme (PROCFish/C/CoFish). Secretariat of the Pacific Community (SPC): Noumea, New Caledonia. 349 pp.

**Supplementary Table 2** - Herbivorous fish recorded within surveys, classified into functional groups as specified in Heenan et al. (2016).

<b>Functional group</b>	<b>Species</b>
Browsers	<i>Calotomus carolinus</i>
Browsers	<i>Calotomus spinidens</i>
Browsers	<i>Kyphosus cinerascens</i>
Browsers	<i>Kyphosus</i> sp.
Browsers	<i>Kyphosus vaigiensis</i>
Browsers	<i>Leptoscarus vaigiensis</i>
Browsers	<i>Naso annulatus</i> <sup>1</sup>
Browsers	<i>Naso brachycentron</i>
Browsers	<i>Naso brevirostris</i> <sup>1</sup>
Browsers	<i>Naso lituratus</i>
Browsers	<i>Naso</i> sp.
Browsers	<i>Naso tuberosus</i>
Browsers	<i>Naso unicornis</i>
Browsers	<i>Platax orbicularis</i>
Browsers	<i>Platax</i> sp.
Browsers	<i>Platax teira</i>
Browsers	<i>Siganus canaliculatus</i>
Detritivores	<i>Ctenochaetus binotatus</i>
Detritivores	<i>Ctenochaetus cyanocheilus</i>
Detritivores	<i>Ctenochaetus flavicauda</i>
Detritivores	<i>Ctenochaetus hawaiiensis</i>
Detritivores	<i>Ctenochaetus marginatus</i>
Detritivores	<i>Ctenochaetus</i> sp.
Detritivores	<i>Ctenochaetus striatus</i>
Detritivores	<i>Ctenochaetus strigosus</i>
Detritivores	<i>Ctenochaetus tominiensis</i>
Grazers	<i>Acanthurus achilles</i>
Grazers	<i>Acanthurus auranticavus</i>
Grazers	<i>Acanthurus blochii</i>
Grazers	<i>Acanthurus dussumieri</i>
Grazers	<i>Acanthurus fowleri</i>
Grazers	<i>Acanthurus guttatus</i>
Grazers	<i>Acanthurus leucocheilus</i>
Grazers	<i>Acanthurus leucopareius</i>
Grazers	<i>Acanthurus lineatus</i>
Grazers	<i>Acanthurus maculiceps</i>
Grazers	<i>Acanthurus nigricans</i>
Grazers	<i>Acanthurus nigricauda</i>
Grazers	<i>Acanthurus nigrofuscus</i>
Grazers	<i>Acanthurus nigroris</i>
Grazers	<i>Acanthurus olivaceus</i>
Grazers	<i>Acanthurus pyroferus</i>
Grazers	<i>Acanthurus</i> sp.

<b>Functional group</b>	<b>Species</b>
Grazers	<i>Acanthurus triostegus</i>
Grazers	<i>Acanthurus xanthopterus</i>
Grazers	<i>Centropyge bicolor</i>
Grazers	<i>Centropyge bispinosa</i>
Grazers	<i>Centropyge flavissima</i>
Grazers	<i>Centropyge loricula</i>
Grazers	<i>Centropyge sp.</i>
Grazers	<i>Centropyge tibicen</i>
Grazers	<i>Centropyge vrolikii</i>
Grazers	<i>Siganus argenteus</i>
Grazers	<i>Siganus corallinus</i>
Grazers	<i>Siganus doliatus</i>
Grazers	<i>Siganus fuscescens</i>
Grazers	<i>Siganus guttatus</i>
Grazers	<i>Siganus lineatus</i>
Grazers	<i>Siganus niger</i>
Grazers	<i>Siganus puellus</i>
Grazers	<i>Siganus punctatissimus</i>
Grazers	<i>Siganus punctatus</i>
Grazers	<i>Siganus randalli</i>
Grazers	<i>Siganus sp.</i>
Grazers	<i>Siganus spinus</i>
Grazers	<i>Siganus stellatus</i>
Grazers	<i>Siganus uspi</i>
Grazers	<i>Siganus vermiculatus</i>
Grazers	<i>Siganus vulpinus</i>
Grazers	<i>Zebrasoma flavescens</i>
Grazers	<i>Zebrasoma rostratum</i>
Grazers	<i>Zebrasoma scopas</i>
Grazers	<i>Zebrasoma velifer</i>
Scrapers / small excavators	<i>Chlorurus bleekeri</i>
Scrapers / small excavators	<i>Chlorurus bowersi</i>
Scrapers / small excavators	<i>Chlorurus japanensis</i>
Scrapers / small excavators	<i>Chlorurus sordidus</i>
Scrapers / small excavators	<i>Chlorurus sp.</i>
Scrapers / small excavators	<i>Hipposcarus longiceps</i>
Scrapers / small excavators	<i>Scarus altipinnis</i>
Scrapers / small excavators	<i>Scarus chameleon</i>
Scrapers / small excavators	<i>Scarus dimidiatus</i>
Scrapers / small excavators	<i>Scarus festivus</i>
Scrapers / small excavators	<i>Scarus flavipectoralis</i>
Scrapers / small excavators	<i>Scarus forsteni</i>
Scrapers / small excavators	<i>Scarus frenatus</i>
Scrapers / small excavators	<i>Scarus ghobban</i>
Scrapers / small excavators	<i>Scarus globiceps</i>
Scrapers / small excavators	<i>Scarus hypselopterus</i>

<b>Functional group</b>	<b>Species</b>
Scrapers / small excavators	<i>Scarus longipinnis</i>
Scrapers / small excavators	<i>Scarus niger</i>
Scrapers / small excavators	<i>Scarus oviceps</i>
Scrapers / small excavators	<i>Scarus prasiognathos</i>
Scrapers / small excavators	<i>Scarus psittacus</i>
Scrapers / small excavators	<i>Scarus quoyi</i>
Scrapers / small excavators	<i>Scarus rivulatus</i>
Scrapers / small excavators	<i>Scarus schlegeli</i>
Scrapers / small excavators	<i>Scarus sp.</i>
Scrapers / small excavators	<i>Scarus spinus</i>
Scrapers / small excavators	<i>Scarus tricolor</i>
Scrapers / small excavators	<i>Scarus xanthopleura</i>
Large excavators / bioeroders	<i>Bolbometopon muricatum</i>
Large excavators / bioeroders	<i>Cetoscarus ocellatus</i> <sup>2</sup>
Large excavators / bioeroders	<i>Chlorurus frontalis</i> <sup>2</sup>
Large excavators / bioeroders	<i>Chlorurus microrhinos</i> <sup>2</sup>
Large excavators / bioeroders	<i>Scarus rubroviolaceus</i> <sup>2</sup>

<sup>1</sup> classified as 'browser' until reaches 20 cm in length

<sup>2</sup> classified as 'scraper/small excavator' when smaller than 35 cm in length, and 'large excavator/bioeroder' when larger than 35 cm in length

## Reference

Heenan A, Hoey AS, Williams GJ, Williams ID (2016) Natural bounds of herbivorous coral reef fishes. *Proc. R. Soc. B* **283**: 20161716. <http://dx.doi.org/10.1098/rspb.2016.1716>



**Supplementary Table 3** - As part of a sensitivity analysis, ‘spatio-physical’ models were additionally run using  $\pm 5$  and  $\pm 10$  humans  $\text{km}^{-2}$  reef as a threshold from which to categorise ‘low’ and ‘high’ impact sites. Notably, site allocations were the same for 30 and 35 humans  $\text{km}^{-2}$  reef (i.e. no sites had values between 30 and 35 humans  $\text{km}^{-2}$  reef). Values in tables represent adjusted- $R^2$  values from GAMM models with structure as described in the manuscript. All outcomes were consistent with those found for the 25 humans  $\text{km}^{-2}$  reef threshold, indicating the findings are robust.

	Threshold = 15 humans $\text{km}^{-2}$ reef		Threshold = 20 humans $\text{km}^{-2}$ reef	
	low (n=25)	high (n=37)	low (n=27)	high (n=35)
Live hard coral	0.69	0.00	0.49	0.00
Branching	0.66	0.09	0.63	0.07
Massive	0.41	0.00	0.55	0.02
Encrusting	0.09	0.00	0.04	0.00
Turf algae	0.00	0.00	0.00	0.00
Macroalgae	0.63	0.19	0.63	0.16
Dead hard coral	0.21	0.09	0.23	0.03
CCA	0.00	0.00	0.00	0.00
<b>Average</b>	<b>0.34</b>	<b>0.05</b>	<b>0.32</b>	<b>0.04</b>
<b>t-test outcome</b>	<b>0.019</b>		<b>0.015</b>	

	Threshold = 25 humans $\text{km}^{-2}$ reef		Threshold = 30/35 humans $\text{km}^{-2}$ reef	
	low (n=29)	high (n=33)	low (n=34)	high (n=28)
Live hard coral	0.52	0.00	0.50	0.00
Branching	0.64	0.17	0.64	0.01
Massive	0.50	0.10	0.47	0.00
Encrusting	0.19	0.00	0.08	0.06
Turf algae	0.00	0.00	0.00	0.00
Macroalgae	0.59	0.26	0.55	0.23
Dead hard coral	0.33	0.24	0.37	0.17
CCA	0.02	0.00	0.05	0.00
<b>Average</b>	<b>0.35</b>	<b>0.10</b>	<b>0.33</b>	<b>0.06</b>
<b>t-test outcome</b>	<b>0.010</b>		<b>0.015</b>	

**Supplementary Table 4** - As an additional analysis, ‘spatio-physical’ models were run for human density as calculated per *outer reef area* only (i.e. human density per outer reef) using 60, 80 and 100 humans km<sup>-2</sup> outer reef as thresholds from which to categorise ‘low’ and ‘high’ impact sites (note: the thresholds are higher as the area of reef relative to humans is smaller when considering only outer reef). Values in tables represent adjusted-R<sup>2</sup> values from GAMM models with structure as described in the manuscript. All outcomes were consistent with those found for the 25 humans km<sup>-2</sup> reef threshold, indicating the findings are robust. While certain target fish species only occur on the outer reefs, some of the fishing and gleaning activity to meet protein demands would be focused on areas other than outer reefs. Furthermore, additional impacts by local human populations (in particular nutrient input) would be diluted in cases where large areas of backreef occur. Thus, even though there are trade-offs involved with either calculation of human density, we feel that human density per total reef area is the most appropriate proxy of local human impact, and thus present those values in the main manuscript.

	Threshold = 60 humans km <sup>-2</sup> outer reef		Threshold = 80 humans km <sup>-2</sup> outer reef	
	low (n=25)	high (n=37)	low (n=32)	high (n=30)
Live hard coral	0.48	0.18	0.38	0.01
Branching	0.33	0.12	0.33	0.20
Massive	0.49	0.08	0.47	0.09
Encrusting	0.00	0.00	0.00	0.00
Turf algae	0.23	0.00	0.15	0.00
Macroalgae	0.39	0.31	0.39	0.21
Dead hard coral	0.28	0.00	0.20	0.02
CCA	0.00	0.03	0.00	0.00
<b>Average</b>	<b>0.27</b>	<b>0.09</b>	<b>0.24</b>	<b>0.07</b>
<b>t-test outcome</b>	<b>0.012</b>		<b>0.011</b>	

	Threshold = 100 humans km <sup>-2</sup> outer reef	
	low (n=35)	high (n=27)
Live hard coral	0.37	0.37
Branching	0.35	0.35
Massive	0.47	0.47
Encrusting	0.00	0.00
Turf algae	0.07	0.07
Macroalgae	0.45	0.45
Dead hard coral	0.20	0.20
CCA	0.01	0.01
<b>Average</b>	<b>0.24</b>	<b>0.02</b>
<b>t-test outcome</b>	<b>0.009</b>	

**Supplementary Table 5** - Best-fit model outcomes based on GAMM models (with island cluster as a random effect for all response variables, and surveyor bias as an additional random effect for turf algae). A ‘+’ indicates that the predictor was selected by model selection procedures for the respective row’s model. RI values in grey boxes under model selection output (boldface if predictor was selected by model selection procedures for models with Akaike weight >0.05). (A) outcomes for low local impact sites (< 25 humans km<sup>-2</sup> reef), (B) outcomes for high local impact sites (> 25 humans km<sup>-2</sup> reef).

(A)

	relief	browsers	depth	detrivores	excavators	gravity	grazers	latitude	scrapers	storms	adj-R2	adj-R2 conditional	df	logLik	AICc	delta	weight
LIVE CORAL	-0.24		+								0.50	0.70	6.00	-13.18	42.20	0.00	0.27
											0.09	0.13	3.00	-18.29	43.50	1.36	0.14
			+	+					+		0.51	0.70	7.00	-12.19	43.70	1.54	0.13
											0.31	0.43	5.00	-15.83	44.30	2.10	0.10
		+									0.33	0.45	5.00	-15.93	44.50	2.29	0.09
	-0.19										0.21	0.29	4.00	-17.86	45.40	3.22	0.05
	+	+								0.48	0.67	7.00	-13.12	45.60	3.39	0.05	
	<b>0.42</b>	<b>0.18</b>	<b>0.66</b>	0.01	0.01	0.01	0.01	0.01	<b>0.20</b>	0.01							
BRANCHING CORAL										+	0.55	0.59	5.00	-31.52	75.70	0.00	0.49
				+						+	0.64	0.68	7.00	-28.77	76.90	1.23	0.26
			+							+	0.60	0.63	7.00	-30.50	80.30	4.67	0.05
	0.06	0.02	<b>0.06</b>	<b>0.30</b>	0.05	0.01	0.02	0.04	0.02	<b>1.00</b>							
MASSIVE CORAL										+	0.48	0.51	5.00	-30.20	73.00	0.00	0.32
										+	0.60	0.64	7.00	-27.49	74.30	1.31	0.17
		+									0.37	0.40	5.00	-31.87	76.30	3.33	0.06
	-0.21									+	0.50	0.54	6.00	-30.33	76.50	3.46	0.06
	-0.43										0.34	0.37	4.00	-33.51	76.70	3.67	0.05
	<b>0.22</b>	<b>0.13</b>	0.03	0.06	0.01	0.05	0.05	0.03	<b>0.73</b>	<b>0.27</b>							
ENCRUSTING CORAL											0.10	0.12	3.00	-25.28	57.50	0.00	0.63
	0.09				+						0.20	0.24	5.00	-24.48	61.60	4.05	0.08
	<b>0.10</b>	0.01	0.02	0.05	<b>0.11</b>	0.01	0.04	0.02	0.02	0.05							
TURF ALGAE											0.18	0.18	4.00	-49.83	109.30	0.00	0.49
				+							0.25	0.26	6.00	-48.22	112.30	2.93	0.11
	-0.19										0.19	0.19	5.00	-49.95	112.50	3.19	0.10
					+						0.21	0.21	6.00	-48.66	113.10	3.81	0.07
	<b>0.15</b>	0.03	0.02	<b>0.15</b>	<b>0.10</b>	0.05	0.03	0.03	0.03	0.02							
FLESHY ALGAE	-0.83									+	0.50	0.51	6.00	-47.75	111.30	0.00	0.25
	-0.96		+							+	0.64	0.65	8.00	-44.34	111.90	0.58	0.19
	-1.00										0.31	0.32	4.00	-51.89	113.50	2.14	0.09
										+	0.36	0.37	5.00	-50.75	114.10	2.80	0.06
	<b>0.72</b>	0.13	<b>0.25</b>	0.03	0.07	0.02	0.02	0.13	0.07	<b>0.79</b>							
CCA	-0.25										0.00	0.00	3.00	-38.85	84.70	0.00	0.55
								+			0.06	0.07	4.00	-38.72	87.10	2.45	0.16
											0.03	0.03	5.00	-38.65	89.90	5.26	0.04
	<b>0.22</b>	0.03	0.05	0.02	0.02	0.02	0.06	0.04	0.04	0.03							
DEAD CORAL										+	0.39	0.48	5.00	-20.21	53.00	0.00	0.58
		+								+	0.52	0.64	7.00	-18.26	55.80	2.82	0.14
										+	0.51	0.63	7.00	-18.93	57.20	4.16	0.07
	0.06	<b>0.16</b>	0.02	0.01	0.01	0.01	0.01	0.01	<b>0.89</b>	<b>0.13</b>							

(B)

	relief	browsers	depth	detritivores	excavators	gravity	grazers	latitude	scrapers	storms	adj-R2	adj-R2 conditional	df	logLik	AICc	delta	weight
LIVE CORAL											0.06	0.08	3.00	-23.00	52.90	0.00	0.44
	-0.25										0.15	0.20	4.00	-22.47	54.40	1.57	0.20
							+				0.21	0.28	5.00	-21.28	54.90	2.00	0.16
BRANCHING CORAL	<b>0.25</b>	0.03	0.04	0.02	0.01	0.05	<b>0.23</b>	0.04	0.01	0.01							
											0.03	0.04	3.00	-50.81	108.50	0.00	0.32
	-0.51										0.11	0.11	4.00	-49.80	109.10	0.60	0.23
MASSIVE CORAL				+							0.11	0.11	5.00	-49.71	111.70	3.25	0.06
	<b>0.41</b>	0.03	0.03	<b>0.13</b>	0.03	0.06	0.05	0.04	0.03	0.12							
											0.34	0.35	3.00	-47.54	101.90	0.00	0.34
									+		0.42	0.43	5.00	-46.27	104.80	2.91	0.08
	0.19										0.35	0.36	4.00	-47.71	104.90	2.96	0.08
								+			0.40	0.42	5.00	-46.42	105.10	3.20	0.07
ENCRUSTING CORAL				+							0.38	0.39	5.00	-46.42	105.10	3.21	0.07
											0.38	0.39	5.00	-46.46	105.20	3.29	0.07
	<b>0.16</b>	0.03	<b>0.12</b>	<b>0.12</b>	0.07	0.02	<b>0.12</b>	0.08	<b>0.14</b>	0.03							
											0.03	0.03	3.00	-40.01	86.90	0.00	0.25
							+				0.25	0.27	5.00	-37.35	87.00	0.13	0.24
					+						0.21	0.23	5.00	-38.19	88.70	1.81	0.10
TURF ALGAE											0.40	0.44	7.00	-35.19	89.00	2.17	0.09
	0.20										0.06	0.07	4.00	-40.20	89.90	3.01	0.06
	<b>0.15</b>	<b>0.15</b>	0.02	<b>0.19</b>	0.03	<b>0.42</b>	0.02	0.03	0.03	0.02							
	-0.04										0.53	0.55	4.00	-44.78	99.00	0.00	0.62
FLESHY ALGAE											0.53	0.55	5.00	-45.42	103.20	4.10	0.08
							+				0.58	0.60	6.00	-44.01	103.40	4.33	0.07
	<b>0.11</b>	0.02	0.06	0.04	0.03	0.02	<b>0.10</b>	0.03	0.02	0.02							
CCA											0.06	0.06	3.00	-54.46	115.80	0.00	0.33
	-0.54										0.15	0.15	4.00	-53.49	116.50	0.69	0.23
											0.11	0.11	5.00	-53.43	119.20	3.39	0.06
DEAD CORAL	<b>0.36</b>	<b>0.10</b>	0.03	0.07	0.03	0.04	0.04	0.07	0.06	0.07							
											0.05	0.06	3.00	-28.76	64.40	0.00	0.44
											0.26	0.31	5.00	-26.36	65.00	0.65	0.32
DEAD CORAL	-0.07										0.06	0.07	4.00	-29.70	68.90	4.51	0.05
	<b>0.08</b>	0.38	0.02	0.02	0.02	0.05	0.01	0.02	0.02	0.02							
	0.00										0.00	0.00	3.00	-8.94	24.70	0.00	0.79
DEAD CORAL											0.00	0.00	4.00	-10.73	30.90	6.20	0.04
	0.04	0.03	0.02	0.01	0.02	0.01	0.03	0.04	0.02	0.01	0.10	0.28	5.00	-9.38	31.10	6.33	0.03