

Logics of formal inconsistency arising from systems of fuzzy logic

Marcelo Coniglio¹ Francesc Esteva²
Lluís Godo²

¹ CLE and Department of Philosophy
State University of Campinas
13083-859 Campinas, Brazil
email: coniglio@cle.unicamp.br

² Artificial Intelligence Research Institute (IIIA), CSIC
Campus UAB, 08193 Bellaterra, Spain
email: {esteva,godo}@iiia.csic.es

Abstract

This paper proposes the meeting of fuzzy logic with paraconsistency in a very precise and foundational way. Specifically, in this paper we introduce expansions of the fuzzy logic MTL by means of primitive operators for consistency and inconsistency in the style of the so-called Logics of Formal Inconsistency (LFIs). The main novelty of the present approach is the definition of postulates for this type of operators over MTL-algebras, leading to the definition and axiomatization of a family of logics, expansions of MTL, whose degree-preserving counterpart are paraconsistent and moreover LFIs.

1 Introduction

The well-known *Sorites paradox* is representative of the problems arising from the use of vague predicates, that is, predicates whose extension is unclear such as ‘tall’ and ‘bald’. According to Charles S. Peirce,

A proposition is vague when there are possible states of things concerning which it is intrinsically uncertain whether, had they been contemplated by the speaker, he would have regarded them as excluded or allowed by the proposition. By intrinsically uncertain we mean not uncertain in consequence of any ignorance of the interpreter, but because the speaker’s habits of language were indeterminate; so that one day he would regard the proposition as excluding, another as admitting, those states of things. ([25])

Besides being an instigating topic for Philosophy, vagueness is also studied from the mathematical and logical point of view. For instance, the so-called *Mathematical Fuzzy Logic* (MFL), inspired by the paradigm of *Fuzzy Set Theory* introduced in 1965 by L. Zadeh (cf. [28]), studies the question of vagueness from a foundational point of view based on many-valued logics. In this sense, MFL can be considered as a degree-based approach to vagueness.¹ Some systems like Łukasiewicz and Gödel-Dummett infinitely valued logics are, just like fuzzy sets, valued over the real interval $[0, 1]$. This supports the idea of MFL being as a kind of foundational counterpart of fuzzy set theory (which is a discipline mainly devoted to engineering applications). The book [19] by P. Hájek is the first monograph dedicated to a broad study of the new subject of MFL. In that book the so-called Basic fuzzy logic BL is introduced as the residuated many-valued logic with the semantics on the real unit interval induced by all *continuous* t-norms and their residua. BL generalizes three prominent fuzzy logics, Łukasiewicz, Gödel-Dummett and Product logics, each one capturing the semantics determined by three particular continuous t-norms, namely Łukasiewicz, minimum and product t-norms respectively. The so-called Monoidal t-norm based logic (MTL was introduced in [16] as a generalization of BL to capture the semantics induced by *left continuous* t-norms and their residua, in fact, as it was proved in [23] the theorems of MTL correspond to the common tautologies of all many-valued calculi defined by a left-continuous t-norm and its residuum. This logic, the most general residuated fuzzy logic whose semantics is based on t-norms, will be the starting point of our investigations in the present paper.

Frequently, vagueness is associated to a phenomenon of ‘underdetermination of truth’. However, vagueness could be seen from an opposite perspective: if a is a borderline case of a vague predicate P , the sentences ‘ a is P ’ and ‘ a is not P ’ can be both true (at least to some extent). This leads to an interpretation of vagueness as ‘overdetermination of truth’, instead of underdetermination. Being so, a sentence A and its negation can simultaneously be both true, without trivializing (as much we assume that not every sentence is true). This perspective, known as *Paraconsistent Vagueness*, connects vagueness to the subject of *Paraconsistent Logic* (see, for instance, [22] and [13]).

Paraconsistency is devoted to the study of logic systems with a negation operator, say \neg , such that not every contradictory set of premises $\{\varphi, \neg\varphi\}$ trivializes the system. Thus, any paraconsistent logic contains at least a contradictory but non-trivial theory. There exist several systematic approaches to paraconsistency, for instance: N. da Costa’s hierarchy of C-systems C_n , for $n > 0$, introduced in 1963 (see [14]); Relevance (or Relevant) logics,

¹See e.g. [10] for several discussions on degree-based approaches (and in particular fuzzy logic approaches) to vagueness.

introduced by A. Anderson and N. Belnap in 1975 (see [1]); the Adaptive Logics programme, developed by D. Batens and his group; R. Routley and G. Priest’s philosophical school of Dialetheism, with Priest’s logic LP as its formalized counterpart (see, for instance, [26]); and the Logics of Formal Inconsistency (LFIs), introduced by W. Carnielli and J. Marcos in 2000 (see [7] and [6]), and also studied e.g. by Avron et al. [3, 2]. The main characteristic of the latter logics is that they internalize in the object language the notions of consistency and inconsistency by means of specific connectives (primitive or not). This constitutes a generalization of da Costa’s C-systems.

The present paper proposes the meeting of fuzzy logic with paraconsistency in a very precise and foundational way. Specifically, we introduce extensions of the fuzzy logic MTL by means of primitive operators for consistency and inconsistency, defining so LFIs based on (extensions of) MTL. An important feature of this approach is that the LFIs defined in this manner are not based on (positive) classical logic, as in the case of most LFIs studied in the literature, including da Costa’s C-systems. In particular, the LFIs proposed here do not satisfy the law of excluded middle: $\varphi \vee \neg\varphi$ is not a valid schema, in general.

The main novelty of the present approach is the definition of postulates for primitive consistency and inconsistency fuzzy operators over the algebras associated to (extensions of) MTL; in particular, we show how to define consistency and inconsistency operators over MTL-algebras. This generalizes the previous approach to fuzzy LFIs introduced in [15], where it was shown that a consistency operator can be defined in MTL_Δ , the expansion of MTL with the Monteiro-Baaz projection connective Δ . However, this consistency operator is not primitive, but it is defined in terms of the operator Δ together with other operators of MTL. At this point, it is important to observe that MTL, as well as its extensions, are *not* paraconsistent logics, provided that the usual truth-preserving consequence relation is considered: from $\{\varphi, \neg\varphi\}$ every other formula can be derived. On the other hand, if a degree-preserving consequence relation is adopted, MTL as well as some of its extensions become paraconsistent (see Section 2).

The organization of this paper is as follows. In Sections 2 and 3, the basic notions about fuzzy logics and LFIs are introduced. Then Section 4 contains the main definitions and technical results. In particular, we introduce the notion of consistency operators on MTL-algebras and axiomatize several classes of them as expansions of MTL. In this framework, the question about how the consistency operator propagates with respect to the MTL connectives is studied in Section 5. In its turn, in Section 6 we propose a fuzzy LFI able to recover classical logic by considering additional hypothesis on the consistency operator. The dual case of inconsistency operators is briefly analyzed in Section 7. We end up with some concluding remarks in Section 8.

2 Preliminaries I: truth-preserving and degree-preserving fuzzy logics

In the framework of Mathematical Fuzzy Logic there are two different families of fuzzy logics according to how the logical consequence is defined, namely truth-preserving and degree-preserving logics. In this section we review the main definitions and properties of these two families of logics.

Truth-preserving fuzzy logics. Most well known and studied systems of mathematical fuzzy logic are the so-called *t-norm based fuzzy logics*, corresponding to formal many-valued calculi with truth-values in the real unit interval $[0, 1]$ and with a conjunction and an implication interpreted respectively by a (left-) continuous t-norm and its residuum respectively, and thus, including e.g. the well-known Łukasiewicz and Gödel infinitely-valued logics, corresponding to the calculi defined by Łukasiewicz and min t-norms respectively. The weakest t-norm based fuzzy logic is the logic MTL (monoidal t-norm based logic) introduced in [16], whose theorems correspond to the common tautologies of all many-valued calculi defined by a left-continuous t-norm and its residuum [23].

The language of MTL consists of denumerably many propositional variables p_1, p_2, \dots , binary connectives $\wedge, \&, \rightarrow$, and the truth constant $\bar{0}$. Formulas, which will be denoted by lower case greek letters $\varphi, \psi, \chi, \dots$, are defined by induction as usual. Further connectives and constants are definable; in particular, $\neg\varphi$ stands for $\varphi \rightarrow \bar{0}$, $\bar{1}$ stands for $\neg\bar{0}$, $\varphi \vee \psi$ stands for $((\varphi \rightarrow \psi) \rightarrow \psi) \wedge ((\psi \rightarrow \varphi) \rightarrow \varphi)$, and $\varphi \leftrightarrow \psi$ stands for $(\varphi \rightarrow \psi) \wedge (\psi \rightarrow \varphi)$. A Hilbert-style calculus for MTL was introduced in [16] with the following set of axioms:

- (A1) $(\varphi \rightarrow \psi) \rightarrow ((\psi \rightarrow \chi) \rightarrow (\varphi \rightarrow \chi))$
- (A2) $\varphi \& \psi \rightarrow \varphi$
- (A3) $\varphi \& \psi \rightarrow \psi \& \varphi$
- (A4) $\varphi \wedge \psi \rightarrow \varphi$
- (A5) $\varphi \wedge \psi \rightarrow \psi \wedge \varphi$
- (A6) $\varphi \& (\varphi \rightarrow \psi) \rightarrow \varphi \wedge \psi$
- (A7a) $(\varphi \rightarrow (\psi \rightarrow \chi)) \rightarrow (\varphi \& \psi \rightarrow \chi)$
- (A7b) $(\varphi \& \psi \rightarrow \chi) \rightarrow (\varphi \rightarrow (\psi \rightarrow \chi))$
- (A8) $((\varphi \rightarrow \psi) \rightarrow \chi) \rightarrow (((\psi \rightarrow \varphi) \rightarrow \chi) \rightarrow \chi)$
- (A9) $\bar{0} \rightarrow \varphi$

and whose unique rule of inference is *modus ponens*: from φ and $\varphi \rightarrow \psi$ derive ψ .

MTL is an algebraizable logic in the sense of Blok and Pigozzi [4] and its equivalent algebraic semantics is given by the class of MTL-algebras, that is indeed a variety; call it \mathbf{MTL} . MTL-algebras can be equivalently introduced as commutative, bounded, integral residuated lattices $\langle A, \wedge, \vee, \&, \rightarrow, 0, 1 \rangle$ further satisfying the following *prelinearity* equation:

$$(x \rightarrow y) \vee (y \rightarrow x) = \bar{1}.$$

Given an MTL-algebra $\mathbf{A} = \langle A, \wedge^{\mathbf{A}}, \vee^{\mathbf{A}}, \&^{\mathbf{A}}, \rightarrow^{\mathbf{A}}, 0^{\mathbf{A}}, 1^{\mathbf{A}} \rangle$, an \mathbf{A} -evaluation is any function mapping each propositional variable into A , $e(\bar{0}) = 0^{\mathbf{A}}$ and such that, for formulas φ and ψ , $e(\varphi \wedge \psi) = e(\varphi) \wedge^{\mathbf{A}} e(\psi)$; $e(\varphi \vee \psi) = e(\varphi) \vee^{\mathbf{A}} e(\psi)$; $e(\varphi \& \psi) = e(\varphi) \&^{\mathbf{A}} e(\psi)$; $e(\varphi \rightarrow \psi) = e(\varphi) \rightarrow^{\mathbf{A}} e(\psi)$. An evaluation e is said to be a *model* for a set of formulas Γ , if $e(\gamma) = 1^{\mathbf{A}}$ for each $\gamma \in \Gamma$.

We shall henceforth adopt a lighter notation dropping the superscript \mathbf{A} . The distinction between a syntactic object and its interpretation in an algebraic structure will always be clear by the context.

The algebraizability gives the following strong completeness theorem:

For every set $\Gamma \cup \{\varphi\}$ of formulas, $\Gamma \vdash_{\mathbf{MTL}} \varphi$ iff for every $\mathbf{A} \in \mathbf{MTL}$ and every \mathbf{A} -evaluation e , if e is a model of Γ then e is a model of φ as well.

For this reason, since the consequence relation amounts to preservation of the truth-constant $\bar{1}$, MTL can be called a (*full*) *truth-preserving* logic.

Actually, the algebraizability is preserved for any logic L that is a (finitary) expansion of MTL satisfying the following congruence property

$$\begin{aligned} (\text{Cong}) \quad \varphi \rightarrow \psi, \psi \rightarrow \varphi \vdash_L \quad & c(\chi_1, \dots, \chi_i, \varphi, \chi_{i+2}, \dots, \chi_n) \\ & \rightarrow c(\chi_1, \dots, \chi_i, \psi, \chi_{i+2}, \dots, \chi_n) \end{aligned}$$

for any possible new n -ary connective c and each $i < n$. These expansions, that we will call *core expansions* of MTL (in accordance with [11]), are in fact *Rasiowa-implicative logics* (cf. [27]). As proved in [12], every Rasiowa-implicative logic L is algebraizable and, if it is finitary, its equivalent algebraic semantics, the class \mathbb{L} of L -algebras, is a quasivariety. *Axiomatic* expansions of MTL, i.e. without any further inference rule, satisfying (Cong) are called *core fuzzy logics* in the literature (see e.g. [12]), and their associated quasi-varieties of algebras are in fact varieties.

As a consequence, any logic L which is a core expansion of MTL, in particular any core fuzzy logic, enjoys the same kind of the above strong completeness theorem with respect to the whole class of corresponding L -algebras. But for core fuzzy logics we can say more than that. Indeed, for any core fuzzy logic L , the variety of L -algebras can also be shown to be

generated by the subclass of all its linearly ordered members [12].² This means that any core fuzzy logic L is strongly complete with respect to the class of L-chains, that is, core fuzzy logics are *semilinear*.

All core fuzzy logics enjoy a form of local deduction theorem. As usual, φ^n will be used as a shorthand for $\varphi \& .^n. \& \varphi$, where $\varphi^0 = \bar{1}$. Using this notation one can write the following *local deduction theorem* for any core fuzzy logics L: for each set of formulas $\Sigma \cup \{\varphi, \psi\}$ the following holds:

$$\Sigma, \varphi \vdash_L \psi \text{ iff there is } n \geq 0 \text{ such that } \Sigma \vdash_L \varphi^n \rightarrow \psi.$$

Interesting axiomatic extensions of MTL used in the paper are the ones given in Table 2, but first we list in Table 1 the axioms needed to define these extensions of MTL.

Axiom schema	Name
$\neg\neg\varphi \rightarrow \varphi$	Involution (Inv)
$\neg\varphi \vee ((\varphi \rightarrow \varphi \& \psi) \rightarrow \psi)$	Cancellation (C)
$\varphi \rightarrow \varphi \& \varphi$	Contraction (Con)
$\varphi \wedge \psi \rightarrow \varphi \& (\varphi \rightarrow \psi)$	Divisibility (Div)
$\varphi \wedge \neg\varphi \rightarrow \bar{0}$	Pseudo-complementation (PC)
$(\varphi \& \psi \rightarrow \bar{0}) \vee (\varphi \wedge \psi \rightarrow \varphi \& \psi)$	Weak Nilpotent Minimum (WNM)

Table 1: Some usual axiom schemata in fuzzy logics.

MTL can be considered in fact as the logic of left-continuous t-norms [23] and BL as the logic of continuous t-norms [8], in the sense that theorems of these logics coincide with common tautologies of interpretations on the MTL (respectively BL) chains defined on the real unit interval $[0, 1]$ by left-continuous (respectively continuous) t-norms and their residua.

Another interesting family of fuzzy logics are the so-called *logics of a (left-continuous) t-norm*. Given a left-continuous t-norm $*$, define the real (or standard) algebra $[0, 1]_* = ([0, 1], \min, \max, *, \rightarrow_*, 0, 1)$ where \rightarrow_* is the residuum of $*$. Then define the logic of the t-norm $*$ as the logic L_* whose (semantical) notion of consequence relation is as follows: φ is a consequence of a set of formulas Γ iff for every evaluation v over $[0, 1]_*$ such that $v(\gamma) = 1$ for each $\gamma \in \Gamma$, then $v(\varphi) = 1$. When $*$ is a continuous t-norm, L_* has been proved finitely axiomatizable as extension of BL (see [18]).

²Moreover, for a number of core fuzzy logics, including MTL, it has been shown that their corresponding varieties are also generated by the subclass of MTL-chains defined on the real unit interval, indistinctively called in the literature as *standard* or *real* chains. For instance, MTL is also complete wrt real MTL-chains, that are of the form $[0, 1]_* = \langle [0, 1], \min, \max, *, \rightarrow_*, 0, 1 \rangle$ of type $\langle 2, 2, 2, 2, 0, 0 \rangle$, where $*$ denotes a left-continuous t-norm and \rightarrow_* is its residuum [23].

Logic	Additional axiom schemata	References
Strict MTL (SMTL)	(PC)	[20]
Involutive MTL (IMTL)	(Inv)	[16]
Weak Nilpotent Minimum (WNM)	(WNM)	[16]
Nilpotent Minimum (NM)	(Inv) and (WNM)	[16]
Basic Logic (BL)	(Div)	[19]
Strict Basic Logic (SBL)	(Div) and (PC)	[17]
Lukasiewicz Logic (\mathbb{L})	(Div) and (Inv)	[19]
Product Logic (Π)	(Div) and (C)	[21]
Gödel Logic (G)	(Con)	[19]

Table 2: Some axiomatic extensions of MTL obtained by adding the corresponding additional axiom schemata, and the references where they have been introduced (in the context of fuzzy logics).

As we have mentioned, all the axiomatic expansions of MTL (i.e. all core fuzzy logics) are semilinear and enjoy the local deduction detachment theorem. Another very interesting class of fuzzy logics arise from the (non-axiomatic) expansion of MTL with the Monteiro-Baaz projection connective Δ , obtaining again a finitary Rasiowa-implicative semilinear logic MTL_Δ . Indeed, MTL_Δ is axiomatized by adding to the Hilbert-style system of MTL the deduction rule of necessitation (from φ infer $\Delta\varphi$) and the following axiom schemata:

- (Δ 1) $\Delta\varphi \vee \neg\Delta\varphi$
- (Δ 2) $\Delta(\varphi \vee \psi) \rightarrow (\Delta\varphi \vee \Delta\psi)$
- (Δ 3) $\Delta\varphi \rightarrow \varphi$
- (Δ 4) $\Delta\varphi \rightarrow \Delta\Delta\varphi$
- (Δ 5) $\Delta(\varphi \rightarrow \psi) \rightarrow (\Delta\varphi \rightarrow \Delta\psi)$

Then, one analogously defines the class of Δ -core fuzzy logics as the axiomatic expansions of MTL_Δ satisfying (Cong) for any possible new connective. They satisfy the *global deduction theorem* in the following way: for any Δ -core fuzzy logic L , and each set of formulas $\Sigma \cup \{\varphi, \psi\}$, the following holds:

$$\Sigma, \varphi \vdash_L \psi \text{ iff } \Sigma \vdash_L \Delta\varphi \rightarrow \psi.$$

Semilinearity can also be inherited by many expansions of (Δ -)core fuzzy logics with new (finitary) inference rules. Indeed, in [12] it is shown that an expansion L of a (Δ -)core fuzzy logic is semilinear iff for each newly added finitary inference rule

(R) from Γ derive φ ,

its corresponding \vee -form

(\mathbf{R}^\vee) from $\Gamma \vee p$ derive $\varphi \vee p$

is derivable in L as well, where p is an arbitrary propositional variable not appearing in $\Gamma \cup \{\varphi\}$.

In this paper we will use the following notions of completeness of a logic L with respect to the class *real* L -chains. Although we will mainly focus on core fuzzy logics, we formulate them for the more general case of logics that are semilinear expansions of MTL whose class of *real* L -chains is non-empty.

Definition 2.1 ($\mathcal{R}C$, $FSRC$, SRC) *Let L be a semilinear core expansion of MTL and let \mathcal{R} be the class of real L -chains, i.e. L -chains whose support is the real unit interval $[0, 1]$. We say that L has the (finitely) strong \mathcal{R} -completeness property, (F) SRC for short, when for every (finite) set of formulas T and every formula φ it holds that $T \vdash_L \varphi$ iff $e(\varphi) = \bar{1}^{\mathbf{A}}$ for each \mathbf{A} -evaluation such that $e[T] \subseteq \{\bar{1}^{\mathbf{A}}\}$ for every L -algebra $\mathbf{A} \in \mathcal{R}$. We say that L has the \mathcal{R} -completeness property, $\mathcal{R}C$ for short, when the equivalence is true for $T = \emptyset$.*

Of course, the SRC implies the $FSRC$, and the $FSRC$ implies the $\mathcal{R}C$. The SRC and $FSRC$ have traditionally been proved for many fuzzy logics by showing an embeddability property, namely by showing in the first case that every countable L -chain is embeddable into a chain of \mathcal{R} , and in the second case by showing that every countable L -chain is *partially* embeddable into a chain of \mathcal{R} (i.e. for every finite partial algebra of a countable L -chain there is a one-to-one mapping into some L -chain over $[0, 1]$ preserving the defined operations). In [9] it was shown that, for (Δ) -core fuzzy logics these sufficient conditions are also necessary (under a weak condition). This was further generalized in [12], where Cintula and Noguera show that these conditions are also necessary for a more general class of logics, including semilinear core expansions of MTL.

Theorem 2.2 ([9, 12] Characterization of completeness properties)

Let L be a semilinear core expansion of MTL. Then:

- *L has the SRC iff every countable L -chain is embeddable into some chain of \mathcal{R} .*
- *If the language of L is finite, then L has the $FSRC$ iff every countable L -chain is partially embeddable into some chain of \mathcal{R} .*

Degree-preserving fuzzy logics. It is clear that (Δ) -core fuzzy logics, like MTL, are (full) truth-preserving fuzzy logics. But besides the truth-preserving paradigm that we have so far considered, one can find an alternative approach in the literature. Given a (Δ) -core fuzzy logic L , and based on the definitions in [5], we can introduce a variant of L that we shall denote

by L^\leq , whose associated deducibility relation has the following semantics: for every set of formulas $\Gamma \cup \{\varphi\}$, $\Gamma \vdash_{L^\leq} \varphi$ iff for every L-chain \mathbf{A} , every $a \in A$, and every \mathbf{A} -evaluation v , if $a \leq v(\psi)$ for every $\psi \in \Gamma$, then $a \leq v(\varphi)$. For this reason L^\leq is known as a fuzzy logic *preserving degrees of truth*, or the *degree-preserving companion* of L. In this paper, we often use generic statements about “every logic L^\leq ” referring to “the degree-preserving companion of any (Δ -)core fuzzy logic (or even of any semilinear core expansion of MTL) L”.

As regards to axiomatization, if L is a core fuzzy logic, i.e. with Modus Ponens as the unique inference rule, then the logic L^\leq admits a Hilbert-style axiomatization having the same axioms as L and the following deduction rules [5]:

(Adj- \wedge) from φ and ψ derive $\varphi \wedge \psi$

(MP- r) if $\vdash_L \varphi \rightarrow \psi$ (i.e. if $\varphi \rightarrow \psi$ is a theorem of L), then from φ derive ψ

Note that if the set of theorems of L is decidable, then the above is in fact a recursive Hilbert-style axiomatization of L^\leq .

In general, let L be a semilinear core expansion of MTL with a set of new inference rules,

(\mathbf{R}_i) from Γ_i derive φ_i , for each $i \in I$.

Then L^\leq is axiomatized by adding to the axioms of L the above two inference rules plus the following restricted rules

(\mathbf{R}_i - r) if $\vdash_L \Gamma_i$, then derive φ_i .

Moreover, if L is a Δ -core fuzzy logic, then the only rule one should add to L^\leq is the following restricted necessitation rule for Δ :

(Δ - r) if $\vdash_L \varphi$, then derive $\Delta\varphi$.

The key relationship between L and L^\leq is given by the following equivalence: for any formulas $\varphi_1, \dots, \varphi_n, \psi$, it holds

$$\varphi_1, \dots, \varphi_n \vdash_{L^\leq} \psi \text{ iff } \vdash_L (\varphi_1 \wedge \dots \wedge \varphi_n) \rightarrow \psi.$$

This relation points out that, indeed, deductions from a finite set of premises in L^\leq exactly correspond to theorems in L. In particular, both logics share the same theorems: $\vdash_L \varphi$ iff $\vdash_{L^\leq} \varphi$. Moreover, this also implies that if L' is a conservative expansion of L, then L'^\leq is also a conservative expansion of L^\leq .

3 Preliminaries II: logics of formal inconsistency

Paraconsistency is the study of logics having a negation operator \neg such that it is not explosive with respect to \neg , that is, there exists at least a formula φ such that from $\{\varphi, \neg\varphi\}$ it does not follow any formula. In other words, a paraconsistent logic is a logic having at least a contradictory, non-trivial theory.

Among the plethora of paraconsistent logics proposed in the literature, the *Logics of Formal Inconsistency* (LFIs), proposed in [7] (see also [6]), play an important role, since they internalize in the object language the very notions of consistency and inconsistency by means of specific connectives (primitives or not).³ This generalizes the strategy of N. da Costa, which introduced in [14] the well-known hierarchy of systems C_n , for $n > 0$. Besides being able to distinguish between contradiction and inconsistency, on the one hand, and non-contradiction and consistency, on the other, LFIs are non-explosive logics, that is, paraconsistent: in general, a contradiction does not entail arbitrary statements, and so the Principle of Explosion (for all φ, ψ it holds $\varphi, \neg\varphi \vdash \psi$) does not hold. However, LFIs are *gently explosive*, in the sense that, adjoining the additional requirement of consistency, then contradictoriness does cause explosion: $\circ(\varphi), \varphi, \neg\varphi \vdash \psi$ for every φ and ψ . Here, $\circ(\varphi)$ denotes that φ is consistent. The general definition of LFIs we will adopt here, slightly modified from the original one proposed in [7] and [6], is the following:

Definition 3.1 *Let L be a logic defined in a language \mathcal{L} containing a negation \neg , and let $\circ(p)$ be a nonempty set of formulas depending exactly on the propositional variable p . Then L is an LFI (with respect to \neg and $\circ(p)$) if the following holds (here, $\circ(\varphi) = \{\psi[p/\varphi] : \psi(p) \in \circ(p)\}$ and $\psi[p/\varphi]$ denotes the formula obtained from ψ by replacing every occurrence of the variable p by the formula φ):*

- (i) $\varphi, \neg\varphi \not\vdash \psi$ for some φ and ψ , i.e., L is not explosive w.r.t. \neg ;
- (ii) $\circ(\varphi), \varphi \not\vdash \psi$ for some φ and ψ ;
- (iii) $\circ(\varphi), \neg\varphi \not\vdash \psi$ for some φ and ψ ; and
- (iv) $\circ(\varphi), \varphi, \neg\varphi \vdash \psi$ for every φ and ψ , i.e., L is gently explosive w.r.t. \neg and $\circ(p)$.

In the case that $\circ(\varphi)$ is a singleton (which will be the usual situation), its element will be denoted by $\circ\varphi$, and \circ will be called a *consistency* operator in L with respect to \neg . A consistency operator can be primitive (as in the

³We should warn the reader that in the frame of LFIs, the term *consistency* is used to refer to formulas that basically exhibit a classical, explosive behaviour rather than for referring to formulas being (classically) satisfiable.

case of most of the systems treated in [7] and [6]) or, on the contrary, it can be defined in terms of the other connectives of the language. For instance, in the well-known system C_1 by da Costa, consistency is defined by the formula $\circ\varphi = \neg(\varphi \wedge \neg\varphi)$ (see [14]).

Given a consistency operator \circ , an *inconsistency* operator \bullet is naturally defined as $\bullet\varphi = \neg\circ\varphi$. In the stronger LFIs, the other way round holds, and so \circ can be defined from a given \bullet as $\circ\varphi = \neg\bullet\varphi$.

All the LFIs proposed in [7] and [6] are extensions of positive classical logic, therein called CPL^+ . The weaker system considered there is called mbC, defined in a language containing $\wedge, \vee, \rightarrow, \neg$ and \circ , and it is obtained from CPL^+ by adding the schema axioms $\varphi \vee \neg\varphi$ and $\circ\varphi \rightarrow (\varphi \rightarrow (\neg\varphi \rightarrow \psi))$.

As we shall see in the next section, the definition of LFIs can be generalized to the algebraic framework of MTLs, constituting an interesting approach to paraconsistency under the perspective of LFIs, but without the requirement of being an extension of CPL^+ .

4 Axiomatizing expansions of paraconsistent fuzzy logics with consistency operators \circ

As observed in [15], truth preserving fuzzy logics are not paraconsistent since from $\varphi, \neg\varphi$ we obtain $\varphi \& \neg\varphi$, that is equivalent to the truth-constant $\bar{0}$, and thus they are explosive. However in the case of degree-preserving fuzzy logics, from $\varphi, \neg\varphi$ one cannot always derive the truth-constant $\bar{0}$, and hence there are paraconsistent degree-preserving fuzzy logics. Indeed we have the following scenario.

Proposition 4.1 *Let L be a semilinear core expansion of MTL. The following conditions hold:*

1. L is explosive, and hence it is not paraconsistent
2. L^{\leq} is paraconsistent iff L is not pseudo-complemented, i.e. if L does not prove the law $\neg(\varphi \wedge \neg\varphi)$.

The proof of the second item is easy since $\varphi, \neg\varphi \vdash_{L^{\leq}} \bar{0}$ does not hold only in the case L does not prove $(\varphi \wedge \neg\varphi) \rightarrow \bar{0}$, or in other words, only in the case L is not an expansion of SMTL.

As a consequence, from now on L will refer to any semilinear core expansion of MTL which is not a SMTL logic (not satisfying axiom (PC)). Indeed we are interested in the expansion with a consistency operator \circ of a logic L^{\leq} (when L is not a SMTL logic). In order to axiomatize these expansions, we need first to axiomatize the expansion of the truth-preserving L with such an operator \circ and from them, as explained in Section 2, we can then obtain the desired axiomatizations.

4.1 Expansions of truth-preserving fuzzy logics with consistency operators \circ

Having in mind the properties that a consistency operator has to verify in a paraconsistent logic (recall Definition 3.1), and taking into account that any semilinear core expansion of MTL is complete with respect to the chains of the corresponding varieties, it seems reasonable to define a consistency operator over a non-SMTL chain \mathbf{A} as a unary operator $\circ : A \rightarrow A$ satisfying the following conditions:

- (i) $x \wedge \circ(x) \neq 0$ for some $x \in A$;
- (ii) $\neg x \wedge \circ(x) \neq 0$ for some $x \in A$;
- (iii) $x \wedge \neg x \wedge \circ(x) = 0$ for every $x \in A$.

Such an operator \circ can be indeed considered as the algebraic counterpart of a *consistency operator* in the sense of Definition 3.1. Actually, we can think about the value $\circ(x)$ as denoting the (fuzzy) degree of ‘classicality’ (or ‘reliability’, or ‘robustness’) of x with respect to the satisfaction of the law of explosion, namely $x \wedge \neg x = 0$.

Let us have a closer look at how operators \circ on a non-SMTL chain satisfying the above conditions (i), (ii) and (iii) may look like. Let us consider the set $N(\mathbf{A}) = \{x \in A \setminus \{1\} \mid \neg x = 0\}$. Notice that either $N(\mathbf{A}) = \emptyset$ (for example, this is the case of IMTL chains) or $N(\mathbf{A}) \in \{[a, 1), (a, 1)\}$ where $a = \bigwedge N(\mathbf{A})$. If $x \notin N(A) \cup \{0, 1\}$ then by (iii) we have $\circ(x) = 0$. Thus since $\neg(x) = 0$ for $x \in N(A)$, (ii) implies that $\circ(0) > 0$. On the other hand by (i), $\circ(x) > 0$ for some $x \in N(A) \cup \{1\}$. Therefore, any operator \circ verifying (i), (ii) and (iii) must satisfy the following minimal conditions:

$$\begin{cases} \circ(0) > 0, \\ \circ(x) = 0, & \text{if } x \in (0, 1) \setminus N(\mathbf{A}) \\ \circ(x) > 0, & \text{for some } x \in N(A) \cup \{1\} \end{cases}$$

However, since in our setting the intended meaning of $\circ(x)$ is the (fuzzy) degree of ‘classicality’ or ‘reliability’, or ‘robustness’ of x , we propose the following stronger postulates for such a consistency operator on non-SMTL chains \mathbf{A} :

- (c1) If $x \wedge \neg x \neq 0$ then $\circ(x) = 0$;
- (c2) If $x \in \{0, 1\}$ then $\circ(x) = 1$;
- (c3) If $\neg x = 0$ and $x \leq y$ then $\circ(x) \leq \circ(y)$.

Clause (c1) just guarantees that condition (iii) for consistency operators is satisfied by \circ . In the classical case, both truth-values 0 and 1 satisfy the explosion law $x \wedge \neg x = 0$ and so $\circ(x) = 1$ for every truth-value x . Since \circ intends to extend the classical case, clause (c2) reflects this situation (another justification for (c2) is that 0 and 1 are classical truth-values with fuzzy degree 1). Moreover, clause (c2) ensures that conditions (i) and (ii) for consistency operators are satisfied. Finally, clause (c3) ensures the coherency of \circ : in $N(\mathbf{A})$, the segment of the chain where \circ is positive, the consistency operator \circ is monotonic, in accordance with the idea that $\circ(x)$ is the fuzzy degree of classicality, from the perspective of the explosion law: “the closer is x to 1, the more classical is x ”. In Figure 1, we depict in blue (dashed-line) the graph of the negation \neg in the real BL-chain $[0, 1]_*$, where $*$ is the ordinal sum of Łukasiewicz t-norm in $[0, a]$ and another arbitrary t-norm on the interval $[a, 1]$ and in red (bold line) the graph of a \circ operator compatible with the above postulates.




Figure 1: Graph of the negation in a BL-algebra of the form $[0, a]_{\mathbb{L}} \oplus [a, 1]_*$ and a graph of a \circ operator satisfying postulates (c1), (c2) and (c3).

As a consequence, we propose the following definition.

Definition 4.2 *Let \mathbb{L} be any semilinear core expansion of MTL. Given an axiomatization of \mathbb{L} , we define the logic \mathbb{L}_{\circ} as the expansion of \mathbb{L} in a language which incorporates a new unary connective \circ with the following axioms:*

- (A1) $\neg(\varphi \wedge \neg\varphi \wedge \circ\varphi)$
- (A2) $\circ\bar{1}$
- (A3) $\circ\bar{0}$

and the following inference rules:

$$\text{(Cong)} \quad \frac{(\varphi \leftrightarrow \psi) \vee \delta}{(\circ\varphi \leftrightarrow \circ\psi) \vee \delta} \qquad \text{(Coh)} \quad \frac{(\neg\neg\varphi \wedge (\varphi \rightarrow \psi)) \vee \delta}{(\circ\varphi \rightarrow \circ\psi) \vee \delta}$$

Due to the presence of the rule (Cong), L_\circ is a Rasiowa-implicative logic, and thus it is algebraizable in the sense of Blok and Pigozzi, and its algebraic semantics is given by L_\circ -algebras.

Definition 4.3 L_\circ -algebras are expansions of L -algebras with a new unary operation \circ satisfying the following conditions, for all $x, y, z \in A$:

$$(\circ 1) \quad x \wedge \neg x \wedge \circ(x) = 0$$

$$(\circ 2) \quad \circ(1) = \circ(0) = 1$$

$$(\circ 3) \quad \text{if } (\neg\neg x \wedge (x \rightarrow y)) \vee z = 1 \text{ then } (\circ(x) \rightarrow \circ(y)) \vee z = 1$$

Thus, the class \mathcal{L}_\circ of L_\circ -algebras is a quasivariety, and since it is the equivalent algebraic semantics of the logic L_\circ , L_\circ is (strongly) complete with respect to \mathcal{L}_\circ . But since the inference rules (Cong) and (Coh) are closed under \vee -forms, we know (see Section 2) that L_\circ is also semilinear and hence it is complete with respect to the class of L_\circ -chains.

Proposition 4.4 (Chain completeness) *The logic L_\circ is strongly complete with respect to the class of L_\circ -chains.*

It is worth pointing out that the above conditions on \circ in a linearly ordered L_\circ -algebra faithfully capture the three intended properties (c1)-(c3) that were required to such \circ operator at the beginning of this section. Indeed, one can easily show the following lemma.

Lemma 4.5 *Let \mathbf{A} be a L -chain and let $\circ : A \rightarrow A$ a mapping. Then \circ satisfies conditions (c1), (c2) and (c3) iff \mathbf{A} expanded with \circ is a L_\circ -chain.*

Proof: The implication from left to right is immediate since each condition (ci) implies condition ($\circ i$) for $i = 1, 2, 3$, actually (c2) = ($\circ 2$). For the other direction, it is enough to observe that in a chain it holds that $x \wedge y = 0$ iff $x = 0$ or $y = 0$, and $x \vee y = 1$ iff either $x = 1$ or $y = 1$. Then it is obvious that ($\circ 1$) and ($\circ 3$) are indeed equivalent to (c1) and (c3) respectively. \square

Example 4.6 (1) *Let L be the logic of a t -norm which is an ordinal sum of a Lukasiewicz component and a Gödel component with an idempotent separating point $0 < a < 1$ (a non-SMTL chain denoted $L \oplus G$ such that $N(L \oplus G) = [a, 1)$). Then an \circ operator in the corresponding standard algebra is any function $\circ : [0, 1] \rightarrow [0, 1]$ such that :*

$$(i) \quad \circ(x) = 1 \text{ if } x \in \{0, 1\},$$

$$(ii) \quad \circ(x) = 0 \text{ if } x \in (0, a) \text{ (where } x \wedge \neg x \neq 0),$$

$$(iii) \quad \circ \text{ is not decreasing in } N(L \oplus G) = [a, 1) \text{ (where } \neg x = 0).$$

Therefore there are as many consistency operators as non-decreasing functions over the interval $[a, 1]$ with values in $[0, 1]$.

(2) Let $L = L$ be Lukasiewicz logic, i.e. the logic of the Lukasiewicz t -norm complete with respect to the standard chain $[0, 1]_L$. Since the negation is involutive, we have $N([0, 1]_L) = \emptyset$, and thus there is a unique \circ operator definable on the Lukasiewicz standard chain: the one defined as $\circ(x) = 1$ if $x \in \{0, 1\}$, and $\circ(x) = 0$ otherwise.

We can now prove that the logic L_\circ is a conservative expansion of L in the following strong sense.

Proposition 4.7 (Conservative expansion) *Let \mathcal{L} be the language of L . For every set $\Gamma \cup \{\varphi\}$ of \mathcal{L} -formulas, $\Gamma \vdash_{L_\circ} \varphi$ iff $\Gamma \vdash_L \varphi$.*

Proof: One implication is trivial. For the other one, assume that $\Gamma \not\vdash_L \varphi$. Then there exists an L -chain \mathbf{A} and an \mathbf{A} -evaluation e such that $e[\Gamma] \subseteq \{\bar{1}\}$ and $e(\varphi) \neq \bar{1}$. \mathbf{A} can be expanded to an L_\circ -chain \mathbf{A}' e.g. by defining $\circ(1) = \circ(0) = 1$ and $\circ(x) = 0$ for every $x \in A \setminus \{0, 1\}$. Then \mathbf{A}' and e provide a counterexample in the expanded language showing that $\Gamma \not\vdash_{L_\circ} \varphi$. \square

Theorem 4.8 (Strong real completeness) *The logic L_\circ has the SRC if, and only if, L has the SRC.*

Proof: Again, one implication just follows from the fact that L_\circ is a conservative expansion of L . For the converse one assume that L has the SRC. We have to show that any countable L_\circ -chain can be embedded into a standard L_\circ -chain. Let \mathbf{A} be a countable L_\circ -chain. By Theorem 2.2, we know that the \circ -free reduct of \mathbf{A} is embeddable into a (standard) L -chain \mathbf{B} on $[0, 1]$. Denote this embedding by f and define $\circ': [0, 1] \rightarrow [0, 1]$ in the following way:

- (i) $\circ'(0) = \circ'(1) = 1$
- (ii) $\circ'(x) = 0$ for x such that $x > 0$ and $\neg x > 0$.
- (iii) \circ' restricted to the interval $\{x \in (0, 1) \mid \neg x = 0\}$ is defined as $\circ'(z) = \sup\{f(\circ(x)) : x \in A, f(x) \leq z\}$

So defined, \circ' is non-decreasing on $\{x \in [0, 1] : \neg x = 0\}$ such that $\circ'(f(x)) = f(\circ(x))$ for any $x \in A$ and hence \mathbf{B} expanded with \circ' is a standard L_\circ -chain where \mathbf{A} is embedded. \square

Taking into account that for a (semilinear core expansion of MTL) logic L being finite strong real completeness is equivalent to the fact that every countable L -chain is partially embeddable into some L -chain over $[0, 1]$ (see Theorem 2.2), the following corollary can be easily proved by the same technique used in the above Theorem 4.8.

Corollary 4.9 (Finite strong standard completeness) *A logic L has the FSRC if, and only if, L_\circ has the FSRC.*

4.2 Some interesting extensions of the logics L_\circ

As shown in the examples above of operators \circ in L_\circ -chains \mathbf{A} , these operators are completely determined over the set $\{x \in A : x \wedge \neg x \neq 0\}$, but they can be defined in different ways in the interval where $\neg x = 0$. In this section we first consider adding a consistency operator \circ to logics whose associated chains have no elements $x < 1$ such that $\neg x = 0$ (chains \mathbf{A} such that $N(\mathbf{A}) = \emptyset$). These logics can be obtained from any L by adding a suitable inference rule, and will be denoted as $L^{\neg\neg}$. In the second subsection we focus on logics L_\circ where \circ is crisp, and in particular we consider the two extremal cases of these operators, namely those such that $\circ(x) = 0$ for all $x \in N(\mathbf{A})$ and those such that $\circ(x) = 1$ for all $x \in N(\mathbf{A})$.

4.2.1 The case of $L^{\neg\neg}$ logics with \circ operators

In this subsection we study the case of logics L_\circ whose associated L -chains are those where $\neg x = 0$ necessarily implies that $x = 1$. First, from a logic L we will define the logic $L^{\neg\neg}$ and then we will add the consistency operator.

The logic $L^{\neg\neg}$ is defined as the extension of L by adding the following rule:

$$(\neg\neg) \quad \frac{\neg\neg\varphi}{\varphi}$$

Obviously $L^{\neg\neg}$ is complete with respect to the corresponding quasi-variety of $L^{\neg\neg}$ -algebras, that is, the class of L -algebras satisfying the quasi-equation “If $\neg\neg x = 1$ then $x = 1$ ”, or equivalently the quasi-equation “If $\neg x = 0$ then $x = 1$ ”.

Remark 4.10 *In general, the class of $L^{\neg\neg}$ -algebras is not a variety. For instance, in [24] it is proved that the class of weak nilpotent minimum algebras satisfying the quasi-equation $\neg x = 0 \Rightarrow x = 1$ is a quasi-variety that is not a variety. For instance, take the WNM-chain \mathbf{C} over the real unit interval defined by the negation:*

$$n(x) = \begin{cases} 1 - x & \text{if } x \in [0, \frac{1}{5}] \cup [\frac{4}{5}, 1] \\ \frac{1}{5}, & \text{if } x \in [\frac{3}{5}, \frac{4}{5}] \\ \frac{4}{5} - x, & \text{if } x \in [\frac{1}{5}, \frac{3}{5}] \end{cases}$$

Take the filter $F = [\frac{4}{5}, 1]$. Then an easy computation shows that the quotient algebra \mathbf{C}/\equiv_F is isomorphic to the standard WNM-chain \mathbf{C}_F defined by the negation:

$$n_F(x) = \begin{cases} 0, & \text{if } x \in [\frac{2}{3}, 1] \\ \frac{2}{3} - x, & \text{if } x \in (0, \frac{2}{3}] \end{cases}$$

Clearly $\{x \in C : \neg x = 0\} = \{1\}$ but $\{x \in C_F : \neg x = 0\} \neq \{1\}$, i.e. \mathbf{C} belongs to the quasi-variety of $WNM^{\neg\neg}$ -algebras but \mathbf{C}_F does not, so the class of $L^{\neg\neg}$ -algebras is not closed by homomorphisms.

Moreover $L^{\neg\neg}$ is a semilinear logic since it satisfies the following proposition.

Lemma 4.11 *The following rule*

$$(\neg\neg)^\vee \frac{\neg\neg\varphi \vee \delta}{\varphi \vee \delta}$$

is derivable in $L^{\neg\neg}$.

Proof: Since $\delta \rightarrow \neg\neg\delta$ is a theorem of MTL, it is clear that $\neg\neg\varphi \vee \delta \vdash_L \neg\neg\varphi \vee \neg\neg\delta$, and so $\neg\neg\varphi \vee \delta \vdash_L \neg\neg(\varphi \vee \delta)$ as well. Then, by using the rule $(\neg\neg)$ we have that $\varphi \vee \delta$ is derivable in $L^{\neg\neg}$ from the premise $\neg\neg\varphi \vee \delta$. \square

Corollary 4.12 (Chain completeness) *The logic $L^{\neg\neg}$ is semilinear and thus strongly complete with respect to the class of $L^{\neg\neg}$ -chains.*

Proof: Since the inference rule $(\neg\neg)$ is closed under \vee -forms, we know that $L^{\neg\neg}$ is also semilinear (see Section 2) and hence it is complete with respect to the class of $L^{\neg\neg}$ -chains \square

Remark 4.13 *Obviously, if L is an IMTL logic (i.e. a logic where its negation is involutive), then $L^{\neg\neg} = L$. Also, for interested readers, we could notice that $BL^{\neg\neg}$ is actually Łukasiewicz logic L since the only BL -chains satisfying the quasi-equation “if $\neg x = 0$ then $x = 1$ ” are the involutive BL -chains, i.e. MV -chains. This is not the case for $MTL^{\neg\neg}$ which is not equivalent to IMTL (see the WNM logic defined in the previous remark, that satisfy rule $(\neg\neg)$ and it is not IMTL.).*

Now we add the consistency operator \circ to the logic $L^{\neg\neg}$. By this we mean to expand the language with an unary connective \circ and to add the axioms (A1), (A2) and (A3) and the inference rules (Cong) and (Coh). Obviously, the resulting logic $L_\circ^{\neg\neg}$ is complete with respect to the quasi-variety of $L_\circ^{\neg\neg}$ -algebras and with respect to the class of chains of the quasi-variety. The completeness theorems with respect to real chains also apply to $L_\circ^{\neg\neg}$. Moreover we can easily prove that the following schemes and inference rule are provable and derivable respectively in $L_\circ^{\neg\neg}$:

- (B1) $\neg \circ\varphi \vee \varphi \vee \neg\varphi$,
- (B2) $\circ(\varphi \leftrightarrow \psi) \rightarrow (\circ\varphi \leftrightarrow \circ\psi)$,
- (B3) $\circ(\varphi \vee \psi) \rightarrow \circ\varphi \vee \psi$,
- (B4) $\circ\bar{0}$
- (\circ Nec) $\frac{\varphi}{\circ\varphi}$

These properties allows us to provide a simpler axiomatization of $L_{\circ}^{\neg\neg}$.

Theorem 4.14 $L_{\circ}^{\neg\neg}$ can be axiomatized by adding to the axiomatization of $L^{\neg\neg}$ the axioms (B1)-(B4) and the rule (\circ Nec).

Proof: Let us denote by L_{\circ}^+ the resulting new system in the expanded language with \circ obtained from $L^{\neg\neg}$ by adding the axioms (B1)-(B4) and the rule (\circ Nec). The axioms (B1)-(B4) and the rule (\circ Nec) are clearly sound wrt $L_{\circ}^{\neg\neg}$ -algebras. Thus we need only to prove that axioms of $L_{\circ}^{\neg\neg}$ are provable in the new system L_{\circ}^+ , and that the rules (Cong) and (Coh) are also admissible in L_{\circ}^+ . It is obvious that from (B1) we can obtain (A1), since $\neg\circ\varphi \vee \varphi \vee \neg\varphi$ implies $\neg\circ\varphi \vee \neg\neg\varphi \vee \neg\varphi$ and the latter is equivalent to (A1). (A2) is an easy consequence of rule (\circ Nec), and (A3) is (B4). Thus it only remains to prove that (Cong) and (Coh) are derivable in L_{\circ}^+ (in what follows \vdash stands for $\vdash_{L_{\circ}^+}$).

On the one hand, from $(\varphi \leftrightarrow \psi) \vee \delta$, using rule (\circ Nec), we obtain $\circ((\varphi \leftrightarrow \psi) \vee \delta)$, and by (B3) and MP, $\circ(\varphi \leftrightarrow \psi) \vee \delta$. Finally, by (B2), MP and taking into account the monotonicity of \vee , we get $(\circ\varphi \leftrightarrow \circ\psi) \vee \delta$. Hence (Cong) is derivable.

On the other hand, from $\neg\neg\varphi \vee \delta$ and $(\varphi \rightarrow \psi) \vee \delta$, using $(\neg\neg)^{\vee}$, we obtain $\varphi \vee \delta$ and $(\varphi \rightarrow \psi) \vee \delta$, and thus $(\varphi \vee \delta) \& ((\varphi \rightarrow \psi) \vee \delta)$ as well. Therefore by properties of $\&$, we get $(\varphi \& ((\varphi \rightarrow \psi) \vee \delta)) \vee (\delta \& ((\varphi \rightarrow \psi) \vee \delta))$, and by MP and monotonicity for $\&$, we obtain $\psi \vee \delta$. Now, by (\circ Nec) it follows $\circ(\psi \vee \delta)$ and by (B3), $\circ\psi \vee \delta$. Since MTL proves the schema $\alpha \rightarrow (\beta \rightarrow \alpha)$ then $\circ\psi \rightarrow (\circ\varphi \rightarrow \circ\psi)$ is a theorem of L_{\circ}^+ . Thus, by monotonicity of \vee and modus ponens, we obtain $(\circ\varphi \rightarrow \circ\psi) \vee \delta$. Therefore (Coh) is a derivable rule in $\text{MTL}_{\circ}^{\neg\neg}$, and hence in $L_{\circ}^{\neg\neg}$ as desired. \square

Taking into account that $L_{\circ}^{\neg\neg}$ is chain-complete, it is interesting to check how operators \circ can be defined in a $L^{\neg\neg}$ -chain \mathbf{A} . Indeed, since in this case $N(\mathbf{A}) = \emptyset$, the \circ operator is completely determined and defined as:

$$\circ(x) = \begin{cases} 1, & \text{if } x \in \{0, 1\} \\ 0, & \text{otherwise} \end{cases}$$

The interested reader will have observed that such an operator can also be defined in the algebras of the logic L_{Δ} (the expansion of L with the

Monteiro-Baaz Δ operator) as $\circ(x) = \Delta(x \vee \neg x)$ (cf. [15]). And conversely, in $L_{\circ}^{\neg\neg}$ -algebras the Δ operator is also definable as $\Delta x = \circ(x) \wedge x$. Therefore the following result is easy to prove using chain completeness results for both logics.

Corollary 4.15 *$L_{\circ}^{\neg\neg}$ -algebras and $(L_{\Delta})^{\neg\neg}$ -algebras are termwise equivalent, hence the logics $L_{\circ}^{\neg\neg}$ and $(L_{\Delta})^{\neg\neg}$ themselves are equivalent.*

As a consequence, let us mention that, unlike $L^{\neg\neg}$, the class of $L_{\circ}^{\neg\neg}$ -algebras is always a variety, since this is clearly the case of $(L_{\Delta})^{\neg\neg}$: indeed, the rule $(\neg\neg)$ can be equivalently expressed in $(L_{\Delta})^{\neg\neg}$ as the axiom $\Delta(\neg\neg\varphi) \rightarrow \varphi$.

4.2.2 Logics with crisp consistency operators: minimal and maximal consistency operators

As previously observed, the consistency operator \circ is non-decreasing in the segments of the chains where $\neg x = 0$, producing a kind of ‘fuzzy degree of classicality’. In the previous section we have analyzed a special case where the operator \circ is crisp in the sense that it takes only the values 0 and 1. The aim of this section is to study the general case where \circ is crisp.

Definition 4.16 *Let L_{\circ}^c be the logic obtained from L_{\circ} by adding the following axiom:*

$$(c) \quad \circ\varphi \vee \neg\circ\varphi$$

A L_{\circ}^c -algebra \mathbf{A} is a L_{\circ} -algebra such that $\circ(x) \vee \neg\circ(x) = 1$ for every $x \in A$.

Since it is an axiomatic extension of the logic L_{\circ} , it turns out that L_{\circ}^c is algebraizable, whose equivalent algebraic semantics is given by the quasi-variety of L_{\circ}^c -algebras, and semilinear as well, and thus complete with respect to the class of L_{\circ}^c -chains. From the definition above, it is clear that the operator \circ in any L_{\circ}^c -chain A is such that $\circ(x) \in \{0, 1\}$ for every $x \in A$. Moreover, this implies that the set $\{x \in A \setminus \{0\} : \circ(x) = 1\}$ is an interval containing $N(A) \cup \{1\}$.

Let us consider now the logics corresponding to the minimal and maximal (pointwisely) consistency operators, as announced in the introduction of Section 4.2. First, consider L_{\circ}^{min} to be the axiomatic extension of the logic L_{\circ} with the following axiom:

$$(A4) \quad \varphi \vee \neg\varphi \vee \neg\circ\varphi$$

Since it is an axiomatic extension of L_{\circ} , L_{\circ}^{min} is complete with respect to the class of L_{\circ}^{min} -chains, i.e. L_{\circ} -chains satisfying the equation $x \vee \neg x \vee \neg\circ(x) =$

1. One can readily check that the equation $x \vee \neg x \vee \neg \circ(x) = 1$ holds in an L_\circ -chain only in the case that $\circ(x) = 0$ when $0 < x < 1$. Indeed, if $\min(x, \neg x) > 0$ it is clear that $\circ(x)$ has to be 0, while if $x < 1$ and $\neg x = 0$ then (A4) forces $\neg \circ(x) = 1$, that is $\circ(x) = 0$. Therefore, the \circ -operator in any L_\circ^{min} -chain is completely determined, and it is indeed the (pointwisely) minimal one definable in a L_\circ -chain.

Proposition 4.17 *The logic L_\circ^{min} is complete with respect to the class of L_\circ^{min} -chains, i.e. L_\circ -chains where the \circ operator is the minimal one.*

Since L_\circ^{min} are a special kind of L_\circ^c -chains, this proposition yields that L_\circ^{min} must be an axiomatic extension of L_\circ^c . Moreover, it turns out that, for all x in a L_\circ^{min} -chain, $\circ(x)$ coincides with $\Delta(x \vee \neg x)$, where Δ is the Baaz-Monteiro projection operator, as it happened in the case of $L_\circ^{\neg\neg}$ -chains. Using the fact that both logics L_\circ^{min} and L_Δ are chain-complete, it follows that they are inter-definable.

Proposition 4.18 *The logics L_\circ^{min} and L_Δ are inter-definable by means of the following translations:*

(i) from L_\circ^{min} to L_Δ : define $\Delta\varphi$ as $\varphi \wedge \circ\varphi$

(ii) from L_Δ to L_\circ^{min} : define $\circ\varphi$ as $\Delta(\varphi \vee \neg\varphi)$.

By (ii) of the above proposition $\circ\varphi$ is equivalent to the formula $\Delta(\varphi \vee \neg\varphi)$ in L_Δ . Thus by axiom ($\Delta 1$) the following result (proving the axiom of L_\circ^c) is obvious.

Lemma 4.19 *L_\circ^{min} proves the axiom (c), i.e. $\circ\varphi \vee \neg\circ\varphi$.*

Finally, consider the logic L_\circ^{max} to be the extension of the logic L_\circ with the following inference rule:

$$(\neg\neg_\circ) \quad \frac{\neg\neg\varphi \vee \delta}{\circ\varphi \vee \delta}$$

Again, since $(\neg\neg_\circ)^\vee$ is closed under disjunction, L_\circ^{max} is complete with respect to L_\circ^{max} -chains, i.e. L_\circ -chains where the following condition holds: if $\neg x = 0$ then $\circ(x) = 1$. Since $\circ(x) = 0$ for all $x > 0$ such that $\neg x > 0$, then it is clear that \circ is completely determined in such a chain and defined as: $\circ(x) = 0$ if $0 < x \wedge \neg x$ and $\circ(x) = 1$ otherwise (i.e. if $x \in \{0, 1\}$ or $\neg x = 0$). Hence \circ is the maximal (pointwisely) consistency operator definable in a L -chain.

Proposition 4.20 *The logic L_\circ^{max} is complete with respect to the class of L_\circ^{max} -chains, i.e. L_\circ -chains where the \circ operator is the maximal one.*

As a final remark, we notice that in case L is an extension of the basic fuzzy logic BL, the above rule $(\neg\neg\circ)$ can be equivalently replaced by the following axiom:

$$(\neg\neg\varphi \rightarrow \varphi) \vee \circ\varphi$$

Indeed, it is not difficult to check that, given the special features of negations in BL-chains, a consistency operator \circ in a BL-chain \mathbf{A} satisfies this axiom iff $\circ(x) = 0$ for x such that $0 < \min(x, \neg x)$ and $\circ(x) = 1$ otherwise. Therefore, the quasivariety of L_{\circ}^{max} -algebras is in fact a variety when L is a BL-extension, but whether the class of L_{\circ}^{max} -algebras is a variety in a more general case remains as an open problem.

Figure 2 gathers the axiomatizations (relative to L) of the logic L_{\circ} and of the different extensions we have defined in Section 4.2.

Remark 4.21 *Taking into account the graphs of the \circ operators on real chains associated to the logics $L_{\circ}^{\neg\neg}$, L_{\circ}^c , L_{\circ}^{min} and L_{\circ}^{max} (see Figure 2), and the proof of conservativeness of L_{\circ} with respect to L in Proposition 4.7, it is clear that the same kind of proof also applies to all these extensions of L_{\circ} . Hence $L_{\circ}^{\neg\neg}$ is a conservative expansion of $L^{\neg\neg}$ and L_{\circ}^c , L_{\circ}^{min} and L_{\circ}^{max} are conservative expansions of L .*

4.3 Axiomatizing paraconsistent fuzzy logics L^{\leq} with \circ

As mentioned in the introduction of this section we know that the only paraconsistent fuzzy logic are the logics L^{\leq} when L is not an SMTL logic. Thus our ultimate goal is the axiomatization of the expansion of paraconsistent logics L^{\leq} with a consistency operator \circ , that will be denoted L_{\circ}^{\leq} . But from results of this section we know how to axiomatize the logics L_{\circ} and, as shown in Section 2 we know how to get an axiomatization of L_{\circ}^{\leq} from the one of L_{\circ} . Indeed the axiomatization of L_{\circ}^{\leq} is obtained by taking the same axioms of L_{\circ} and adding the following inference rules:

(Adj- \wedge) from φ and ψ deduce $\varphi \wedge \psi$

(MP- r) if $\vdash_{L_{\circ}} \varphi \rightarrow \psi$ (i.e. if $\varphi \rightarrow \psi$ is a theorem of L_{\circ}), then from φ derive ψ

(Cong- r) if $\vdash_{L_{\circ}} (\varphi \leftrightarrow \psi) \vee \delta$ then derive $(\circ\varphi \leftrightarrow \circ\psi) \vee \delta$

(Coh- r) if $\vdash_{L_{\circ}} (\neg\neg\varphi \wedge (\varphi \rightarrow \psi)) \vee \delta$ then derive $(\circ\varphi \rightarrow \circ\psi) \vee \delta$

In the same way we could obtain axiomatizations of the logics L_{\circ}^{\leq} when L is any of the logics studied in this section. Then axiomatizations of the logics $(L_{\circ}^{\neg\neg})^{\leq}$, $(L_{\circ}^c)^{\leq}$, $(L_{\circ}^{min})^{\leq}$ and $(L_{\circ}^{max})^{\leq}$ are easily obtained. Figure 3

Logic	Definition	Operator \circ
L_{\circ}	$L +$ (A1) $\neg(\varphi \wedge \neg\varphi \wedge \circ\varphi)$ (A2) $\circ\bar{1}$ (A3) $\circ\bar{0}$ (Cong) $\frac{(\varphi \leftrightarrow \psi) \vee \delta}{(\circ\varphi \leftrightarrow \circ\psi) \vee \delta}$ (Coh) $\frac{(\neg\neg\varphi \wedge (\varphi \rightarrow \psi)) \vee \delta}{(\circ\varphi \rightarrow \circ\psi) \vee \delta}$	
$L_{\circ}^{\neg\neg}$	$L_{\circ} +$ $(\neg\neg)$ $\frac{\neg\neg\varphi}{\varphi}$	
L_{\circ}^c	$L_{\circ} +$ (c) $\circ\varphi \vee \neg\circ\varphi$	
L_{\circ}^{min}	$L_{\circ} +$ (A4) $\varphi \vee \neg\varphi \vee \neg\circ\varphi$	
L_{\circ}^{max}	$L_{\circ} +$ $(\neg\neg_{\circ})$ $\frac{\neg\neg\varphi \vee \delta}{\circ\varphi \vee \delta}$	

Figure 2: Summary of the axiomatizations of the logics L_{\circ} , $L_{\circ}^{\neg\neg}$, L_{\circ}^c , L_{\circ}^{min} , L_{\circ}^{max} , and corresponding graphs of the operators \circ on a standard BL-chain, where $0 < a \leq 1$.

gathers the inference rules of these logics (recall that the axioms coincide with those of the corresponding truth-preserving logics). Therefore we have defined and axiomatized a general family of LFIs based on fuzzy logics and some of its extensions.

Logic	Inference rules
L_{\circ}^{\leq}	rules of L^{\leq} + (Cong-r) $\frac{\vdash_{L_{\circ}} (\varphi \leftrightarrow \psi) \vee \delta}{(\circ\varphi \leftrightarrow \circ\psi) \vee \delta}$ (Coh-r) $\frac{\vdash_{L_{\circ}} (\neg\neg\varphi \wedge (\varphi \rightarrow \psi)) \vee \delta}{(\circ\varphi \rightarrow \circ\psi) \vee \delta}$
$(L_{\circ}^{\neg\neg})^{\leq}$	rules of L_{\circ}^{\leq} + $(\neg\neg\text{-}r)$ $\frac{\vdash_{L_{\circ}^{\neg\neg}} \neg\neg\varphi}{\varphi}$
$(L_{\circ}^c)^{\leq}$	rules of L_{\circ}^{\leq}
$(L_{\circ}^{\min})^{\leq}$	rules of L_{\circ}^{\leq}
$(L_{\circ}^{\max})^{\leq}$	rules of L_{\circ}^{\leq} + $(\neg\neg\text{-}r)$ $\frac{\vdash_{L_{\circ}^{\max}} \neg\neg\varphi \vee \delta}{\circ\varphi \vee \delta}$

Figure 3: Summary of the inference rules of the logics L_{\circ}^{\leq} , $(L_{\circ}^{\neg\neg})^{\leq}$, $(L_{\circ}^c)^{\leq}$, $(L_{\circ}^{\min})^{\leq}$ and $(L_{\circ}^{\max})^{\leq}$.

Finally, recall that, as observed at the end of Section 2, if a logic L' is a conservative expansion of another L , then L'^{\leq} is also a conservative expansion of L^{\leq} . Therefore, taking into account Proposition 4.7 and Remark 4.21, we get the following immediate result.

Proposition 4.22 *The logics L_{\circ}^{\leq} , $(L_{\circ}^c)^{\leq}$, $(L_{\circ}^{\min})^{\leq}$ and $(L_{\circ}^{\max})^{\leq}$ are conservative expansions of L^{\leq} , while $(L_{\circ}^{\neg\neg})^{\leq}$ is a conservative expansion of $(L^{\neg\neg})^{\leq}$.*

5 About the propagation property

One of the distinctive features of da Costa's C-systems is the so-called *propagation property* of the consistency connective \circ , which states that consistency (or well-behavior, according to da Costa's terminology) is propagated

in the following sense: from $\circ\varphi$ it follows $\circ\neg\varphi$, and from $\{\circ\varphi, \circ\psi\}$ it follows $\circ(\varphi\#\psi)$, for every binary connective $\#$. We can adapt this property to our setting, and study conditions which ensure its validity.

Definition 5.1 *Let L be a paraconsistent fuzzy logic with a consistency operator \circ . Then we say that \circ satisfies the propagation property in L with respect to a subset X of connectives of the language of L if*

$$\circ\varphi_1, \dots, \circ\varphi_n \vdash_L \circ\#(\varphi_1, \dots, \varphi_n),$$

for every n -nary connective $\# \in X$ and formulas $\varphi_1, \dots, \varphi_n$ built with connectives from X .

Observe that in the case $n = 0$, $\#$ is a constant and that the above condition requires $\vdash_L \circ\#$.

The paraconsistent fuzzy logics studied in this paper are logics L_{\circ}^{\leq} and some extensions, where L is a truth-preserving fuzzy logic. Knowing the relation between truth-preserving and degree-preserving fuzzy logics, \circ satisfies the *propagation property* in a logic L_{\circ}^{\leq} with respect to some subset of connectives $X \subseteq \{\bar{0}, \wedge, \&, \rightarrow\}$ ⁴ whenever:

$$\begin{cases} \vdash_{L_{\circ}} \circ\bar{0} & \text{if } \bar{0} \in X \\ \vdash_{L_{\circ}} (\circ\varphi \wedge \circ\psi) \rightarrow \circ(\varphi\#\psi), & \text{for each binary } \# \in X \end{cases} \quad (\text{Prop}^*)$$

In such a case we will say that \circ satisfies the propagation property (Prop*) in the logic L_{\circ}^{\leq} with respect to the set of connectives X . Actually since in the logics L_{\circ} we have $\vdash_L \circ\bar{0}$, the first condition is always satisfied and it can be dropped from (Prop*).

Proposition 5.2 *The following conditions hold:*

1. \circ satisfies (Prop*) in any fuzzy logic L_{\circ} with respect to the set of primitive connectives $\{\wedge, \rightarrow\}$.
2. \circ satisfies (Prop*) in any fuzzy logic of the families $L_{\circ}^{\neg\neg}$, L_{\circ}^{\min} or L_{\circ}^{\max} with respect to the set of primitive connectives $\{\wedge, \&, \rightarrow\}$.

Proof: Due to chain completeness of the logics involved, the whole proof is done by algebraic means. Let \mathbf{A} be a L_{\circ} -chain. The first item for the connective \wedge is a consequence of the non-decreasing property of \circ in $A \setminus \{0\}$ combined with the fact that $\circ(0) = 1$. Before dealing with the property for \rightarrow , we first consider the property for the negation \neg , that is proved by cases. If $x \in \{0, 1\}$ the property is obvious. If $x \in N(\mathbf{A})$ then $\neg x = 0$, and

⁴We are assuming here to work with a core fuzzy logic, and in core fuzzy logics these are the primitive connectives, the rest are definable from them.

thus $\circ(\neg x) = 1$ and the property obviously holds. Otherwise, if x does not belong to the previous cases $x \wedge \neg x > 0$ then $\circ(x) = 0$ and also the property is obviously satisfied. Finally, for \rightarrow the proof is easy since, remembering that a residuated implication satisfies the inequality $y \leq x \rightarrow y$, assuming $y > 0$ we have $\circ(x) \wedge \circ(y) \leq \circ(y) \leq \circ(x \rightarrow y)$; if $y = 0$ then we are back to the case of the negation.

For the second item we only need to deal with the case of $\&$. First observe that if $\neg x = 0$ and $\neg y = 0$, then $\neg(x \& y) = 0$ as well, since $\neg(x \& y) = (x \& y) \rightarrow 0 = x \rightarrow (y \rightarrow 0) = x \rightarrow \neg y = x \rightarrow 0 = \neg x = 0$. Second observe that in the chains of the considered logics, the image of \circ is $\{0, 1\}$. From there the proof is easy. If one of the values x, y is 0 or 1 the result is obvious. If one of the values x, y is in $(0, 1) \setminus N(A)$ then $\circ(x) \wedge \circ(y) = 0$, and the implication is trivially valid. Finally if $x, y \in N(A)$ the result follows from the fact that \circ is non-decreasing taking into account the first observation above. \square

The first item of Proposition 5.2 can not be improved in the sense that L_o does not prove $\circ(\varphi) \wedge \circ(\psi) \rightarrow \circ(\varphi \& \psi)$, as the following example shows.

Example 5.3 *Let L be the logic of the t -norm \otimes that is the ordinal sum of a Lukasiewicz component and a product component, with $\frac{1}{2}$ being the idempotent separating point. Then consider the L_o -chain where the consistency operator \circ is defined by $\circ(x) = 0$ if $x \in (0, \frac{3}{4})$ and $\circ(x) = 1$ otherwise. Take now $x = \frac{5}{6}$ and $y = \frac{3}{4}$. Then $\frac{5}{6} \otimes \frac{3}{4} < \frac{3}{4}$, and clearly $\circ(x) = \circ(y) = 1$ while $\circ(\frac{5}{6} \otimes \frac{3}{4}) = 0$.*

6 Recovering Classical Logic

In the context of LFIs, it is a desirable property to recover the classical reasoning by means of the consistency connective \circ (see [6]). Specifically, let **CPL** be classical propositional logic. If L is a given LFI such that its reduct to the language of **CPL** is a sublogic of **CPL**, then a DAT (Derivability Adjustment Theorem) for L with respect to **CPL** is as follows: for every finite set of formulas $\Gamma \cup \{\varphi\}$ in the language of **CPL**, there exists a finite set of formulas Θ in the language of L , whose variables occur in formulas of $\Gamma \cup \{\varphi\}$, such that

$$(DAT) \quad \Gamma \vdash_{\mathbf{CPL}} \varphi \text{ iff } \circ(\Theta), \Gamma \vdash_L \varphi.$$

When the operator \circ enjoys the propagation property in the logic L with respect to the classical connectives (see the previous section) then the DAT takes the following, simplified form: for every finite set of formulas $\Gamma \cup \{\varphi\}$ in the language of **CPL**,

$$(PDAT) \quad \Gamma \vdash_{\mathbf{CPL}} \varphi \text{ iff } \circ p_1, \dots, \circ p_n, \Gamma \vdash_L \varphi$$

where $\{p_1, \dots, p_n\}$ is the set of propositional variables occurring in $\Gamma \cup \{\varphi\}$.

Here we are interested in investigating whether we can expect some form of the (PDAT) for the logics L_\circ^\leq and, as we have noted before, \circ has the propagation property iff certain formulas are theorems in L_\circ . Thus in terms of theoremhood, to have a DAT when \circ propagates in L_\circ^\leq is equivalent to prove the following:

$$(\text{PDAT}^*) \quad \vdash_{\text{CPL}} \varphi \text{ iff } \vdash_{L_\circ} \left(\bigwedge_{i=1}^n \circ p_i \right) \rightarrow \varphi \text{ iff } \bigwedge_{i=1}^n \circ p_i \vdash_{L_\circ^\leq} \varphi$$

where $\{p_1, \dots, p_n\}$ is the set of propositional variables occurring in φ (obviously, when this set is empty, that is, when $n = 0$, then $\bigwedge_{i=1}^n \circ p_i$ is set to be $\bar{1}$). Arguably, (PDAT) (or (PDAT*)) is more interesting than (DAT). For instance, the hierarchy $(C_n)_{n \geq 1}$ of paraconsistent logics introduced by da Costa satisfies (PDAT).

Since \circ satisfies the propagation property in logics L_\circ with respect to the classical signature (cf. Proposition 5.2), we try to prove (PDAT*) for them. However, in the general setting of logics L_\circ (PDAT*) does not always hold. Indeed it is easy to see that $\vdash_{\text{CPL}} p \vee \neg p$ but $\not\vdash_{L_\circ} \circ p \rightarrow (p \vee \neg p)$, i.e. $\circ p \rightarrow (p \vee \neg p)$ is not a tautology over all L_\circ -chains. Take for example the L_\circ -chain defined on a L -chain \mathbf{A} by defining \circ as follows: $\circ(x) = 1$ if $x \in \{0, 1\} \cup N(\mathbf{A})$ and $\circ(x) = 0$ otherwise. Then it is easy to see that if $N(\mathbf{A}) \neq \emptyset$ then $e(\circ p \rightarrow (p \vee \neg p)) \neq 1$ for any evaluation e on \mathbf{A} such that $e(p) \in N(\mathbf{A})$.

This example is significative since the principle $\varphi \vee \neg \varphi$ is enough to collapse MTL-logic with classical logic. In fact we propose the following definition.

Definition 6.1 *Let L_\circ^{dat} be the logic obtained from L_\circ by adding the following axiom:*

$$(\circ\text{EM}) \quad \circ \varphi \rightarrow (\varphi \vee \neg \varphi)$$

By the same argument as above, L_\circ^{dat} is algebraizable and its algebraic semantics is given by the class of L_\circ^{dat} -algebras.

Definition 6.2 *A L_\circ^{dat} -algebra \mathbf{A} is a L_\circ -algebra such that $\circ(x) \leq x \vee \neg x$ for every $x \in A$.*

Therefore L_\circ^{dat} is complete with respect to the variety of L_\circ^{dat} -algebras and, more important, with respect to the chains of the variety (since the logic is an axiomatic extension and thus it is semilinear as L_\circ).

Moreover since L_\circ^{dat} extends L_\circ , it follows that \circ satisfies the propagation property in L_\circ^{dat} with respect to the classical signature.

However, since L_{\circ}^{dat} does not satisfy contraction, property (PDAT*) will be hardly satisfied: it should be intuitively clear that, in some situations, it could be necessary to use the ‘consistency assumption’ (and so the law of excluded middle) more than once in order to obtain a given tautology. We show next that a slightly modified form of DAT indeed holds for L_{\circ}^{dat} .

Proposition 6.3 *The logic L_{\circ}^{dat} satisfies the following form of DAT:*

$$(PDAT^{**}) \quad \vdash_{\mathbf{CPL}} \varphi \text{ iff there is } k \geq 1 \text{ such that } \vdash_{L_{\circ}^{dat}} \left(\bigwedge_{i=1}^n \circ p_i \right)^k \rightarrow \varphi$$

where $\{p_1, \dots, p_n\}$ is the set of propositional variables occurring in φ and ψ^k is as a shorthand for $\psi \& \dots \& \psi$.

Proof: Let φ be a formula in the language of **CPL** and suppose that p_1, p_2, \dots, p_n are the propositional variables appearing in φ . If $\vdash_{\mathbf{CPL}} \varphi$ then $\{p_i \vee \neg p_i : i = 1, 2, \dots, n\} \vdash_L \varphi$, since for any evaluation e in any L-chain, $e(p_i \vee \neg p_i) = 1$ iff $e(p_i)$ is either 0 or 1. Then by the local deduction-detachment theorem of L, there is a natural k such that $\vdash_L (\bigwedge_{i=1}^n (p_i \vee \neg p_i))^k \rightarrow \varphi$, and this theorem is also valid in L_{\circ}^{dat} . Then, by axiom (\circ EM), this implies $\vdash_{L_{\circ}^{dat}} (\bigwedge_{i=1}^n \circ p_i)^k \rightarrow \varphi$, and hence (PDAT**) holds.

Conversely, assume that $\vdash_{L_{\circ}^{dat}} (\bigwedge_{i=1}^n \circ p_i)^k \rightarrow \varphi$ for some $k \geq 1$, and let e be any evaluation on the 2-element Boolean algebra \mathbf{B}_2 . Since \mathbf{B}_2 can be considered as a L_{\circ}^{dat} -chain where $\circ(0) = \circ(1) = 1$, then we have $e((\bigwedge_{i=1}^n \circ p_i)^k \rightarrow \varphi) = 1$. But then we necessarily have $e(\varphi) = 1$, because $e(\bigwedge_{i=1}^n \circ p_i) = 1$. Therefore φ is a **CPL**-tautology and so $\vdash_{\mathbf{CPL}} \varphi$. \square

An easy reasoning shows an analogous result when we have an arbitrary set of premises built from a finite set of propositional variables:

$$\Gamma \vdash_{\mathbf{CPL}} \varphi \text{ iff there is } k \geq 1 \text{ such that } \Gamma \vdash_{L_{\circ}^{dat}} \left(\bigwedge_{i=1}^m \circ p_i \right)^k \rightarrow \varphi$$

where now p_1, \dots, p_m are the propositional variables appearing in $\Gamma \cup \{\varphi\}$.

One interesting question is to know examples where (PDAT**) is only true for some $k \geq 2$. This is an open question in general but we have the following illustrative example. Consider the formula $\varphi = (p \vee \neg p)^k$ with $k > 1$. Since in a logic L_{\circ} both the weak and strong conjunctions \wedge and $\&$ are many-valued generalizations of the classical conjunction, it is clear that $(p \vee \neg p)^k$ is equivalent to the classical tautology $p \vee \neg p$ when we restrict to Boolean evaluations. However there are extensions of BL logic where (\circ EM) is a tautology but $\circ p \rightarrow (p \vee \neg p)^k$ is not. For instance let L be the logic of the BL-chain $[0, 1]_{L \oplus L}$ where $L \oplus L$ is the ordinal sum of two

copies of Łukasiewicz standard algebra $[0, 1]_{\mathbb{L}}$, and let a be the idempotent element separating the two components. Further, take the operator \circ in $[0, 1]$ defined by $\circ(0) = \circ(1) = 1$, $\circ(x) = x$ if $x \in [a, 1]$, and $\circ(x) = 0$ otherwise. An easy computation shows that $\circ p \rightarrow (p \vee \neg p)^2$ is already not a tautology in L_{\circ}^{dat} . Therefore, if we use $\&$ as conjunction symbol in the language of **CPL**, (PDAT*) is not valid in L_{\circ}^{dat} , while obviously (PDAT**) is so. Nevertheless we have not been able to find a similar example when we use \wedge as conjunction symbol the language of **CPL**.

Finally, notice that axiom (\circ EM) is a theorem of the logics $L_{\circ}^{\neg\neg}$ and L_{\circ}^{\min} . Therefore, in these logics $\circ p_i \vee \neg \circ p_i$ is a theorem, hence it is clear that $\circ p_i$ is equivalent to $(\circ p_i)^k$ for any k , and thus we have the following direct corollary.

Corollary 6.4 *Let $\Gamma \cup \{\varphi\}$ be a finite set of formulas in the language of **CPL** and let $\{p_1, \dots, p_m\}$ the set of propositional variables appearing in $\Gamma \cup \{\varphi\}$. Then*

$$\Gamma \vdash_{\mathbf{CPL}} \varphi \text{ iff } \Gamma \vdash_{L_{\circ}^+} \left(\bigwedge_{i=1}^n \circ p_i \right) \rightarrow \varphi$$

where $+ \in \{\neg\neg, \min\}$.

7 Inconsistency operators in the logic MTL

As recalled in Section 3, within the LFIs framework one can also consider an *inconsistency operator* \bullet , dual to the consistency operator \circ , where $\bullet\varphi$ has the intended meaning of $\neg\circ\varphi$ (see [6]).

In this section we show how to add inconsistency operators to MTL-algebras, as well as to its logical counterparts, based on the content of the previous sections in terms of consistency operators.

Definition 7.1 *Given a logic L that is a semilinear core expansion of MTL but not SMTL, we define the logic L_{\bullet} as the expansion of L in a language which incorporates a new unary connective \bullet with the following axioms:*

$$\begin{aligned} (A1') & \quad \neg(\varphi \wedge \neg\varphi) \vee \bullet\varphi \\ (A2') & \quad \neg\bullet\bar{1} \\ (A3') & \quad \neg\bullet\bar{0} \end{aligned}$$

and the following inference rules:

$$\begin{aligned} (\text{Cong}') & \quad \frac{(\varphi \leftrightarrow \psi) \vee \delta}{(\bullet\varphi \leftrightarrow \bullet\psi) \vee \delta} & (\text{Coh}') & \quad \frac{(\neg\neg\varphi \wedge (\varphi \rightarrow \psi)) \vee \delta}{(\bullet\psi \rightarrow \bullet\varphi) \vee \delta} \end{aligned}$$

As in the case of L_{\circ} , due to the presence of the rule (Cong'), L_{\bullet} is a Rasiowa-implicative logic, and thus it is also algebraizable in the sense of Blok and Pigozzi and its algebraic semantics is given by L_{\bullet} -algebras.

Definition 7.2 A L_\bullet -algebra \mathbf{A} is an expansion of a L -algebra with a new unary operation $\bullet : A \rightarrow A$ satisfying the following conditions for all $x, y, z \in A$:

- (•1) $\neg(x \wedge \neg x) \vee \bullet(x) = 1$
- (•2) $\bullet(1) = \bullet(0) = 0$
- (•3) if $(\neg\neg x \wedge (x \rightarrow y)) \vee z = 1$ then $(\bullet(y) \rightarrow \bullet(x)) \vee z = 1$.

Again, since the rules (Cong') and (Coh') are closed under \vee -forms, L_\bullet is complete with respect to the class of L_\bullet -chains. Obviously, the \bullet operators in L -chains have the dual form of the \circ operators (described in Figure 1), and we will not go into further details.

The intended duality between both operators \circ and \bullet is made explicit in the following results.

Proposition 7.3 Let t be a translation map from the language of L_\circ to the language of L_\bullet which replaces \circ by $\neg\bullet$. Conversely, let t' be the translation map in the opposite direction, which replaces \bullet by $\neg\circ$. Then $\Gamma \vdash_{L_\circ} \varphi$ implies that $t(\Gamma) \vdash_{L_\bullet} t(\varphi)$ and $\Gamma' \vdash_{L_\bullet} \varphi'$ implies that $t'(\Gamma') \vdash_{L_\circ} t'(\varphi')$.

Proof: It is enough to prove that: (i) the translation of each axiom of the source logic can be derived in the target logic, and (ii) the translation of each inference rule of the source logic is an inference rule which is derivable in the target logic. The proof easily follows by using that in MTL the following formulas are theorems: $\neg(\varphi \wedge \psi) \leftrightarrow (\neg\varphi \vee \neg\psi)$, $\varphi \rightarrow \neg\neg\varphi$ and $(\varphi \rightarrow \psi) \rightarrow (\neg\psi \rightarrow \neg\varphi)$. \square

Notice however that the above translations do not yield that both logics are equivalent: indeed, the translations t and t' are not in general each other's inverse. This is due to the fact that, e.g. in L_\circ , $\circ\varphi$ in general is not equivalent to $\neg\neg\circ\varphi$. However, in the frame of the logic L_\circ^c and its extensions, where $\circ\varphi$ is Boolean, one can prove $\circ\varphi \leftrightarrow \neg\neg\circ\varphi$, and one can establish their equivalence with their \bullet -dual corresponding logics. Notice that if L is an IMTL logic (i.e. whose negation is involutive), then L_\circ^c coincides with L_\circ itself.

In particular, if we define dual counterparts of the logics $L_\circ^{\neg\neg}$, L_\circ^c , L_\circ^{\min} or L_\circ^{\max} as:

- $L_\bullet^{\neg\neg}$: is the extension of L_\bullet with the rule “from $\neg\neg\varphi$ infer φ ”
- L_\bullet^c : is the axiomatic extension of L_\bullet with the axiom $\bullet\varphi \vee \neg\bullet\varphi$
- L_\bullet^{\max} : is the axiomatic extension of L_\bullet with the axiom $\varphi \vee \neg\varphi \vee \bullet\varphi$
- L_\bullet^{\min} : is the extension of L_\bullet with the rule “from $\neg\neg\varphi \vee \delta$ infer $\neg\bullet\varphi \vee \delta$ ”

then we can list the following equivalences (denoted by \equiv) among logics via the translations t and t' : $L_{\circ}^{\neg\neg} \equiv L_{\bullet}^{\neg\neg}$, $L_{\circ}^c \equiv L_{\bullet}^c$, $L_{\circ}^{\min} \equiv L_{\bullet}^{\max}$ and $L_{\circ}^{\max} \equiv L_{\bullet}^{\min}$.⁵ As a consequence, the (quasi) varieties associated to pairs of equivalent logics are termwise equivalent.

To conclude, just to point out that, as in the case of the consistency operators, the paraconsistent versions of the above logics with the inconsistency operators \bullet would correspond to their degree-preserving counterparts, namely the logics L_{\bullet}^{\leq} , $(L_{\bullet}^{\neg\neg})^{\leq}$, $(L_{\bullet}^c)^{\leq}$, $(L_{\bullet}^{\min})^{\leq}$ and $(L_{\bullet}^{\max})^{\leq}$.

8 Concluding remarks

In this paper we have investigated the possibility of defining paraconsistent logics of formal inconsistency (LFIs) based on systems of mathematical fuzzy logic, in particular by first expanding axiomatic extensions of the fuzzy logic MTL with the characteristic consistency and inconsistency operators of LFIs, and then by considering their degree-preserving versions, that are paraconsistent. Actually, in the same line of [15] and based on a novel perspective, this paper intends to contribute to the study and understanding of the relationships between paraconsistency and fuzziness.

Acknowledgments

The authors are grateful to the anonymous reviewers for their useful suggestions and comments. The authors have been partially supported by the FP7-PEOPLE-2009-IRSES project MaToMUVI (PIRSES-GA-2009-247584). Coniglio was also supported by FAPESP (Thematic Project Log-Cons 2010/51038-0), and by a research grant from CNPq (PQ 305237/2011-0). Esteva and Godo also acknowledge partial support by the MINECO project TIN2012-39348-C02-01.

References

- [1] Alan Ross Anderson and Nuel D. Belnap. *Entailment: The Logic of Relevance and Necessity*, volume 1. Princeton University Press, Princeton, 1975.
- [2] Ofer Arieli, Arnon Avron, and Anna Zamansky. Ideal paraconsistent logics. *Studia Logica*, 99(1-3):31–60, 2011.
- [3] Arnon Avron and Anna Zamansky. Many-valued non-deterministic semantics for first-order Logics of Formal (In)consistency. In Stefano

⁵Keep in mind that, since we have kept the superscripts min and max for the logics that respectively correspond to minimum and maximum \bullet operators, the dual L_{\circ}^{\min} is L_{\bullet}^{\max} and viceversa, the dual of L_{\circ}^{\max} is L_{\bullet}^{\min} .

- Aguzzoli, Agata Ciabattoni, Brunella Gerla, Corrado Manara, and Vincenzo Marra, editors, *Algebraic and Proof-theoretic Aspects of Non-classical Logics. Papers in Honor of Daniele Mundici on the Occasion of His 60th Birthday*, volume 4460 of *Lecture Notes in Computer Science*, pages 1–24. Springer, 2007.
- [4] Willem J. Blok and Don L. Pigozzi. *Algebraizable Logics*, volume 396 of *Memoirs of the American Mathematical Society*. American Mathematical Society, Providence, RI, 1989. Freely downloadable from <http://orion.math.iastate.edu/dpigozzi/>.
- [5] Félix Bou, Francesc Esteva, Josep Maria Font, Àngel Gil, Lluís Godo, Antoni Torrens, and Ventura Verdú. Logics preserving degrees of truth from varieties of residuated lattices. *Journal of Logic and Computation*, 19(6):1031–1069, 2009.
- [6] Walter A. Carnielli, Marcelo E. Coniglio, and João Marcos. Logics of Formal Inconsistency. In Dov Gabbay and Franz Guenther, editors, *Handbook of Philosophical Logic (2nd. edition)*, volume 14, pages 1–93. Springer, 2007.
- [7] Walter A. Carnielli and João Marcos. A taxonomy of \mathbf{C} -systems. In Walter A. Carnielli, Marcelo E. Coniglio, and Itala M. L. D’Ottaviano, editors, *Paraconsistency - the Logical Way to the Inconsistent*, volume 228 of *Lecture Notes in Pure and Applied Mathematics*, pages 1–94, New York, 2002. Marcel Dekker.
- [8] Roberto Cignoli, Francesc Esteva, Lluís Godo, and Antoni Torrens. Basic fuzzy logic is the logic of continuous t-norms and their residua. *Soft Computing*, 4(2):106–112, 2000.
- [9] Petr Cintula, Francesc Esteva, Joan Gispert, Lluís Godo, Franco Montagna, and Carles Noguera. Distinguished algebraic semantics for t-norm based fuzzy logics: Methods and algebraic equivalencies. *Annals of Pure and Applied Logic*, 160(1):53–81, 2009.
- [10] Petr Cintula, Christian G. Fermüller, Lluís Godo, and Petr Hájek, editors. *Understanding Vagueness*, volume 36 of *Studies in Logic, Mathematical Logic and Foundations*. College Publications, London, 2011.
- [11] Petr Cintula, Rostislav Horčík, and Carles Noguera. The quest for the basic fuzzy logic. In Franco Montagna, editor, *Petr Hájek on Mathematical Fuzzy Logic*, Trends in Logic. Springer, 2014. To appear.
- [12] Petr Cintula and Carles Noguera. A general framework for mathematical fuzzy logic. In Petr Cintula, Petr Hájek, and Carles Noguera, editors, *Handbook of Mathematical Fuzzy Logic - Volume 1*, volume 37

of *Studies in Logic, Mathematical Logic and Foundations*. 103–207, London, 2011.

- [13] Pablo Cobreros. Paraconsistent vagueness: a positive argument. *Synthese*, 183(2):211–227, 2011.
- [14] Newton C. A da Costa. Sistemas formais inconsistentes (Inconsistent formal systems, in Portuguese). Habilitation thesis, Universidade Federal do Paraná, Curitiba, Brazil, 1963. Republished by Editora UFPR, Curitiba, Brazil, 1993.
- [15] Rodolfo Ertola, Francesc Esteva, Tommaso Flaminio, Lluís Godo, and Carles Noguera. Exploring paraconsistency in degree-preserving fuzzy logics. In Gabriella Pasi, Javier Montero, and Davide Ciucci, editors, *Proceedings of the 8th conference of the European Society for Fuzzy Logic and Technology EUSFLAT 2013*, pages 117–124, Milan, 2013. Atlantis Press.
- [16] Francesc Esteva and Lluís Godo. Monoidal t-norm based logic: Towards a logic for left-continuous t-norms. *Fuzzy Sets and Systems*, 124(3):271–288, 2001.
- [17] Francesc Esteva, Lluís Godo, Petr Hájek, and Mirko Navara. Residuated fuzzy logics with an involutive negation. *Archive for Mathematical Logic*, 39(2):103–124, 2000.
- [18] Francesc Esteva, Lluís Godo, and Franco Montagna. Equational characterization of subvarieties of BL-algebras generated by t-norm algebras. *Studia Logica*, 76(2):161–200, 2004.
- [19] Petr Hájek. *Metamathematics of Fuzzy Logic*, volume 4 of *Trends in Logic*. Kluwer, Dordrecht, 1998.
- [20] Petr Hájek. Observations on the monoidal t-norm logic. *Fuzzy Sets and Systems*, 132(1):107–112, 2002.
- [21] Petr Hájek, Lluís Godo, and Francesc Esteva. A complete many-valued logic with product conjunction. *Archive for Mathematical Logic*, 35(3):191–208, 1996.
- [22] Dominic Hyde and Mark Colyvan. Paraconsistent vagueness: Why not? *Australasian Journal of Logic*, 6:107–121, 2008.
- [23] Sándor Jenei and Franco Montagna. A proof of standard completeness for Esteva and Godo’s logic MTL. *Studia Logica*, 70(2):183–192, 2002.
- [24] Carles Noguera, Francesc Esteva, and Joan Gispert. On some varieties of MTL-algebras. *Logic Journal of the Interest Group of Pure and Applied Logic*, 13(4):443–466, 2005.

- [25] Charles S. Peirce. Vague. In James M. Baldwin, editor, *Dictionary of Philosophy and Psychology*, volume 2, page 748, New York, 1902. MacMillan.
- [26] Graham Priest, Richard Routley, and Jean Norman, editors. *Paraconsistent Logic: Essays on the Inconsistent*. Philosophia Verlag, München, 1989.
- [27] Helena Rasiowa. *An Algebraic Approach to Non-Classical Logics*. North-Holland, Amsterdam, 1974.
- [28] Lotfi A. Zadeh. Fuzzy sets. *Information and Control*, 8(3):338–353, 1965.