

Article

Long-chained 4-Aminoquinolines as Quorum Sensing Inhibitors in *Serratia marcescens* and *Pseudomonas aeruginosa*

Ivana Aleksi#, Sandra Šegan, Filip Andric, Mario Zlatovi#,
Ivana Moric, Dejan M. Opsenica, and Lidija Senerovic

ACS Chem. Biol., **Just Accepted Manuscript** • DOI: 10.1021/acscchembio.6b01149 • Publication Date (Web): 28 Mar 2017

Downloaded from <http://pubs.acs.org> on March 31, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1
2
3
4 **Long-chained 4-Aminoquinolines as Quorum Sensing Inhibitors in *Serratia marcescens***
5
6 **and *Pseudomonas aeruginosa***
7
8

9
10
11 Ivana Aleksić[†], Sandra Šegan[‡], Filip Andrić[‡], Mario Zlatović[§], Ivana Moric[†], Dejan M.
12 Opsenica^{*†}, Lidija Senerovic^{†*}
13
14

15 [†]Institute of Molecular Genetics and Genetic Engineering, University of Belgrade, Vojvode
16 Stepe 444a, P.O. Box 23, 11010 Belgrade, Serbia
17
18

19 [‡]Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoševa 12,
20 P.O. Box 473, 11000, Belgrade, Serbia
21
22

23 [§] Faculty of Chemistry, University of Belgrade, Studentski trg 12-16, P.O. Box 51, 11158,
24 Belgrade, Serbia
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Antibiotic resistance has become serious global threat to public health; therefore, the improved strategies and structurally novel antimicrobials are urgently needed to combat infectious diseases. Here we report a new type of highly potent 4-aminoquinoline derivatives as quorum sensing inhibitors in *Serratia marcescens* and *Pseudomonas aeruginosa*, exhibiting weak bactericidal activities (MIC > 400 μ M). Through detailed structure-activity study we have identified 7-Cl and 7-CF₃ substituted *N*-dodecylamino-4-aminoquinolines (**5** and **10**) as biofilm formation inhibitors with 50% biofilm inhibition at 69 μ M and 63 μ M in *S. marcescens* and *P. aeruginosa*, respectively. These two compounds, **5** and **10**, are the first quinoline derivatives with anti-biofilm formation activity reported in *S. marcescens*. QSAR analysis identified structural descriptors such as Wiener indices, HDPI, MTC, TCI, and $\log D(o/w)_{exp}$ as the most influential in biofilm inhibition in this bacterial species. Derivative **10** is one of the most potent quinoline type inhibitors of pyocyanin production described so far (IC₅₀=2.5 μ M). While we have demonstrated that **5** and **10** act as *Pseudomonas* quinolone system (PQS) antagonists, the mechanism of inhibition of *S. marcescens* biofilm formation with these compounds remains open since signaling similar to *P. aeruginosa* PQS system has not yet been described in *Serratia* and activity of these compounds on AHL signalling has not been detected. Our data show that 7-Cl and 7-CF₃ substituted *N*-dodecylamino-4-aminoquinolines present the promising scaffolds for developing anti-virulence and anti-biofilm formation agents against multidrug-resistant bacterial species.

INTRODUCTION

The rising problem of microbial resistance to current antibiotics and high spreading rate of resistant bacterial species has become the major public health concern. Resistance to most antibiotics has emerged only few years after their introduction into clinical practice; therefore, improved strategies and new antimicrobials are urgently needed to control infectious diseases. Many bacteria employ cell density-dependent communication system called quorum sensing (QS) to control their virulence factor production, motility, biofilm formation, bioluminescence, sporulation, and conjugation.¹ Quorum sensing system allows bacteria to monitor their cell density through the release of signaling molecules called autoinducers. At a high cell density, autoinducers reach threshold concentrations and initiate the signaling cascade that regulates expression of genes required for microbial pathogenicity.

Biofilms are complex communities of bacteria embedded in a self-produced matrix of polysaccharides, proteins, and extracellular DNA that strongly adhere to the surfaces of both organic or inorganic structures.² The biofilm-embedded cells are highly resistant to antimicrobial drug therapy, difficult to eradicate, and often cause serious life-threatening infections.³ Biofilms formed in the tissues and medical devices associated to human body (e.g., catheters, naso-laryngeal tubes, or stents) account for 70% of nosocomial infections.⁴ The colonization of a patient's tissue or the surfaces of indwelling medical devices with biofilm-forming bacteria usually leads to persistent infections which fail to resolve despite of aggressive antibiotic therapies.

Biofilms play an important role in virulence of many bacteria including Gram-negative pathogens *Serratia marcescens* and *Pseudomonas aeruginosa*.² *S. marcescens* as important nosocomial healthcare-associated pathogen has been recognised only in the last four decades.⁵ These bacteria are particularly involved in catheter-associated bacteremia, urinary tract infections, and wound infections. Infections caused by this opportunistic pathogen may be difficult to treat due to their resistance to multiple antimicrobial agents such

1
2
3
4 as β -lactams, aminoglycosides, and fluoroquinolones.⁶ On the other hand, *P. aeruginosa* is
5 well-known opportunistic pathogen that poses a significant health threat for
6 immunocompromised patients such as burn victims, cancer, cystic fibrosis (CF), or AIDS
7 patients.⁷ Therapeutics against *P. aeruginosa* are increasingly limited due to the continued
8 emergence and spreading of antibiotic resistant strains causing high mortality.⁸ At the same
9 time, multidrug-resistant strains of *S. marcescens* have been isolated from both clinical and
10 environmental settings⁹, suggesting that antibiotic usage should be restricted and the
11 alternative therapeutic approach should be implemented to treat these infections. Both *S.*
12 *marcescens* and *P. aeruginosa* control their virulence factor production including biofilm
13 formation using QS system. Therefore, attenuation of virulence and biofilm formation
14 through interference with QS could be a strategy of choice to treat these multidrug-resistant
15 bacterial infections. Disrupting cell-to-cell communication instead of killing the bacteria
16 would result in less selective pressure comparing to conventional antibacterial agents and
17 thus could circumvent the problem of resistance.¹⁰

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Numerous small organic molecules with QS inhibitory activity influencing bacterial
virulence in tissue culture and in animal models have been described.^{11, 12} Majority of these
compounds are non-natural synthetic or semi-synthetic derivatives of acylhomoserine lactone
(AHL). The most active biofilm inhibitors are synthetic or semi-synthetic antimicrobial
peptides¹³ but they show low stability and significant toxicity, thus making small organic QS
inhibitors more attractive therapeutic options against persistent bacterial infections.

Quinoline moiety is a well-known pharmacophore responsible for broad-spectrum of
biological activities¹⁴ including antimalarial and inhibitory activity against botulinum
neurotoxins.¹⁵ A number of quinolone derivatives with strong bactericidal activity have been
synthesized, affecting both planktonic and biofilm-associated bacteria.¹⁶⁻²⁰ Some of them
have demonstrated QS inhibitory activity such as antibiotic nitroxoline, which in micromolar
concentrations reduces formation of *P. aeruginosa* biofilm up to 80%.¹⁹ Recently, the new

1
2
3
4 quinolines have been described with strong anti-QS^{21,22} or anti-biofilm activities, including
5 halogen derivatives¹⁶ and nitroxaline amino derivatives¹⁷, quinoline β -amino alcohol¹⁸,
6 imines²³, phthalazine-quinoline²⁴ or 2-alkyl-4(3H)-quinazolinone derivatives²⁵.
7
8

9
10
11 Considering that various quinoline derivatives affect both Gram(-) and Gram(+) bacterial
12 species, together with their good anti-QS and anti-biofilm properties, in this study we have
13 examined antimicrobial activity of structurally less complex series of 4,7-disubstituted
14 aminoquinoline derivatives.
15
16
17
18

19 20 21 22 RESULTS AND DISCUSSION

23
24 4-Aminoquinoline derivatives (4-AQs) are clinically used antimalarial drugs. While their
25 antibacterial activity has been investigated at some extent^{26, 27}, to the best of our knowledge
26 their anti-virulence activity has not yet been reported. Numerous small inhibitors of *P.*
27 *aeruginosa* virulence factor production have been synthesized but reports on synthetic biofilm
28 inhibitors of *S. marcescens* remain scarce.²⁸ Here we address antibacterial and anti-virulence
29 potentials of series of twenty two 4,7-disubstituted AQs (Figure 1) against these two common
30 nosocomial opportunistic pathogens. Two derivatives (**9** and **10**) were synthesized for the first
31 time in this study (Figure 1). Synthesis procedures with corresponding spectral data and
32 copies of NMR spectra are given in Supplemental Information 1 (SI1, Scheme 1S).
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Chemical structures of tested compounds.

Inhibition of quorum sensing in *Serratia marcescens* by 7 Cl-aminquinolines.

Firstly we have tested antibacterial activity of N-substituted derivatives of 4-amino-7-chloroquinoline containing aliphatic aminoalkyl normal chain with different lengths, i.e., C2, C4, C6, C8, and C12 (**1 – 5**, Figure 1). All tested compounds showed low antibacterial

activities against *S. marcescens* with minimal inhibitory concentrations (MICs) ranging from 565 μM for **1** to 1800 μM for **3** (Table 1). Lengthening the aliphatic chain to C8 or C12 (**4** and **5**, respectively) slightly increased antibacterial effects of the compounds and observed order of bactericidal activity was **1** > **2** > **3** < **4** < **5** (Table 1). Obtained MIC values were similar to MIC values reported for 1-methyl-3-sulfonylthio-4-aminoquinolinium salts against *S. marcescens*²⁶, but were at least one order of magnitude higher than MIC values reported for C(2) substituted bromo-quinolines against *S. aureus* and *S. epidermidis*^{16,17} or quinoline amino alcohols against *Vibrio cholera*.¹⁸

Table 1. Minimal inhibitory concentration of tested compounds against *Serratia marcescens* and *Pseudomonas aeruginosa*.

Compound	MIC ^a (μM)	
	<i>S. marcescens</i>	<i>P. aeruginosa</i>
1	565	565
2	1001	1000
3	1800	1800
4	1635	1635
5	690	1380
6	1335	1335
7	2320	2320
8	980	1960
9	185	1475
10	315	1265

1			
2			
3			
4			
5	11	970	970
6			
7	12	700	1400
8			
9			
10	13	3870	3870
11			
12			
13	14	3055	3055
14			
15	15	765	3055
16			
17			
18	16	2525	2525
19			
20			
21	17	2160	2160
22			
23			
24	18	1390	2785
25			
26	19	430	860
27			
28			
29	20	1750	1750
30			
31			
32	21	2245	2245
33			
34	22	1130	1130
35			
36			
37	23	8320	8320
38			
39			
40	24	5670	5670
41			
42			
43	25	4300	4300
44			
45	26	3465	3465
46			
47			
48	27	2495	2495
49			

^a MIC – concentration of tested compound that cause 100 % inhibition of bacterial growth.

Since bactericidal effects were negligible, we have next examined anti-quorum sensing (anti-QS) activity of these compounds. *S. marcescens* has *N*-acylhomoserine lactones

1
2
3
4 (AHL)- dependent QS system and produces at least four AHLs (3-oxo-C6-HSL, C6-HSL,
5 C7-HSL, and C8-HSL), which regulate production of prodigiosin, carbapenem antibiotic
6 resistance, extracellular and cell-associated enzymes, and virulence factors such as motility
7 and biofilm formation.²⁹ The interference of *N*-alkylamino-7-Cl AQ-derivatives with QS in *S.*
8 *marcescens* was examined by measuring inhibition of prodigiosin production in a disk assay
9 and inhibitory effect of tested compounds was detected as appearance of colourless halo
10 around a disk. All compounds except **5** caused inhibition of prodigiosin synthesis. The zones
11 of prodigiosin inhibition were larger in the presence of derivatives with shorter aliphatic
12 chains ranging from 14 to 22 mm in following order **1**>**2**=**3**=**4** (Table 2, Table 1S). All the
13 compounds (except **7**) also inhibited violacein production in *Chromobacterium violaceum*
14 CV026 disk assay, suggesting that 4-AQ derivatives can interfere with AHL signalling (Table
15 1S).
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 Therefore, we have then tested whether 7-Cl AQ derivatives show anti-biofilm
32 formation activity, since biofilm formation is the major virulence factor regulated by QS in *S.*
33 *marcescens*.³⁰ Surprisingly, only derivatives without prodigiosin inhibitory activity, **5** and **10**,
34 exhibited anti-biofilm formation activity with BFIC₅₀* = 69 μM and BFIC₅₀ = 63 μM
35 respectively (i.e., 50% inhibition at 25 μg/mL, Table 2). More importantly, **5** showed
36 significant difference between anti-biofilm formation and antibacterial activities exhibiting
37 10 fold lower BFIC₅₀ than MIC value. Together, these results indicated that *N*-alkylamino-7-
38 Cl AQ-derivatives could target different processes and thus exhibit different activities against
39 *S. marcescens*.
40
41
42
43
44
45
46
47
48
49
50

51 **SAR analysis of quinoline derivatives as biofilm formation inhibitors *Serratia*** 52 ***marcescens*** 53 54

55 To investigate an effect of structural changes to observed bioactivities, we examined
56 influence of substituents at C(7) and C(4) on bactericidal, anti-QS, and anti-biofilm formation
57
58
59

60 * BFIC₅₀ - concentration of compound that inhibited biofilm formation by 50%

1
2
3
4 activities. Replacing the chlorine atom at C(7) with hydrogen resulted in the reduction of
5
6 bactericidal activities as demonstrated by twofold decrease in activities of derivatives **6** (MIC
7
8 = 1335 μ M) and **7** (MIC = 2320 μ M) in comparison to their 7-chloro analogues **1** and **2**,
9
10 respectively (Table 1). These weak bactericidal activities fulfilled the requirements of ideal
11
12 anti-QS agents exhibiting low selective pressure on bacteria and thus, reducing possibility to
13
14 develop resistance. However, these des-chloro derivatives demonstrated low effect on
15
16 prodigiosin production and stimulated biofilm formation (Tables 2). Introduction of strong
17
18 electron withdrawing CF₃-group at C(7) position was followed by the increase of bactericidal
19
20 activity of **8** (MIC = 980 μ M) in comparison to des-chloro derivative **6**, but still **8** was two
21
22 times less active in comparison to **1**. However, while biofilm formation was stimulated in the
23
24 presence of small concentrations of **1** (147 % and 135 % at 10 and 25 μ g/ml, respectively;
25
26 Table 2) biofilms remained at the same level (around 95%) in the presence of all three tested
27
28 concentrations of **8**. Introduction of C8 and C12 chains to 7-CF₃ derivatives caused
29
30 significant increase in bactericidal activity of the compounds against *S. marcescens* (Table 1).
31
32 Derivatives **9** and **10**, with MIC values of 184 μ M and 316 μ M were 5 and 3 time,
33
34 respectively, more active than **8**. Furthermore, we have shown that 7-CF₃ derivatives with
35
36 long aminoalkyl chain were significantly more active in comparison to 7-Cl derivatives.
37
38 These results suggest that C(7) substituents have a strong influence on bactericidal activity of
39
40 long chain C(4)-amino(alkylamino) substituted quinoline derivatives, with **9** being the most
41
42 active derivative within the group. Inhibition of prodigiosin production was detected only in
43
44 the presence of **9** (Table 2, Table 1S), while **10** was the sole compound that inhibited biofilm
45
46 formation in *S. marcescens* (Table 2), with BFIC₅₀ value (69 μ M) which was five times lower
47
48 than its MIC value. Importantly, the inhibition of biofilm formation with derivatives **5** and **10**
49
50 occurs without effect on bacterial viability as confirmed by fluorescence and scanning
51
52 electron microscopy (Figure 1S).
53
54
55
56
57
58
59
60

Only 7-Cl and 7-CF₃ derivatives substituted with C12 alkyl chain showed effective inhibition of biofilm formation in *S. marcescens*. Derivatives with shorter diaminoalkyl chains either showed no effect or stimulated biofilm formation. These results have indicated that the length of aminoalkyl chain rather than the type of the substituent at C(7) is the key parameter for efficient anti-biofilm formation activity of 4-AQ derivate against *S. marcescens*.

Table 2. Inhibition of prodigiosin synthesis and biofilm formation in *Serratia marcescens* in the presence of quinoline derivatives and experimentally determined logD values.

Compound	Prodigiosin inhibition ^a	Biofilm formation (%)			logD(o/w) _{exp}
		Concentration of applied compounds			
		10 [μg/mL]	25 [μg/mL]	50 [μg/mL]	
1	20±2	147±10	135±8	95±10	1.03
2	14±2	109±7	107±10	140±12	1.22
3	14±1	137±12	137±12	147±10	1.43
4	14±1	98±10	98±10	99±10	1.81
5	n.a.	88±15	48±5	39±5	2.81
6	10±1	138±12	125±10	134±15	0.45
7	n.a.	128±10	117±8	169±15	0.71
8	n.a.	93±10	96±2	90±10	1.17
9	12±1	103±15	70±5	80±10	1.91
10	n.a.	110±6	43±2	40±4	2.94
11	16±2	118±10	130±15	144±15	1.73

1						
2						
3						
4						
5	12	n.a.	124±15	110±15	110±8	1.17
6						
7	13	22±2	155±15	165±8	116±6	1.62
8						
9						
10	14	n.a.	100±12	100±15	126±7	1.51
11						
12						
13	15	n.a.	129±15	136±12	158±20	1.43
14						
15	16	n.a.	122±12	101±5	170±15	1.91
16						
17						
18	17	n.a.	138±8	133±7	110±10	2.57
19						
20						
21	18	n.a.	88±10	95±8	99±10	1.77
22						
23						
24	19	28±3	125±10	119±10	96±8	1.13
25						
26	20	28±2	139±25	159±10	136±15	1.81
27						
28						
29	21	n.a.	150±10	170±20	132±15	1.43
30						
31						
32	22	8±1	134±10	140±15	139±12	1.70
33						
34	23	n.a.	98±12	104±10	108±15	n.d.
35						
36						
37	24	n.a.	89±12	85±10	94±15	n.d.
38						
39						
40	25	n.a.	104±6	119±10	139±10	n.d.
41						
42						
43	26	n.a.	94±10	85±12	112±15	n.d.
44						
45	27	14±1	105±10	124±12	141±10	n.d.
46						
47						

^a Zones of inhibition (mm). Inhibition of pigment production was determined in the presence of 250 µg of tested compound per disk; n.a. – not active; n.d. – not determined.

The importance of the presence of substituent at C(4) and its character were explored using a range of quinoline derivatives (Tables 1 and 2). Substantial changes in the structure, from derivatives without substituent, i.e., with hydrogen at C(4), like **13** and **15**, through 4-

1
2
3
4 chloroquinolines **14**, **16**, **17**, 4-hydroxyquinoline **18**, 4-amino-7-chloroquinoline **12** and 4-
5 amino-*N*-(substituted) derivatives **11**, **20**, **21** and **22** did not significantly affect either
6 bactericidal activity or prodigiosin production and biofilm formation in *S. marcescens*. The
7 strongest bactericidal activity exhibited derivative **19** with MIC = 430 μ M, already known as
8 good Fe(III), Zn(II), and Ca(II) N,O-chelating ligand³¹⁻³³.
9

10
11 Diaminoalkanes **23**, **24**, **25**, **26**, and **27** did not perform any antibacterial (MIC > 2500
12 μ M), anti-QS or anti-biofilm formation activity (Tables 1 and 2), thus demonstrating that
13 quinoline core has been indispensable component in establishing QS and biofilm formation
14 inhibitory activities within tested group of compounds.
15
16
17
18
19
20
21
22
23
24
25
26

27 **QSAR modeling of biofilm formation in *Serratia marcescens* with 4-aminoquinolines**

28
29 Chromatographic parameters obtained under reversed-phase (RP) conditions can be
30 correlated with biological activity of compounds, such as transport through cell membranes,
31 binding to plasma proteins or drug-target interactions.^{34, 35} Most of the investigated 4-AQ
32 derivatives exhibit strong retention on octadecyl (C18) stationary phase, using mobile phase
33 pH \geq 5. On the contrary, with mobile phase of pH \leq 1, their retentions were much weaker,
34 and typical RP mechanism of retention, based on hydrophobic interactions with nonpolar
35 stationary phase, was established. The partition coefficients of examined quinoline
36 derivatives, denoted as $\log D(o/w)_{exp}$ (Table 2), were determined using reversed-phase thin-
37 layer (RP-TLC) method as described in SI1.
38
39
40
41
42
43
44
45
46
47
48

49 In order to quantitatively correlate structural features of 4-AQ derivatives with their
50 potential to affect biofilm formation in *S. marcescens*, QSAR models were built using Partial
51 Least Square (PLS) regression.³⁶ The entire set of 180 molecular descriptors (SI2) and
52 $\log D(o/w)_{exp}$ values were used as independent variables, while logarithms of biofilm
53 formation values measured at different concentration (10, 25, and 50 μ g/mL), denoted as
54 $\log BI_{10}$, $\log BI_{25}$ and $\log BI_{50}$, respectively, were used as dependent ones. The number of
55
56
57
58
59
60

variables in final models was reduced to 52 – 59 by a double fold PLS regression, hence significantly increasing predictive performance. Predictive ability and optimal model complexity (number of PLS components) were estimated through the double cross-validation procedure proposed by Warmuza and Filzmoser.³⁷ Detailed computation and modeling procedures are described in the SI1 and SI2. In the case of $\log BI_{50}$ and $\log BI_{25}$ models of satisfactory predictive ability were achieved with R^2 values not falling below 0.645 for prediction, and 0.817 for calibration purposes, and errors not exceeding 0.1 log units (Table 3). On the other hand, in the case of $\log BI_{10}$, a narrow range of biofilm formation percentages (0.25 log units), was insufficient to build a reliable QSAR ($R^2_{\text{Cal}} = 0.649$, $R^2_{\text{CV}} = 0.304$, $R^2_{\text{Pred}} = 0.371$).

Table 3. Statistical performance of QSAR models connecting the most contributing molecular descriptors with inhibition of the biofilm formation in *Serratia marcescens*.

Dependent variable	Statistical performance parameters	The most contributing variables (VIP scores > 1)
$\log BI_{50}$	$RMSE_{\text{Cal}} = 0.037$, $RMSE_{\text{CV}} = 0.080$, $RMSE_{\text{Pred}} = 0.100$ $R^2_{\text{Cal}} = 0.950$, $R^2_{\text{CV}} = 0.784$, $R^2_{\text{Pred}} = 0.645$	QPllogBB, W, SMT, MDDD, GMT, GMTV, AVDD, UP, CENT, VAR,
	$n(\text{PLS components}) = 4$	ECCc, ECC, QW, FM,
	Percent of variance captured by each latent variable in X and Y domain respectively	TCI8-10, MTCI9,10, HDPI, Whet(Z, e, m, v, and p), logD
	PLS1: 56.10% and 62.20%	
	PLS2: 11.71% and 19.34%	
$\log BI_{25}$	$RMSE_{\text{Cal}} = 0.064$, $RMSE_{\text{CV}} = 0.081$, $RMSE_{\text{Pred}} = 0.084$ $R^2_{\text{Cal}} = 0.817$, $R^2_{\text{CV}} = 0.720$, $R^2_{\text{Pred}} = 0.690$	IP, Dipole_PC_X, Dipole_S_X, W, MW, MSDB, SMT, SMTV,
	$n(\text{PLS components}) = 2$	MDDD, GMT, GMTV, AVDD, UP, CENT, VAR,
	Percent of variance captured by each latent variable in X and Y domain respectively	ECCc, ECC, AECC,
	PLS3: 6.60% and 11.67%	
	PLS4: 6.12% and 1.76%	

PLS1: 63.47% and 67.53%

PLS2: 14.26% and 14.13%

DECC, AvX5, QW, FM,
SM, HDPI, Whet(Z, e, m,
v, and p), ACIX2, KAMS3,
RSIpw5

Obtained QSAR models showed that branching and the size of the molecules are the key topological descriptors responsible for modulation of biofilm formation. The most contributing variables ($VIP > 1$), for $\log BI_{50}$ and $\log BI_{25}$ were: Wiener topological index (W), MW, WhetZ, Whete, Whetm, Whetv, and Whetp, GMT and GMTV, MTCI9 and MTCI 10, ECCECCc, HDPI, UP, molecular centrality (CENT), etc (for abbreviation and complete list see in SI2). As common contributing variables for $\log BI_{50}$ and $\log BI_{25}$ are the Wiener indices associated to molecular branching and size. They are reciprocally related to branching degree, i.e., molecules with smaller values have more branched substituents.³⁸ All of the Wiener indices have negative regression coefficients (Figure 2), which means that introduction of long unbranched hydrocarbon chains to 4-AQs core results in an increase of anti-QS activity. Derivatives **5** and **10** have significantly higher values of Wiener indices (SI2), corresponding to other examined 4-AQs, which coincide with the highest biofilm formation inhibition, even when they are compared to structurally closest derivatives **4** and **9**. Similarly, HDPI show negative contribution to $\log BI_{50}$ and $\log BI_{25}$. Again, most active derivatives **5** and **10** have the highest values of HDPI. The highest positive contribution on $\log BI_{50}$ have MTCI10 index. The lowest values of MTCI10 (SI2) have most potent biofilm inhibitors **5** and **10**, with **4** and **9** as closest followers. Lipophilicity ($\log D(o/w)_{exp}$) may also be a significant contributor to anti-QS and biofilm formation inhibitory activity of 4-AQ derivatives. Although less important than the most of topological descriptors, $\log D(o/w)_{exp}$ negatively affects $\log BI_{50}$ (Figure 2a), which means that the increase of the lipophilicity of 4-AQs leads to increased anti-biofilm formation potency. As it was expected, linear dependence between alkyl-chain length and $\log D(o/w)_{exp}$ was found. In addition, compounds

1
2
3
4 with strong electron withdrawing CF₃-group at C(7) are more lipophilic than corresponding
5
6 7-chloro derivatives, and following order of lipophilicity 7-CF₃ > 7-Cl > 7-H was observed.
7
8
9 Derivative **10** have the highest lipophilicity value (Table 2), while **6** have the lowest. Indices
10
11 TCIs (order **8**, **9** and **10**; Figure 2a) and MTCIs (order **9** and **10**; Figure 2a) have positive
12
13 impact on log*BI*₅₀. Those indices describe the ability of a charge transfer throughout the
14
15 molecule. Derivatives with 7-CF₃ group have 1.5 – 2 times higher values of TCIs indices in
16
17 comparison to corresponding 7-Cl analogues, and obtained order 7-CF₃ > 7-Cl > 7-H clearly
18
19 indicates stronger influence of substituent in C(7) position than length of alkyl chain on TCIs
20
21 values. That is additionally confirmed with very low values of TCIs for diaminoalkanes. On
22
23 the other hand, MTCIs values are more influenced by the length of alkyl chain than by the
24
25 nature of C(7) substituents, i.e., derivatives with longer diaminoalkyl chain have lower
26
27 MTCIs values. However, correlation between TCIs and either anti-QS or anti-biofilm
28
29 formation values have not been observed. Descriptors defining specific interactions such as
30
31 acceptHB, NO, FISA and PSA, QPlogS and QPlogBB, have significantly lower impact on
32
33 log*BI*. This clearly implies that shape and size of molecule, especially introduction of long
34
35 unbranching substituents, are the most important factors in establishing ligand-receptor
36
37 interactions that affects QS signaling pathways.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 or significantly higher than previously reported 1- methyl-3-sulfonylthio-4-aminoquinolinium
5 salts.²⁶
6
7

8
9 Two compounds with anti-biofilm formation activity in *S. marcescens* derivatives **5**
10 and **10** were examined for their anti-QS activity in *P. aeruginosa*. Derivative **4**, which
11 demonstrated weak anti-QS activity and had no effect on biofilm formation in *S. marcescens*
12 was also tested for the comparison. QS inhibition in *P. aeruginosa* was followed in biofilm
13 formation and pyocyanin production assays in the presence of these derivatives. Derivative **4**
14 inhibited *P. aeruginosa* biofilm formation by approximately 60 %, even at 10 µg/mL (33 µM,
15 Table 4). Derivatives **5** and **10** also inhibited biofilm formation, with decrease ranging from
16 20 – 40 % and 30 – 50 %, respectively (Figure 1S c and d), depending on applied dose and
17 C(7) substituent. On the other hand, while **5** and **10** were able to reduce pyocyanin production
18 with respective IC₅₀ of 140 and 2.5 µM, in comparison to DMSO treated controls, **4** induced
19 pyocyanin overproduction (Table 4 and Figure 2S). Pyocyanin inhibitory activity of
20 derivative **10** was similar to recently reported activity of 3-carboxamido-2-heptyl-4-
21 quinolinon antagonist (IC₅₀ = 2 µM)³⁹, but twenty times stronger than 2-alkyl-4(3H)-
22 quinazolinone derivatives²⁵. However, derivative **10** was ten fold less potent in pyocyanin
23 inhibition than benzamide-benzimidazole derivatives which is one of the most potent
24 pyocyanin inhibitors described so far⁴⁰. Derivative **17**, which was also used as a control
25 compound, exhibited no QS inhibitory activity. Similar results showing the influence of alkyl
26 chain length on *P. aeruginosa* QS system with agonistic or antagonistic activities were
27 reported for some 4-quinolon analogues^{21, 41}.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 The QS network of *P. aeruginosa* is organized in a multi-layered hierarchy consisting
52 of at least four interconnected signaling pathways: Las, Rhl, the PqsR-controlled quinolone
53 system (PQS), and Integrated QS system (IQS). Three autoinducer synthases LasI, RhlI, and
54 PqsABCDH produce autoinducers 3-oxo-C12-homoserine lactone (HSL), C4-HSL, and
55 2-heptyl-3-hydroxy-4-quinolone (PQS), respectively, which regulate formation of biofilms
56
57
58
59
60

and production of virulence factors.⁴² Inhibition of specific QS pathway in *P. aeruginosa* PAO1 by **4**, **5**, and **10** was quantified using three biosensors: *P. aeruginosa* PA14-R3, used to measure 3OC12-HSL production (LasI activity), *P. aeruginosa* PAOJP2/pKD-*rhlA*, used for measurements of C4-HSL (RhlI activity), and *P. aeruginosa* PAO1Δ*pqsA* (CTX *lux::pqsA*), used for evaluation of PQS production (PqsABCDH activity). Compound **17** with no effect on pyocyanin production and biofilm formation was used as negative control.

Table 4. Effects of selected quinoline derivatives on biofilm formation and inhibition of pyocyanin production in *Pseudomonas aeruginosa*.

Compound	Biofilm formation (%)			Pyocyanin production ^a (%)
	Concentration of compounds			
	10 [μg/mL]	25 [μg/mL]	50 [μg/mL]	
4	44±2	48±5	38±2	155±10
5	79±2	58±7	60±6	50±3
10	67±3	46±5	52±5	10±1
17	83±8	82±10	98±10	103±1

^a Production of pyocyanin was measured in the presence of a compound at concentration of 50 μg/mL.

Derivative **4** reduced Las signalling by 45 %, while C12-amino derivatives, **5** and **10**, induced overproduction of C12-AHL by 316 % and 118 %, respectively (Table 5). Threefold stronger effect of **5** over **10** on LasI activity is due to different C(7)-substituent. Derivatives **5** and **10** did not have any effect on prodigiosin production in *S. marcescens* (Table 2), which is regulated by short-chain AHLs⁴³, suggesting the absence of interference with C4-AHL signalling. The assumption was confirmed by their weak effect on RhlI activity, manifested

as 10 % and 20 % inhibition, respectively. Derivative **4**, which showed small zone of prodigiosin inhibition (Table 1), reduced RhlI activity by 23 %.

Table 5. Influence of tested derivatives on *Pseudomonas aeruginosa* QS pathways.

Production of autoinducers in <i>P. aeruginosa</i> PAO1 (%)			
Compound ^a	3oxoC12-HSL	C4-HSL	PQS
4	55±6	77±5	91±8
5	316±16	91±15	16±2
10	118±10	80±12	26±4
17	148±12	86±9	69±3

^a Bacteria were incubated with tested compounds at 50 µg/mL; values are given relative to control samples where bacteria were grown with DMSO and are average of three independent experiments ± SD.

Significant difference between tested derivatives were observed when comparing their activities against *P. aeruginosa* PQS. While **4** showed a minor effect on quinolone signalling with only 9 % inhibition, both **5** and **10** inhibited PQS by 85 % and 75 %, respectively. Derivative **17** showed 1.5-fold stimulatory effect on LasI activity and caused 30 % reduction of PQS signalling.

Taken together, these results suggest that 4-AQs with *N*-dodecylamino substituent inhibit biofilm formation and virulence factor production in *P. aeruginosa* through inhibition of PQS signalling. These data are consistent with previous reports showing that PQS signalling regulates the production of diverse virulence factors including pyocyanin, in

1
2
3
4 addition to affecting biofilm formation.⁴⁴ Although **4** exhibited stronger anti-biofilm
5 formation activity in *P. aeruginosa* than **5** and **10**, it also had stimulatory effect on pyocyanin
6 production (Table 4). Considering detrimental effects that pyocyanin can cause in infected
7 host tissues through interference with cellular functions such as electron transport, cellular
8 respiration, energy metabolism, gene expression, and innate immune mechanisms⁴⁵ our
9 results show that C12 substituted derivatives have more preferred anti-virulence
10 characteristics comparing to compound **4**.
11
12
13
14
15
16
17
18
19
20
21

22 CONCLUSION

23
24 In this study we have examined 22 quinoline derivatives as potential antimicrobial agents
25 against two pathogens, *S. marcescens* and *P. aeruginosa*, and identified new quinoline
26 derivatives with strong anti-QS activity inhibiting biofilm formation and without bactericidal
27 effect. We have demonstrated that the efficient QS inhibition by these compounds depends
28 on the presence of C(7)-substituent, basic amino group, and the long-chain *N*-alkylamino
29 substituent at C(4) of quinoline core, with C12 chain derivatives being the most active against
30 biofilm formation. Consistently with previous report from Klein and coworkers⁴⁶, differences
31 observed for C8 and C12 AQs related to inhibition of bacterial pigments production and
32 biofilms formation in both examined bacteria clearly showed that relatively small structural
33 changes cause significant differences in QS modulating activity. Derivative **10** reduced *P.*
34 *aeruginosa* pyocyanin production with a potency belonging to the group of the most active
35 quinoline type inhibitors described so far. Notably, the compounds **5** and **10** are the first
36 quinoline derivatives with demonstrated anti-biofilm formation activity in *S. marcescens*. The
37 important question opened in this study but not yet clarified, is the mechanism of *S.*
38 *marcescens* biofilm formation inhibition with C12 4-AQs since signalling similar to *P.*
39 *aeruginosa* PQS system has not been described in *S. marcescens*, and activity of these
40 compounds on AHL signalling was not detected.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS

Compounds Synthesis. Compounds **11**, **13**, **14**, **15**, **16**, **17**, **19**, **23**, **24**, **25**, **26** and **27** were obtained from Sigma-Aldrich Co (Sigma Aldrich, Germany) and used without further purification. Compound **18** was obtained according to procedure described by Terzic and coworkers⁴⁷ and all spectra were identical. Compounds **9** and **10** were synthesized for the first time, while other tested compounds were synthesized according to previously published procedures (SI 1).

IR spectra were recorded on a Perkin-Elmer spectrophotometer FT-IR 1725X. ¹H and ¹³C NMR spectra were recorded on a Varian Gemini-200 spectrometer (at 200 and 50 MHz, respectively), and on a Bruker Ultrashield Advance III spectrometer (at 500 and 125 MHz, respectively) employing indicated solvents (*vide infra*) using TMS as the internal standard. Chemical shifts are expressed in ppm (δ) values and coupling constants (J) in Hz. ESI-MS spectra were recorded on Agilent Technologies 6210 Time-Of-Flight LC-MS instrument in positive ion mode with CH₃CN/H₂O 1/1 with 0.2 % HCOOH as the carrying solvent solution. Samples were dissolved in CH₃CN or MeOH (HPLC grade purity). The selected values were as follows: capillary voltage = 4 kV, gas temperature = 350 °C, drying gas = 12.1 min⁻¹, nebulizer pressure = 45 psig, fragmentator voltage = 70 V. The elemental analysis was performed on the Vario EL III- C,H,N,S/O Elemental Analyzer (Elementar Analysensysteme GmbH, Hanau-Germany). Thin-layer chromatography (TLC) was performed on precoated Merck silica gel 60 F254 and RP-18 F254 plates.

Microbial Strains and Growth Conditions. *Pseudomonas aeruginosa* PAO1 NCTC 10332 and *Serratia marcescens* ATCC 27117 were used in this study. Bacteria were grown in Luria Bertani (LB) broth on a rotary shaker at 180 rpm.

Antimicrobial Susceptibility Tests for Planktonic Cells. The minimum inhibitory concentrations of 4-AQ derivatives were determined according to standard broth microdilution assays recommended by the Clinical and Laboratory Standards Institute (M07-

1
2
3
4 A9). Stock solutions of 4-AQ derivatives were prepared in DMSO (50 g/L, w/v). The highest
5 tested concentration of any compound was 500 mg/L. The inoculums were 10^5 colony
6 forming units (CFU)/mL. The MIC value corresponds to the lowest concentration that
7 inhibited the growth after 20 h at 30 °C for *S. marcescens* or 37 °C for *P. aeruginosa*.
8
9

10
11 **Antimicrobial Susceptibility Tests for Biofilms.** Biofilm quantification assays were
12 performed in 96-well microtiter plates using a crystal violet (CV) method to stain adherent
13 cells⁴⁸. Biofilms formed for 24 h in the presence or absence of compound at 30 °C for *S.*
14 *marcescens* or 37 °C for *P. aeruginosa* were washed and adherent cells stained with 0.1%
15 (v/v) CV. Each biofilm formation assay was performed in six wells and repeated three times.
16
17

18
19 ***Serratia marcescens* Disk Assay.** Overnight culture of *S. marcescens* was diluted 100-fold in
20 molten semi-solid LB agar (0.3 % w/v) and poured over solid LB medium. Cellulose disks
21 containing compounds (250 µg/disk) were placed on solidified agar and incubated for 24 h at
22 30 °C. Inhibition of prodigiosin synthesis was identified by the absence of red colour around
23 the disk.
24
25

26
27 **Pyocyanin Assay** was performed with *P. aeruginosa* PA14 indicator strain as reported
28 previously¹². Pyocyanin in the supernatant was quantified using UV–vis spectrophotometer
29 Ultrospec 3300pro (Amersham Biosciences, USA) at 695 nm. All experiments were
30 performed in triplicate and repeated at least three times.
31
32

33
34 **AHL Production Assays.** Production of 3OC12-HSL and C4-HSL were determined in
35 supernatants of *P. aeruginosa* PAO1 culture grown for 6 h in the presence of selected
36 compounds or DMSO as previously reported.⁴⁹ Aliquots of *P. aeruginosa* PAO1 supernatants
37 were added to *P. aeruginosa* PA14-R3 ($\Delta lasI PrsA::lux$)⁵⁰ or PAOJP2/pKD-*rhlA* ($\Delta rhlA$
38 *PrhIA::lux*)⁵¹ biosensors' cultures and cell density (A_{600}) and bioluminescence (light counts
39 per second, LCPS) were simultaneously measured after 4 h of incubation using Tecan
40 Infinite200 multiplate-reader (Tecan Group Ltd., Switzerland). Luminescence values were
41 normalized per cell density.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **PQS Measurements.** PQS measurements were performed according to Fletcher et al⁵² with
5 some modifications. *P. aeruginosa* PAO1 stationary phase cultures (10 ml) grown with
6 selected compounds or DMSO were extracted with the same volume of acidified ethyl
7 acetate. After vortexing and centrifugation, organic phase was transferred to a fresh tube and
8 dried out under a stream of nitrogen gas. The residue was resuspended in 50 μ l of methanol
9 for subsequent PQS measurements. Overnight cultures of *P. aeruginosa* PAO1 $\Delta pqsA$ (CTX
10 *lux::pqsA*)⁵² were diluted 1:1000 in fresh LB medium, and 0.2 mL of cultures were grown in
11 microtiter plates in the presence of 5 μ l of extracts. Cell density and bioluminescence were
12 measured as described above. All assays were carried out in triplicate at least two times.
13
14
15
16
17
18
19
20
21
22
23

24 **Molecular and QSAR.** All structures were built using the Maestro 10.1 from Schrödinger
25 Suite 2015-1 (Maestro, version 10.1, Schrödinger, LLC, New York, 2015). QikProp version
26 4.3 (QikProp, version 4.3, Schrödinger, LLC, New York, 2015) was used for the calculation
27 of physically significant molecular descriptors and pharmaceutically relevant properties.
28 Single point calculations using the RM1 method⁵³ from Semiempirical NDDO module of
29 Schrödinger Suite 2015-1 was used for semi-empirical parameters. QSAR models were built
30 by Partial Least Square regression using PLS_Toolbox software package (v. 5.7 Eigenvectors
31 Inc.) for MATLAB (v. 7.8.0 R2009; MathWorks, Natick, USA). Detailed procedure is
32 described in the SI1.
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **ASSOCIATED CONTENT**

47
48 The Supporting Information is available free of charge on the ACS Publications website at
49 DOI:
50

51
52 Supporting Tables S1 and S2, Figures S1 and S2, Scheme S1, experimental and synthetic
53 procedures, compounds characterizations, QSAR computations, NMR spectra (SI1, PDF);
54
55 Molecular descriptors and Details of PLS model (SI2, XLSX).
56
57
58
59
60

AUTHOR INFORMATION

Corresponding Authors:

*E-mail: dopsen@chem.bg.ac.rs

*E-mail: lidijasenerovic@imgge.bg.ac.rs

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

This study has been funded by a Research Grant 2015 by the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) to L. Senerovic and by the Ministry of Education, Science and Technological Development, Republic of Serbia (Grants No. 173048 and No.172008). We gratefully acknowledge L. Leoni, Department of Biology, University Roma Tre, Italy for providing *P. aeruginosa* biosensor strains.

REFERENCES

1. Camilli, A. and Bassler, B. L. (2006) Bacterial small-molecule signaling pathways. *Science*. *311*, 1113-1116.
2. Hall-Stoodley, L., Costerton, J. W. and Stoodley, P. (2004) Bacterial biofilms: from the natural environment to infectious diseases. *Nat Rev Microbiol*. *2*, 95-108.
3. Oppenheimer-Shaanan, Y., Steinberg, N. and Kolodkin-Gal, I. (2013) Small molecules are natural triggers for the disassembly of biofilms. *Trends Microbiol*. *21*, 594-601.
4. Bryers, J. D. (2008) Medical biofilms. *Biotechnol Bioeng*. *100*, 1-18.
5. Merkier, A. K., Rodriguez, M. C., Togneri, A., Brengi, S., Osuna, C., Pichel, M., Cassini, M. H., *Serratia marcescens* Argentinean Collaborative Group, and Centron, D. (2013) Outbreak of a cluster with epidemic behavior due to *Serratia marcescens* after colistin administration in a hospital setting. *J Clin Microbiol*. *51*, 2295-2302.
6. Van Houdt, R., Givskov, M. and Michiels, C. W. (2007) Quorum sensing in *Serratia*. *FEMS Microbiol Rev*. *31*, 407-424.
7. Lyczak, J. B., Cannon, C. L. and Pier, G. B. (2000) Establishment of *Pseudomonas aeruginosa* infection: lessons from a versatile opportunist. *Microbes Infect*. *2*, 1051-1060.
8. Gellatly, S. L. and Hancock, R. E. (2013) *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathog Dis*. *67*, 159-173.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
9. Yang, H. F., Cheng, J., Hu, L. F., Ye, Y. and Li, J. B. (2012) Identification of a *Serratia marcescens* clinical isolate with multiple quinolone resistance mechanisms from China. *Antimicrob Agents Chemother.* *56*, 5426-5427.
 10. Hentzer, M. and Givskov, M. (2003) Pharmacological inhibition of quorum sensing for the treatment of chronic bacterial infections. *J Clin Invest.* *112*, 1300-1307.
 11. Stacy, D. M., Welsh, M. A., Rather, P. N. and Blackwell, H. E. (2012) Attenuation of quorum sensing in the pathogen *Acinetobacter baumannii* using non-native N-Acyl homoserine lactones. *ACS Chem Biol.* *7*, 1719-1728.
 12. O'Loughlin, C. T., Miller, L. C., Siryaporn, A., Drescher, K., Semmelhack, M. F. and Bassler, B. L. (2013) A quorum-sensing inhibitor blocks *Pseudomonas aeruginosa* virulence and biofilm formation. *Proc Natl Acad Sci U S A.* *110*, 17981-17986.
 13. Melvin, J. A., Montelaro, R. C. and Bomberger, J. M. (2016) Clinical potential of engineered cationic antimicrobial peptides against drug resistant biofilms. *Expert Rev Anti Infect Ther.* 1-3.
 14. Chung P.-Y., Bian Z.-X., Pun H.-Y., Chan D., Chan A. S.-C., Chui C.-H., Tang J. C.-O. and K.-H., L. (2015) Recent advances in research of natural and synthetic bioactive quinolines. *Future Med. Chem.* *7*, 947-967.
 15. Videnovic, M., Opsenica, D. M., Burnett, J. C., Gomba, L., Nuss, J. E., Selakovic, Z., Konstantinovic, J., Krstic, M., Segan, S., Zlatovic, M., Sciotti, R. J., Bavari, S. and Solaja, B. A. (2014) Second generation steroidal 4-aminoquinolines are potent, dual-target inhibitors of the botulinum neurotoxin serotype A metalloprotease and *P. falciparum* malaria. *J Med Chem.* *57*, 4134-4153.
 16. Abouelhassan, Y., Garrison, A. T., Burch, G. M., Wong, W., Norwood, V. M. t. and Huigens, R. W., 3rd (2014) Discovery of quinoline small molecules with potent dispersal activity against methicillin-resistant *Staphylococcus aureus* and *Staphylococcus epidermidis* biofilms using a scaffold hopping strategy. *Bioorg Med Chem Lett.* *24*, 5076-5080.
 17. Basak, A., Abouelhassan, Y. and Huigens, R. W., 3rd (2015) Halogenated quinolines discovered through reductive amination with potent eradication activities against MRSA, MRSE and VRE biofilms. *Org Biomol Chem.* *13*, 10290-10294.
 18. Leon, B., Haeckl, F. P. and Linington, R. G. (2015) Optimized quinoline amino alcohols as disruptors and dispersal agents of *Vibrio cholerae* biofilms. *Org Biomol Chem.* *13*, 8495-8499.
 19. Sobke, A., Klinger, M., Hermann, B., Sachse, S., Nietzsche, S., Makarewicz, O., Keller, P. M., Pfister, W. and Straube, E. (2012) The urinary antibiotic 5-nitro-8-hydroxyquinoline (Nitroxoline) reduces the formation and induces the dispersal of *Pseudomonas aeruginosa* biofilms by chelation of iron and zinc. *Antimicrob Agents Chemother.* *56*, 6021-6025.
 20. Eswaran, S., Adhikari, A. V., Chowdhury, I. H., Pal, N. K. and Thomas, K. D. (2010) New quinoline derivatives: synthesis and investigation of antibacterial and antituberculosis properties. *Eur J Med Chem.* *45*, 3374-3383.
 21. Lu, C., Kirsch, B., Zimmer, C., de Jong, J. C., Henn, C., Maurer, C. K., Musken, M., Haussler, S., Steinbach, A. and Hartmann, R. W. (2012) Discovery of antagonists of PqsR, a key player in 2-alkyl-4-quinolone-dependent quorum sensing in *Pseudomonas aeruginosa*. *Chem Biol.* *19*, 381-390.
 22. Lu, C., Kirsch, B., Maurer, C. K., de Jong, J. C., Braunshausen, A., Steinbach, A. and Hartmann, R. W. (2014) Optimization of anti-virulence PqsR antagonists regarding aqueous solubility and biological properties resulting in new insights in structure-activity relationships. *Eur J Med Chem.* *79*, 173-183.
 23. Sangshetti, J. N., Khan, F. A., Patil, R. H., Marathe, S. D., Gade, W. N. and Shinde, D. B. (2015) Biofilm inhibition of linezolid-like Schiff bases: synthesis, biological activity, molecular docking and in silico ADME prediction. *Bioorg Med Chem Lett.* *25*, 874-880.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
24. Zaheer, Z., Khan, F. A., Sangshetti, J. N., Patil, R. H. and Lohar, K. S. (2016) Novel amalgamation of phthalazine-quinolines as biofilm inhibitors: One-pot synthesis, biological evaluation and in silico ADME prediction with favorable metabolic fate. *Bioorg Med Chem Lett.* *26*, 1696-1703.
25. Ilangovan, A., Fletcher, M., Rampioni, G., Pustelny, C., Rumbaugh, K., Heeb, S., Camara, M., Truman, A., Chhabra, S. R., Emsley, J. and Williams, P. (2013) Structural basis for native agonist and synthetic inhibitor recognition by the *Pseudomonas aeruginosa* quorum sensing regulator PqsR (MvfR). *PLoS Pathog.* *9*, e1003508.
26. Zieba, A., Wojtyczka, R. D., Idzik, D. and Kepa, M. (2013) Synthesis and in vitro antimicrobial activity of 1-methyl-3-sulfonylthio-4-aminoquinolinium chlorides. *Acta Pol Pharm.* *70*, 163-166.
27. Medapi, B., Suryadevara, P., Renuka, J., Sridevi, J. P., Yogeewari, P. and Sriram, D. (2015) 4-Aminoquinoline derivatives as novel *Mycobacterium tuberculosis* GyrB inhibitors: Structural optimization, synthesis and biological evaluation. *Eur J Med Chem.* *103*, 1-16.
28. Morohoshi, T., Shiono, T., Takidouchi, K., Kato, M., Kato, N., Kato, J. and Ikeda, T. (2007) Inhibition of quorum sensing in *Serratia marcescens* AS-1 by synthetic analogs of N-acylhomoserine lactone. *Appl Environ Microbiol.* *73*, 6339-6344.
29. Wei, J. R. and Lai, H. C. (2006) N-acylhomoserine lactone-dependent cell-to-cell communication and social behavior in the genus *Serratia*. *Int J Med Microbiol.* *296*, 117-124.
30. Rice, S. A., Koh, K. S., Queck, S. Y., Labbate, M., Lam, K. W. and Kjelleberg, S. (2005) Biofilm formation and sloughing in *Serratia marcescens* are controlled by quorum sensing and nutrient cues. *J Bacteriol.* *187*, 3477-3485.
31. Mitchell, K. J., Abboud, K. A. and Christou, G. (2016) Magnetostructural Correlation for High-Nuclearity Iron(III)/Oxo Complexes and Application to Fe₅, Fe₆, and Fe₈ Clusters. *Inorg Chem.* *55*, 6597-6608.
32. Thorson, M. K., Puerta, D. T., Cohen, S. M. and Barrios, A. M. (2014) Inhibition of the lymphoid tyrosine phosphatase: the effect of zinc(II) ions and chelating ligand fragments on enzymatic activity. *Bioorg Med Chem Lett.* *24*, 4019-4022.
33. Rahier, R., Noiriél, A. and Abousalham, A. (2016) Development of a Direct and Continuous Phospholipase D Assay Based on the Chelation-Enhanced Fluorescence Property of 8-Hydroxyquinoline. *Anal Chem.* *88*, 666-674.
34. Segan, S., Terzic-Jovanovic, N., Milojkovic-Opsenica, D., Trifkovic, J., Solaja, B. and Opsenica, D. (2014) Correlation study of retention data and antimalarial activity of 1,2,4,5-mixed tetraoxanes with their molecular structure descriptors and LSER parameters. *J Pharm Biomed Anal.* *97*, 178-183.
35. Milosevic, N. P., Stojanovic, S. Z., Penov-Gasi, K., Perisic-Janjic, N. and Kaliszan, R. (2014) Reversed- and normal-phase liquid chromatography in quantitative structure retention-property relationships of newly synthesized seco-androstene derivatives. *J Pharm Biomed Anal.* *88*, 636-642.
36. Sabet, R. and Fassihi, A. (2008) QSAR study of antimicrobial 3-hydroxypyridine-4-one and 3-hydroxypyran-4-one derivatives using different chemometric tools. *Int J Mol Sci.* *9*, 2407-2423.
37. Varmuza, K. and Filzmoser, P. (2009) Introduction to multivariate statistical analysis in chemometrics. Taylor & Francis Group, Boca Raton, Florida, USA.
38. Todeschini, R. and Cosonni, V. (2000) Handbook of Molecular Descriptors, Methods and Principles in Medicinal Chemistry. Vol. 11, Wiley-VCH, GmbH, Weinheim, Germany.
39. Lu, C., Maurer, C. K., Kirsch, B., Steinbach, A. and Hartmann, R. W. (2014) Overcoming the unexpected functional inversion of a PqsR antagonist in *Pseudomonas aeruginosa*: an in vivo potent antivirulence agent targeting pqs quorum sensing. *Angew Chem Int Ed Engl.* *53*, 1109-1112.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
40. Starkey, M., Lepine, F., Maura, D., Bandyopadhaya, A., Lesic, B., He, J., Kitao, T., Righi, V., Milot, S., Tzika, A. and Rahme, L. (2014) Identification of anti-virulence compounds that disrupt quorum-sensing regulated acute and persistent pathogenicity. *PLoS Pathog.* *10*, e1004321.
41. Hodgkinson, J., Bowden, S. D., Galloway, W. R., Spring, D. R. and Welch, M. (2010) Structure-activity analysis of the *Pseudomonas* quinolone signal molecule. *J Bacteriol.* *192*, 3833-3837.
42. Lee, J. and Zhang, L. (2015) The hierarchy quorum sensing network in *Pseudomonas aeruginosa*. *Protein Cell.* *6*, 26-41.
43. Thomson, N. R., Crow, M. A., McGowan, S. J., Cox, A. and Salmond, G. P. (2000) Biosynthesis of carbapenem antibiotic and prodigiosin pigment in *Serratia* is under quorum sensing control. *Mol Microbiol.* *36*, 539-556.
44. Dubern, J. F. and Diggle, S. P. (2008) Quorum sensing by 2-alkyl-4-quinolones in *Pseudomonas aeruginosa* and other bacterial species. *Mol Biosyst.* *4*, 882-888.
45. Rada, B. and Leto, T. L. (2013) Pyocyanin effects on respiratory epithelium: relevance in *Pseudomonas aeruginosa* airway infections. *Trends Microbiol.* *21*, 73-81.
46. Klein, T., Henn, C., de Jong, J. C., Zimmer, C., Kirsch, B., Maurer, C. K., Pistorius, D., Muller, R., Steinbach, A. and Hartmann, R. W. (2012) Identification of small-molecule antagonists of the *Pseudomonas aeruginosa* transcriptional regulator PqsR: biophysically guided hit discovery and optimization. *ACS Chem Biol.* *7*, 1496-1501.
47. Terzic, N., Konstantinovic, J., Tot, M., Burojevic, J., Djurkovic-Djakovic, O., Srbljanovic, J., Stajner, T., Verbic, T., Zlatovic, M., Machado, M., Albuquerque, I. S., Prudencio, M., Sciotti, R. J., Pecic, S., D'Alessandro, S., Taramelli, D. and Solaja, B. A. (2016) Reinvestigating Old Pharmacophores: Are 4-Aminoquinolines and Tetraoxanes Potential Two-Stage Antimalarials? *J Med Chem.* *59*, 264-281.
48. Merritt, J. H., Kadouri, D. E. and O'Toole, G. A. (2005) Growing and analyzing static biofilms. *Curr Protoc Microbiol. Chapter 1*, Unit 1B 1.
49. Pekmezovic, M., Aleksic, I., Barac, A., Arsic-Arsenijevic, V., Vasiljevic, B., Nikodinovic-Runic, J. and Senerovic, L. (2016) Prevention of polymicrobial biofilms composed of *Pseudomonas aeruginosa* and pathogenic fungi by essential oils from selected Citrus species *Pathog Dis.* *74*, ftw102.
50. Massai, F., Imperi, F., Quattrucci, S., Zennaro, E., Visca, P. and Leoni, L. (2011) A multitask biosensor for micro-volumetric detection of N-3-oxo-dodecanoyl-homoserine lactone quorum sensing signal. *Biosens Bioelectron.* *26*, 3444-3449.
51. Duan, K. and Surette, M. G. (2007) Environmental regulation of *Pseudomonas aeruginosa* PAO1 Las and Rhl quorum-sensing systems. *J Bacteriol.* *189*, 4827-4836.
52. Fletcher, M. P., Diggle, S. P., Crusz, S. A., Chhabra, S. R., Camara, M. and Williams, P. (2007) A dual biosensor for 2-alkyl-4-quinolone quorum-sensing signal molecules. *Environ Microbiol.* *9*, 2683-2693.
53. Rocha, G. B., Freire, R. O., Simas, A. M. and Stewart, J. J. (2006) RM1: a reparameterization of AM1 for H, C, N, O, P, S, F, Cl, Br, and I. *J Comput Chem.* *27*, 1101-1111.

1
2
3
4 **Figure legends:**
5

6 **Figure 1.** Chemical structures of tested compounds.
7

8 **Figure 2.** Regression coefficients corresponding to the QSAR models of $\log BI_{50}$ (a), and
9 $\log BI_{25}$ (b). Variables with VIP score > 1 are marked in bold.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Chemical structures of tested compounds.

182x218mm (300 x 300 DPI)





Figure 2. Regression coefficients corresponding to the QSAR models of logBI₅₀ (a), and logBI₂₅ (b). Variables with VIP score > 1 are marked in bold.

140x150mm (300 x 300 DPI)



74x39mm (300 x 300 DPI)