

 Open access • Journal Article • DOI:10.1038/NATURE04787

Long γ -ray bursts and core-collapse supernovae have different environments

— [Source link](#) 

A. S. Fruchter, Andrew J. Levan, Andrew J. Levan, Andrew J. Levan ...+38 more authors

Institutions: Space Telescope Science Institute, University of Leicester, University of Hertfordshire, Western Kentucky University ...+17 more institutions

Published on: 25 May 2006 - Nature (Nature Publishing Group)

Topics: Gamma-ray burst, Galaxy, Gravitational collapse, Supernova and Stars

Related papers:

- [A very energetic supernova associated with the \$\gamma\$ -ray burst of 29 March 2003](#)
- [Gamma-ray bursts from stellar mass accretion disks around black holes](#)
- [The Supernova Gamma-Ray Burst Connection](#)
- [Collapsars: Gamma-ray bursts and explosions in 'failed supernovae'](#)
- [The Observed Offset Distribution of Gamma-Ray Bursts from Their Host Galaxies: A Robust Clue to the Nature of the Progenitors* **](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/long-g-ray-bursts-and-core-collapse-supernovae-have-15xy3xemt6>

LJMU Research Online

Fruchter, AS, Levan, AJ, Strolger, L, Vreeswijk, PM, Thorsett, SE, Bersier, D, Burud, I, Castro Ceron, JM, Castro-Tirado, AJ, Conselice, C, Dahlen, T, Ferguson, HC, Fynbo, JPU, Garnavich, PM, Gibbons, RA, Gorosabel, J, Gull, TR, Hjorth, J, Holland, ST, Kouveliotou, C, Levay, Z, Livio, M, Metzger, MR, Nugent, PE, Petro, L, Pian, E, Rhoads, JE, Riess, AG, Sahu, KC, Smette, A, Tanvir, NR, Wijers, RAMJ and Woosley, SE

Long gamma-ray bursts and core-collapse supernovae have different environments

<http://researchonline.ljmu.ac.uk/id/eprint/4882/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Fruchter, AS, Levan, AJ, Strolger, L, Vreeswijk, PM, Thorsett, SE, Bersier, D, Burud, I, Castro Ceron, JM, Castro-Tirado, AJ, Conselice, C, Dahlen, T, Ferguson, HC, Fynbo, JPU, Garnavich, PM, Gibbons, RA, Gorosabel, J, Gull, TR, Hjorth, J, Holland, ST, Kouveliotou, C, Levay, Z, Livio, M, Metzger, MR.

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

<http://researchonline.ljmu.ac.uk/>

For more information please contact researchonline@ljmu.ac.uk

Long γ -ray bursts and core-collapse supernovae have different environments

A.S. Fruchter¹, A.J. Levan^{1,2,3}, L. Strolger^{1,4}, P.M. Vreeswijk⁵, S.E. Thorsett⁶, D. Bersier^{1,7},
I. Burud^{1,8}, J.M. Castro Cerón^{1,9}, A.J. Castro-Tirado¹⁰, C. Conselice^{11,12}, T. Dahlen¹³, H.C. Ferguson¹,
J.P.U. Fynbo⁹, P.M. Garnavich¹⁴, R.A. Gibbons^{1,15}, J. Gorosabel^{1,10}, T.R. Gull¹⁶, J. Hjorth⁹, S.T. Holland¹⁷,
C. Kouveliotou¹⁸, Z. Levay¹, M. Livio¹, M.R. Metzger¹⁹, P.E. Nugent²⁰, L. Petro¹, E. Pian²¹,
J.E. Rhoads¹, A.G. Riess¹, K.C. Sahu¹, A. Smette⁵, N.R. Tanvir³, R.A.M.J. Wijers²², S.E. Woosley⁶

¹*Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA*

²*Department of Physics and Astronomy, University of Leicester, University Road, Leicester, LE1 7RH, UK*

³*Centre for Astrophysics Research, University of Hertfordshire, College Lane, Hatfield, AL10 9AB, UK*

⁴*Physics & Astronomy, TCCW 246, Western Kentucky University, 1 Big Red Way, Bowling Green, KY 42101, USA*

⁵*European Southern Observatory, Alonso de Córdova 3107, Casilla 19001, Santiago, Chile*

⁶*Dept of Astronomy & Astrophysics, University of California, 1156 High St, Santa Cruz, CA 95064, USA*

⁷*Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead, CH41 1LD*

⁸*Norwegian Meteorological Institute, P.O. Box 43, Blindern, N-0313 Oslo, Norway*

⁹*Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, DK-2100 Copenhagen,*

Denmark

¹⁰*Instituto de Astrofísica de Andalucía (CSIC), Camino Bajo de Huétor, 50, 18008 Granada, Spain.*

¹¹*California Institute of Technology, Mail Code 105-24, Pasadena, CA 91125, USA*

¹²*School of Physics and Astronomy, University of Nottingham, University Park, United Kingdom,
NG7 2RD*

¹³*Department of Physics, Stockholm University, SE-106 91 Stockholm, Sweden*

¹⁴*Physics Department, University of Notre Dame, 225 Nieuwland Hall, Notre Dame, IN 46556,
USA*

¹⁵*Vanderbilt University, Dept. of Physics and Astronomy, 6301 Stevenson Center, Nashville, TN
37235, USA*

¹⁶*Code 667 Extraterrestrial Planets and Stellar Astrophysics, Exploration of the Universe Division,
Goddard Space Flight Center, Greenbelt, MD 20771, USA*

¹⁷*Code 660.1, NASA's GSFC, Greenbelt, MD 20771, USA*

¹⁸*NASA/Marshall Space Flight Center, VP-62, National Space Science & Technology Center, 320
Sparkman Drive, Huntsville, AL 35805, USA*

¹⁹*Renaissance Technologies Corporation, 600 Route 25A, East Setauket, New York 11733*

²⁰*Lawrence Berkeley National Laboratory, M.S. 50F-1650, 1 Cyclotron Road, Berkeley, CA
94720, USA*

²¹*INAF, Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, I-34131 Trieste, Italy*

²²*Astronomical Institute 'Anton Pannekoek', University of Amsterdam, Kruislaan 403, NL-1098
SJ Amsterdam, The Netherlands*

This paper has been accepted for publication in Nature. In accordance with that journal's editorial policy we ask that you do not discuss this work with the Press until it appears in Nature either online or in print. Thank you.

When massive stars exhaust their fuel they collapse and often produce the extraordinarily bright explosions known as core-collapse supernovae. On occasion, this stellar collapse also powers an even more brilliant relativistic explosion known as a long-duration γ -ray burst. One would then expect that long γ -ray bursts and core-collapse supernovae should be found in similar galactic environments. Here we show that this expectation is wrong. We find that the long γ -ray bursts are far more concentrated on the very brightest regions of their host galaxies than are the core-collapse supernovae. Furthermore, the host galaxies of the long γ -ray bursts are significantly fainter and more irregular than the hosts of the core-collapse supernovae. Together these results suggest that long-duration γ -ray bursts are associated with the most massive stars and may be restricted to galaxies of limited chemical evolution. Our results directly imply that long γ -ray bursts are relatively rare in galaxies such as our own Milky Way.

It is an irony of astrophysics that stellar birth is most spectacularly marked by the deaths of massive stars. Massive stars burn brighter and hotter than smaller stars, and exhaust their fuel far more rapidly. Therefore a region of star formation filled with low mass stars still early in their lives, and in some cases still forming, may also host massive stars already collapsing and producing supernovae. Indeed, with the exception of the now famous Type Ia supernovae, which

have been so successfully used for cosmological studies^{1,2} and which are thought to be formed by the uncontrolled nuclear burning of stellar remnants comparable in mass to the sun³, all supernovae are thought to be produced by the collapse of massive stars. The collapse of the very most massive stars (tens of solar masses) is thought to leave behind either black holes or neutron stars, depending largely on the state of chemical evolution of the material that formed the star, while the demise of stars between approximately 8 and 20 solar masses produces only neutron stars⁴.

Gamma-ray bursts (GRBs), like supernovae, are a heterogeneous population. GRBs can be divided into two classes: short, hard bursts, which last between milliseconds and about two seconds and have hard high-energy spectra, and long, soft bursts, which last between two and tens of seconds, and have softer high-energy spectra⁵. Only very recently have a few of the short bursts been well localized, and initial studies of their apparent hosts suggest that these bursts may be formed by the binary merger of stellar remnants^{6,7}. In contrast, the afterglows of over eighty long GRBs (LGRBs) have been detected in the optical and/or radio. And as a result of these detections, it has become clear that LGRBs, like core-collapse supernovae, are related to the deaths of young, massive stars. It is these objects, born of the deaths of massive stars, that we study here.

LGRBs are generally found in extremely blue host galaxies^{8–11} which exhibit strong emission lines^{12,13} suggesting a significant abundance of young, very massive stars. Furthermore while the light curves of the optical transients (OTs) associated with LGRBs are often dominated by radiation from the relativistic outflow of the GRB, numerous LGRBs have shown late-time “bumps” in their light-curves consistent with the presence of an underlying SN^{14–16}. In several cases spectroscopic

evidence has provided confirmation of the light of a SN superposed on the OT^{17–20}. Indeed, given the large variations in the brightnesses of OTs and supernovae, and the limited observations on some GRBs, it seems plausible that *all* LGRBs have an underlying SN²¹. Furthermore, while the energy released in a LGRB often appears to the observer to be orders of magnitude larger than that of a supernovae, there is now good evidence suggesting that most LGRBs are highly collimated and often illuminate only a few percent of the sky^{22,23}. When one takes this into account, the energy released in LGRBs more closely resembles that of energetic supernovae. However, not all core-collapse supernovae may be candidates for the production of GRBs. The supernovae with good spectroscopic identifications so far associated with GRBs have been Type Ic – that is cc supernovae which show no evidence of hydrogen or helium in their spectra. (Type Ib supernovae, which are often studied together with Type Ic, have spectra which are also largely devoid of hydrogen lines but show strong helium features.) A star may therefore need to lose its outer envelope if a GRB is to be able to burn its way through the stellar atmosphere²⁴. Studies which have compared the locations Type Ib/c supernovae with the more numerous Type II supernovae (cc supernovae showing hydrogen lines) in local galaxies so far show no differences in either the type of host or the placement of the explosion on the host^{25,26}. This result led Ref. 25 to argue cc supernovae all come from the same mass range of progenitor stars, but that Type Ib/c supernovae may have had their envelopes stripped by interaction with a binary stellar companion. Whether Type Ic supernovae come from single stars, or binary stars, or both, it is very likely that only small fraction of these supernovae produce GRBs²⁷.

Given the common massive stellar origins of cc supernovae and LGRBs, one might expect

that their hosts and local environments might be quite similar. It has long been argued that cc supernovae should track the blue light in the universe (the light from massive stars is blue), both in their distribution among galaxies and within their host galaxies themselves. One would expect similar behavior from LGRBs, and indeed rough evidence for such a correlation has been reported²⁸. Here we use the high resolution available from Hubble Space Telescope (*HST*) images, and an analytical technique developed by us that is independent of galaxy morphology, to study the correlation between these objects and the light of their hosts. We also compare the sizes, morphologies and brightnesses of the LGRB hosts with those of the supernovae. Our results reveal surprising and substantial differences between the birth places of these cosmic explosions. We find that while cc supernovae trace the blue-light of their hosts, GRBs are far more concentrated on the brightest regions of their hosts. Furthermore, while the hosts of cc supernovae are approximately equally divided between spiral and irregular galaxies, the overwhelming majority of GRBs are on irregulars, even when we restrict the GRB sample to the same redshift range as the SN sample. We argue that these results may be best understood if GRBs are formed from the collapse of extremely massive, low-metallicity stars.

1 The Sample

Over forty LGRBs have been observed with *HST* at various times after outburst. *HST* is unique in its capability to easily resolve the distant hosts of these objects. Shown in Figure 1 is a mosaic of *HST* images of the hosts of forty-two bursts. These are all LGRBs with public data which had an afterglow detected with better than three-sigma significance and a position sufficiently well

localized to determine a host galaxy. A list of all the GRBs used in this work can be found in Tables 1—3 of the Supplementary Material.

The supernovae discussed in this *Article* were all discovered as part of the *Hubble* Higher z Supernova Search^{29,30}, which was done in cooperation with the *HST* GOODS survey³¹. The GOODS survey observed two ~ 150 sq. arcminute patches of sky five times each in epochs separated by forty-five days. Supernovae were identified by image subtraction. In this paper we discuss only the cc supernovae identified in this survey. A list of the supernovae used is presented in Table 4 of the Supplementary Material, and images of the supernovae hosts can be seen in Figure 2.

2 Positions of GRBs and supernovae on their Hosts

If LGRBs do in fact trace massive star formation, then in the absence of strong extinction we should find a close correlation between their position on their host galaxies and the blue light of those galaxies. However, many of the GRB hosts and quite a few of the supernovae hosts are irregular galaxies made up of more than one bright component. As a result the common astronomical procedure of identifying the centroid of the galaxy’s light, and then determining the distance of the object in question from the centroid is not particularly appropriate for these galaxies – the centroid of light may in fact lie on a rather faint region of the host (examine GRBs 000926 and 020903 in Figure 1 for excellent illustrations of this effect). We therefore have developed a method which is independent of galaxy morphology. We sort all of the pixels of the host galaxy image from faintest to brightest and ask what fraction of the total light of the host is contained in pixels fainter than or

equal to the pixel containing the explosion. If the explosions track the distribution of light, then the fraction determined by this method should be uniformly distributed between zero and one. (A detailed exposition of this method can be found in the supplementary materials).

As can be seen in Figure 3, the cc supernovae do track the light of their hosts as well as could be expected given their small number statistics. A Kolmogorov-Smirnov (KS) test finds that the distribution of the supernovae is indistinguishable from the distribution of the underlying light. The situation is clearly different for LGRBs. As can be seen in Figure 3, the GRBs do not simply trace the blue light of the hosts, rather they are far more concentrated on the peaks of light in the hosts than the light itself. A KS test rejects the hypothesis that GRBs are distributed as the light of their hosts with a probability greater than 99.98%. Furthermore, this result is robust: it shows no dependence on GRB host size or magnitude. And in spite of the relatively small number of SN hosts on which a comparison can be made, the two populations are found by the KS test to be drawn from different distributions with $\sim 99\%$ certainty. In the next section of this paper we show that the surprising differences in the locations of these objects on the underlying light of their hosts may be due not only to the nature of their progenitor stars but also that of their hosts.

3 A Comparison of the Host Populations

An examination of the mosaics of the GRB and SN hosts (Figures 1 and 2) immediately shows a remarkable contrast – only one GRB host in this set of 42 galaxies is a grand-design spiral, while nearly half of the SN hosts are grand-design spirals. One might wonder if this effect is due to a

difference in redshift distribution – the cc supernovae discovered by the GOODS collaboration all lie at $z < 1.2$, while LGRBs can be found at much larger redshifts where grand-design spirals are rare to non-existent. Yet if we restrict the GRB population to $z < 1.2$ (and thus produce a population with a nearly identical mean and standard deviation in redshift space compared to the GOODS cc supernovae), the situation remains essentially unchanged: only one out of the eighteen GRB hosts is a grand-design spiral. (For a detailed comparison of GRB hosts to field galaxies, rather than the SN selected galaxies shown here, see Ref. 32).

Were the difference in spiral fraction the only indication of a difference in the host populations, we could not rule out random chance – given the small number statistics both populations are barely consistent with each other and a spiral fraction of $\sim 25\%$. However, the host populations differ strongly in ways other than morphology.

In Figure 4 we compare the 80% light radius (r_{80}) and absolute magnitude distributions of the GRB and supernovae hosts. Included in the comparison are all LGRBs with known redshifts $z < 1.2$ at the time of submission and the 16 cc supernovae of GOODS with spectroscopic or photometric redshifts (See the Supplementary Tables for a complete list of the GRBs, supernovae and associated parameters used in this study). The small minority of GRB hosts in this redshift range without *HST* imaging are compared only in absolute magnitude and not in size. The absolute magnitudes have been derived from the observed photometry using a cosmology of $\Omega_m = 0.27$, $\Lambda = 0.73$, and $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, and the magnitudes have been corrected for foreground Galactic extinction³³. For a technical discussion of the determination of the magnitude

and size of individual objects, please see the Supplementary Materials.

As can be readily seen the two host populations differ substantially both in their intrinsic magnitudes and sizes. The GRB hosts are fainter and smaller than the SN hosts. Indeed KS tests reject the hypothesis that these two populations are drawn from the same population with greater than 98.6% and 99.7% certainty for the magnitude and size distributions, respectively.

4 Discussion

Although the evidence is now overwhelming that both cc supernovae and LGRBs are formed by the collapse of massive stars, our observations show that the distribution of LGRBs and cc supernovae on their hosts, and the nature of their hosts themselves are substantially different. How then can this be? We propose here that these surprising findings are the result of the dependence of the probability of GRB formation on the state of the chemical evolution of massive stars in a galaxy.

Even before the association of LGRBs with massive stars had been established, a number of theorists had suggested that these objects could be formed by the collapse of massive stars, which would leave behind rapidly spinning black holes. An accretion disk about the black hole would power the GRB jet. These models, sometimes referred to as “hypernovae” or “collapsar” models implicitly require very massive stars, since only stars greater than about 18 solar masses form black holes. But in fact it was widely suspected that even more massive stars would be required – if only to provide the required large energies, and to limit the numbers of supernovae progressing to GRBs.

We conclude that LGRBs do indeed form from the very most massive stars and this is the reason that they are even more concentrated on the blue light of their hosts than the light itself. The most massive stars (O stars) are frequently found in large associations. These associations can be extremely bright, and can indeed provide the peak of the light of a galaxy – particularly if that galaxy is a faint, blue irregular, as are the GRB hosts in general. Indeed, a connection of LGRBs with O-stars (and perhaps Wolf-Rayet stars) is a natural one – given the strong emission lines (including Ne [III]) seen in many of these hosts^{12,13} and the evidence for possible strong winds off of the progenitors of the GRBs seen in absorption in some LGRB spectra^{34,35}.

However, O stars are found in galaxies of all sizes. Indeed, studies of the Magellanic clouds suggest the initial distribution of masses of stars at formation in these dwarf galaxies is essentially identical to that in our much larger spiral, the Milky Way³⁶. Therefore, a difference in the initial mass function of stars is unlikely to be responsible for the differences between the hosts. We propose that the fundamental differences between the LGRB and SN host populations is not their size or luminosity, but rather their metallicity, or chemical evolution. Some evidence of this already exists. The hosts of seven LGRBs (GRBs 980425 (P. M. Vreeswijk, personal communication), 990712¹³, 020903³⁷, 030323³⁸, 030329¹⁷, 031203³⁹ and 050730⁴⁰) have measurements of or limits on their metallicity, and in all cases the metallicity is less than one-third solar. The small size and low luminosity of the GRB hosts is then a result of the well known correlation between galaxy mass and metallicity (see Ref. 41 and references therein).

But why do LGRBs choose low-metallicity galaxies? This may be a direct result of the evo-

lution of the most massive stars. It has recently been proposed that metal rich stars with masses of tens of solar masses have such large winds off their surfaces (due to the photon pressure on their metal rich atmospheres) that they lose most of their mass before they collapse and produce supernovae⁴. As a result they leave behind neutron stars, not the black holes necessary for LGRB formation. Ironically, stars of 15-30 solar masses may still form black holes, as they do not possess radiation pressure sufficient to drive off their outer envelopes. Direct evidence for this scenario comes from recent work showing that the Galactic soft gamma-ray repeater, SGR 1820-06, is in a cluster of extremely young stars of which the most massive have only started to collapse⁴² – yet, the progenitor of SGR 1820-06 collapsed to a neutron star, not a black hole. Recent observations of winds from very massive (Wolf-Rayet) stars provide further support for this scenario: outflows from the low-metallicity stars in the LMC are substantially smaller than those seen from more metal-rich Galactic stars⁴³. The possible importance of metallicity in LGRB formation has therefore not escaped the notice of theorists^{44,45}.

A preference for low-metallicity may also explain one of the most puzzling results of GRB host studies. None of the LGRB hosts is a red, sub-millimeter bright galaxy. These highly dust-enshrouded galaxies at redshifts of $\sim 1 - 3$ are believed to be the site of a large fraction of the star formation in the distant universe⁴⁶. And while some LGRB hosts do show sub-mm emission, none has the red colors characteristic of the majority of this population. However, it is likely that these red dusty galaxies have substantial metallicities at all redshifts. The low-metallicity of hosts may also help explain the fact that a substantial fraction of high-redshift LGRB hosts display strong Lyman-alpha emission⁴⁷.

All well classified supernovae associated with LGRBs are Types Ic, presumably because the presence of a hydrogen envelope about the collapsing core can block the emergence of a GRB jet²⁴. Thus only those supernovae whose progenitors have lost some, but not too much mass, appear to be candidates for the formation of a GRB. Given the large numbers of Type Ic supernovae in comparison to the estimated numbers of LGRBs however, it is likely that only a small fraction of Ic supernovae produce LGRBs. Indeed, even the number of unusually energetic Type Ib/Ic supernovae appears to dwarf the LGRB population⁴⁸. Another process, perhaps the spin-up of the progenitor in a binary²⁷, may decide which Type Ic supernovae produce LGRBs. Interestingly, it was the similar distribution of supernovae on their hosts, and particularly the fact that Type Ib/Ic were no more correlated than Type II supernovae with the UV bright regions of their hosts, that led Ref 25 to the conclusion that Type Ib/Ic form from binaries. LGRBs clearly track light differently than the general Type Ic population. However the samples used by Refs 25, 26 were from supernovae largely discovered on nearby massive galaxies – dwarf irregular hosts are underrepresented in these samples. It will be particularly interesting to see whether large unbiased SN surveys presently underway produce similar locations for their supernovae.

We do not know, however, what separates the small fraction of low-metallicity Type Ic supernovae which turn into LGRBs from the rest of the population. Potentially, the answer is the amount of angular momentum available in the core to form the jet. In this case, the preference for low-metallicity may indicate that single star evolution dominates over binary interaction in forming LGRBs. Deep, high spectral resolution studies of LGRB afterglows may provide insight here, by allowing a studies of the winds off of the progenitor and any binary companion.

Only a small fraction of LGRBs are found in spiral galaxies, even in LGRBs with redshifts $z < 1$ where spirals are much more common. However, the local metallicity in spirals is known to be anti-correlated with distance from the center of the galaxy. Thus one might expect LGRBs in spirals to violate the trend we have seen for the general LGRB population and avoid the bright central regions of their hosts. The present number of LGRBs known in spirals is still too small to test this prediction. But a sample size a few times larger should begin to allow such a test. Additionally, a survey of the metallicity of the hosts of the GOODS supernovae should find a higher average metallicity than that seen in GRB hosts. Finally, if low-metallicity is indeed the primary variable in determining whether LGRBs are produced, then as we observe higher redshifts, where metallicities are lower than in most local galaxies, LGRBs should be more uniformly distributed among star-forming galaxies. Indeed, some evidence of this may already be present in the data³². LGRBs, however, are potentially visible to redshifts as high as $z \sim 10$. At significant redshifts, where the metallicities of even relatively large galaxies are expected to be low, we may find that LGRBs do become nearly unbiased tracers of star formation.

References

1. Riess, A. G. *et al.* Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. *Astron. J.* **116**, 1009–1038 (1998).
2. Perlmutter, S. *et al.* Measurements of Omega and Lambda from 42 High-Redshift Supernovae. *Astrophys. J.* **517**, 565–586 (1999).
3. Branch, D., Livio, M., Yungelson, L. R., Boffi, F. R. & Baron, E. In Search of the Progenitors

- of Type IA Supernovae. *Publ. Astr. Soc. Pacific* **107**, 1019–1028 (1995).
4. Heger, A., Fryer, C. L., Woosley, S. E., Langer, N. & Hartmann, D. H. How Massive Single Stars End Their Life. *Astrophys. J.* **591**, 288–300 (2003).
 5. Kouveliotou, C. *et al.* Identification of two classes of gamma-ray bursts. *Astrophys. J.* **413**, 101–104 (1993).
 6. Gehrels, N. *et al.* A short γ -ray burst apparently associated with an elliptical galaxy at redshift $z = 0.225$. *Nature* **437**, 851–854 (2005).
 7. Prochaska, J. X. *et al.* The galaxy hosts and large-scale environments of short-hard γ -ray bursts. *Astrophys. J.* (2006). Accepted, astro-ph/0510022.
 8. Fruchter, A. S. *et al.* HST and Palomar imaging of GRB 990123: Implications for the nature of gamma-ray bursts and their hosts. *Astrophys. J.* **519**, 13–16 (1999).
 9. Sokolov, V. V. *et al.* Host galaxies of gamma-ray bursts: Spectral energy distributions and internal extinction. *Astr. Astrophys.* **372**, 438–455 (2001).
 10. Le Floc'h, E. *et al.* Are the hosts of gamma-ray bursts sub-luminous and blue galaxies? *Astr. Astrophys.* **400**, 499–510 (2003).
 11. Christensen, L., Hjorth, J. & Gorosabel, J. UV star-formation rates of GRB host galaxies. *Astr. Astrophys.* **425**, 913–926 (2004).
 12. Bloom, J. S., Djorgovski, S. G., Kulkarni, S. R. & Frail, D. A. The host galaxy of GRB 970508. *Astrophys. J.* **507**, L25–L28 (1998).

13. Vreeswijk, P. M. *et al.* VLT spectroscopy of GRB 990510 and GRB 990712: Probing the faint and bright ends of the gamma-ray burst host galaxy population. *Astrophys. J.* **546**, 672–680 (2001).
14. Bloom, J. S. *et al.* The unusual afterglow of the gamma-ray burst of 26 March 1998 as evidence for a supernova connection. *Nature* **401**, 453–456 (1999).
15. Galama, T. J. *et al.* Evidence for a supernova in reanalyzed optical and near-infrared images of GRB 970228. *Astrophys. J.* **536**, 185–194 (2000).
16. Levan, A. *et al.* GRB 020410: A Gamma-Ray Burst Afterglow Discovered by Its Supernova Light. *Astrophys. J.* **624**, 880–888 (2005).
17. Hjorth, J. *et al.* A very energetic supernova associated with the γ -ray burst of 29 March 2003. *Nature* **423**, 847–850 (2003).
18. Stanek, K. Z. *et al.* Spectroscopic Discovery of the Supernova 2003dh Associated with GRB 030329. *Astrophys. J.* **591**, L17–L20 (2003).
19. Della Valle, M. *et al.* Evidence for supernova signatures in the spectrum of the late-time bump of the optical afterglow of GRB 021211. *Astr. Astrophys.* **406**, L33–L37 (2003).
20. Malesani, D. *et al.* SN 2003lw and GRB 031203: A Bright Supernova for a Faint Gamma-Ray Burst. *Astrophys. J.* **609**, L5–L8 (2004).
21. Zeh, A., Klose, S. & Hartmann, D. H. A Systematic Analysis of Supernova Light in Gamma-Ray Burst Afterglows. *Astrophys. J.* **609**, 952–961 (2004).

22. Panaitescu, A. & Kumar, P. Fundamental Physical Parameters of Collimated Gamma-Ray Burst Afterglows. *Astrophys. J.* **560**, L49–L53 (2001).
23. Frail, D. A. *et al.* Beaming in Gamma-Ray Bursts: Evidence for a Standard Energy Reservoir. *Astrophys. J.* **562**, L55–58 (2001).
24. MacFadyen, A. I., Woosley, S. E. & Heger, A. Supernovae, Jets, and Collapsars. *Astrophys. J.* **550**, 410–425 (2001).
25. van Dyk, S. D., Hamuy, M. & Filippenko, A. V. Supernovae and Massive Star Formation Regions. *Astron. J.* **111**, 2017–2027 (1996).
26. van den Bergh, S., Li, W. & Filippenko, A. V. Classifications of the Host Galaxies of Supernovae, Set III. *Publ. Astr. Soc. Pacific* **117**, 773–782 (2005).
27. Podsiadlowski, P., Mazzali, P. A., Nomoto, K., Lazzati, D. & Cappellaro, E. The Rates of Hypernovae and Gamma-Ray Bursts: Implications for Their Progenitors. *Astrophys. J.* **607**, L17–L20 (2004).
28. Bloom, J. S., Kulkarni, S. R. & Djorgovski, S. G. The Observed Offset Distribution of Gamma-Ray Bursts from Their Host Galaxies: A Robust Clue to the Nature of the Progenitors. *Astron. J.* **123**, 1111–1148 (2002).
29. Riess, A. G. *et al.* Identification of Type Ia Supernovae at Redshift 1.3 and Beyond with the Advanced Camera for Surveys on the Hubble Space Telescope. *Astrophys. J.* **600**, L163–L166 (2004).

30. Strolger, L.-G. *et al.* The Hubble Higher z Supernova Search: Supernovae to $z \sim 1.6$ and Constraints on Type Ia Progenitor Models. *Astrophys. J.* **613**, 200–223 (2004).
31. Giavalisco, M. *et al.* The Great Observatories Origins Deep Survey: Initial Results from Optical and Near-Infrared Imaging. *Astrophys. J.* **600**, L93–L98 (2004).
32. Conselice, C. J. *et al.* Gamma-Ray Burst Selected High Redshift Galaxies: Comparison to Field Galaxy Populations to $z \sim 3$. *Astrophys. J.* **633**, 29–40 (2005).
33. Schlegel, D. J., Finkbeiner, D. P. & Davis, M. Maps of dust infrared emission for use in estimation of reddening and cosmic microwave background radiation foregrounds. *Astrophys. J.* **500**, 525–553 (1998).
34. Mirabal, N. *et al.* GRB 021004: A Possible Shell Nebula around a Wolf-Rayet Star Gamma-Ray Burst Progenitor. *Astrophys. J.* **595**, 935–949 (2003).
35. Klose, S. *et al.* Probing a Gamma-Ray Burst Progenitor at a Redshift of $z = 2$: A Comprehensive Observing Campaign of the Afterglow of GRB 030226. *Astron. J.* **128**, 1942–1954 (2004).
36. Weidner, C. & Kroupa, P. Variations of the imf. In Corbelli, E., Pila, F. & Zinnecker, H. (eds.) *The Initial Mass Function 50 Year Later*, 125–186 (Springer, Dordrecht, The Netherlands, 2005).
37. Bersier, D. *et al.* Evidence for a supernova associated with the x-ray flash 020903. *Astrophys. J.* (2005). Submitted to Ap. J.

38. Vreeswijk, P. M. *et al.* The host of GRB 030323 at $z=3.372$: A very high column density DLA system with a low metallicity. *Astr. Astrophys.* **419**, 927–940 (2004).
39. Prochaska, J. X. *et al.* The Host Galaxy of GRB 031203: Implications of Its Low Metallicity, Low Redshift, and Starburst Nature. *Astrophys. J.* **611**, 200–207 (2004).
40. Chen, H.-W., Prochaska, J. X., Bloom, J. S. & Thompson, I. B. Echelle Spectroscopy of a GRB Afterglow at $z=3.969$: A New Probe of the Interstellar and Intergalactic Media in the Young Universe. *Astrophys. J.* **634**, L25–L28 (2005).
41. Kobulnicky, H. A. & Kewley, L. J. Metallicities of galaxies in the GOODS-North Field. *Astrophys. J.* **617**, 240–261 (2004).
42. Figer, D. F., Najarro, F., Geballe, T. R., Blum, R. D. & Kudritzki, R. P. Massive Stars in the SGR 1806-20 Cluster. *Astrophys. J.* **622**, L49–L52 (2005).
43. Crowther, P. A. & Hadfield, L. J. Reduced wolf-rayet line luminosities at low metallicity. *Astr. Astrophys.* 711–722 (2006).
44. Woosley, S. & Heger, A. The Progenitor Stars of Gamma-Ray Bursts. *Astrophys. J.* **637**, 914–921 (2006).
45. Yoon, S.-C. & Langer, N. Evolution of rapidly rotating metal-poor massive stars towards gamma-ray bursts. *Astr. Astrophys.* 643–648 (2005).
46. Chapman, S. C., Blain, A. W., Smail, I. & Ivison, R. J. A Redshift Survey of the Submillimeter Galaxy Population. *Astrophys. J.* **622**, 772–796 (2005).

47. Fynbo, J. P. U. *et al.* On the Ly α emission from gamma-ray burst host galaxies: Evidence for low metallicities. *Astr. Astrophys.* **406**, L63–L66 (2003).
48. Soderberg, A. M., Nakar, E., Kulkarni, S. R. & Berger, E. Late-time Radio Observations of 68 Type Ibc Supernovae: Strong Constraints on Off-Axis Gamma-ray Bursts. *Astrophys. J.* **638**, 930–937 (2006).
49. van den Bergh, S. & Tammann, G. A. Galactic and extragalactic supernova rates. *Ann. Rev. Astr. Ap.* **29**, 363–407 (1991).
50. Mannucci, F. *et al.* The supernova rate per unit mass. *Astr. Astrophys.* **433**, 807–814 (2005).

Figure I: A mosaic of GRB host galaxies imaged by *HST*. Each individual image corresponds to a square region on the sky $3''.75$ on a side. These images were taken with the Space Telescope Imaging Spectrograph (STIS), Wide-Field and Planetary Camera 2 (WFPC2) and the Advanced Camera for Surveys (ACS) on *HST*. In cases where the location of the GRB on the host is known to better than $0''.15$, the position of GRB is shown by a green mark. If the positional error is smaller than the point spread function of the image ($0''.07$ for STIS and ACS, $0''.13$ for WFPC2) the position is marked by a cross-hair, otherwise the positional error is indicated by a circle. The STIS images were all taken in white light (no filter) and in most cases the WFPC2 and ACS image are in the F606W filter (though in a few cases where images in this filter were not available we have used images in F555W or F775W). The STIS and F606W images can be thought of as broad "V" or visual images, and are, for galaxies exhibiting typical colors of GRB hosts, the single most sensitive settings for these cameras. F555W is close to the ground-based Johnson V-band, and F775W corresponds to the ground-based Johnson I-band. Due to the redshifts of the hosts, these images generally correspond to blue or ultra-violet images of the hosts in their rest frame, and thus detect light largely produced by the massive stars in the hosts.

Figure 2: A mosaic of cc SN host galaxies imaged with *HST* as part of the GOODS program. Each image in the mosaic has a width of $7''.5$ on the sky, and thus two times the field-of-view of each image in the GRB mosaic. The position of each SN on its host galaxy is marked. In all cases, these positions are known to sub-pixel accuracy. supernovae in the GOODS sample were identified by [30] as either Type Ia or cc supernovae based on their colors, luminosities and light curves, as data allowed (a SN going off near the beginning or end of one of the multi-epoch observing runs

would have much less data, and sometimes poor color information). Thus bright Type Ib and Ic supernovae, which have colors and luminosities similar to Type Ia supernovae, would have likely been classified as Type Ia (unless a grism spectrum was taken – however only a small fraction of objects were observed spectroscopically). On the other hand fainter Type Ib and Ic supernovae ($M_B \gtrsim -18$) could in principle be identified from photometric data; however, in practice the data were rarely sufficient for a clear separation from other cc supernovae. Based on surveys of nearby galaxies, one might expect approximately 20% of the cc supernovae to be Type Ib or Ic^{49,50}.

Figure 3: The locations of the explosions in comparison to the host light. For each object an arrow indicates the fraction of total host light in pixels fainter than or equal to the light in the pixel at the location of the transient. The cumulative fraction of GRBs or supernovae found at a given fraction of the total light is shown as a histogram. The blue arrows and histogram correspond to the GRBs and the red arrows and histogram correspond to the supernovae. Were the GRBs and supernovae to track the light identically, their histograms would follow the diagonal line. While the supernovae positions do follow the light within the statistical error, the GRBs are far more concentrated on the brightest regions of their hosts. Thus while the probability of a SN exploding in a particular pixel is roughly proportional to the surface brightness of the galaxy at that pixel, the probability of a GRB at a given location effectively goes as a higher power of the local surface brightness.

Figure 4: A comparison of the absolute magnitude and size distributions of the GRB and SN hosts. In the main panel, the cc SN hosts are represented as red squares and the LGRB hosts as blue

circles. The absolute magnitudes of the hosts are shown on the x-axis and the lengths of the semi-major axes of the hosts on the y-axis. The plot is then projected onto the two side panels where a histogram is displayed for each host population in each of the dimensions - absolute magnitude and semi-major axis. Shown as blue arrows are the absolute magnitudes of GRB hosts with $z < 1.2$ that have been detected from the ground but have not yet been observed by HST. These hosts are only included in the absolute magnitude histogram. The hosts of GRBs are both smaller and fainter than those of supernovae.

Acknowledgements Support for this research was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc. Observations analyzed in this work were taken by the NASA/ESA Hubble Space Telescope under programs: 7785, 7863, 7966, 8189, 8588, 9074 and 9405 (PI: Fruchter); 7964, 8688, 9180, and 10135 (PI S. R. Kulkarni); 8640 (PI: S. T. Holland). The authors wish to thank Nino Panagia, Nolan Walborn and Alicia Soderberg for informative conversations. We also thank Alex Filippenko and collaborators for early-time images of GRB 980326, and Josh Bloom and collaborators for making their early observations of GRB 020322 public.

Competing Interests The authors declare that they have no competing financial interests.

Correspondence Correspondence and requests for materials should be addressed to Andrew Fruchter (email: fruchter@stsci.edu).




Figure 1




Figure 2




Figure 3




Figure 4