

Long Time Behavior of Solutions to Nernst-Planck and Debye-Hückel Drift-Diffusion Systems

Piotr Biler and Jean Dolbeault

Abstract. We study the convergence rates of solutions to drift-diffusion systems (arising from plasma, semiconductors and electrolytes theories) to their self-similar or steady states. This analysis involves entropy-type Lyapunov functionals and logarithmic Sobolev inequalities.

1 Introduction

We consider the long time asymptotics of solutions to drift-diffusion systems

$$u_t = \nabla \cdot (\nabla u + u \nabla \phi), \quad (1.1)$$

$$v_t = \nabla \cdot (\nabla v - v \nabla \phi), \quad (1.2)$$

$$\Delta \phi = v - u, \quad (1.3)$$

where u, v denote densities of negatively, respectively positively, charged particles. The Poisson equation (1.3) defines the electric potential ϕ coupling the equations (1.1)-(1.2) for the temporal evolution of charge distributions. The system (1.1)-(1.3) was formulated by W. Nernst and M. Planck at the end of the nineteenth century as a basic model for electrodiffusion of ions in electrolytes filling the whole space \mathbb{R}^3 . Note that the case of multicharged particles is also covered by (1.1)-(1.3) since u and v denote the charge densities.

Supplemented with the no-flux boundary conditions

$$\frac{\partial u}{\partial \nu} + u \frac{\partial \phi}{\partial \nu} = 0, \quad (1.4)$$

$$\frac{\partial v}{\partial \nu} - v \frac{\partial \phi}{\partial \nu} = 0 \quad (1.5)$$

on the boundary of a bounded domain $\Omega \subset \mathbb{R}^d$, $d \leq 3$, and either

$$\phi = 0 \quad \text{on } \partial\Omega, \quad (1.6)$$

or

$$\phi = E_d * (v - u), \quad (1.7)$$

where E_d is the fundamental solution of the Laplacian in \mathbb{R}^d , the system (1.1)-(1.3) was also studied by P. Debye and E. Hückel in the 1920's. (1.6) signifies a

conducting boundary of the container, while in the case of a bounded domain the “free” boundary condition (1.7) corresponds to a container immersed in a medium with the same dielectric constant as the solute.

These equations, together with their generalizations including e.g. an exterior potential, known as drift-diffusion Poisson systems, also appear in plasma physics and (supplemented with some mixed linear boundary conditions instead of (1.4)-(1.5)) in semiconductor device modelling.

To determine completely the evolution, the initial conditions

$$u(x,0) = u_0(x), \quad v(x,0) = v_0(x) \quad (1.8)$$

are added. Obviously, positivity of $u_0 \geq 0, v_0 \geq 0$ is conserved: $u(x,t) \geq 0, v(x,t) \geq 0$, as well as the total charges

$$M_u = \int u_0(x) dx = \int u(x,t) dx, \quad M_v = \int v_0(x) dx = \int v(x,t) dx. \quad (1.9)$$

Here M_u, M_v are not necessarily the same, *i.e.* the electroneutrality condition

$$M_u = M_v \quad (1.10)$$

is not, in general, required. Condition (1.10) must be satisfied in the case of the homogeneous Neumann boundary conditions $\frac{\partial \phi}{\partial \nu} = 0$ (*i.e.* an isolated wall of the container) leading together with (1.4)-(1.5) to

$$\frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = \frac{\partial \phi}{\partial \nu} = 0.$$

Our results (Theorem 1.2 below) are valid in that case, with even a simpler proof.

The asymptotic properties of solutions to (1.1)-(1.3), (1.7) have been studied recently in [1]. The authors proved that (for $d \geq 3$, $M_u = M_v = 1$ and u_0 and v_0 regular enough) u, v tend to their self-similar asymptotic states at an algebraic rate. We improve these results by relaxing assumptions on the initial data and showing a stronger (still algebraic) decay rate, which we expect to be optimal (see Theorem 1.1 below).

In the case of a bounded domain, the convergence (with no specific speed) in the L^1 -norm of u and v solving (1.1)-(1.5) to their corresponding steady states has been proved in [5] (as well as the L^∞ -convergence for more regular u_0, v_0). Here we prove the exponential convergence towards the steady states with a decay rate depending on $\Omega \subset \mathbb{R}^d$, $d \geq 2$, and the initial value of the entropy functional only (see Theorem 1.2 below).

Notation. The L^p -norm in \mathbb{R}^d or $\Omega \subset \subset \mathbb{R}^d$ is denoted by $|\cdot|_p$, and inessential constants (which may vary from line to line) are denoted generically by C .

Define the asymptotic states in \mathbb{R}^d by

$$u_{as}(x, t) = \frac{M_u}{(2\pi(2t+1))^{d/2}} \exp\left(-\frac{|x|^2}{2(2t+1)}\right), \tag{1.11}$$

$$v_{as}(x, t) = \frac{M_v}{(2\pi(2t+1))^{d/2}} \exp\left(-\frac{|x|^2}{2(2t+1)}\right), \tag{1.12}$$

where the charges of the solution $\langle u, v \rangle$ of (1.1)-(1.2) are given by (1.9), and the entropy functional by

$$L(t) = \int u(x, t) \log\left(\frac{u(x, t)}{u_{as}(x, t)}\right) dx + \int v(x, t) \log\left(\frac{v(x, t)}{v_{as}(x, t)}\right) dx + \frac{1}{2} |\nabla\phi(t)|_2^2. \tag{1.13}$$

Theorem 1.1 *There exists a constant $C = C(d, M_u, M_v, L_0)$ such that for each solution $\langle u, v \rangle$ of (1.1)-(1.3), (1.7)-(1.8) in \mathbb{R}^d , $d \geq 3$, if $L(0) = L_0$, then for all $t \geq 0$,*

$$L(t) \leq C H(t) \tag{1.14}$$

and

$$|u(t) - u_{as}(t)|_1^2 + |v(t) - v_{as}(t)|_1^2 + |\nabla\phi(t)|_2^2 \leq C H(t), \tag{1.15}$$

where

$$H(t) = \begin{cases} (2t+1)^{-1/2}, & d=3, \\ (2t+1)^{-1} (\log(2t+1) + 1), & d=4, \\ (2t+1)^{-1}, & d>4. \end{cases}$$

Moreover if $M_u = M_v$, then $H(t) = (2t+1)^{-1}$ for any $d \geq 3$.

In the case of a bounded domain, define the entropy functional

$$\begin{aligned} W(t) &= \int u(x, t) \log u(x, t) dx - \int U(x) \log U(x) dx \\ &+ \int v(x, t) \log v(x, t) dx - \int V(x) \log V(x) dx \\ &+ \frac{1}{2} \int (u-v)\phi dx - \frac{1}{2} \int (U-V)\Phi dx, \end{aligned} \tag{1.16}$$

for the solution $\langle u, v, \phi \rangle$ of (1.1)-(1.5), (1.6) or (1.7), (1.8) and the unique steady state $\langle U, V, \Phi \rangle$ of the Debye-Hückel system with

$$M_u = \int U(x) dx, \quad M_v = \int V(x) dx. \tag{1.17}$$

Note that for the condition (1.6) the fifth and the sixth terms in $W(t)$ take the form $\frac{1}{2} |\nabla\phi|_2^2 - \frac{1}{2} |\nabla\Phi|_2^2$.

Theorem 1.2 *If $d \geq 2$, then there exist two constants $\lambda = \lambda(\Omega) > 0$ and $C = C(M_u, M_v, W_0)$ such that for each solution $\langle u, v, \phi \rangle$ of (1.1)-(1.6), (1.8) in a bounded uniformly convex domain Ω , if $W(0) = W_0$, then for all $t \geq 0$,*

$$W(t) \leq W(0) e^{-\lambda t}, \tag{1.18}$$

and

$$|u(t) - U|_1^2 + |v(t) - V|_1^2 + |\nabla(\phi - \Phi)|_2^2 \leq C e^{-\lambda t}. \tag{1.19}$$

2 Proof of Theorem 1.1

We begin with a rescaling of the system (1.1)-(1.3) which will lead to a system with a quadratic confinement potential, and therefore (eliminating the dispersion) to the expected exponential convergence to the steady states. This idea was applied in [8] and [7], as well as in [1], to a variety of problems ranging from kinetic equations to porous media equations.

Let $\bar{x} \in \mathbb{R}^d$, $\tau > 0$, be the new variables defined by

$$\bar{x} = \frac{x}{R(t)}, \quad \tau = \log R(t), \quad R(t) = (2t + 1)^{1/2}, \tag{2.1}$$

and consider the rescaled functions $\bar{u}, \bar{v}, \bar{\phi}$ such that

$$\begin{aligned} u(x, t) &= \frac{1}{R^d(t)} \bar{u}(\bar{x}, \tau), \\ v(x, t) &= \frac{1}{R^d(t)} \bar{v}(\bar{x}, \tau), \\ \phi(x, t) &= \bar{\phi}(\bar{x}, \tau). \end{aligned} \tag{2.2}$$

This whole section will deal with the rescaled system, so omitting the bars over x, u, v, ϕ will not lead to confusions with the original system, which now takes, after rescaling, the form

$$u_\tau = \nabla \cdot (\nabla u + ux + u \nabla \phi), \tag{2.3}$$

$$v_\tau = \nabla \cdot (\nabla v + vx - v \nabla \phi), \tag{2.4}$$

$$\Delta \phi = e^{-\tau(d-2)}(v - u). \tag{2.5}$$

The scaling (2.2) preserves the L^1 -norms, so the rescaled initial data u_0, v_0 still satisfy

$$M_u = \int u_0(x) dx = \int u(x, \tau) dx, \quad M_v = \int v_0(x) dx = \int v(x, \tau) dx. \tag{2.6}$$

Denote by $\langle u_\infty, v_\infty \rangle$ the steady state of (2.3)-(2.4), that is

$$u_\infty(x) = \frac{M_u}{(2\pi)^{d/2}} \exp\left(-\frac{|x|^2}{2}\right), \tag{2.7}$$

$$v_\infty(x) = \frac{M_v}{(2\pi)^{d/2}} \exp\left(-\frac{|x|^2}{2}\right). \tag{2.8}$$

Of course, going back to the original variables x, t , $\langle u_\infty, v_\infty \rangle$ corresponds to the asymptotic state $\langle u_{as}, v_{as} \rangle$ defined by (1.11)-(1.12). Writing $\phi = \beta\psi$ with $\beta = \beta(\tau) = e^{-\tau(d-2)} \rightarrow 0$ as $\tau \rightarrow +\infty$, we introduce the relative entropy

$$W(\tau) = \int u \log\left(\frac{u}{u_\infty}\right) dx + \int v \log\left(\frac{v}{v_\infty}\right) dx + \frac{\beta}{2} |\nabla\psi|_2^2 \tag{2.9}$$

corresponding to the original entropy functional L in (1.13). The evolution of W is given by

$$\frac{dW}{d\tau} = - \int u \left| \nabla \left(\log \frac{u}{U} \right) \right|^2 dx - \int v \left| \nabla \left(\log \frac{v}{V} \right) \right|^2 dx - \left(\frac{d}{2} - 1 \right) \beta |\nabla\psi|_2^2, \tag{2.10}$$

with U, V denoting the local Maxwellians

$$U(x, \tau) = M_u \frac{\exp\left(-\frac{1}{2}|x|^2 - \phi(x, \tau)\right)}{\int \exp\left(-\frac{1}{2}|y|^2 - \phi(y, \tau)\right) dy}, \tag{2.11}$$

$$V(x, \tau) = M_v \frac{\exp\left(-\frac{1}{2}|x|^2 + \phi(x, \tau)\right)}{\int \exp\left(-\frac{1}{2}|y|^2 + \phi(y, \tau)\right) dy}, \tag{2.12}$$

so that $\nabla U/U = -(x + \nabla\phi)$, $\nabla V/V = -(x - \nabla\phi)$. Using the notation

$$J = \frac{1}{2} \int u \left| \frac{\nabla u}{u} + x \right|^2 dx + \frac{1}{2} \int v \left| \frac{\nabla v}{v} + x \right|^2 dx, \tag{2.13}$$

(2.10) can be rewritten as

$$\begin{aligned} \frac{dW}{d\tau} &= -2J - 2 \int (\nabla u - \nabla v) \cdot \nabla\phi dx - 2 \int (u - v) x \cdot \nabla\phi dx \\ &\quad - \int (u + v) |\nabla\phi|^2 dx - \left(\frac{d}{2} - 1 \right) \beta |\nabla\psi|_2^2 \\ &= -2J - \beta^2 \int (u + v) |\nabla\psi|^2 dx - 2\beta |u - v|_2^2 + \left(\frac{d}{2} - 1 \right) \beta |\nabla\psi|_2^2. \end{aligned} \tag{2.14}$$

The quantity J in (2.13) can be estimated from below using the Gross logarithmic Sobolev inequality

$$\int f \log\left(\frac{f}{|f|_1}\right) dx + d \left(1 + \frac{1}{2} \log(2\pi a)\right) |f|_1 \leq \frac{a}{2} \int \frac{|\nabla f|^2}{f} dx \tag{2.15}$$

valid for each $a > 0$, see e.g. [11] or a thorough discussion of different versions of logarithmic Sobolev inequalities in [2]. (2.15) becomes an equality if and only if $f(x) = C \exp(-|x|^2/(2a))$ (up to a translation).

Taking $a = 1$ in (2.15), the relation (2.14) leads to

$$-\left(\frac{dW}{d\tau} + 2W\right) \geq 2\beta|u - v|_2^2 - \beta\frac{d}{2}|\nabla\psi|_2^2 \geq -C\beta(M_u + M_v)^2 \tag{2.16}$$

with a constant $C = C(d) = \frac{2}{d} \left(\frac{d-2}{4}\right)^{(d-2)/2} \Sigma^{d/2}$, because by the Hardy-Littlewood-Sobolev inequality and an interpolation

$$|\nabla\psi|_2^2 \leq \Sigma|u - v|_{2d/(d+2)}^2 \leq \Sigma|u - v|_1^{4/d}|u - v|_2^{2-4/d} \leq \frac{4}{d}|u - v|_2^2 + C|u - v|_1^2.$$

Clearly, (2.16) implies

$$\frac{d}{d\tau} (e^{2\tau}W(\tau)) \leq C(M_u + M_v)^2 e^{\tau(4-d)}$$

and, after one integration, we obtain

$$W(\tau) \leq \left(W(0)e^{-\tau} + C(M_u + M_v)^2\right)e^{-\tau} \tag{2.17}$$

in the case $d = 3$,

$$W(\tau) \leq \left(W(0) + C(M_u + M_v)^2\tau\right)e^{-2\tau} \tag{2.18}$$

if $d = 4$, and finally for all $d > 4$

$$W(\tau) \leq \left(W(0) + C(M_u + M_v)^2\right)e^{-2\tau}. \tag{2.19}$$

Since from the Csiszár-Kullback inequality (cf. (1.9) in [2], App. D in [7], [6] or [10]) $W(\tau)$ controls the L^1 -norm of $u - u_\infty$ and $v - v_\infty$, we get the same decay rates as in (2.17)-(2.19) for

$$|u(\tau) - u_\infty|_1^2 + |v(\tau) - v_\infty|_1^2 + \beta|\nabla\psi(\tau)|_2^2 \leq 2\left(\max(M_u, M_v) + 1\right)W(\tau). \tag{2.20}$$

Returning to the original variables x, t , this implies, of course, the estimates (1.14)-(1.15) of Theorem 1.1 in the general case.

In the electroneutrality case (1.10): $M_u = M_v$, since $u_\infty = v_\infty$, so for $d = 3$, $|u - v|_1^2 = |u - u_\infty + v_\infty - v|_1^2 \leq Ce^{-\tau}$. Next, a modification of (2.16) reads

$$\frac{d}{d\tau} \left(e^{2\tau}W(\tau)\right) \leq Ce^{2\tau}\beta|u - v|_1^2 \leq C,$$

and this leads to $W(\tau) \leq C(1 + \tau)e^{-2\tau}$. Inserting this into (2.20) and (2.16) once again implies

$$\frac{d}{d\tau} \left(e^{2\tau} W(\tau) \right) \leq C(1 + \tau)e^{-\tau},$$

so that $W(\tau) \leq Ce^{-2\tau}$. If $d=4$, the same reasoning once again applies providing also the same improved decay rate.

Remark. 2.1 *Note that the constant C in (1.15) depends on d, M_u, M_v and $L(0)$ only, and is independent of e.g. $|u_0|_r, |v_0|_r$ with some $r > d/2$ — as it was in fact in [1]. Conditions like $|u_0|_r + |v_0|_r < \infty$ are sufficient for (local in time) existence of solutions to the considered systems (cf. Theorem 2 in [5]), but they can be relaxed — as it was done for a related parabolic-elliptic system describing the gravitational interaction of particles in [4]. Thus, compared to [1], Theorem 1.1 gives not only an improvement of the exponents but also gets rid of the unnecessary dependence on quantities other than $L(0), M_u, M_v$. We do not know if the exponents in Theorem 1.1 are optimal, but such a conjecture is supported by the calculations in the proof of the following*

Proposition 2.2 *There exists a constant $\lambda > 0$ depending only on d with $\lambda \geq \lambda(d) = (d-2) \left(\sqrt{(d-1)^2 + 3} - (d-1) \right)$, such that*

$$W(\tau) \leq W(0) e^{-\lambda\tau} \tag{2.21}$$

and hence

$$L(t) \leq L(0) (2t+1)^{-\lambda/2}$$

for each solution $\langle u, v \rangle$ to the Nernst-Planck system.

Remark. 2.3 *The interest of this proposition is that the constants controlling the convergence of $W(t), L(t)$, and hence $|u - u_{as}|_1, |v - v_{as}|_1$ in (1.15), depend on the initial values of $W(0), L(0)$ only (and not on $|u|_1 = M_u, |v|_1 = M_v$, which are quantities not comparable with, say, $\int u \log u \, dx, \int v \log v \, dx$ in the whole \mathbb{R}^d space case). However, the exponent λ — which is evaluated explicitly — is not as good as the one in Theorem 1.1.*

Proof of Proposition 2.2. Using (2.9), (2.13), (2.14), we may write for any positive λ

$$\begin{aligned} -\left(\frac{dW}{d\tau} + \lambda W \right) &= \lambda \left(J - \int u \log \left(\frac{u}{u_\infty} \right) - \int v \log \left(\frac{v}{v_\infty} \right) \right) \\ &+ (2 - \lambda)J + B + 2E - \mu F, \end{aligned} \tag{2.22}$$

where

$$B = \beta^2 \int (u + v) |\nabla \psi|^2 \, dx,$$

$$\begin{aligned} E &= \beta |u - v|_2^2, \\ F &= \left(\frac{d}{2} - 1\right) \beta |\nabla \psi|_2^2, \\ \mu &= 1 + \frac{\lambda}{d-2}. \end{aligned}$$

Observe that if we define

$$G_1 = \int u \left(\frac{\nabla u}{u} + x \right) \cdot \nabla \phi \, dx, \quad G_2 = \int v \left(\frac{\nabla v}{v} + x \right) \cdot \nabla \phi \, dx,$$

then

$$G_1 - G_2 = \int \nabla(u - v) \cdot \nabla \phi \, dx + \int (u - v)(x \cdot \nabla \phi) \, dx = E - F.$$

Define now

$$\begin{aligned} f_1 &= \sqrt{2-\lambda} \cdot \sqrt{u} \left(\frac{\nabla u}{u} + x \right), & g_1 &= \sqrt{u} \nabla \phi, \\ f_2 &= \sqrt{2-\lambda} \cdot \sqrt{v} \left(\frac{\nabla v}{v} + x \right), & g_2 &= \sqrt{v} \nabla \phi, \\ a_1 &= |f_1|_2, & b_1 &= |g_1|_2, & a_2 &= |f_2|_2, & b_2 &= |g_2|_2. \end{aligned}$$

By the Cauchy-Schwarz inequality we have

$$\begin{aligned} (2-\lambda)^{1/2} |E - F| &= (2-\lambda)^{1/2} |G_1 - G_2| \\ &= \left| \int (f_1 g_1 - f_2 g_2) \, dx \right| \\ &\leq a_1 b_1 + a_2 b_2. \end{aligned}$$

But

$$0 \leq (a_1 b_2 - a_2 b_1)^2 = (a_1^2 + a_2^2)(b_1^2 + b_2^2) - (a_1 b_1 + a_2 b_2)^2,$$

$$\begin{aligned} a_1 b_1 + a_2 b_2 &\leq \sqrt{2} \sqrt{(a_1^2 + a_2^2)/2} \sqrt{b_1^2 + b_2^2} \\ &\leq \frac{1}{\sqrt{2}} \left(\frac{1}{2} (a_1^2 + a_2^2) + (b_1^2 + b_2^2) \right) \\ &= \frac{1}{\sqrt{2}} \left((2-\lambda)J + B \right), \end{aligned}$$

and thus

$$(2-\lambda)^{1/2} |E - F| \leq \frac{1}{\sqrt{2}} \left((2-\lambda)J + B \right).$$

Using (2.22) we get

$$\begin{aligned}
 -\left(\frac{dW}{d\tau} + \lambda W\right) &\geq \sqrt{2(2-\lambda)}|E-F| + 2E - \mu F \\
 &= F \cdot \left(\sqrt{2(2-\lambda)}|X-1| + 2X - \mu\right) \tag{2.23}
 \end{aligned}$$

with $X = E/F \geq 0$. For either $d \geq 4$ and $\lambda \leq 2$, or $d = 3$ and $\lambda \leq 1$, we have $\mu \leq 2$. The right hand side of (2.23) (positive for $X \geq \mu/2$) equals (for $X \leq \mu/2 \leq 1$)

$$\sqrt{2(2-\lambda)}(1-X) + 2X - \mu = \left(2 - \sqrt{2(2-\lambda)}\right)X + \sqrt{2(2-\lambda)} - \mu,$$

so that

$$\sqrt{2(2-\lambda)} \geq \mu \tag{2.24}$$

guarantees $\frac{dW}{d\tau} + \lambda W \leq 0$, which implies (2.21). The condition (2.24) is equivalent to $\lambda \leq \lambda(d)$. In particular, $\lambda(d)$ is an increasing function of d , $\lambda(3) = \sqrt{7} - 2 < 1$, $\lambda(4) = 4\sqrt{3} - 6$ and $\lim_{d \rightarrow +\infty} \lambda(d) = \frac{3}{2}$.

Remark. 2.4 *In the case of one species of particles, i.e. $v \equiv 0$ as was in [3] and [4], the result of Proposition 2.2 still holds.*

Finally, we remark that there is, in general, no hope to have $\lambda > 2$ in nontrivial cases. This can be inferred from the formula (2.22), where for each $\chi > 1$, $J - \chi \left(\int u \log\left(\frac{u}{u_\infty}\right) dx + \int v \log\left(\frac{v}{v_\infty}\right) dx\right)$ could be negative and dominate the other terms (for instance, in the limit $M_u, M_v \rightarrow 0^+$).

3 Proof of Theorem 1.2

First, we recall that steady states $\langle U, V, \Phi \rangle$ of (1.1)-(1.3) satisfy the relations

$$\nabla \cdot (e^{-\Phi} \nabla (e^\Phi U)) = 0, \quad \nabla \cdot (e^\Phi \nabla (e^{-\Phi} V)) = 0,$$

hence

$$U = M_u \frac{e^{-\Phi}}{\int e^{-\Phi} dx}, \quad V = M_v \frac{e^\Phi}{\int e^\Phi dx}. \tag{3.1}$$

Together with (1.3) this leads to the Poisson-Boltzmann equation

$$\Delta \Phi = M_v \frac{e^\Phi}{\int e^\Phi dx} - M_u \frac{e^{-\Phi}}{\int e^{-\Phi} dx}. \tag{3.2}$$

This equation, supplemented with the Dirichlet boundary condition (1.6) or the free condition (1.7), for every $M_u, M_v \geq 0$, has a unique (weak) solution Φ , see [9] or Proposition 2 in [5] (and this solution is classical whenever $\partial\Omega$ is of class $C^{1+\epsilon}$ for some $\epsilon > 0$).

The evolution of the Lyapunov functional defined by (1.16) in the case of the Dirichlet boundary condition (1.6) or in the case (1.7) is given by

$$\frac{dW}{dt} = - \int u |\nabla(\log u + \phi)|^2 dx - \int v |\nabla(\log v - \phi)|^2 dx, \tag{3.3}$$

cf. (35) in [5], where the above relation is obtained for weak solutions to the Debye-Hückel system.

Concerning the global in time existence of solutions to the Debye-Hückel system with nonlinear boundary conditions (1.4)-(1.5), we note that this was proved for $d=2$ only in Theorem 3 of [5]. Thus, in higher dimensions $d \geq 3$, we assume that $\langle u(t), v(t) \rangle$ exists for all $t \geq 0$. If equations (1.1)-(1.3) are supplemented with linear type boundary conditions (as it is the case in semiconductor modelling), the assumption $u_0, v_0 \in L^r(\Omega)$ with an exponent $r > d/2$ (cf. Theorem 2 (ii) in [5] and [1] for the case of the whole space \mathbb{R}^d) guarantees the existence of $\langle u(t), v(t) \rangle$ for all $t \geq 0$.

First, we represent the entropy production terms in (3.3) as

$$\int u |\nabla(\log(ue^\phi))|^2 dx = \int ue^\phi |\nabla(\log(ue^\phi))|^2 \frac{e^{-\phi}}{\int e^{-\phi} dx} dx \cdot \int e^{-\phi} dx, \tag{3.4}$$

with an obvious modification for the second term. Note that (1.4)-(1.5) read $\frac{\partial}{\partial \nu}(ue^\phi) = \frac{\partial}{\partial \nu}(ve^{-\phi}) = 0$. Then we recall Remark 3.7 of [2], where counterparts of the logarithmic Sobolev inequality (2.15) (or Poincaré-type inequalities) are discussed in the case of a bounded uniformly convex domain. We apply this remark to the domain Ω and the probability measure

$$\rho_0(x) = \frac{e^{-\phi}}{\int e^{-\phi} dx}$$

in the first entropy production term in (3.3) written as in (3.4). This implies the existence of a constant $C(\Omega) > 0$ such that

$$\int \Psi\left(\frac{f}{\int f d\rho_0}\right) d\rho_0 \leq C(\Omega) \int \Psi''\left(\frac{f}{\int f d\rho_0}\right) \frac{|\nabla f|^2}{(\int f d\rho_0)^2} dx,$$

where $\Psi(s) = 1 - s + s \log s$ and $f = ue^\phi$. Here we have

$$\int u |\nabla(\log(ue^\phi))|^2 dx = M_u \int \Psi''\left(\frac{f}{\int f d\rho_0}\right) \frac{|\nabla f|^2}{(\int f d\rho_0)^2} dx$$

since $\int f d\rho_0 = \int ue^\phi d\rho_0 = \frac{M_u}{\int e^{-\phi} dx}$. Thus we arrive at

$$\int u |\nabla(\log(ue^\phi))|^2 dx \geq \frac{M_u}{C(\Omega)} \int \left(\frac{f}{\int f d\rho_0} \log\left(\frac{f}{\int f d\rho_0}\right) + 1 - \frac{f}{\int f d\rho_0} \right) d\rho_0,$$

or

$$\int u |\nabla (\log(ue^\phi))|^2 dx \geq \frac{1}{C(\Omega)} \int u \log \left(\frac{ue^\phi}{\frac{M_u}{\int e^{-\phi} dx}} \right) dx. \tag{3.5}$$

Similarly, we have

$$\int v |\nabla (\log(v e^{-\phi}))|^2 dx \geq \frac{1}{C(\Omega)} \int v \log \left(\frac{v e^{-\phi}}{\frac{M_v}{\int e^\phi dx}} \right) dx. \tag{3.6}$$

Now we compute the expression

$$\delta = \int u \log \left(\frac{ue^\phi}{\frac{M_u}{\int e^{-\phi} dx}} \right) dx + \int v \log \left(\frac{v e^{-\phi}}{\frac{M_v}{\int e^\phi dx}} \right) dx.$$

If $\langle U, V, \Phi \rangle$ is the solution of the Poisson-Boltzmann equation (3.2) with the homogeneous Dirichlet boundary conditions for Φ , then it can be checked that

$$\delta = W + J[\phi] - J[\Phi], \tag{3.7}$$

where

$$\begin{aligned} W &= \int u \log u dx + \int v \log v dx + \frac{1}{2} \int |\nabla \phi|^2 dx \\ &\quad - \int U \log U dx - \int V \log V dx - \frac{1}{2} \int |\nabla \Phi|^2 dx \end{aligned}$$

is as in (1.16), and

$$J[\phi] = \frac{1}{2} \int |\nabla \phi|^2 dx + M_u \log \left(\int e^{-\phi} dx \right) + M_v \log \left(\int e^\phi dx \right)$$

is a strictly convex functional reaching its minimum at Φ .

Now it is clear from (3.3), (3.5)-(3.6) and (3.7) that for some $\lambda = \lambda(\Omega) > 0$ $\frac{dW}{dt} + \lambda W \leq 0$, i.e. $W(t)$ decays exponentially in t

$$W(t) \leq W(0) e^{-\lambda t}. \tag{3.8}$$

By the Csiszár-Kullback inequality (as was in Section 2), $W(t)$ controls the L^1 -convergence to the unique steady state, so the conclusion (1.19) of Theorem 1.2 follows from (3.8). This improves (34) in Theorem 6 of [5] in two ways. First, there is an exponential decay rate. Second, (34) is proved under the assumption $W(0) < \infty$, which is much weaker than the assumption on the L^2 -boundedness in time of the solution $\langle u, v \rangle$ in Theorem 6 of [5]. Evidently, this result is also valid for one species case (M_u or M_v equal to 0), so Theorem 2 in [3] is also improved.

Acknowledgements. This research was partially supported by the grants POLONIUM 98111 and KBN 324/P03/97/12. The second author thanks the program on Charged Particle Kinetics at the Erwin Schrödinger Institute and the TMR Kinetic Equations for partial support too.

References

- [1] A. Arnold, P. Markowich, G. Toscani, On large time asymptotics for drift-diffusion Poisson systems, preprint ESI no. 655 (1999), 1–11.
- [2] A. Arnold, P. Markowich, G. Toscani, A. Unterreiter, On logarithmic Sobolev inequalities, Csiszár-Kullback inequalities, and the rate of convergence to equilibrium for Fokker-Planck type equations, preprint TMR “Asymptotic Methods in Kinetic Theory” **12** (1998), 1–77.
- [3] P. Biler, Existence and asymptotics of solutions for a parabolic-elliptic system with nonlinear no-flux boundary conditions, *Nonlinear Analysis T.M.A.* **19** (1992), 1121–1136.
- [4] P. Biler, The Cauchy problem and self-similar solutions for a nonlinear parabolic equation, *Studia Math.* **114** (1995), 181–205.
- [5] P. Biler, W. Hebisch, T. Nadzieja, The Debye system: existence and large time behavior of solutions, *Nonlinear Analysis T.M.A.* **23** (1994), 1189–1209.
- [6] I. Csiszár, Information-type measures of difference of probability distributions and indirect observations, *Studia Sci. Math. Hungar.* **2** (1967), 299–318.
- [7] J. Dolbeault, M. del Pino, Generalized Sobolev inequalities and asymptotic behaviour in fast diffusion and porous medium problems, preprint Ceremade no. 9905 / ESI no. 704 (1999), 1–45.
- [8] J. Dolbeault, G. Rein, Time-dependent rescalings and Lyapunov functionals for the Vlasov-Poisson and Euler-Poisson systems, and for related models of kinetic equations, fluid dynamics and quantum physics, preprint Ceremade no. 9945 / TMR “Asymptotic Methods in Kinetic Theory” no. 102 (1999), 1–34, to appear in *Math. Mod. Appl. Sci.*
- [9] D. Gogny, P.-L. Lions, Sur les états d’équilibre pour les densités électroniques dans les plasmas, *RAIRO Modél. Math. Anal. Numér.* **23** (1989), 137–153.
- [10] S. Kullback, A lower bound for discrimination information in terms of variation, *IEEE Trans. Information Theory* **13** (1967), 126–127.
- [11] G. Toscani, Sur l’inégalité logarithmique de Sobolev, *C. R. Acad. Sci. Paris Sér. I Math.* **324** (1997), 689–694.

Piotr Biler
 Mathematical Institute
 University of Wrocław
 pl. Grunwaldzki 2/4
 50-384 Wrocław, Poland
 e-mail : Piotr.Biler@math.uni.wroc.pl

Jean Dolbeault
 Ceremade, U.M.R. C.N.R.S. no. 7534
 Université Paris IX-Dauphine
 Pl. du Marchal de Lattre de Tassigny
 F-75775 Paris Cedex 16, France
 e-mail : dolbeaul@ceremade.dauphine.fr

Communicated by J. Bellissard
 submitted 19/03/99 ; accepted 07/09/99