

## Low upper limits on the O<sub>2</sub> abundance from the Odin satellite<sup>★</sup>

L. Pagani<sup>1</sup>, A. O. H. Olofsson<sup>2</sup>, P. Bergman<sup>2</sup>, P. Bernath<sup>3</sup>, J. H. Black<sup>2</sup>, R. S. Booth<sup>2</sup>, V. Buat<sup>4</sup>, J. Crovisier<sup>12</sup>, C. L. Curry<sup>3</sup>, P. J. Encrenaz<sup>1</sup>, E. Falgarone<sup>5</sup>, P. A. Feldman<sup>6</sup>, M. Fich<sup>3</sup>, H. G. Floren<sup>7</sup>, U. Frisk<sup>8</sup>, M. Gerin<sup>5</sup>, E. M. Gregersen<sup>9</sup>, J. Harju<sup>10</sup>, T. Hasegawa<sup>11</sup>, Å. Hjalmarsen<sup>2</sup>, L. E. B. Johansson<sup>2</sup>, S. Kwok<sup>11</sup>, B. Larsson<sup>7</sup>, A. Lecacheux<sup>12</sup>, T. Liljeström<sup>13</sup>, M. Lindqvist<sup>2</sup>, R. Liseau<sup>7</sup>, K. Mattila<sup>10</sup>, G. F. Mitchell<sup>14</sup>, L. H. Nordh<sup>15</sup>, M. Olberg<sup>2</sup>, G. Olofsson<sup>7</sup>, I. Ristorcelli<sup>16</sup>, Aa. Sandqvist<sup>7</sup>, F. von Scheele<sup>8</sup>, G. Serra<sup>16</sup>, N. F. Tothill<sup>14</sup>, K. Volk<sup>11</sup>, T. Wiklind<sup>2</sup>, and C. D. Wilson<sup>9</sup>

<sup>1</sup> LERMA & FRE 2460 du CNRS, Observatoire de Paris, 61 Av. de l'Observatoire, 75140 Paris, France

<sup>2</sup> Onsala Space Observatory, 439 92 Onsala, Sweden

<sup>3</sup> Department of Physics, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada

<sup>4</sup> Laboratoire d'Astronomie Spatiale, BP 8, 13376 Marseille Cedex 12, France

<sup>5</sup> LERMA & FRE 2460 du CNRS, École Normale Supérieure, 24 rue Lhomond, 75005 Paris, France

<sup>6</sup> Herzberg Institute of Astrophysics, NRC of Canada, 5071 W. Saanich Rd, Victoria, BC V9E 2E7, Canada

<sup>7</sup> Stockholm Observatory, SCFAB, Roslagstullsbacken 21, 106 91 Stockholm, Sweden

<sup>8</sup> Swedish Space Corporation, PO Box 4207, 171 04 Solna, Sweden

<sup>9</sup> Department of Physics and Astronomy, McMaster University, Hamilton, Ontario L8S 4M1, Canada

<sup>10</sup> Observatory, PO Box 14, University of Helsinki, 00014 Helsinki, Finland

<sup>11</sup> Department of Physics and Astronomy, University of Calgary, Calgary, Alberta T2N 1N4, Canada

<sup>12</sup> LESIA, Observatoire de Paris, Section de Meudon, 5 place Jules Janssen, 92195 Meudon Cedex, France

<sup>13</sup> Metsähovi Radio Observatory, Helsinki University of Technology, Otakaari 5A, 02150 Espoo, Finland

<sup>14</sup> Department of Astronomy and Physics, Saint Mary's University, Halifax, Nova Scotia B3H 3C3, Canada

<sup>15</sup> Swedish National Space Board, PO Box 4006, 171 04 Solna, Sweden

<sup>16</sup> CESR, 9 avenue du Colonel Roche, BP 4346, 31029 Toulouse, France

Received 2 December 2002 / Accepted 25 January 2003

**Abstract.** For the first time, a search has been conducted in our Galaxy for the 119 GHz transition connecting to the ground state of O<sub>2</sub>, using the Odin satellite. Equipped with a sensitive 3 mm receiver ( $T_{\text{sys}}(\text{SSB}) = 600$  K), Odin has reached unprecedented upper limits on the abundance of O<sub>2</sub>, especially in cold dark clouds where the excited state levels involved in the 487 GHz transition are not expected to be significantly populated. Here we report upper limits for a dozen sources. In cold dark clouds we improve upon the published SWAS upper limits by more than an order of magnitude, reaching  $N(\text{O}_2)/N(\text{H}_2) \leq 10^{-7}$  in half of the sources. While standard chemical models are definitively ruled out by these new limits, our results are compatible with several recent studies that derive lower O<sub>2</sub> abundances. Goldsmith et al. (2002) recently reported a SWAS tentative detection of the 487 GHz transition of O<sub>2</sub> in an outflow wing towards  $\rho$  Oph A in a combination of 7 beams covering approximately  $10' \times 14'$ . In a brief (1.3 hour integration time) and partial covering of the SWAS region ( $\approx 65\%$  if we exclude their central position), we did not detect the corresponding 119 GHz line. Our 3 sigma upper limit on the O<sub>2</sub> column density is  $7.3 \times 10^{15} \text{ cm}^{-2}$ . We presently cannot exclude the possibility that the SWAS signal lies mostly outside of the 9' Odin beam and has escaped our sensitive detector.


**Key words.** radio lines: ISM – ISM: molecules – Galaxy: abundances

### 1. Introduction

Oxygen, the third most abundant cosmic element, is a key species in the chemistry of the interstellar medium (ISM). Our knowledge of its prevalence and distribution depends on the ability to measure the abundance of the main oxygen-bearing species: O, CO, CO<sub>2</sub>, OH, H<sub>2</sub>O, CH<sub>3</sub>OH and O<sub>2</sub>. The task is difficult due to telluric blocking of CO<sub>2</sub>, H<sub>2</sub>O and O<sub>2</sub>, in addition to important transitions of OH and O. Standard chemical

Send offprint requests to: L. Pagani,  
e-mail: laurent.pagani@obspm.fr


<sup>★</sup> Based on observations with Odin, a Swedish-led satellite project funded jointly by the Swedish National Space Board (SNSB), the Canadian Space Agency (CSA), the National Technology Agency of Finland (Tekes) and Centre National d'Études Spatiales (CNES). The Swedish Space Corporation was the industrial prime contractor and is operating Odin.



**Fig. 1.** Receiver stability. Over a period of 24 hours of integration, the noise keeps decreasing steadily.

models predicted that the abundance of O<sub>2</sub> should be comparable to that of CO inside well-shielded cloud cores (e.g., Herbst & Klamperer 1973; Graedel et al. 1982; Maréchal et al. 1997). Consequently, O<sub>2</sub> was considered an important cooling species for clouds (Goldsmith & Langer 1978). It has been actively searched for from the ground (redshifted 119 GHz transition, e.g. Liszt 1985, 1992, Combes et al. 1996; <sup>16</sup>O<sup>18</sup>O rare isotope transition at 1.3 mm, e.g. Goldsmith et al. 1985, Maréchal et al. 1997), from high above the troposphere by balloon (Olofsson et al. 1998), and most recently by satellite (SWAS, Goldsmith et al. 2000). None of these efforts has produced a convincing detection of O<sub>2</sub>. SWAS reported the lowest upper limits yet:  $N(\text{O}_2)/N(\text{C}^{18}\text{O}) \leq 1.5$  ( $3\sigma$ ) and  $N(\text{O}_2)/N(\text{H}_2) \leq 2.6 \times 10^{-7}$  ( $3\sigma$ ) in sources with star-forming activity and  $N(\text{O}_2)/N(\text{H}_2) \leq 3 \times 10^{-6}$  ( $3\sigma$ ) in cold dark clouds (TMC-1 and L183/L134N). More recently, the SWAS team reported a possible detection of the 487 GHz O<sub>2</sub> line in the  $\rho$  Oph A outflow (Goldsmith et al. 2002) with  $N(\text{O}_2)/N(\text{H}_2) \approx 10^{-5}$ . Despite its diminished importance as a cloud coolant, O<sub>2</sub> still attracts interest because it provides strong constraints on chemical models.

Odin, a 1.1 m spaceborne millimeter and submillimeter telescope (Frisk et al. 2003; Olberg et al. 2003), is equipped with two receivers having O<sub>2</sub> search capabilities: a cryogenically cooled 3 mm HEMT preamplifier (fixed-) tuned to the 118.750343 GHz ( $N_J: 1_1 - 1_0$ ) transition of O<sub>2</sub> for which the beam diameter is 9'; and a submm receiver tunable to the 487 GHz O<sub>2</sub> line with a beamsize of 2.4'. Since the 119 GHz line is easy to thermalize, stronger than any of the submm transitions at temperatures below 100 K (Maréchal et al. 1997), and has an upper level only 5.7 K above the ground state, it is probably the best line to search for, especially in cold dark clouds. At 10 K, the 119 GHz line is 20 times stronger than the 487 GHz line. With a 119 GHz system temperature ( $T_{\text{sys}}(\text{SSB}) = 600$  K) 8 times lower than that of SWAS, we gain 2 orders of magnitude in sensitivity in extended cold clouds and gain a factor of 40 even if the source is smaller than the SWAS beam.



**Fig. 2.** O<sub>2</sub> spectra with unscaled C<sup>18</sup>O spectra. The noise peaks at source velocity are  $1.5\sigma$  (L134N) and  $2.5\sigma$  (Orion A). The horizontal axis is LSR velocity in km s<sup>-1</sup>, the vertical axis,  $T_A$  in K (uncorrected for beam efficiency).


In this paper, we present the observations in Sect. 2, the data reduction in Sect. 3, and discuss briefly the intriguing results in Sect. 4.

## 2. Observations

The first useful Odin observations were obtained towards Orion on 2001-08-19 and all reported sources but TMC-1 were subsequently observed until 2001-10-04. At that time, the phase-lock system became unstable and was lost after 2001-10-14. The phase-lock was recovered no later than 2002-03-09. Data taken towards TMC-1 (2002-03-23 to 2002-03-25) were correctly phase-locked. The absolute pointing uncertainty is within 30'', which is negligible for our beam size of 9'. In some sources, we allowed offsets up to 2' to be included in the average, which still represents less than 25% error in pointing. The autocorrelator spectral sampling was 0.125 MHz ( $\delta v = 0.315$  km s<sup>-1</sup>) in a 100 MHz bandwidth for all sources except G0.26-0.01, for which we used 1 MHz ( $\delta v = 2.52$  km s<sup>-1</sup>) sampling (in a bandwidth of 800 MHz). The autocorrelator resolution is twice as wide as its sampling. In this Letter, we adopt  $\eta_{\text{MB}} = 0.9$ , as measured at 557 GHz towards Jupiter. Details of the calibration are presented by Olberg et al. (2003).

## 3. Data reduction

Our first concern was to establish the exact center frequency of the O<sub>2</sub> receiver band at all times and check its stability. The Local Oscillator phase-lock reference signal is not thermally controlled and is thus allowed to drift somewhat around its standard frequency. The receiver tuning is checked by measuring the telluric O<sub>2</sub> line position in the backend during the 40% of each orbit Odin spends pointing at Earth's atmosphere. The tuning was extremely stable from mid-August to the 4th



**Fig. 3.** The SWAS and Odin beams towards  $\rho$  Oph A. The dashed circle represents the SWAS central beam, not taken into account in their detection (Goldsmith et al. 2002).

of October 2001, with high and low resolution data requiring corrections of  $4.85 \pm 0.1$  MHz and  $-0.3$  to  $-0.1$  MHz, respectively. In March 2002, the high resolution data correction was 0.12 MHz.


The observing mode (“Dicke” switching against blank sky with a  $4.4^\circ$  FWHM reference beam, off-axis by  $20^\circ$ ) gives reasonably good baselines at high resolution but rather bad ones for large bandwidth. For the large-bandwidth G0.26-0.01 observation, the baselines were improved by subtracting a Dicke-switched off-source observation ( $+30'$  in declination, with no C<sup>18</sup>O emission). After long integrations in high resolution mode, a sinusoidal ripple of low amplitude (a few mK) could sometimes be seen and was removed with a sinusoidal baseline fit.

The stability of the receiver is clearly demonstrated in the decrease of the rms noise versus time shown in Fig. 1. Some of our observational results are displayed in Fig. 2.

## 4. Results and discussion

### 4.1. Analysis

Table 1 gives the list of sources with their final noise limit, the C<sup>18</sup>O ( $J: 1-0$ ) integrated intensity convolved in a  $9'$  Gaussian beam taken from literature, and the final O<sub>2</sub> relative abundance upper limit ( $3\sigma$ ). To derive O<sub>2</sub> column density upper limits, we assumed that the 119 GHz transition is thermalized (LTE), since the upper energy level is only 5.7 K and the spontaneous transition probability is  $A_{ul} = 4.46 \times 10^{-9} \text{ s}^{-1}$ , one order of magnitude lower than that of C<sup>18</sup>O (see Goldsmith et al. 2000). Though the upper state cannot for quantum mechanical reasons be directly collisionally excited (Bergman 1995), the line should be thermalized above densities  $\approx 10^3 \text{ cm}^{-3}$ , which is satisfied by all of the observed clouds. To estimate the O<sub>2</sub> integrated intensity upper limit, we took  $I \leq 3\sigma \sqrt{\delta\nu \Delta\nu}$  where  $\Delta\nu$  is the C<sup>18</sup>O linewidth. The C<sup>18</sup>O ( $J: 1-0$ ) line is rather easy to



**Fig. 4.** Smoothed  $\rho$  Oph A O<sub>2</sub> spectrum, with the signal expected at 119 GHz from the SWAS O<sub>2</sub> column density estimate superimposed. Axes are the same as in Fig. 2.

thermalize and is relatively insensitive to the density and temperature ranges spanned by the observed sources. For a  $9'$  beam size, depletion and saturation effects should be of limited importance. To derive  $N(\text{C}^{18}\text{O})$  we also assumed LTE and we took  $N(\text{C}^{18}\text{O})/N(\text{H}_2) = 1.7 \times 10^{-7}$  (Frerking et al. 1982) for local clouds and modulated this value as a function of Galactocentric distance following the <sup>16</sup>O/<sup>18</sup>O variation described by Wilson & Rood (1994). The C<sup>18</sup>O data are all in the  $T_R^*$  scale as defined by Kutner & Ulich (1981), which is the most appropriate for extended sources.

### 4.2. $\rho$ Oph A

In a recent paper (Goldsmith et al. 2002), the SWAS team reported a possible detection of 487 GHz O<sub>2</sub> in the red wing of the  $\rho$  Oph A outflow. Their tentative detection was made by adding 7 different SWAS positions surrounding, but excluding, the source itself (cf. our Fig. 3). Our larger beam, with a slightly different target position ( $\rho$  Oph A-VLA) covers about 65% of the SWAS-mapped region, not including their central beam. With the same assumptions as in Goldsmith et al. (2002), we find a  $3\sigma$  upper limit of  $N(\text{O}_2) \leq 6.7 \times 10^{15} \text{ cm}^{-2}$ , corresponding to  $X(\text{O}_2) \leq 4.4 \times 10^{-6}$ , a factor of 3 below their estimate ( $N(\text{O}_2) = 2.3 \times 10^{16} \text{ cm}^{-2}$ ,  $X(\text{O}_2) = 1.5 \times 10^{-5}$ ). Using their estimated O<sub>2</sub> column density, we have computed the expected Gaussian line at 119 GHz and overlaid it on our spectrum in Fig. 4 (the Gaussian line has been combined with our own noise spectrum to simulate real observing conditions). A clear signal should have been detected if O<sub>2</sub> is evenly distributed among the SWAS mapped region. Because the Odin beam does not completely cover that region, we cannot exclude the possibility that most of the emission lies outside our beam and thus escaped detection at 119 GHz. Further Odin observations are underway to settle this point.

**Table 1.** Derived upper limits on O<sub>2</sub> column density relative to H<sub>2</sub>. The 1 channel rms noise is given in the T<sub>A</sub> scale, but all computations have been done assuming a 90% coupling to the main beam. Resolution is 0.3 km<sup>-1</sup> except for G0.26-0.01 (2.5 km<sup>-1</sup>). 6.8(14) means 6.8 × 10<sup>14</sup>. The “Ref.” column provides references for the C<sup>18</sup>O data.

Source	$\alpha_{2000}$	$\delta_{2000}$	O <sub>2</sub> rms (K)	$\int T_r^{*}(\text{C}^{18}\text{O})dv$ (K km s <sup>-1</sup> )	$\Delta v(\text{C}^{18}\text{O})$ (km s <sup>-1</sup> )	T <sub>kin</sub> (K)	N(O <sub>2</sub> ) (cm <sup>-2</sup> )	N(C <sup>18</sup> O) (cm <sup>-2</sup> )	X(O <sub>2</sub> )	Ref.
TMC1 – NH <sub>3</sub>	4 <sup>h</sup> 41 <sup>m</sup> 16 <sup>s</sup>	25°48′44″	0.010	1.6	1.1	10	≤6.8(14)	1.5(15)	≤7.7(-8)	1
OrionA	5 <sup>h</sup> 35 <sup>m</sup> 14 <sup>s</sup> .36	-5°22′30″	0.006	2.7	3.8	50	≤1.9(15)	3.6(15)	≤8.9(-8)	2
NGC2071	5 <sup>h</sup> 47 <sup>m</sup> 4 <sup>s</sup> .1	0°21′42″.8	0.026	2.86	1.8	20	≤2.6(15)	2.8(15)	≤1.5(-7)	3
L134N – NH <sub>3</sub>	15 <sup>h</sup> 54 <sup>m</sup> 08 <sup>s</sup> .52	-2°52′48″	0.024	1.16	0.6	10	≤1.1(15)	1.1(15)	≤1.7(-7)	4
$\rho$ OphA	16 <sup>h</sup> 26 <sup>m</sup> 26 <sup>s</sup> .4	-24°24′30″.5	0.027	5.17	1.53	30	≤3.4(15)	6.2(15)	≤9.3(-8)	5
IRAS16293	16 <sup>h</sup> 32 <sup>m</sup> 22 <sup>s</sup> .8	-24°28′35″	0.017	2.07	0.94	30	≤1.7(15)	2.5(15)	≤1.2(-7)	5
NGC6334I	17 <sup>h</sup> 20 <sup>m</sup> 53 <sup>s</sup> .38	-35°47′1″.5	0.017	30. <sup>a</sup>	4.0	50	≤5.0(15)	<sup>d</sup>	≤7.1(-8) <sup>e</sup>	6–8
(G0.26 – 0.01) <sup>c</sup>	17 <sup>h</sup> 46 <sup>m</sup> 11 <sup>s</sup>	-28°42′39″	0.021	11.75	(164.0) <sup>c</sup>	20?	≤5.6(16)	1.2(16)	≤7.6(-7)	9, 10
M17SW	18 <sup>h</sup> 20 <sup>m</sup> 23 <sup>s</sup> .11	-16°12′47″.2	0.024	2.84	4.3	50	≤7.3(15)	3.0(15)	≤5.7(-7)	11
S68FIRS1	18 <sup>h</sup> 29 <sup>m</sup> 50 <sup>s</sup> .3	1°15′18″.6	0.016	2.63	1.4	25	≤1.6(15)	2.8(15)	≤9.7(-8)	12
G34.3 + 0.2	18 <sup>h</sup> 53 <sup>m</sup> 18 <sup>s</sup> .34	1°14′58″.4	0.021	<sup>b</sup>	6.0	30	≤5.2(15)	<sup>d</sup>	≤5.2(-8)	13–16

(a) Core only: 2′ × 3′; (b) No extended C<sup>18</sup>O measurements available; (c) *Line of sight towards* G0.26-0.01, containing many clouds over a large velocity range; (d) From <sup>13</sup>CO, N(H<sub>2</sub>) ≈ 10<sup>23</sup> cm<sup>-3</sup>; (e) ≤1.7 × 10<sup>-7</sup> C<sup>18</sup>O core only.

Refs.: (1) Onishi et al. (1996), (2) Dutrey et al. (1996), (3) Aoyama et al. (2001), (4) Pagani et al. (2001), (5) Tachihara et al. (2000), (6) Kraemer & Jackson (1999), (7) Schwartz et al. (1989), (8) Dickel et al. (1977), (9) Dahmen et al. (1997), (10) Dahmen et al. (1998), (11) Ando et al. (2002), (12) McMullin et al. (2000), (13) Churchwell et al. (1992), (14) Matthews et al. (1987), (15) Little et al. (1994), (16) Kim et al. (2001).

#### 4.3. The absence of O<sub>2</sub>

Odin has improved upon the SWAS upper limits by a factor of a few in star-forming regions, and by factors of 20 and 40 for L134N and TMC-1, respectively, yet still has not detected O<sub>2</sub>. It is now well established (since the work of Olofsson et al. 1998) that standard, steady state, chemical models are ruled out by this absence of O<sub>2</sub>. In the meantime, more sophisticated models have been developed which can accommodate O<sub>2</sub> abundances compatible with these new upper limits. Goldsmith et al. (2000) and Bergin et al. (2000) discuss those models which appeared up to the time of the SWAS observations. More recent work includes Casu et al. (2001), Spaan & van Dishoeck (2001), Viti et al. (2001), Roberts & Herbst (2002), and Willacy et al. (2002). All propose different reasons for O<sub>2</sub> to be rare, and much work remains to be done to test which model(s) best represent reality (and also predict the observed abundances of other molecules). The detection of O<sub>2</sub> in one or more sources, even at a low level, would greatly help in this task.

*Acknowledgements.* We thank E. Herbst and P. Goldsmith for sending us their manuscripts prior to publication. We also thank the NANTEN group, G. Dahmen, J. McMullin and J. Bally for kindly providing us with their C<sup>18</sup>O data. Generous financial support from the Research Councils and Space Agencies in Canada, Finland, France and Sweden is gratefully acknowledged.

#### References

- Ando, M., Nagata, T., Sato, S., et al. 2002, ApJ, 574, 187  
Aoyama, H., Mizuno, N., Yamamoto, H., et al. 2001, PASJ, 53, 1053  
Bergin, E. A., Melnick, G. J., Stauffer, J. R., et al. 2000, ApJ, 539, L129  
Bergman, P. 1995, ApJ, 445, L167  
Casu, S., Cecchi-Pestellini, C.-C., & Aiello, S. 2001, MNRAS, 325, 826  
Churchwell, E., Walmsley, C. M., & Wood, D. O. S. 1992, A&A, 253, 541  
Combes, F., Wiklind, T., & Nakai, N. 1996, A&A, 327, L17  
Dahmen, G., Hüttemeister, S., Wilson, T. L., et al. 1997, A&AS, 126, 197  
Dahmen, G., Hüttemeister, S., Wilson, T. L., & Mauersberger, R. 1998, A&A, 331, 959  
Dickel, H. R., Dickel, J. R., & Wilson, W. J. 1977, ApJ, 217, 56  
Dutrey, A., Duvert, G., Castets, A., et al. 1993, A&A, 270, 468  
Frerking, M. A., Langer, W. D., & Wilson, R. W. 1982, ApJ, 262, 590  
Frisk, U., Hagström, M., Ala-Laurinaho, J., et al. 2003, A&A, 402, L27  
Goldsmith, P. F., & Langer, W. D. 1978, ApJ, 222, 881  
Goldsmith, P. F., Snell, R. L., Erickson N. R., et al. 1985, ApJ, 289, 613  
Goldsmith, P. F., Melnick, G. J., Bergin, E. A., et al. 2000, ApJ, 539, L123  
Goldsmith, P. F., Li, D., Bergin, E. A., et al. 2002, ApJ, 576, 814  
Graedel, T. E., Langer, W. D., & Frerking, M. A. 1982, ApJS, 48, 321  
Herbst, E., & Klamperer, W. 1973, ApJ, 185, 505  
Kim, H.-D., Cho, S.-H., Chung, H.-S., et al. 2000, ApJS, 131, 483  
Kraemer, K. E., & Jackson, J. M. 1999, ApJS, 124, 439  
Kutner, M. L., & Ulich, B. L. 1981, ApJ, 250, 341

- Liszt, H. S. 1985, *ApJ*, 298, 281  
Liszt, H. S. 1992, *ApJ*, 386, 139  
Little, L. T., Gibb, A. G., Heaton, B. D., Ellison, B. N., & Claude, S. M. X. 1994, *MNRAS*, 271, 649  
Maréchal, P., Viala, Y.-P., & Benayoun, J. J. 1997a, *A&A*, 324, 221  
Maréchal, P., Pagani, L., Langer, W. D., & Castets, A. 1997b, *A&A*, 318, 252  
Matthews, N., Little, L. T., MacDonald, G. H., et al. 1987, *A&A*, 184, 284  
McMullin, J. P., Mundy, L. G., Blake, G. A., et al. 2000, *ApJ*, 536, 845  
Olberg, M., Frisk, U., Lecacheux, A., et al. 2003, *A&A*, 402, L35  
Olofsson, G., Pagani, L., Tauber, J., et al. 1998, *A&A*, 339, L81  
Onishi, T., Mizuno, A., Kawamura, A., Ogawa, H., & Fukui, Y. 1996, *ApJ*, 465, 815  
Pagani, L., Pardo, J. R., Fich, M., Motte, F., & Stepnik, B. 2001, in *Infrared and submillimeter Space Astronomy*, ed. M. Giard et al. (EDP Sciences), *EAS Publ. Ser.*, 4, 145  
Roberts, H., & Herbst, E. 2002, *A&A*, 395, 233  
Spaans, M., & van Dishoeck, E. F. 2001, *ApJ*, 548, L517  
Tachihara, K., Mizuno, A., & Fukui, Y. 2000, *ApJ*, 528, 817  
Schwartz, P. R., Snell, R. L., & Schloerb, F. P. 1989, *ApJ*, 336, 519  
Viti, S., Roueff, E., Hartquist, T. W., Pineau des Forêts, G., & Williams, D. A. 2001, *A&A*, 370, 557  
Willacy, K., Langer, W. D., & Allen, M. 2002, *ApJ*, 573, L119  
Wilson, T. L., & Rood, R. T. 1994, *ARA&A*, 32, 191