



OPEN

Lower regional gray matter volume in the absence of higher cortical amyloid burden in late-life depression

Akihiro Takamiya^{1,2,3,10}, Thomas Vande Castele^{1,2,10}, Michel Koole⁴, François-Laurent De Winter^{1,2}, Filip Bouckaert^{1,2}, Jan Van den Stock^{1,2}, Stefan Sunaert^{5,6}, Patrick Dupont^{7,8}, Rik Vandenberghe^{7,8,9}, Koen Van Laere⁴, Mathieu Vandenbulcke^{1,2}✉ & Louise Emsell^{1,2,5}

Late-life depression (LLD) is associated with a risk of developing Alzheimer's disease (AD). However, the role of AD-pathophysiology in LLD, and its association with clinical symptoms and cognitive function are elusive. In this study, one hundred subjects underwent amyloid positron emission tomography (PET) imaging with [¹⁸F]-flutemetamol and structural MRI: 48 severely depressed elderly subjects (age 74.1 ± 7.5 years, 33 female) and 52 age-/gender-matched healthy controls (72.4 ± 6.4 years, 37 female). The Geriatric Depression Scale (GDS) and Rey Auditory Verbal Learning Test (RAVLT) were used to assess the severity of depressive symptoms and episodic memory function respectively. Amyloid deposition was quantified using the standardized uptake value ratio. Whole-brain voxel-wise comparisons of amyloid deposition and gray matter volume (GMV) between LLD and controls were performed. Multivariate analysis of covariance was conducted to investigate the association of regional differences in amyloid deposition and GMV with clinical factors, including GDS and RAVLT. As a result, there were no significant group differences in amyloid deposition. In contrast, LLD showed significant lower GMV in the left temporal and parietal region. GMV reduction in the left temporal region was associated with episodic memory dysfunction, but not with depression severity. Regional GMV reduction was not associated with amyloid deposition. LLD is associated with lower GMV in regions that overlap with AD-pathophysiology, and which are associated with episodic memory function. The lack of corresponding associations with amyloid suggests that lower GMV driven by non-amyloid pathology may play a central role in the neurobiology of LLD presenting as a psychiatric disorder.

Trial registration: European Union Drug Regulating Authorities Clinical Trials identifier: EudraCT 2009-018064-95.

Late-life depression (LLD) is a common but disabling psychiatric disorder with a high risk of mortality¹. Given the global aging population, LLD has become a significant public health concern. LLD is associated with cognitive dysfunction, including episodic memory impairment^{2–4}, which may persist even after the depressive episode has been treated successfully^{5,6}. This and multiple lines of evidence associating LLD with an increased risk of Alzheimer's disease (AD)^{7,8} suggest shared neuropathophysiology between the two disorders. The main neurobiological markers in this context are increased cerebral amyloid deposition, tau deposition, and gray matter (GM) atrophy, particularly in the medial temporal lobe (MTL).

¹Neuropsychiatry, Department of Neurosciences, Leuven Brain Institute, KU Leuven, Leuven, Belgium. ²Geriatric Psychiatry, University Psychiatric Center KU Leuven, Leuven, Belgium. ³Department of Neuropsychiatry, Keio University School of Medicine, Tokyo, Japan. ⁴Nuclear Medicine and Molecular Imaging, Department of Imaging & Pathology, KU Leuven, Leuven, Belgium. ⁵Department of Imaging & Pathology, Translational MRI, KU Leuven, Leuven, Belgium. ⁶Department of Radiology, University Hospitals Leuven (UZ Leuven), Leuven, Belgium. ⁷Department of Neurosciences, Laboratory for Cognitive Neurology, Leuven Brain Institute, KU Leuven, Leuven, Belgium. ⁸Alzheimer Research Centre KU Leuven, Leuven Brain Institute, KU Leuven, Leuven, Belgium. ⁹Neurology Department, University Hospitals Leuven (UZ Leuven), Leuven, Belgium. ¹⁰These authors contributed equally: Akihiro Takamiya and Thomas Vande Castele. ✉email: mathieu.vandenbulcke@uzleuven.be

	Patients with LLD	Healthy controls	Statistical values	p-value
Number	48	52		
Age, years (mean, SD)	74.1 (7.5)	72.4 (6.4)	t = 1.2, df=98	0.21
Female, n (%)	33 (68.5%)	37 (71.2%)	$\chi^2 = 0.07$, df = 1	0.79
ApoE $\epsilon 4$ carrier, n (%)	12 (25.0%)	22 (42.3%)	$\chi^2 = 3.08$, df = 1	0.08
Psychotic depression, n (%)	16 (33.3%)			
Late-onset depression, n (%)	28 (58.3%)			
Geriatric Depression Scale (mean, SD)	23.1 (4.3)	3.0 (3.1)	t = 24.4, df = 80	<0.001
Episodic memory domain scores (mean, SD)	-3.8 (3.0)	0 (2.7)	t = 6.4, df = 90	<0.001

Table 1. Clinical characteristics of participants. *SD* standard deviation.

Previous research into the relationship between amyloid deposition and depression has yielded mixed results. For example, neuropathological studies have reported both significant associations of amyloid deposition with depressive symptoms or a history of depression in patients with dementia^{9,10}, and no associations^{11,12}. Longitudinal positron emission tomography (PET) imaging studies in cognitively normal subjects have consistently shown baseline amyloid burden to be associated with emerging mild depressive symptoms during follow-up^{13–15}, whereas others did not find any associations^{16–18}. Notably, a recent large multisite florbetapir-PET study from the ADNI consortium reported lower, not higher, amyloid accumulation in LLD with mild depressive symptoms¹⁹.

Both LLD and AD are associated with lower gray matter volume (GMV). In AD, GM atrophy is widespread with the most pronounced GM loss in the temporal and parietal cortex^{20,21}. Findings in LLD are inconsistent: lower GMV was reported in temporal^{22,23}, prefrontal^{22,24}, and parietal²³ regions, and in the hippocampus^{24,25}, thalamus, and putamen²⁴, whereas some studies did not find any significant GMV reduction^{26,27}. This inconsistency might be due to the heterogeneity of LLD. For example, late-onset depression (LOD) may be related to vascular or neurodegenerative changes²⁸, whereas early-onset depression (EOD) may be more familial with a genetic predisposition²⁹. In addition, psychotic depression, which is common in LLD³⁰, may be considered a distinct entity from non-psychotic depression³¹.

Studies that have assessed the relationship between GMV and amyloid deposition in LLD are scarce. In a previous study, we reported lower hippocampal volume in LLD and found no association with amyloid deposition in regions associated with AD nor with episodic memory scores³².

In sum, most neuropathological and neuroimaging studies so far have focused on the relationship between amyloid deposition and depressive symptoms or a history of depression in cognitively normal subjects and subjects who had already been diagnosed with MCI or AD. Although results from these studies provide some insight into the relationship between amyloid burden and depressive symptoms or with previous depressive episodes, they are less informative regarding the neurobiology of LLD as a psychiatric disorder. The distinction here is important, as depressive symptoms in healthy subjects or subjects with neurodegenerative disorders may have a different underlying neural substrate to the mood symptoms that present alongside the more severe and wide-ranging neuropsychological and psychomotor deficits that characterize LLD.

In this study, we extend our previous work by applying whole-brain voxel-wise analyses to identify regional differences in amyloid deposition and GMV between patients with LLD and healthy subjects. Additionally, we investigate how these regional differences relate to clinical factors, including episodic memory scores, the age at onset of depression, and the presence of psychosis.

Materials and methods

Participants. This study was approved by the ethical committee of the University Hospitals Leuven (protocol S52151) and registered as a clinical trial (EudraCT Number 2009-018064-95). All participants provided written informed consent prior to enrollment in the study in accordance with the Declaration of Helsinki. The cohort of this study is the same as the previous study³².

Late-life depression. The late-life depression group included forty-eight participants over 55 years of age with a diagnosis of major depressive disorder (MDD) according to the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV-TR) were recruited at the University Psychiatric Center KU Leuven, Belgium. The exclusion criteria were: primary referral for assessment of cognitive impairment; comorbid major psychiatric illness; previous or current alcohol or drug dependence; neurological illness, including dementia; illness or medication precluding cognitive testing; metal implants precluding MRI scanning and ECT in the past 6 months. The characteristics of the LLD group are summarized in Table 1. The proportion of the ApoE $\epsilon 4$ carriers was not different between LOD (21.4%) and EOD (35.0%) ($p = 0.34$), nor between psychotic depression (37.5%) and non-psychotic depression (21.9%) ($p = 0.31$).

Healthy controls. Fifty-two age- and gender-matched healthy comparison subjects were included, of which 46 were recruited from another [¹⁸F]-flutemetamol study at our center³³. At inclusion, these subjects had a Clinical Dementia Rating score of 0 and Mini-Mental State Examination (MMSE) score > 26 and were stratified for ApoE $\epsilon 4$ allele. By design almost half of this cohort (44.6%) was an ApoE $\epsilon 4$ carrier³³. However, the ApoE $\epsilon 4$ status of the stratified comparison subjects did not differ from those of the patient group or the non-stratified

comparison group, nor was there a difference in the proportion of amyloid positive participants (10 patients with LLD and 12 controls) defined by a global standardized uptake value ratio (SUVR) value > 1.38 derived from the mean uptake across five bilateral volumes of interest (lateral frontal, parietal, temporal cortex and anterior and posterior cingulate cortex)³².

Clinical assessment. The Geriatric Depression Scale (GDS)-30 and Rey Auditory Verbal Learning Test (RAVLT) were used to assess the severity of depressive symptoms and episodic memory, respectively. Raw scores of neuropsychological data were converted to standardized z scores based on the mean and standard deviation of the healthy control group. The episodic memory domain score was calculated by summing the z-scores of total recall score (the total number of correctly recalled during the five trials), delayed recall score, and recognition score (the number of correct hits minus the number of false positives)³⁴. This domain score was used to investigate the association between episodic memory and the imaging measures.

MRI acquisition and image processing. High-resolution 3D turbo field echo (3DTFE) T1-weighted images were acquired using a 3-T Philips Intera scanner with an 8-channel head coil (Best, the Netherlands) with parameters: TR = 9.6 s, TE = 4.6 ms, flip angle = 8 degrees, Slice thickness = 1.2 mm, in-plane voxel-size = $0.98 \times 0.98 \times 1.2 \text{ mm}^3$, 182 axial slices. Due to decommissioning of the 3 T Intera scanner, 19 controls were scanned with identical sequence parameters on a 3 T Philips Achieva scanner using a 32-channel head coil. To account for the effect of the different scanner on the analysis, we included scanner type as a covariate in the analysis, and we reanalyzed data after excluding these 19 subjects' data.

All T1-images were processed using the default pipeline of the Computational Anatomy Toolbox (CAT12, <http://dbm.neuro.uni-jena.de/cat/>), a toolbox for Statistical Parametric Mapping software (SPM12, <http://www.fil.ion.ucl.ac.uk/spm>). Prior to preprocessing, all data were manually aligned with the anterior commissure–posterior commissure (AC-PC) line. Preprocessing included bias-correction, segmentation into GM, white matter (WM), and cerebrospinal fluid (CSF), spatial normalization using the DARTEL algorithm, and modulation. A study-specific DARTEL template was created from all participants' data in this study. The pre-processed images were smoothed with a Gaussian kernel of 8-mm full-width at half maximum (FWHM). Total intracranial volume (TIV) was calculated using CAT12.

Amyloid imaging. [¹⁸F]-Flutemetamol PET imaging was acquired on a 16-slice Siemens Biograph PET/CT scanner (Siemens, Erlangen, Germany) at the University Hospitals Leuven, as described previously^{32,35}. The PET tracer was injected intravenously as a bolus in an antecubital vein. There were no differences in the radioactivity between LLD (mean 149.5 MBq, SD = 4.7, range = 139–159) and controls (mean 150.6 MBq, SD = 5.1, range = 139–164) ($p = 0.28$). Image acquisition started 90 min after tracer injection and lasted for 30 min in list mode. Prior to the PET scan, a low-dose computed tomography scan was performed for attenuation correction. Random and scatter corrections were also applied. Images were reconstructed using Ordered Subsets Expectation Maximization (4 iterations \times 16 subsets).

All PET frames were analyzed using an in-house Matlab® script (run on Matlab 2017b) and SPM12. All six PET frames were realigned to the first frame to correct for potential head motion, and then averaged to create a single PET image. Because both limited spatial resolution of the PET scanner and GM atrophy in our population may lead to a partial volume effect (PVE)³⁶, PET images were subsequently corrected for PVE using the modified method of Muller-Gartner et al.³⁷ as implemented in our in-house Matlab script. After coregistration of the individual PET images to their respective T1-weighted MRI images, PET images were normalized to Montreal Neurological Institute space by using the transformation parameters calculated during T1 normalization. SUVR images were calculated using cerebellar GM as reference region. The cerebellar GM volume of interest was derived from the Automated Anatomical Labeling (AAL) atlas and was masked with the normalized subject-specific segmented GM map. SUVR images were smoothed with an 8 mm FWHM Gaussian kernel.

Statistical analysis. Descriptive statistics were used to analyze the clinical data. Distributions of all variables were inspected using histograms, q-q plots, and Shapiro–Wilk tests. Whole-brain analyses were conducted using SPM12, and other analyses were conducted by using R version 3.4.3.

To investigate any differences in GMV and amyloid deposition between groups, whole-brain voxel-wise comparisons of GMV and [¹⁸F]-flutemetamol uptake between LLD and healthy controls were performed by using a “Two-sample t-test” design in SPM12, with age as a covariate. TIV and scanner type were included as additional covariates in the GMV analysis. ApoE $\epsilon 4$ carrier status was included as an additional covariate in the amyloid analysis. The significance threshold for the voxel-wise whole-brain analyses was set at a cluster-level family-wise error (FWE)-corrected $p < 0.05$ with an individual voxel threshold of $p = 0.001$. Only voxel values greater than 0.1 were analyzed using absolute threshold masking in the GMV analysis.

Mean voxel values within the clusters determined to be significantly different between groups were extracted and used in subsequent analyses. First, group comparisons of amyloid deposition in regions of significantly lower GMV in LLD were conducted by using an independent t test and a linear regression model controlling for age and ApoE $\epsilon 4$ carrier status. Bonferroni correction was conducted to control for multiple comparisons ($\alpha = 0.05/\text{the number of regions with significantly lower GMV in LLD}$).

Second, to investigate the associations between regional GMV differences based on the results in the whole-brain analysis and clinical factors in LLD, we applied four multivariate analysis of covariance (MANCOVA) (Pillai's trace) models. We used the MANCOVAs to avoid multiple comparisons and to account for any relationships among dependent variables. Dependent variables included the regional GMV differences detected in the whole-brain analysis. In each model, the independent variables were (1) episodic memory domain score,

(2) GDS, (3) late onset (> 55 years) or early onset of depression, and (4) the presence of psychosis. Age and TIV were included as covariates in all models. For these analyses, we set $p < 0.0125$ ($\alpha = 0.05/4$) as the statistically significant threshold. We also conducted the same sequential MANCOVAs including age and ApoE $\epsilon 4$ carrier status as covariates and using SUVR values from the regions of significantly lower GMV in LLD to investigate potential correlations between regional amyloid deposition and clinical factors.

Finally, a linear regression model including age, TIV and ApoE $\epsilon 4$ carrier status as covariates, was used to investigate the relationship between GMV and amyloid deposition in LLD. As a proxy for amyloid deposition, we used the mean voxel SUVR values within the clusters extracted from regions of significantly lower GMV in LLD identified in the whole-brain analysis. Bonferroni correction was conducted to control for multiple comparisons.

Results

We included 48 patients with LLD (mean age 74.1, SD = 7.5; 33 females; 28 late-onset depression; 16 psychotic depression), and 52 healthy controls (mean age 72.4, SD = 6.4). The mean GDS score was 23.1 (SD = 4.3). LLD showed significant lower episodic memory scores than healthy controls (means of episodic memory scores in LLD and HC are -3.84 (SE = 0.3) and 0 (SE = 0.5), $t = -6.4$, $df = 90$, $p < 0.001$). Episodic memory scores did not correlate with GDS scores (Spearman's $\rho = -0.08$ [95%CI: -0.21 to 0.36], $p = 0.63$) in LLD. Clinical characteristics of the participants are presented in Table 1.

Regional differences in amyloid deposition and GMV. Whole-brain voxel-wise analysis revealed that patients with LLD showed significantly lower GMV in the left posterior temporal region (mainly ventral and lateral cortex), left parietal region (mainly at the temporo-parietal junction), and right ventral occipital region compared to healthy controls (Fig. 1A–E; Table 2). Brain regions showing higher GMV in LLD than healthy controls were not detected. These findings were minimally dependent on template, smoothing extent, scanner effect or amyloid positivity status (Supplementary information). In contrast, no statistically significant group differences in amyloid deposition were detected in the whole-brain analysis. Moreover, SUVR values in the regions of statistically significant lower GMV in LLD did not differ between groups (Fig. 1F–H; Supplementary information). For comparison with previous studies, we also conducted additional analyses without PVE correction, and the findings were the same (Supplementary information).

Associations of clinical factors with GMV/amyloid deposition in LLD. There was a significant main effect of episodic memory scores on regional GMV differences ($V = 0.29$, $F_{3,34} = 4.6$, $p = 0.008$). Separate ANCOVAs revealed that episodic memory scores were significantly associated with GMV in the left temporal region ($F_{1,36} = 12.3$, $p = 0.001$). The association between episodic memory and GMV did not survive after multiple comparison correction in the right occipital region ($F_{1,36} = 6.5$, $p = 0.015$), and there was no association with GMV in the left parietal region ($F_{1,36} = 0.96$, $p = 0.33$). There were no main effects of GDS ($V = 0.01$, $F_{3,33} = 0.15$, $p = 0.93$), age at onset ($V = 0.005$, $F_{3,42} = 0.10$, $p = 0.97$), or the presence of psychosis ($V = 0.02$, $F_{3,42} = 0.30$, $p = 0.82$). To illustrate the results, scatter plots of clinical factors and GMV in the left temporal region are presented in Fig. 2. There were no significant main effects of any clinical factors on regional amyloid deposition (Supplementary information). As an additional analysis, an association between the executive function domain score, GMV and amyloid deposition in LLD was also investigated; however, no significant associations were identified (Supplementary information).

Associations between GMV and amyloid deposition. Within the LLD cohort, there were no associations between GMV and SUVR in the left temporal ($p = 0.44$), left parietal ($p = 0.10$) and right occipital regions ($p = 0.76$) (Supplementary Fig. 1). In addition to these local associations between GMV and amyloid deposition, we also investigated the effect of amyloid deposition on different brain regions using a global measure of amyloid deposition; however, no associations were detected in the whole-brain analysis (Supplementary information).

Discussion

In this study we aimed to extend the literature on the relationship between LLD and AD-pathophysiology by applying multimodal whole-brain assessments comparing patients with LLD to healthy controls. We found that LLD was associated with lower regional GMV but not with differences in amyloid deposition compared to healthy controls. Additionally, lower left temporal lobe GMV was associated with episodic memory impairment in LLD.

AD-pathophysiology in LLD. Amyloid deposition is considered to be one of the earliest neuropathological hallmarks in AD³⁸, along with tau deposition, which triggers downstream effects, such as gray matter atrophy and cognitive impairment. Although amyloid pathology is hypothesized to be also related to the neurobiology of LLD³⁹, we did not find higher amyloid deposition in LLD compared to healthy controls in the whole brain analysis. Surprisingly, recent work¹⁹ reported lower amyloid deposition in LLD. The discrepancy could be due to considerable differences in methodology. For example, Mackin et al.¹⁹ analyzed data from multiple sites (33 imaging sites with 9 different scanner types) which could introduce methodological variability that is less readily controlled compared to a monocentric study. Additionally, the severity of depressive symptoms between our cohorts was significantly different (mean GDS of 7.3 in their study vs. 23.1 in our study: mild vs. severe depressive symptoms), and there were important differences in the quantification methods of amyloid burden (different PET tracers and the use of partial volume correction in our data). Another recent work, including 21 moderately depressed elders with limited or no prior antidepressant treatment history, reported increased amyloid deposition in the left parietal cortex in LLD⁴⁰ and no differences in GMV, which is in contrast with




Figure 1. Brain regions showing significantly lower GMV in patients with late-life depression compared to healthy controls following whole-brain voxel-wise analysis. Three regions were identified: the left temporal, left parietal and right occipital regions. Areas associated with lower GMV in LLD are shown in yellow in (A) the left lateral view and (B) the bottom view. Boxplots of GMVs and SUVRs in each region are also presented: GMV in (C) the left temporal, (D) the left parietal, and (E) the right occipital regions; SUVR in (F) the left temporal region, (G) the left parietal region, and (H) the right occipital region. GMVs in these regions were significantly lower in LLD than in healthy controls (C–E), whereas SUVRs in these regions did not differ between groups (F–H). Y-axis represents mean voxel values in each cluster. *HC* healthy control, *LLD* late-life depression.




Figure 1. (continued)

Cluster size (voxels)	Brain Regions	Peak MNI coordinates			T values	Cluster-level p (FWE-corrected)	Voxel-level p (FWE-corrected)
		x	y	z			
3295	Left temporal cluster					< 0.001	
	Left middle temporal gyrus	-64	-50	6	5.51		0.004
		-60	-57	-4	4.58		0.12
	Left inferior temporal gyrus	-50	-62	-8	4.81		0.055
		-38	-46	-14	4.14		0.40
		-51	-58	-22	3.96		0.57
		-50	-51	-8	3.77		0.77
		-46	-50	-27	3.39		0.98
1399	Left fusiform gyrus	-22	-44	-14	4.61		0.11
		-34	-62	-9	3.99		0.54
	Left parietal cluster					0.005	
	Left middle temporal gyrus	-51	-50	22	5.42		0.006
1564	Left supramarginal gyrus	-56	-46	32	4.53		0.14
		-58	-57	28	3.62		0.89
	Right occipital cluster					0.003	
	Right lingual gyrus	14	-86	-10	4.37		0.22
	Right middle occipital gyrus	32	-100	-2	4.09		0.44
	Right inferior occipital gyrus	24	-93	-6	3.99		0.54
		30	-86	-10	3.79		0.75
	44	-75	-14	3.77		0.77	
	Right fusiform gyrus	30	-75	-16	3.91		0.63
		46	-58	-20	3.86		0.68

Table 2. Regions of significantly lower GMV in LLD detected in the whole-brain GMV analysis. Peak coordinates were local maxima more than 10 mm apart.

our findings. It is possible that these subjects represent a different clinical phenotype to ours given their lack of prior medication use and preserved cognitive functioning relative to controls. Whilst the inconsistent findings in group differences in amyloid burden in LLD challenges the notion that amyloid pathology plays a central role in the disorder, it does not necessarily exclude a neurobiological link between LLD and AD.

Although hippocampal volume reduction has been the main focus in LLD and AD literature, GMV reduction beyond hippocampus has also been reported^{22,24}. Our findings of GMV decrease in LLD are in line with other studies, which reported GMV reduction in the left temporoparietal region^{23,41} and occipital region⁴². Temporoparietal and to a lesser degree occipital atrophy are also reported in AD literature^{20,43–45}. Therefore, the current




Figure 2. Associations of clinical factors with regional GMV decrease in the left temporal region. Episodic memory scores correlated with lower GMV in the left temporal region. To illustrate the results of the MANCOVAs, scatter plots of (A) episodic memory scores and (B) GDS, and boxplots of (C) onset age and (D) the presence of psychosis are presented. Episodic memory scores (A) had a significant main effect on regional GMV difference, whereas other clinical factors (B–D) did not have significant main effect. In all figures, data from healthy controls are also overlaid for reference. Y-axis in (C,D) represents mean voxel values in the left temporal cluster. *EOD* early-onset depression, *GDS* geriatric depression scale, *GMV* gray matter volume, *HC* healthy control, *LLD* late-life depression, *LOD* late-onset depression, *NPMD* non-psychotic major depression, *PMD* psychotic major depression.

results of GMV reduction in LLD partly overlap with atrophic patterns in AD. In the current whole-brain analysis using the stringent cluster-level FWE correction, the left hippocampal volume reduction in LLD we reported in our previous ROI analysis³² was not detected. However, a supplementary ROI analysis confirmed our previous finding (Supplementary information). We did not find any differences in frontal GMV between LLD and healthy controls. Some, but not all studies using whole-brain voxel-wise analysis found frontal GMV reduction²². This discrepancy may be due to a difference in clinical characteristics in study participants regarding the severity of depressive symptoms and the rate of psychotic patients (i.e., our cohort includes more severe psychotic patients than previous studies).

The etiology of GMV reduction in LLD. Stress has long been considered to play a major role in depression^{46,47} and stress-related neurotoxic effects, such as elevated cortisol levels, excess oxidative stress, and reduced neurotrophic factors, could contribute to GMV reduction⁴⁸. If stress-mediated effects on the brain were the main explanation of GMV reduction, one would expect that patients with EOD would show lower GMV than those with LOD⁴⁹. However, we did not find any associations between regional GMV and age at onset of depression. Another possible explanation is that lower GMV reduction arises from accelerated aging^{50,51}. Notably, consistent with a previous study²³, we found a negative correlation between age and GMV in our healthy cohort, which was not present in LLD i.e. the GMV differences were most pronounced between the younger

patients compared to the younger controls and these differences diminished with increasing age (Supplementary information).

Our findings of GMV reduction unrelated to amyloid pathology may also be related to a condition called suspected non-Alzheimer's disease or non-amyloid pathophysiology (SNAP)^{52,53}, which is related to an increased risk of cognitive decline and progression to dementia⁵³, and is estimated to affect almost a quarter of adults over 65 years of age⁵³. Moreover, tau deposition in the MTL in the absence of cortical amyloid deposition was postulated as primary age-related tauopathy (PART)⁵⁴. Tau deposition has been associated with temporal and parietal cortical atrophy in cognitively normal subjects⁵⁵ and left hippocampal atrophy in subjects with PART⁵⁶. Depressive symptoms have also been related to tau accumulation in the entorhinal cortex and inferior temporal lobe in cognitively normal older adults using tau [¹⁸F] AV-1451 PET imaging⁵⁷. A recent meta-analysis found limited evidence for tau pathology in LLD, but concluded it could not be ruled out given the lack of available data⁵⁸. As others have hypothesized, it is plausible that a combination of neurobiological processes such as proteinopathies, vascular pathology and stress-mediated damage interact to lower the threshold for the development of neuropsychiatric symptoms in vulnerable individuals⁵⁹. Conversely, in some individuals, reactive depressive symptoms may develop due to environmental stressors and contribute to transient cognitive dysfunction unrelated to neuropathology. Further research into non-amyloid pathology in LLD, and its association with stress, age at onset of depression, and GMV reduction may help to elucidate the underlying neurobiology of LLD.

Episodic memory dysfunction in LLD. Episodic memory dysfunction is considered as the first clinical sign of AD²¹, and it is also reported in LLD^{2–4}. We found episodic memory dysfunction in LLD, and this was associated with lower GMV in the left temporal region, but not with amyloid burden. Our results are supported by previous studies, showing the associations of episodic memory dysfunction with reduced hippocampal or parahippocampal volume^{34,60}. The identified temporal region in the current study included lateral temporal regions as well as the parahippocampal gyrus. Interestingly, recent whole-brain functional imaging studies have revealed that large-scale distributed networks, including lateral temporal or temporoparietal regions, were associated with episodic memory^{61–63}. Notably, the auditory verbal learning task used in our study involves various cognitive processes, including semantic processing and working memory⁶⁴. Auditory verbal short-term memory has been associated with activations in left-lateralized posterior superior temporal regions and inferior parietal/supramarginal gyrus⁶⁵ as well as with left lateral temporal and temporoparietal regions in patients with AD⁶⁴.

We found that regional GMV differences in LLD correlated with impairments in episodic memory but not with depression severity, and that episodic memory scores did not correlate with depression severity. This is in contrast to a previous meta-analysis of depression at all ages which found an association between depression severity and cognitive function⁶⁶. However, our negative finding is consistent with previous studies which focused specifically on LLD^{2,34}, suggesting that the effect of depression severity on cognitive function may be age-dependent. Moreover, our results suggest that the neural correlates mediating affective symptoms and cognitive symptoms in LLD might be different.

Strengths and limitations. It is important to emphasize the uniqueness of our cohort when comparing our analysis with previous studies. Our main focus was to explore the contribution of AD-associated pathophysiology to the neurobiology of LLD. Therefore, we recruited actively depressed patients without a diagnosis of MCI or dementia. In contrast, most previous amyloid-PET studies have focused on a history of depression or depressive symptoms in healthy subjects^{13–15} or subjects who were already diagnosed with MCI or AD at the time of scanning^{67,68}. However, there is a possibility that the underlying neurobiology of a psychiatric disorder (e.g. LLD) with/without cognitive dysfunction and a neurodegenerative disorder (e.g. AD) with depressive symptoms are different. In this context, Mackin et al.¹⁹ and our previous study³² have investigated LLD patients with mild or severe depressive symptoms respectively, and neither study found higher amyloid to be associated with LLD, whilst Smith et al.⁴⁰ reported higher amyloid in LLD with moderate depressive symptoms without prior antidepressant treatments in a smaller cohort of 21 patients and 21 controls.

One potential limitation of our study is the cross-sectional design, therefore there is still a possibility that amyloid deposition is related to longitudinal GM atrophy, onset of depression, and cognitive dysfunction. Second, most participants took antidepressant medications, which might have an effect on amyloid accumulation⁶⁹, although no association between the length of lifetime antidepressant treatments and amyloid burden in LLD was reported recently¹⁹. Third, whilst we attempted to mitigate any bias due to including data from 19 control subjects scanned on a different scanner, in the absence of formal test data from a subject scanned on both scanners to confirm this, we cannot rule out such effects. Fourth, we did not find any differences in GMV or amyloid deposition between psychotic depression and non-psychotic depression, nor those between EOD and LOD. These negative findings might be due to the limited sample size in each group. In addition, the negative finding related to amyloid deposition might be a consequence of the heterogeneity of our cohort and therefore may not be generalized to all LLD as a whole. Future investigation of a more homogenous LLD sample could provide further insights into the role of amyloid in specific LLD subtypes.

In conclusion, LLD was associated with lower GMV in regions also implicated in AD. Left temporal GMV reduction was associated with episodic memory dysfunction in LLD. However, amyloid deposition did not differ between LLD and healthy subjects, and it did not correlate with clinical symptoms nor clinical characteristics of LLD. Our results suggest that amyloid deposition might not play a central or proximal role in the neurobiology of LLD.

Data availability

The datasets generated during and/or analyzed during the current study are not publicly available.

Received: 11 February 2021; Accepted: 21 June 2021

Published online: 05 August 2021

References

1. Jeuring, H. W. *et al.* A six-year prospective study of the prognosis and predictors in patients with late-life depression. *Am. J. Geriatr. Psychiatry* **26**, 985–997 (2018).
2. Sheline, Y. I. *et al.* Cognitive function in late life depression: Relationships to depression severity, cerebrovascular risk factors and processing speed. *Biol. Psychiatry* **60**, 58–65 (2006).
3. Koenig, A. M. *et al.* Neuropsychological functioning in the acute and remitted states of late-life depression. *J. Alzheimer's Dis.* **45**, 175–185 (2015).
4. Riddle, M. *et al.* Longitudinal cognitive outcomes of clinical phenotypes of late-life depression. *Am. J. Geriatr. Psychiatry* **25**, 1123–1134 (2017).
5. Nebes, R. D. *et al.* Persistence of cognitive impairment in geriatric patients following antidepressant treatment: A randomized, double-blind clinical trial with nortriptyline and paroxetine. *J. Psychiatr. Res.* **37**, 99–108 (2003).
6. O'Brien, J. T., Lloyd, A., McKeith, I., Gholkar, A. & Ferrier, N. A longitudinal study of hippocampal volume, cortisol levels, and cognition in older depressed subjects. *Am. J. Psychiatry* **161**, 2081–2090 (2004).
7. Da Silva, J., Gonçalves-Pereira, M., Xavier, M. & Mukaetova-Ladinska, E. B. Affective disorders and risk of developing dementia: Systematic review. *Br. J. Psychiatry* **202**, 177–186 (2013).
8. Diniz, B. S., Butters, M. A., Albert, S. M., Dew, M. A. & Reynolds, C. F. Late-life depression and risk of vascular dementia and Alzheimer's disease: Systematic review and meta-analysis of community-based cohort studies. *Br. J. Psychiatry* **202**, 329–335 (2013).
9. Sweet, R. A. *et al.* Neuropathologic correlates of late-onset major depression. *Neuropsychopharmacology* **29**, 2242–2250 (2004).
10. Rapp, M. A. *et al.* Increased neurofibrillary tangles in patients with Alzheimer disease with comorbid depression. *Am. J. Geriatr. Psychiatry* **16**, 168–174 (2008).
11. Tsopelas, C. *et al.* Neuropathological correlates of late-life depression in older people. *Br. J. Psychiatry* **198**, 109–114 (2011).
12. Wilson, R. S., Capuano, A. W. & Boyle, P. A. Clinical-pathologic study of depressive symptoms and cognitive decline in old age. *Neurology* **83**, 702–709 (2014).
13. Babulal, G. M. *et al.* Mood changes in cognitively normal older adults are linked to Alzheimer disease biomarker levels. *Am. J. Geriatr. Psychiatry* **24**, 1095–1104 (2016).
14. Harrington, K. D. *et al.* Amyloid burden and incident depressive symptoms in cognitively normal older adults. *Int. J. Geriatr. Psychiatry* **32**, 455–463 (2017).
15. Donovan, N. J. *et al.* Longitudinal association of amyloid beta and anxious-depressive symptoms in cognitively normal older adults. *Am. J. Psychiatry* **175**, 530–537 (2018).
16. Madsen, K. *et al.* Lack of association between prior depressive episodes and cerebral [¹¹C] PiB binding. *Neurobiol. Aging* **33**, 2334–2342 (2012).
17. Tateno, A. *et al.* Amyloid imaging with [¹⁸F] florbetapir in geriatric depression: Early-onset versus late-onset. *Int. J. Geriatr. Psychiatry* **30**, 720–728 (2015).
18. Wu, K. Y. *et al.* Diversity of neurodegenerative pathophysiology in nondemented patients with major depressive disorder: Evidence of cerebral amyloidosis and hippocampal atrophy. *Brain Behav.* **8**, 1–10 (2018).
19. Mackin, R. S. *et al.* Late life depression is associated with reduced cortical amyloid burden: Findings from the ADNI Depression Project. *Biol. Psychiatry* **89**, 757–765. <https://doi.org/10.1016/j.biopsych.2020.06.017> (2021).
20. Jack, C. R. & Holtzman, D. M. Biomarker modeling of Alzheimer's disease. *Neuron* **80**, 1347–1358 (2013).
21. Jagust, W. Imaging the evolution and pathophysiology of Alzheimer disease. *Nat. Rev. Neurosci.* **19**, 687–700 (2018).
22. Du, M. *et al.* Brain grey matter volume alterations in late-life depression. *J. Psychiatry Neurosci.* **39**, 397–406 (2014).
23. Lebedeva, A. *et al.* Neuroanatomical correlates of late-life depression and associated cognitive changes. *Neurobiol. Aging* **36**, 3090–3099 (2015).
24. Sexton, C. E., MacKay, C. E. & Ebmeier, K. P. A systematic review and meta-analysis of magnetic resonance imaging studies in late-life depression. *Am. J. Geriatr. Psychiatry* **21**, 184–195 (2013).
25. Geerlings, M. I. & Gerritsen, L. Late-life depression, hippocampal volumes, and hypothalamic-pituitary-adrenal axis regulation: A systematic review and meta-analysis. *Biol. Psychiatry* **82**, 339–350 (2017).
26. Colloby, S. J. *et al.* Cortical thickness and VBM-DARTEL in late-life depression. *J. Affect. Disord.* **133**, 158–164 (2011).
27. Sexton, C. E. *et al.* Magnetic resonance imaging in late-life depression: multimodal examination of network disruption. *Arch. Gen. Psychiatry* **69**, 680–689 (2016).
28. Naismith, S. L., Norrie, L. M., Mowszowski, L. & Hickie, I. B. The neurobiology of depression in later-life: Clinical, neuropsychological, neuroimaging and pathophysiological features. *Prog. Neurobiol.* **98**, 99–143 (2012).
29. Baldwin, R. C. & Tomenson, B. Depression in later life. A comparison of symptoms and risk factors in early and late onset cases. *Br. J. Psychiatry* **167**, 649–652 (1995).
30. Brodaty, H. *et al.* Increased rate of psychosis and psychomotor change in depression with age. *Psychol. Med.* **27**, 1205–1213 (1997).
31. Nelson, J. C., Bickford, D., Delucchi, K., Fiedorowicz, J. G. & Coryell, W. H. Risk of psychosis in recurrent episodes of psychotic and nonpsychotic major depressive disorder: A systematic review and meta-analysis. *Am. J. Psychiatry* **175**, 897–904 (2018).
32. De Winter, F. L. *et al.* No association of lower Hippocampal volume with Alzheimer's disease pathology in late-life depression. *Am. J. Psychiatry* **174**, 237–245 (2017).
33. Adamczuk, K. *et al.* Functional changes in the language network in response to increased amyloid B deposition in cognitively intact older adults. *Cereb. Cortex* **26**, 358–373 (2016).
34. Sexton, C. E. *et al.* Exploring the pattern and neural correlates of neuropsychological impairment in late-life depression. *Psychol. Med.* **42**, 1195–1202 (2012).
35. Vandenbergh, R. *et al.* 18F-flutemetamol amyloid imaging in Alzheimer disease and mild cognitive impairment a phase 2 trial. *Ann. Neurol.* **68**, 319–329 (2010).
36. Thomas, B. A. *et al.* The importance of appropriate partial volume correction for PET quantification in Alzheimer's disease. *Eur. J. Nucl. Med. Mol. Imaging* **38**, 1104–1119 (2011).
37. Müller-Gärtner, H. W. *et al.* Measurement of radiotracer concentration in brain gray matter using positron emission tomography: MRI-based correction for partial volume effects. *J. Cereb. Blood Flow Metab.* **12**, 571–583 (1992).
38. Jack, C. R. *et al.* Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol.* **9**, 119–128 (2010).
39. Mahgoub, N. & Alexopoulos, G. S. Amyloid hypothesis: Is there a role for anti-amyloid treatment in late-life depression?. *Am. J. Geriatr. Psychiatry* **24**, 239–247 (2016).
40. Smith GS, Kuwabara H, Nandi A, Gould NF, Nassery M, Savonenko A, *et al.* Molecular Imaging of Beta-Amyloid Deposition in Late-life Depression. *Neurobiol. Aging* **101**, 85–93 (2021). <https://doi.org/10.1016/j.neurobiolaging.2021.01.002>
41. Smith, G. S. *et al.* The functional neuroanatomy of geriatric depression. *Int. J. Geriatr. Psychiatry* **24**, 798–808 (2009).
42. Xie, C. *et al.* The co-existence of geriatric depression and amnesic mild cognitive impairment detrimentally affect gray matter volumes: Voxel-based morphometry study. *Behav. Brain Res.* **235**, 244–250 (2012).

43. Dickerson, B. C. *et al.* The cortical signature of Alzheimer's disease: Regionally specific cortical thinning relates to symptom severity in very mild to mild AD dementia and is detectable in asymptomatic amyloid-positive individuals. *Cereb. Cortex* **19**, 497–510 (2009).
44. Whitwell, J. L. *et al.* Temporoparietal atrophy: A marker of AD pathology independent of clinical diagnosis. *Neurobiol. Aging* **32**, 1531–1541 (2011).
45. Whitwell, J. L. *et al.* Neuroimaging correlates of pathologically defined subtypes of Alzheimer's disease: A case-control study. *Lancet Neurol.* **11**, 868–877 (2012).
46. Monroe, S. M. & Harkness, K. L. Life stress, the “kindling” hypothesis, and the recurrence of depression: Considerations from a life stress perspective. *Psychol. Rev.* **112**, 417–445 (2005).
47. Ritchie, K. *et al.* Association of adverse childhood environment and 5-HTTLPR genotype with late-life depression. *J. Clin. Psychiatry* **70**, 1281–1288 (2009).
48. Belleau, E. L., Treadway, M. T. & Pizzagalli, D. A. The impact of stress and major depressive disorder on hippocampal and medial prefrontal cortex morphology. *Biol. Psychiatry* **85**, 443–453 (2019).
49. Schmaal, L. *et al.* Subcortical brain alterations in major depressive disorder: Findings from the ENIGMA Major Depressive Disorder working group. *Mol. Psychiatry* **21**, 806–812 (2016).
50. Wolkowitz, O. W., Epel, E. S., Reus, V. I. & Mellon, S. H. Depression gets old fast: Do stress and depression accelerate cell aging?. *Depress Anxiety* **27**, 327–338 (2010).
51. Freret, T., Gaudreau, P., Schumann-Bard, P., Billard, J. M. & Popa-Wagner, A. Mechanisms underlying the neuroprotective effect of brain reserve against late life depression. *J. Neural Transm.* **122**, 55–61 (2015).
52. Jack, C. R. *et al.* An operational approach to National Institute on Aging-Alzheimer's Association criteria for preclinical Alzheimer disease. *Ann. Neurol.* **71**, 765–775 (2012).
53. Jack, C. R. *et al.* Suspected non-Alzheimer disease pathophysiology—concept and controversy. *Nat Rev Neurol* **12**, 117–124 (2016).
54. Cray, J. F. *et al.* Primary age-related tauopathy (PART): A common pathology associated with human aging. *Acta Neuropathol.* **128**, 755–766 (2014).
55. LaPoint, M. R. *et al.* The association between tau PET and retrospective cortical thinning in clinically normal elderly. *Neuroimage* **157**, 612–622 (2017).
56. Josephs, K. A. *et al.* Tau aggregation influences cognition and hippocampal atrophy in the absence of beta-amyloid: A clinico-imaging-pathological study of primary age-related tauopathy (PART). *Acta Neuropathol.* **133**, 705–715 (2017).
57. Gatchel, J. R. *et al.* Depressive symptoms and tau accumulation in the inferior temporal lobe and entorhinal cortex in cognitively normal older adults: A pilot study. *J. Alzheimer Dis.* **59**, 975–985 (2017).
58. Brown, E. E., Iwata, Y., Chung, J. K., Gerresten, P. & Graff-Guerrero, A. Tau in late-life depression: A systematic review and meta-analysis. *J. Alzheimer Dis.* **54**, 615–633 (2016).
59. Weisenbach, S. L., Kim, J., Hammers, D., Konopacki, K. & Koppelmans, V. Linking late life depression and Alzheimer's disease: Mechanisms and resilience. *Curr. Behav. Neurosci. Rep.* **6**, 103–112 (2019).
60. Ballmaier, M. *et al.* Hippocampal morphology and distinguishing late-onset from early-onset elderly depression. *Am. J. Psychiatry* **165**, 229–237 (2008).
61. Dickerson, B. C. & Eichenbaum, H. The episodic memory system: Neurocircuitry and disorders. *Neuropsychopharmacology* **35**, 86–104 (2010).
62. Epelbaum, S. *et al.* Neural correlates of episodic memory in the Memento cohort. *Alzheimer's Dement Transl. Res. Clin. Interv.* **4**, 224–233 (2018).
63. Rugg, M. D. & King, D. R. Ventral lateral parietal cortex and episodic memory retrieval. *Cortex* **107**, 238–250 (2018).
64. Wolk, D. A. & Dickerson, B. C. Fractionating verbal episodic memory in Alzheimer's disease. *Neuroimage* **54**, 1530–1539 (2011).
65. Buchsbaum, B. R. & D'Esposito, M. The search for the phonological store: From loop to convolution. *J. Cogn. Neurosci.* **20**, 762–778 (2008).
66. McDermott, L. M. & Ebmeier, K. P. A meta-analysis of depression severity and cognitive function. *J. Affect. Disord.* **119**, 1–8 (2009).
67. Chung, J. K. *et al.* Cortical amyloid β deposition and current depressive symptoms in Alzheimer disease and mild cognitive impairment. *J. Geriatr. Psychiatry Neurol.* **29**, 149–159 (2016).
68. Goukasian, N. *et al.* Association of brain amyloidosis with the incidence and frequency of neuropsychiatric symptoms in ADNI: A multisite observational cohort study. *BMJ Open* **9**, 1–10 (2019).
69. Cirrito, J. R. *et al.* Serotonin signaling is associated with lower amyloid- β levels and plaques in transgenic mice and humans. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 14968–14973 (2011).

Acknowledgements

We would like to thank Dr. Kristof Vansteelandt for providing help for the statistical analyses.

Author contributions

M.V., L.E., J.Van den.S., F.L.De.W., F.B., A.T. and T.V.C. designed the study. A.T., T.V.C., L.E., and M.K. analyzed the data. A.T. and T.V.C. wrote the first draft. M.K., F.L.De.W., F.B., J.Van den.S., S.S., P.D., R.V., K.Van.L., M.V., and L.E. contributed to the interpretation of the results and revised the first draft. All authors contributed to and have approved the final manuscript.

Funding

This study was supported by Research Foundation Flanders (FWO) project G.0746.09 (M. Vandenbulcke), G0C0319N (M. Vandenbulcke, F. Bouckaert, L. Emsell), Research Grant Old Age Psychiatry UZ Leuven R94859 (M. Vandenbulcke), KU Leuven C24/18/095 (M. Vandenbulcke, F. Bouckaert, J. Van den Stock, L. Emsell), KU Leuven Sequoia Fund. The tracer [^{18}F]-flutemetamol was provided free of charge by GE for this investigator led study. A. Takamiya has received grants from Keio University Medical Science Fund and Kanagawa Foundation for the Promotion of Medical Science to study in Belgium and is supported by AMED (Grant Number: JP20dm0307102h0003).

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-021-95206-0>.

Correspondence and requests for materials should be addressed to M.V.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021