



LRP6 Knockdown Ameliorates Insulin Resistance via Modulation of Autophagy by Regulating GSK3 β Signaling in Human LO2 Hepatocytes

Lei Li^{1,2}, Jing Xue¹, Jipeng Wan¹, Qian Zhou¹, Shan Wang¹, Yu Zhou¹, Heyong Zhao^{3*} and Xietong Wang^{1,3,4*}

¹ Department of Obstetrics and Gynaecology, Shandong Provincial Hospital Affiliated to Shandong University, Jinan, China, ² Department of Obstetrics and Gynecology, The First Affiliated Hospital of Chongqing Medical University, Chongqing, China, ³ Maternal and Child Health Care of Shandong Province, Jinan, China, ⁴ Key Laboratory of Birth Regulation and Control Technology of National Health and Family Planning Commission of China, Jinan, China

OPEN ACCESS

Edited by:

Timo Dirk Müller,
Helmholtz Center Munich-German
Research Center for Environmental
Health, Germany

Reviewed by:

Fouad Janat,
Independent Researcher, Westery, RI,
United States
Benyi Li,
University of Kansas Medical Center,
United States

*Correspondence:

Heyong Zhao
zhaohyong@126.com
Xietong Wang
xietongwangg@sina.com

Specialty section:

This article was submitted to
Cellular Endocrinology,
a section of the journal
Frontiers in Endocrinology

Received: 29 August 2018

Accepted: 25 January 2019

Published: 12 February 2019

Citation:

Li L, Xue J, Wan J, Zhou Q, Wang S,
Zhou Y, Zhao H and Wang X (2019)
LRP6 Knockdown Ameliorates Insulin
Resistance via Modulation of
Autophagy by Regulating GSK3 β
Signaling in Human LO2 Hepatocytes.
Front. Endocrinol. 10:73.
doi: 10.3389/fendo.2019.00073

Recent studies suggest that autophagy is highly involved in insulin resistance (IR). Inhibition of the PI3K/AKT/mTOR signaling pathway induces autophagy activation. Additionally, depletion of LRP6 has been shown to increase insulin sensitivity but its mechanism is still not clear. We hypothesized that LRP6 contributes to IR by regulating mTOR mediated autophagy through GSK3 β in hepatocytes. LO2 hepatocytes were treated with palmitate (PA) and insulin to induced IR. Levels of LRP6 mRNA and protein expression were measured by real time-PCR and western blot analysis. LRP6 knock down was achieved by adenovirus mediated Si-LRP6 expression and its roles in IR, glucose, GSK3 β , mTOR signaling, and autophagy were explored. Finally, GSK3 β was overexpressed and its involvement in autophagy and IR was examined. We found that PA treatment led to a reduced glucose uptake and IR in hepatocytes, which was accompanied by an upregulation of LRP6 expression. Knocking down of LRP6 enhanced glucose uptake and insulin sensitivity in PA treated cells, probably through increasing GSK3 β activity. Overexpression of GSK3 β mimicked LRP6 reduction by enhancing autophagy and ameliorating IR. Our study revealed a significant molecular mechanism connecting LRP6 to insulin sensitivity through GSK3 β -mTOR mediated autophagy.

Keywords: LRP6, insulin resistance, autophagy, GSK3 β , hepatocytes

INTRODUCTION

Insulin resistance (IR) plays a major role in the pathogenesis of several metabolic disorders, including type 2 diabetes, obesity, and hyperlipidemia. Although reduced physical activity and increased food, especially fat consumption, have been shown to contribute to IR, IR is a complicated condition with its exact mechanism still yet to be clarified (1, 2). Recently, accumulating evidence has pointed autophagy as an important player in the development of IR.

Autophagy is an important cellular process, responsible for removing intracellular debris and protein turnover (3). It is usually activated in stressed status such as starvation (4). In obese mice, IR is associated with downregulation of autophagy in the liver (5). Restoring autophagy enhances glucose tolerance and insulin sensitivity in obese mice. Increasing physical activity is associated with increased basal autophagy that improves glucose handling in diet-induced obese mice (6–8).

Mammalian target of rapamycin (mTOR), a protein whose activity is largely dependent on the nutrient availability such as amino acid and glucose, is a negative regulator of autophagy (9). mTOR is activated by the PI3K-Akt pathway which is negatively regulated by GSK3 β . mTOR phosphorylates key components such as Atg13 in the autophagy machinery. Inhibition of mTOR by rapamycin or nutrient starvation dephosphorylates Atg13 in mammalian cells, leading to induction of autophagy, accompanied by upregulation of Beclin 1 and LC3 II/I (10, 11).

A co-receptor of Wnt/ β -catenin signaling pathway, low-density-lipoprotein-related protein 6 (LRP6) is involved in insulin resistance. Mice heterozygous for LRP6 are less prone to diet-induced obesity and are protected from insulin resistance (12). However, the detailed mechanism underlying this regulation is not clearly characterized. Previously, LRP6 has been shown to attenuate GSK3 β activity (13). Although a link between LRP and autophagy has not been explored before, it is possible that LRP6 could potentially regulate mTOR signaling through modulating GSK3 β activity, which may then negatively impact autophagy.

Palmitate (PA) is a saturated fatty acid that induces IR in various scenarios, including hepatocytes, although the molecular mechanism remains elusive (14). We hypothesized that PA treatment leads to differentiated expression of LRP6 that alters the activity of GSK3 β and contributes to enhanced autophagy through mTOR in hepatocytes. In this current study, we explored the functions of LRP6 and GSK3 β in PA induced IR in LO2 hepatocytes. Our finding revealed a significant molecular mechanism linking the function of LRP6 to IR through autophagy.

MATERIALS AND METHODS

Cell Culture and Drug Treatment

LO2 hepatocytes were acquired from American Type Culture Collection (ATCC, Manassas, VA) and maintained according to standard procedure in high glucose Dulbecco's modified eagle medium (DMEM) supplemented with 10% fetal bovine serum, 100,000 U/L penicillin and 100 mg/L streptomycin. For palmitate (PA) and insulin treatment, cells were first cultured in 100 mm dishes to 80–85% confluence, washed with PBS three times, and then incubated in serum-free DMEM with 250 μ M PA dissolved in ethanol and complexed to BSA at a 2:1 molar ratio for 24 h. Cells were then washed with PBS for three times and incubated in the absence or presence of 0.1 μ M insulin for 10 min. An equivalent volume of ethanol added to 2% fatty-acid free BSA-DMEM was used as control treatment.

Western Blot Analysis

Cells were harvested and homogenized with RIPA buffer (50 mM Tris pH 8.0, 150 mM NaCl, 1% NP-40, 5 mM EDTA, 0.5% sodium deoxycholate, 0.1% SDS) supplemented with protease inhibitors and phosphatase inhibitors (Sigma, St. Louis, MO). Lysates with an equivalent amount of proteins were analyzed by

Western blot using the following antibodies: anti-phospho-IRS1-S307, anti-IRS1, anti-phospho-AKT-Thr308, anti-AKT, anti-LRP6, anti-Beclin-1, anti-LC3, anti-phospho-GSK3 β -Ser9, anti-GSK3 β , and anti- β -actin. Antibodies were purchased from Abcam (Cambridge, MA, USA) unless it was specified. Western blot was quantified by densitometry using ImageQuant TL 1D v8.1 software (GE Healthcare Life Sciences, Little Chalfont, Buckinghamshire, United Kingdom). Data were normalized to control treatment.

2-NBDG Assay

2-NBDG assay was performed to assess glucose uptake in cells as previously described (15). Briefly, LO2 hepatocytes were first treated with control or 250 μ M PA for 24 h and washed with PBS. Cells were then incubated in DMEM supplemented with 1 g/L glucose and 2% fatty acid free BSA for 3 h before being treated with or without 0.1 μ M insulin for 10 min. After washing with Krebs buffer containing 0.5% BSA, cells were treated with 150 μ g/ml 2-[N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl) amino]-2-deoxyglucose (2-NBDG), a fluorescent 2-deoxyglucose glucose analog (Cayman Chemicals, Arbor, MI, USA) for 40 min. A microplate reader (Perkin Elmer VictorX4, Waltham, MA, USA) was then used to measure the fluorescent signal representing the amount of 2-NBDG uptake at a excitation wavelength of 485 nm and emission wavelength of 535 nm.

Real Time-PCR (RT-PCR)



Lrp6 mRNA levels were assessed by real time-PCR. Total RNA was prepared with TRIzol[®] RNA Isolation Reagents (company, catalog number). The same amount of RNA was used to synthesize cDNA through iScript[™] cDNA Synthesis Kit (Life Technology, Pleasanton, CA, USA). RT-PCR was performed using SsoAdvanced[™] Universal SYBR[®] Green Supermix. Primers for *Lrp6* are: 5'-CACTTACTTCCCTGC AATTTTGAACC-3', and 5'-TGGCCTGTAGCTGTATGACCT ATG -3'. *Lrp6* mRNA levels were normalized to control treatment.

Recombinant Adenovirus Preparation and Infection

Expression vectors for recombinant adenovirus were constructed according to the Gateway cloning system (Invitrogen, Carlsbad, CA, USA) as previously described (16). Briefly, Si-LRP6, GFP-LC3, GSK3 β were cloned into the pAD/CMB/V5-DEST vector and packaged by ViraPower Adenoviral expression system (Invitrogen, Carlsbad, CA, USA). LO2 hepatocytes were infected with adenoviruses that express Si-LRP6, GFP-LC3 or GSK3 β at multiplicity of infection of 100 and harvested for further analysis after 48 h of infection.

GSK3 β Activity Assay

GSK3 β activity was assessed by GSK3 activity assay kit (Sigma, CS0990) according to manufacturer's instruction as described previously (17). Briefly, GSK3 β in LO2 hepatocytes was immunoprecipitated with an anti-GSK3 β antibody bound protein A/G affinity gel and incubated with γ -³²P-ATP. GSK3 β activity was measured by the incorporated ³²P.



Statistical Analysis

Data were represented as mean \pm SEM. Two-tailed unpaired *t*-test with equal variance was performed to determine the statistical difference between two groups. Significance was set at p -value < 0.05 .

RESULTS

PA Disrupts Insulin Signaling and Blocks Glucose Uptake in LO2 Hepatocytes

We first investigated the effects of PA treatment on insulin signaling and glucose uptake in LO2 hepatocytes. We found that PA treatment alone significantly reduced the levels of phosphorylated IRS1 (p-IRS1) and AKT (p-AKT) compared to the control treatment (**Figures 1A,B**). Additionally, it also suppressed insulin induced upregulation of p-IRS1 and p-AKT. These results indicate that PA treatment leads to

insulin resistance in LO2 hepatocytes. To further confirm this observation, we examined glucose uptake by 2-NBDG assay. We found that insulin significantly increased glucose uptake which was blocked by PA treatment (**Figure 1C**). In fact, PA treatment alone significantly reduced glucose uptake compared to that of control treatment.

PA Treatment Upregulates LRP6 Expression

Since LRP6 has previously been shown to regulate insulin sensitivity (12), we hypothesized that PA induced insulin resistance through LRP6. To test this hypothesis, we examined the mRNA and protein levels of LRP6 in LO2 hepatocytes treated with PA. We found that while insulin slightly increased both mRNA and protein levels of LRP6, PA led to a substantial upregulation of these levels following 24 h treatment (**Figures 2A,B**).




FIGURE 3 | Knock down of LRP6 attenuated PA-induced IR in LO2 hepatocytes. LO2 hepatocytes were infected with Ad-Si-LRP6 to knock down LRP6 protein expression. **(A)** LRP6 protein expression was reduced upon Ad-Si-LRP6 infection. $*p < 0.05$ vs. control ($n = 6$). **(B)** Western blot analysis of insulin signal pathway following LRP6 knock down. **(C)** Quantification of western blot results. Data were normalized to control treatment for each protein. $*p < 0.05$ vs. no-PA treatment control; $\#p < 0.05$ vs. PA treatment group ($n = 6$). **(D)** LRP6 knock down enhanced glucose uptake in PA treated LO2 hepatocytes in the presence of insulin. $*p < 0.05$ vs. no-insulin stimulated control; $\#p < 0.05$ vs. PA treatment group ($n = 6$).

LRP6 Knock Down Suppresses PA Induced Insulin Resistance

To determine if PA induced insulin resistance is indeed through upregulated LRP6, we investigated whether insulin resistance could be reversed by suppressing LRP6 expression. We knocked down LRP6 by adenovirus mediated Si-LRP6 expression and confirmed reduction of LRP6 protein in LO2 hepatocytes (Figure 3A). We found that LRP6 knock down significantly enhanced the insulin signaling pathway in the absence of PA treatment (Figures 3B,C). Additionally, LRP6 knock down also blocked PA-induced insulin resistance by increasing p-IRS1 and p-AKT levels. We further examined glucose uptake and found that LRP6 knock down significantly improved 2-NBDG uptake in the presence of PA treatment (Figure 3D).

LRP6 Knock Down Activates Autophagy in the Presence of PA Treatment

Since increasing autophagy has resulted in alleviating insulin resistance in hepatocytes, we investigated whether the induction

of LRP6 expression after PA treatment was involved in regulating autophagy. We found that PA treatment significantly suppressed autophagy as indicated by reduced expression of autophagy markers Beclin-1 and LC3II/I (Figures 4A,B). Knocking down LRP6 significantly increased Beclin-1 and LC3II/I protein expression in the presence of PA treatment. Furthermore, we also examined autophagosomes by expressing LC3-GFP. We found that LC3-GFP autophagosome formation was abolished by PA treatment and reversed by knocking down LRP6 (Figures 4C,D). These results indicate that LRP6 downregulation attenuates insulin resistance by activating autophagy.

LRP6 Knock Down Induces Autophagy Through Modulating GSK3 β Activity

Finally, we investigated GSK3 β and mTOR signaling in PA treated LO2 hepatocytes. We found that PA treatment significantly reduced phosphorylated GSK3 β (p-GSK3 β) levels in the presence of insulin. LRP6 knock down increased the level of GSK3 β , comparable to control treatment. Interestingly, we found




FIGURE 4 | Knock down of LRP improved insulin sensitivity through activation of autophagy in LO2 hepatocytes. **(A,B)** Western blot analysis **(A)** and quantification **(B)** showing that LRP6 knock down induced autophagy activation in LO2 hepatocytes ($n = 6$). **(C,D)** Representative images and statistical data of LO2 hepatocytes expressing GFP-LC3 via adenovirus infection upon on control, PA and/or Ad-Si-LRP6 treatment. * $p < 0.05$ vs. no-PA treatment group; # $p < 0.05$ vs. PA treatment group.

that PA treatment suppressed mTOR phosphorylation (p-mTOR) (Figure 5A). In fact, LRP6 knock down further reduced p-mTOR levels after PA treatment. We asked whether LRP6 knock down induced GSK3 β activation independent on phosphorylation. We examined GSK3 β activity and found that PA treatment inhibited GSK3 β activation. LRP6 knock out significantly enhanced GSK3 β activity (Figure 5B).

We then overexpressed GSK3 β in LO2 hepatocytes and examined the effects on autophagy and insulin signaling. We found that GSK3 β overexpression enhanced autophagy as indicated by increased levels of Beclin-1 and LC3 II/I in the presence of PA compared to that of PA treatment alone (Figure 5C). Additionally, we also found that GSK3 β overexpression alleviated PA induced insulin resistance by upregulating p-IRS1 and p-AKT (Figure 5D).

DISCUSSION

Understanding the molecular mechanism underlying insulin resistance will have significant implications in the development of therapeutic strategies against multiple metabolic disorders including type II diabetes. In the current study, we investigated the signaling pathway connecting LRP6 to IR in hepatocytes. We first confirmed that PA treatment in hepatocytes induced IR by suppressing glucose uptake and phosphorylation of key components in insulin signaling including IRS1 and AKT.

We found that LRP6 expression was substantially upregulated as a result of PA treatment and knock down of LRP6 restored IRS1 and AKT phosphorylation and glucose uptake. This correlates with an induction of autophagy which was suppressed by PA treatment alone. Importantly, we found that LRP6 regulated insulin sensitivity through GSK3 β and knock down of LRP6 enhanced GSK3 β activity. Consistent with this observation, we found that overexpression of GSK3 β in PA treated hepatocytes also restored autophagy induction and insulin signaling, as indicated by rescued IRS1 and AKT phosphorylation. To our knowledge, this is the first study linking LRP6 function to fat induced insulin resistance through autophagy.

Wnt signaling has been implicated in the pathogenesis of metabolic disorders including type 2 diabetes (18, 19). A single missense mutation in LRP6 that impairs Wnt signaling was identified in patients with metabolic syndrome (19). A previous study using heterozygous mice showed that LRP6 was involved in insulin resistance (12). Mice that were missing one copy of the LRP6 gene were protected from high fat induced obesity and insulin resistance, indicating that LRP6 may be negatively correlated with insulin sensitivity. However, it is not clear whether fat induced insulin resistance involves differential regulation of LRP6 expression. Our study revealed that the saturated fatty acid PA significantly enhanced both LRP6 mRNA and protein levels. Consistent with the mouse LRP6 heterozygous




FIGURE 5 | GSK3 β activation was involved in mediating the effects of LRP6 knock down on autophagy and insulin signaling. **(A)** LRP6 knock down inhibited mTOR in LO2 hepatocytes as shown by western blot analysis and quantifications of indicated proteins in the bar graph. * $p < 0.05$ vs. no-PA treatment group; # $p < 0.05$ vs. PA treatment group ($n = 6$). **(B)** LRP6 activated GSK3 β independent of inhibitory phosphorylation. * $p < 0.05$ vs. no-PA treatment group; # $p < 0.05$ vs. PA treatment group ($n = 6$). **(C)** Overexpression of GSK3 β activated autophagy in LO2 hepatocytes. Protein quantification was shown in the bar graph. * $p < 0.05$ vs. control group; # $p < 0.05$ vs. PA treatment group ($n = 6$). **(D)** Overexpression of GSK3 β attenuated insulin signal pathway disruption induced by PA treatment as shown by western blot analysis and quantifications of indicated proteins in the bar graph. * $p < 0.05$ vs. control group; # $p < 0.05$ vs. PA treatment group ($n = 6$).

study, our results using adenovirus mediated LRP6 knock down strategy indicated that modulating LRP6 levels may be sufficient to overcome insulin resistance in human hepatocytes. Although not frequent, a loss-of-function mutation (*LRP6*_{R611C}) has been identified in some patients with impaired insulin sensitivity (20). This study together with our finding suggest that a low expression level of LRP6 with normal function is both required and sufficient to maintain insulin reactivity.

A loss-of-function mutation of LRP6 resulted in a reduction of GSK3 β phosphorylation in response to insulin stimulation, as compared to cells with wild type LRP6 (20). Interestingly, our study also found enhanced GSK3 β phosphorylation in LRP6 knock down hepatocytes. However, we observed an increased GSK3 β activity independent on phosphorylation, which was accompanied by reduced mTOR activation. mTOR complex 1 (mTORC1) activity was reduced in LRP6 heterozygous mice

and increased in cells carrying a LRP6 loss-of-function mutation (12, 20, 21). mTOR activation prevents apoptosis while inhibition of mTOR signaling is necessary to activate autophagy to remove intracellular debris (22). Interestingly, a link between LRP6 and autophagy has not been established prior to our study.

Autophagy is an important cellular process that protects cells by removing unnecessary proteins. Deletion of a critical autophagy gene *Atg7* in the liver leads to accumulation of hepatic fat, indicating that autophagy is involved in fat metabolism (23). Recent accumulating evidence has identified autophagy as an important player in insulin signaling and glucose metabolism (24). Autophagy has been shown to be important for the maintenance of the normal function of insulin targeting tissues such as the liver, and the homeostasis of pancreatic β -cells. β -cell dysfunction is associated with impaired autophagy (25). However, consistent results have not been reached. Some studies showed that defective autophagy promoted insulin resistance while some reports pointed to the opposite, showing that autophagy deficiency actually protects from insulin resistance (5, 26, 27). We believe the function of autophagy in insulin resistance may be cell type specific. Consistent with the study in liver (5), we found that PA treatment-induced insulin resistance was accompanied by suppressed autophagy in hepatocytes. Importantly, we found that knock down of LRP6 rescued autophagy in PA treated hepatocytes which is in line with a function of autophagy in maintaining insulin sensitivity.

As previously stated, our results indicate that LRP6 regulates GSK3 β activity. Since GSK3 β is a negative regulator of mTOR signaling, we further explored the relation between GSK3 β and autophagy in PA treated hepatocytes. We found that adenovirus-mediated GSK3 β overexpression not only restored the levels of autophagy related proteins back to control status, but also enhanced insulin sensitivity. Thus, our results suggest that enhancing GSK3 β expression alone is also sufficient to suppress fat induced insulin resistance in hepatocytes.

Although beyond the scope of our current study, further detailed study is necessary to characterize the potential adverse effects when developing therapeutic strategies. Another limitation is that this study was focused on one cell type. One future direction for this research will be to characterize this signaling pathway in more cell types. Nevertheless, our study revealed a significant link among LRP6, GSK3 β , autophagy, and fat induced insulin resistance.

CONCLUSION

In summary, our findings suggest that fat induced insulin resistance may be caused by enhanced LRP6 expression, which leads to suppressed autophagy. Our results point to GSK3 β being a bridge connecting LRP6 to insulin resistance. Since modulating LRP6 and GSK3 β expression levels is sufficient to alter the status of insulin sensitivity in hepatocytes, altering LRP6 and GSK3 β expression may represent a potential therapeutic strategy for insulin resistance associated metabolism disorders.

AUTHOR CONTRIBUTIONS

LL, JX, JW, QZ, SW, and YZ, collected and analyzed the data. HZ and XW designed the study and wrote the manuscript.

FUNDING

This study was supported by Shandong Provincial Natural Science Foundation, China (ZR201702200727 and ZR2017BH046); Shandong Medical and Health Technology Development Program, China (2016WS0412 and 2016WS0411); Jinan Science and Technology Development Program (201602161); National Natural Science Foundation of China (81741037, 81801473, and 81571461); China Postdoctoral Science Foundation (2018M631067); Special Project Fund of Taishan Scholars.

REFERENCES

- Kitahara CM, Trabert B, Katki HA, Chaturvedi AK, Kemp TJ, Pinto LA, et al. Body mass index, physical activity, and serum markers of inflammation, immunity, and insulin resistance. *Cancer Epidemiol Biomarkers Prev.* (2014) 23:2840–9. doi: 10.1158/1055-9965.EPI-14-0699-T
- Sears B, Perry M. The role of fatty acids in insulin resistance. *Lipids Health Dis.* (2015) 14:121. doi: 10.1186/s12944-015-0123-1
- White E, Mehnert JM, Chan CS. Autophagy, metabolism, and cancer. *Clin Cancer Res.* (2015) 21:5037–46. doi: 10.1158/1078-0432.CCR-15-0490
- Lee WS, Yoo WH, Chae HJ. ER stress and autophagy. *Curr Mol Med.* (2015) 15:735–45. doi: 10.2174/1566524015666150921105453
- Yang L, Li P, Fu S, Calay ES, Hotamisligil GS. Defective hepatic autophagy in obesity promotes ER stress and causes insulin resistance. *Cell Metab.* (2010) 11:467–78. doi: 10.1016/j.cmet.2010.04.005
- Ghareghani P, Shanaki M, Ahmadi S, Khoshdel AR, Rezvan N, Meshkani R, et al. Aerobic endurance training improves nonalcoholic fatty liver disease (NAFLD) features via miR-33 dependent autophagy induction in high fat diet fed mice. *Obes Res Clin Pract.* (2018) 12:80–9. doi: 10.1016/j.orcp.2017.01.004
- Rosa-Caldwell ME, Brown JL, Lee DE, Blackwell TA, Turner KW, Brown LA, et al. Autophagy activation, not peroxisome proliferator-activated receptor gamma coactivator 1alpha, may mediate exercise-induced improvements in glucose handling during diet-induced obesity. *Exp Physiol.* (2017) 102:1194–207. doi: 10.1113/EP086406
- Rosa-Caldwell ME, Lee DE, Brown JL, Brown LA, Perry RA Jr, Greene ES, et al. Moderate physical activity promotes basal hepatic autophagy in diet-induced obese mice. *Appl Physiol Nutr Metab.* (2017) 42:148–56. doi: 10.1139/apnm-2016-0280
- Hara K, Yonezawa K, Weng QP, Kozlowski MT, Belham C, Avruch J. Amino acid sufficiency and mTOR regulate p70 S6 kinase and eIF-4E BP1 through a common effector mechanism. *J Biol Chem.* (1998) 273:14484–94. doi: 10.1074/jbc.273.23.14484
- Jiang P, Mizushima N. LC3- and p62-based biochemical methods for the analysis of autophagy progression in mammalian cells. *Methods* (2015) 75:13–8. doi: 10.1016/j.ymeth.2014.11.021
- Kang R, Zeh HJ, Lotze MT, Tang D. The Beclin 1 network regulates autophagy and apoptosis. *Cell Death Differ.* (2011) 18:571–80. doi: 10.1038/cdd.2010.191
- Liu W, Singh R, Choi CS, Lee HY, Keramati AR, Samuel VT, et al. Low density lipoprotein (LDL) receptor-related protein 6 (LRP6) regulates body fat and glucose homeostasis by modulating nutrient sensing pathways

- and mitochondrial energy expenditure. *J Biol Chem.* (2012) 287:7213–23. doi: 10.1074/jbc.M111.286724
13. Mi K, Dolan PJ, Johnson GV. The low density lipoprotein receptor-related protein 6 interacts with glycogen synthase kinase 3 and attenuates activity. *J Biol Chem.* (2006) 281:4787–94. doi: 10.1074/jbc.M508657200
 14. Ishii M, Maeda A, Tani S, Akagawa M. Palmitate induces insulin resistance in human HepG2 hepatocytes by enhancing ubiquitination and proteasomal degradation of key insulin signaling molecules. *Arch Biochem Biophys.* (2015) 566:26–35. doi: 10.1016/j.abb.2014.12.009
 15. Li H, Liu S, Yuan H, Niu Y, Fu L. Sestrin 2 induces autophagy and attenuates insulin resistance by regulating AMPK signaling in C2C12 myotubes. *Exp Cell Res.* (2017) 354:18–24. doi: 10.1016/j.yexcr.2017.03.023
 16. Yuan H, Wang T, Niu Y, Liu X, Fu L. AMP-activated protein kinase-mediated expression of heat shock protein beta 1 enhanced insulin sensitivity in the skeletal muscle. *FEBS Lett.* (2017) 591:97–108. doi: 10.1002/1873-3468.12516
 17. Sarikhani M, Mishra S, Maity S, Kotyada C, Wolfgeher D, Gupta MP, et al. SIRT2 deacetylase regulates the activity of GSK3 isoforms independent of inhibitory phosphorylation. *Elife* (2018) 7:e32952. doi: 10.7554/eLife.32952
 18. Fujino T, Asaba H, Kang MJ, Ikeda Y, Sone H, Takada S, et al. Low-density lipoprotein receptor-related protein 5 (LRP5) is essential for normal cholesterol metabolism and glucose-induced insulin secretion. *Proc Natl Acad Sci USA.* (2003) 100:229–34. doi: 10.1073/pnas.0133792100
 19. Mani A, Radhakrishnan J, Wang H, Mani A, Mani MA, Nelson-Williams C, et al. LRP6 mutation in a family with early coronary disease and metabolic risk factors. *Science* (2007) 315:1278–82. doi: 10.1126/science.1136370
 20. Singh R, De Aguiar RB, Naik S, Mani S, Ostadsharif K, Wencker D, et al. LRP6 enhances glucose metabolism by promoting TCF7L2-dependent insulin receptor expression and IGF receptor stabilization in humans. *Cell Metab.* (2013) 17:197–209. doi: 10.1016/j.cmet.2013.01.009
 21. Go GW, Srivastava R, Hernandez-Ono A, Gang G, Smith SB, Booth CJ, et al. The combined hyperlipidemia caused by impaired Wnt-LRP6 signaling is reversed by Wnt3a rescue. *Cell Metab.* (2014) 19:209–20. doi: 10.1016/j.cmet.2013.11.023
 22. Maiese K. Targeting molecules to medicine with mTOR, autophagy and neurodegenerative disorders. *Br J Clin Pharmacol.* (2016) 82:1245–66. doi: 10.1111/bcp.12804
 23. Martinez-Lopez N, Singh R. Autophagy and lipid droplets in the liver. *Annu Rev Nutr.* (2015) 35:215–37. doi: 10.1146/annurev-nutr-071813-105336
 24. Barlow AD, Thomas DC. Autophagy in diabetes: beta-cell dysfunction, insulin resistance, and complications. *DNA Cell Biol.* (2015) 34:252–60. doi: 10.1089/dna.2014.2755
 25. Ebato C, Uchida T, Arakawa M, Komatsu M, Ueno T, Komiya K, et al. Autophagy is important in islet homeostasis and compensatory increase of beta cell mass in response to high-fat diet. *Cell Metab.* (2008) 8:325–32. doi: 10.1016/j.cmet.2008.08.009
 26. Kim KH, Jeong YT, Oh H, Kim SH, Cho JM, Kim YN, et al. Autophagy deficiency leads to protection from obesity and insulin resistance by inducing Fgf21 as a mitokine. *Nat Med.* (2013) 19:83–92. doi: 10.1038/nm.3014
 27. Mellor KM, Bell JR, Young MJ, Ritchie RH, Delbridge LM. Myocardial autophagy activation and suppressed survival signaling is associated with insulin resistance in fructose-fed mice. *J Mol Cell Cardiol.* (2011) 50:1035–43. doi: 10.1016/j.yjmcc.2011.03.002

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Li, Xue, Wan, Zhou, Wang, Zhou, Zhao and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.