

## Macrofungi of manorial park in Turew near Poznań

ANNA BUJAKIEWICZ<sup>1</sup> and ANNA KUJAWA<sup>2</sup>

<sup>1</sup> Department of Plant Ecology and Environment Protection, Adam Mickiewicz University  
Al. Niepodległości 14, PL-61-713 Poznań, Poland

<sup>2</sup> Research Centre for Agricultural and Forest Environment of Polish Academy  
of Sciences, Field Station  
Szkołna 4, PL-64-003 Turew, Poland

Bujakiewicz A., Kujawa A.: Macrofungi of manorial park in Turew near Poznań. *Acta Mycol.* 35 (2): 183–195, 2000.

The aim of this study was to describe the species composition of macrofungi in a manorial park in Turew and to evaluate the importance of the park as a refuge habitat. The study carried out in 1997–1999 resulted in 171 taxa (*Basidiomycetes* – 84.7%, *Ascomycetes* – 15.3%). Among them 23 species were recognized to be protected or listed in the Polish Red List. It turned out that the manorial park in Turew is an important refuge area for many species of macrofungi occurring in agricultural landscape.

**Key words:** macrofungi, agricultural landscape, manorial park, conservation of fungi.

### INTRODUCTION

In an agricultural landscape which prevails in Polish lowlands the impoverishment and disappearance of natural, semi-natural and other non-farmed habitats is observed (Loster 1991; Ryszkowski and Bałazy 1995) because of deforestation, wetlands draining, water bodies eutrophication and area pollution which are closely linked to the development of intensive agriculture. In such intensively used farmlands many species are able to survive due to the presence of small-sized landscape elements such as forest islands, village parks and mid-field afforestations (Ryszkowski and Bałazy 1991; Dąbrowska-Prot 1998; Ratyńska and Szwed 1998 a, b). The importance of village parks for biodiversity protection in agricultural landscape was pointed out by many authors who showed that for plants as well as for animals village parks are refuge areas, especially on the areas where the forest was cut or changed into a one-species tree stand (Tomiałojć 1970; Olaczek 1972;

Kujawa 1992). For the effective conservation of fungi species richness the protection of their natural habitats is particularly important because the fungi cannot be held (like most vascular plant species and many animal species) in "mycological gardens" (Wojewoda 1976; Wojewoda and Ławrynowicz 1992). Though the adaptation of some species to habitats altered by man is sometimes observed (Wojewoda and Ławrynowicz 1992; Wojewoda 1996), most species of macrofungi are significantly endangered or even disappearing. Ca. 25% of the potential number of macrofungi occurring in Poland is listed in the Polish Red List (Wojewoda and Ławrynowicz 1992).

In this paper an attempt was taken to answer the question whether the village parks can play a role of refuge areas for macrofungi. Such a role of village parks was earlier confirmed in the case of plants and animals. The macrofungi of village parks in Poland have been poorly recognized so far. The studies were carried out in the parks in Radojewo (Lisiewska and Ratyńska 1984) and Uniejów (Lisiewska and Rybak 1990) as well as in arboreta in Goluchów (Lisiewska and Plączek 1993) and Kórnik (Lisiewska and Nowicka 1979).

The aim of the present study was to recognize the macrofungi species composition in a village park in Turew and to evaluate the importance of the park as a refuge habitat.

## STUDY AREA AND METHODS

The study was carried out in 1997–1999 in a manorial park in Turew established in the 18th century, located in the central part of the General Dezydery Chłpowski Landscape Park, ca. 50 km south of Poznań. The park in Turew is shaped as a "landscape" park with rather "open arrangement" (Błaszyk 1977) and covers the area of 21.86 ha (Wendlandt 1992). It comprises relatively well developed fragments of oak-hornbeam forest (*Galio sylvatici-Carpinetum*) and elm-alder forest (*Ficario-Ulmetum*) as well as a small enclave of ash-alder forest (*Circaco-Alnetum*). A small part of the park is kept as a walking area for visitors. Some cultivating activities (raking the litter, mowing the grass and herbs, removing the seedlings of trees and shrubs) are undertaken there. A major part of the park is not available for visitors. Here the cultivating activities are limited to removing some trees knocked down by the wind (Fig. 1 and 2). The tree stand of the park is characterized by a number of old trees. Ca. 80 trees fulfill the criteria of a "monumental tree" and 40 others will fulfill them within several years (Karg and Kujawa 1996).

The study was carried out with the aid of a transect method. Transects were located in fragments of oak-hornbeam forest, elm-alder forest and ash-alder forest. Additionally, fungi were also collected in other parts of the



Fig. 2. Fragment of the park in Turów with blooming *Lescocleis vernus* (Phot. Anna Kujawa)



Fig. 3. *Meriphan gyanus* growing on the roots of living *Ficus zylvestica* (Phot. Anna Kujawa)




Fig. 1. Plan of the manorial park in Turew (after Karg, Kujawa 1996)

park. Transects were visited 35 times altogether at least once a month from April to November and occasionally in other months. During each visit the number of fruit-bodies was recorded as well as the substrate, which the fungus was recorded on.

Recorded fungi were classified to three basic ecological groups: **terrestrial species** – i.e. fungi occurring on soil (saprotrophic and mycorrhizal ones); **litter decomposing species** – i.e. saprotrophs growing on litter (fallen leaves, dead herbs stems, fruits and little twigs); **lignicolous species** – i.e. saprotrophs occurring on dead branches, bark, stumps and logs as well as parasites growing on living trees.

#### LIST OF SPECIES

Collected macrofungi were determined according to several monographs and identification guides (among them the most important were: M o s e r

1963, 1983; Lisiewska 1987; Nespiak 1990; Domański 1991; Rudnicka-Jezierska 1991; Skirgiello 1991; Antonin and Noordeloos 1997). The nomenclature and systematic order of macrofungi follow those by Hawksworth et al. 1995.

Explanations:

GC	- Galio-Carpinetum	t	- terrestrial species
FU	- Ficario-Ulmietum	l	- litter decomposing species
CA	- Circaeo-Ulmietum	lg	- lignicolous species
L	- lawna	P	- protected species
O	- orchards	V	- valuable species
A	- allotments	I	- indeterminate (category) species
C	- charted species	R	- rare species

After Nespiak (1959), the abundance of fruit-bodies was measured with the aid of the Moser's scale (+, 1, 2, ...5) taking into account the maximum number recorded.

## ASCOMYCETES

### Leotiales

- Ascocoryne cylichnium* (Tul.) Korf - lg (on stump); FU - Oct. 1998; 2  
*Bulgaria inquinans* Fr. - lg (on dead *Carpinus betulus*); GC - July 1999; 4; R; C  
*Ciboria batsekiana* (Zopf.) Buchw. - l (on acorns); GC - Sept. 1999; 2  
*C. viridifusca* (Fuck.) von Höhnel. - l (on fruits of *Alnus glutinosa*); CA - Dec. 1997; 1  
*Cyathicula coronata* (Bull. ex Mérat) de Notaris - l (on stems of perennials); GC - Oct. 1999; 2  
*Hymenoscyphus fructigenus* (Bull.: Fr.) S. F. Gray - l (on acorns), FU - July, Sept. 1997, Aug. 1999; GC - Sept., Oct. 1997, Sept. 1998, Sept., Oct. 1999; 4  
*H. scutula* (Pers.: Fr.) Phill. - l (on stems of perennials and dry blades); FU - Aug. 1999; 2  
*Mollisia amenticola* (Sacc.) Rehm. - l (on fruits of *Alnus glutinosa*); CA - Dec. 1997; 1  
*M. cinerea* (Batsch. ex Mérat) Karst. - l (on fruits of *Fagus sylvatica*); FU - Oct. 1999; 1  
*Orbilia xanthostigma* (Fr.) Fr. - lg (on stump); FU - Sept. 1997; 3  
*Rutstroemia conformata* (Karst.) Nannf. - l (on leaves of *Alnus glutinosa*); CA - Oct. 1999; 2  
*R. futeovirescens* (Rob. in Desm.) White - l (on leaves of trees); FU - Oct. 1999; 2  
*R. sydowiana* (Rehm) White - l (on leaves of *Quercus robur*); GC - Sept. 1999; 2  
*Sclerotinia tuberosa* (Hedw.) Fuckel - t; FU - May 1999; 1

*Pezizales*

- Aleuria aurantia* (Fr.) Fuckel — t (among roots of knocked down *Fagus sylvatica* and on walks); FU — Oct. 1997, Sept. 1998, GC — Oct. 1999; 3  
*Helvella crispa* Fr. — t (on rotten wood in the place of wood accumulating and close to *Fagus sylvatica*); GC — Oct. 1997, FU — Oct. 1997; 2  
*Humaria hemisphaerica* (Wigg.: Fr.) Nannf. — t; FU — July 1998; 2  
*Leptopodla elastica* (Bull.) Boud. — t; FU — July 1998; 2  
*Macroscyphus macropus* (Pers.: Fr.) S. F. Gray — t; FU — Sept. 1998; +  
*Morchella gigas* (Batsch) Pers.: Fr. — t; O — May 1998, Apr. 1999, L — Apr. 1999; 2; P, C, 1  
*Otidea alutacea* (Pers.) Massoe — t; FU — Oct. 1997, June 1999; 2  
*Peziza micropus* Pers.: Pers. — lg (on log); GC — Oct. 1997, May 1998; 1  
*Scutellinia scutellata* (L.: St. Amans) Lambotte — t; CA — July 1997, May, June 1998, June 1999; 2  
*Tarzetta cupularis* (L.: Fr.) Lambotte ss. Dennis — t; GC — Sept. 1998; 1

*Sphaeriales*

- Hypoxylon fragiforme* (Pers.: Fr.) Fr. — lg (on dead *Fagus sylvatica*); FU — Oct. 1998; 2  
*Xylaria hypoxylon* (L.: Hook.) Grev. — lg (on stump); FU — Aug., Dec. 1997, Nov. 1998; 2  
*X. polymorpha* (Pers.) Grev. — lg (on stump); FU — June 1999; 4

**BASIDIOMYCETES***Auriculariales and Tremellales*

- Exidia truncata* Fr. — lg (on *Corylus avellana*); FU — June 1999; 2  
*Hirneola auricula-judae* (Bull.) Berk. — lg (on *Sambucus nigra*, *Fraxinus excelsior* and *Ribes nigrum*); GC — Oct. 1997, Oct., Nov. 1998; CA — May, June 1998; O — Apr. 2000; 2; C

*Aphylophorales*

- Auriculariopsis ampla* (Lév.) Maire — lg (on twigs of broad-leaved trees); GC — Sept. 1997; CA — May 1999; 2  
*Bjerkandera adusta* (Willd.: Fr.) P. Karst. — lg (on dead *Fagus sylvatica*); FU — Oct. 1999; 4  
*Byssomerulius corium* (Fr.) Parm. — lg (on twigs of broad-leaved trees); CA — Aug. 1997; 1  
*Chondrostereum purpureum* (Pers.: Fr.) Pouzar — lg (on log); GC — Sept. 1999; 2

- Clavulina cinerea* (Fr.) Schroet. - t; GC - Sept. 1997, July 1998; 1  
*Fistulina hepatica* (Schff.) Fr. - lg (on old living and dead *Quercus robur*);  
 GC - Sept. 1997, Oct. 1998, Sept. 1999; 2; V; C  
*Fomes fomentarius* (L.: Fr.) Kickx - lg (on dead *Fagus sylvatica*); GC - June  
 1997; 2; C  
*Ganoderma applanatum* (Pers.: Wallr.) Pat. - lg (on dead *Ulmus minor*); FU -  
 Sept. 1998; 2; C  
*Gloeophyllum odoratum* (Wulf.: Fr.) Imaz. - lg (on stump); GC - Sept. 1997,  
 Apr. 2000; 1  
*Laetiporus sulphureus* (Bull.: Fr.) - lg (on dead *Quercus robur*); GC - June  
 1997, Sept. 1999; 1  
*Meripilus giganteus* (Pers.: Fr.) P. Karst. - lg (about dead *Quercus robur* and  
 on roots of living *Fagus sylvatica*); FU - Oct. 1999, GC - Oct. 1999;  
 2; P; C (Fig. 3)  
*Merulius tremellanus* Fr. - lg (on branches of dead *Quercus robur*); GC -  
 Oct. 1999; 2  
*Piptoporus betulinus* (Bull.: Fr.) P. Karst. - lg (on dead *Betula pendula*); FU -  
 Jan. 1999; +; C  
*Polyporus arcularius* (Batsch) Fr. - lg (on twigs); FU - Oct. 1998; +  
*P. squamosus* (Huds.) Fr. - lg (on living *Ulmus minor* and dead *Acer  
 platanoides*); GC - May 1997, FU - May 1998; 2  
*P. varius* Fr. - lg (on dead *Ulmus minor*); FU - Dec. 1997; 1  
*Postia subcaesia* (David) Jülich - lg (on stump); FU - Oct. 1997; 2  
*Ramaria stricta* (Fr.) Quél. - t; GC - Sept. 1998; +  
*Schizophyllum commune* Fr. - lg (on dead *Fraxinus excelsior*, *Quercus robur*  
 and *Juglans regia*); FU - March 1998; GC - Oct. 1998; 4; C  
*Stereum hirsutum* (Wülld.: Fr.) S.F. Gray - lg (on roots of knocked down *Tilia  
 cordata*); GC - Nov. 1998; 4  
*S. subtomentosum* Pouz. - lg (on dead branch); CA - Sept. 1997; 1; R  
*Typhula erythropus* Fr. - l (on leaves and twigs); CA - Oct. 1999; 2

#### Boletales

- Paxillus atrotomentosus* (Batsch) Fr. - lg (about stump); GC - Sept. 1998; 1  
*P. filamentosus* Fr. - t; FU - Sept., Oct. 1997, Sept. 1998; 2; R  
*P. involutus* (Batsch) Fr. - t; GC - Sept. 1997; 1  
*Xerocomus badius* (Fr.) Kühner ex Gilb. - t; GC - Oct. 1997, Oct. 1999; 2  
*X. chrysenteron* (Bull. ex St. Amans) Quél. - t; GC - Aug., Sept. 1997;  
 Aug. 1998; 1  
*X. subtomentosus* (L.: Fr.) Quél. - t; July 1998; 1

#### Agaricales

- Agaricus arvensis* Schff.: Fr. - t; GC - Oct. 1999; 2  
*A. semotus* Fr. - t; FU - Sept. 1998; +

- Agrocybe cylindracea* (DC.: Fr.) Mre. - 1 (on branch buried in the ground); GC - May 1999; 1; R
- Amanita citrina* (Schff.) S.F. Gray - t; GC - Oct. 1997; 1; C
- A. phalloides* (Vaill.) Socr. - t; FU - Sept., Oct. 1997, Aug. - Oct. 1999; 1; C
- A. rubescens* (Pers.: Fr.) S.F. Gray - t; GC - Oct. 1997; 2
- Armillaria mellea* s.l. - t; GC - Oct. 1997, Oct. 1998; 4; C
- Clitocybe clavipes* (Pers.: Fr.) Kummer - t; GC - Oct. 1997; 2
- C. flaccida* (Sow.: Fr.) Kummer - 1 (on leaves); GC - Oct. 1997; 2
- C. lignatilis* (Pers.: Fr.) Karst. - lg (on log); FU - Aug. 1997; 1, V
- Conocybe mairei* (Kühner ex) Watling - t; FU - July 1999; 1
- C. rickeniana* P.D. Orton - t; GC - Oct. 1997; 2
- Coprinus atramentarius* (Bull.: Fr.) Fr. - t; GC - Apr. 1998, Oct. 1999; 2
- C. cortinatus* Lge. - t; FU - Sept. 1999; 1
- C. disseminatus* (Pers.: Fr.) S.F. Gray - lg (on and about stump); FU - July, Sept. 1998, GC - June 1999; 4
- C. hemerobius* Fr. - t; FU - July, Oct. 1998, Sept. 1999; 2
- C. impatiens* (Fr.) Quéf. - t; FU - Oct. 1997; 1
- C. micaceus* (Bull.: Fr.) Fr. - lg (about living *Carpinus betulus*); GC - Nov. 1998; 2
- C. plicatilis* (Curt.: Fr.) Fr. - t; FU - July, Sept. 1998, Aug. 1999, CA - May 1998; 2
- Cortinarius abietorum* (Vel.) Mos. - t; FU - Oct. 1998; CA - Oct. 1998; 2
- C. helveticoides* (Fr.) Fr. - t; CA - Oct. 1998; 2
- Crepidotus variabilis* (Pers.: Fr.) Kummer - t (on branch); GC - Sept., Oct. 1999; 2
- Cystolepista sistrata* (Fr.) Sing. - t; GC - Sept. 1998; 1
- Entoloma araneosum* (Quéf.) Mos. - t; FU - Oct. 1997, Oct. 1998, Aug. 1999; 1
- E. hyssedum* (Pers.: Fr.) Donk - lg (on branch); CA - Sept. 1999; +; 1
- E. nidarosum* (Fr.) Quéf. - t; FU - Aug. 1999; 2
- E. rhodopolium* (Fr.) Kummer - t; GC - Sept. 1997; 2
- Flanmulinia velutipes* (Curt.: Fr.) Sing. - lg (on dead *Ulmus minor*); FU - Nov. 1998; 2
- Galerina ulicolor* (Fr.) Sing. - lg (on log); FU - Oct. 1998; 2
- Gymnopis confluens* (Pers.: Fr.) Antonin (*Collybia confluens* (Pers.: Fr.) Kummer) - t (often in fairy rings); FU - Aug. 1998, GC - Sept., Oct. 1997, Oct. 1998, June, July, Sept. 1999; 4
- G. dryophilus* (Bull.: Fr.) Murrill (*Collybia dryophila* (Bull.: Fr.) Kummer - 1 (litter); FU - Sept. 1998; GC - June, Aug., Sept. 1997; 1
- G. oclor* (Pers.) Antonin et Noordel. (*Collybia exculpta* (Fr.) Gill.) - t; FU - June 1999; 1
- G. peronata* (Bolt.: Fr.) Antonin (*Collybia peronata* (Bolt.: Fr.) Sing.) - 1 (litter); FU - Sept. 1998, Aug. 1999, GC - Sept., Oct. 1997, Sept. 1999; 1
- Hebeloma sacchariolens* Quéf. - t; FU - Aug. 1999; +
- H. testaceum* (Batsch: Fr.) Quéf. - t; FU - Oct. 1999; 1



- Hohenbuehelia myxotricha* (Lév.) Sing. - lg (on branches of dead *Fraxinus excelsior*); CA - Sept. 1997, 1999; 2; R
- Hypholoma fasciculare* (Huds.: Fr.) Kummer - lg (about dead *Quercus robur*); GC - Aug., Oct. 1997, May, June 1999; 2
- Inocybe fastigiata* (Schff.: Fr.) Quéf. - t; GC - July 1998; 1
- I. geophylla* (Sow.: Fr.) Kummer - t; GC - Sept. 1997, Aug. 1998; 2
- I. geophylla* (Sow.: Fr.) Kummer var. *violacea* Pat. - t; GC - Oct. 1997, July 1998; 2
- I. patouillardii* Bres. - t; GC - June 1999; 1; C
- I. perlata* Cooke - t; FU - Oct. 1997, July 1998, Aug. 1999, GC - Oct. 1997; 2
- I. petiginosa* (Fr.: Fr.) Gill. - t (close to *Fagus sylvatica*); FU - Sept., Oct. 1997, 1998, 1999; 2
- Kuehneromyces mutabilis* (Schaeff.: Fr.) Sing. ex Smith - lg (on log and roots of knocked down tree); GC - Sept. 1997, Sept., Nov. 1998, Apr., Sept. 1999; 4; C
- Laccaria amethystina* (Bolt. ex Hook.) Murr. - t; FU - Oct. 1997, 1998, 1999; 2; C
- L. laccata* (Scop.: Fr.) Bk. et Br. - t; FU - Aug. 1999, GC - Oct. 1997, 1998, 1999; 2
- L. tortilis* (Bolt.) S.F. Gray - t, FU - Sept., Oct. 1999; 2
- Lepiota aspera* (Pers. in Hofm.) Quéf. - t (on rotten wood in the place of wood accumulation); FU - Oct. 1997; 2
- L. cristata* (A. et S.: Fr.) Kummer - t, GC - Sept. 1997; +
- L. pseudofelina* Lge. - t; FU - Sept. 1998; 1; I
- Lepista nebularis* (Fr.) Harmaja - t; GC - Oct. 1997, 1998; 2
- L. nuda* (Bull.: Fr.) Cke - t; GC - Oct. 1997, 1998; 2
- Macrolepiota procera* (Scop.: Fr.) Sing. - t; GC - Aug. 1998; 2; I
- M. rhacodes* (Vitt.) Sing. - t; GC - Oct. 1998; 2; I
- Marasmius oreades* (Bolt.: Fr.) Fr. - t; L - Sept. 1997; 2
- M. setosus* (Sow.) Noordel. (*Marasmius recubans* Quéf.) - l (on leaves, twigs and acorns of *Quercus robur*); FU - Oct. 1997, 1998, 1999; 2; I
- M. rotula* (Scop.: Fr.) Fr. - l (on twigs); FU - July - Oct. 1997, June 1998, June - Aug. 1999, GC - June 1997, Oct. 1998, VI, Sept. 1999; 2
- M. scorodanius* (Fr.) Fr. - l (on twigs); GC - Sept. 1999; 1
- Melanoleuca cognata* (Fr.) K. et M. - t; GC - Sept. 1998, Oct. 1999; 1
- Mycena acicula* (Schff.: Fr.) Kummer - l (on twigs); FU - Aug. 1997, June, July 1999, GC - Sept. 1997, June 1999; 1
- M. aestives* (Fr.) Quéf. - l (on leaves); GC - Oct. 1997, CA - Aug. 1997; 3
- M. alcalina* (Fr.) Kummer - lg (on stump); GC - Oct. 1998; 2
- M. aurantiomarginata* (Fr.) Quéf. - l (on leaves); FU - Oct. 1999; 1; R
- M. citrinomarginata* Gill. - l (on leaves); FU - June 1999; +
- M. chlorinella* (Lge.) Sing. - l (on leaves); CA - Oct. 1999, GC - Oct. 1998, Sept. 1999; 2
- M. flavescens* Vel. - l (on leaves of *Fagus sylvatica*); FU - Oct. 1999; 2; R

- M. galericulata* (Scoop: Fr.) S.F. Gray - l-ig (on twigs and pieces of wood); CA - Oct. 1999, FU - Oct. 1997, GC - Sept., Oct. 1997, Sept., Oct. 1998, Sept., Oct. 1999; 2
- M. galopus* (Pers.: Fr.) Kummer - l (on twigs under the ground); CA - June 1999, FU - Oct. 1997, June, July, Oct. 1999, GC - Sept. 1997, July, Sept., Oct. 1998, June, Sept. 1999; 2
- M. lineata* (Fr.) Quéf. ss. Lange - lg (on stump); FU - Sept. 1998; 1
- M. polygramma* (Bull.: Fr.) S.F. Gray - l (on twigs buried in the ground), GC - Oct. 1997, Sept. 1999; 2
- M. pseudocorticicola* Kühn. - lg (on mossy butt of *Ulmus minor*, *Fraxinus excelsior* and *Quercus robur*); FU - Dec. 1997, GC - Dec. 1997, Oct. 1999; 2; R
- M. pura* (Pers.: Fr.) Kummer - t; GC - Oct. 1997, Sept. 1998, Oct. 1999; 2
- M. purpureofusca* (Peck) Sacc. - lg (on stump); GC - Oct. 1997, 1998; +; V
- M. sanguinolenta* (Alb. et Schw.: Fr.) Kummer - l (on leaves); Fu - June 1999, GC - Oct. 1998; 2
- M. speirea* (Fr.: Fr.) Gill. - l (on twigs); FU - Sept. 1997, Sept. 1998, June, Aug., Oct. 1999; 2
- M. tintinnabulum* (Fr.) Quéf. - lg (on stump); FU - Aug. 1999, GC - Dec. 1997; 4
- M. vitilis* (Fr.) Quéf. - l (on leaves); CA - Oct. 1998, GC - Oct. 1997, Oct. 1998, May 1999; 2
- M. zephyrus* (Fr.: Fr.) Kummer - l (on leaves); GC - Oct. 1998, Oct. 1999; 1
- Naucoris abietorum* (R. Mre.) Kühner et Romagn. - l (on twigs); CA - Oct. 1998; 1
- N. excharoides* (Fr.: Fr.) Kummer - l (on twigs); CA - Sept. 1998, Oct. 1999, FU - Oct. 1997, Sept. 1998, Sept., Oct. 1999; 2
- N. scolecina* (Fr.) Quéf. - t; CA - Aug., Oct. 1997, May, June, Sept., Oct. 1998, May 1999, FU - Oct. 1997, Aug., Oct. 1999; 3
- N. subconspersa* Kühner - l (on twigs buried in the ground); CA - Sept. 1998; 2
- Oudemansiella radicata* (Rehhan: Fr.) Sing. - t; GC - Aug., Oct. 1997, Sept. 1999; 1
- Pholiota aurivella* (Batsch: Fr.) Kummer - lg (on roots of knocked down *Fagus sylvatica*); GC - Oct. 1997; 3
- Ph. squarrosa* (Pers.: Fr.) Kummer - lg (on stump); GC - Sept. 1998; 2
- Pleurotus ostreatus* (Jacq.: Fr.) Kummer - lg (on dead *Ulmus minor*); FU - Dec. 1997, Nov. 1998; 3
- Pluteus atricapillus* (Secr.) Sing. - lg (on stump); CA - Aug. 1999, FU - Sept., Oct. 1997, June 1998; Aug. 1999, GC - June, Aug., Oct. 1997, Aug., Oct. 1998, Sept. 1999; 1
- P. podospileus* Sacc. ex Cub. - l (on twigs buried in the ground); CA - Sept., Oct. 1999; 1
- P. romellii* (Britz.) Laplanche - t; FU - Aug. 1999; +

- P. semibulbosus* (Lasch sp. Fr.) Gill (*P. houdieri* Orton) – lg (on stump); FU – Sept. 1998; +
- P. umbrinus* (Pers.: Fr.) Kummer – lg (on log); FU – Oct. 1997, Nov. 1998; +
- Psathyrella candolleana* (Fr.) Mre. – l (on pieces of wood); CA – Sept. 1997, June, Sept. 1998, June, July, Sept. 1999; 3
- P. gracilis* (Fr.) Quél. – l (on pieces of wood); CA – Oct. 1999; 2
- P. hydrophila* (Bull.: Merat) R. Mre. – lg (about and on stump); GC – Sept. 1997, Sept. 1998, Oct. 1999; 4
- P. prona* (Fr.) Gill. – t; FU – Oct. 1997; 2
- P. spadiceo-grisea* (Fr.) R. Mre. – t; CA – Apr. 1998, Apr., May 1999, FU – March 1998; 1
- P. subatrata* (Batsch.: Fr.) Gill. – t; FU – Oct. 1997; 2
- P. velutina* (Pers.: Fr.) Sing. – t; FU – Sept. 1998, Aug. 1999; 2
- Rhodocollybia butyracea* (Bull.: Fr.) Lennox (*Collybia butyracea* (Bull.: Fr.) Quél.) – t; GC – Oct. 1997, Sept., Oct. 1998; 2
- Simocybe rubi* (Berk.) Sing. – t (on twigs); FU – Aug. 1997; 1
- Stropharia aeruginosa* (Curt.: Fr.) Quél. – t; GC – Oct. 1997, 1998; 2
- S. coronilla* (Bull.: Fr.) Quél. – t; T – Sept. 1998; +
- Tricholoma sulphureum* (Bull.: Fr.) Kummer – t; FU – Oct. 1999; 1; C
- T. terreum* (Schff.: Fr.) Kummer – t; GC – Oct. 1999; 1
- Volvariella taylori* (Berk.) Sing. – t; FU – Aug., Sept. 1999; +; I
- V. volvacea* (Bull.: Fr.) Sing. – lg (on stump); GC – June 1999; 1; I

#### *Russulales*

- Lactarius blennius* Fr. – t (close to *Fagus sylvatica*); FU – Sept. 1997, Oct. 1998, Oct. 1999; 2
- Russula cyanoxantha* (Schff.) Fr. – t (close to *Quercus robur*); GC – Aug. – Oct. 1997, July 1998; 1
- R. delicata* Fr. – t; GC – Oct. 1997, July 1998; 1
- R. grisea* (Pers.) Fr. ss. Gill. – t; GC – Sept. 1999; 2
- R. mairei* Sing. – t (close to *Fagus sylvatica*); Aug. – Oct. 1997, Oct. 1998, July, Sept., Oct. 1999; 2
- R. pumila* Rouzeau et Massatr. – t; CA – Oct. 1998; +; I
- R. solaris* Ferd. et Winge – t (close to *Fagus sylvatica*); FU – Oct. 1997, Sept. 1998; 2

#### *Sclerodermatales*

- Scleroderma verrucosum* Pers. – t; GC – Oct. 1997, June 1998; 1

#### *Nidulariales*

- Crucibulum laeve* (Huds. ex Relh.) – lg (on branches); GC – July 1998; 2
- Cyathus olla* (Batsch): Pers. – l (on dead stems of *Rubus idaeus*); A – March 1999; 2; C

*Lycoperdales*

*Langermannia gigantea* (Batsch.: Pers.) Rostk. — t; GC — Oct. 1997; 1; P; C

*Lycoperdon perlatum* Pers. — t; GC — Oct. 1997; 2

*L. pyriforme* Schff.: Pers. — lg (about stump); GC — Oct. 1997; 2

*Phallales*

*Phallus impudicus* L.: Pers. — t; GC — Oct. 1997; 1; P; C

## RESULTS AND CONCLUSIONS

1. In the manorial park in Turew 177 taxa (176 species and 1 variety) were recorded in total.
2. Terrestrial species constitute the most numerous group of species (47.5% of all taxa). A group of lignicolous species make 31.6% and a group of litter decomposing species — 20.9%.
3. The proportion of *Ascomycetes* was relatively high and constitutes 15.3%. Most species belonged to *Basidiomycetes* (84.7%). Genera with the highest number of species were: *Mycena* — 19 spp., *Coprinus* — 7, *Psathyrella* — 7, *Russula* — 6, *Inocybe* — 5 + 1 var., *Pluteus* — 5.
4. Among the recorded macrofungi 23 species (13% of all) are protected by law in Poland: *Langermannia gigantea*, *Meripilus giganteus*, *Morchella gigas* and *Phallus impudicus* or listed in Polish Red List (Wojewoda and Ławrynowicz 1992): *Agrocybe cylindracea*, *Bulgaria inquinans*, *Clitocybe lignatilis*, *Entoloma byssisedum*, *Fistulina hepatica*, *Hohenbuehelia myxotricha*, *Lepiota pseudofelina*, *Macrolepiota procera*, *M. rhacodes*, *Marasmius setosus*, *Morchella gigas*, *Mycena aurantiomarginata*, *M. flavescens*, *M. pseudocorticola*, *M. purpureofusca*, *Paxillus filamentosus*, *Russula pomila*, *Stereum subtomentosum*, *Volvariella taylora*, *V. volvacea*. Besides, four other rare species were found: *Ascocoryne cylichnium*, *Coprinus coriñatus*, *Exidia truncata* and *Pluteus podospileus*.
5. The geographical distribution in Europe of 19 species occurring in the park in Turew was mapped by Skirgiello (1965–1984).
6. The majority of species were recorded in an oak-hornbeam forest (93 spp., incl. 68 exclusive spp.), which covers the largest area. 85 spp. occurred in an elm-alder forest (60 exclusive spp.), and 29 spp. — in an ash-alder forest (16 exclusive), covering the smallest area in the park. The species number distribution is probably partially influenced by the differentiated size of a studied habitat (and the length of transect). Only three species were found in the all studied types of forests: *Mycena galericulata*, *M. galopus* and *Pluteus atricapillus*.
7. The results show that the manorial park in Turew is an important refuge area for many species of macrofungi occurring in agricultural landscape.

## REFERENCES

- Antonin V., Noordeloos M. E. 1997. A Monograph of *Marasmius*, *Collybia* and related genera in Europe. Part 2: *Collybia*, *Gyrovopse*, *Rhodescollybia*, *Crispella*, *Chaetocaulon* and additions to *Marasmius*. Libri Botanici 17. IHM-Verlag.
- Błaszczak H. 1977. Ewidencja parku pałacowego w Turwi (mac).
- Dąbrowska-Prot E. 1998. Ekologiczne problemy wysp środowiskowych w krajobrazie ze szczególnym uwzględnieniem wysp leśnych. In: J. Banaszak (ed). Ekologia wysp leśnych. WSP Bydgoszcz: 177-192.
- Domasński S. 1991. Flora Polska. Grzyby (*Mycota*) 21: *Basidiomycetes*, *Aphyllophorales*, *Stereaceae*, *Podocyphaceae*. PWN. Warszawa-Kraków.
- Grzywacz A. 1989. Grzyby chronione. PWRiL. Warszawa.
- Hawksworth D. L., Kirk P. M., Sutton B. C., Pegler D. N. 1995. Ainsworth & Bisby's Dictionary of the Fungi. (8 ed.). CAB International University Press, Cambridge, UK.
- Karg J., Kujawa A. 1996. Zabytkowy park pałacowy w Turwi - porównanie stanu drzewostanu na przestrzeni 15 lat. Biol. Parków Krajobrazowych Wielkopolski. Poznań 1 (3): 51-61.
- Kujawa K. 1992. Znaczenie parków podworskich dla zachowania bogactwa awifauny lęgowej w krajobrazie rolniczym. In: S. Błaszczak, L. Ryszkowski (eds). Produkcja pierwotna, zasoby zwierząt i wymywanie materii organicznej w krajobrazie rolniczym. Zakład Badań Środowiska Rolniczego i Leśnego PAN. Poznań: 137-150.
- Lisiewska M. 1987. Flora Polska. Grzyby (*Mycota*) 17: *Mycota*. PWN. Warszawa-Kraków.
- Lisiewska M., Nowicka D. 1979. *Macromycetes Arborescentium Kórnickiego Arboretum Kórnickie* 24: 339-371.
- Lisiewska M., Piączek U. 1993. *Macromycetes* zbiorowisk leśnych parku w Gołuchowie (województwo kaliszkie). Bad. Fizjograf. Pol. Zach. 42, Bot: 47-75.
- Lisiewska M., Ratyńska H. 1984. *Macromycetes* na tle zbiorowisk leśnych parku w Radziejowie koło Poznania. Bad. Fizjograf. Pol. Zach. 35, Bot: 5-23.
- Lisiewska M., Rybak M. 1990. Udział makromycetów w zespołach leśnych w Uniejowie. Bad. Fizjograf. Pol. Zach. 40, Bot: 5-28.
- Luster S. 1991. Różnorodność florystyczna w krajobrazie rolniczym i znaczenie dla niej naturalnych i półnaturalnych zbiorowisk wyspowych. *Fragm. Flor. Geobot.* 36 (2): 427-457.
- Moser M. 1963. *Ascomycetes. Kleine Kryptogamenflora*. 2a. Jena.
- Moser M. 1983. *Die Röhrlinge und Blätterpilze. Kleine Kryptogamenflora*. II h/2. *Basidiomycetes 2*. Gustav Fischer Verlag, Stuttgart, New York.
- Nespiak A. 1959. Studia nad udziałem grzybów kapeluszowych w zespołach leśnych na terenie Białowieckiego Parku Narodowego. *Monogr. Bot.* 8: 2-141.
- Nespiak A. 1990. Flora Polska. Grzyby (*Mycota*) 19. *Inocybe*. PWN. Warszawa-Kraków.
- Olszczek R. 1972. Parki wiejskie ostoją rodzimjej flory leśnej. *Chrośny Przyr.* Ojcz. 28 (2): 5-22.
- Olszczek R. 1976. Zmiany w szacie roślinnej Polski od połowy XIX wieku do lat bieżących. *Zesz. Probl. Post. Nauk Roln.* 177: 369-408.
- Ratyńska-Nowak H. 1986. Charakterystyka fitosocjologiczna i florystyczna zbiorowisk leśnych i zaroślowych na obszarze projektowanego Parku Agrokologicznego im. gen. D. Chłapowskiego (mac).
- Ratyńska H., Szwed W. 1998a. Wyspy leśne i zadrzewienia jako refugia. Flora i zbiorowiska roślinne. In: J. Banaszak (ed). Ekologia wysp leśnych. WSP Bydgoszcz: 227-230.
- Ratyńska H., Szwed W. 1998b. Problemy zagospodarowania krajobrazu rolniczego. In: J. Banaszak (ed). Ekologia wysp leśnych. Wyd. WSP Bydgoszcz: 315-326.
- Rudnicka-Jeziarska W. 1991. Flora Polska. Grzyby (*Mycota*) 23: *Lycoperdales*, *Sclerotiumatales*, *Tubulomatales*, *Nidulariales*, *Phalloles*, *Podzales*. PWN. Warszawa-Kraków.

- Ryszkowski L., Bałazy S. 1991. Strategia ochrony żywych zasobów przyrody w Polsce. Zakład Badań Środowiska Rolniczego i Leśnego PAN, Poznań.
- Ryszkowski L., Bałazy S. 1995. Strategia ochrony środowiska i przyrody na obszarach wiejskich. In: L. Ryszkowski, S. Bałazy (eds), *Zasady ekopolityki w rozwoju obszarów wiejskich. Materiały z Sejmu Wyjazdowej Rady Ekologicznej przy Prezydencie Rzeczypospolitej Polskiej Poznań—Leszno 28—30.06. 1994*. Zakład Badań Środowiska Rolniczego i Leśnego PAN, Poznań: 49—64.
- Skirgiełło A. 1965—84. Materiały do rozmieszczenia geograficznego grzybów wyższych w Europie. 1—VI. *Acta Mycol.* 1: 23—26, 3: 243—249, 4: 101—123, 8 (2): 191—218, 12 (2): 155—189, 20 (1): 129—157.
- Skirgiełło A. 1991. Flora Polska. Grzyby (*Mycota*) 20: *Rusalski*. PWN, Warszawa—Kraków.
- Tomiałojć L. 1970. Badania ilościowe nad syntantropijną awifauną Legnicy i okolic. *Acta Orn.* 12.
- Wendlandt J. 1992. Parki i ogrody zabytkowe w Polsce. Ogrody 1. Zarząd Ochrony i Komerwacji Zespołów Pałacowo-Ogrodowych, Warszawa.
- Wojewoda W., Ławrynowicz M. 1986. Red list of threatened macrofungi in Poland. In: K. Zarzycki, W. Wojewoda, Z. Heinrich (eds), *List of threatened plants in Poland*. PWN, Warszawa: 45—52.
- Wojewoda W., Ławrynowicz M. 1992. Red list of threatened macrofungi in Poland. In: K. Zarzycki, W. Wojewoda, Z. Heinrich (eds), *List of threatened plants in Poland*. 2 ed.: 27—56. Pol. Acad. Sci. W. Szafer Inst. Bot. Kraków.

## Grzyby wielkoowocnikowe parku pałacowego w Turwi, koło Poznania

### Streszczenie

Badania prowadzono w latach 1997—99 na terenie parku pałacowego w Turwi (Wielkopolska, Park Krajobrazowy im. gen. D. Chłapowskiego) w celu poznania składu gatunkowego grzybów wielkoowocnikowych tego parku i określenia jego roli jako środowiska refugialnego w krajobrazie rolniczym. Ogółem zebrano 176 gatunków i 1 odmianę grzybów wielkoowocnikowych. Dominowały *Basidiomyces* — 84,7%, a *Ascomycetes* stanowiły 15,3%. 23 gatunki (prawie 13%) to grzyby objęte ochroną gatunkową i/lub umieszczone na czerwonej liście z różnym stopniem zagrożenia. W krajobrazie rolniczym, jako dominujące w Parku Krajobrazowym im. gen. D. Chłapowskiego, park pałacowy w Turwi stanowi ostoję dla wielu gatunków grzybów wielkoowocnikowych.