


Making an impact on kidney disease in people with type 2 diabetes: the importance of screening for albuminuria

Janet B McGill,¹ Hermann Haller,² Prabir Roy-Chaudhury,^{3,4} Andrea Cherrington,⁵ Takashi Wada,⁶ Christoph Wanner,⁷ Linong Ji,⁸ Peter Rossing ^{9,10}

To cite: McGill JB, Haller H, Roy-Chaudhury P, *et al.* Making an impact on kidney disease in people with type 2 diabetes: the importance of screening for albuminuria. *BMJ Open Diab Res Care* 2022;**10**:e002806. doi:10.1136/bmjdr-2022-002806

Received 8 February 2022
Accepted 9 May 2022

ABSTRACT

Albuminuria is useful for early screening and diagnosis of kidney impairment, especially in people with pre-diabetes or type 2 diabetes (T2D), which is the leading cause of chronic kidney disease (CKD) and end-stage kidney disease (ESKD), associated with increased mortality, poor cardiovascular outcomes, and high economic burden. Identifying patients with CKD who are most likely to progress to ESKD permits timely implementation of appropriate interventions. The early stages of CKD are asymptomatic, which means identification of CKD relies on routine assessment of kidney damage and function. Both albuminuria and estimated glomerular filtration rate are measures of kidney function. This review discusses albuminuria as a marker of kidney damage and cardiorenal risk, highlights the importance of early screening and routine testing for albuminuria in people with T2D, and provides new insights on the optimum management of CKD in T2D using albuminuria as a target in a proposed algorithm. Elevated urine albumin can be used to detect CKD in people with T2D and monitor its progression; however, obstacles preventing early detection exist, including lack of awareness of CKD in the general population, poor adherence to clinical guidelines, and country-level variations in screening and treatment incentives. With albuminuria being used as an entry criterion and a surrogate endpoint for kidney failure in clinical trials, and with novel treatment interventions available to prevent CKD progression, there is an urgent need for early screening and diagnosis of kidney function decline in people with T2D or pre-diabetes.

INTRODUCTION

Type 2 diabetes (T2D) is the leading cause of chronic kidney disease (CKD) and end-stage kidney disease (ESKD).¹⁻³ More than one-third of people with T2D also have CKD,⁴ and this population is associated with a 10-fold or greater increase in all-cause mortality compared with T2D alone.⁵ Furthermore, CKD progression leads to ESKD, which is irreversible and fatal in the absence of kidney replacement therapy.⁶ CKD and ESKD are associated with high economic burden,

accounting for 22.3% (US\$81.8 billion) and 7.2% (US\$36.6 billion), respectively, of all Medicare fee-for-service spending in 2018.^{7,8} Medicare expenditures for people with CKD have risen at a rate higher than expenditures for the general Medicare population and have been found costlier for people with CKD and comorbid heart failure or diabetes (type 1 or 2), highlighting clear clinical and economic rationales for early identification and treatment intervention to limit CKD progression in all populations, particularly in people with T2D and cardiovascular risk factors.⁷

Early diagnosis of CKD may prove vital in people with T2D, allowing for more immediate clinical care, and identification of the risks for progression to ESKD. This may be particularly important in Afro-Caribbean/African-American, Hispanic and Asian populations, who tend to have a higher risk of CKD progression relative to White people; the increased risk may be explained in part by factors such as higher blood pressure and increased prevalence of T2D and cardiovascular disease in these populations.⁹ However, diagnosis of CKD is challenging because of the lack of symptoms during its early stages.¹⁰ Other potential factors that hinder early detection of CKD include a lack of awareness of CKD in the general population, poor adherence to clinical guidelines, and the variation in screening and treatment incentives by practice and by country.¹⁰

Diagnosis and determination of the severity of CKD relies on routine measurement of reduced estimated glomerular filtration rate (eGFR; <60 mL/min/1.73 m²) for at least 3 months and/or persistently elevated albuminuria (urine albumin-to-creatinine ratio (UACR)), to assess the overall kidney function and the presence of kidney damage, respectively.^{12,11} As such, the Kidney Disease: Improving Global Outcomes



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Janet B McGill;
jmcgill@wustl.edu

				Persistent albuminuria categories		
				Description and range		
				A1	A2	A3
				Normal to mildly increased	Moderately increased	Severely increased
				<30 mg/g <3 mg/mmol	30-300 mg/g 3-30 mg/mmol	>300 mg/g >30 mg/mmol
				GFR categories (mL/min/1.73 m ²) Description and range	G1	Normal or high
G2	Mildly decreased	60-89				
G3a	Mildly to moderately decreased	45-59				
G3b	Moderately to severely decreased	30-44				
G4	Severely decreased	15-29				
G5	Kidney failure	<15				

Figure 1 Prognosis of CKD by GFR and albuminuria categories. Green: low risk (if no other markers of kidney disease, no CKD); yellow: moderately increased risk; orange: high risk; red: very high risk. Reproduced with permission from Kidney Disease: Improving Global Outcomes (KDIGO) 2020 Clinical Practice Guideline for Diabetes Management in Chronic Kidney Disease.¹ A, albuminuria category; CKD, chronic kidney disease; G, GFR category; GFR, glomerular filtration rate.

(KDIGO) CKD working group uses a combination of eGFR and UACR categories in its risk stratification tool for predicting CKD outcomes (figure 1).² Clinical guidelines in Europe, USA and other countries recommend yearly screening of albuminuria and eGFR in people with T2D.^{11 12} However, real-world data suggest that rates of albuminuria testing in clinical practice are suboptimal,⁴ resulting in a gross underestimation of the risk of CKD in people with T2D. Routine assessment of both screening for CKD prognosis and management was further emphasized in a recent systematic review that showed an association between the prevalence of comorbidities in CKD and an increase in albuminuria severity.¹³ While CKD is common, only a small proportion of patients were classified as high risk or very high risk according to the KDIGO classification, highlighting key gaps in the burden and outcomes of CKD defined by the KDIGO 2012 guideline.¹³ With novel treatment interventions now available that can slow CKD progression, there is an urgent need for the early screening and diagnosis of kidney function decline in people with T2D. To this end, this review discusses albuminuria as a marker of kidney damage and cardiorenal risk, and the importance of earlier and more frequent screening.

ALBUMINURIA AS A MARKER OF KIDNEY DAMAGE AND CARDIORENAL RISK

UACR can be easily measured by spot urine samples, preferably obtained from the first morning void.² Despite strong recommendations to screen for UACR in patients

with T2D, studies show testing for UACR is underused compared with eGFR testing: the Awareness, Detection and Drug Therapy in Type 2 Diabetes and Chronic Kidney Disease study reports 85% of patients undergo eGFR testing versus 47% with UACR determination¹⁴; similar patterns for underutilization of eGFR and UACR testing were noted in the Center for Kidney Disease Research, Education, and Hope (CURE-CKD) registry of people with T2D at risk for CKD.^{15 16} Addressing this discord in eGFR and UACR testing is important because detectable increases in albuminuria generally occur before a decline in eGFR and early changes could indicate signs of kidney disease.^{17 18} Similarly, albuminuria has been associated with an elevated risk of hospitalization in older adults with and without diabetes (type 1 or 2), further emphasizing the importance of albuminuria measurement as a marker.¹⁹

As well as being useful for the early detection of CKD in people with T2D, elevated UACR levels can also be used to monitor CKD progression.² The ability of albuminuria to predict the progression of kidney disease is irrespective of eGFR,²⁰ with the relative risks of ESKD, acute kidney injury, and progressive CKD increasing with elevating levels of albuminuria.^{17 20 21} In general, lower eGFR and higher levels of albuminuria (>300 mg/g) independently predict faster progression to ESKD among patients with stage 3 CKD (eGFR 30–60 mL/min/1.73 m²).²⁰ While kidney function decline is a continuum, recognized albuminuria categories in

CKD include: A1 (normal to mildly increased albuminuria), defined as UACR <30 mg/g; A2 (moderately increased albuminuria), defined as UACR 30–300 mg/g; and A3 (severely increased albuminuria), defined as UACR >300 mg/g (figure 1).²

Large interindividual variability in kidney decline has been documented. The UK Prospective Diabetes Study found that 38% of people with T2D developed albuminuria and 29% developed renal impairment, with many developing one outcome but not the other.²² Thus, some patients with T2D may have CKD without the presence of albuminuria,²² suggesting that the assessment of albuminuria alone may not optimally identify people with T2D who are at high risk of kidney impairment. An additional challenge in screening for CKD in people with T2D is the occurrence of hyperfiltration with normal creatinine/eGFR and without albuminuria. Results from a large observational study indicate that in T2D, both hyperfiltration and high-normal eGFR levels are associated with increased risk of mortality, independent of confounders that may directly impact mortality or GFR estimation.²³ Routine assessment of both albuminuria and eGFR is warranted in these patients. Also, it is important to note that elevated albumin in the urine may be indicative of other disorders. Albuminuria is the earliest marker of other glomerular diseases, and may be elevated in hypertensive nephrosclerosis; it is also associated with hypertension, obesity, and vascular disease.² Furthermore, exercise within 24 hours of albuminuria assessment, infection, fever, congestive heart failure, marked hyperglycemia, menstruation, and marked

hypertension may elevate the UACR independent of kidney damage.¹¹

While elevated albuminuria can be indicative of CKD, incremental increases in albuminuria are associated with corresponding increases in cardiorenal risk.²⁴ Concomitant CKD and T2D is an independent risk factor for cardiovascular events, with this population being approximately three times more likely to die from cardiovascular-related causes than those with T2D alone, and almost six times more likely to die from cardiovascular-related causes than those with neither T2D nor CKD.⁵ Based on a report from the Framingham Heart Study, the presence of albuminuria is associated with a higher risk of incident heart failure, specifically heart failure with reduced ejection fraction, in the general population. In patients with T2D and kidney disease, patients with an increased level of baseline albuminuria had a higher risk for heart failure compared with those with low albuminuria. In addition, the Reduction of Endpoints in NIDDM with the Angiotensin II Antagonist Losartan (RENAAL) study showed that reduction in albuminuria by 50% was associated with reduced risks of cardiovascular events by 18% and heart failure by 27%.²⁴ People with T2D and moderately to severely increased albuminuria have an approximately threefold, fourfold, and fivefold greater risk of myocardial infarction, cardiovascular death and hospitalization for heart failure, respectively, than those with T2D alone.²⁵ A recent study (n=512 700) showed that early CKD in people with diabetes may shorten life expectancy by 16 years, compared with 6 years for early CKD without diabetes




Figure 2 Life expectancy in people with early CKD, T2D, and early CKD in T2D. The reference group consisted of participants with neither diabetes nor CKD. Early CKD was defined as CKD stages 1–3 without diabetes. Diabetes was defined as T2D without CKD. Early CKD in T2D was defined as diabetes with early CKD stages 1–3. Reproduced with permission from Kidney International.²⁶ CKD, chronic kidney disease; T2D, type 2 diabetes.

(either type 1 or type 2), and 10 years for diabetes alone (either type 1 or type 2; [figure 2](#)).²⁶ Patients with high albuminuria may die of cardiovascular events before developing ESKD.²⁰

TREATMENT AND PREVENTION OF CKD

The ability to identify patients with CKD most likely to progress to ESKD would permit timely implementation of appropriate interventions. Progression of CKD poses a significant impact on the health and quality of life of affected patients,^{27 28} as well as a large financial burden.⁷ Efforts to improve care of patients with CKD should focus on reducing the incidence of kidney failure (often defined as a composite of outcomes including sustained low/declining GFR and clinical outcomes of transplantation, dialysis or death from kidney failure), and should provide more affordable treatment options, as well as improving access to kidney transplantation and home dialysis.^{27 28}

Early intensive glycemic control can reduce the risk of developing high albuminuria by one-third compared with the standard of care in people with type 1 diabetes²⁹ and significantly reduce the rate of progression from normoalbuminuria to microalbuminuria in newly diagnosed T2D.³⁰ Current pharmacologic recommendations for managing CKD in T2D include reducing hyperglycemia and hypertension, reducing cardiovascular risk through management of dyslipidemia with statin therapy, using an ACE inhibitor or angiotensin receptor blocker (ARB) and most recently, using sodium-glucose co-transporter-2 inhibitors (SGLT2is) or glucagon-like peptide-1 receptor agonists (GLP-1RAs).² ACE inhibitors and ARBs are not recommended for the primary prevention of CKD; however, they are recommended as separate treatments to help preserve kidney function in all people with T2D and persistent albuminuria (ie, persistent UACR >30 mg/g).¹² ACE inhibitors and ARBs are potent antihypertensive agents that counter the vasoconstrictive effects of angiotensin II, causing selectively greater vasodilatation of the efferent arterioles of the glomeruli, thereby reducing intraglomerular pressure.¹ ACE inhibitors and ARBs have also been shown to provide kidney protection independent of blood pressure control.^{31 32}

SGLT2is and GLP-1RAs have demonstrated reduced risk of CKD progression in people with T2D in several clinical trials.^{33–38} Improved kidney outcomes in people with T2D, eGFR <60 mL/min/1.73 m², and either established cardiovascular disease or cardiovascular risk factors,³⁷ and reduced risk of kidney failure and cardiovascular events in patients with CKD regardless of T2D status,^{38 39} have resulted in SGLT2is being the recommended first-line glucose-lowering medication, with or without metformin and regardless of glucose control levels in people with CKD and T2D when eGFR is ≥30 mL/min/1.73 m² and UACR is >300 mg/g.^{1 40}

A proposed mechanism for SGLT2is is improving the intrarenal hemodynamics, which protects the glomeruli from high-pressure damage. This occurs via

tubuloglomerular feedback from the macula densa, leading to constriction of the afferent arteriole or dilation of the efferent arteriole.⁴¹ Novel approaches to slowing the progression of CKD in people with T2D involve the pharmacologic blockade of mineralocorticoid receptor (MR) overactivation in the kidney.⁴² MR overactivation is known to exacerbate inflammation and fibrosis in the kidneys, the heart, and the vascular system.⁴² Finerenone, a novel, selective, non-steroidal, MR antagonist that blocks MR overactivation, has demonstrated significant dose-dependent reductions in albuminuria in people with T2D and CKD,⁴³ as well as significantly reducing the risk of kidney failure, sustained ≥40% decline in eGFR, or renal death (composite endpoint), and other cardiovascular outcomes in people with CKD and T2D in Finerenone in Reducing Kidney Failure and Disease Progression in Diabetic Kidney Disease (FIDELIO-DKD).⁴⁴ In addition, UACR reductions were demonstrated with finerenone versus placebo (−34.7% vs −4.7%).⁴⁴ Finerenone in Reducing Cardiovascular Mortality and Morbidity in Diabetic Kidney Disease (FIGARO-DKD), a recently completed successful companion phase III trial to FIDELIO-DKD, met its composite primary endpoint of cardiovascular death and non-fatal cardiovascular events versus placebo in people with CKD and T2D.^{44 45} Despite also showing reduced albuminuria in people with T2D and CKD,¹ the steroidal MR antagonists spironolactone and eplerenone are currently underused because of difficulties in managing severe hyperkalemia and worsening of kidney function.⁴⁶

Ethnic minority representation varies across these clinical trials but is consistently low compared with Caucasian representation; this is not CKD-specific but a global and societal issue that needs to be addressed across all forms of healthcare. For example, African Americans account for 35% of people with CKD,⁴⁷ yet in large CKD studies such as Canagliflozin and Renal Events in Diabetes with Established Nephropathy Clinical Evaluation (CREDENCE), Dapagliflozin and Prevention of Adverse Outcomes in Chronic Kidney Disease (DAPA-CKD), and FIDELIO-DKD, these patients accounted for 5%, 4%, and 5% of the trial population, respectively.^{38 39 44}

Assessment of albuminuria is an important component of entry criteria for clinical trials, and its use as a predictor of efficacy should be assessed. Clinical trials in people with CKD and T2D typically include UACR measurements as part of the study entry criteria; however, criteria often vary hugely between studies.^{38 39 43 44} A recent meta-analysis supported the role of albuminuria as a surrogate endpoint in clinical trials of progression of CKD, with a reduction in albuminuria of >30% associated with substantial reductions in the risk of ESKD.⁴⁸ Proteinuria may contribute to worsening kidney function by overloading the tubular epithelial cells leading to intrarenal activation of complement, resulting in interstitial inflammation predominantly mediated by macrophages and sustained fibrogenesis.⁴⁹ As such, remission of albuminuria (defined as reversal of UACR

to normoalbuminuria or <30 mg/g creatinine and a reduction in UACR by $\geq 30\%$ from baseline at two consecutive time points) was used as the primary endpoint in the Esaxerenone (CS-3150) in Patients with Type 2 Diabetes and Microalbuminuria trial, which investigated the use of esaxerenone, a non-steroidal MR antagonist.⁵⁰ Despite evidence for the utility of albuminuria, regulatory authorities place more emphasis on GFR thresholds over UACR, currently accepting a 30%–40% GFR decline as a surrogate endpoint for kidney failure in clinical trials of kidney disease progression. However, this may not be appropriate for drugs targeted at early stages of CKD or with potential hemodynamic effects.¹⁷

ALBUMINURIA AS A TARGET FOR INTERVENTION

Microalbuminuria (UACR <300 mg/g) has been reported in association with impaired fasting glucose and impaired glucose tolerance,⁵¹ with a prevalence of 15.5% in people with pre-diabetes.⁵¹ Correlation between the presence of microalbuminuria in people with pre-diabetic conditions and progression towards T2D suggests that screening of albuminuria in the general population may be beneficial for the early prevention of kidney damage.⁵¹

Reducing albuminuria may slow progression of CKD and should, therefore, be considered as a separate target for kidney-protective therapy. A meta-analysis of data from

41 clinical trials that included participants with diabetes (71%), glomerular disease (4.4%) and other CKD (25.2%) demonstrated that decreasing albuminuria is associated with a reduced risk for developing the clinical endpoint (a composite of ESKD, doubling of serum creatinine concentration, or eGFR <15 mL/min/1.73 m²), predictive of CKD progression.⁴⁸ This is in addition to monitoring cardiovascular risks using well-established surrogate endpoints, such as glycated hemoglobin for glucose lowering, blood pressure for control of hypertension, and low-density lipoprotein cholesterol for efficacy of lipid lowering.⁵²

Optimum patient management requires regular screening of people with T2D to ensure early identification of CKD, followed by targeted intervention to reduce albuminuria (ie, with the use of SGLT2is and GLP-1RAs) and stabilize eGFR, along with continued regular monitoring.^{1,53} A proposed algorithm for the optimum management of CKD in T2D using albuminuria as a target is presented in figure 3. However, barriers to albuminuria testing exist, including confusion caused by the similarity of test names and results. For example, the typical range of results for UACR and urine albumin (UALB) is similar, but the UACR test measures albumin content relative to creatine (in units of mg/g) while the UALB test measures total albumin in urine (in units of mg/dL).⁵⁴ Other barriers include a lack of understanding of testing guidelines and




Figure 3 Proposed algorithm for the optimum management of CKD in T2D using albuminuria as a target.^{2,65} †Exercise (strenuous exercise within 24 hours of sample collection), infection (septicemia or other conditions increasing vascular permeability, symptomatic urinary tract infection), fever, congestive heart failure, marked hyperglycemia, menstruation, upright posture (orthostatic proteinuria causing transient elevation in UACR), and marked hypertension may elevate UACR independent of kidney damage. A, albuminuria category; ACEi, ACE inhibitor; ARB, angiotensin receptor blocker; CKD, chronic kidney disease; DPP-4i, dipeptidyl peptidase-4 inhibitor; eGFR, estimated glomerular filtration rate; G, GFR category; GLP-1RA, glucagon-like peptide-1 receptor agonist; RAS, renin–aldosterone system; SGLT2i, sodium-glucose co-transporter-2 inhibitor; T2D, type 2 diabetes; UACR, urine albumin-to-creatinine ratio.

Table 1 Methods for testing albuminuria as recommended by KDIGO and ADA guidelines^{2 11 54 57}

Method	Urine albumin-to-creatinine ratio; screening*†	Urine albumin (24-hour urine albumin test)	Urine dipsticks sensitive for albuminuria‡	Total protein measurements (urine protein-to-creatinine ratio)§
Pros	<ul style="list-style-type: none"> ▶ Guideline-preferred method. ▶ Capable of specific and precise quantification at low concentrations and of producing quantitative results over the clinically relevant range. ▶ Provides a more specific and sensitive measure of changes in glomerular permeability than urinary total protein. 	<ul style="list-style-type: none"> ▶ Capable of specific and precise quantification at low concentrations, and of producing quantitative results over the clinically relevant range. 	<ul style="list-style-type: none"> ▶ Convenient. ▶ Less expensive. 	<ul style="list-style-type: none"> ▶ Less expensive.
Cons	<ul style="list-style-type: none"> ▶ Measurement may be influenced by several factors;¶ therefore, repeat tests are required, with 2 of 3 abnormal measurements reported within a period of 3–6 months before a patient is considered to have high albuminuria (figure 3). 	<ul style="list-style-type: none"> ▶ Method may be burdensome and time consuming and adds little to prediction or accuracy. ▶ Measurement may be influenced by several factors;¶ therefore, repeat tests are required, with 2 of 3 abnormal measurements reported within a period of 3–6 months before a patient is considered to have high albuminuria (figure 3). 	<ul style="list-style-type: none"> ▶ Method of measuring albumin alone. <ul style="list-style-type: none"> – Measurement of albumin alone without simultaneously measuring urine creatinine is susceptible to false-negative and false-positive determinations as a result of variation in urine concentration due to hydration. – Simultaneous measurement of creatinine allows more accurate results. 	<ul style="list-style-type: none"> ▶ Do not give equal analytical specificity and sensitivity for all proteins that can contribute to diverse estimates of proteinuria prevalence. ▶ Not as specific or sensitive a measure of changes in glomerular permeability as UACR, thus not as accurate in detecting low concentrations of albumin indicative of the early stages of kidney damage or kidney disease progression.

*Screening, as measured by a spot urine sample, is done annually; however, patients with UALB >30 mg/g creatinine or an eGFR <60 mL/min/1.73 m² should be monitored twice annually to guide therapy.¹¹

†Albumin is measured using immunologic assays by diagnostic laboratories and albumin concentration reported as a ratio to urinary creatinine concentration (mg/mmol or mg/g).¹¹

‡Urine dipstick test results may vary depending on the manufacturer and are affected by the urine pH. The results are also operator dependent and may be affected by colored compounds in urine (ie, bilirubin and drugs such as ciprofloxacin, chloroquine). This test cannot reliably distinguish between proteinuria categories and has a low diagnostic accuracy for proteinuria detection.²

§Most laboratories use either turbidimetric or colorimetric methods to measure total protein.²

¶Exercise (strenuous exercise within 24 hours of sample collection), infection (septicemia or other conditions increasing vascular permeability, symptomatic urinary tract infection), fever, congestive heart failure, marked hyperglycemia, menstruation, upright posture (orthostatic proteinuria causing transient elevation in UACR), and marked hypertension may elevate UACR independent of kidney damage.¹¹

ADA, American Diabetes Association; eGFR, estimated glomerular filtration rate; KDIGO, Kidney Disease: Improving Global Outcomes; UACR, urine albumin-to-creatinine ratio; UALB, urine albumin.

which tests to order; infrequent physician visits by patients or avoidance of treatment; and methodologic issues, such as urine collection instructions, requirement for specific urine cups, or an inadequate patient recall system when patients fail to provide a urine sample.^{54 55} The guideline-preferred method of screening is annual testing of albuminuria using the accurate and cost-effective UACR test (table 1).^{2 11 56}

Methods of measuring albuminuria are not standardized; however, KDIGO guidelines recommend

standardization against a serum-based calibrant (CRM 470), with albumin measured using immunologic assays capable of specific and precise quantification at low concentrations and of producing quantitative results over the clinically relevant range.² Alternate methods, such as a 24-hour UALB test, are available; however, this test is more burdensome (table 1).^{2 11} The use of urine dipsticks offers a convenient, cost-effective method of measuring albuminuria alone (table 1).^{2 11} However, the use of urine

dipsticks is susceptible to false results with poor sensitivity for UACR ≥ 30 mg/g detection (~62%; 95% CI 50.9 to 72.0).^{2 57 58} Combining urine dipstick measurements for albumin with simultaneous measurement of creatinine could ensure a higher level of accuracy as this will correct for variations in albumin concentration that is highly dependent on hydration status.¹¹ In addition, assays that measure total protein, such as colorimetric protein assays, may be used.^{2 11} However, it is important to note that UALB measurements provide a more specific and sensitive measure of changes in glomerular permeability than urinary total protein, and they enable early detection of low concentrations of albumin as a marker of kidney damage and the progression of kidney disease (table 1).² Therefore, as kidney function declines, it is recommended that testing be carried out more frequently,² with repeat testing of albuminuria in addition to eGFR measurements recommended to assess disease progression or response to treatment (figure 3).²

DISCUSSION AND CONCLUSIONS

Albuminuria is a key marker of kidney damage and cardiorenal risk, and it is important to perform periodic tests for albuminuria in people with T2D, both to detect CKD and to monitor its progression. The CREDENCE (CKD and T2D population) and DAPA-CKD (CKD and mixed T2D and non-T2D population) trials of SGLT2is have shown that the presence of high albuminuria at baseline is predictive of elevated risk for future hard kidney outcomes;^{38 39} the Heart Outcomes Prevention Evaluation (HOPE) study identified albuminuria as a continuous risk factor for all-cause mortality and cardiovascular mortality,⁵⁹ and in a meta-analysis of clinical trial data, lowering of albuminuria with SGLT2is correlated with delayed onset of hard kidney outcomes such as kidney replacement therapy.^{60 61} An increased focus on routine systematic testing for albuminuria in clinical practice is important to address the current suboptimal rates and overcome existing barriers for albuminuria testing. There has been a long-standing perception of no utility in screening for CKD. For decades, there were no interventions apart from ACE inhibitors and ARBs; often prescribed for hypertension or heart failure that could address a decline in kidney function. However, this is changing with an evolving landscape: SGLT2is are now indicated for CKD in people with T2D⁶² and positive outcomes are being reported for new agents that specifically target kidney damage (eg, non-steroidal MR antagonists). The KDIGO and American Diabetes Association guidelines now recommend that people with CKD and T2D, and an eGFR ≥ 30 mL/min/1.73 m² would benefit from the use of SGLT2is, with or without metformin; GLP-IRAs recommended as a preferred additional therapy based on patient preferences, comorbidities, cost, and eGFR status¹ because these therapies appear to reduce the risk of CKD progression, cardiovascular events, and hypoglycemia.¹¹

The importance of education and awareness around albuminuria screening is underscored by the launch of a new public awareness initiative by the US Department of Health and Human Services, the National Kidney Foundation, and the American Society of Nephrology, which aim to provide education about the risks of CKD and promote the early detection, treatment and management of CKD to improve patient outcomes.⁶³ Furthermore, a majority of CKD burden is concentrated in geographies with the lowest sociodemographic index,⁶⁴ where early detection using cost-effective methodologies and early intervention could have a significant impact on reducing disease burden.

We strongly believe that the time is right for a far more aggressive and interventional approach to CKD in T2D that goes all the way from education and awareness to appropriate and regular screening and follow-up. This is particularly relevant in view of both the recent executive order on kidney diseases and the availability, for the first time, of novel therapies that could effectively prevent the progression of CKD.

Author affiliations

¹Division of Endocrinology, Metabolism and Lipid Research, Washington University School of Medicine, Saint Louis, Missouri, USA

²Department of Nephrology, Hannover Medical School, Hannover, Germany

³Division of Nephrology and Hypertension, Department of Medicine, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, North Carolina, USA

⁴Department of Medicine, WG Hefner VA Medical Center, Salisbury, North Carolina, USA

⁵Division of Preventive Medicine, Department of Medicine, The University of Alabama at Birmingham, Birmingham, Alabama, USA

⁶Department of Nephrology and Laboratory Medicine, Kanazawa University, Kanazawa, Ishikawa, Japan

⁷Division of Nephrology and Hypertension, University Hospital of Würzburg, Würzburg, Germany

⁸Department of Endocrinology and Metabolism, Peking University People's Hospital, Beijing, China

⁹Steno Diabetes Center Copenhagen, Herlev, Denmark

¹⁰Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

Acknowledgements Medical writing assistance was provided by Cindy Jenner, PhD, of Chameleon Communications International, and was funded by Bayer AG.

Contributors JBM and PR conceived the manuscript, approved the outline, edited outlines and all drafts and approved the final manuscript content. HH, PR-C, AC, TW, CW, LJ added expertise, edited the outline and drafts and approved the final draft. JBM is the guarantor of the work.

Funding This study was funded by Bayer.

Competing interests JBM reports non-financial support from Boehringer Ingelheim; personal fees from Bayer, Boehringer Ingelheim, MannKind, Novo Nordisk and Provention Bio; grants from Beta Bionics, Dexcom, Medtronic, and Novo Nordisk. PR-C is a consultant/advisor for Akebia, Bayer, Becton Dickinson, Cormedix, Humacyte, Medtronic, Vifor-Relypsa, and WL Gore. He is also the founder and CSO of Inovasc. AC is a consultant/advisor for Bayer. LJ reports receiving consulting and lecture fees from Bayer, Boehringer Ingelheim, Bristol Myers Squibb, Eli Lilly, Merck, Novartis, Novo Nordisk, Roche, Sanofi, and Takeda; and research grants from Roche and Sanofi. HH reports personal fees from Alexion Pharma, AstraZeneca, Bayer, Boehringer Ingelheim, and Vifor Pharma, outside the submitted work. TW reports grants, personal fees and other from Kyowa Kirin; grants and personal fees from Astellas Pharma, Daiichi Sankyo, Kissei Pharmaceutical, Mitsubishi Tanabe Pharma, Sanofi, and Takeda Pharmaceutical; grants from Chugai Pharmaceutical and MSD; and personal fees from AstraZeneca, Bayer, Boehringer Ingelheim, Eli Lilly Japan, Kowa, Nippon, Ono Pharmaceutical, Sanwa Chemistry,

and Taisho Pharma, outside the submitted work. CW reports consultancy and/or speaking fees from AstraZeneca, Bayer, Boehringer Ingelheim, Gilead, GSK, Eli Lilly, MSD, and Mundipharma; and research grants from Boehringer Ingelheim and Sanofi Genzyme outside this body of work. PR reports lecture fees from AstraZeneca, Bayer, Boehringer Ingelheim, Merck, and Novo Nordisk; consultancy fees from AbbVie, Astellas, AstraZeneca, Bayer, Boehringer Ingelheim, Eli Lilly, Gilead, Mundipharma, and Novo Nordisk, with all fees given to Steno Diabetes Center Copenhagen; and equity interest in Novo Nordisk.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. The data supporting the findings of this study are not currently available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Peter Rossing <http://orcid.org/0000-0002-1531-4294>

REFERENCES

- Kidney Disease: Improving Global Outcomes (KDIGO) Diabetes Work Group. KDIGO 2020 clinical practice guideline for diabetes management in chronic kidney disease. *Kidney Int* 2020;98:S1–115.
- Kidney Disease: Improving Global Outcomes (KDIGO) Diabetes Work Group. KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl* 2013;3:1–150.
- Alicic RZ, Rooney MT, Tuttle KR. Diabetic kidney disease: challenges, progress, and possibilities. *Clin J Am Soc Nephrol* 2017;12:2032–45.
- Stempniewicz N, Vassalotti JA, Cuddeback JK, et al. Chronic kidney disease testing among primary care patients with type 2 diabetes across 24 U.S. health care organizations. *Diabetes Care* 2021;44:2000–9.
- Afkarian M, Sachs MC, Kestenbaum B, et al. Kidney disease and increased mortality risk in type 2 diabetes. *J Am Soc Nephrol* 2013;24:302–8.
- Abbasi M, Chertow G, Hall Y. End-stage renal disease. *Am Fam Physician* 2010;82:1512.
- United States Renal Data System. Chapter 6: healthcare expenditures for persons with CKD. USRDS annual data report volume 1: chronic kidney disease. Bethesda, MD: National Institutes of Health, National Institute of diabetes and digestive and kidney diseases, 2020. Available: <https://adr.usrds.org/2020/chronic-kidney-disease/6-healthcare-expenditures-for-persons-with-ckd> [Accessed 15 Nov 2021].
- United States Renal Data System. Chapter 9: healthcare expenditures for persons with ESRD. USRDS annual data report volume 2: end stage renal disease. Bethesda, MD: National Institutes of Health, National Institute of diabetes and digestive and kidney diseases, 2020. Available: <https://adr.usrds.org/2020/end-stage-renal-disease/9-healthcare-expenditures-for-persons-with-esrd> [Accessed 15 Nov 2021].
- Hounkpatin HO, Fraser SDS, Honney R, et al. Ethnic minority disparities in progression and mortality of pre-dialysis chronic kidney disease: a systematic scoping review. *BMC Nephrol* 2020;21:217.
- Plantinga LC, Tuot DS, Powe NR. Awareness of chronic kidney disease among patients and providers. *Adv Chronic Kidney Dis* 2010;17:225–36.
- American Diabetes Association. 11. Microvascular Complications and Foot Care: *Standards of Medical Care in Diabetes-2020*. *Diabetes Care* 2020;43:S135–51.
- Cosentino F, Grant PJ, Aboyans V. ESC guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD. *Eur Heart J* 2019;2020:255–323.
- Murton M, Goff-Leggett D, Bobrowska A, et al. Burden of chronic kidney disease by KDIGO categories of glomerular filtration rate and albuminuria: a systematic review. *Adv Ther* 2021;38:180–200.
- Szczeczek LA, Stewart RC, Su H-L, et al. Primary care detection of chronic kidney disease in adults with type-2 diabetes: the ADD-CKD study (awareness, detection and drug therapy in type 2 diabetes and chronic kidney disease). *PLoS One* 2014;9:e110535.
- Tuttle KR, Alicic RZ, Duru OK, et al. Clinical characteristics of and risk factors for chronic kidney disease among adults and children: an analysis of the CURE-CKD registry. *JAMA Netw Open* 2019;2:e1918169.
- Norris KC, Duru OK, Alicic RZ, et al. Rationale and design of a multicenter chronic kidney disease (CKD) and at-risk for CKD electronic health records-based registry: CURE-CKD. *BMC Nephrol* 2019;20:416.
- Levey AS, Gansevoort RT, Coresh J, et al. Change in albuminuria and GFR as end points for clinical trials in early stages of CKD: a scientific workshop sponsored by the National kidney Foundation in collaboration with the US food and drug administration and European medicines Agency. *Am J Kidney Dis* 2020;75:84–104.
- Coresh J, Levey AS. A combination of change in albuminuria and GFR as a surrogate end point for progression of CKD. *Clin J Am Soc Nephrol* 2019;14:792–4.
- Barzilay JI, Buzkova P, Shlipak MG, et al. Hospitalization rates in older adults with albuminuria: the cardiovascular health study. *J Gerontol A Biol Sci Med Sci* 2020;75:2426–33.
- Bakris GL, Molitch M. Microalbuminuria as a risk predictor in diabetes: the continuing saga. *Diabetes Care* 2014;37:867–75.
- Heerspink HJL, Parving H-H, Andress DL, et al. Atrasentan and renal events in patients with type 2 diabetes and chronic kidney disease (Sonar): a double-blind, randomised, placebo-controlled trial. *Lancet* 2019;393:1937–47.
- Retnakaran R, Cull CA, Thorne KI, et al. Risk factors for renal dysfunction in type 2 diabetes: U.K. prospective diabetes study 74. *Diabetes* 2006;55:1832–9.
- Penno G, Orsi E, Solini A, et al. Renal hyperfiltration is independently associated with increased all-cause mortality in individuals with type 2 diabetes: a prospective cohort study. *BMJ Open Diabetes Res Care* 2020;8:e001481.
- Pafundi PC, Garofalo C, Galiero R, et al. Role of albuminuria in detecting Cardio-Renal risk and outcome in diabetic subjects. *Diagnostics* 2021;11:290.
- Scirica BM, Mosenzon O, Bhatt DL, et al. Cardiovascular outcomes according to urinary albumin and kidney disease in patients with type 2 diabetes at high cardiovascular risk: observations from the SAVOR-TIMI 53 trial. *JAMA Cardiol* 2018;3:155–63.
- Wen CP, Chang CH, Tsai MK, et al. Diabetes with early kidney involvement may shorten life expectancy by 16 years. *Kidney Int* 2017;92:388–96.
- U.S. Department of Health and Human Services. Advancing American kidney health, 2019. Available: <https://aspe.hhs.gov/system/files/pdf/262046/AdvancingAmericanKidneyHealth.pdf> [Accessed 15 Nov 2021].
- Mendu ML, Weiner DE. Health policy and kidney care in the United States: core curriculum 2020. *Am J Kidney Dis* 2020;76:720–30.
- Diabetes Control and Complications (DCCT) Research Group. Effect of intensive therapy on the development and progression of diabetic nephropathy in the diabetes control and complications trial. The diabetes control and complications (DCCT) Research Group. *Kidney Int* 1995;47:1703–20.
- UK Prospective Diabetes Study (UKPDS) Group. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK prospective diabetes study (UKPDS) group. *Lancet* 1998;352:837–53.
- Heart Outcomes Prevention Evaluation Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the hope study and MICRO-HOPE substudy. heart outcomes prevention evaluation study Investigators. *Lancet* 2000;355:253–9.
- Lewis EJ, Hunsicker LG, Clarke WR, et al. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med* 2001;345:851–60.
- Marso SP, Bain SC, Consoli A, et al. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2016;375:1834–44.
- Perkovic V, de Zeeuw D, Mahaffey KW, et al. Canagliflozin and renal outcomes in type 2 diabetes: results from the canvas program randomised clinical trials. *Lancet Diabetes Endocrinol* 2018;6:691–704.
- Marso SP, Daniels GH, Brown-Frandsen K, et al. Liraglutide and cardiovascular outcomes in type 2 diabetes. *N Engl J Med* 2016;375:311–22.
- Tuttle KR, Lakshmanan MC, Rayner B, et al. Dulaglutide versus insulin glargine in patients with type 2 diabetes and moderate-to-severe chronic kidney disease (AWARD-7): a multicentre, open-label, randomised trial. *Lancet Diabetes Endocrinol* 2018;6:605–17.

- 37 Lo KB, Gul F, Ram P, *et al*. The effects of SGLT2 inhibitors on cardiovascular and renal outcomes in diabetic patients: a systematic review and meta-analysis. *Cardiorenal Med* 2020;10:1–10.
- 38 Perkovic V, Jardine MJ, Neal B, *et al*. Canagliflozin and renal outcomes in type 2 diabetes and nephropathy. *N Engl J Med* 2019;380:2295–306.
- 39 Heerspink HJL, Stefánsson BV, Correa-Rotter R, *et al*. Dapagliflozin in patients with chronic kidney disease. *N Engl J Med* 2020;383:1436–46.
- 40 American Diabetes Association. 11. Microvascular complications and foot care: standards of Medical Care in Diabetes-2021. *Diabetes Care* 2021;44:S151–67.
- 41 DeFronzo RA, Reeves WB, Awad AS. Pathophysiology of diabetic kidney disease: impact of SGLT2 inhibitors. *Nat Rev Nephrol* 2021;17:319–34.
- 42 Agarwal R, Kolkhof P, Bakris G, *et al*. Steroidal and non-steroidal mineralocorticoid receptor antagonists in cardiorenal medicine. *Eur Heart J* 2021;42:152–61.
- 43 Bakris GL, Agarwal R, Chan JC, *et al*. Effect of finerenone on albuminuria in patients with diabetic nephropathy: a randomized clinical trial. *JAMA* 2015;314:884–94.
- 44 Bakris GL, Agarwal R, Anker SD, *et al*. Effect of finerenone on chronic kidney disease outcomes in type 2 diabetes. *N Engl J Med* 2020;383:2219–29.
- 45 Bayer. Bayer's finerenone meets primary endpoint in Phase III FIGARO-DKD cardiovascular outcomes study in patients with chronic kidney disease and type 2 diabetes, 2021. Available: <https://media.bayer.com/baynews/baynews.nsf/id/Bayers-finerenone-meets-primary-endpoint-Phase-III-FIGARO-DKD-cardiovascular-outcomes-study-patients> [Accessed 15 Nov 2021].
- 46 Kolkhof P, Jaisser F, Kim S-Y, *et al*. Steroidal and novel non-steroidal mineralocorticoid receptor antagonists in heart failure and cardiorenal diseases: comparison at bench and bedside. *Handb Exp Pharmacol* 2017;243:271–305.
- 47 U.S. Department of Health and Human Services. Race, ethnicity, & kidney disease, 2014. Available: <https://www.niddk.nih.gov/health-information/kidney-disease/race-ethnicity> [Accessed 15 Nov 2021].
- 48 Heerspink HJL, Greene T, Tighiouart H, *et al*. Change in albuminuria as a surrogate endpoint for progression of kidney disease: a meta-analysis of treatment effects in randomised clinical trials. *Lancet Diabetes Endocrinol* 2019;7:128–39.
- 49 Abbate M, Zoja C, Remuzzi G. How does proteinuria cause progressive renal damage? *J Am Soc Nephrol* 2006;17:2974–84.
- 50 Ito S, Kashihara N, Shikata K. Esaxerenone (CS-3150) in patients with type 2 diabetes and microalbuminuria (ESAX-DN). *Clin J Am Soc Nephrol* 2020;15:1715–27.
- 51 Bahar A, Makhlough A, Yousefi A, *et al*. Correlation between prediabetes conditions and microalbuminuria. *Nephrourol Mon* 2013;5:741–4.
- 52 Wieczorek A, Rys P, Skrzekowska-Baran I, *et al*. The role of surrogate endpoints in the evaluation of efficacy and safety of therapeutic interventions in diabetes mellitus. *Rev Diabet Stud* 2008;5:128–35.
- 53 Cherney DZI, Heerspink HJL, Frederich R, *et al*. Effects of ertugliflozin on renal function over 104 weeks of treatment: a post hoc analysis of two randomised controlled trials. *Diabetologia* 2020;63:1128–40.
- 54 MacLean CD, MacCaskey M, Littenberg B. Improving testing for proteinuria in diabetes using decision support: role of laboratory ordering systems. *Lab Med* 2013;44:353–7.
- 55 Hellemons ME, Denig P, de Zeeuw D, *et al*. Is albuminuria screening and treatment optimal in patients with type 2 diabetes in primary care? observational data of the GIANTT cohort. *Nephrol Dial Transplant* 2013;28:706–15.
- 56 Lepore G, Maglio ML, Nosari I, *et al*. Cost-effectiveness of two screening programs for microalbuminuria in type 2 diabetes. *Diabetes Care* 2002;25:2103–4.
- 57 Park JI, Baek H, Kim BR, *et al*. Comparison of urine dipstick and albumin:creatinine ratio for chronic kidney disease screening: A population-based study. *PLoS One* 2017;12:e0171106.
- 58 Sumida K, Nadkarni GN, Grams ME, *et al*. Conversion of Urine Protein-Creatinine Ratio or Urine Dipstick Protein to Urine Albumin-Creatinine Ratio for Use in Chronic Kidney Disease Screening and Prognosis : An Individual Participant-Based Meta-analysis. *Ann Intern Med* 2020;173:426–35.
- 59 Gerstein HC, Mann JF, Yi Q, *et al*. Albuminuria and risk of cardiovascular events, death, and heart failure in diabetic and nondiabetic individuals. *JAMA* 2001;286:421–6.
- 60 Christofides EA, Desai N. Optimal early diagnosis and monitoring of diabetic kidney disease in type 2 diabetes mellitus: addressing the barriers to albuminuria testing. *J Prim Care Community Health* 2021;12:21501327211003683.
- 61 Delanaye P, Wissing KM, Scheen AJ. SGLT2 inhibitors: renal outcomes according to baseline albuminuria. *Clin Kidney J* 2021.
- 62 Janssen Pharmaceuticals Inc. Invokana® (canagliflozin) prescribing information., 2020. Available: <http://www.janssenlabels.com/package-insert/product-monograph/prescribing-information/INVOKANA-pi.pdf> [Accessed 15 Nov 2021].
- 63 National Kidney Foundation. Public awareness initiative for AAKH announced as a public-private partnership with NKF and ASN, 2019. Available: <https://www.kidney.org/news/aakh-announced-public-private-partnership-nkf-asn> [Accessed 15 Nov 2021].
- 64 GBD Chronic Kidney Disease Collaboration. Global, regional, and national burden of chronic kidney disease, 1990-2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2020;395:709–33.
- 65 KDOQI. KDOQI clinical practice guidelines and clinical practice recommendations for diabetes and chronic kidney disease. *Am J Kidney Dis* 2007;49:S12–154.