

Mammographic Density Phenotypes and Risk of Breast Cancer: A Meta-analysis

Andreas Pettersson, Rebecca E. Graff, Giske Ursin, Isabel dos Santos Silva, Valerie McCormack, Laura Baglietto, Celine Vachon, Marije F. Bakker, Graham G. Giles, Kee Seng Chia, Kamila Czene, Louise Eriksson, Per Hall, Mikael Hartman, Ruth M. L. Warren, Greg Hislop, Anna M. Chiarelli, John L. Hopper, Kavitha Krishnan, Jingmei Li, Qing Li, Ian Pagano, Bernard A. Rosner, Chia Siong Wong, Christopher Scott, Jennifer Stone, Gertraud Maskarinec, Norman F. Boyd, Carla H. van Gils, Rulla M. Tamimi

Manuscript received July 23, 2013; revised February 3, 2014; accepted February 24, 2014.

Correspondence to: Andreas Pettersson, MD, PhD, Department of Epidemiology, Harvard School of Public Health, 677 Huntington Avenue, Boston, MA 02115 (e-mail: apetters@hsph.harvard.edu) and Rulla M. Tamimi, ScD, Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, 181 Longwood Avenue, Boston, MA 02115 (e-mail: rulla.tamimi@channing.harvard.edu).

Background Fibroglandular breast tissue appears dense on mammogram, whereas fat appears nondense. It is unclear whether absolute or percentage dense area more strongly predicts breast cancer risk and whether absolute nondense area is independently associated with risk.

Methods We conducted a meta-analysis of 13 case-control studies providing results from logistic regressions for associations between one standard deviation (SD) increments in mammographic density phenotypes and breast cancer risk. We used random-effects models to calculate pooled odds ratios and 95% confidence intervals (CIs). All tests were two-sided with *P* less than .05 considered to be statistically significant.

Results Among premenopausal women (*n* = 1776 case patients; *n* = 2834 control subjects), summary odds ratios were 1.37 (95% CI = 1.29 to 1.47) for absolute dense area, 0.78 (95% CI = 0.71 to 0.86) for absolute nondense area, and 1.52 (95% CI = 1.39 to 1.66) for percentage dense area when pooling estimates adjusted for age, body mass index, and parity. Corresponding odds ratios among postmenopausal women (*n* = 6643 case patients; *n* = 11 187 control subjects) were 1.38 (95% CI = 1.31 to 1.44), 0.79 (95% CI = 0.73 to 0.85), and 1.53 (95% CI = 1.44 to 1.64). After additional adjustment for absolute dense area, associations between absolute nondense area and breast cancer became attenuated or null in several studies and summary odds ratios became 0.82 (95% CI = 0.71 to 0.94; $P_{\text{heterogeneity}} = .02$) for premenopausal and 0.85 (95% CI = 0.75 to 0.96; $P_{\text{heterogeneity}} < .01$) for postmenopausal women.

Conclusions The results suggest that percentage dense area is a stronger breast cancer risk factor than absolute dense area. Absolute nondense area was inversely associated with breast cancer risk, but it is unclear whether the association is independent of absolute dense area.

JNCI J Natl Cancer Inst (2014) 106(5): dju078 doi:10.1093/jnci/dju078

Fibroglandular breast tissue (epithelial cells, fibroblasts, and connective tissue) appears radio-dense on a mammogram, whereas breast fat appears transparent or nondense. Dense mammographic tissue, quantified as either the absolute or percentage dense area on a mammogram, is a strong risk factor for breast cancer. According to one meta-analysis, women with 75% or more dense area are at a 4.6-fold increased risk of breast cancer relative to women with less than 5% dense area (1).

It is possible that women with high mammographic density are at increased risk of breast cancer because they have large amounts of fibroglandular breast tissue at risk (2). In line with this hypothesis, some studies have reported greater magnitudes of breast cancer risk for absolute dense area than percentage dense area (3). Several other studies, however, have reported greater magnitudes of risk for percentage than absolute dense area (1,4). The latter findings

suggest that the ratio of fibroglandular to fatty tissue is a stronger breast cancer risk factor, implying that either the ratio itself reflects an underlying biological mechanism associated with breast cancer risk or that absolute nondense area, which contributes to the denominator of percentage dense area, is inversely associated with breast cancer risk. Two recent nested case-control studies examining the association between absolute nondense area and breast cancer risk yielded conflicting results (5,6). In statistical models including both absolute dense and nondense area, in one study (5) there was a statistically significant lower risk of breast cancer with increasing absolute nondense area, whereas in the other study (6) there was a statistically significant increased risk of breast cancer.

It is critical to establish the relationship between the different mammographic density phenotypes (absolute dense area, absolute nondense area, and percentage dense area) and breast cancer risk to

uncover underlying biological mechanisms and to improve upon breast cancer risk prediction modeling. We thus conducted a meta-analysis of 13 case-control studies that examined the associations between mammographic density phenotypes and risk of breast cancer; we benefitted from a reanalysis of each individual study for the purpose of this meta-analysis such that the study-specific associations were consistently analyzed using the same adjustments and categorizations.

Methods

Invited Studies, Inclusion Criteria, and Participating Studies

Eligible studies for this meta-analysis included those partaking in the DENSE consortium (7), an international collaboration of 19 epidemiological studies with data on breast cancer susceptibility genetic variants and mammographic density, as well as any others led by DENSE consortium principal investigators. To be eligible for inclusion in the meta-analysis, we required that studies used a case-control design (nested or not) and had digitized prediagnostic film mammograms, mammographic density assessed using a computer-assisted thresholding technique, and relevant covariable data measured at the time (within a few years) of mammography. In total, 13 studies provided data for the meta-analysis (Table 1) (5,6,8–17). Twelve were case-control studies nested within existing prospective cohorts, trials, or fully enumerated registries, and one was a population-based case-control study.

Mammographic density studies within the AGE Trial were approved by the UK South East Research Ethics Committee (REC 05/MRE01/77). Approval for the Cancer and Hormones Replacement Study (CAHRES) study was given by the ethical review boards in the respective regions in which the subjects were based in Sweden. The Canadian Breast Density Study (CBDS) study was approved by the ethics committees at the University of Toronto, the University Health Network (Toronto), the Ontario Breast Screening Program (OBSP), and the University of British Columbia. The European Prospective Investigation into Cancer and Nutrition study--Netherlands (EPIC-NL) study was approved by the institutional review board of the University Medical Centre Utrecht. The Melbourne Collaborative Cohort Study (MCCS) study was approved by the Cancer Council Victoria's human research ethics committee. The Mayo Clinic Mammography Study (MCMAM) and Mayo Mammography Health Study (MMHS) studies were approved by the Mayo Clinic Institutional Review Board. The Multiethnic Cohort (MEC) study was approved by the Committee on Human Studies at the University of Hawaii. The Nurses' Health Study 1 (NHS1) and Nurses' Health Study 2 (NHS2) studies were approved by the Committee on the Use of Human Subjects in Research at Brigham and Women's Hospital. The Singapore Breast Cancer Screening Project (SBSP) study was approved by the National University of Singapore Institutional Review Board. The Norwich & Cambridge, UK National Health Service Breast Cancer Screening Program (UK-NHS) study was approved by the Norfolk Local Research Ethics Committee. The University of Southern California (USC) study was approved by the institutional review board at the University of Southern California. Participants of all studies except the UK-NHS provided informed consent. The UK-NHS was a medical records study, so direct consent from the patients was not required.

Mammographic Density Measures

All studies had prediagnostic film mammograms of the craniocaudal (CC; $n = 9$ studies) or mediolateral oblique (MLO; $n = 4$ studies) view (Table 1). [The MCMAM study provided data for both the CC and MLO views; data from the CC view were used in all analyses unless otherwise specified because they have been used in previous publications (16).] Mammographic density was measured using the computer-assisted thresholding technique Cumulus (18) ($n = 12$ studies) or Madena (19) ($n = 1$ study). Using these techniques, two grayscale thresholds are selected on the digitized mammograms. One threshold separates the breast from the background (or alternatively the breast edge is manually delimited), and the other classifies the breast tissue into dense and nondense area. The percentage dense area is calculated as $100 \times (\text{absolute dense area} / (\text{absolute dense area} + \text{absolute nondense area}))$. Trained readers blinded to case-control status read the mammograms and selected the thresholds in all studies.

Statistical Analysis

Study Level. Each study provided study-specific parameter estimates (β s) and standard errors (SEs) from conditional or unconditional logistic regression models conducted specifically for this meta-analysis. Because menopausal status is associated with both mammographic density and breast cancer risk, all analyses were conducted separately for premenopausal and postmenopausal status at the time of mammography. Menopausal status was self-reported in 10 studies and based on age (≤ 50 years premenopausal, > 50 years postmenopausal) at the time of mammography in three studies. Each mammographic density phenotype was divided into quartiles based on its distribution among the control subjects. Quartiles were modeled as categorical variables using the lowest quartile as the reference. Because the distributions of the mammographic density phenotypes differed across studies (Table 1), each study transformed the mammographic density phenotypes so that they were approximately normally distributed and then provided the β s and their standard errors associated with one standard deviation (SD) increments in each mammographic density phenotype. We were thus able to compare the magnitude of the associations between the different mammographic density phenotypes and breast cancer risk.

The following covariables, measured at the time of mammography, were included in the analyses: age (available for all studies; self-reported in all studies), body mass index (BMI; kg/m^2 ; available for 12 studies; self-reported in 7 studies and technician-measured in 5 studies), parity (available for 12 studies; self-reported in all studies), and, among postmenopausal women only, use of hormonal replacement therapy (HRT; ever use available for 11 studies and current use for 9 studies; self-reported in all studies). The covariables age and BMI were modeled as continuous variables, whereas parity (nulliparous, 1–2 children, ≥ 3 children) and both ever and current HRT use (yes, no) were modeled as categorical variables.

For each density phenotype, as detailed in Tables 2 and 3, statistical models were run adjusting 1) for age at mammography, and 2) additionally for BMI and parity (ie, the fully adjusted model for percentage dense area). For dense and nondense area, models were also fitted adjusting 3) for age and the other absolute phenotype (dense or nondense area, as appropriate) and 4) additionally for BMI and parity (ie, the fully adjusted model for absolute dense and nondense area).

Table 1. Characteristics at or around the time of mammography among case patients and control subjects for each study included in the meta-analysis*

Study name	AGE		CAHRES		CBDS		EPIC-NL		MCCS		MCMAM	
	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study
Study type	UK	Sweden	Canada	Netherlands	Australia	US	White	White	White	White	White	White
Country	White	White	White	White	White	White	White	White	White	White	White	White
Ethnicity†	MLO	MLO	CC	MLO	CC	CC (MLO)	MLO	MLO	CC	CC	CC (MLO)	CC (MLO)
Mammogram view‡	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus
Threshold technique	1991–2005	1993–1995	1980–1999	1990–2007	1990–2007	1990–2001	1990–1999	1990–2007	1990–2007	1990–2007	1990–2001	1990–2001
Study period	1988	1986	1984	1989	1991	1991	1989	1991	1991	1991	1991	1991
Year of screening program initiation in the country of study	Both	Both	Both	Both	Both	Both	Both	Both	Both	Both	Both	Both
Screening or interval detected breast cancer	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported
BMI ascertainment	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported
Menopausal status ascertainment	Self-reported	Based on age	Self-reported	Self-reported	Based on age	Measured	Measured	Measured	Measured	Measured	Measured	Measured
Reference	17	9	10	5	16	11	5	16	16	11	11	11
Include carcinoma in situ cases	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
	Cases	Cases	Cases	Cases	Cases	Cases	Cases	Cases	Cases	Cases	Cases	Cases
	Controls	Controls	Controls	Controls	Controls	Controls	Controls	Controls	Controls	Controls	Controls	Controls
Premenopausal												
Number	317	–	292	95	44	58	276	44	60	58	119	119
Age, y, mean	41.1	–	47.3	52.8	46.0	46.5	52.2	46.0	46.0	46.5	47.3	47.3
Absolute dense area, cm ² , mean	54.9	–	47.1	36.9	36.4	43.7 (49.4)	30.0	36.4	26.4	43.7 (49.4)	39.8 (43.4)	39.8 (43.4)
Absolute nondense area, cm ² , mean	77.3	–	68.8	90.4	85.2	67.3 (81.2)	101.6	85.2	103.9	67.3 (81.2)	79.0 (93.0)	79.0 (93.0)
Percentage dense area, %, mean	44.8	–	44.2	30.5	31.9	41.1 (39.0)	25.8	31.9	24.0	41.1 (39.0)	35.2 (33.3)	35.2 (33.3)
BMI, kg/m ² , mean	24.9	–	24.2	25.5	25.6	26.1	25.4	25.6	26.4	26.1	27.2	27.2
Parity, No., mean§	2.1	–	2.5	2.0	2.0	2.5	2.1	2.0	2.0	2.5	2.8	2.8
Postmenopausal												
Number	–	1,511	872	358	380	313	859	380	964	313	558	558
Age, y, mean	–	62.5	59.7	60.1	61.0	63.6	59.1	61.0	61.0	63.6	63.7	63.7
Absolute dense area, cm ² , mean	–	28.5	36.0	22.5	20.5	35.0 (36.0)	20.3	20.5	13.8	35.0 (36.0)	28.5 (30.6)	28.5 (30.6)
Absolute nondense area, cm ² , mean	–	143.1	103.1	127.3	121.0	96.7 (114.0)	119.6	121.0	122.4	96.7 (114.0)	99.9 (115.9)	99.9 (115.9)
Percentage dense area, %, mean	–	18.0	28.7	16.9	16.4	28.8 (25.8)	16.6	16.4	11.6	28.8 (25.8)	24.7 (22.6)	24.7 (22.6)
BMI, kg/m ² , mean	–	25.7	25.2	26.6	27.7	27.9	26.0	27.7	26.5	27.9	27.0	27.0
Parity, No., mean¶	–	2.2	3.0	2.3	3.0	3.2	2.5	3.0	3.0	3.2	3.3	3.3
Ever HRT use, %, mean	–	41.8	45.5	27.7	38.4	26.3	25.3	38.4	33.1	26.3	31.2	31.2
Current HRT use, %, mean	–	36.5	25.6	–	27.6	–	–	27.6	21.8	–	–	–

(Table continues)

Table 1 (Continued).

Study name	MEC	MMHS	NHS1	NHS2	SBSP	UK NHS	USC
Study type	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Nested case-control study	Case-control study
Country	US-Hawaii	US	US	US	Singapore	UK	US
Ethnicity†	Mixed	White	White	White	Asian	White	Mixed
Mammogram view‡	CC	CC	CC	CC	CC	MLO	CC
Threshold technique	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Cumulus	Madena
Study period	1993-2000	2003-2011	1989-2004	1995-2003	1994-1997	1995-2004	1989-1998
Year of screening program initiation in the country of study	1991	1991	1991	1991	2002	1988	1991
Screening or interval detected breast cancer	Both	Both	Both	Both	Both	Both	Both
BMI ascertainment	Self-reported	Measured	Self-reported	Self-reported	Measured	-	Self-reported
Menopausal status ascertainment	Self-reported	Self-reported	Self-reported	Self-reported	Self-reported	Based on age	Self-reported
Reference	8	12	6	6	13	14	15
Include carcinoma in situ cases	Yes	Yes	Yes	Yes	Yes	No	No
	Cases	Cases	Cases	Cases	Cases	Cases	Cases
	Controls	Controls	Controls	Controls	Controls	Controls	Controls
Premenopausal							
Number	59	99	284	172	69	-	287
Age, y, mean	49.1	46.3	48.9	44.4	53.1	-	42.0
Absolute dense area, cm ² , mean	46.7	29.8	65.2	107.1	74.1	-	51.9
Absolute nondense area, cm ² , mean	67.4	106.2	117.8	131.3	78.2	-	91.4
Percentage dense area, %, mean	46.8	24.4	44.9	48.2	51.6	-	42.6
BMI, kg/m ² , mean	24.6	27.1	27.9	24.9	24.8	-	24.7
Parity, No., mean§	1.5	2.4	2.4	1.7	2.3	-	1.6
Postmenopausal							
Number	548	349	903	37	400	634	338
Age, y, mean	62.8	64.3	63.7	61.9	58.0	573	54.7
Absolute dense area, cm ² , mean	34.7	22.8	48.2	103.2	58.2	46.2	44.7
Absolute nondense area, cm ² , mean	83.9	136.0	137.7	188.0	111.8	123.9	107.6
Percent dense area, %, mean	33.8	15.8	33.9	38.4	39.4	29.3	33.1
BMI, kg/m ² , mean	24.9	29.0	28.0	27.1	25.2	-	25.4
Parity, No., mean¶	2.5	3.0	3.0	1.6	2.8	-	2.3
Ever HRT use, %, mean	79.9	64.8	65.8	89.2	18.5	-	27.8
Current HRT use, %, mean	59.7	29.2	28.9	78.4	10.5	-	0.0

* Results are presented separately for women who were premenopausal and postmenopausal at the time of mammography. Em dashes (-) indicate lack of data availability. AGE = UK AGE Trial nested case-control study; BMI = body mass index; CAHRES = Cancer and Hormones Replacement in Sweden study; CBDS = Canadian Breast Density Study; CC = craniocaudal; EPIC-NL = Dutch contribution to the European Prospective Investigation into Cancer and Nutrition study; HRT = hormone replacement therapy; MCCS = Melbourne Collaborative Cohort Study; MCMAM = Mayo Clinic Mammography Study; MEC = Hawaii component of the Multiethnic Cohort Study; ML = mediolateral oblique; MMHS = Mayo Mammography Health Study; NHS1 = Nurses' Health Study 1; NHS2 = Nurses' Health Study 2; SBSP = Singapore Breast Cancer Screening Project; UK NHS = Norwich & Cambridge UK NHS Breast Cancer Screening Program; USC = University of Southern California mammographic density study.

† All studies with ethnicity listed as white include >90% whites. The MEC study includes 30% whites, 46% Asians, and 24% of other ethnicity. The USC study includes 47% whites, 20% Asians, and 33% blacks. The SBSP study includes 100% Asians.

‡ Data provided for the MCMAM cohort are based on data from the craniocaudal mammogram view; data from the mediolateral oblique mammogram view are given in parentheses. For the mediolateral oblique view, data on mammographic density were only available for 309 case patients and 556 control subjects.

§ Among parous women.

Table 2. Summary odds ratios (ORs) and 95% confidence intervals (CIs) for mammographic density phenotypes and risk of breast cancer among women who were premenopausal at the time of mammography

Main exposure	Covariables in model*	Summary OR† (95% CI)	P for heterogeneity by			
			Study	Age	BMI	Ethnicity
Absolute dense area	Age	1.38 (1.30 to 1.48)	.61	.44	.99	.73
Absolute dense area	Age, absolute nondense area	1.38 (1.29 to 1.48)	.89	.84	.86	.39
Absolute dense area	Age, BMI, parity	1.37 (1.29 to 1.47)	.50	.38	.94	.68
Absolute dense area	Age, BMI, parity, absolute nondense area	1.39 (1.30 to 1.49)	.73	.76	.92	.38
Absolute nondense area	Age	0.81 (0.76 to 0.86)	.79	.26	.84	.07
Absolute nondense area	Age, absolute dense area	0.84 (0.78 to 0.91)	.28	.06	.79	.03
Absolute nondense area	Age, BMI, parity	0.78 (0.71 to 0.86)	.20	.08	.32	.01
Absolute nondense area	Age, BMI, parity, absolute dense area	0.82 (0.71 to 0.94)	.02	.03	.38	.01
Percentage dense area	Age	1.45 (1.35 to 1.55)	.54	.40	.75	.04
Percentage dense area	Age, BMI, parity	1.52 (1.39 to 1.66)	.27	.31	.46	.01

* Covariables are measured at (within a few years) the time of mammography. Absolute dense area (per standard deviation), absolute nondense area (per standard deviation), percentage dense area (per standard deviation), age (per year), and body mass index (BMI; per kg/m²) are modeled as continuous variables, and parity is modeled as a categorical variable (nulliparous, 1–2, ≥3 children). Random effects models. All statistical tests were two-sided.

† Odds ratios represent one standard deviation increases in the mammographic density phenotypes.

Table 3. Summary odds ratios (ORs) and 95% confidence intervals (CIs) for different mammographic density phenotypes and risk of breast cancer among women who were postmenopausal at the time of mammography

Main exposure	Covariables in model*	Summary OR† (95% CI)	P for heterogeneity by			
			Study	Age	BMI	Ethnicity
Absolute dense area	Age	1.37 (1.33 to 1.40)	.58	.41	.59	.87
Absolute dense area	Age, absolute nondense area	1.37 (1.33 to 1.41)	.49	.60	.83	.74
Absolute dense area	Age, BMI, parity	1.38 (1.31 to 1.44)	.15	.80	.19	.76
Absolute dense area	Age, BMI, parity, absolute nondense area	1.38 (1.32 to 1.45)	.14	.46	.43	1.00
Absolute nondense area	Age	0.89 (0.84 to 0.94)	<.01	.25	.15	.61
Absolute nondense area	Age, absolute dense area	0.96 (0.88 to 1.03)	<.01	.64	.21	.33
Absolute nondense area	Age, BMI, parity	0.79 (0.73 to 0.85)	<.01	.95	.81	.41
Absolute nondense area	Age, BMI, parity, absolute dense area	0.85 (0.75 to 0.96)	<.01	.76	.74	.27
Percentage dense area	Age	1.37 (1.32 to 1.42)	.07	.24	.36	.30
Percentage dense area	Age, BMI, parity	1.53 (1.44 to 1.64)	.01	.45	.69	.27

* Covariables are measured at (within a few years) the time of mammography. Absolute dense area (per standard deviation), absolute nondense area (per standard deviation), percentage dense area (per standard deviation), age (per year), and body mass index (BMI; per kg/m²) are modeled as continuous variables, and parity is modeled as a categorical variable (nulliparous, 1–2, ≥3 children). Random effects models. All statistical tests were two-sided.

† Odds ratios represent one standard deviation increases in mammographic density phenotypes.

Finally, for postmenopausal women, we examined models additionally adjusted for ever HRT use and, separately, for current HRT use.

Meta-analysis. For each of the analyses described above, we pooled the study-specific β s using DerSimonian and Laird random-effects models to obtain a combined estimate, which we then exponentiated to arrive at a pooled odds ratio (OR) and its confidence interval (CI) (20). This method was conducted for the associations for quartiles 2, 3, and 4 (vs 1) and for the continuous effects of one standard deviation increases. For all analyses, we assessed heterogeneity across studies using Cochran's Q test. We further investigated heterogeneity using the I^2 statistic, which describes the percentage of total variation across studies resulting from heterogeneity rather than chance (21,22). We conducted all statistical analyses of categorical exposures using MIX version 2.0 Professional (BIOSTATXL) (23,24) and analyses of continuous exposures using the METAANAL macro (<http://www.hsph.harvard.edu/faculty/donna-spiegelman/files/metaanal>) for SAS version 9 (SAS Institute, Cary, NC). Forest plots were created using Microsoft Excel 2010 for Windows.

We used meta-regressions to assess effect modification by the following factors, chosen a priori: average age at mammography among control subjects, average BMI at mammography among control subjects, method of BMI ascertainment (self-reported or technician-measured), mammogram view (CC or MLO), ethnicity (white, Asian, or mixed), percentage ever having used HRT among control subjects (among postmenopausal women only), and percentage currently using HRT (among postmenopausal women only) among control subjects. Analyses of effect modification were conducted using STATA/IC 10.0 for Mac (STATA Corp, College Station, TX).

All tests were two-sided with P less than .05 considered to be statistically significant.

Results

Of the 13 studies in the meta-analysis (Table 1), 11 contributed data for premenopausal women ($n = 1776$ breast cancer cases; $n = 2834$ control subjects), and 12 contributed data for postmenopausal women ($n = 6643$ breast cancer cases; $n = 11\,187$ control subjects). Ten studies included almost exclusively (>90%) white women from the United States, Canada, Europe, or Australia, two studies included women of various ethnicities from the United States (including 20% and 46% Asians, respectively), and one study (the SBSP cohort) included only Asian women from Singapore. With the exception of the SBSP cohort, and to some extent the studies including women of various ethnicities, all studies were conducted in populations with a generally high risk of breast cancer (<http://globocan.iarc.fr>). The mean ages at mammography among premenopausal case patients and control subjects were 46 and 47 years, respectively, whereas the mean age among both postmenopausal case patients and control subjects was 61 years. In all studies, at both premenopausal and postmenopausal ages, absolute dense areas were higher among case patients than control subjects by between 2 and 13 cm², with one study (NHS2 at postmenopausal ages) finding a greater difference of 30 cm². Mean study-specific percentage dense area was also consistently higher in case patients than in control subjects by between 2 to 10 absolute percentage points, with the exception of the EPIC-NL study at postmenopausal ages where no appreciable difference was

observed. In contrast, the distributions were largely reversed for absolute nondense area, where study-specific mean nondense areas were between 1 and 30 cm² lower in case patients than control subjects, again with the exception of the EPIC-NL study at postmenopausal ages (Table 1). The SBSP cohort, which included only Asian women, had the highest percentage dense area among case patients and control subjects, both among premenopausal and postmenopausal women. The within-study correlation between absolute dense and nondense area varied across studies, ranging from -0.49 to 0.20 among premenopausal women, and from -0.48 to 0.03 among postmenopausal women (Supplementary Table 1, available online).

Premenopausal Women

Results for premenopausal women ($n = 1776$ breast cancer cases; $n = 2834$ control subjects) are presented in Table 2 and Figure 1. The age-adjusted summary odds ratio for a one standard deviation increment was 1.38 (95% CI = 1.30 to 1.48) for absolute dense area, 1.45 (95% CI = 1.35 to 1.55) for percentage dense area, and 0.81 (95% CI = 0.76 to 0.86) for absolute nondense area. Additional adjustment for BMI and parity changed the odds ratio for percentage dense area to 1.52 (95% CI = 1.39 to 1.66). Additional adjustment for BMI and parity did not materially change the results for absolute dense area (OR = 1.37; 95% CI = 1.29 to 1.47) or nondense area (OR = 0.78; 95% CI = 0.71 to 0.86). Results for nondense area were similar after further adjustment for absolute dense area (OR = 0.82; 95% CI = 0.71 to 0.94).

In the fully adjusted model, we observed statistically significant heterogeneity between studies ($P = .02$) for absolute nondense area (Table 2). After exclusion of the USC study, the heterogeneity was reduced (summary OR = 0.78; 95% CI = 0.69 to 0.89; $P_{\text{heterogeneity}} = .09$). In the fully adjusted models, there was also statistically significant heterogeneity by ethnicity for both absolute nondense and percentage dense area (both $P = .01$). In analyses restricted to studies including almost exclusively white women, the summary odds ratio for absolute nondense area was 0.78 (95% CI = 0.71 to 0.87; $P_{\text{heterogeneity}} = .36$), and for percentage dense area it was 1.60 (95% CI = 1.47 to 1.75; $P_{\text{heterogeneity}} = .99$) (Supplementary Table 3, available online). The corresponding odds ratios for the three studies including non-white women were 0.92 (95% CI = 0.53 to 1.57; $P_{\text{heterogeneity}} = .01$) and 1.31 (95% CI = 0.97 to 1.78; $P_{\text{heterogeneity}} = .11$). There was also statistically significant heterogeneity by age at the time of mammography for absolute nondense area in the fully adjusted model ($P = .03$); in general, the inverse association between the absolute nondense area and breast cancer risk increased with higher average age of each study's participants at the time of mammography (Supplementary Figure 1, available online). In analyses of the fully adjusted models restricted to studies including almost exclusively white women, the P for heterogeneity by age at the time of mammography was .09.

Postmenopausal Women

In postmenopausal women ($n = 6643$ breast cancer cases; $n = 11\,187$ control subjects) (Table 3; Figure 2), the age-adjusted summary odds ratio for a one standard deviation increment in absolute dense area was 1.37 (95% CI = 1.33 to 1.40). Much like among premenopausal women, additional adjustment for BMI and parity did not materially change the results (OR = 1.38; 95% CI = 1.31 to 1.44), nor did additional adjustment for absolute nondense area




Figure 1. Odds ratios (ORs) and 95% confidence intervals (CIs) for mammographic density phenotype and risk of breast cancer among women who were premenopausal at the time of mammography. Random effects models. All statistical tests were two-sided. Additional analytical details are in [Supplementary Table 2](#) (available online). AGE = UK AGE Trial nested case-control study; CAHRES = Cancer and Hormones Replacement in Sweden study; CBDS = Canadian Breast Density Study; CC = craniocaudal; EPIC-NL = Dutch contribution to the European Prospective Investigation

into Cancer and Nutrition study; HRT = hormone replacement therapy; MCCS = Melbourne Collaborative Cohort Study; MCMAM = Mayo Clinic Mammography Study; MEC = Hawaii component of the Multiethnic Cohort Study; ML = mediolateral oblique; MMHS = Mayo Mammography Health Study; NHS1 = Nurses' Health Study 1; NHS2 = Nurses' Health Study 2; SBSP = Singapore Breast Cancer Screening Project; UK NHS = Norwich & Cambridge UK NHS Breast Cancer Screening Program; USC = University of Southern California mammographic density study.




Figure 2. Odds ratios (ORs) and 95% confidence intervals (CIs) for mammographic density phenotypes and risk of breast cancer among women who were postmenopausal at the time of mammography. Random effects models. All statistical tests were two-sided. Additional analytical details are in [Supplementary Table 4](#) (available online). AGE = UK AGE Trial nested case-control study; CAHRES = Cancer and Hormones Replacement in Sweden study; CBDS = Canadian Breast Density Screening Program; CC = craniocaudal; EPIC-NL = Dutch contribution to the European Prospective Investigation into Cancer and Nutrition study; HRT = hormone replacement therapy; MCCS = Melbourne Collaborative Cohort Study; MCMAM = Mayo Clinic Mammography Study; MEC = Hawaii component of the Multiethnic Cohort Study; ML = mediolateral oblique; MMHS = Mayo Mammography Health Study; NHS1 = Nurses' Health Study 1; NHS2 = Nurses' Health Study 2; SBSP = Singapore Breast Cancer Screening Project; UK NHS = Norwich & Cambridge UK NHS Breast Cancer Screening Program; USC = University of Southern California mammographic density study.

(OR = 1.38; 95% CI = 1.32 to 1.45). For percentage dense area, the age-adjusted summary odds ratio was 1.37 (95% CI = 1.32 to 1.42), which increased to 1.53 (95% CI = 1.44 to 1.64) after additional adjustment for BMI and parity. For absolute nondense area, the age-adjusted summary odds ratio for a one standard deviation increment was 0.89 (95% CI = 0.84 to 0.94). Additional adjustment for BMI and parity changed the odds ratio to 0.79 (95% CI = 0.73 to 0.85). In the model further adjusted for absolute dense area, the corresponding summary odds ratio was 0.85 (95% CI = 0.75 to 0.96). Additionally adjusting for ever or current use of HRT did not materially change the results (data not shown).

In the fully adjusted models, there was statistically significant between-study heterogeneity for absolute nondense area ($P < .01$) and for percentage dense area ($P = .01$) with respect to breast cancer risk (Table 3). Between-study heterogeneity remained statistically significant in all analyses excluding the studies one by one (data not shown). In the fully adjusted models, there was also statistically significant heterogeneity by mammogram view for both absolute nondense area ($P = .04$) and percentage dense area ($P = .03$). For both mammographic density phenotypes, the associations were stronger in studies using CC vs MLO view mammograms (Supplementary Table 5, available online). For absolute nondense area, the summary odds ratio among studies using CC views was 0.79 (95% CI = 0.70 to 0.88; $P_{\text{heterogeneity}} < .01$), and the summary odds ratio among studies using MLO views (including the MLO view for the MCMAM study) was 1.00 (95% CI = 0.81 to 1.23; $P_{\text{heterogeneity}} < .01$). For percentage dense area, the summary odds ratio among studies using CC views was 1.59 (95% CI = 1.49 to 1.69; $P_{\text{heterogeneity}} = .12$), and the summary odds ratio among studies using MLO views (including the MLO view for the MCMAM study) was 1.40 (95% CI = 1.28 to 1.54; $P_{\text{heterogeneity}} = .20$). For the MCMAM study, results were generally similar using data from either the CC or MLO view (data not shown).

Discussion

In this meta-analysis of 13 case-control studies assessing the association between different mammographic density phenotypes and breast cancer risk, we confirmed that both absolute and percentage dense area are strong risk factors for breast cancer. The risk estimates for breast cancer were higher for percentage than for absolute dense area in the fully adjusted models. Although the confidence intervals for absolute and percentage dense area overlapped, this finding suggests that percentage dense area is the stronger of the two breast cancer risk factors. Results for absolute nondense area were less consistent across studies; most reported a lower risk of breast cancer with increasing absolute nondense area, but in several studies the association disappeared after adjustment for absolute dense area.

Among premenopausal women, the risk estimate for percentage dense area was higher than for absolute dense area in analyses adjusted for age and higher yet after further adjustment for BMI and parity (Table 2). Among postmenopausal women, the risk estimate for percentage dense area was identical to that for absolute dense area in analyses adjusted for age but substantially stronger in analyses additionally adjusted for BMI and parity (Table 3). Higher BMI is associated with a decreased risk of premenopausal breast cancer and an increased risk of postmenopausal breast cancer (25). Higher BMI is also associated with lower percentage dense area.

Prior studies suggest that percentage dense area and BMI are independent risk factors for breast cancer that confound each other (26,27). Our results confirm that BMI is an important confounder of the association between percentage dense area (and absolute nondense area) and breast cancer risk, especially among postmenopausal women. Our results also suggest that BMI is not an important confounder for the association between absolute dense area and breast cancer risk. These results underscore the importance of having data available on BMI when assessing the association between percentage dense area (and absolute nondense area) and breast cancer risk, including in risk prediction modeling studies comparing the predictive ability of the different mammographic density phenotypes. If data on BMI are unavailable, the association between absolute dense area and breast cancer risk should still represent an unbiased estimate (ie, unconfounded by BMI) of the association between dense area and breast cancer risk.

Most studies in this meta-analysis reported a lower risk of breast cancer with increasing absolute nondense area, but the associations disappeared after adjustment for absolute dense area in studies with the strongest negative correlation between absolute dense and nondense area. It is unclear why the correlation between absolute dense and nondense area (and thus the association of nondense area and breast cancer risk adjusted for dense area) varies substantially across studies. A possible explanation is measurement error. That is, if dense area is misclassified as nondense area (or vice versa) because of delineation issues with the thresholding technique, this will create a negative correlation between the absolute dense and nondense area. Another possible explanation is that there is a true negative correlation between the absolute dense and nondense area, perhaps because of lobular involution, which varies between populations and studies. Regardless of the underlying mechanism, the between-study variation in the correlation between nondense and dense area in this meta-analysis and the corresponding statistically significant between-study heterogeneity for absolute nondense area in several analyses make it difficult to determine whether absolute nondense area is an independent protective factor for breast cancer. This meta-analysis was limited by the information available to assess between-study heterogeneity. We were able to assess the effect modifiers which we considered to be relevant a priori (eg, BMI, ethnicity, mammogram view). We were unable, however, to determine the extent to which differences across studies in the associations between the mammographic density phenotypes and breast cancer risk were explained by study differences in exposure to other confounding factors, host biology among study participants, underlying breast cancer risk in the study populations, and/or technical differences such as the use of different mammogram machines, digitizers, and readers.

Although the confidence intervals for absolute and percentage dense area overlapped, our results suggest that percentage dense area is the stronger of the two breast cancer risk factors. If correct, this suggests that the ratio of fibroglandular to fatty tissue may be important in breast cancer development. Alternatively, assuming that women with high absolute or percentage mammographic density are at increased risk for breast cancer only because they have large amounts of fibroglandular breast tissue, measurement error may explain the weaker association for absolute dense area, and percentage dense area may be the most accurate estimate of the total amount of fibroglandular breast tissue. The current literature,

however, suggests that percentage dense area on a mammogram has a weak relationship with the total amount of fibroglandular breast tissue. Shepherd and colleagues estimated the fibroglandular volume from digitized film mammograms using single x-ray absorptiometry (28) and reported virtually no association between percentage dense area (measured using a software similar to Cumulus) and fibroglandular volume ($r^2 = 0.01$). Interestingly, other research has shown a high correlation between standard percentage density and calibrated percentage density measures accounting for acquisition parameters, including breast compression thickness (29). Still, additional studies are needed to determine whether absolute or percentage dense area on a mammogram most accurately reflects the total volume of fibroglandular breast tissue.

There are several biological mechanisms that can explain the association between the different mammographic density phenotypes and breast cancer risk. Absolute and percentage dense area are likely positively associated with breast cancer risk at least partially because they are positively correlated with the number of epithelial cells at risk of malignant transformation. Percentage dense area additionally reflects the amounts of fibroblasts and other stromal cells, connective tissue, and fat cells in the breasts, all which may affect breast cancer risk (30). It has been proposed, for example, that high extracellular matrix content and tissue stiffness or quantitative or structural changes of the stromal collagen such as cross-linking may partly explain the association between percentage dense area and breast cancer risk (31–33). Larger amounts of breast fat, which is the major component of absolute nondense area and inversely associated with percentage dense area, may also decrease breast cancer risk (6,34). The adipocytes that form the breast adipose tissue arise from the differentiation of stromal preadipocytes. Aromatase activity in the breast provides a source of estrogen production that may stimulate tumor growth. Aromatase activity in adipose tissue is primarily in stromal preadipocytes, and activity diminishes with differentiation to mature adipocytes (35,36). The loss of this source of estrogen after the differentiation might contribute to the inverse association between absolute nondense area and breast cancer risk. It should be noted, however, that breast fat has also been suggested to increase breast cancer risk (5,37). Finally, lobular involution is inversely associated with breast cancer risk, positively associated with absolute nondense area, and negatively associated with percentage dense area (38). Thus it is possible that nondense area is inversely associated with breast cancer risk because it reflects the degree of lobular involution.

In conclusion, we confirm that both absolute and percentage dense area on a mammogram are strong predictors of breast cancer risk. Our results also suggest that percentage dense area is a stronger breast cancer risk factor than absolute dense area and that absolute nondense area may be inversely associated with breast cancer risk.

References

1. McCormack VA, dos Santos Silva I. Breast density and parenchymal patterns as markers of breast cancer risk: a meta-analysis. *Cancer Epidemiol Biomarkers Prev.* 2006;15(6):1159–1169.
2. Trichopoulos D, Lipman RD. Mammary gland mass and breast cancer risk. *Epidemiology.* 1992;3(6):523–526.
3. Kato I, Beinat C, Bleich A, Su S, Kim M, Toniolo PG. A nested case-control study of mammographic patterns, breast volume, and breast cancer. *Cancer Causes Control.* 1995;6(5):431–438.

4. Boyd N, Martin L, Gunasekara A, et al. Mammographic density and breast cancer risk: evaluation of a novel method of measuring breast tissue volumes. *Cancer Epidemiol Biomarkers Prev.* 2009;18(6):1754–1762.
5. Lokate M, Peeters PH, Peelen LM, Haars G, Veldhuis WB, van Gils CH. Mammographic density and breast cancer risk: the role of the fat surrounding the fibroglandular tissue. *Breast Cancer Res.* 2011;13(5):R103.
6. Pettersson A, Hankinson SE, Willett WC, Lagiou P, Trichopoulos D, Tamimi RM. Nondense mammographic area and risk of breast cancer. *Breast Cancer Res.* 2011;13(5):R100.
7. Vachon CM, Scott CG, Fasching PA, et al. Common breast cancer susceptibility variants in LSP1 and RAD51L1 are associated with mammographic density measures that predict breast cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2012;21(7):1156–1166.
8. Baglietto L, Krishnan K, Stone J, et al. Associations of mammographic dense and nondense areas and body mass index with risk of breast cancer. *Am J Epidemiol.* 2014;179(4):475–483.
9. Boyd NE, Guo H, Martin LJ, et al. Mammographic density and the risk and detection of breast cancer. *N Engl J Med.* 2007;356(3):227–236.
10. Maskarinec G, Pagano I, Lurie G, Wilkens LR, Kolonel LN. Mammographic density and breast cancer risk: the multiethnic cohort study. *Am J Epidemiol.* 2005;162(8):743–752.
11. Olson JE, Sellers TA, Scott CG, et al. The influence of mammogram acquisition on the mammographic density and breast cancer association in the Mayo Mammography Health Study cohort. *Breast Cancer Res.* 2012;14(6):R147.
12. Pinto Pereira SM, McCormack VA, Hipwell JH, et al. Localized fibroglandular tissue as a predictor of future tumor location within the breast. *Cancer Epidemiol Biomarkers Prev.* 2011;20(8):1718–1725.
13. Stone J, Ding J, Warren RM, Duffy SW, Hopper JL. Using mammographic density to predict breast cancer risk: dense area or percentage dense area. *Breast Cancer Res.* 2010;12(6):R97.
14. Tamimi RM, Eriksson L, Lagiou P, et al. Birth weight and mammographic density among postmenopausal women in Sweden. *Int J Cancer.* 2010;126(4):985–991.
15. Ursin G, Ma H, Wu AH, et al. Mammographic density and breast cancer in three ethnic groups. *Cancer Epidemiol Biomarkers Prev.* 2003;12(4):332–338.
16. Vachon CM, Brandt KR, Ghosh K, et al. Mammographic breast density as a general marker of breast cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2007;16(1):43–49.
17. Wong CS, Lim GH, Gao F, et al. Mammographic density and its interaction with other breast cancer risk factors in an Asian population. *Br J Cancer.* 2011;104(5):871–874.
18. Byng JW, Boyd NE, Fishell E, Jong RA, Yaffe MJ. The quantitative analysis of mammographic densities. *Phys Med Biol.* 1994;39(10):1629–1638.
19. Ursin G, Astrahan MA, Salane M, et al. The detection of changes in mammographic densities. *Cancer Epidemiol Biomarkers Prev.* 1998;7(1):43–47.
20. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials.* 1986;7(3):177–188.
21. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med.* 2002;21(11):1539–1558.
22. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327(7414):557–560.
23. Bax L, Yu LM, Ikeda N, Moons KG. A systematic comparison of software dedicated to meta-analysis of causal studies. *BMC Med Res Methodol.* 2007;7(40).
24. Bax L, Yu LM, Ikeda N, Tsuruta H, Moons KG. Development and validation of MIX: comprehensive free software for meta-analysis of causal research data. *BMC Med Res Methodol.* 2006;6(50).
25. Renehan AG, Tyson M, Egger M, Heller RF, Zwahlen M. Body-mass index and incidence of cancer: a systematic review and meta-analysis of prospective observational studies. *Lancet.* 2008;371(9612):569–578.
26. Boyd NE, Martin LJ, Sun L, et al. Body size, mammographic density, and breast cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2006;15(11):2086–2092.
27. Harris HR, Tamimi RM, Willett WC, Hankinson SE, Michels KB. Body size across the life course, mammographic density, and risk of breast cancer. *Am J Epidemiol.* 2011;174(8):909–918.
28. Shepherd JA, Kerlikowske K, Ma L, et al. Volume of mammographic density and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev.* 2011;20(7):1473–1482.

29. Heine JJ, Cao K, Rollison DE, Tiffenberg G, Thomas JA. A quantitative description of the percentage of breast density measurement using full-field digital mammography. *Acad Radiol*. 2011;18(5):556–564.
30. Boyd NF, Martin LJ, Bronskill M, Yaffe MJ, Duric N, Minkin S. Breast tissue composition and susceptibility to breast cancer. *J Natl Cancer Inst*. 2010;102(16):1224–1237.
31. Brower V. Homing in on mechanisms linking breast density to breast cancer risk. *J Natl Cancer Inst*. 2010;102(12):843–845.
32. Butcher DT, Alliston T, Weaver VM. A tense situation: forcing tumour progression. *Nat Rev Cancer*. 2009;9(2):108–122.
33. Provenzano PP, Inman DR, Eliceiri KW, Keely PJ. Matrix density-induced mechanoregulation of breast cell phenotype, signaling and gene expression through a FAK-ERK linkage. *Oncogene*. 2009;28(49):4326–4343.
34. Pettersson A, Tamimi RM. Breast fat and breast cancer. *Breast Cancer Res Treat*. 2012;135(1):321–323.
35. McInnes KJ, Brown KA, Knowler KC, Chand AL, Clyne CD, Simpson ER. Characterisation of aromatase expression in the human adipocyte cell line SGBS. *Breast Cancer Res Treat*. 2008;112(3):429–435.
36. Simpson ER, Clyne C, Rubin G, et al. Aromatase—a brief overview. *Annu Rev Physiol*. 2002;64:93–127.
37. Beer AE, Billingham RE. Adipose tissue, a neglected factor in aetiology of breast cancer? *Lancet*. 1978;2(8084):296.
38. Ghosh K, Hartmann LC, Reynolds C, et al. Association between mammographic density and age-related lobular involution of the breast. *J Clin Oncol*. 2010;28(13):2207–2212.

Funding

This work was supported by the National Cancer Institute at the National Institutes of Health (R01 CA124865, R01 CA131332, P01 CA87969, R01 CA50385, R01 CA085265, R03 CA135699, R37 CA054281, R01 CA140286, R01 CA128931, R01 CA97396, P50 CA116201); Department of Defense grant DAMD 17-00-1-0331; the National Institutes of Health (R25 CA098566 to REG); the Märta and Hans Rausing's Initiative Against Breast Cancer; the W81XWH-05-1-0314 Innovator Award; the United States Department of Defense Breast Cancer Research Program; the Office of the Congressionally Directed Medical Research Programs; the Agency for Science, Technology and Research; the Swedish Cancer Society (5128-B07-01PAF to KC); the National Cancer Institute of Canada (to NFB); the SingHealth Foundation (SHF/FG450S/2009); the National Medical Research Council (NMRC/0711/2002); the National University Cancer Institute, Singapore (NMRC/CG/NCIS/2010); the MammoGrid Project (to RMLW); the Victorian Breast Cancer Consortium; the National Breast Cancer Foundation; the Cancer Council Victoria; the National Health and Medical Research Council (251533, 209057, 504711, 454671 to JS); the case-control study within the EPIC-NL cohort was funded by the Dutch Cancer Society (KWF 2008–4071 and KWF h1-4348); the EPIC-NL cohort was funded by the 'Europe against Cancer' program of the European Commission (SANCO); the Dutch Ministry of

Health, Welfare and Sports (WVS) and ZonMW; Breast Cancer Campaign (2007MayPR23 to VM).

Notes

The authors contributed to the design of the study, analysis, and interpretation of results, the decision to submit the manuscript, and the writing of the manuscript.

We would like to thank the participants and staff of the NHS and NHS2 for their valuable contributions as well as the following state cancer registries for their help: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, WY. In addition, this study was approved by the Connecticut Department of Public Health (DPH) Human Investigations Committee. Certain data used in this publication were obtained from the DPH. The authors assume full responsibility for analyses and interpretation of these data.

Affiliations of authors: Department of Epidemiology (AP, REG, RMT), and Department of Biostatistics (BAR), Harvard School of Public Health, Boston, MA; Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Oslo, Norway (GU); Department of Preventive Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA (GU); Department of Non-Communicable Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK (IdSS); Section of Environment and Radiation, International Agency for Research on Cancer, Lyon, France (VM); Cancer Epidemiology Centre, Cancer Council Victoria, Melbourne, Australia (LB, GGG, KK); Centre for Molecular, Environmental, Genetic and Analytical Epidemiology, University of Melbourne, Melbourne, Australia (LB, GGG, JLH, KK, JS); Division of Epidemiology (CV) and Division of Biomedical Statistics and Informatics (CS), Department of Health Sciences Research, Mayo Clinic, Rochester, MN; Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, The Netherlands (MFB, CHvG); Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, Australia (GGG); Saw Swee Hock School of Public Health (KSC, MH, CSW), and Department of Surgery, Yong Loo Lin School of Medicine (MH), National University of Singapore, National University Health System, Singapore, Singapore; Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden (KC, LE, PH, MH); Department of Radiology, Addenbrooke's Hospital, Cambridge, UK (RMLW); School of Population and Public Health, University of British Columbia, Vancouver, BC, Canada (GH); Prevention and Cancer Control, Cancer Care, Toronto, ON, Canada (AMC); Human Genetics, Genome Institute of Singapore, Singapore, Singapore (JL); Campbell Family Institute for Breast Cancer Research, Ontario Cancer Institute, Toronto, ON, Canada (QL, NFB); University of Hawaii Cancer Center, Honolulu, HI (IP, GM); Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA (BAR, RMT).