



Published in final edited form as:

Nature. 2010 October 28; 467(7319): 1128–1132. doi:10.1038/nature09465.

Maternal mRNA deadenylation and decay by the piRNA pathway in the early *Drosophila* embryo

Christel ROUGET¹, Catherine PAPIN¹, Anthony BOUREUX², Anne-Cécile MEUNIER, Bénédicte FRANCO, Nicolas ROBINE³, Eric C. LAI³, Alain PELISSON⁴, and Martine SIMONELIG*

mRNA Regulation and Development, Institute of Human Genetics, CNRS UPR1142, 141 rue de la Cardonille, 34396 Montpellier Cedex 5, France

²CRBM, UMR5237, Université Montpellier II, CNRS, 1919 route de Mende, 34293 Montpellier, France

³Sloan-Kettering Institute, Department of Developmental Biology, 1017 Rockefeller Research Laboratories, 1275 York Avenue, New York, New York 10065, USA

⁴Mechanisms and Control of Retrotransposition, Institute of Human Genetics, CNRS UPR1142, 141 rue de la Cardonille, 34396 Montpellier Cedex 5, France

Keywords

CCR4 deadenylase; mRNA decay; *nanos*; piRNA; RNA silencing; Translational control

Piwi-associated RNAs (piRNAs), a specific class of 24- to 30-nucleotide-long RNAs produced by the Piwi-type of Argonaute proteins have a specific germline function in repressing transposable elements. This repression is thought to involve heterochromatin formation, transcriptional and posttranscriptional silencing^{1–6}. The piRNA pathway has other essential functions in germline stem cell maintenance⁷ and in maintaining germline DNA integrity^{8–10}. Here, we uncover an unexpected function of the piRNA pathway in the decay of maternal mRNAs and in translational repression in the early embryo. A subset of maternal mRNAs is degraded in the embryo, at the maternal to zygotic transition. In *Drosophila*, maternal mRNA degradation depends on the RNA binding protein Smaug (Smg) and the deadenylase CCR4^{11–13}, as well as the zygotic expression of a microRNA (miRNA) cluster¹⁴. Using mRNA encoding the embryonic posterior morphogen Nanos (Nos) as a paradigm to study maternal mRNA decay, we found that CCR4-mediated deadenylation of *nos* depends on components of the piRNA pathway including piRNAs

Reprints and permissions information is available at www.nature.com/reprints.

*Corresponding author: Martine Simonelig, Tel: 33 4 99 61 99 59, Fax: 33 4 99 61 99 57, Martine.Simonelig@igh.cnrs.fr.

¹Equal first authors

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Author Contributions C.R. and C.P. designed and performed the experiments, analysed the data and contributed equally to the study. A.C.M. contributed to the generation of DNA constructs, B.F. contributed to PAT assays in Fig. 1. A.B., N.R. and E.C.L. performed the bioinformatic analyses. A.P. performed northern blots. M.S. designed the study, analysed data and wrote the paper. All authors discussed the results and commented on the manuscript.

complementary to a specific region in *nos* 3'-UTR. Reduced deadenylation when piRNA-induced regulation is impaired, correlates with *nos* mRNA stabilization and translational derepression in the embryo, resulting in head development defects. Aubergine (Aub), one of the Argonaute proteins in the piRNA pathway, is present in a complex with Smg, CCR4, *nos* mRNA and piRNAs that target *nos* 3'-UTR, in the bulk of the embryo. We propose that piRNAs and their associated proteins act together with Smg to recruit the CCR4 deadenylation complex to specific mRNAs, thus promoting their decay. Because piRNAs involved in this regulation are produced from transposable elements, this identifies a direct developmental function for transposable elements in the regulation of gene expression.

In *Drosophila* embryos, Nos is expressed as a gradient which emanates from the posterior pole and organizes abdominal segmentation¹⁵. The majority of *nos* mRNA is distributed throughout the bulk cytoplasm, translationally repressed¹⁶, and subsequently degraded during the first 2–3 hours of development. This repression is essential for head and thorax segmentation^{16,17}. A small amount of *nos* transcripts, localised at the posterior pole of the embryo, escapes degradation and is actively translated, giving rise to the Nos protein gradient. *nos* mRNA decay in the bulk cytoplasm depends on the CCR4-NOT deadenylation complex and its recruitment onto *nos* by Smg. This contributes to translational repression in the bulk of the embryo and is required for embryonic antero-posterior patterning¹³.

Smg was suggested not to be the only activator of *nos* mRNA decay during early embryogenesis^{11,12}. Zygotically-expressed miRNAs have been reported to activate maternal mRNA deadenylation in zebrafish embryos¹⁸ and decay in *Drosophila* embryos¹⁴. We investigated the potential involvement of other classes of small RNAs in mRNA deadenylation and decay before zygotic expression. Since piRNAs are expressed maternally in the germline and present in early embryos^{19,20}, we analysed the possible role of the piRNA pathway in maternal mRNA deadenylation. Piwi, Aub and Ago3 are specific Argonaute proteins^{1,3,21,22}, Armitage (Armi) and Spindle-E (Spn-E) are RNA helicases, and Squash (Squ) is a nuclease^{2,10,23,24} involved in piRNA biogenesis and function. Poly(A) test (PAT) assays were performed to measure *nos* mRNA poly(A) tail length in embryos spanning one hour intervals during the first four hours of embryogenesis. In contrast to the progressive shortening of *nos* mRNA poly(A) tails observed in wild-type embryos correlating with mRNA decay during this period, *nos* poly(A) tail shortening was affected in embryos from females mutant for the piRNA pathway (herein referred to as mutant embryos) (Figure 1a, Supplementary Figures 1a, 2, 12). This defect in deadenylation correlated with higher amounts of *nos* mRNA in mutant embryos, as quantified by RT-QPCR (Figure 1b). *in situ* hybridization revealed stabilized *nos* mRNA in the bulk cytoplasm of mutant embryos where it is normally degraded in the wild-type (Figure 1c, Supplementary Figure 1b). Consistent with previous data showing that *nos* mRNA deadenylation is required for translational repression¹³, defective deadenylation in mutant embryos resulted in the presence of ectopic Nos protein throughout the embryo (Figure 1d, Supplementary Figure 1c). The presence of Nos in the anterior region results in the repression of *bicoid* and *hunchback* mRNA translation and in affected head skeleton. Consistent with earlier mentioned defects⁷, we found that the *piwi*¹ mutant embryos which were able to produce a cuticle had strong head defects (Figure 1e).

The piRNA pathway plays a role during early oogenesis in preventing DNA damage, possibly through the repression of transposable element transposition. DNA double-strand breaks arising in mutants of the piRNA pathway result in affected embryonic axis specification, and this developmental defect is suppressed by mutations in the Chk2 DNA damage signal transduction pathway^{9,10}. We found that defects in *nos* mRNA deadenylation and decay observed in *aub* or *armi* mutants were not suppressed by Chk2 (*mnk^{P6}*) mutations, indicating that these defects did not result from activation of the Chk2 pathway earlier during oogenesis (Supplementary Figure 3a–c). Moreover, affected deadenylation of *nos* mRNA in piRNA pathway mutants did not depend on *oskar* (Supplementary Figure 3d).

We addressed a potential direct role of the piRNA pathway in the regulation of *nos* mRNA deadenylation and decay in the embryo. Aub and Piwi accumulate in the pole plasm and in pole cells of the embryo^{25,26}. However, we found lower levels of Aub and Piwi throughout the entire embryo (Figure 2a, Supplementary Figures 4, 5). Ago3 was also present throughout the embryo (Supplementary Figure 6a, c). Aub and Ago3 were cytoplasmic and accumulated in discrete foci, a distribution similar to those of CCR4 and Smg (Figure 2b, Supplementary Figure 6b). CCR4 and Smg were reported to partially colocalise in small cytoplasmic foci¹³. Aub and Ago3 also partially colocalised with Smg and CCR4 in the bulk of syncytial embryos, in both cytoplasmic foci and a diffusely distributed cytoplasmic pool (Figure 2b, Supplementary Figure 6b). Importantly, the distributions of CCR4 and Smg depended on the piRNA pathway, as they were strongly affected in *aub* and *spn-E* mutant embryos. Although global amounts of CCR4 and Smg did not decrease in mutant embryos, CCR4 foci strongly increased in size, whereas Smg foci decreased in size or disappeared (Figure 2c, d). This suggests different functions for a subset of CCR4 and Smg foci and that deadenylation could take place diffusely in the cytoplasm. These results demonstrate a functional link between CCR4-mediated deadenylation and the piRNA pathway.

Co-immunoprecipitation experiments showed that Aub co-precipitated Smg, CCR4 and Ago3 in the absence of RNA, suggesting the presence of these proteins in a common complex (Figure 3a, Supplementary Figure 7a, b). Smg also co-precipitated CCR4, Aub and Ago3 (Figure 3b, Supplementary Figure 7c). Piwi, however, was not found to co-precipitate Smg or CCR4 (data not shown). Importantly, Smg, CCR4 and Ago3 also co-precipitated with Aub in *osk⁵⁴* mutant embryos that are defective in pole plasm assembly²⁷, indicating the presence of this complex outside the pole plasm (Figure 3a). We next showed that *nos* mRNA co-precipitated with Aub in both wild-type and *osk⁵⁴* embryos. The amounts of *nos* mRNA were similar in Aub and Smg immunoprecipitates (Figure 3c).

These findings show that the Argonaute proteins Aub and Ago3 associate with Smg and the CCR4 deadenylase complex to directly regulate *nos* mRNA in the bulk cytoplasm of early embryos.

nos 3'-UTR contains Smg binding sites (SRE) located in its 5'-most region (referred to as TCE)¹⁶. We searched for piRNAs, sequenced from early embryos, presumed capable of targeting *nos* 3'-UTR based on their sequence complementarity. Strikingly, a specific region located in the 3'-most part of the 3'-UTR could be targeted by over 200 copies of piRNAs originating from two transposable elements, *412* and *roo* (Figure 4a, Supplementary Figure

8). piRNAs complementary to *nos* 3'-UTR were visualized by northern blots. In addition, piRNAs predicted to target *nos* 3'-UTR co-immunoprecipitated with Aub (Figure 4b). We used *nos* genomic transgenes deleted for different parts of the 3'-UTR¹⁶ to address the requirement of the corresponding regions for *nos* mRNA deadenylation. We have shown previously that the TCE (nt 1–184) is required for *nos* mRNA poly(A) tail shortening, consistent with the role of Smg in this process¹³. Deletion of region 184–403 (*nos*($\Delta 1$)) had no effect, whereas poly(A) tails from the transgene deleted of the region 403–618 (*nos*($\Delta 2$)) were elongated in 3–4 hour embryos (Figure 4c, Supplementary Figure 12). This could indicate a regulation by the miRNA predicted by miRBase to target this region. Deletion of 618–844 in *nos* 3'-UTR (*nos*($\Delta 3$)) had a strong effect on *nos* deadenylation (Figure 4c, Supplementary Figures 9, 12). Consistent with this, *nos* mRNA levels produced by this transgene remained mostly stable (Figure 4d). This resulted in defects in embryo patterning: A total of 35% (n=1894) of embryos from *nos*($\Delta 3$) females did not hatch and among them 86% (n=28) showed head skeleton defects (Figure 3e). We next deleted specific sequences complementary to *412* (15 nt) and *roo* (11 nt) retrotransposon piRNAs (Supplementary Figure 8). These short deletions, either independently or in combination, affected *nos* mRNA deadenylation (Figure 4f, Supplementary Figure 12).

To further support the role of retrotransposon piRNAs in *nos* mRNA regulation, we blocked *412* and *roo* piRNAs by injecting specific 2' O-methyl anti-piRNA in embryos²⁸, and recorded cuticles as a functional assay of Nos ectopic synthesis at the anterior pole. Injection of anti-piRNA(*412*) or anti-piRNA(*roo*) resulted in specific head development defects (Figure 4g).

Together, these results provide strong evidence that an interaction between piRNAs and *nos* mRNA is required for *nos* mRNA deadenylation and translational repression in the first hours of embryogenesis.

We have identified a new function of the piRNA pathway in the regulation of maternal mRNAs. Recently, piRNAs derived from 3'-UTR of cellular transcripts have been identified in gonadal somatic cells, although their biological role has not been clarified^{29,30}. Here, we propose that piRNAs, in complex with Piwi-type Argonaute proteins Aub and Ago3, target *nos* maternal mRNAs and recruit or stabilize the CCR4-NOT deadenylation complex together with Smg (Supplementary Figure 10). These interactions induce rapid mRNA deadenylation and decay. Thus, activation of mRNA deadenylation represents a new direct mechanism of action for the piRNA pathway with an essential developmental function during the first steps of embryogenesis.

Smg is a general factor for mRNA decay during early embryogenesis¹². Because Aub and Ago3 are present in a complex with Smg in early embryos, a proportion of Smg mRNA targets could be regulated by the piRNA pathway. Consistent with this, other maternal mRNAs that are destabilized during early embryogenesis, are targeted by abundant piRNAs and their deadenylation depends on the piRNA pathway (Supplementary Figure 11).

These piRNAs involved in gene regulation are generated from transposable element sequences. Although transposable elements have been described to be essential for genome

dynamics and evolution, their immediate function within an organism has remained rather elusive. This study provides evidence for a co-evolution between transposable elements and the host genome and reveals the direct developmental function of transposable elements in embryonic patterning, through the regulation of gene expression.

Methods summary

RNA and proteins were manipulated using methods described previously and reported in the Methods section. **Embryo injections.** Injections of embryos were performed laterally with 400 μM of 2' O-methyl oligonucleotides as reported previously²⁸. Injection buffer was 0.5 mM NaPO_4 , 5 mM KCl. Sequences of 2' O-methyl oligonucleotides are indicated in Methods. **Bioinformatics.** A total of 29,108,987 piRNAs sequenced from 0–1 hour embryos (GSM286613 and GSM286604 data sets¹⁹) and from 0–2 hour embryos (GSM327625, GSM327626, GSM327627, GSM327628 and GSM327629 data sets²⁰) was blasted against *nos* 3'-UTR using the following parameters: NCBI blast with an E value of 100 and a 14-nt match and WU-blast with an E value of 10 and a 11-nt match. Regions potentially targeted by piRNAs with an occurrence of less than 10 were not considered.

Methods

Drosophila stocks and genetics

The *w*¹¹¹⁸ stock was used as a control. Mutant stocks were *w*; *Sp aub*^{N11} *bw*/CyO, *aub*^{HN2} *cn*¹ *bw*¹/CyO, *aub*^{QC42} *cn*¹ *bw*¹/CyO³¹, *mnk*^{P6}, *mnk*^{P6} *aub*^{HN2}/CyO, *mnk*^{P6} *aub*^{QC42}/CyO⁹, *mnk*^{P6}; *armi*¹/SM6-TM6B, *mnk*^{P6}; *armi*^{72.1}/SM6-TM6B⁹, *piwi neo-FRT40A*/CyO⁷, *spn-E*¹/TM3, *ry*⁵⁰⁶ *spn-E*^{hls-03987} *e*/TM3³², *armi*¹/TM3, *armi*^{72.1}/TM3²³, *cn*¹ *bw*¹ *squ*^{HE47}/CyO, *cn*¹ *bw*¹ *squ*^{PP32}/CyO¹⁰, *bw*; *st ago*^{3t1}/TM6B *Tb*, *bw*; *st ago*^{3t3}/TM6B *Tb*²², *osk*⁵⁴ *spn-E*¹/TM3, *osk*⁵⁴ *spn-E*^{hls-03987} *e*/TM3. *piwi*¹ mutant embryos are from germline clones that were induced with two 1.5-hour heat shocks at 37°C during the second- and third-instar larval stage, using the FLP recombinase-dominant female sterile technique³³. *smg* mutants were *smg*¹ and a deficiency overlapping *smg*, *Df*(*Scj*^{R6})¹⁷. *nos*^{BN} mutant does not produce *nos* mRNA³⁴. *osk*⁵⁴ is a null allele. GFP-Aub was expressed following crosses between the germline driver *nos-Gal4:VP16*³⁵ and *UASp-GFP-Aub*²⁵. *nos*(Δ) stocks are transgenic lines containing a *nos* genomic transgene in which different parts of the 3'UTR have been removed¹⁶. *nos*(Δ) stocks were a gift from R. Wharton. The *nos*(Δ *pi412*) and *nos*(Δ *piroo*) transgenes, in which 15 nt (TATATTTATTCAATT) and 11 nt (AACACACATAT) have been deleted, respectively were generated as follows. The **pBSKS-R5561** (containing a 5.7 kb *nos* genomic fragment, gift from R. Wharton) was used as a template for PCR reactions to produce the deletion. For each construct, two PCR reactions were performed using the following primers: For *nos*(Δ *pi412*), 5'CATTCCGATCAAAGCTGGGTTAACC (Primer 1) and 5'AAATTGATCAATGGTAAACAATAACATATATATATAT which contains the 15 nt deletion, and 5'TATATATATATATATATATGTTATTGTTTACCATTGATCAATTT which contains the 15 nt deletion and 5'CTCCACCGCGGTGGCGGCCGC (Primer 2). For *nos*(Δ *piroo*), Primer 1 and 5'TATATATATATATATATATAGGAAATGAATACTTGCATACATA which contains the 11 nt deletion, and 5'TGTATCGCAAGTATTCATTTCCCTATATATATATATATATATA which contains the 11 nt deletion and Primer 2. For each construct, the two PCR products

were annealed and used as a template for a third PCR reaction using Primers 1 and 2. This third PCR product containing either the 15 nt or the 11 nt deletion surrounded by the restriction sites *Bgl*III and *Not*I was cloned into the TAcloning vector (**pCRII**) (Invitrogen) and sequenced. For the *nos*(Δ *piroo*- Δ *pi412*) transgene, the PCR generating the Δ *piroo* deletion was done using **pCRII** containing the *nos* Δ *pi412* deletion as a template. The *Bgl*III-*Not*I fragment containing the deletion was used to replace the *Bgl*III-*Not*I fragment in the original **pBSKS-R5561**. An *Eco*RI-*Not*I fragment containing the whole genomic fragment with the deletion was cloned into the **pCaSpeR4**. Transformant stocks were produced by BestGene Inc.

Embryo injections

Sequences of 2' O-methyl oligonucleotides were as follows. anti-*pi*(412):
UCGGGCUGACAUAUAUUUAUCAAUU anti-*pi*(*roo*):
UCCAAACACACAUAUAUAUAUAAAUA anti-miR129-1:
GCAAGCCCAGACCGCAAAAAG (human miR129-1 is not conserved in *Drosophila*).

RNA

PAT assays, RT-PCR and RT-QPCR were performed as described previously^{13,36}, and were made from two to four independent RNA preparations. For *nos*(Δ 3) transgene, a different *nos* specific primer was used (5'GTCGTCGGCTACGCATTCATTGT), as the region normally amplified in *nos* PAT assays is deleted in this transgene. We verified by sequencing that the poly(A) site used in mRNA from this transgene is identical to the one used in *nos* endogenous mRNA. Real-time PCR (QPCR) were performed with the LightCycler System (Roche Molecular Biochemical) using *rp49* as a control mRNA¹³. For quantification of *nos* mRNA in 2–3 hour and 3–4 hour embryos, the levels were normalized with the levels of *nos* mRNA in 0–1 hour embryos that were set to 100% for each genotype. Northern blots were performed as described previously³⁷. The sequence of the riboprobe specific to 412 piRNA was 5'GGGCUGACAUAUAUUUAUCAAUU.

RNA *in situ* hybridization and cuticle preparations

Whole mount *in situ* hybridization and cuticle preparations were performed by standard methods. The probe for *in situ* hybridization was an antisense RNA made from the **pN5** *nos* cDNA clone.

Antibodies, western blots, immunostaining and immunoprecipitations

Immunoprecipitations were performed as described previously¹³ using 0–2 hour embryos, and mouse anti-Aub (4D10²¹), mock IPs: mouse anti-HA (12CA5 Developmental Studies Hybridoma Bank, for wild-type embryos) and mouse IgG (sc-2025 Santa Cruz Biotechnology, for *osk*⁵⁴ embryos), rabbit anti-Aub (Abcam, ab17724), mock IPs: rabbit IgG (sc-2027 Santa Cruz Biotechnology), mouse anti-GFP (mAb 3E6 Invitrogen), guinea pig anti-Smg (gift from C. Smibert), mock IP: guinea pig pre-immune serum. Protein coimmunoprecipitations were performed in the presence of 0.1 mg/ml RNase A. Western blots and immunostaining were performed as reported^{38,39}. Antibodies for western blots were used at the following dilutions: guinea pig anti-Smg 1/5000, anti-CCR4 1/1000⁴⁰, anti-

Piwi 1/20 (P4D2¹), anti-Aub 1/1500 (4D10²¹) and anti-Ago3 1/500 (9G3²¹). Antibodies for immunostaining were used at the following dilutions: rabbit anti-Nos 1/1000 (gift from A. Nakamura), guinea pig anti-Smg 1/1000, anti-CCR4 1/300, anti-Aub 1/1500 (4D10), anti-Piwi 1/1 (P4D2) and anti-Ago3 1/300 (9G3).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments


We are very grateful to A. Nakamura, M. and H. Siomi, C. Smibert, H. Lin, P. Macdonald, T. Schüpbach, W. Theurkauf, R. Wharton and P. Zamore, for their gifts of antibodies or *Drosophila* stocks. We thank M. Benkirane for the gift of 2' O-methyl anti-miR129. This work was supported by the CNRS UPR1142, ANR Blanche (CONTRA), FRM (Equipe FRM 2007) and ARC Libre 2009 to M.S. and by the NIH to E.C.L. C.R., A.C.M. and B.F. held salaries from ANR Blanche.

References


1. Saito K, et al. Specific association of Piwi with rasiRNAs derived from retrotransposon and heterochromatic regions in the *Drosophila* genome. *Genes Dev.* 2006; 20:2214–22. [PubMed: 16882972]
2. Vagin VV, et al. A distinct small RNA pathway silences selfish genetic elements in the germline. *Science.* 2006; 313:320–4. [PubMed: 16809489]
3. Brennecke J, et al. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell.* 2007; 128:1089–103. [PubMed: 17346786]
4. Yin H, Lin H. An epigenetic activation role of Piwi and a Piwi-associated piRNA in *Drosophila melanogaster*. *Nature.* 2007; 450:304–8. [PubMed: 17952056]
5. Aravin AA, et al. A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol Cell.* 2008; 31:785–99. [PubMed: 18922463]
6. Lim AK, Tao L, Kai T. piRNAs mediate posttranscriptional retroelement silencing and localization to pi-bodies in the *Drosophila* germline. *J Cell Biol.* 2009; 186:333–42. [PubMed: 19651888]
7. Cox DN, et al. A novel class of evolutionarily conserved genes defined by piwi are essential for stem cell self-renewal. *Genes Dev.* 1998; 12:3715–27. [PubMed: 9851978]
8. Chen Y, Pane A, Schupbach T. Cutoff and aubergine mutations result in retrotransposon upregulation and checkpoint activation in *Drosophila*. *Curr Biol.* 2007; 17:637–42. [PubMed: 17363252]
9. Klattenhoff C, et al. *Drosophila* rasiRNA pathway mutations disrupt embryonic axis specification through activation of an ATR/Chk2 DNA damage response. *Dev Cell.* 2007; 12:45–55. [PubMed: 17199040]
10. Pane A, Wehr K, Schupbach T. zucchini and squash encode two putative nucleases required for rasiRNA production in the *Drosophila* germline. *Dev Cell.* 2007; 12:851–62. [PubMed: 17543859]
11. Semotok JL, et al. Smaug recruits the CCR4/POP2/NOT deadenylase complex to trigger maternal transcript localization in the early *Drosophila* embryo. *Curr Biol.* 2005; 15:284–94. [PubMed: 15723788]
12. Tadros W, et al. SMAUG is a major regulator of maternal mRNA destabilization in *Drosophila* and its translation is activated by the PAN GU kinase. *Dev Cell.* 2007; 12:143–55. [PubMed: 17199047]
13. Zaessinger S, Busseau I, Simonelig M. Oskar allows nanos mRNA translation in *Drosophila* embryos by preventing its deadenylation by Smaug/CCR4. *Development.* 2006; 133:4573–83. [PubMed: 17050620]

14. Bushati N, Stark A, Brennecke J, Cohen SM. Temporal reciprocity of miRNAs and their targets during the maternal-to-zygotic transition in *Drosophila*. *Curr Biol*. 2008; 18:501–6. [PubMed: 18394895]
15. Gavis ER, Lehmann R. Translational regulation of nanos by RNA localization. *Nature*. 1994; 369:315–8. [PubMed: 7514276]
16. Dahanukar A, Wharton RP. The Nanos gradient in *Drosophila* embryos is generated by translational regulation. *Genes Dev*. 1996; 10:2610–20. [PubMed: 8895662]
17. Dahanukar A, Walker JA, Wharton RP. Smaug, a novel RNA-binding protein that operates a translational switch in *Drosophila*. *Mol Cell*. 1999; 4:209–18. [PubMed: 10488336]
18. Giraldez AJ, et al. Zebrafish MiR-430 Promotes Deadenylation and Clearance of Maternal mRNAs. *Science*. 2006
19. Chung WJ, Okamura K, Martin R, Lai EC. Endogenous RNA interference provides a somatic defense against *Drosophila* transposons. *Curr Biol*. 2008; 18:795–802. [PubMed: 18501606]
20. Brennecke J, et al. An epigenetic role for maternally inherited piRNAs in transposon silencing. *Science*. 2008; 322:1387–92. [PubMed: 19039138]
21. Gunawardane LS, et al. A slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science*. 2007; 315:1587–90. [PubMed: 17322028]
22. Li C, et al. Collapse of germline piRNAs in the absence of Argonaute3 reveals somatic piRNAs in flies. *Cell*. 2009; 137:509–21. [PubMed: 19395009]
23. Cook HA, Koppetsch BS, Wu J, Theurkauf WE. The *Drosophila* SDE3 homolog armitage is required for oskar mRNA silencing and embryonic axis specification. *Cell*. 2004; 116:817–29. [PubMed: 15035984]
24. Malone CD, et al. Specialized piRNA pathways act in germline and somatic tissues of the *Drosophila* ovary. *Cell*. 2009; 137:522–35. [PubMed: 19395010]
25. Harris AN, Macdonald PM. Aubergine encodes a *Drosophila* polar granule component required for pole cell formation and related to eIF2C. *Development*. 2001; 128:2823–32. [PubMed: 11526087]
26. Megosh HB, Cox DN, Campbell C, Lin H. The role of PIWI and the miRNA machinery in *Drosophila* germline determination. *Curr Biol*. 2006; 16:1884–94. [PubMed: 16949822]
27. Lehmann R, Nusslein-Volhard C. The maternal gene nanos has a central role in posterior pattern formation of the *Drosophila* embryo. *Development*. 1991; 112:679–91. [PubMed: 1935684]
28. Leaman D, et al. Antisense-mediated depletion reveals essential and specific functions of microRNAs in *Drosophila* development. *Cell*. 2005; 121:1097–108. [PubMed: 15989958]
29. Saito K, et al. A regulatory circuit for piwi by the large Maf gene traffic jam in *Drosophila*. *Nature*. 2009; 461:1296–9. [PubMed: 19812547]
30. Robine N, et al. A broadly conserved pathway generates 3'UTR-directed primary piRNAs. *Curr Biol*. 2009; 19:2066–76. [PubMed: 20022248]
31. Schupbach T, Wieschaus E. Female sterile mutations on the second chromosome of *Drosophila melanogaster*. II. Mutations blocking oogenesis or altering egg morphology. *Genetics*. 1991; 129:1119–36. [PubMed: 1783295]
32. Gillespie DE, Berg CA. Homeless is required for RNA localization in *Drosophila* oogenesis and encodes a new member of the DE-H family of RNA-dependent ATPases. *Genes Dev*. 1995; 9:2495–508. [PubMed: 7590230]
33. Chou TB, Perrimon N. The autosomal FLP-DFS technique for generating germline mosaics in *Drosophila melanogaster*. *Genetics*. 1996; 144:1673–1679. [PubMed: 8978054]
34. Wang C, Dickinson LK, Lehmann R. Genetics of nanos localization in *Drosophila*. *Dev Dyn*. 1994; 199:103–15. [PubMed: 7515724]
35. Rorth P. Gal4 in the *Drosophila* female germline. *Mech Dev*. 1998; 78:113–118. [PubMed: 9858703]
36. Juge F, Zaessinger S, Temme C, Wahle E, Simonelig M. Control of poly(A) polymerase level is essential to cytoplasmic polyadenylation and early development in *Drosophila*. *EMBO J*. 2002; 21:6603–6613. [PubMed: 12456666]

37. Pelisson A, Sarot E, Payen-Groschene G, Bucheton A. A novel repeat-associated small interfering RNA-mediated silencing pathway downregulates complementary sense gypsy transcripts in somatic cells of the *Drosophila* ovary. *J Virol.* 2007; 81:1951–60. [PubMed: 17135323]
38. Benoit B, et al. An essential cytoplasmic function for the nuclear poly(A) binding protein, PABP2, in poly(A) tail length control and early development in *Drosophila*. *Dev Cell.* 2005; 9:511–522. [PubMed: 16198293]
39. Benoit B, et al. The *Drosophila* poly(A)-binding protein II is ubiquitous throughout *Drosophila* development and has the same function in mRNA polyadenylation as its bovine homolog *in vitro*. *Nucleic Acids Res.* 1999; 27:3771–3778. [PubMed: 10481015]
40. Temme C, Zaessinger S, Meyer S, Simonelig M, Wahle E. A complex containing the CCR4 and CAF1 proteins is involved in mRNA deadenylation in *Drosophila*. *EMBO J.* 2004; 23:2862–2871. [PubMed: 15215893]

**Figure 1.**

The piRNA pathway is required for *nos* mRNA deadenylation and decay as well as translational repression in the bulk cytoplasm of the embryo. **(a, b)** PAT assays and RT-QPCR of *nos* mRNA. Mutant females of the indicated genotypes were crossed with wild-type males. The *sop* mRNA was used as a control in **a**. **(b)** Levels of *nos* mRNA in 2–3 hour and 3–4 hour embryos. Mean value of three quantifications, error bars correspond to s.d. **(c)** *In situ* hybridizations of *nos* mRNA. **(d)** Immunostaining of embryos with anti-Nos antibody. **(e)** Cuticle preparations of *piwi¹* embryos showing head defects (rudimentary head skeleton (top panel), head skeleton replaced by a hole (bottom panel)). 2% of embryos from *piwi¹* germline clones produced a cuticle (n=1060), among those, 22/23 had strong head defects. No embryos from *aub^{N11}/aub^{HN2}* (n=1230) or *aub^{QC42}/aub^{HN2}* (n=813) females produced a cuticle.

**Figure 2.**

Aub is present in the bulk of the embryo and the piRNA pathway is required for CCR4 and Smg cytoplasmic distributions. **(a)** Confocal images of cytoplasmic expression of Aub in the embryo. Syncytial blastoderm embryo at nuclear cycle 11, anterior is to the left. Pole cells of the same embryo, at the same setting (middle Aub panel) and at lower intensity (right Aub panel)^{20,25}. DAPI staining (right panel). **(b)** Double immunostaining of embryos at nuclear cycles 11/12 with anti-Aub and anti-Smg, or anti-Aub and anti-CCR4. Arrows indicate examples of small foci showing colocalisation in **b** and **c**. **(c)** Smg and CCR4 cytoplasmic


distributions are affected in *aub* and *spn-E* mutant embryos. Double immunostaining of embryos at nuclear cycle 11 with anti-CCR4 and anti-Smg. **(d)** Western blots of proteins from 0–2 hour embryos revealed with anti-Smg and anti-CCR4. α -tubulin (Tub) was used as a loading control.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Figure 3.**

Aub, Ago3, Smg, CCR4 and *nos* mRNA are present in a common complex in the bulk of the embryo. **(a)** Co-immunoprecipitations of Smg, CCR4 and Ago3 with Aub in 0–2 hour embryo extracts. Anti-Aub and anti-GFP were used for immunoprecipitations in wild-type, *osk*⁵⁴ and GFP-Aub expressing embryos, respectively. **(b)** Co-immunoprecipitations of CCR4, Aub and Ago3 with Smg in 0–2 hour wild-type embryo extracts. The asterisks indicate immunoglobulins. **(c)** Quantification of *nos* mRNA enrichment in Aub and Smg immunoprecipitations. Extracts from 0–2 hour wild-type or *osk*⁵⁴ embryos were immunoprecipitated with anti-Aub (rabbit), or anti-Smg. For quantifications performed by RT-QPCR, the ratio of *nos* mRNA/*rp49* mRNA was set to 1 in the mock immunoprecipitation. Mean value of three quantifications, error bars correspond to s.d. *rp49* was used as a control mRNA.




Figure 4. piRNAs target a specific region in *nos* 3'-UTR which is required for *nos* mRNA deadenylation. (a) Schematic representation of *nos* 3'-UTR. The regions deleted in *nos* genomic transgenes (*nos*(Δ)) are indicated on the upper line¹⁶. SRE, piRNA and miRNA target sites are indicated. Predictions of miRNA targeted regions are from miRBase (miR-31a, miR-314 and miR-263b from proximal to distal). piRNA occurrences in the data sets^{19,20} are indicated. (b) Northern blots of 0–2 hour embryos probed with riboprobes corresponding to sense *nos* 3'-UTR (position 403–844) (left panel) and to antisense 412 piRNA (right panel). Anti-GFP immunoprecipitations (GFP IP) were performed using wild-type and GFP-Aub expressing embryos. (c) *nos* PAT assays. For *nos*(Δ 3) the fragment

amplified in the PAT assay is shorter than the fragment amplified in the other *nos* PAT assays (Supplementary Figure 9). **(d)** Quantification of *nos* mRNA levels from the *nos*($\Delta 3$) transgene by RT-QPCR. Mean value of three quantifications, error bars correspond to s.d. **(e)** Cuticle preparations of embryos from *nos*($\Delta 3$) females (lack of head skeleton). **(f)** PAT assays of embryos with *nos* genomic transgenes in which sequences complementary to *412* piRNA, *roo* piRNA, or both sequences have been deleted. The *sop* mRNA was used as a control in **e** and **f**. **(g)** Injection of 2' O-methyl anti-piRNA in embryos. Control injections were with injection buffer alone or with the irrelevant anti-miR129. Examples of cuticles following injections of anti-miR129 (wild-type head skeleton), anti-pi(*412*) and anti-pi(*roo*) (affected head skeleton).