MATRIX LI-YAU-HAMILTON ESTIMATES UNDER RICCI FLOW AND PARABOLIC FREQUENCY

XIAOLONG LI AND QI S. ZHANG

ABSTRACT. We prove matrix Li-Yau-Hamilton estimates for positive solutions to the heat equation and the backward conjugate heat equation, both coupled with the Ricci flow. We then apply these estimates to establish the monotonicity of parabolic frequencies up to correction factors. As applications, we obtain some unique continuation results under the nonnegativity of sectional or complex sectional curvature.

1. INTRODUCTION

1.1. Matrix Li-Yau-Hamilton Estimates. In their seminal paper [LY86], P. Li and S.-T. Yau developed fundamental gradient estimates for positive solutions to the heat equation on a Riemannian manifold. In particular, they proved that if $u: M^n \times [0, \infty) \to \mathbb{R}$ is a positive solution to the heat equation

(1.1)
$$u_t - \Delta_\sigma u = 0$$

on an *n*-dimensional complete Riemannian manifold (M^n, g) with nonnegative Ricci curvature, then

(1.2)
$$\frac{u_t}{u} - \frac{|\nabla u|^2}{u^2} + \frac{n}{2t} = \Delta \log u + \frac{n}{2t} \ge 0$$

for all $(x,t) \in M \times (0,\infty)$. Remarkably, the equality in (1.2) is achieved on the heat kernel on the Euclidean space \mathbb{R}^n . The Li-Yau estimate is also called a differential Harnack inequality since integrating it yields a sharp version of the classical Harnack inequality originated from Moser [Mos64]. The Li-Yau estimate and its generalizations in various settings provide a versatile tool for studying the analytical, topological, and geometrical properties of manifolds (see for instance the classical books [SY94] and [Li12]).

Under the stronger assumption that (M^n, g) has nonnegative sectional curvature and parallel Ricci curvature, Hamilton [Ham93b] extended the Li-Yau estimate to the full matrix version

(1.3)
$$\nabla_i \nabla_j \log u + \frac{1}{2t} g_{ij} \ge 0.$$

²⁰²⁰ Mathematics Subject Classification. 53E20 (Primary), 58J35, 35K05 (Secondary).

Key words and phrases. Li-Yau-Hamilton estimates, matrix Harnack inequality, parabolic frequency, unique continuation, conjugate heat equation, Ricci flow.

The first author's research is partially supported by Simons Collaboration Grant #962228 and a start-up grant at Wichita State University.

The second author's research is partially supported by Simons Collaboration Grant #710364.

LI AND ZHANG

Note that the trace of (1.3) is (1.2). Later on, Chow and Hamilton [CH97] further extended (1.2) and (1.3) to the constrained case under the same curvature assumptions, and discovered new linear Harnack estimates.

When (M^n, g) is a complete Kähler manifold with nonnegative bisectional curvature, Cao and Ni [CN05] proved the matrix inequality

(1.4)
$$\nabla_{\alpha}\nabla_{\bar{\beta}}\log u + \frac{1}{t}g_{\alpha\bar{\beta}} \ge 0,$$

and they called it a matrix Li-Yau-Hamilton estimate. Inspired by Chow's interpolation consideration [Cho98] of Li-Yau's and Hamilton's Harnack inequalities on a surface, Chow and Ni [Ni07, Theorem 2.2] proved that if $(M^n, g(t))$ is a complete solution to Kähler-Ricci flow with bounded nonnegative bisectional curvature and if u is a positive solution to the forward conjugate heat equation

(1.5)
$$u_t - \Delta_{a(t)} u = R u$$

then

(1.6)
$$R_{\alpha\bar{\beta}} + \nabla_{\alpha}\nabla_{\bar{\beta}}\log u + \frac{1}{t}g_{\alpha\bar{\beta}} \ge 0.$$

The equality holds if and only if $(M^n, g(t))$ is an expanding Kähler-Ricci soliton. These results were generalized to the constrained case by Ren, Yao, Shen, and Zhang [RYSZ15].

When the metrics are evolved by the Ricci flow (see [Ham82] or [CLN06])

$$\partial_t g = -2 \operatorname{Ric},$$

Perelman [Per02] discovered a spectacular differential Harnack estimate for the fundamental solution to the backward conjugate heat equation

(1.7)
$$u_t + \Delta_{q(t)} u = Ru,$$

where R denotes the scalar curvature of $(M^n, g(t))$. Astoundingly, his estimate did not require any curvature conditions. For more information on matrix Li-Yau-Hamilton estimates and differential Harnack estimates, as well as their important applications in geometry, we refer the reader to Chow's survey [Cho22], Ni's survey [Ni08], and the monographs [CCG⁺08, Chapters 15-16] and [CCG⁺10, Chapters 23-26] by Chow, etc.

In this paper, we first extend Hamilton's matrix estimate (1.3) for static metrics to the Ricci flow case. Our estimate does not require the parallel Ricci curvature condition and thus should be more applicable.

Theorem 1.1. Let $(M^n, g(t)), t \in [0, T]$, be a complete solution to the Ricci flow. Let $u : M^n \times [0, T] \to \mathbb{R}$ be a positive solution to the heat equation

(1.8)
$$u_t - \Delta_{g(t)} u = 0$$

Suppose that $(M^n, g(t))$ has nonnegative sectional curvature and $\text{Ric} \leq \kappa g$ for some constant $\kappa > 0$. Then

(1.9)
$$\nabla_i \nabla_j \log u + \frac{\kappa}{1 - e^{-2\kappa t}} g_{ij} \ge 0,$$

for all $(x,t) \in M \times (0,T)$.

Remark 1.1. Note that (1.9) can be restated equivalently as

$$\nabla_i \nabla_j u + \frac{\kappa u}{1 - e^{-2\kappa t}} g_{ij} + \nabla_i u V_j + \nabla_j u V_i + u V_i V_j \ge 0$$

for any vector field V by choosing the optimal vector field $V = -\nabla \log u$. Other matrix Li-Yau-Hamilton estimates also admit such equivalent restatements.

Remark 1.2. Note that (1.9) is asymptotically sharp as $t \to 0^+$. To see this, one notices

(1.10)
$$\frac{1}{2t} < \frac{\kappa}{1 - e^{-2\kappa t}} < \frac{1}{2t} + \kappa$$

for all t > 0 and $\kappa > 0$. Applying Theorem 1.1 to $(M^n, g) = (\mathbb{R}^n, \delta_{ij})$ and letting $\kappa \to 0^+$ produce $\nabla_i \nabla_j \log u \ge \frac{1}{2t} \delta_{ij}$, for which the equality is achieved when u is the heat kernel on \mathbb{R}^n .

Remark 1.3. For proving the unique continuation result in Corollary 1.8, it is important to obtain a lower bound for $\nabla_i \nabla_j \log u$ that is asymptotic to $\frac{1}{2t}g_{ij}$ as $t \to 0^+$.

Tracing (1.9) yields a gradient estimate.

Corollary 1.2. Under the same assumptions as in Theorem 1.1, we have

(1.11)
$$\frac{u_t}{u} - \frac{|\nabla u|^2}{u^2} + \frac{n\kappa}{1 - e^{-2\kappa t}} = \Delta \log u + \frac{n\kappa}{1 - e^{-2\kappa t}} \ge 0.$$

On a general compact manifold, Hamilton [Ham93b] proved that for any positive solution u to (1.1), there exist constants B and C depending only on the geometry of M (in particular the diameter, the volume, and the curvature and covariant derivative of the Ricci curvature) such that $t^{n/2}u \leq B$ and

(1.12)
$$\nabla_i \nabla_j \log u + \frac{1}{2t} g_{ij} + C \left(1 + \log \left(\frac{B}{t^{n/2} u} \right) \right) g_{ij} \ge 0.$$

Here, we also establish such a result for a general compact Ricci flow.

Theorem 1.3. Let $(M^n, g(t)), t \in [0, T]$, be a compact solution to the Ricci flow. Let $u : M^n \times [0, T] \to \mathbb{R}$ be a positive solution to the heat equation (1.8). Suppose the sectional curvatures of $(M^n, g(t))$ are bounded by K for some K > 0. Then

(1.13)
$$\nabla_i \nabla_j \log u + \left(\frac{1}{2t} + \frac{1}{t}\beta(t, n, K)\right) g_{ij} \ge 0,$$

where

$$\beta(t, n, K) = 4\sqrt{nKt} + C_2(K+1)t + C_1\sqrt{K} \operatorname{diam}$$

Here $C_1 > 0$ is a numerical constant, $C_2 > 0$ depends only on the dimension and the non-collapsing constant $v_0 = \inf\{|B(x, 1, g(0))|_{g(0)} : x \in M^n\}$, and

diam :=
$$\sup_{t \in [0,T]} \operatorname{diam}(M^n, g(t)).$$

Compared to (1.12) for static metrics, our estimate (1.13) does not depend on the covariant derivatives of Ricci curvature. We have also made the dependence of the constants on curvature and the diameter more explicit. The proof of Theorem 1.3 is more involved than that of Theorem 1.1, and the key steps are to establish bounds for heat kernel under Ricci flow, $\nabla \log u$, and $\Delta \log u$. In contrast to (1.12)

LI AND ZHANG

having the term $\log B/u$ whose order is not clear, (1.13) implies a lower bound of $\nabla_i \nabla_j \log u$ that is asymptotic to C/t as $t \to 0^+$, where $C = \frac{1}{2} + C_1 \sqrt{K}$ diam. Therefore, (1.13) can be regarded as a sharp version of Hamilton's estimate (1.12), just like [Zha21, Theorem 1.1] is a sharp version of the Li-Yau estimate [LY86, Theorem 1.3]. Finally, we remark that the C/t-type bound does not hold for noncompact manifolds in general, such as on \mathbb{H}^3 , the three-dimensional hyperbolic space, as explained in [Zha21].

A similar argument yields an improvement of Hamilton's classical matrix Harnack inequality (1.12) in the static case on compact manifolds.

Theorem 1.4. Let (M^n, g) be a closed Riemannian manifold and let $u : M^n \times [0,T] \to \mathbb{R}$ be a positive solution to the heat equation (1.1). Suppose that the sectional curvatures of M are bounded by K and $|\nabla \operatorname{Ric}| \leq L$, for some K, L > 0. Then

$$(1.14)\nabla_i \nabla_j \log u + \left(\frac{1}{2t} + (2n-1)K + \frac{\sqrt{3}}{2}L^{\frac{2}{3}} + \frac{1}{2t}\gamma(t,n,K,L)\right)g_{ij} \ge 0$$

for all $(x,t) \in M \times (0,T)$, where

$$\begin{split} \gamma(t,n,K,L) &:= \sqrt{nKt(2+(n-1)Kt)} + \sqrt{C_3(K+L^{\frac{2}{3}})t(1+Kt)(1+K+Kt)} \\ &+ \left(2K(2+(n-1)Kt) + \frac{3}{2}L^{\frac{2}{3}}(1+(n-1)Kt)\right) \text{Diam}, \end{split}$$

 $C_3 > 0$ depends only on the dimension n, and Diam denotes the diameter of (M^n, g) .

Notice that Hamilton's original inequality (1.12) has the term $C \log \left(\frac{B}{t^{\frac{2}{2}}u}\right)$, where B and C depend on the geometry of the manifold and B is greater than $t^{\frac{n}{2}}u$, which itself is an additional assumption. The constant C is equal to zero only when M has nonnegative sectional curvature and parallel Ricci curvature. Otherwise, for this log term, we do not have any definite control on the order q of $-t^{-q}$ coming out of this term, for general positive solutions, making this lower bound less practical. In Theorem 1.4, we manage to replace this term with a C/t term, with C depending only on K, L, and Diam, which is of the correct order for t.

Next, we prove a matrix Li-Yau-Hamilton estimate for positive solutions to the backward conjugate heat equation coupled with the Ricci flow.

Theorem 1.5. Let $(M^n, g(t))$, $t \in [0, T]$, be a solution to the Ricci flow with nonnegative complex sectional curvature and Ric $\leq \kappa g$ for some $\kappa > 0$. Let u : $M^n \times [0,T] \to \mathbb{R}$ be a positive solution to the backward conjugate heat equation (1.7) on $M \times [0,T]$. Suppose $\eta : (0,T) \to (0,\infty)$ is a C^1 function satisfying the ordinary differential inequality

(1.15)
$$\eta' \le 2\eta^2 - 2\kappa\eta - \frac{\kappa}{t}$$

on (0,T) and that $\eta(t) \to \infty$ as $t \to T$. Then

(1.16)
$$R_{ij} - \nabla_i \nabla_j \log u - \eta g_{ij} \le 0$$

for all $(x,t) \in M \times (0,T)$, where R_{ij} denotes the Ricci curvature.

Remark 1.4. On ancient Ricci flows, we can get rid of the $\frac{\kappa}{t}$ term in (1.15) and prove the cleaner estimate

$$R_{ij} - \nabla_i \nabla_j \log u - \frac{\kappa}{1 - e^{-2\kappa(T-t)}} g_{ij} \le 0.$$

See Theorem 6.3 for details.

We give an explicit choice of $\eta(t)$.

Corollary 1.6. Under the same assumptions as in Theorem 1.5, we have

(1.17)
$$R_{ij} - \nabla_i \nabla_j \log u - \left(\frac{\kappa}{1 - e^{-2\kappa(T-t)}} + \sqrt{\frac{\kappa}{2t}}\right) g_{ij} \le 0.$$

One motivation for bounding $R_{ij} - \nabla_i \nabla_j \log u$ from above comes from the work of Baldauf and Kim [BK22]. They defined a parabolic frequency under Ricci flow and proved its monotonicity with a correction factor that depends on the upper bound of $R_{ij} - \nabla_i \nabla_j \log K$, where K is the fundamental solution to (1.7). As an application of (1.17), we give an explicit correction factor in Proposition 1.10 in the nonnegative complex sectional curvature case. In addition, we believe such matrix Li-Yau-Hamilton estimates are of their own interest and should be useful in other situations.

We need to assume nonnegative complex sectional curvature in Theorem 1.5 because the proof uses Brendle's generalization [Bre09] of Hamilton's Harnack inequality for the Ricci flow [Ham93a]. This feature is shared by the proof of (1.6) in [Ni07], which makes use of H.D. Cao's Harnack estimate for the Kähler-Ricci flow [Cao92]. We also note that it suffices to assume (M, g(0)) has bounded nonnegative complex sectional curvature, as nonnegative complex sectional curvature is preserved by Ricci flows with bounded curvature (see Brendle and Schoen [BS09] and Ni and Wolfson [NW07]).

Finally, we would like to mention that since the pioneer works of Li and Yau [LY86], Hamilton [Ham93b], Perelman [Per02], and others, various gradient and Hessian estimates for positive solutions to heat-type equations, with either fixed or time-dependent metrics, have been established by many authors, including Guen-ther [Gue02], Ni [Ni04b, Ni04a], Cao and Ni [CN05], Ni [Ni07], Kotschwar [Kot07], the second author [Zha06], Souplet and the second author [SZ06], Kuang and the second author [KZ08], Cao [Cao08], X. Cao and Hamilton [CH09], Liu [Liu09], Bailesteanu, X. Cao, and Pulemotov [BCP10], X. Cao and the second author [CZ11], Li and Xu [LX11], Han and the second author [HZ16], Zhu and the second author [ZZ17, ZZ18], Huang [Hua21], and Yu and Zhao [YZ20], just to name a few. Our matrix Li-Yau-Hamilton estimates are new additions to the literature.

1.2. **Parabolic Frequency.** The elliptic frequency

$$I_A(r) = \frac{r \int_{B_r(p)} |\nabla u|^2 dx}{\int_{\partial B_r(p)} u^2 dA}$$

for a harmonic function u on \mathbb{R}^n was introduced by Almgren [Alm79]. He used its monotonicity to study the local regularity of (multiple-valued) harmonic functions and minimal surfaces. The monotonicity of $I_A(r)$ also played an important role in studying unique continuation properties of elliptic operators by Garafalo and Lin

LI AND ZHANG

[GL86a, GL87] and in estimating the size of nodal sets of solutions to elliptic and parabolic equations by Lin [Lin91]. When \mathbb{R}^n is replaced by a Riemannian manifold, Garofalo and Lin [GL86b] proved that there exist constants R_0 and Λ , depending only on the Riemannian metric, such that $e^{\Lambda r} I_A(r)$ is monotone nondecreasing in $(0, R_0)$ (see also Mangoubi [Man13, Theorem 2.2]). Frequency monotonicity is also crucial in the work of Logunov [Log18a, Log18b] in estimating the size of nodal sets for harmonic functions and eigenfunctions on manifolds. In addition, frequency functions also play a crucial role in studying the dimension of the space of harmonic functions of polynomial growth on complete noncompact manifolds; see Colding and Minicozzi [CM97a, CM97b], G. Xu [Xu16], J.Y. Wu and P. Wu [WW23], Mai and Ou [MO22], and the references therein. For more applications, we refer the reader to the books [HL] and [Zel08].

Poon [Poo96] introduced the parabolic frequency

$$I_P(t) = \frac{t \int_{\mathbb{R}^n} |\nabla u|^2(x, T - t) G(x, x_0, t) \, dx}{\int_{\mathbb{R}^n} u^2(x, T - t) G(x, x_0, t) \, dx},$$

where u solves the heat equation on $\mathbb{R}^n \times [0, T]$ and $G(x, x_0, t)$ is the heat kernel with a pole at $(x_0, 0)$. He proved that $I_P(t)$ is monotone nondecreasing and derived some unique continuation results out of it. The monotonicity of $I_P(t)$ remains valid when \mathbb{R}^n is replaced by a complete Riemannian manifold with nonnegative sectional curvature and parallel Ricci curvature, as remarked by Poon [Poo96, page 530] and proved independently by Ni [Ni15]. The curvature conditions are needed to use Hamilton's matrix estimate (1.3).

Without assuming the restrictive parallel Ricci condition, Wang and the first author [LW19] showed that $te^{\sqrt{t}}I_P(t)$ is monotone nondecreasing for a short period of time on compact manifolds with nonnegative sectional curvature, which also produces a unique continuation result. They also defined the parabolic frequency

$$I_{LW}(t) = \frac{t \int_M |\nabla v(x,t)|^2 R d\mu_{g(t)}}{\int_M v^2(x,t) R d\mu_{g(t)}}$$

where v solves the backward heat equation $v_t + \Delta_{g(t)}v = 0$ coupled with a twodimensional Ricci flow with positive scalar curvature. Using that R satisfies the forward heat equation (1.5) and admits a matrix Li-Yau-Hamilton estimate due to Hamilton (see [CLN06, Proposition 10.20]), they showed that $I_{LW}(t)$ is monotone nondecreasing.

Colding and Minicozzi [CM22] proved that the parabolic frequency

$$I_{CM}(t) = \frac{\int_M |\nabla u(x,t)|^2 e^{-f} d\mu_g}{\int_M u^2(x,t) e^{-f} d\mu_g},$$

where f is a smooth function on a Riemannian manifold M^n and $u: M^n \times [0, T] \to \mathbb{R}^N$ solves the weighted heat equation $u_t - \Delta_f u = 0$, is monotone nonincreasing without any curvature assumptions. The special case $f \equiv 1$ and M^n being a bounded domain in \mathbb{R}^n was treated in [Eva10, pages 61-62] to prove the backward uniqueness of the heat equation with specified boundary values. They also defined a parabolic frequency for shrinking gradient Ricci solitons and showed its monotonicity with no curvature restrictions.

For a general Ricci flow, Baldauf and Kim [BK22] defined the parabolic frequency

$$I_{BK}(t) := \frac{(T-t)\int_M |\nabla u(x,t)|^2 K(x,x_0,t) d\mu_{g(t)}}{\int_M u^2(x,t) K(x,x_0,t) d\mu_{g(t)}}$$

where K is the backward conjugate heat kernel with a pole at (x_0, T) and u solves the heat equation (1.8). They were able to show that $e^{\int \frac{1-k(t)}{T-t}dt} \cdot I_{BK}(t)$ is monotone nonincreasing, where k(t) is a time-dependent function such that

$$\operatorname{Ric} -\nabla^2 \log K \le \frac{k(t)}{2(T-t)}$$

More recently, C. Li, Y. Li, and K. Xu [LLX22] studied the monotonicity of $I_{BK}(t)$ and its generalizations under the Ricci flow and the Ricci-harmonic flow. They obtained monotonicity formulas with correction factors depending on the bounds of the Bakry-Émery Ricci curvature or the Ricci curvature. It is also worth mentioning that H.Y. Liu and P. Xu [LX22] investigated the monotonicity of a parabolic frequency for weighted *p*-Laplacian heat equation with $p \ge 2$ on Riemannian manifolds and obtained generalizations of [CM22]. Using (1.6), they generalized a frequency monotonicity formula of Ni [Ni15] on Kähler manifolds to the setting of Kähler-Ricci flow.

In this paper, we define a parabolic frequency for solutions to the backward conjugate heat equation (1.7) coupled with the Ricci flow and prove its monotonicity up to certain correction factors. We shall use $G(x, x_0, t)$, the heat kernel with a pole at $(x_0, 0)$, as a weight and define the following quantities:

$$\begin{split} I(t) &= \int_{M} u^{2}(x,t) G(x,x_{0},t) d\mu_{g(t)}, \\ D(t) &= \int_{M} |\nabla u(x,t)|^{2} G(x,x_{0},t) d\mu_{g(t)}, \\ S(t) &= \int_{M} u^{2}(x,t) R(x,t) G(x,x_{0},t) d\mu_{g(t)} \end{split}$$

The first two quantities are direct generalizations of the terms in Poon's parabolic frequency $I_P(t)$ in the static case. The third one is new due to the Ricci flow. A natural generalization of Poon's frequency $I_P(t)$ is

(1.18)
$$F(t) := \frac{I'(t)}{I(t)} = \frac{2D(t) + S(t)}{I(t)}.$$

In the nonnegative sectional curvature case, we prove that

Theorem 1.7. Let $(M^n, g(t))$, $t \in [0, T]$, be a complete Ricci flow and let $u : M^n \times [0, T] \to \mathbb{R}$ be a solution to the backward conjugate heat equation (1.7). Suppose that $(M^n, g(t))$ has nonnegative sectional curvature and Ric $\leq \kappa g$ for some constant $\kappa > 0$. Then

(1.19)
$$e^{(n-2)\kappa t}(1-e^{-2\kappa t})F(t)$$

where F(t) is defined in (1.18), in monotone nondecreasing on [0, T].

We point out that Theorem 1.7 covers, by letting $\kappa \to 0^+$, Poon's frequency monotonicity on \mathbb{R}^n in [Poo96]. It also implies a unique continuation result (see

Lin [Lin90], Poon [Poo96], and Vessella [Ves03] for unique continuation results for parabolic equations).

Corollary 1.8. Let $(M^n, g(t)), t \in [0, T]$, be a complete Ricci flow with nonnegative sectional curvature and Ric $\leq \kappa g$ for some $\kappa > 0$. Suppose that a solution u(x, t) of the backward conjugate heat equation (1.7) on $M^n \times [0, T]$ vanishes of infinity order at $(x_0, t_0) \in M \times (0, T)$, in the sense that

$$|u(x,t)| \le O\left(d_t^2(x,x_0) + |t-t_0|\right)^N$$

for all positive integer N and all (x,t) near (x_0,t_0) . Then $u \equiv 0$ in $M \times [0,T]$.

Under general compact Ricci flows, we prove that F(t) is monotone up to an implicit correction factor.

Theorem 1.9. Let $(M^n, g(t)), t \in [0, T]$, be a compact solution to the Ricci flow with sectional curvatures bounded by K for some K > 0. Let $u : M^n \times [0, T] \to \mathbb{R}$ be a solution to the backward conjugate heat equation (1.7). Then, for any T > 0, there is a power $p = p(T, n, K, v_0, \text{diam}) > 0$ such that

(1.20)
$$t^p(F(t) + Z_0),$$

where F(t) is defined in (1.18), is monotone nondecreasing on [0,T]. Here $Z_0 = Z_0(T, n, K, v_0, \text{diam})$ is any sufficiently large number.

Extra curvature terms arise due to the Ricci flow when proving Theorem 1.9. We handle them using some cancellation property and some Li-Yau estimates for the heat kernel under the Ricci flow, together with the matrix Harnack inequality in Theorem 1.3. Besides the above-mentioned results, we also prove the monotonicity of a parabolic frequency without weight at the end of Section 5 assuming nonnegative Ricci curvature; see Theorem 5.1.

Finally, we apply Theorem 1.5 to prove that

Proposition 1.10. Let $(M^n, g(t))$, $t \in [0, T]$, be a solution to the Ricci flow with nonnegative complex sectional curvature and Ric $\leq \kappa g$ for some $\kappa > 0$. Let u : $M^n \times [0, T] \to \mathbb{R}$ be a solution to the heat equation (1.8) and let $w : M^n \times [0, T] \to$ $(0, \infty)$ be a positive solution to the backward conjugate heat equation (1.7). Then the quantity

(1.21)
$$(e^{2\kappa(T-t)} - 1)e^{-\sqrt{8\kappa t}} \cdot \frac{\int_M |\nabla u(x,t)|^2 w(x,t) d\mu_{g(t)}}{\int_M u^2(x,t) w(x,t) d\mu_{g(t)}}$$

is monotone nonincreasing on [0, T].

As mentioned before, Baldauf and Kim [BK22] proved the monotonicity of $e^{\int \frac{1-k(t)}{T-t} dt} I_{BK}(t)$ under Ricci flow, where k(t) is a time-dependent function such that

$$\operatorname{Ric} -\nabla^2 \log K \le \frac{k(t)}{2(T-t)}$$

However, it is not clear whether such k(t) exists in the complete noncompact case. In the compact case, the existence of k(t) is shown by Huang [Hua21] and it depends on |Rm|, $|\nabla Rm|$ and $|\nabla^2 R|$, but no explicit k(t) is known. Theorem 1.5 and Corollary 1.6 provide an explicit k(t) in the nonnegative complex sectional curvature case. Proposition 1.10 then gives an explicit correction factor in the monotonicity of $I_{BK}(t)$ and it is also applicable to complete noncompact Ricci flows with bounded nonnegative complex sectional curvature. In addition, we also get a unique continuation result in this case (see Corollary 6.6).

Note: Throughout the paper, we assume either M is compact or M is complete with bounded curvature/geometry, and the functions satisfy certain growth conditions so that the integrals are finite and all integration by parts can be justified.

The rest of this article is organized as follows. In Section 2, we derive the evolution equation satisfied by the Hessian of $\log u$, where u is a positive solution to heat-type equations. Section 3 deals with the nonnegative sectional case and proves Theorem 1.1. Section 4 gives the proof of Theorem 1.3. Section 5 is devoted to studying the parabolic frequency and proving Theorem 1.7 and Theorem 1.9. In Section 6, we prove Theorem 1.5 and Proposition 1.10. In Section 7, we prove Theorem 1.4.

2. Evolution Equations

Let $(M^n, g(t)), t \in [0, T]$, be a solution to the Ricci flow. Let $u: M^n \times [0, T] \to \mathbb{R}$ be a positive solution to the heat-type equation

(2.1)
$$(\partial_t - \varepsilon \Delta_{q(t)})u = \delta Ru,$$

where ε and δ are real parameters. We are mainly interested in the heat equation corresponding to $\varepsilon = 1$ and $\delta = 0$ and the backward conjugate heat equation corresponding to $\varepsilon = -1$ and $\delta = 1$, but the calculations in this section are valid for all $\varepsilon, \delta \in \mathbb{R}$.

The main result of this section is the evolution equation satisfied by

$$H_{ij} := \nabla_i \nabla_j \log u.$$

Proposition 2.1. In the setting described above, we have

$$(2.2) \qquad (\partial_t - \varepsilon \Delta) H_{ij} \\ = \delta \nabla_i \nabla_j R + 2\varepsilon \left(H_{ij}^2 + R_{ikjl} \nabla_k v \nabla_l v + \nabla_k H_{ij} \nabla_k v \right) \\ + \varepsilon (2R_{ikjl} H_{kl} - R_{ik} H_{jk} - R_{jk} H_{ik}) \\ + (1 - \varepsilon) (\nabla_i R_{jk} + \nabla_j R_{ik} - \nabla_k R_{ij}) \nabla_k v,$$

where $H_{ij}^2 := H_{ik}H_{jk}$.

We first prove a commutator formula for $\partial_t - \varepsilon \Delta_L$ and $\nabla_i \nabla_j$, where Δ_L denotes the Lichnerowicz Laplacian acting on symmetric two-tensors via

$$\Delta_L h_{ij} = \Delta h_{ij} + 2R_{ikjl}h_{kl} - R_{ik}h_{jk} - R_{jk}h_{ik}.$$

Lemma 2.1. Under the Ricci flow, it holds that

(2.3)
$$(\partial_t - \varepsilon \Delta_L) (\nabla_i \nabla_j f)$$

= $\nabla_i \nabla_j (\partial_t - \varepsilon \Delta) f + (1 - \varepsilon) (\nabla_i R_{jk} + \nabla_j R_{ik} - \nabla_k R_{ij}) \nabla_k f$

for any smooth function f(x,t).

Proof. The cases $\varepsilon = \pm 1$ are proved in [CLN06], so we only do a slight generalization here. The time derivatives of the Christoffel symbols Γ_{ij}^k under the Ricci flow are given by (see [CLN06, page 108])

$$\partial_t \Gamma_{ij}^k = -g^{kl} (\nabla_i R_{jl} + \nabla_j R_{il} - \nabla_l R_{ij}).$$

It follows that

$$\partial_t (\nabla_i \nabla_j f) = \nabla_i \nabla_j (\partial_t f) + (\nabla_i R_{jk} + \nabla_j R_{ik} - \nabla_l R_{ij}) \nabla_k f.$$

Second, we have

$$\Delta_L \nabla_i \nabla_j f = \nabla_i \nabla_j \Delta f + (\nabla_i R_{il} + \nabla_j R_{il} - \nabla_l R_{ij}) \nabla_l f.$$

This can be seen by commuting covariant derivatives as follows

$$\begin{split} \nabla_i \nabla_j \Delta f &= \nabla_i \nabla_j \nabla_k \nabla_k f \\ &= \nabla_i \nabla_k \nabla_k \nabla_j f - \nabla_i (R_{jl} \nabla_l f) \\ &= \nabla_k \nabla_i \nabla_k \nabla_j f - R_{il} \nabla_l \nabla_l f + R_{ikjl} \nabla_k \nabla_l f \\ &- \nabla_i R_{jl} \nabla_l f - R_{jl} \nabla_i \nabla_l f \\ &= \nabla_k \nabla_k \nabla_i \nabla_j f + \nabla_k (R_{ikjl} \nabla_l f) - R_{il} \nabla_l \nabla_j f + R_{ikjl} \nabla_k \nabla_l f \\ &- \nabla_i R_{jl} \nabla_l f - R_{jl} \nabla_i \nabla_l f \\ &= \Delta \nabla_i \nabla_j f + 2 R_{ikjl} \nabla_k \nabla_l f + \nabla_l R_{ij} \nabla_l f - \nabla_j R_{il} \nabla_l f \\ &- R_{il} \nabla_l \nabla_j f - (\nabla_i R_{jl} \nabla_l f - R_{jl} \nabla_i \nabla_l f) \\ &= \Delta_L \nabla_i \nabla_j f - (\nabla_i R_{jl} + \nabla_j R_{il} - \nabla_l R_{ij}) \nabla_l f, \end{split}$$

where we have used the contracted Bianchi identity

$$\nabla_k R_{ikjl} = \nabla_l R_{ij} - \nabla_j R_{il}.$$

Combining the above two calculations, we obtain (2.3).

We now prove Proposition 2.1.

Proof of Proposition 2.1. For convenience, we write $v = \log u$. One derives from (2.1) that v satisfies the equation

$$(\partial_t - \varepsilon \Delta)v = \varepsilon |\nabla v|^2 + \delta R.$$

We compute that

$$\begin{aligned} \nabla_i \nabla_j |\nabla v|^2 &= 2\nabla_i (\nabla_j \nabla_k v) \nabla_k v + 2\nabla_i \nabla_k v \nabla_j \nabla_k v \\ &= 2\nabla_k (\nabla_i \nabla_j v) \nabla_k v + 2H_{ij}^2 + 2R_{ikjl} \nabla_k v \nabla_l v \\ &= 2H_{ij}^2 + 2R_{ikjl} \nabla_k v \nabla_l v + 2\nabla_k H_{ij} \nabla_k v, \end{aligned}$$

where $H_{ij}^2 := H_{ik}H_{jk}$. Applying the identity (2.3) to f = v yields

$$\begin{aligned} &(\partial_t - \varepsilon \Delta_L) H_{ij} \\ &= \nabla_i \nabla_j (\varepsilon |\nabla v|^2 + \delta R) \\ &+ (1 - \varepsilon) (\nabla_i R_{jk} + \nabla_j R_{ik} - \nabla_k R_{ij}) \nabla_k v \\ &= \delta \nabla_i \nabla_j R + 2\varepsilon \left(H_{ij}^2 + R_{ikjl} \nabla_k v \nabla_l v + \nabla_k H_{ij} \nabla_k v \right) \\ &+ (1 - \varepsilon) (\nabla_i R_{jk} + \nabla_j R_{ik} - \nabla_k R_{ij}) \nabla_k v. \end{aligned}$$

Finally, (2.2) follows from the above identity and

$$\Delta_L H_{ij} = \Delta_L H_{ij} + 2R_{ikjl}H_{kl} - R_{ik}H_{jk} - R_{jk}H_{ik}.$$

The proof is complete.

3. MATRIX HARNACK FOR THE HEAT EQUATION

In this section, we prove Theorem 1.1. The proof of Theorem 1.1 divides into two cases: the compact case and the complete noncompact case.

3.1. The compact case.

Proof of Theorem 1.1. In the compact case, we use Hamilton's tensor maximum principle to prove Theorem 1.1.

Setting $\varepsilon = 1$ and $\delta = 0$ in (2.2), we get that $H_{ij} := \nabla_i \nabla_j \log u$ satisfies

(3.1)
$$(\partial_t - \Delta)H_{ij} = 2H_{ij}^2 + (2R_{ikjl}H_{kl} - R_{ik}H_{jk} - R_{jk}H_{ik}) + 2R_{ikjl}\nabla_k v \nabla_l v + 2\nabla_k H_{ij}\nabla_k v$$

where $H_{ij}^2 := H_{ik}H_{jk}$. We write

(3.2)
$$c(t) := \frac{\kappa}{1 - e^{-2\kappa t}}$$

and define

$$Z_{ij} := H_{ij} + c(t)g_{ij}.$$

Direct calculations using (3.1) and the identity

$$2H_{ij}^{2} + 2R_{ikjl}H_{kl} - R_{ik}H_{jk} - R_{jk}H_{ik}$$

= $2Z_{ij}^{2} - 4cZ_{ij} + 2c^{2}g_{ij} + 2R_{ikjl}Z_{kl} - R_{ik}Z_{jk} - R_{jk}Z_{ik}$

show that

(3.3)
$$(\partial_t - \Delta)Z_{ij} = 2Z_{ij}^2 - 4cZ_{ij} + 2R_{ikjl}Z_{kl} - R_{ik}Z_{jk} - R_{jk}Z_{ik} + 2R_{ikjl}\nabla_k v \nabla_l v + 2\nabla_k Z_{ij}\nabla_k v + (c' + 2c^2 - 2\kappa c)g_{ij} + 2c(\kappa g_{ij} - R_{ij}).$$

Noting that $2R_{ikjl}\nabla_k v\nabla_l v \ge 0$, $R_{ij} \le \kappa g_{ij}$, and c(t) solves the ODE

$$c' = -2c^2 + 2\kappa c,$$

we derive from (3.3) that

$$(3.4) \qquad (\partial_t - \Delta)Z_{ij} \geq 2Z_{ij}^2 - 4cZ_{ij} + 2R_{ikjl}Z_{kl} - R_{ik}Z_{jk} - R_{jk}Z_{ik} + 2\nabla_k Z_{ij}\nabla_k v.$$

Since M is compact and $c(t) \to \infty$ as $t \to 0^+$, we have $Z_{ij} \ge 0$ as $t \to 0^+$. Then the tensor maximum principle of Hamilton [Ham86] implies that $Z_{ij} \ge 0$ for all $t \in [0, T]$, as it is clear that

$$2Z_{ij}^2 - 4cZ_{ij} + 2R_{ikjl}Z_{kl} - R_{ik}Z_{jk} - R_{jk}Z_{ik}$$

is nonnegative at a null-eigenvector of Z_{ij} . The proof is complete.

LI AND ZHANG

3.2. The complete noncompact case. Now we deal with the case that $(M^n, q(t))$, $t \in [0,T]$, is a complete noncompact Ricci flow with nonnegative sectional curvature and Ric $\leq \kappa g$. We note that the uniqueness of solutions to the heat equation (1.8) fails to be true on a complete noncompact manifold. In order to apply Hamilton's tensor maximum principle (see for instance [CCG⁺08, Theorem 12.33] for a version on complete noncompact Ricci flows) to Z_{ij} , one needs to impose some growth condition on the function u and its first and second derivatives. Using an idea in [CN05] and [Ni07], we can, however, get away without assuming any growth conditions on u. The key is that we are working with a positive solution of the heat equation and we can make use of the Li-Yau estimate for u under the Ricci flow (see [CCG⁺10, Theorem 25.9 and Corollary 25.13]) to obtain required growth estimates at any positive time. Then, we can get integral bounds on the first and second derivatives of u via integration by parts. Finally, we use the idea of working with the smallest eigenvalue of the symmetric two-tensor tuZ_{ij} to use a maximum principle for scalar heat equation (see $[CCG^+08, Theorem 12.22]$), avoiding the tensor maximum principle which requires a more restrictive growth condition.

We first prove a growth estimate for u on a slightly smaller time interval using the Li-Yau estimate and its resulting Harnack inequality.

Lemma 3.1. Let $(M^n, g(t))$, $t \in [0, T]$, be a complete solution of the Ricci flow with $-\kappa g \leq \text{Ric} \leq \kappa g$. Let u be a positive solution to the heat equation 1.8 on $M \times [0,T]$. Fix $p \in M$. For any $\delta \in (0,T/3)$, there exist a positive constant A_1 , depending on n, κ, T, δ , and u(p, T) such that

(3.5)
$$u(x,t) \le \exp\left(A_1(d_0^2(x,p)+1)\right)$$

for all $x \in M$ and $t \in [\delta, T - \delta]$, where $d_0(\cdot, \cdot)$ is the distance function with respect to g(0).

Proof. Since $-\kappa g \leq \text{Ric} \leq \kappa g$, we have

$$e^{-2\kappa T}g(0) \le g(t) \le e^{2\kappa T}g(0)$$

for all $t \in [0,T]$. In the Li-Yau estimate stated in [CCG⁺10, Theorem 25.9] and the Harnack inequality stated in [CCG⁺10, Corollary 25.13], we can take $\tilde{g} = g(0)$, $\tilde{C}_0 = e^{2\kappa T}$, Q = 0, and $\varepsilon = \frac{1}{3}$ and let $R \to \infty$. Then, we conclude that there exist positive constants $B_1 = B_1(n,\kappa)$ and $B_2 = B_2(\kappa,T)$, such that

(3.6)
$$\frac{u(x_2, t_2)}{u(x_1, t_1)} \ge e^{-B_1(t_2 - t_1)} \left(\frac{t_2}{t_1}\right)^{-n} \exp\left(-B_2 \frac{d_0^2(x_1, x_2)}{t_2 - t_1}\right)$$

for any $x_1, x_2 \in M$ and $0 < t_1 < t_2 \leq T$. Applying (3.6) with $x = x_1, x_2 = p$, $t_1 = t$, and $t_2 = T$, we get

(3.7)
$$u(x,t) \le u(p,T)e^{B_1(T-t)} \left(\frac{T}{t}\right)^n \exp\left(B_2 \frac{d_0^2(x,p)}{T-t}\right)$$

for all $(x,t) \in M \times (0,T)$. This implies that, for $(x,t) \in M \times [\delta, T-\delta]$, there exists a constant A_1 depending on n, κ, T, δ , and u(p,T) such that (3.5) holds. \Box

Next, we obtain integral bounds for the first and second derivatives of u.

Lemma 3.2. Let $(M^n, g(t))$ and u be the same as in Lemma 3.1. Assume further that $(M^n, g(t))$ has nonnegative scalar curvature. Then there exists $A_2 \ge A_1$ such that

(3.8)
$$\int_{\delta}^{T-\delta} \int_{M} |\nabla u(x,t)|^2 \exp\left(-A_2(d_0^2(x,p)+1)\right) dx dt < \infty$$

and

(3.9)
$$\int_{\delta}^{T-\delta} \int_{M} |\nabla^2 u(x,t)|^2 \exp\left(-A_2(d_0^2(x,p)+1)\right) dx dt < \infty.$$

Proof. We derive from $u_t = \Delta u$ that

$$(\partial_t - \Delta)u^2 = -2|\nabla u|^2.$$

Multiplying both sides by a cut-off function φ^2 (independent of time) and integrating by parts yield

$$\begin{split} & 2\int_{\delta}^{T-\delta}\int_{M}\varphi^{2}|\nabla u|^{2}dxdt \\ = & -\int_{\delta}^{T-\delta}\int_{M}\varphi^{2}(\partial_{t}-\Delta)u^{2}dxdt \\ \leq & \int_{M}\varphi^{2}u^{2}(x,\delta)dx + 4\int_{\delta}^{T-\delta}\int_{M}\varphi u|\nabla\varphi||\nabla u|dxdt \\ \leq & \int_{M}\varphi^{2}u^{2}(x,\delta)dx + 4\int_{\delta}^{T-\delta}\int_{M}|\nabla\varphi|^{2}u^{2}dxdt \\ & +\int_{\delta}^{T-\delta}\int_{M}\varphi^{2}|\nabla u|^{2}dxdt. \end{split}$$

Now (3.8) follows from (3.5). Applying the same argument to

$$(\partial_t - \Delta)|\nabla u|^2 = 2|\nabla^2 u|^2$$

produces (3.9).

Proof of Theorem 1.1: the complete noncompact case. Recall that

$$Z_{ij} := H_{ij} + c(t)g_{ij},$$

where $c(t)=\frac{\kappa}{1-e^{-2\kappa t}},$ satisfies (3.4) under our curvature assumptions. Define

$$\widetilde{Z}_{ij} := t u Z_{ij}$$

Using (3.4), we derive that

$$(3.10) \qquad (\partial_t - \Delta)\widetilde{Z}_{ij} = \frac{1}{t}\widetilde{Z}_{ij} + \frac{2}{tu}\widetilde{Z}_{ij}^2 - 4c(t)\widetilde{Z}_{ij} + 2R_{ikjl}\widetilde{Z}_{kl} - R_{ik}\widetilde{Z}_{jk} - R_{jk}\widetilde{Z}_{ik}.$$

Next, let's consider the function $\alpha(x,t)$ on $M \times [0,T]$ defined by

$$\alpha(x,t) = \inf\{s \ge 0 : \widetilde{Z}_{ij}(x,t) + sg_{ij}(x,t) \ge 0\}.$$

In other words,

(3.11)
$$\alpha(x,t) = \max\{0, -\lambda_1(x,t)\}\$$

on $M \times (0,T)$, where $\lambda_1(x,t)$ is the smallest eigenvalue of \widetilde{Z}_{ij} at (x,t). The key is to show that

$$(3.12) \qquad \qquad (\partial_t - \Delta)\alpha \le 0$$

holds in the following barrier sense: for any $(x,t) \in M \times (0,T)$, we can find a neighborhood $U \subset M \times (0,T)$ of (x,t) and a smooth (lower barrier) function $\phi : U \to \mathbb{R}$ such that $\phi \leq \alpha$ on U, with equality at (x,t), and

$$(3.13)\qquad\qquad (\partial_t - \Delta)\phi \le 0$$

at (x, t). Note that inequality (3.12) holds also in the viscosity sense and in the sense of distributions by standard arguments (see [MMU14] for an elliptic version).

To prove (3.12), we consider $Y_{ij}(x,t) := \tilde{Z}_{ij}(x,t) + \alpha(x,t)g_{ij}(x,t)$. Fix $(x,t) \in M \times (0,T)$. By the definition of $\alpha(x,t)$, we have $Y_{ij} \ge 0$ on $M \times [0,T]$ and there exists a unit vector $e_1 \in T_x M$ such that $Y(e_1, e_1) = 0$. We extend e_1 to an orthonormal basis $\{e_i\}_{i=1}^n$ of $T_x M$ consisting of eigenvector of \tilde{Z}_{ij} such that $\tilde{Z}(e_i) = \lambda_i e_i$ with $\lambda_1 \le \cdots \le \lambda_n$. Next, we extend $\{e_i\}_{i=1}^n$ smoothly in a neighborhood U of (x,t) by parallel translation along radial geodesics using $\nabla^{g(t)}$ and regard the resulting vector fields, still denoted by $\{e_i\}_{i=1}^n$, as stationary in time in the sense that $\partial_t e_i = 0$ for each $1 \le i \le n$.

If $\lambda_1(x,t) > 0$, then $\alpha \equiv 0$ near (x,t) and the barrier function $\phi \equiv 0$ will fulfill (3.13). If $\lambda_1(x,t) \leq 0$, then the function $\phi(x,t) = -\frac{\tilde{Z}(e_1,e_1)}{g(t)(e_1,e_1)}$ is defined in U and gives a lower barrier for $\alpha(x,t)$ in that neighborhood. Therefore, we get using (3.10) that at (x,t),

$$(\partial_t - \Delta)\phi = -(\partial_t - \Delta)\frac{\widetilde{Z}(e_1, e_1)}{g(t)(e_1, e_1)}$$

$$= -\frac{1}{t}\lambda_1 - \frac{2}{tu}\lambda_1^2 + 4c(t)\lambda_1 - 2R_{1k1k}\lambda_k$$

$$\leq \frac{1}{t}\alpha - \frac{2}{tu}\alpha^2 - 4c(t)\alpha + 2\kappa\alpha$$

$$\leq \alpha \left(\frac{1}{t} - 4c(t) + 2\kappa\right)$$

$$\leq 0,$$

where we have used (3.11), the estimate

 $R_{1k1k}\lambda_k = R_{1k1k}(\lambda_k - \lambda_1) + R_{11}\lambda_1 \ge -\alpha R_{11} \ge -\alpha \kappa,$

and the elementary inequality

$$\frac{1}{t} - 4c(t) + 2\kappa < 0 \text{ for } t > 0.$$

Hence, we have proved $(\partial_t - \Delta)\alpha \leq 0$ in the barrier sense.

Without loss of generality, we may assume $u \ge \varepsilon > 0$. This is because once the estimate has been established for $u_{\varepsilon} := u + \varepsilon$, one can then let $\varepsilon \to 0$ and get the estimate for any positive u. By shifting the time from t to $t + \delta$, we have the growth bounds (3.5), (3.8), and (3.9), which implies that there exists b > 0 such that

(3.14)
$$\int_0^T \int_M \exp\left(-bd_0^2(x,p)\right)\alpha^2(x,t)dxdt < \infty,$$

Since $\alpha(x, 0) = 0$ for all $x \in M$, we can use the maximum principle (see [CCG⁺08, Theorem 12.22]) to conclude that $\alpha(x, t) \leq 0$ on $M \times [0, T]$.

4. MATRIX HARNACK FOR THE HEAT EQUATION: THE GENERAL CASE

In this section, we prove Theorem 1.3. Without the nonnegativity of sectional curvatures, we have to estimate the terms involving curvature and derivatives of u and the proof becomes much more involved. Here we employ an idea that has has been used in [LWH⁺], [YZ22], and [Zha21], namely we first prove the estimate for the heat kernel and then derive the estimate for any positive solution to the heat equation.

Proof of Theorem 1.3. The proof is divided into three steps.

Step 1. We derive a partial differential inequality satisfied by the smallest eigenvalue of $Q_{ij} := tH_{ij}$, where $H_{ij} = \nabla_i \nabla_j \log u$ as before.

A straightforward computation using (3.1) shows that Q_{ij} satisfies

$$(4.1) \qquad (\partial_t - \Delta)Q_{ij} = \frac{1}{t}Q_{ij} + \frac{2}{t}Q_{ij}^2 + 2R_{ikjl}Q_{kl} - R_{ik}Q_{jk} - R_{jk}Q_{ik} + 2tR_{ikjl}\nabla_k v\nabla_l v + 2\nabla_k Q_{ij}\nabla_k v.$$

Here and through out this section $v = \log u$.

Let λ_1 be the minimum negative eigenvalue of Q_{ij} in $M^n \times [0, t_0]$ which is reached at the point (x_0, t_0) . Note that we are done with the proof if no such λ_1 exists. Our task is to find a lower bound for λ_1 . Let η be a unit eigenvector with respect to the metric $g(t_0)$ at x_0 . Using parallel transport, we extend η along geodesic rays starting from x_0 so that it becomes a parallel unit vector field in a neighborhood of x_0 with respect to $g(t_0)$. This vector field, still denoted by $\eta = \eta(x)$, is regarded as stationary in the time interval $[0, t_0]$. Now consider the vector field

$$\xi = \xi(x, t) = \frac{\eta(x)}{\|\eta(x)\|_{g(x, t)}}$$

which is a unit one with respect to g(t). In local coordinates, we write $\xi = (\xi_1, ..., \xi_n)$ and we also introduce the scalar function

$$\Lambda = \Lambda(x, t) = \xi_i Q_{ij} \xi_j = \xi^T(Q_{ij}) \xi.$$

Notice that Λ is a smooth function defined in a neighborhood of x_0 on the time interval $[0, t_0]$ and reaches its minimum value λ_1 at the point (x_0, t_0) . Using (4.1), we find that

$$\begin{aligned} &\xi_i[(\partial_t - \Delta)Q_{ij}]\xi_j \\ &= \frac{1}{t}\xi_i Q_{ij}\xi_j + \frac{2}{t}\xi_i Q_{ij}^2\xi_j + 2R_{ikjl}Q_{kl}\xi_i\xi_j - R_{ik}Q_{jk}\xi_i\xi_j - R_{jk}Q_{ik}\xi_i\xi_j \\ &+ 2tR_{ikjl}\nabla_k v\nabla_l v\xi_i\xi_j + 2\nabla_k Q_{ij}\nabla_k v\xi_i\xi_j. \end{aligned}$$

Recall that at $t = t_0$, ξ is a parallel vector field and

$$\partial_t(\xi_i Q_{ij}\xi_j) = \partial_t \left(\frac{\eta_i Q_{ij}\eta_j}{g_{ij}\eta_i\eta_j}\right) = \xi_i \partial_t Q_{ij}\xi_j + 2\xi_i Q_{ij}\xi_j R_{kl}\xi_k\xi_l.$$

Combining the above two identities, we deduce, at $(x, t) = (x_0, t_0)$, that

(4.2)
$$(\partial_t - \Delta)\Lambda = \frac{1}{t}\Lambda + \frac{2}{t}\Lambda^2 + 2R_{ikjl}Q_{kl}\xi_i\xi_j + 2tR_{ikjl}\nabla_k v\nabla_l v\xi_i\xi_j + 2\nabla_k\Lambda\nabla_k v.$$

LI AND ZHANG

Notice the terms involving the Ricci curvature are canceled. We remark that this equation may not be satisfied for $t < t_0$ but the proof uses this equation only at $(x,t) = (x_0, t_0)$. As mentioned, Λ reaches its minimum value at (x_0, t_0) . Therefore, (4.2) implies, at (x_0, t_0)

(4.3)
$$2\lambda_1^2 \le -\lambda_1 - 2tR_{ikjl}Q_{kl}\xi_i\xi_j - 2t^2R_{ikjl}\nabla_k v\nabla_l v\xi_i\xi_j.$$

Next, we aim to bound the curvature terms on the right-hand side of (4.3). First, we write

$$\begin{aligned} 2R_{ikjl}Q_{kl} &= 2(R_{ikjl} + K(g_{ij}g_{kl} - g_{il}g_{jk}))Q_{kl} - 2K(g_{ij}g_{kl} - g_{il}g_{jk})Q_{kl} \\ &:= 2w_{ikjl}Q_{kl} - 2K(g_{ij}g_{kl} - g_{il}g_{jk})Q_{kl}. \end{aligned}$$

Besides the lowest negative eigenvalue λ_1 , let λ_2 , ..., λ_n be other eigenvalues of (Q_{ij}) at (x_0, t_0) arranged in increasing order. After diagonalizing (Q_{ij}) at (x_0, t_0) with an orthonormal basis $\{\xi, ...\}$, we deduce

$$R_{ikjl}Q_{kl}\xi_i\xi_j = \sum_k w_{1k1k}\lambda_k - K(g_{ij}g_{kl} - g_{il}g_{jk})Q_{kl}\xi_i\xi_j$$

$$= \sum_{\lambda_k \ge 0} w_{1k1k}\lambda_k + \sum_{\lambda_k < 0} w_{1k1k}\lambda_k - Kt\Delta v + K\lambda_1$$

$$\ge \sum_{\lambda_k < 0} w_{1k1k}\lambda_1 - Kt\Delta v + K\lambda_1.$$

Here we just used the assumption on sectional curvature

$$R_{ikjl} \ge -K(g_{ij}g_{kl} - g_{il}g_{jk})$$

or $w_{ikjl} \geq 0$ and the identity

$$\operatorname{tr}(Q_{ij}) = t\Delta v.$$

Using the upper bound of the sectional curvature

$$R_{ikjl} \le K(g_{ij}g_{kl} - g_{il}g_{jk}),$$

we see that

$$\sum_{k} w_{1k1k} \le 2K \sum_{k} (g_{11}g_{kk} - g_{1k}g_{1k}) = 2K(n-1)$$

and we arrive at, via $\lambda_1 < 0$, that

(4.4)
$$R_{ikjl}Q_{kl}\xi_i\xi_j \ge (2n-1)K\lambda_1 - Kt\Delta v.$$

Using the lower bound on the sectional curvatures again, noticing $\xi = (1, 0, ..., 0)$ and $g_{ij} = \delta_{ij}$ at (x_0, t_0) by our choice of the orthonormal coordinates, we have that

$$(4.5) R_{ikjl}\nabla_k v\nabla_l v\xi_i\xi_j = R_{1k1l}\nabla_k v\nabla_l v$$

$$\geq -K(g_{kl} - g_{1l}g_{1k})\nabla_k v\nabla_l v$$

$$= -K|\nabla v|^2 + K|\nabla_1 v|^2$$

$$\geq -K|\nabla v|^2.$$

Substituting (4.4), (4.5) into (4.3), we deduce, for $v = \log u$,

(4.6)
$$2\lambda_1^2 \le -\lambda_1 - 2t(2n-1)K\lambda_1 + 2Kt^2\Delta v + 2t^2K|\nabla v|^2.$$

Step 2. We need to bound the right hand side of (4.6). This might be difficult for all positive solutions u but doable for the heat kernel.

In this step, we assume u(x,t) = G(x,t,y) := G(x,t;y,0) is the heat kernel with a pole at $y \in M, t = 0$ and $v = \log u$. It is known that the following curvature-free bound holds:

$$(4.7) \ (t - (t_0/2))^2 |\nabla \log u|^2 \le (t - (t_0/2)) \log \frac{\sup\{u(x,t) : (x,t) \in M \times [t_0/2, t_0]\}}{u}$$

for all $(x,t) \in M \times [t_0/2, t_0]$ (see [Zha06] or [CH09]). Under the condition of bounded sectional curvature, the upper and lower bound for the heat kernel can be obtained in a classical way in any finite time interval and hence are more or less known. By now, we know that only the pointwise bound on the scalar curvature and initial volume non-collapsing condition are needed for the heat kernel bounds to hold (see [BZ17, Theorem 1.4]). Using that theorem repeatedly over fixed time intervals and taking advantage of the reproducing formula of the heat kernel, we know that the following bounds hold: there exists a numerical positive constant C_1 and another positive constant C_2 depending only on the volume non-collapsing constant of g_0 and the dimension such that

(4.8)
$$\frac{1}{C_2 t^{n/2}} e^{-C_2 K t - C_1 d^2(x, y, t)/t} \le G(x, t, y) \le \frac{C_2}{t^{n/2}} e^{C_2 K t - d^2(x, y, t)/(C_1 t)}.$$

Alternatively, since the sectional curvature is bounded, one can just follow the classical method by Li-Yau to obtain such bounds. The above bounds are far from optimal for large times. Since the manifold is compact, the large-time behavior of the heat kernel is relatively simple since positive solutions tend to be constant. But we will not pursue an optimal large-time bound this time.

Substituting (4.8) to (4.7), we obtain, for all $t_0 > 0$

(4.9)
$$t_0^2 |\nabla \log u|^2(x, t_0) \le C_3(K+1)t_0^2 + 2C_1 \sup_{t \in [t_0/2, t_0]} d^2(x, t, y).$$

Here C_3 depends only on the volume noncollapsing constant of g_0 and the dimension.

Next, we need to find an upper bound for the term $t^2 \Delta \log u = t^2 (\frac{\Delta u}{u} - \frac{|\nabla u|^2}{u^2})$. In the stationary case, this is done in Hamilton [Ham93b, Lemma 4.1]. Following that proof, the Ricci flow produces one extra term involving the Ricci curvature. Since the sectional curvature is bounded, we can treat this term without much difficulty.

Let L be the operator

(4.10)
$$L = \Delta + 2\nabla \log u\nabla - \partial_t.$$

The following identities are well known and also follow from the calculations in Section 2.

(4.11)
$$L(\Delta u/u) = 2R_{ij}\nabla_i\nabla_j u/u, \quad L(|\nabla \log u|^2) = 2|\nabla_i\nabla_j \log u|^2.$$

Therefore,

$$L\left(\frac{\Delta u}{u} + |\nabla \log u|^{2}\right)$$

$$= 2|\nabla_{i}\nabla_{j}\log u|^{2} - 2R_{ij}\left(\frac{\nabla_{i}\nabla_{j}u}{u} - \frac{\nabla_{i}u\nabla_{j}u}{u^{2}}\right) - 2R_{ij}\frac{\nabla_{i}u\nabla_{j}u}{u^{2}}$$

$$= 2|\nabla_{i}\nabla_{j}\log u|^{2} - 2R_{ij}\nabla_{i}\nabla_{j}\log u - 2R_{ij}\frac{\nabla_{i}u\nabla_{j}u}{u^{2}}$$

$$= |\nabla_{i}\nabla_{j}\log u|^{2} + |\nabla_{i}\nabla_{j}\log u - R_{ij}|^{2} - R_{ij}^{2} - 2R_{ij}\frac{\nabla_{i}u\nabla_{j}u}{u^{2}}$$

$$\geq \frac{1}{n}|\Delta \log u|^{2} + \frac{1}{n}|\Delta \log u - R|^{2} - C_{n}K^{2} - C_{n}K|\nabla \log u|^{2}.$$

Here, C_n is a dimensional constant and the assumption $|R_{ijkl}| \le K(g_{ik}g_{jl} - g_{il}g_{jk})$ has been used. Writing

$$Y = \frac{\Delta u}{u} + |\nabla \log u|^2,$$

then the above inequality implies

$$LY \geq \frac{1}{n} |Y - 2|\nabla \log u|^2|^2 + \frac{1}{n} |Y - 2|\nabla \log u|^2 - R|^2 - C_n K^2 - C_n K |\nabla \log u|^2.$$

Therefore,

(4.12)
$$L(t^{2}Y) \geq \frac{1}{nt^{2}} |t^{2}Y - 2t^{2}|\nabla \log u|^{2}|^{2} + \frac{1}{nt^{2}} |t^{2}Y - 2t^{2}|\nabla \log u|^{2} - t^{2}R|^{2} - C_{n}t^{2}K^{2} - C_{n}Kt^{2}|\nabla \log u|^{2} - 2\frac{t^{2}Y}{t}$$

Let T > 0 be any fixed time and (x_0, t_0) be a maximal point of t^2Y in $M^n \times (0, T]$ where t^2Y reaches a positive maximum value. Note that t_0 may be less than T but the argument by maximum principle together with (4.9) is good enough to bound t^2Y up to T and hence for all time. By (4.12), we know, at (x_0, t_0) , the following inequality holds:

$$\frac{1}{nt^2} \left| t^2 Y - 2t^2 |\nabla \log u|^2 \right|^2 \le C_n t^2 K^2 + C_n K t^2 |\nabla \log u|^2 + 2\frac{t^2 Y}{t}.$$

Hence for all $t \in (0, T]$, we have

$$\frac{1}{2nt^2} |t^2 Y|^2 \le \left(\frac{1}{nt^2} |2t^2|\nabla \log u|^2|^2 + C_n t^2 K^2 + C_n K t^2 |\nabla \log u|^2 + 2\frac{t^2 Y}{t}\right)_{(x_0, t_0)}$$

Now we take u to be the heat kernel G(x, t, 0). Using (4.9) we conclude that

$$t^2 Y \le 4nt + C_3(K+1)t^2 + 2C_1 \operatorname{diam}^2$$

which yields, since $Y = \frac{\Delta u}{u} + |\nabla \log u|^2$, that

(4.13)
$$t^2 \frac{\Delta u}{u} \le 4nt + C_3(K+1)t^2 + 2C_1 \operatorname{diam}^2.$$

From (4.6) and the relation $\Delta v = \Delta \log u = \frac{\Delta u}{u} - |\nabla \log u|^2$, we see that

(4.14)
$$2\lambda_1^2 \le -\lambda_1 - 2t(2n-1)K\lambda_1 + 2Kt^2\frac{\Delta u}{u}.$$

Using (4.13), we deduce that

(4.15)
$$2\lambda_1^2 \leq -\lambda_1 - 2t(2n-1)K\lambda_1 \\ + 2K[4nt + C_3(K+1)t^2 + 2C_1 \operatorname{diam}^2].$$

Since $\lambda_1 < 0$ by assumption, we conclude, after elementary estimates,

$$(4.16) \quad t(H_{ij}) \geq \lambda_1(g_{ij})$$

$$\geq \left(-\frac{1}{2} - 4\sqrt{nKt} - C_2(K+1)t - C_1\sqrt{K}\operatorname{diam}\right)(g_{ij})$$

$$:= \left(-\frac{1}{2} - \beta(t, n, K)\right)(g_{ij}),$$

where C_1 is a numerical constant and C_2 depends only on the non-collapsing constant v_0 of g_0 and the dimension. Note that we have renamed C_3 to C_2 for consistency with the statement of the theorem.

Step 3. Finally, we show the matrix Harnack estimate holds for any positive solution u(x,t) to the heat equation.

Note that

$$u(x,t) = \int_M G(x,t,y) u_0(y) dy.$$

Differentiating under the integral yields

$$\begin{aligned} \nabla_{j} u(x,t) &= \int_{M} \nabla_{j} G(x,t,y) u_{0}(y) dy, \\ \nabla_{i} \nabla_{j} u(x,t) &= \int_{M} \nabla_{i} \nabla_{j} G(x,t,y) u_{0}(y) dy. \end{aligned}$$

Here and later in the step dy = dg(0)(y) etc. Therefore,

$$\begin{aligned} tu^2 \nabla_i \nabla_j \log u(x,t) \\ &= t(u \nabla_i \nabla_j u - \nabla_i u \nabla_j u)(x,t) \\ &= \int_M tG(x,t,z) u_0(z) dz \int_M \nabla_i \nabla_j G(x,t,y) u_0(y) dy \\ &- \int_M t \nabla_j G(x,t,z) u_0(z) dz \int_M \nabla_j G(x,t,y) u_0(y) dy. \end{aligned}$$

Hence,

$$(4.17) tu^2 \nabla_i \nabla_j \log u(x,t) \\ = \int_M \int_M tG(x,t,z) \nabla_i \nabla_j G(x,t,y) u_0(z) u_0(y) dz dy \\ - \int_M \int_M t \nabla_j G(x,t,z) \nabla_j G(x,t,y) u_0(z) u_0(y) dz dy.$$

Fixing a space time point (x, t), t > 0. Let us diagonalize $(Q_{ij}) = t(\nabla_i \nabla_j \log u(x, t))$ using its orthonormal eigenvectors $\{e_1, ..., e_n\}$ such that e_1 corresponds to the smallest eigenvalue λ_1 . By (4.17), we have

(4.18)
$$tu^{2}\nabla_{1}\nabla_{1}\log u(x,t)$$
$$= \int_{M}\int_{M}tG(x,t,z)\nabla_{1}\nabla_{1}G(x,t,y)u_{0}(z)u_{0}(y)dzdy$$
$$-\int_{M}\int_{M}t\nabla_{1}G(x,t,z)\nabla_{1}G(x,t,y)u_{0}(z)u_{0}(y)dzdy.$$

According to (4.16) in Step 2, the following holds

$$t\left(\frac{\nabla_1 \nabla_1 G(x,t,y)}{G(x,t,y)} - \frac{|\nabla_1 G(x,t,y)|^2}{G^2(x,t,y)}\right) \ge -\frac{1}{2} - \beta(t,n,K)$$

so that

$$\begin{split} t\nabla_1\nabla_1G(x,t,y) &\geq t\frac{|\nabla_1G(x,t,y)|^2}{G(x,t,y)} \\ &\quad -\left(\frac{1}{2}+\beta(t,n,K)\right)G(x,t,y). \end{split}$$

Substituting the last inequality into (4.18) and regrouping the third term on the right-hand side, we deduce

$$(4.19) tu^{2} \nabla_{1} \nabla_{1} \log u(x,t) \\ \geq \int_{M} \int_{M} G(x,t,z) t \frac{|\nabla_{1}G(x,t,y)|^{2}}{G(x,t,y)} u_{0}(z) u_{0}(y) dz dy \\ -[\frac{1}{2} + \beta(t,n,K)] \int_{M} \int_{M} G(x,t,z) G(x,t,y) u_{0}(z) u_{0}(y) dz dy \\ - \int_{M} \int_{M} \frac{\sqrt{t} \nabla_{1}G(x,t,z) \sqrt{G(x,t,y)}}{\sqrt{G(x,t,z)}} \sqrt{u_{0}(z) u_{0}(y)} \\ \times \frac{\sqrt{t} \nabla_{1}G(x,t,y) \sqrt{G(x,t,z)}}{\sqrt{G(x,t,y)}} \sqrt{u_{0}(z) u_{0}(y)} dz dy.$$

Observe that the first term on the right-hand side dominates the third term due to the Cauchy-Schwarz inequality and the integral in the second term is $u^2(x,t)$. Hence we have proven

$$t\nabla_1 \nabla_1 \log u(x,t) \ge -\frac{1}{2} - \beta(t,n,K)$$

Since the left-hand side is the smallest eigenvalue of (Q_{ij}) , the proof is done. \Box

5. PARABOLIC FREQUENCY MONOTONICITY

5.1. **Parabolic Frequency.** Let $(M^n, g(t)), t \in [0, T]$, be a complete solution to the Ricci flow. Let u be a solution to the (backward) conjugate heat equation $(\partial_t + \Delta)u = Ru$. Let $G(x, x_0, t)$ be the heat kernel of the heat equation (1.8) with

the pole at $(x_0, 0)$. We defined in the Introduction section that

(5.1)
$$I(t) := \int_{M} u^{2}(x,t)G(x,x_{0},t)dg(t)(x),$$

(5.2)
$$D(t) := \int_{M} |\nabla u(x,t)|^2 G(x,x_0,t) dg(t)(x),$$

(5.3)
$$S(t) := \int_{M} u^{2}(x,t)R(x,t)G(x,x_{0},t)dg(t)(x).$$

Here $dg(t)(x) := d\mu_{g(t)}$ is the Riemannian measure induced by g(t).

Below we will simply write $I(t)=\int_M u^2 G dg$ and similar notations for other integrals if no confusion arises.

Lemma 5.1. I(t) defined in (5.1) satisfies

(5.4)
$$I'(t) = 2D(t) + S(t).$$

Proof. Under the Ricci flow, the measure dg(t) evolves by $\partial_t(dg(t)) = -Rdg(t)$. A straightforward computation shows that

$$\begin{split} I'(t) &= \int_{M} 2uu_{t}Gdg + \int_{M} u^{2}G_{t}dg - \int_{M} u^{2}RGdg \\ &= \int_{M} 2u(Ru - \Delta u)Gdg + \int_{M} u^{2}\Delta Gdg - \int_{M} u^{2}RGdg \\ &= \int_{M} u^{2}RGdg - \int_{M} (\Delta u^{2} - 2|\nabla u|^{2})Gdg + \int_{M} \Delta u^{2}Gdg \\ &= 2\int_{M} |\nabla u|^{2}Gdg + \int_{M} u^{2}RGdg \\ &= 2D(t) + S(t). \end{split}$$

Lemma 5.2. D(t) defined in (5.2) satisfies

(5.5)
$$D'(t) = 2 \int_{M} (\operatorname{Ric} -\nabla^{2} f) (\nabla u, \nabla u) G dg + 2 \int_{M} (\Delta_{f} u)^{2} G dg$$
$$-2 \int_{M} Ru(\Delta_{f} u) G dg - \int_{M} |\nabla u|^{2} RG dg,$$

where $f = -\log G$ and $\Delta_f := \Delta - \langle \nabla f, \nabla \cdot \rangle$.

Proof. Noticing $\partial_t |\nabla u|^2 = 2 \operatorname{Ric}(\nabla u, \nabla u) + 2 \langle \nabla u, \nabla u_t \rangle$ and $\partial_t dg(t) = -Rdg(t)$, we get by differentiating under the integral that

$$(5.6) \qquad D'(t) = 2 \int_{M} \operatorname{Ric}(\nabla u, \nabla u) G dg + 2 \int_{M} \langle \nabla u, \nabla u_{t} \rangle G dg \\ + \int_{M} |\nabla u|^{2} G_{t} dg - \int_{M} |\nabla u|^{2} R G dg \\ = 2 \int_{M} \operatorname{Ric}(\nabla u, \nabla u) G dg + 2 \int_{M} \langle \nabla u, \nabla (Ru - \Delta u) \rangle G dg \\ + \int_{M} |\nabla u|^{2} \Delta G dg - \int_{M} |\nabla u|^{2} R G dg \\ = 2 \int_{M} \operatorname{Ric}(\nabla u, \nabla u) G dg + 2 \int_{M} \langle \nabla u, \nabla (Ru) \rangle G dg \\ - 2 \int_{M} \langle \nabla u, \nabla (\Delta u) \rangle G dg + \int_{M} \Delta |\nabla u|^{2} G dg \\ - \int_{M} |\nabla u|^{2} R G dg.$$

In view of the integration by parts

$$\int_{M} \langle \nabla u, \nabla (Ru) \rangle G dg = -\int_{M} Ru \Delta_{f} u G dg$$

and the Bochner formula

$$\Delta |\nabla u|^2 = 2|\nabla^2 u|^2 + 2\langle \nabla u, \nabla \Delta u \rangle + 2\operatorname{Ric}(\nabla u, \nabla u),$$

we derive from (5.6) that

(5.7)
$$D'(t) = 4 \int_{M} \operatorname{Ric}(\nabla u, \nabla u) G dg + 2 \int_{M} |\nabla^{2} u|^{2} G dg$$
$$+ 2 \int_{M} R u \Delta_{f} u G dg - \int_{M} |\nabla u|^{2} R G dg,$$

The weighted Bochner formula

$$\Delta_f |\nabla u|^2 = 2|\nabla^2 u|^2 + 2\langle \nabla u, \nabla \Delta_f u \rangle + 2\operatorname{Ric}(\nabla u, \nabla u) + 2\nabla^2 f(\nabla u, \nabla u),$$

implies that

$$2\int_{M} \operatorname{Ric}(\nabla u, \nabla u)G + 2\int_{M} |\nabla^{2}u|^{2}Gdg$$

= $-2\int_{M} \langle \nabla u, \nabla \Delta_{f}u \rangle Gdg - 2\int_{M} \nabla^{2}f(\nabla u, \nabla u)dg$
= $2\int_{M} (\Delta_{f}u)^{2}Gdg - 2\int_{M} \nabla^{2}f(\nabla u, \nabla u)dg.$

Substituting the above identity into (5.7) produces

$$D'(t) = 2 \int_{M} (\operatorname{Ric} -\nabla^{2} f) (\nabla u, \nabla u) G dg + 2 \int_{M} (\Delta_{f} u)^{2} G dg$$
$$-2 \int_{M} Ru(\Delta_{f} u) G dg - \int_{M} |\nabla u|^{2} RG dg.$$

This proves (5.5).

Lemma 5.3. S(t) defined in (5.3) satisfies

(5.8)
$$S'(t) = \int_{M} u^{2} R^{2} G dg + 2 \int_{M} u^{2} |\operatorname{Ric}|^{2} G dg + 2 \int_{M} |\nabla u|^{2} R G dg$$
$$+ 2 \int_{M} u^{2} R \Delta G dg + 4 \int_{M} u R \langle \nabla u, \nabla G \rangle dg.$$

Proof. We compute using $\partial_t R = \Delta R + 2|\operatorname{Ric}|^2$ that

$$\begin{split} S'(t) &= \int_{M} 2uu_{t}RGdg + \int u^{2}R_{t}Gdg \\ &+ \int_{M} u^{2}RG_{t}dg - \int u^{2}R^{2}Gdg \\ &= \int_{M} 2u(Ru - \Delta u)RGdg + \int u^{2}(\Delta R + 2|\operatorname{Ric}|^{2})Gdg \\ &+ \int_{M} u^{2}R\Delta Gdg - \int u^{2}R^{2}Gdg \\ &= \int_{M} u^{2}R^{2}Gdg - 2\int_{M} u\Delta uRGdg + \int u^{2}\Delta RGdg \\ &+ 2\int_{M} u^{2}|\operatorname{Ric}|^{2}Gdg + \int_{M} u^{2}R\Delta Gdg. \end{split}$$

Observing

$$\begin{split} \int_{M} u^{2} \Delta RG dg &= \int_{M} R\Delta(u^{2}G) dg \\ &= \int_{M} R\left(\Delta u^{2}G + u^{2}\Delta G + 2\langle \nabla u^{2}, \nabla G \rangle\right) dg \\ &= 2\int_{M} u\Delta uRG dg + 2\int_{M} |\nabla u|^{2}RG dg \\ &+ \int_{M} u^{2}R\Delta G dg + 4\int_{M} uR\langle \nabla u, \nabla G \rangle dg, \end{split}$$

we deduce that

$$S'(t) = \int_{M} u^{2} R^{2} G dg + 2 \int_{M} u^{2} |\operatorname{Ric}|^{2} G + 2 \int_{M} |\nabla u|^{2} R G dg$$
$$+ 2 \int_{M} u^{2} R \Delta G dg + 4 \int_{M} u R \langle \nabla u, \nabla G \rangle dg.$$

Lemma 5.4. For I(t) defined in (5.1), we have

(5.9)
$$I''(t) = 4 \int_{M} (\operatorname{Ric} -\nabla^{2} f) (\nabla u, \nabla u) G dg + \int_{M} (2\Delta_{f} u - Ru)^{2} G dg + 2 \int_{M} u^{2} |\operatorname{Ric}|^{2} G dg + 2 \int_{M} u^{2} R \Delta G dg + 4 \int_{M} Ru \langle \nabla u, \nabla G \rangle dg.$$

Proof. By (5.4), we have I'(t) = 2D(t) + S(t). Using (5.5) and (5.8), we calculate

$$\begin{split} I''(t) &= 2D'(t) + S'(t) \\ &= 4\int_{M}(\operatorname{Ric} -\nabla^{2}f)(\nabla u, \nabla u)Gdg + 4\int_{M}(\Delta_{f}u)^{2}Gdg \\ &- 4\int_{M}Ru(\Delta_{f}u)Gdg - 2\int_{M}|\nabla u|^{2}RGdg \\ &+ \int_{M}u^{2}R^{2}Gdg + 2\int_{M}u^{2}|\operatorname{Ric}|^{2}G + 2\int_{M}|\nabla u|^{2}RGdg \\ &+ 2\int_{M}u^{2}R\Delta Gdg + 4\int_{M}Ru\langle\nabla u, \nabla G\rangle dg \\ &= 4\int_{M}(\operatorname{Ric} -\nabla^{2}f)(\nabla u, \nabla u)Gdg + \int_{M}(2\Delta_{f}u - Ru)^{2}Gdg \\ &+ 2\int_{M}u^{2}|\operatorname{Ric}|^{2}Gdg + 2\int_{M}u^{2}R\Delta Gdg + 4\int_{M}Ru\langle\nabla u, \nabla G\rangle dg. \end{split}$$

5.2. The nonnegative sectional curvature case. We prove Theorem 1.7 and Corollary 1.8 in this subsection.

Proof of Theorem 1.7. We need to estimate I''(t) in (5.9) from below. Using Theorem 1.1 and Ric ≥ 0 , we get

(5.10)
$$\int_{M} (\operatorname{Ric} + \nabla^{2} \log G) (\nabla u, \nabla u) G dg \ge -\frac{\kappa}{1 - e^{-2\kappa t}} D(t).$$

By Corollary 1.2, we have

$$\Delta G \ge |\nabla G|^2 G^{-1} - \frac{n\kappa}{1 - e^{-2\kappa t}} G.$$

Since $R \ge 0$, we obtain

(5.11)
$$\int_{M} u^2 R \Delta G \ge \int_{M} u^2 R |\nabla G|^2 G^{-1} dg - \frac{n\kappa}{1 - e^{-2\kappa t}} S(t)$$

Noticing $0 \le R \le n\kappa$, we estimate that

(5.12)
$$2\int_{M} Ru \langle \nabla u, \nabla G \rangle dg$$
$$\geq -\int_{M} Ru^{2} |\nabla G|^{2} G^{-1} dg - \int_{M} R |\nabla u|^{2} G dg$$
$$\geq -\int_{M} Ru^{2} |\nabla G|^{2} G^{-1} dg - n\kappa D(t)$$

Inserting the estimates (5.10), (5.11), and (5.12) into (5.9) yields

(5.13)
$$I''(t) \ge \int_M \left(2\Delta_f u - Ru\right)^2 Gdg - \left(\frac{2\kappa}{1 - e^{-2\kappa t}} + n\kappa\right) I'(t)$$

where we have used $S(t) \ge 0$ and (5.4).

Using (5.13) and the Cauchy-Schwarz inequality,

$$(I'(t))^2 = \left(\int_M (2\Delta_f u - Ru)Gdg\right)^2$$

$$\leq \int_M u^2Gdg \cdot \int_M (2\Delta_f u - Ru)^2 Gdg$$

we obtain that for $F(t) := (\log I(t))'$,

$$F'(t) = I(t)^{-2} \left(I''(t) - (I'(t))^2 \right)$$

$$\geq -\left(\frac{2\kappa}{1 - e^{-2\kappa t}} + n\kappa \right) F(t).$$

Therefore, the quantity

$$e^{(n-2)\kappa t}(1-e^{-2\kappa t})F(t)$$

is monotone nondecreasing.

Next, we prove the unique continuation property.

Proof of Corollary 1.8. The key point to achieve unique continuation is that the correction factor in (1.19) is asymptotic to t as $t \to 0$.

Suppose a solution u = u(x,t) of the conjugate heat equation in $M \times [0,T)$ vanishes at infinity order at $(x_0,t_0) \in M \times (0,T)$. Since the quantity $e^{(n-2)\kappa t}(1-e^{-2\kappa t})F(t)$ is monotone nondecreasing on [0,T], we have

$$(\log I(t))' = F(t) \ge F(T)e^{(n-2)\kappa(T-t)}\frac{1-e^{-2\kappa T}}{1-e^{-2\kappa t}} \ge \frac{C}{t}$$

for all $t \in (t_0, T)$, where $C = F(T)(1 - e^{-2\kappa T})/(2\kappa)$. Hence,

(5.14)
$$I(t) \ge \left(\frac{t}{t_1}\right)^{\mathbb{C}} I(t_1)$$

The rest of the proof is standard since the heat kernel G has Gaussian upper bound and the distance $d(x, x_0, t)$ are comparable in short time due to our assumption. This Gaussian bound and the assumption on the infinite vanishing order of u at $(x_0, 0)$ implies, for all small t > 0,

$$I(t) = \int_{M} u^{2}(x,t)G(x,x_{0},t)dg(t)(x) \le C_{N}t^{N}$$

for any positive integer N, which is a contradiction to (5.14) unless $u \equiv 0$.

5.3. The general case. Since our assumption implies that $|Ric| \leq c_n K$, taking $\alpha = 2$ and $\rho = \infty$ in [BCP10, Theorem 2.7], we have the following Li-Yau bound

(5.15)
$$\frac{|\nabla G|^2}{G^2} - 2\frac{G_t}{G} \le c(n)\left(\frac{1}{t} + K\right),$$

where c(n) is a dimensional constant. Then

(5.16)
$$\Delta G = G_t \ge \frac{|\nabla G|^2}{2G} - \left(\frac{c(n)}{t} + K\right)G.$$

Note that this also follows from the matrix Harnack inequality in Section 4 after taking the trace.

Let $s_0 = \inf_{x \in M^n} R(x, 0)$. It is well known that $R(x, t) \ge s_0$ for all $t \in [0, T]$. Since $R - s_0 \ge 0$ and

$$(5.17) \qquad \int_{M} u^{2} R \Delta G dg + 2 \int_{M} R u \langle \nabla u, \nabla G \rangle dg$$
$$= \int_{M} u^{2} (R - s_{0}) \Delta G dg + \int_{M} u^{2} s_{0} \Delta G dg + 2 \int_{M} R u \langle \nabla u, \nabla G \rangle dg$$
$$= \int_{M} u^{2} (R - s_{0}) \Delta G dg + 2 \int_{M} (R - s_{0}) u \langle \nabla u, \nabla G \rangle dg,$$

we can apply (5.16) on the right-hand hand side of (6.9) to reach

$$\int_{M} u^{2} R \Delta G dg + 2 \int_{M} R u \langle \nabla u, \nabla G \rangle dg$$

$$\geq \int_{M} u^{2} (R - s_{0}) \left[\frac{|\nabla G|^{2}}{2G} - \left(\frac{c(n)}{t} + K \right) G \right] dg$$

$$+ 2 \int_{M} (R - s_{0}) u \langle \nabla u, \nabla G \rangle dg.$$

Using Cauchy-Schwarz inequality, we have

$$\begin{split} & 2\int_M (R-s_0)u\langle \nabla u, \nabla G\rangle dg \\ \geq & -\frac{1}{2}\int (R-s_0)u^2 |\nabla \log G|^2 G dg - 2\int_M (R-s_0)|\nabla u|^2 G dg. \end{split}$$

Hence, we obtain the estimate

$$\begin{split} &\int_{M} Ru^{2} \Delta G dg 2 + \int_{M} Ru \langle \nabla u, \nabla G \rangle dg \\ \geq & -\left(\frac{c(n)}{t} + K\right) \int_{M} (R-s_{0}) u^{2} G dg \\ & -2 \int_{M} (R-s_{0}) |\nabla u|^{2} G dg. \end{split}$$

Using this inequality, the matrix Harnack inequality in Theorem 1.3,

(5.18)
$$-\nabla^2 f \ge -c_1(t)g_{ij}, \quad c_1(t) := \frac{1}{2t} + \frac{1}{t}\beta(t, n, K)$$

we deduce, from Lemma 5.4, that

$$\begin{split} I''(t) &\geq \int_{M} (2\Delta_{f}u - Ru)^{2} \, Gdg + 2 \int u^{2} |\operatorname{Ric}|^{2} Gdg \\ &- 4(c_{1}(t) + c(n)K) \int_{M} |\nabla u|^{2} Gdg - 4 \int_{M} (R - s_{0}) |\nabla u|^{2} Gdg \\ &- 2 \left(\frac{c(n)}{t} + K \right) \int_{M} (R - s_{0}) u^{2} Gdg. \end{split}$$

Here we have used $\operatorname{Ric}(\nabla u, \nabla u) \geq -c(n)K|\nabla u|^2$. Using the fact that $R - s_0 \leq c(n)K$ and adjusting the dimensional constant c(n), we deduce

$$I''(t) \geq \int_{M} (2\Delta_{f}u - Ru)^{2} Gdg + 2 \int u^{2} |\operatorname{Ric}|^{2} Gdg$$
$$-2(c_{1}(t) + c(n)K) \left[\int_{M} 2|\nabla u|^{2} Gdg + \int_{M} Ru^{2} Gdg \right]$$
$$+(c_{1}(t) + c(n)K) \int_{M} Ru^{2} Gdg$$
$$-2 \left(\frac{c(n)}{t} + K \right) \int_{M} (R - s_{0})u^{2} Gdg.$$

Therefore

$$\begin{split} I''(t) &\geq \int_M \left(2\Delta_f u - Ru\right)^2 Gdg + 2\int u^2 |\operatorname{Ric}|^2 Gdg \\ &- 2(c_1(t) + c(n)K) \left[\int_M 2|\nabla u|^2 Gdg + \int_M Ru^2 Gdg\right] \\ &- c_2(t) \int_M u^2 Gdg, \end{split}$$

where

(5.19)
$$c_2(t) = c(n)K\left[2(c_1(t) + c(n)K) + 2\left(\frac{c(n)}{t} + K\right)\right].$$

This implies that

$$I^{2}(t)(\log I(t))''$$

$$= I''(t)I(t) - (I'(t))^{2}$$

$$\geq -2(c_{1}(t) + c(n)K)I(t)\left(2\int_{M}|\nabla u|^{2}Gdg + \int_{M}u^{2}Gdg\right)$$

$$-c_{2}(t)I^{2}(t) + I(t)\int_{M}(2\Delta_{f}u - Ru)^{2}Gdg - (2D(t) + S(t))^{2}.$$

Using integration by parts, we see that

$$2D(t) + S(t) = \int_M (-2\Delta_f u + Ru) u G dg.$$

Therefore the difference of the last two terms in the preceding inequality is nonnegative by Cauchy-Schwarz inequality, giving us:

$$(5.20) I^2(t)(\log I(t))'' \ge -2(c_1(t) + c(n)K)I(t)\left[2D(t) + S(t)\right] - c_2(t)I^2(t).$$

Hence

(5.21)
$$(\log I(t))'' \ge -2(c_1(t) + c(n)K)\frac{2D(t) + S(t)}{I(t)} - c_2(t).$$

Let us recall from (5.18) and Theorem 1.3,

(5.22)
$$c_1(t) := \frac{1}{2t} + \frac{1}{t}\beta(t, n, K)$$

= $\frac{1}{2t} + \frac{1}{t} \left[4\sqrt{nKt} + C_2(K+1)t + C_1\sqrt{Kt} \text{ diam} \right]$

and from (5.19)

$$c_2(t) = c(n)K\left[2(c_1(t) + c(n)K) + 2\left(\frac{c(n)}{t} + K\right)\right].$$

Let

$$Z_0 = \sup_{t \in (0,T]} [tc_2(t)].$$

Inequality (5.21) infers for $F(t) := (\log I(t))' = \frac{2D(t) + S(t)}{I(t)}$,

(5.23)
$$F'(t) \ge -2(c_1(t) + c(n)K)F(t) - \frac{Z_0}{t}.$$

Let

(5.24)
$$p = p(n, K, v_0, T, diam) = \sup_{t \in (0,T]} [t2(c_1(t) + c(n)K)]$$

From the preceding inequality, we see that

$$tF'(t) \ge -pF(t) - Z_0.$$

Therefore

$$\left(t^p F(t) + \frac{Z_0}{p} t^p\right)' \ge 0$$

Thus, we have proved Theorem 1.9.

5.4. The unweighted case. For a solution u(x,t) to the heat equation on \mathbb{R}^n , the monotonicity of the unweighted frequency

$$\frac{\int_{\mathbb{R}^n} |\nabla u(x,t)|^2 dx}{\int_{\mathbb{R}^n} u^2(x,t) dx}$$

is equivalent to the log convexity of the energy $\int u^2 dx$. This is a classical result that can be used to prove the uniqueness of the backward heat equation (see for instance [Joh82]). Here we extend this result to the conjugate heat equation coupled with the Ricci flow. Compared with the weighted case in this section, the curvature assumption is Ric ≥ 0 and no upper bound of any curvature is needed. The unweighted monotonicity, however, is not strong enough to prove the unique continuation property.

Theorem 5.1. Let $(M^n, g(t))$, $t \in [0, T]$, be a compact Ricci flow. Let u be a solution to the backward conjugate heat equation (1.7). Define

$$I(t) = \int u^2(x,t) d\mu_{g(t)}.$$

If $(M^n, g(t))$ has nonnegative Ricci curvature, then

$$(\log I(t))'' \ge 0.$$

Proof. As before, we also define

$$D(t) = \int |\nabla u(x,t)|^2 d\mu_{g(t)},$$

$$S(t) = \int u^2(x,t) R(x,t) d\mu_{g(t)},$$

and write $I(t) = \int_M u^2 dg$ for short and similar notations for other integrals.

By direct computation as for the weighted case, we have

$$I'(t) = 2 \int |\nabla u|^2 dg + \int u^2 R dg = 2D(t) + S(t),$$

$$D'(t) = 2 \int \langle \nabla u, \nabla u_t \rangle dg + 2 \int \operatorname{Ric}(\nabla u, \nabla u) dg - \int |\nabla u|^2 R dg$$
$$= -2 \int u_t \Delta u dg + 2 \int \operatorname{Ric}(\nabla u, \nabla u) dg - \int |\nabla u|^2 R dg,$$

and

$$S'(t) = 2 \int u u_t R dg + \int u^2 (\Delta R + 2|\operatorname{Ric}|^2) dg - \int u^2 R^2 dg$$

= 2 \int u u_t R dg + \int R (2u \Delta u + 2|\nabla u|^2)
+2 \int u^2 |\operatorname{Ric}|^2 dg - \int u^2 R^2 dg.

Using $\Delta u = Ru - u_t$, we deduce

$$\begin{split} I''(t) &= 2D'(t) + S'(t) \\ &= -4\int u_t \Delta u dg + 4\int \operatorname{Ric}(\nabla u, \nabla u) dg \\ &+ 2\int Ru \Delta u dg + 2\int u u_t R + 2\int u^2 |\operatorname{Ric}|^2 dg - \int u^2 R^2 dg \\ &= 4\int u_t^2 dg - 4\int u u_t R dg + \int u^2 R^2 dg \\ &+ 4\int \operatorname{Ric}(\nabla u, \nabla u) dg + 2\int u^2 |\operatorname{Ric}|^2 dg \\ &= \int (2u_t - uR)^2 dg + 4\int \operatorname{Ric}(\nabla u, \nabla u) dg + 2\int u^2 |\operatorname{Ric}|^2 dg. \end{split}$$

Using $I'(t) = \int u(2u_t - uR)$, we then get

$$I^{2}(\log I)''$$

$$= \int u^{2} dg \cdot \int (2u_{t} - uR)^{2} dg - \left(\int u(2u_{t} - uR) dg\right)^{2}$$

$$+ \int u^{2} dg \left(4 \int \operatorname{Ric}(\nabla u, \nabla u) dg + 2 \int u^{2} |\operatorname{Ric}|^{2} dg\right)$$

The first line on the right-hand side of the above equation is nonnegative by the Cauchy-Schwarz inequality and the second line is nonnegative since $\text{Ric} \ge 0$. Therefore, we have proved the log convexity of the energy I(t).

6. MATRIX HARNACK FOR THE CONJUGATE HEAT EQUATION

In this section, we prove Theorem 1.5. Let's first recall the Harnack estimate for the Ricci flow since it will be used in the proof.

Proposition 6.1. Let $(M^n, g(t))$, $t \in (0, T)$, be a complete solution to the Ricci flow with bounded nonnegative complex sectional curvature. Define

(6.1)
$$M_{ij} := \Delta R_{ij} - \frac{1}{2} \nabla_i \nabla_j R + 2R_{ikjl} R_{kl} - R_{ik} R_{jk} + \frac{1}{2t} R_{ij},$$

and

$$(6.2) P_{kij} := \nabla_k R_{ij} - \nabla_i R_{kj}.$$

Then we have

(6.3)
$$M(w,w) + 2P(v,w,w) + \operatorname{Rm}(v,w,v,w) \ge 0$$

for all $(x,t) \in M \times (0,T)$ and all vectors $v, w \in T_x M$.

Proof. The Harnack estimate for the Ricci flow was originally proved by Hamilton [Ham93a] under the nonnegative curvature operator condition. This version stated here is a generalization due to Brendle [Bre09]. Notice that M has nonnegative complex sectional curvature if and only $M \times \mathbb{R}^2$ has nonnegative isotropic curvature, which is an observation of Ni and Wolfson [NW07].

The next step is to derive an evolution inequality for

(6.4)
$$Z_{ij} := R_{ij} - \nabla_i \nabla_j \log u - \eta(t) g_{ij}.$$

Proposition 6.2. Let $(M^n, g(t))$, u, and $\eta(t)$ be the same as in Theorem 1.5. Then Z_{ij} defined in (6.4) satisfies

(6.5)
$$\frac{1}{2}(\partial_t + \Delta)Z_{ij} \ge Z_{ij}^2 - R_{ikjl}Z_{kl} - \frac{1}{2}R_{ik}Z_{jk} - \frac{1}{2}R_{jk}Z_{ik} + 2\eta Z_{ij}.$$

Proof of Proposition 6.2. For simplicity, we write $v = \log u$ and

$$H_{ij} = \nabla_i \nabla_j \log u.$$

Then the calculations in Section 2 with $\varepsilon = -1$ and $\delta = 1$ apply to this setting and we obtain from (2.2) that

$$(6.6) \qquad (\partial_t + \Delta)H_{ij} = \nabla_i \nabla_j R - (2R_{ikjl}H_{kl} - R_{ik}H_{jk} - R_{jk}H_{ik}) -2 \left(H_{ij}^2 + R_{ikjl}\nabla_k v \nabla_l v + \nabla_k H_{ij}\nabla_k v\right) +2 (\nabla_i R_{jk} + \nabla_j R_{ik} - \nabla_k R_{ij})\nabla_k v.$$

Under the Ricci flow, we have (see [CLN06, page 112])

$$(\partial_t - \Delta)R_{ij} = 2R_{ikjl}R_{kl} - 2R_{ik}R_{jk}.$$

Hence,

(6.7)
$$(\partial_t + \Delta)R_{ij} = 2\Delta R_{ij} + 2R_{ikjl}R_{kl} - 2R_{ik}R_{jk}.$$

We also notice that

(6.8)
$$(\partial_t + \Delta)(\eta(t)g_{ij}) = \eta'(t)g_{ij} - 2\eta(t)R_{ij}.$$

Combining (6.6), (6.7), and (6.8) together, we obtain that

$$\begin{split} \frac{1}{2}(\partial_t + \Delta)Z_{ij} \\ &= \frac{1}{2}(\partial_t + \Delta)R_{ij} - \frac{1}{2}(\partial_t + \Delta)H_{ij} - \frac{1}{2}(\partial_t + \Delta)(\eta(t)g_{ij}) \\ &= \Delta R_{ij} + R_{ikjl}R_{kl} - R_{ik}R_{jk} - \frac{1}{2}\nabla_i\nabla_jR + H_{ij}^2 \\ &+ R_{ikjl}H_{kl} - \frac{1}{2}R_{ik}H_{jk} - \frac{1}{2}R_{jk}H_{ik} + R_{ikjl}\nabla_kv\nabla_lv \\ &- \nabla_k(Z_{ij} - R_{ij})\nabla_kv - (\nabla_iR_{jk} + \nabla_jR_{ik} - \nabla_kR_{ij})\nabla_kv \\ &- \frac{\eta'(t)}{2}g_{ij} + \eta(t)R_{ij} \\ &= \Delta R_{ij} - \frac{1}{2}\nabla_i\nabla_jR + 2R_{ikjl}R_{kl} - R_{ik}R_{jk} + R_{ikjl}\nabla_kv\nabla_lv \\ &+ H_{ij}^2 - R_{ikjl}R_{kl} + R_{ikjl}H_{kl} - \frac{1}{2}R_{ik}H_{jk} - \frac{1}{2}R_{jk}H_{ik} \\ &- \nabla_kZ_{ij}\nabla_kv - (\nabla_iR_{jk} + \nabla_jR_{ik} - 2\nabla_kR_{ij})\nabla_kv \\ &- \frac{\eta'(t)}{2}g_{ij} + \eta(t)R_{ij}. \end{split}$$

Using (6.1), (6.2), and

$$\begin{aligned} &-(\nabla_i R_{jk} + \nabla_j R_{ik} - 2\nabla_k R_{ij})\nabla_k v \\ &= (\nabla_k R_{ij} - \nabla_i R_{jk})\nabla_k v + (\nabla_k R_{ij} - \nabla_j R_{ik})\nabla_k v \\ &= (P_{kij} + P_{kji})\nabla_k v, \end{aligned}$$

we get

(6.9)
$$\frac{1}{2}(\partial_t + \Delta)Z_{ij} = M_{ij} - \frac{1}{2t}R_{ij} + (P_{kij} + P_{kji})\nabla_k v + R_{ikjl}\nabla_k v\nabla_l v - \nabla_k Z_{ij}\nabla_k v + H_{ij}^2 - R_{ikjl}R_{kl} + R_{ikjl}H_{kl} + \frac{1}{2}R_{ik}H_{jk} + \frac{1}{2}R_{jk}H_{ik} - \frac{\eta'(t)}{2}g_{ij} + \eta(t)R_{ij}.$$

Next, we compute using (6.4) that

$$(6.10) H_{ij}^2 - R_{ikjl}R_{kl} + R_{ikjl}H_{kl} + \frac{1}{2}R_{ik}H_{jk} + \frac{1}{2}R_{jk}H_{ik}
= (R_{ik} - Z_{ik} - \eta g_{ik})(R_{jk} - Z_{jk} - \eta g_{jk}) - R_{ikjl}(Z_{kl} + \eta g_{kl})
- \frac{1}{2}R_{ik}(R_{jk} - Z_{jk} - \eta g_{jk}) - \frac{1}{2}R_{jk}(R_{ik} - Z_{ik} - \eta g_{jk})
= Z_{ij}^2 - R_{ikjl}Z_{kl} - \frac{1}{2}R_{ik}Z_{jk} - \frac{1}{2}R_{jk}Z_{ik} + 2\eta Z_{ij} + \eta^2 g_{ij} - 2\eta R_{ij}.$$

Substituting (6.10) into (6.9) produces

(6.11)
$$\frac{1}{2}(\partial_t + \Delta)Z_{ij} = M_{ij} + (P_{kij} + P_{kji})\nabla_k v + R_{ikjl}\nabla_k v\nabla_l v - \nabla_k Z_{ij}\nabla_k v + Z_{ij}^2 - R_{ikjl}Z_{kl} - \frac{1}{2}R_{ik}Z_{jk} - \frac{1}{2}R_{jk}Z_{ik} + 2cZ_{ij} + \eta^2 g_{ij} - \eta R_{ij} - \frac{\eta'(t)}{2}g_{ij} - \frac{1}{2t}R_{ij},$$

Using (1.15) and Ric $\leq \kappa g$, we have

(6.12)
$$\eta^{2} g_{ij} - \eta R_{ij} - \frac{\eta'(t)}{2} g_{ij} - \frac{1}{2t} R_{ij}$$
$$\geq \frac{1}{2t} (\kappa g_{ij} - R_{ij}) + \eta (\kappa g_{ij} - R_{ij}).$$

In view of (6.3) and (6.12), we conclude that

$$\frac{1}{2}(\partial_t + \Delta)Z_{ij} \ge Z_{ij}^2 - R_{ikjl}Z_{kl} - \frac{1}{2}R_{ik}Z_{jk} - \frac{1}{2}R_{jk}Z_{ik} + 2\eta Z_{ij}.$$

Hence, (6.5) is proved.

31

We are ready to prove Theorem 1.5.

Proof of Theorem 1.5. By Proposition 6.2,

(6.13)
$$\frac{1}{2}(\partial_t + \Delta)Z_{ij} \ge Z_{ij}^2 - R_{ikjl}Z_{kl} - \frac{1}{2}R_{ik}Z_{jk} - \frac{1}{2}R_{jk}Z_{ik} + 2\eta Z_{ij}.$$

If (M, g(t)) is compact, it follow from $\eta(t) \to \infty$ as $t \to T$ that $Z_{ij} \leq 0$ as $t \to T^-$. Noticing that each term on the right-hand side of (6.13) satisfies the nulleigenvector condition, we conclude using Hamilton's tensor maximum principle (see [Ham86] or [CLN06, Theorem 3.3]) that $Z_{ij} \leq 0$ on $M \times (0, T)$.

When (M, g(t)) is complete noncompact, one can proceed as in subsection 3.2 and use the maximum principle (see [CCG⁺08, Theorem 12.22] to prove the estimate. We omit the technical details here.

Next, we prove Corollary 1.6.

Proof of Corollary 1.6. Note that the function

$$\eta_0(t) = \frac{\kappa}{1 - e^{-2\kappa(T-t)}} + \sqrt{\frac{\kappa}{2t}}$$

satisfies $\eta'_0 \leq 2\eta_0^2 - 2\kappa\eta_0 - \frac{\kappa}{t}$ and $\eta_0(t) \to \infty$ as $t \to T$. Hence, the inequality (1.17) follows from choosing $\eta = \eta_0$ in (1.16).

Below we present an improvement of Theorem 1.5 when the Ricci flow is ancient.

Theorem 6.3. Let $(M^n, g(t))$, $t \in (-\infty, T)$, be a compact ancient solution to the Ricci flow with bounded nonnegative complex sectional curvature. Let $u : M \times [t_0, T] \to (0, \infty)$, $-\infty \leq t_0 < T$, be a positive solution to the backward conjugate heat equation $u_t + \Delta_{g(t)}u = Ru$. Suppose that $\operatorname{Ric}(x, t) \leq \kappa(x, t)g$ for some $\kappa > 0$ and for all $(x, t) \in M \times [t_0, T]$. Then we have

$$\operatorname{Ric} - \nabla^2 \log u - \frac{\kappa}{1 - e^{-2\kappa(T-t)}} g \le 0,$$

for all $(x,t) \in M \times (t_0,T)$.

Remark 6.1. In Theorem 6.3, it suffices to assume the weaker condition that $M \times \mathbb{R}$ has nonnegative isotropic curvature, in view of [BCRW19] and [LN20, Proposition 6.2].

Proof of Theorem 6.3. The proof is almost identical to that of Theorem 1.5. The difference is that we get the improved Harnack estimate (compared with (6.3))

$$M(w,w) + 2P(v,w,w) + \operatorname{Rm}(v,w,v,w) \ge \frac{1}{2t}\operatorname{Ric}(w,w)$$

on ancient Ricci flows (see [Ham93a] or [Bre09]). As a result, the ordinary differential inequality for $\eta(t)$ becomes

(6.14)
$$\eta' \le 2\eta^2 - 2\kappa\eta.$$

The theorem follows from the fact that the function

$$\eta(t) := \frac{\kappa}{1 - e^{-2\kappa(T-t)}}$$

solves (6.14) with equality on (t_0, T) and satisfies $\eta(t) \to \infty$ as $t \to T$.

We get space-time gradient estimates for $\log u$ by tracing the matrix Li-Yau-Hamilton estimates.

Corollary 6.4. Let $(M^n, g(t))$ and u be the same as in Theorem 1.5. Then

$$R - \Delta \log u - \frac{n\kappa}{1 - e^{-2\kappa(T-t)}} - n\sqrt{\frac{\kappa}{2t}} \le 0$$

for all $(x,t) \in M \times (0,T)$.

Corollary 6.5. Let $(M^n, g(t))$ and u be the same as in Theorem 6.3. Then

$$R - \Delta \log u - \frac{n\kappa}{1 - e^{-2\kappa(T-t)}} \le 0$$

for all $(x,t) \in M \times (0,T)$.

Classical-type Harnack inequalities follow from integrating the above estimates. It is an interesting question whether the above gradient estimates hold under nonnegative Ricci or sectional curvature.

As an application of Theorem 1.5, we prove Proposition 1.10.

Proof of Proposition 1.10. By (1.17), we have $\operatorname{Ric} - \nabla^2 \log u \leq \frac{k(t)}{2(T-t)}$ with

$$\frac{k(t)}{(T-t)} = \frac{2\kappa}{1 - e^{-2\kappa(T-t)}} + \sqrt{\frac{2\kappa}{t}}$$

By the work of Baldauf and Kim [BK22], the correction factor is given by

$$e^{\int \frac{1-k(t)}{T-t}dt} = \frac{1}{T-t} e^{\int \left(\frac{-2\kappa}{1-e^{-2\kappa(T-t)}} - \sqrt{2\kappa}\frac{1}{\sqrt{t}}\right)dt} = \frac{1}{T-t} e^{2\kappa(T-t)-1} e^{-\sqrt{8\kappa t}}.$$

Therefore, we have proved Proposition 1.10.

Proposition 1.10 implies a unique continuation result.

Corollary 6.6. Let $(M^n, g(t))$, $t \in [0, T]$, be a solution to the Ricci flow with nonnegative complex sectional curvature and Ric $\leq \kappa g$ for some $\kappa > 0$. Suppose that a solution u(x,t) of the heat equation (1.8) on $M \times [0,T]$ vanishes of infinity order at $(x_0, t_0) \in M \times (0,T)$. Then $u \equiv 0$ in $M \times [0,T]$.

Proof of Corollary 6.6. The key point is that the correction factor in (1.21) is asymptotic to (T-t) as $t \to T$. The proof is similar to that of Corollary 1.8 and we omit the details.

7. An improvement of Hamilton's matrix estimate

We present the proof of Theorem 1.4 in this section.

Proof of Theorem 1.4. We write $v = \log u$ and $H_{ij} = \nabla_i \nabla_j \log u$. By a straightforward calculation as in Section 2 or [Ham93b], we derive that

$$(\partial_t - \Delta)H_{ij} = 2H_{ij}^2 + 2R_{ikjl}H_{kl} - R_{ik}H_{jk} - R_{jk}H_{ik} + 2R_{ikjl}\nabla_k v \nabla_l v + 2\nabla_k H_{ij}\nabla_k v + (\nabla_l R_{ij} - \nabla_i R_{jl} - \nabla_j R_{il})\nabla_l v.$$

Compared to (3.1) in the Ricci flow case, here we have the addition term $(\nabla_l R_{ij} - \nabla_i R_{jl} - \nabla_j R_{il})\nabla_l v$. As in the proof of Theorem 1.3 in Section 4, we consider $Q_{ij} := tH_{ij}$, which satisfies

(7.1)
$$(\partial_t - \Delta)Q_{ij} = \frac{1}{t}Q_{ij} + \frac{2}{t}Q_{ij}^2 + 2R_{ikjl}Q_{kl} - R_{ik}Q_{jk} - R_{jk}Q_{ik} + 2tR_{ikjl}\nabla_k v\nabla_l v + 2\nabla_k Q_{ij}\nabla_k v + t(\nabla_l R_{ij} - \nabla_i R_{jl} - \nabla_j R_{il})\nabla_l v.$$

Let λ_1 be the smallest eigenvalue of Q_{ij} . Using $|\nabla \operatorname{Ric}| \leq L$, we estimate the last term

$$t(\nabla_{l}R_{ij} - \nabla_{i}R_{jl} - \nabla_{j}R_{il})\nabla_{l}v \ge -3Lt|\nabla v|g_{ij} \ge -3(L^{\frac{2}{3}}|\nabla v|^{2} + L^{\frac{4}{3}})g_{ij}.$$

As in the proof of Theorem 1.3, we have the estimates

$$R_{ikjl}Q_{kl}\xi_i\xi_j \geq (2n-1)K\lambda_1 - Kt\Delta v,$$

$$R_{ikjl}\nabla_k v\nabla_l v\xi_i\xi_j \geq -K|\nabla v|^2,$$

Therefore, we deduce from (7.1) that at a negative minimum point $(x_0, t_0) \in M \times [0, t_0]$ of λ_1 ,

(7.2)
$$2\lambda_1^2 \leq -\lambda_1 - 2(2n-1)Kt\lambda_1 + 2Kt^2\Delta v + 2Kt^2|\nabla v|^2 + 3L^{\frac{2}{3}}t^2|\nabla v|^2 + 3L^{\frac{4}{3}}t^2.$$

From now on, we assume u = G(x, t, y) is the heat kernel and estimate $\Delta \log u$ and $|\nabla \log u|$. First, applying Hamilton's gradient bound [Ham93b, Theorem 1.1] to $u(x, t + t_0, y)$ yields for any $t_0 > 0$ that

(7.3)
$$(t - (t_0/2)) |\nabla \log u|^2 \le (1 + 2(n-1)K(t - t_0/2)) \log \frac{A}{u}$$

where $A = \sup\{u(x,t) : (x,t) \in M \times [t_0/2, t_0]\}$. According to [Zha21], we have

(7.4)
$$\log \frac{A}{u} \leq 2\log C_1 + 4C_2Kt_0 + \frac{d^2(x,y)}{3t_0} + C_3\sqrt{K}\frac{\text{Diam}}{\sqrt{t_0}} \\ \leq C_4(1+K+Kt_0) + \frac{\text{Diam}^2}{2t_0},$$

where C_1 , C_2 , C_3 , and C_4 are dimensional constants. Substituting (7.4) into (7.3) produces

(7.5)
$$t_0 |\nabla \log u|^2 \le 2(1 + (n-1)Kt_0) \left(C_4(1 + K + Kt_0) + \frac{\text{Diam}^2}{2t_0} \right)$$

Next, we apply the Laplacian estimate (see [CCG⁺08, Theorem E.35]) to the function $u(x, t + t_0, y)$ and get

$$(t - t_0/2) \left(\Delta \log u + 2 |\nabla \log u|^2 \right) \le (1 + (n - 1)K(t - t_0/2)) \left(n + 4 \log \frac{A}{u} \right).$$

At $t = t_0$, we obtain using (7.4) that

(7.6)
$$t_0(\Delta \log u + 2|\nabla \log u|^2) \le (2 + (n-1)Kt_0) \left(n + 4C_4(1 + K + Kt_0) + 2\frac{\text{Diam}^2}{t_0}\right)$$

Since $t_0 > 0$ is arbitrary, we conclude that (7.5) and (7.6) are valid for any $t_0 = t > 0$. Combining (7.5) and (7.6), we estimate that

$$B := 2Kt\Delta v + 2Kt|\nabla v|^{2} + 3L^{\frac{2}{3}}t|\nabla v|^{2}$$

$$\leq 2K(2 + (n-1)Kt_{0})\left(n + 4C_{4}(1 + K + Kt_{0}) + 2\frac{\text{Diam}^{2}}{t}\right)$$

$$+ 3L^{\frac{2}{3}}(2(1 + (n-1)Kt_{0}))\left(C_{4}(1 + K + Kt_{0}) + \frac{\text{Diam}^{2}}{2t}\right)$$

$$\leq 2nK(2 + (n-1)Kt) + C_{5}(K + L^{\frac{2}{3}})(1 + Kt)(1 + K + Kt)$$

$$+ \left(4K(2 + (n-1)Kt) + 3L^{\frac{2}{3}}(1 + (n-1)Kt)\right)\frac{\text{Diam}^{2}}{t},$$

where C_5 depends only on the dimension. Now, (7.2) implies,

(7.7)
$$2\lambda_1^2 + \lambda_1 + 2(2n-1)Kt\lambda_1 \le tB + 3L^{\frac{4}{3}}t^2$$

Note that if $ax^2 + bx \le c$ with a > 0, b > 0, and c > 0, then we have the lower bound

$$x \ge \frac{-b - \sqrt{b^2 + 4ac}}{2a} \ge -\frac{b + \sqrt{ac}}{a}.$$

Therefore, we deduce from (7.7) that

$$\begin{aligned} \lambda_1 &\geq -\frac{1}{2} + (2n-1)Kt + \frac{1}{2}\sqrt{2(tB + 3L^{\frac{4}{3}}t^2)} \\ &\geq -\frac{1}{2} + (2n-1)Kt + \frac{1}{2}\sqrt{tB} + \frac{\sqrt{3}}{2}L^{\frac{2}{3}}t. \end{aligned}$$

The desired estimate for u = G(x, t, y) then follows by noting

$$\sqrt{tB} \ge t\gamma(t, n, K, L).$$

Finally, we can follow the same argument in Section 4 or [Zha21] to show that the desired estimate holds from any positive solution to the heat equation. This completes the proof.

Acknowledgments

The authors would like to thank Dr. Jianyu Ou for his interest and for pointing out [MO22]. The second author thanks the Department of Mathematics, Statistics and Physics at Wichita State University for its hospitality during his visit, at which occasion the initial questions for this project were discussed.

CONFLICT OF INTEREST

On behalf of all authors, the corresponding author states that there is no conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

LI AND ZHANG

References

- [Alm79] Frederick J. Almgren, Jr. Dirichlet's problem for multiple valued functions and the regularity of mass minimizing integral currents. In *Minimal submanifolds and* geodesics (Proc. Japan-United States Sem., Tokyo, 1977), pages 1–6. North-Holland, Amsterdam-New York, 1979.
- [BCP10] Mihai Bailesteanu, Xiaodong Cao, and Artem Pulemotov. Gradient estimates for the heat equation under the Ricci flow. J. Funct. Anal., 258(10):3517–3542, 2010.
- [BCRW19] Richard H. Bamler, Esther Cabezas-Rivas, and Burkhard Wilking. The Ricci flow under almost non-negative curvature conditions. *Invent. Math.*, 217(1):95–126, 2019.
- [BK22] Julius Baldauf and Dain Kim. Parabolic frequency on Ricci flows. Int. Math. Res. Not. IMRN, to appear, arXiv:2201.05505v2, 2022.
- [Bre09] Simon Brendle. A generalization of Hamilton's differential Harnack inequality for the Ricci flow. J. Differential Geom., 82(1):207–227, 2009.
- [BS09] Simon Brendle and Richard Schoen. Manifolds with 1/4-pinched curvature are space forms. J. Amer. Math. Soc., 22(1):287–307, 2009.
- [BZ17] Richard H. Bamler and Qi S. Zhang. Heat kernel and curvature bounds in Ricci flows with bounded scalar curvature. Adv. Math., 319:396–450, 2017.
- [Cao92] Huai Dong Cao. On Harnack's inequalities for the Kähler-Ricci flow. Invent. Math., 109(2):247–263, 1992.
- [Cao08] Xiaodong Cao. Differential Harnack estimates for backward heat equations with potentials under the Ricci flow. J. Funct. Anal., 255(4):1024–1038, 2008.
- [CCG⁺08] Bennett Chow, Sun-Chin Chu, David Glickenstein, Christine Guenther, James Isenberg, Tom Ivey, Dan Knopf, Peng Lu, Feng Luo, and Lei Ni. The Ricci flow: techniques and applications. Part II, volume 144 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2008. Analytic aspects.
- [CCG⁺10] Bennett Chow, Sun-Chin Chu, David Glickenstein, Christine Guenther, James Isenberg, Tom Ivey, Dan Knopf, Peng Lu, Feng Luo, and Lei Ni. The Ricci flow: techniques and applications. Part III. Geometric-analytic aspects, volume 163 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 2010.
- [CH97] Bennett Chow and Richard S. Hamilton. Constrained and linear Harnack inequalities for parabolic equations. *Invent. Math.*, 129(2):213–238, 1997.
- [CH09] Xiaodong Cao and Richard S. Hamilton. Differential Harnack estimates for timedependent heat equations with potentials. *Geom. Funct. Anal.*, 19(4):989–1000, 2009.
- [Cho98] Bennett Chow. Interpolating between Li-Yau's and Hamilton's Harnack inequalities on a surface. J. Partial Differential Equations, 11(2):137–140, 1998.
- [Cho22] Bennett Chow. Li-Yau Inequalities in Geometric Analysis dedicated to professor peter li on the occasion of his 70th birthday. J. Geom. Anal., 32(11):Paper No. 271, 10, 2022.
- [CLN06] Bennett Chow, Peng Lu, and Lei Ni. Hamilton's Ricci flow, volume 77 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI; Science Press, New York, 2006.
- [CM97a] Tobias H. Colding and William P. Minicozzi, II. Harmonic functions on manifolds. Ann. of Math. (2), 146(3):725–747, 1997.
- [CM97b] Tobias H. Colding and William P. Minicozzi, II. Harmonic functions with polynomial growth. J. Differential Geom., 46(1):1–77, 1997.
- [CM22] Tobias Holck Colding and William P. Minicozzi, II. Parabolic frequency on manifolds. Int. Math. Res. Not. IMRN, (15):11878–11890, 2022.
- [CN05] Huai-Dong Cao and Lei Ni. Matrix Li-Yau-Hamilton estimates for the heat equation on Kähler manifolds. Math. Ann., 331(4):795–807, 2005.
- [CZ11] Xiaodong Cao and Qi S. Zhang. The conjugate heat equation and ancient solutions of the Ricci flow. Adv. Math., 228(5):2891–2919, 2011.
- [Eva10] Lawrence C. Evans. Partial differential equations, volume 19 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, second edition, 2010.
- [GL86a] Nicola Garofalo and Fang-Hua Lin. Monotonicity properties of variational integrals, A_p weights and unique continuation. *Indiana Univ. Math. J.*, 35(2):245–268, 1986.

- [GL87] Nicola Garofalo and Fang-Hua Lin. Unique continuation for elliptic operators: a geometric-variational approach. Comm. Pure Appl. Math., 40(3):347-366, 1987. [Gue02]Christine M. Guenther. The fundamental solution on manifolds with time-dependent metrics. J. Geom. Anal., 12(3):425-436, 2002. [Ham82] Richard S. Hamilton. Three-manifolds with positive Ricci curvature. J. Differential Geom., 17(2):255-306, 1982. [Ham86] Richard S. Hamilton. Four-manifolds with positive curvature operator. J. Differential Geom., 24(2):153-179, 1986. [Ham93a] Richard S. Hamilton. The Harnack estimate for the Ricci flow. J. Differential Geom., 37(1):225-243, 1993. [Ham93b] Richard S. Hamilton. A matrix Harnack estimate for the heat equation. Comm. Anal. Geom., 1(1):113-126, 1993. [HL] Qing Han and Fang-Hua Lin. Nodal sets of solutions of elliptic differential equation. Book in preparation. Hong Huang. Hessian estimates for the conjugate heat equation coupled with the Ricci [Hua21] flow. arXiv:2108.04761, 2021. [HZ16] Qing Han and Qi S. Zhang. An upper bound for Hessian matrices of positive solutions of heat equations. J. Geom. Anal., 26(2):715-749, 2016. [Joh82] Fritz John. Partial differential equations, volume 1 of Applied Mathematical Sciences. Springer-Verlag, New York, fourth edition, 1982. [Kot07] Brett L. Kotschwar. Hamilton's gradient estimate for the heat kernel on complete manifolds. Proc. Amer. Math. Soc., 135(9):3013-3019, 2007. [KZ08] Shilong Kuang and Qi S. Zhang. A gradient estimate for all positive solutions of the conjugate heat equation under Ricci flow. J. Funct. Anal., 255(4):1008-1023, 2008. [Li12] Peter Li. Geometric analysis, volume 134 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2012. [Lin90] Fang-Hua Lin. A uniqueness theorem for parabolic equations. Comm. Pure Appl. Math., 43(1):127–136, 1990. [Lin91] Fang-Hua Lin. Nodal sets of solutions of elliptic and parabolic equations. Comm. Pure Appl. Math., 44(3):287-308, 1991. [Liu09] Shiping Liu. Gradient estimates for solutions of the heat equation under Ricci flow. Pacific J. Math., 243(1):165-180, 2009. [LLX22] Chuanhuan Li, Yi Li, and Kairui Xu. Parabolic frequency monotonicity on Ricci flow and Ricci-harmonic flow with bounded curvatures. J. Geom. Anal., to appear, arXiv:2205.07702, 2022. [LN20] Xiaolong Li and Lei Ni. Káhler-Ricci shrinkers and ancient solutions with nonnegative orthogonal bisectional curvature. J. Math. Pures Appl. (9), 138:28-45, 2020. [Log18a] Alexander Logunov. Nodal sets of Laplace eigenfunctions: polynomial upper estimates of the Hausdorff measure. Ann. of Math. (2), 187(1):221-239, 2018. Alexander Logunov. Nodal sets of Laplace eigenfunctions: proof of Nadirashvili's con-[Log18b] jecture and of the lower bound in Yau's conjecture. Ann. of Math. (2), 187(1):241-262, 2018.
- [LW19] Xiaolong Li and Kui Wang. Parabolic frequency monotonicity on compact manifolds. Calc. Var. Partial Differential Equations, 58(6):Paper No. 189, 18, 2019.

[LWH⁺] Zhen Lei, Zhiqiang Wang, Bobo Hua, Peng Qu, and Genggeng Huang. Lectures on mathematical and physical equations. Fudan Lecture Notes, in Chinese, unpublished.

- [LX11] Junfang Li and Xiangjin Xu. Differential Harnack inequalities on Riemannian manifolds I: linear heat equation. Adv. Math., 226(5):4456–4491, 2011.
- [LX22] Hao-Yue Liu and Pang Xu. A note on parabolic frequency and a theorem of Hardy-Pólya-Szegö. Internat. J. Math., 33(9):Paper No. 2250064, 15, 2022.
- [LY86] Peter Li and Shing-Tung Yau. On the parabolic kernel of the Schrödinger operator. Acta Math., 156(3-4):153–201, 1986.
- [Man13] Dan Mangoubi. The effect of curvature on convexity properties of harmonic functions and eigenfunctions. J. Lond. Math. Soc. (2), 87(3):645–662, 2013.
- [MMU14] Carlo Mantegazza, Giovanni Mascellani, and Gennady Uraltsev. On the distributional Hessian of the distance function. *Pacific J. Math.*, 270(1):151–166, 2014.
- [MO22] Weixiong Mai and Jianyu Ou. Liouville theorem on Ricci shrinkers with constant scalar curvature and its application. *arXiv:2208.07101*, 2022.

LI AND ZHANG

- [Mos64] Jürgen Moser. A Harnack inequality for parabolic differential equations. Comm. Pure Appl. Math., 17:101–134, 1964.
- [Ni04a] Lei Ni. Addenda to: "The entropy formula for linear heat equation" [J. Geom. Anal. 14 (2004), no. 1, 87–100; mr2030576]. J. Geom. Anal., 14(2):369–374, 2004.
- [Ni04b] Lei Ni. The entropy formula for linear heat equation. J. Geom. Anal., 14(1):87–100, 2004.
- [Ni07] Lei Ni. A matrix Li-Yau-Hamilton estimate for Kähler-Ricci flow. J. Differential Geom., 75(2):303–358, 2007.
- [Ni08] Lei Ni. Monotonicity and Li-Yau-Hamilton inequalities. In Surveys in differential geometry. Vol. XII. Geometric flows, volume 12 of Surv. Differ. Geom., pages 251–301. Int. Press, Somerville, MA, 2008.
- [Ni15] Lei Ni. Parabolic frequency monotonicity and a theorem of Hardy-Pólya-Szegö. In Analysis, complex geometry, and mathematical physics: in honor of Duong H. Phong, volume 644 of Contemp. Math., pages 203–210. Amer. Math. Soc., Providence, RI, 2015.
- [NW07] Lei Ni and Jon Wolfson. Positive complex sectional curvature, Ricci flow and the differential sphere theorem. arXiv:0706.0332, 2007.
- [Per02] Grisha Perelman. The entropy formula for the Ricci flow and its geometric applications. arXiv:math/0211159, 2002.
- [Poo96] Chi-Cheung Poon. Unique continuation for parabolic equations. Comm. Partial Differential Equations, 21(3-4):521–539, 1996.
- [RYSZ15] Xin-An Ren, Sha Yao, Li-Ju Shen, and Guang-Ying Zhang. Constrained matrix Li-Yau-Hamilton estimates on Kähler manifolds. Math. Ann., 361(3-4):927–941, 2015.
- [SY94] R. Schoen and S.-T. Yau. Lectures on differential geometry. Conference Proceedings and Lecture Notes in Geometry and Topology, I. International Press, Cambridge, MA, 1994.
- [SZ06] Philippe Souplet and Qi S. Zhang. Sharp gradient estimate and Yau's Liouville theorem for the heat equation on noncompact manifolds. Bull. London Math. Soc., 38(6):1045–1053, 2006.
- [Ves03] Sergio Vessella. Carleman estimates, optimal three cylinder inequality, and unique continuation properties for solutions to parabolic equations. Comm. Partial Differential Equations, 28(3-4):637–676, 2003.
- [WW23] Jia-Yong Wu and Peng Wu. Harmonic and Schrödinger functions of polynomial growth on gradient shrinking Ricci solitons. *Geom. Dedicata*, 217(4):75, 2023.
- [Xu16] Guoyi Xu. Three circles theorems for harmonic functions. Math. Ann., 366(3-4):1281– 1317, 2016.
- [YZ20] Chengjie Yu and Feifei Zhao. Sharp Li-Yau-type gradient estimates on hyperbolic spaces. J. Geom. Anal., 30(1):54–68, 2020.
- [YZ22] Chengjie Yu and Feifei Zhao. Li-Yau multiplier set and optimal Li-Yau gradient estimate on hyperbolic spaces. *Potential Anal.*, 56(2):191–211, 2022.
- [Zel08] Steve Zelditch. Local and global analysis of eigenfunctions on Riemannian manifolds. In Handbook of geometric analysis. No. 1, volume 7 of Adv. Lect. Math. (ALM), pages 545–658. Int. Press, Somerville, MA, 2008.
- [Zha06] Qi S. Zhang. Some gradient estimates for the heat equation on domains and for an equation by Perelman. *Int. Math. Res. Not.*, pages Art. ID 92314, 39, 2006.
- [Zha21] Qi S. Zhang. A sharp Li-Yau gradient bound on compact manifolds. arXiv:2110.08933, 2021.
- [ZZ17] Qi S. Zhang and Meng Zhu. Li-Yau gradient bound for collapsing manifolds under integral curvature condition. Proc. Amer. Math. Soc., 145(7):3117–3126, 2017.
- [ZZ18] Qi S. Zhang and Meng Zhu. Li-Yau gradient bounds on compact manifolds under nearly optimal curvature conditions. J. Funct. Anal., 275(2):478–515, 2018.

Department of Mathematics, Statistics and Physics, Wichita State University, Wichita, KS, 67260, USA

Email address: xiaolong.li@wichita.edu

Department of Mathematics, University of California, Riverside, Riverside, CA, 92521, USA

 $Email \ address: \verb"qizhang@math.ucr.edu"$