

Maximizing the Potency of siRNA Lipid Nanoparticles for Hepatic Gene Silencing In Vivo**

Muthusamy Jayaraman,* Steven M. Ansell, Barbara L. Mui, Ying K. Tam, Jianxin Chen, Xinyao Du, David Butler, Laxman Eltepu, Shigeo Matsuda, Jayaprakash K. Narayanannair, Kallanthottathil G. Rajeev, Ismail M. Hafez, Akin Akinc, Martin A. Maier, Mark A. Tracy, Pieter R. Cullis, Thomas D. Madden, Muthiah Manoharan, and Michael J. Hope*

RNA interference (RNAi) is a ubiquitous and highly effective biological mechanism to control gene expression. Over the last few years it has been established that synthetic double-stranded, small interfering RNA (siRNA) molecules can be applied as therapeutics that potently harness RNAi in humans and animal models, by demonstrating silencing of target mRNA transcripts in rodents and non-human primates (NHP).^[1] Because of their high molecular weight and polyanionic nature, synthetic siRNAs fail to cross biological membranes by passive diffusion and therefore, generally require transmembrane drug delivery technologies to access the cytoplasm of target cells.^[2,3] Once in the cytoplasm, however, siRNAs readily load into the RNA-induced silencing complex (RISC) and direct sequence-specific cleavage of target mRNA. The resulting reduction in expression of protein, for example, can be profound and durable, lasting several weeks after administration of a single dose.^[4] This new class of therapeutics, with its unique mechanism of action and ability to specifically inhibit previously “undruggable” disease-causing proteins, has prompted research into intracellular delivery technologies suitable for parenteral administration.

Lipid nanoparticles (LNPs) containing ionizable amino lipids that self-assemble into approximately 100 nm particles when mixed with polyanionic oligonucleotides, represent the most advanced delivery platforms for systemic administration of RNAi therapeutics. Substantial progress has been made in

recent years to maximize efficacy and to achieve a favorable therapeutic index for human applications.^[2,4] The ionizable amino lipid plays a dual role in the drug delivery process. First, through electrostatic interaction with polyanionic nucleic acids, it promotes the self-assembly of formulation components into macromolecular nanoparticles encapsulating the siRNA. Second, following endocytosis of LNPs by target cells, it enables siRNA to escape the endosomal compartment and access the cell cytoplasm. In a recently published analysis of the structure–activity relationship of cationic lipid molecules we noted that permanently charged quaternary amine moieties in the hydrophilic head group region were consistently less effective in RNAi-mediated gene knockdown in vivo as compared to ionizable lipids.^[5]

Herein, we report on a detailed study aimed at further investigating the role of the amine head group in the delivery of siRNA in vivo. Through systematic structural modifications to the hydrophilic head group we have modulated the apparent acid dissociation constant (pK_a) of the ionizable amino lipids present in the LNPs and investigated the relationship between the pK_a value and activity, that is, the ability of the LNPs to deliver functionally active siRNAs into hepatocytes for eliciting gene silencing in vivo.

Amino lipid 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA, **1**), has been identified as a highly potent cationic lipid (Figure 1, Table 1)^[5] when




Figure 1. 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane (DLin-KC2-DMA, **1**). No difference in activity between enantiomers (**4**, **5**) or the racemic mixture was observed (Table 1).

incorporated into LNPs encapsulating siRNA to target murine clotting factor VII (FVII) mRNA.^[6] The hydrophobic dilinoleyl chain is known to be optimal for activity,^[5,7] hence this unsaturated alkyl chain configuration was maintained in all the lipids investigated herein and structural modifications were added to the head group region of the molecule. The formulations were prepared by a preformed vesicle method,

[*] Dr. S. M. Ansell, Dr. B. L. Mui, Dr. Y. K. Tam, Dr. J. Chen, Dr. X. Du, Dr. T. D. Madden, Dr. M. J. Hope
AlCana Technologies, Vancouver, BC (Canada)
E-mail: mhope@alcanatech.com
Homepage: <http://www.alnylam.com>

Dr. M. Jayaraman, Dr. D. Butler, Dr. L. Eltepu, Dr. S. Matsuda, Dr. J. K. Narayanannair, Dr. K. G. Rajeev, Dr. A. Akinc, Dr. M. A. Maier, Dr. M. A. Tracy, Dr. M. Manoharan
Alnylam Pharmaceuticals, Cambridge, MA (USA)
E-mail: mjayaraman@alnylam.com

Dr. I. M. Hafez, Dr. P. R. Cullis
Department of Biochemistry and Molecular Biology, University of British Columbia, Vancouver, BC (Canada)

[**] This work was supported in part by the Canadian Institutes for Health Research (CIHR) through U/1 grant FRN 59836. We would like to thank John Maraganore for helpful comments.

Supporting information for this article (experimental details) is available on the WWW under <http://dx.doi.org/10.1002/anie.201203263>.

as described in the Supporting Information (page 38), with the following lipid composition: amino lipid, distearoylphosphatidylcholine (DSPC), cholesterol and (*R*)-2,3-bis(octadecyloxy)propyl-1-(methoxy poly(ethylene glycol)2000)propylcarbamate (PEG-lipid) in the molar ratio 40/10/40/10, respectively, and a FVII siRNA/total lipid ratio of approximately 0.05 (*w/w*).^[5] To ensure a narrow particle size distribution in the range of 70–90 nm and a low polydispersity index of 0.11 ± 0.04 ($n = 56$), the particles were extruded up to three times through 80 nm membranes prior to adding the siRNA.^[5] Amino lipid pK_a values were determined for each LNP by measuring the fluorescence of 2-(*p*-toluidino)-6-naphthalene sulfonic acid (TNS) during titration from pH 3 to 10 in increments of 0.5 pH units (Supporting Information, page 40).^[8–10] The anionic TNS molecule fluoresces when associated with the surface of positively charged membranes, but is not fluorescent when free in solution, and this property has been used extensively to probe positive charge at membrane surfaces.^[8–10]

From the resulting fluorescence titration curve a pH value, at which the charge on the LNP is 50% of maximum (pK_a), can be calculated using curve-fit analysis (Figure 2). Importantly,




Figure 2. Determination of pK_a by in situ TNS fluorescence titration. The four amino lipids shown are examples illustrating the full pK_a range evaluated. Sigmoidal best-fit analyses were applied with pK_a defined as the pH at half-maximal fluorescence intensity. A) **14**, $pK_a = 4.17$; B) **52**, $pK_a = 5.73$; C) **19**, $pK_a = 6.95$; D) **48**, $pK_a = 8.12$ (see Supporting Information page 40 for details).

tantly, this method enables a pK_a determination of amino lipids in situ, which accounts for potential influences of membrane structure and neighboring lipids on the dissociation properties of the amino group. Similar pK_a values were obtained for select formulations using zeta potential measurements over the same pH titration range, but TNS was found to be the more sensitive method (data not shown).

The hydrophilic head-group region of **1** was extensively modified, producing 53 novel amino lipids with pK_a values ranging from 4.17 to 8.12 (Table 1; see the Supporting Information for full experimental details on lipid synthesis). Lipids **1**, **3**, and **53** have been reported earlier and were used as references in this study.^[5,7] The structural modifications introduced were sufficient to affect the pK_a value without significantly altering the molecular dimensions of the hydro-

philic region. This is important because having a small head group is a characteristic that enables amino lipids to adopt inverted, non-bilayer structures, which are hypothesized to be responsible for destabilizing the endosomal membrane, resulting in efficient release of siRNA into the cytoplasm of target cells.^[5,11]

A plot of pK_a value versus ED_{50} (median effective dose for FVII gene silencing in female C57BL/6 mice by i.v. administration) for all 56 amino lipids tested reveals a strong correlation between the in vivo activity and the acid dissociation constant of the amine, with a pK_a optimum seen between 6.2 and 6.5, on either side of which LNP potency rapidly decreases (Figure 3). On the other hand, significant




Figure 3. Plot of in vivo hepatic gene silencing activity vs. pK_a in mice. The 56 amino lipids were formulated in LNPs and subjected to an ED_{50} analysis (described in Supporting Information, page 39) and plotted against their pK_a . For 15 lipids the ED_{50} dose was not achieved, these are indicated by the open diamonds representing the highest dose tested for that lipid. For the remaining lipids (closed diamonds), a polynomial best-fit curve highlights the most active compounds, which exhibit an optimal pK_a between 6.2 and 6.5. Each data point is derived from a four-dose response curve with groups of $n = 4$ mice per dose.


potency differences were found for lipids with similar pK_a values. These results indicate that an optimal pK_a value is a necessary but not sufficient requirement for good in vivo activity and other structural features, such as the nature of the linker between the head group and the lipid tails, can also contribute substantially to the in vivo activity.

To further investigate the pK_a -activity relationship, a selected group of lipids were studied in more detail. These novel lipids were designed around the structure of dilinoleylmethyl-4-dimethylaminobutyrate (**16**, $ED_{50} = 0.03 \text{ mg kg}^{-1}$, $pK_a = 6.44$) also referred to as DLin-MC3-DMA (Figure 4), one of the most active cationic lipids from the group of 56 lipids. In the first group of lipids (**31**, **30**, **38**, and **37**), the distance between ester and amine is maintained at three methylene units and the substitution on the *N*-atom altered to create a pK_a range between 5.44 and 7.62.

In the second group (**14**, **15**, **17**, and **18**), the dimethylamino moiety is maintained but the distance between the

Table 1: Structures of the 56 amino lipids tested for gene silencing activity in vivo. ED₅₀ is for FVII gene silencing in mice (mg siRNA kg⁻¹) and the pK_a of the lipid was measured in situ. Lipids **1**, **3**, and **53** were reported earlier and used as references in this study.^[5,7]

ID	Novel Lipid	ED ₅₀	pK _a	ID	Novel Lipid	ED ₅₀	pK _a	ID	Novel Lipid	ED ₅₀	pK _a
1		0.10	6.68	2		0.10	5.97	3		0.25	5.94
4		0.10	6.65	5		0.10	6.79	6		0.30	6.42
7		0.15	6.43	8		>2	7.29	9		2.50	6.98
10		0.25	6.73	11		2.5	5.65	12		1.5	5.60
13		0.30	6.85	14		>5	4.17	15		0.60	5.64
16		0.03	6.44	17		0.15	6.93	18		0.65	7.16
19		0.70	6.95	20		>1	6.78	21		>1	6.77
22		0.30	6.53	23		>5	4.95	24		1.5	6.66
25		0.50	6.60	26		0.55	6.65	27		>5	7.73
28		0.15	6.60	29		1.4	7.03	30		0.30	6.17
31		4.5	5.44	32		>5	7.88	33		>3	7.25
34		4.0	7.61	35		0.47	6.49	36		0.12	6.52
37		>5	7.62	38		>1	7.31	39		0.14	6.62
40		>5	8.02	41		>5	8.11	42		>5	7.23
43		0.10	6.08	44		2.1	7.21	45		>5	7.07
46		>5	5.50	47		0.03	6.21	48		>5	8.12
49		>5	7.57	50		0.60	6.87	51		2.6	5.49
52		0.25	5.73	53		1	6.38	54		0.25	6.63
55		>1	7.27	56		0.41	6.91				






Figure 4. (6Z,9Z,28Z,31Z)-Heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (**16**, Table 1).

ester and the amine is varied from one to five methylene units to generate lipids with pK_a values ranging from 4.17 to 7.16. The plot of ED_{50} as a function of pK_a value shows that the highest potency is achieved at a pK_a of 6.44 for the siRNA LNP formulation containing lipid **16** (Figure 5, blue line).




Figure 5. Plot of hepatic gene silencing activity in mice, pK_a and calculated protonated/unprotonated amino lipid (% mole fractions) for the set of novel lipids between pK_a 4 and 8.5. Each FVII ED_{50} data point (blue) is derived from a four-dose response ($n=4$ mice per dose level). Open diamonds represent lipids for which the ED_{50} was not achieved and are shown at the highest dose tested. The mole% of each amino lipid estimated to carry a positive charge in blood, assuming pH 7.4, is shown in red and the mole% amino lipid estimated to be uncharged in acidified endosomes at pH 5.5 is shown in green.

Activity substantially decreases as the pK_a value shifts to either the acidic or basic side, underlining the importance of the ionization behavior of LNPs in the functional delivery of siRNA to hepatocytes in mice.

The surprisingly sharp pK_a value dependence of LNP activity provides more insight into the complex in vivo mechanism for this highly potent class of LNPs for siRNA delivery. In brief, it has been proposed that intravenous administration of LNPs with low to zero surface charge will minimize nonspecific binding of LNPs to tissues and proteins thus increasing the likelihood of particles reaching hepatocytes.^[5] In addition, we recently demonstrated that uptake of ionizable LNPs by hepatocytes in vitro and in vivo is mediated by apolipoprotein E (ApoE),^[12] which is also known to preferentially associate with neutral membrane surfaces.^[13]

On the other hand, LNPs must acquire sufficient positive surface charge following endocytosis to promote an electrostatic interaction with the negative surface of the endosome lumen. The resulting close proximity of LNP and membrane allows lipid mixing to occur, leading to the formation of ion pairs between the exogenous amphiphilic amine and endogenous anionic membrane lipids.^[11,14] The ion pairs prefer to

adopt inverted, non-bilayer configurations that disrupt the integrity of the endosome membrane and the siRNA is subsequently released into the cytoplasm.^[5] The ED_{50} - pK_a profile may, therefore, represent a balance between two opposing delivery requirements, namely minimum charge at pH 7.4 to maximize hepatocyte uptake by ApoE-mediated endocytosis and maximum positive charge in the acidified lumen of endosomes to promote membrane disruption.

If potency differences of structurally similar lipids are predominantly a function of pK_a value, we hypothesized that the activity of LNPs containing two or more cationic lipids should reflect the average pK_a of the particle and not the average activity measured for the individual amino lipids. To test this, LNP formulations encapsulating FVII siRNA were prepared in which the 40 mole% amino lipid component consisted of different ratios of lipids **15**, **16**, and **17** thus producing LNPs with average surface pK_a values that increased incrementally from 5.64 (**15**) to 6.93 (**17**). The measured pK_a value for mixed lipid LNPs was shown to be the same (within experimental error) as that calculated from the mole ratio of the mixed amino lipids (Supporting Information, page 41). Each LNP was administered at the same dose (0.1 mg kg^{-1} siRNA) and the concentration of murine plasma FVII protein was determined 24 h later and expressed as a percentage of that measured in a saline-treated control (Figure 6).




Figure 6. Lipid mixing supports pK_a value as the dominant factor in determining in vivo hepatic gene-silencing activity for a given set of amino lipids. Mixtures of amino lipids **15**, **16**, and **17** were made to incrementally shift the average surface pK_a value of the resulting LNP between 5.64 and 6.93. Numbers in parenthesis indicate the amount of each lipid in the mixture to make up the 40 mole% amino lipid component of the LNP. Bars represent group mean ($n=4$) \pm s.d.


As the average pK_a value increases from 5.64 (**15**) or decreases from 6.93 (**17**) by mixing with **16**, LNP activity increases. We did not identify a mixture with greater activity than **16** alone, but importantly, LNPs containing an equimolar mixture of **15** ($ED_{50}=0.6$) and **17** ($ED_{50}=0.15$), lipids that individually are 20- and 5-fold less potent than **16**, respectively, showed an average apparent pK_a value similar to **16** and equivalent activity. These data further confirm that for amino lipids of similar structure, in vivo hepatic gene silencing is driven by pK_a rather than specific head group structure or linker length (the distance between the ester and the amine). We have subsequently shown this to be true for



siRNA-Transport

M. Jayaraman,* S. M. Ansell, B. L. Mui,
Y. K. Tam, J. Chen, X. Du, D. Butler,
L. Eltepu, S. Matsuda,
J. K. Narayanannair, K. G. Rajeev,
I. M. Hafez, A. Akinc, M. A. Maier,
M. A. Tracy, P. R. Cullis, T. D. Madden,
M. Manoharan,
M. J. Hope* ————— ■■■■-■■■■

Maximizing the Potency of siRNA Lipid
Nanoparticles for Hepatic Gene Silencing
In Vivo



Der Einfluss des pK_s-Werts von ionisierbaren Lipiden auf den siRNA-Transport durch Lipidnanopartikel in vivo wurde anhand zahlreicher Modifikationen der Lipid-Kopfgruppen studiert. Dabei wurde ein Zusammenhang zwischen pK_s-Wert und Stummschaltung des Maus-FVII-Gens (FVII ED₅₀) gefunden: pK_s-Werte von 6.2–6.5 erwiesen sich als optimal (siehe Diagramm). Für das wirksamste kationische Lipid betrug ED₅₀ etwa 0.005 mg kg⁻¹ in Mäusen und < 0.03 mg kg⁻¹ in Primaten (außer Menschen).