

RESEARCH

Open Access



Maximum likelihood estimation for stochastic Lotka–Volterra model with jumps

Huiyan Zhao¹, Chongqi Zhang^{1*} and Limin Wen²

*Correspondence:
cqzhang@gzhu.edu.cn
¹School of Economics and Statistics,
Guangzhou University, Guangzhou,
P.R. China
Full list of author information is
available at the end of the article

Abstract

In this paper, we consider the stochastic Lotka–Volterra model with additive jump noises. We show some desired properties of the solution such as existence and uniqueness of positive strong solution, unique stationary distribution, and exponential ergodicity. After that, we investigate the maximum likelihood estimation for the drift coefficients based on continuous time observations. The likelihood function and explicit estimator are derived by using semimartingale theory. In addition, consistency and asymptotic normality of the estimator are proved. Finally, computer simulations are presented to illustrate our results.

Keywords: Stochastic Lotka–Volterra model; Subordinator; Maximum likelihood estimation; Stationary distribution

1 Introduction

The following famous population dynamics

$$dX_t = X_t(a - bX_t) dt$$

is often used to model population growth of a single species, where X_t represents its population size at time t , $a > 0$ is the rate of growth, and $b > 0$ represents the effect of intraspecies interaction. This equation is also known as the Lotka–Volterra model or logistic equation. In this paper, we consider one-dimensional stochastic Lotka–Volterra equation with both multiplicative Brownian noises and additive jump noises, that is,

$$\begin{cases} dX_t = X_t(a - bX_t) dt + \sigma X_t dW_t + r dJ_t, & \text{for all } t \geq 0, \\ X_0 = x_0, & \text{a.s.,} \end{cases} \quad (1.1)$$

where x_0 is a positive initial value, $a, b, \sigma, r \in (0, \infty)$, $(W_t)_{t \geq 0}$ is a one-dimensional Brownian motion (which is also known as the Wiener process), and $(J_t)_{t \geq 0}$ is a one-dimensional subordinator independent of $(W_t)_{t \geq 0}$ (the precise characterization is given below in Sect. 2). The “a.s.” above is the abbreviation of “almost surely”. Suppose that σ and r are known parameters, while a and b are unknown parameters. We will focus on the maximum likelihood estimation (MLE) of the parameter $\theta = (a, b)' \in \mathbb{R}_+^2$ based on the continuous time observations of the path $X^T := (X_t)_{0 \leq t \leq T}$. Here and after, $'$ denotes the transposition of a vector or a matrix.

Stochastic Lotka–Volterra equation, being a reasonable and popular approach to model population dynamics perturbed by random environment, has recently been studied by many authors both from a mathematical perspective and in the context of real biological dynamics. For the mathematical studies, see, for example, [1–8]. In particular, Mao et al. [3] investigated a multi-dimensional stochastic Lotka–Volterra system driven by one-dimensional standard Brownian motion. They revealed that the environmental noise could suppress population explosion. Later, Mao [4] proved a finite second moment of the stationary distribution under Brownian noise, which is very important in application.

The other case is the stochastic dynamics with Lévy noise, which can be used to describe the sudden environmental shocks, e.g., earthquakes, hurricanes, and epidemics. Bao et al. [5] considered a competitive Lotka–Volterra population model with Lévy jumps, see also Bao et al. [6]. Recently, Zhang et al. [8] considered a stochastic Lotka–Volterra model driven by α -stable noise, they got a unique positive strong solution of their model. Moreover, they proved stationary property and exponential ergodicity under relatively small noise and extinction under large enough noise.

We note that our equation (1.1) cannot be covered by [5, 6, 8]. The proof of positive solution in [5, 8] heavily depends on the explicit solution of the corresponding equation. This method does not work for our equation (1.1). We turn to prove that the hitting time of point 0 of the solution is almost surely infinite. We also prove that stationary property and exponential ergodicity do not depend on the weight of the noise, which is different from the conditions needed in [5, 6, 8]. From this point of view, equation (1.1) has its own interest.

On the other hand, the study of the influence of noise is active in the context of real ecosystems. The influence of noise is of paramount importance in open systems, and many noise induced phenomena have been found, like stochastic resonance, noise enhanced stability, noise delayed extinction, and so on. For more details, see, for example, [9–11]. However, in this paper, we shall mainly study our equation (1.1) from the view of mathematics.

We notice that there are huge works in economics and finance considering MLE of jump-diffusion models, where the data is usually observed discretely. In this case, transition densities play an important role, but their closed-form expressions cannot be obtained in general. It will be computationally expensive to conduct MLE. To overcome the difficulty, a popular method is to use closed-form expansions to approximate transition densities. To deepen this topic, we refer the reader to [12–14] and the references therein. The situation of this paper is different from the topic mentioned above. We focus on the MLE of equation (1.1) with regard to the continuous observations. The main difficulty is how to check the existence of the likelihood function. After that, we can get our MLE explicitly.

Our motivation also comes from the problem of parameter estimation for jump-type CIR (Cox–Ingersoll–Ross) process as in Barczy et al. [15] (for related topics, see, e.g., Li et al. [16]). The authors considered the following jump-type CIR process:

$$\begin{cases} dX_t = (a - bX_t) dt + \sqrt{X_t} dW_t + dJ_t, & \text{for all } t \geq 0, \\ X_0 = x_0, & \text{a.s.,} \end{cases}$$

where $(W_t)_{t \geq 0}$ and $(J_t)_{t \geq 0}$ are the same as in equation (1.1). By using the Laplace transform of the process $\int_0^t X_s ds, t \geq 0$, they proved the asymptotic properties of MLE of b under

different cases. As the authors pointed out, the asymptotic property for MLE of a or joint MLE of (a, b) is still open because of the lack of the necessary explicit Laplace transform $\int_0^t 1/X_s ds, t \geq 0$. By studying equation (1.1), we wish to bring some light to this question. For other topics of statical inference for stochastic processes, the reader can refer to the excellent monograph [17].

The rest of this paper is organized as follows. In Sect. 2, we firstly prove the existence of a unique strong positive solution of equation (1.1). After that, we derive the unique stationary distribution and the exponential ergodicity of the solution. In Sect. 3, joint MLE of parameter $\theta = (a, b)'$ is deduced from the theory of semimartingale. We prove the strong consistency and asymptotic normality in Sect. 4. In Sect. 5, we illustrate our results by computer simulations.

2 Preliminaries

Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, \mathbb{P})$ be a filtered probability space with the filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions. Equation (1.1) will be considered in this probability space. Let $(W_t)_{t \geq 0}$ in equation (1.1) be a Wiener process. We assume that the jump process $(J_t)_{t \geq 0}$ in equation (1.1) is a subordinator with zero drift. That is, its characteristic function takes the form

$$\mathbb{E}(e^{iuJ_t}) = t \int_0^\infty (e^{iuz} - 1) \nu(dz), \tag{2.1}$$

where ν is the Lévy measure concentrated on $(0, \infty)$ satisfying

$$\int_0^\infty (z \wedge 1) \nu(dz) < \infty. \tag{2.2}$$

We recall that a subordinator is an increasing Lévy process. For example, the Poisson process, α -stable subordinators, and gamma subordinators are all of this type; for more details, see, e.g., Applebaum [18] p. 52–54. Moreover, we suppose that $(W_t)_{t \geq 0}$ and $(J_t)_{t \geq 0}$ in (1.1) are independent. Let $N(dt, dz)$ be the random measure associated with the subordinator $(J_t)_{t \in \mathbb{R}_+}$, that is,

$$N(dt, dz) := \sum_{u \geq 0} \mathbf{1}_{\{\Delta J_u(\omega) \neq 0\}} \delta_{(u, \Delta J_u(\omega))}(dt, dz),$$

where δ_p is the Dirac measure at point p . Let $\tilde{N}(dt, dz) := N(dt, dz) - \nu(dz) dt$. Then, for $t \in \mathbb{R}_+$, we can write equation (1.1) as

$$\begin{aligned} X_t = x_0 &+ \int_0^t (a - bX_s)X_s ds + \int_0^t \int_0^\infty rz \nu(dz) ds \\ &+ \int_0^t \sigma X_s dW_s + \int_0^t \int_0^\infty rz \tilde{N}(ds, dz). \end{aligned} \tag{2.3}$$

The following assumptions are needed.

(A1) $a, b, \sigma, r \in (0, \infty)$ and $\int_0^\infty z \nu(dz) < \infty$.

(A2) $\int_0^\infty z^2 \nu(dz) < \infty$.

Throughout this paper, we write $\mathbb{R}, \mathbb{R}_+,$ and \mathbb{R}_{++} for real numbers, nonnegative real numbers, and positive real numbers, respectively. The value of constant C with or without

subscript may vary from line to line. First, we prove there is a unique strong positive solution for equation (2.3).

Proposition 2.1 *Assume that (A1) holds. Then, for any $x_0 \in \mathbb{R}_{++}$, there is a unique strong solution $(X_t)_{t \in \mathbb{R}_+}$ of equation (2.3) such that $\mathbb{P}(X_0 = x_0) = 1$ and $\mathbb{P}(X_t \in \mathbb{R}_{++} \text{ for all } t \in \mathbb{R}_+) = 1$.*

Proof Since the coefficients of equation (2.3) are locally Lipschitz continuous, for the given initial value $x_0 \in \mathbb{R}_{++}$, there is a unique solution X_t on $[0, \tau_e)$, where τ_e is the explosion time. In the following, we shall prove that the solution is nonexplosive and positive. The proof below is divided into two steps.

Step 1: We show that the solution of (2.3) is nonexplosive. That is, $\tau_e = \infty$ a.s. To this end, let k_0 be a sufficiently large real number such that $x_0 < k_0$. For each integer $k > k_0$, define the stopping time

$$\tau_k := \inf\{t \in [0, \tau_e) : X_t > k\},$$

and we set $\inf\{\emptyset\} = \infty$ by invention. It is easy to see that τ_k is increasing as $k \rightarrow \infty$. Let $\tau_\infty = \lim_{k \rightarrow \infty} \tau_k$, then $\tau_\infty \leq \tau_e$ a.s. If we can prove $\tau_\infty = \infty$ a.s., then $\tau_e = \infty$ a.s. Let $T > 0$ be arbitrary. For any $0 \leq t \leq T$, we have

$$\begin{aligned} X_{t \wedge \tau_k} &= x_0 + \int_0^{t \wedge \tau_k} (a - bX_s)X_s ds + \int_0^{t \wedge \tau_k} \int_0^\infty rzv(dz) ds \\ &\quad + \int_0^{t \wedge \tau_k} \sigma X_s dW_s + \int_0^{t \wedge \tau_k} \int_0^\infty rz\tilde{N}(ds, dz). \end{aligned}$$

Taking the expectation, we get

$$\begin{aligned} \mathbb{E}(X_{t \wedge \tau_k}) &= x_0 + \int_0^{t \wedge \tau_k} \int_0^\infty rzv(dz) ds + \mathbb{E}\left(\int_0^{t \wedge \tau_k} (a - bX_s)X_s ds\right) \\ &\leq x_0 + rT \int_0^\infty zv(dz) + a \int_0^t \mathbb{E}(X_{s \wedge \tau_k}) ds. \end{aligned}$$

By using Gronwall's inequality

$$\mathbb{E}(X_{T \wedge \tau_k}) \leq \left(x_0 + rT \int_0^\infty zv(dz)\right)e^{aT} \leq C.$$

On the other hand,

$$\mathbb{P}(\tau_k \leq T)k = \mathbb{E}(X_{T \wedge \tau_k} 1_{\{\tau_k \leq T\}}) \leq \mathbb{E}(X_{T \wedge \tau_k}),$$

therefore

$$\mathbb{P}(\tau_k \leq T)k \leq C.$$

Putting $k \rightarrow \infty$ yields

$$\mathbb{P}(\tau_\infty \leq T) = 0.$$

Since T is arbitrary, we get

$$\mathbb{P}(\tau_\infty = \infty) = 1.$$

Step 2: We show that the solution is positive. Let $\tilde{\tau}_0 := \inf\{t \in [0, \infty) : X_t = 0\}$. Let \tilde{k}_0 be a large enough number such that $x_0 > 1/\tilde{k}_0$. For each integer $k > \tilde{k}_0$, define the stopping time

$$\tilde{\tau}_k := \inf\{t \in [0, \infty) : X_t < 1/k\}.$$

Similarly, if we can prove $\tilde{\tau}_\infty := \lim_{k \rightarrow \infty} \tilde{\tau}_k = \infty$ a.s., then we get $\tilde{\tau}_0 = \infty$ a.s., which implies the positive solution. Let $g(x) = x - \log x$, for any $0 \leq t \leq T$, by Itô's formula

$$g(X_{t \wedge \tilde{\tau}_k}) = g(x_0) + \int_0^{t \wedge \tilde{\tau}_k} \mathcal{A}g(X_s) ds + M_{t \wedge \tilde{\tau}_k},$$

where

$$\begin{aligned} \mathcal{A}g(x) &= -bx^2 + (a + b)x + a + \sigma^2/2 \\ &\quad + r \int_0^\infty zv(dz) + \int_0^\infty [\log x - \log(x + rz)]\nu(dz) \end{aligned}$$

and $M_{t \wedge \tilde{\tau}_k}$ is a local martingale defined by

$$\begin{aligned} M_{t \wedge \tilde{\tau}_k} &= \int_0^{t \wedge \tilde{\tau}_k} \int_0^\infty [rz + \log X_{s-} - \log(X_{s-} + rz)]\tilde{N}(ds, dz) \\ &\quad + \int_0^{t \wedge \tilde{\tau}_k} \sigma(X_s - 1) dW_s. \end{aligned}$$

Note that $(M_{t \wedge \tilde{\tau}_k})_{t \in \mathbb{R}_+}$ is a true martingale and $\int_0^\infty [\log x - \log(x + rz)]\nu(dz) \leq 0$. Therefore, there exists a positive number C such that $\mathcal{A}f(x) \leq C$ for all $x \in \mathbb{R}_+$, it follows

$$\mathbb{E}(g(X_{T \wedge \tilde{\tau}_k})) \leq g(x_0) + C.$$

On the other hand,

$$\mathbb{E}(g(X_{T \wedge \tilde{\tau}_k})) \geq \mathbb{P}(\tilde{\tau}_k \leq T)(1/k + \log k).$$

By taking $k \rightarrow \infty$ and $T \rightarrow \infty$, we get

$$\mathbb{P}(\tilde{\tau}_\infty = \infty) = 1.$$

The proof is complete. □

Remark 2.2 From the study of real ecosystems (see, e.g., [19]), it is known that the effects of random fluctuations are proportional to the population size in the presence of multiplicative noise, while they are not proportional to the population size any more in the presence of additive noise. For the latter case, strongly negative values of the noise can

cause negative values of the population size. For our equation, there are in fact two types of noise: one is the multiplicative Brownian noise and the other one is additive positive jump noise. Due to the positivity of the additive noise, our equation has a unique positive solution. Therefore, the phenomena stated above are not in contradiction with our result.

In the following, our aim is to show that under assumption (A1) equation (2.3) has a unique stationary distribution. We need the following lemmas.

Lemma 2.3 *Let assumption (A1) hold. Then there exists a constant $C > 0$ such that*

$$\sup_{t \in \mathbb{R}_+} \mathbb{E}(X_t) \leq C.$$

Proof Applying Itô's formula, we have

$$\mathbb{E}(e^t X_t) = x_0 + \mathbb{E} \int_0^t e^s \left(X_s + aX_s - bX_s^2 + r \int_0^\infty zv(dz) \right) ds.$$

It is easy to see that $(a + 1)x - bx^2 + r \int_0^\infty zv(dz)$ has an upper bound for all $x \in \mathbb{R}_+$. Hence

$$e^t \mathbb{E}(X_t) \leq x_0 + C(e^t - 1),$$

which implies the desired result. □

Lemma 2.4 *Under assumption (A1), equation (2.3) has the Feller property.*

Proof The proof is essentially the same as the proof of Lemma 3.2 of [7], so we omit the proof. □

Based on the standard argument, we can obtain the following result from Lemma 2.3 and Lemma 2.4 (see, e.g., [7, 20]).

Proposition 2.5 *Under assumption (A1), equation (2.3) has a unique stationary distribution.*

Proposition 2.6 *Under assumption (A1), equation (2.3) is exponentially ergodic.*

Proof We define the Lyapunov function $V(x) = x$. Then

$$LV(x) = (a - bx)x + r \int_0^\infty zv(dz),$$

where L is the infinitesimal generator of the solution $(X_t)_{t \in \mathbb{R}_+}$. It is easy to see, for all $x \in \mathbb{R}_{++}$, there exist two positive constants γ and K such that

$$LV(x) + \gamma V(x) = (a + \gamma)x - bx^2 + r \int_0^\infty zv(dz) \leq K,$$

which satisfies the condition for exponential ergodicity in [21]. Then our desired result follows from Theorem 6.1 of [21]. □

Remark 2.7 The results above show that stationary property and exponential ergodicity do not depend on the weight of the noise. These are different from the conditions needed in [5, 6, 8], in which the results only hold under relatively small noise.

Here is a result we will use later to prove the existence of the likelihood function.

Proposition 2.8 *Suppose that assumption (A1) holds, then*

$$\int_0^t X_s^2 ds < \infty \quad a.s.$$

for $t \in \mathbb{R}_+$.

Proof From equation (2.3), for $t \in \mathbb{R}_+$, we have

$$\begin{aligned} X_t + \frac{b}{2} \int_0^t X_s^2 ds &= x_0 + \int_0^t \left(aX_s - \frac{b}{2} X_s^2 + r \int_0^\infty zv(dz) \right) ds \\ &\quad + \int_0^t \sigma X_s dW_s + \int_0^t \int_0^\infty rz \tilde{N}(ds, dz). \end{aligned}$$

By taking the expectation and noting that function $ax - b/2x^2 + r \int_0^\infty zv(dz)$ has an upper bound, we obtain

$$\mathbb{E} \int_0^t X_s^2 ds \leq x_0 + Ct,$$

which implies our result. □

3 Existence and uniqueness of MLE

In this section, we shall deduce our maximum likelihood estimation by using the semi-martingale theory.

Let $\mathbb{D} := D(\mathbb{R}_+, \mathbb{R})$ be the space of càdlàg functions (right-continuous with left limits) from \mathbb{R}_+ to \mathbb{R} . We denote by $(\mathcal{B}_t(\mathbb{D}))_{t \geq 0}$ the canonical filtration on \mathbb{D} . That is, for the canonical process $\eta = (\eta_t)_{t \geq 0}$ defined by

$$\eta_t : \mathbb{D} \ni \omega \rightarrow \omega(t) \in \mathbb{R}.$$

Then

$$\mathcal{B}_t(\mathbb{D}) := \bigcap_{\varepsilon > 0} \sigma(\eta_s; s \leq t + \varepsilon).$$

Let $\mathcal{B}(\mathbb{D})$ be the smallest σ -algebra containing $(\mathcal{B}_t(\mathbb{D}))_{t \geq 0}$. We shall call $(\mathbb{D}, \mathcal{B}(\mathbb{D}), (\mathcal{B}_t(\mathbb{D}))_{t \geq 0})$ the canonical space.

In this section, we denote by $X^\theta = (X_t^\theta)_{t \in \mathbb{R}_+}$ the unique strong solution of equation (2.3) with parameter $\theta = (a, b)'$. Let \mathbb{P}^θ be the probability measures induced by X^θ on the canonical space and \mathbb{P}_t^θ be the restriction probability measure of \mathbb{P}^θ on σ -algebra $\mathcal{B}_t(\mathbb{D})$. We can

write equation (2.3) in the form

$$\begin{aligned}
 X_t &= x_0 + \int_0^t (a - bX_s)X_s \, ds + \int_0^t \int_0^\infty rz \mathbf{1}_{\{rz \leq 1\}} \nu(dz) \, ds \\
 &\quad + \int_0^t \sigma X_s \, dW_s + \int_0^t \int_0^\infty rz \mathbf{1}_{\{rz \leq 1\}} \tilde{N}(ds, dz) \\
 &\quad + \int_0^t \int_0^\infty rz \mathbf{1}_{\{rz > 1\}} N(ds, dz).
 \end{aligned}$$

This form is the so-called Grigelionis decomposition for a semimartingale (see, e.g., [22] Theorem 2.1.2 and [23]). It follows that, under probability measure \mathbb{P}^θ , $(\eta_t)_{t \in \mathbb{R}_+}$ is a semimartingale with semimartingale characteristics $(B^\theta, C^\theta, \mu^\theta)$, where

$$B_t^\theta = \int_0^t \left[(a - b\eta_s)\eta_s + \int_0^\infty rz \mathbf{1}_{\{rz \leq 1\}} \nu(dz) \right] ds, \tag{3.1}$$

$$C_t^\theta = \sigma^2 \int_0^t \eta_s^2 \, ds \tag{3.2}$$

and

$$\mu^\theta(dt, dz) = K(\eta_t, dz) \, dt,$$

where K is a Borel kernel from \mathbb{R}_{++} to \mathbb{R}_{++} given by

$$K(x, A) = \int_0^\infty \mathbf{1}_A(rxz) \nu(dz)$$

for $t \in \mathbb{R}_+$ and $A \in \mathcal{B}(\mathbb{R}_{++})$.

In order to get the likelihood ratio process, we present the following result from [23], see also [15, 24].

Lemma 3.1 *Let Ψ be a parametric space. For $\psi, \tilde{\psi} \in \Psi$, let \mathbb{P}^ψ and $\mathbb{P}^{\tilde{\psi}}$ be two probability measures on the canonical space $(\mathbb{D}, \mathcal{B}(\mathbb{D}), (\mathcal{B}_t(\mathbb{D}))_{t \geq 0})$. We assume that, under these two probability measures, the canonical process $(\eta_t)_{t \in \mathbb{R}_+}$ is a semimartingale with characteristics $(B^\psi, C^\psi, \mu^\psi)$ and $(B^{\tilde{\psi}}, C^{\tilde{\psi}}, \mu^{\tilde{\psi}})$, respectively. We further assume that, for each $\phi \in \{\psi, \tilde{\psi}\}$, there exists a nondecreasing, continuous, and adapted process $(F_t^\phi)_{t \in \mathbb{R}_+}$ with $F_0^\phi = 0$ and a predictable process $(c_t^\phi)_{t \in \mathbb{R}_+}$ such that*

$$C_t^\phi = \int_0^t c_s^\phi \, dF_s^\phi \quad \mathbb{P}^\phi\text{-a.s. for every } t \in \mathbb{R}_+.$$

This can be guaranteed by the condition

$$\text{(B1)} \quad \mathbb{P}^\phi(\mu^\phi(\{t\} \times \mathbb{R}) = 0) = 1 \text{ for each } \phi \in \{\psi, \tilde{\psi}\}.$$

Let \mathcal{P} be the predictable σ -algebra on $\mathbb{D} \times \mathbb{R}_+$. We also assume that there exist a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R})$ -measurable function $V^{\psi, \tilde{\psi}} : \mathbb{D} \times \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}_{++}$ and a predictable \mathbb{R} -valued process $\beta^{\psi, \tilde{\psi}}$ satisfying

$$\text{(B2)} \quad \mu^\psi(dt, dz) = V^{\psi, \tilde{\psi}}(t, z) \mu^{\tilde{\psi}}(dt, dz),$$

$$\text{(B3)} \quad \int_0^t \int_{\mathbb{R}} (\sqrt{V^{\psi, \tilde{\psi}}(s, z)} - 1)^2 \mu^{\tilde{\psi}}(ds, dz) < \infty,$$

$$(B4) \quad B_t^\psi = B_t^{\tilde{\psi}} + \int_0^t c_s^\psi \beta_s^{\psi, \tilde{\psi}} dF_s^\psi + \int_0^t \int_{|z| \leq 1} z(V^{\psi, \tilde{\psi}}(s, z) - 1) \mu^{\tilde{\psi}}(ds, dz),$$

$$(B5) \quad \int_0^t c_s^\psi (\beta_s^{\psi, \tilde{\psi}})^2 dF_s^\psi < \infty$$

\mathbb{P}^ψ -a.s. for every $t \in \mathbb{R}_+$. Moreover, we assume that, for each $\phi \in \{\psi, \tilde{\psi}\}$, local uniqueness holds for the martingale problem on the canonical space corresponding to the triple $(B^\phi, C^\phi, \mu^\phi)$ with the given initial value x_0 , and \mathbb{P}^ϕ is the unique solution. Then, for any $T \in \mathbb{R}_+$, \mathbb{P}_T^ψ is absolutely continuous with respect to $\mathbb{P}_T^{\tilde{\psi}}$. The corresponding Radon–Nikodym derivative is

$$\begin{aligned} \frac{d\mathbb{P}_T^\theta}{d\mathbb{P}_T^{\tilde{\theta}}}(\eta) &= \int_0^T \beta_s^{\psi, \tilde{\psi}} d\eta_s^{\text{cont}} - \frac{1}{2} \int_0^T c_s^\psi (\beta_s^{\psi, \tilde{\psi}})^2 dF_s^\psi \\ &\quad - \int_0^T \int_{\mathbb{R}} (V^{\psi, \tilde{\psi}}(s, z) - 1) \mu^{\tilde{\psi}}(ds, dz) \\ &\quad + \int_0^T \int_{\mathbb{R}} \log(V^{\psi, \tilde{\psi}}(s, z)) N^\eta(ds, dz), \end{aligned}$$

where $(\eta_t^{\text{cont}})_{t \in \mathbb{R}_+}$ is a continuous martingale part of $(\eta_t)_{t \in \mathbb{R}_+}$ under $\mathbb{P}^{\tilde{\psi}}$ and N^η is the random jump measure of process $(\eta_t)_{t \in \mathbb{R}_+}$ defined as

$$N^\eta(\omega; dt, dz) := \sum_u 1_{\{\Delta \eta_u(\omega) \neq 0\}} \delta_{(u, \Delta \eta_u(\omega))}(dt, dz),$$

where δ_p is the Dirac measure at p .

In the following, let $\theta = (a, b)'$, $\tilde{\theta} = (\tilde{a}, \tilde{b})' \in \mathbb{R}_{++}^2$.

Proposition 3.2 *Let assumption (A1) hold, then for all $T \in \mathbb{R}_{++}$, we have*

$$\mathbb{P}_T^\theta \sim \mathbb{P}_T^{\tilde{\theta}}.$$

Moreover, under probability measure $\mathbb{P}^{\tilde{\theta}}$, we have

$$\begin{aligned} \log\left(\frac{d\mathbb{P}_T^\theta}{d\mathbb{P}_T^{\tilde{\theta}}}(\eta)\right) &= \frac{1}{\sigma^2} \int_0^T \left((a - \tilde{a}) \frac{1}{\eta_s} - (b - \tilde{b}) \right) d\eta_s^{\text{cont}} \\ &\quad - \frac{1}{2\sigma^2} \int_0^T ((a - \tilde{a}) - (b - \tilde{b})\eta_s)^2 ds, \end{aligned}$$

where η^{cont} denotes the continuous martingale part of η under probability measure $\mathbb{P}^{\tilde{\theta}}$.

Proof The main task is to check the conditions in Lemma 3.1 and then to apply Lemma 3.1 to get our result. First, it is clear that μ^θ and $\mu^{\tilde{\theta}}$ do not depend on the unknown parameter. Hence

$$\mathbb{P}^\theta(\mu^\theta(\{t\} \times \mathbb{R}) = 0) = \mathbb{P}^{\tilde{\theta}}(\mu^{\tilde{\theta}}(\{t\} \times \mathbb{R}) = 0) = \mathbb{P}^\theta(0 \cdot \nu(\mathbb{R}_{++}) = 0) = 1$$

and $V^{\theta, \tilde{\theta}} \equiv 1$. Therefore, conditions (B1)–(B3) readily hold. From (3.1) and (3.2), we see that, for $t \in \mathbb{R}_+$, $c_t^\theta = \sigma^2 \eta_t^2$ with $F_t^\theta = t$ and

$$\begin{aligned} B_t^\psi - \tilde{B}_t^{\tilde{\psi}} &= \int_0^t ((a - b\eta_s)\eta_s - (\tilde{a} - \tilde{b}\eta_s)\eta_s) ds \\ &= \int_0^t c_s^\theta \frac{1}{\sigma^2} \left(\frac{a - \tilde{a}}{\eta_s} - (b - \tilde{b}) \right) ds. \end{aligned}$$

By choosing $\beta_t^{\theta, \tilde{\theta}} = \frac{1}{\sigma^2} \left(\frac{a - \tilde{a}}{\eta_t} - (b - \tilde{b}) \right)$ for $t \in \mathbb{R}_+$, we get (B4). Now we check (B5), that is, for $t \in \mathbb{R}_+$

$$\mathbb{P}^\theta \left(\int_0^t c_s^\theta (\beta_s^{\theta, \tilde{\theta}})^2 ds < \infty \right) = 1.$$

Note that

$$\begin{aligned} \mathbb{E}_{\mathbb{P}^\theta} \int_0^t c_s^\theta (\beta_s^{\theta, \tilde{\theta}})^2 ds &= \mathbb{E}_{\mathbb{P}^\theta} \int_0^t \frac{1}{\sigma^2} (a - \tilde{a} - (b - \tilde{b})\eta_s)^2 ds \\ &\leq C(a, b, \tilde{a}, \tilde{b}, \sigma^2) \mathbb{E}_{\mathbb{P}^\theta} \int_0^t \eta_s^2 ds \\ &= C(a, b, \tilde{a}, \tilde{b}, \sigma^2) \mathbb{E}_{\mathbb{P}} \int_0^t X_s^2 ds. \end{aligned}$$

According to Proposition 2.8, we see that

$$\mathbb{E}_{\mathbb{P}^\theta} \int_0^t c_s^\theta (\beta_s^{\theta, \tilde{\theta}})^2 ds < \infty$$

for $t \in \mathbb{R}_+$, which implies that (B5) holds. Finally, the local unique property of the corresponding martingale problem comes from the fact that our equation has a unique strong solution. Therefore, all the conditions of Lemma 3.1 are satisfied. For $T \in \mathbb{R}_{++}$, by exchanging the roles of θ and $\tilde{\theta}$, we obtain

$$\mathbb{P}_T^\theta \sim \mathbb{P}_T^{\tilde{\theta}}.$$

The proof is complete. □

In the following, our aim is to estimate the parameter based on the continuous time observations of $X^T := (X_t)_{0 \leq t \leq T}$. Now, we set $\mathbb{P}^{\tilde{\theta}}$ as a fixed reference measure. Since

$$\begin{aligned} d(X_s^{\tilde{\theta}})^{\text{cont}} &= \sigma X_s^{\tilde{\theta}} dW_s \\ &= dX_s^{\tilde{\theta}} - (\tilde{a} - \tilde{b}X_s^{\tilde{\theta}})X_s^{\tilde{\theta}} ds - r dJ_s, \end{aligned}$$

then under \mathbb{P} we have

$$\begin{aligned} \log\left(\frac{d\mathbb{P}_T^\theta}{d\mathbb{P}_T^{\tilde{\theta}}}(X^{\tilde{\theta}})\right) &= \frac{1}{\sigma^2} \int_0^T \left((a - \tilde{a}) \frac{1}{X_s^{\tilde{\theta}}} - (b - \tilde{b}) \right) [dX_s^{\tilde{\theta}} - (\tilde{a} - \tilde{b}X_s^{\tilde{\theta}})X_s^{\tilde{\theta}} ds - r dJ_s] \\ &\quad - \frac{1}{2\sigma^2} \int_0^T ((a - \tilde{a}) - (b - \tilde{b})X_s^{\tilde{\theta}})^2 ds \\ &= \frac{1}{\sigma^2} \left[(a - \tilde{a}) \int_0^T \frac{1}{X_s^{\tilde{\theta}}} (dX_s^{\tilde{\theta}} - r dJ_s) - (b - \tilde{b}) \int_0^T (dX_s^{\tilde{\theta}} - r dJ_s) \right. \\ &\quad \left. - \frac{1}{2}(a^2 - \tilde{a}^2)T - \frac{1}{2}(b^2 - \tilde{b}^2) \int_0^T X_s^{\tilde{\theta}^2} ds + (ab - \tilde{a}\tilde{b}) \int_0^T X_s^{\tilde{\theta}} ds \right]. \end{aligned}$$

Next, we can define the log-likelihood function with respect to the dominated measure $\mathbb{P}^{\tilde{\theta}}$ as

$$l_T(\theta; X^T) = \sigma^2 \log \frac{d\mathbb{P}_t^\theta}{d\mathbb{P}_t^{\tilde{\theta}}}(X^T).$$

Then the maximum likelihood estimator (MLE) $\hat{\theta}_T$ of the unknown parameter θ is defined as

$$\hat{\theta}_T := \arg \max_{\theta \in \mathbb{R}_{++}^2} l_T(\theta; X^T).$$

Proposition 3.3 *If assumption (A1) holds, then for every $T \in \mathbb{R}_{++}$, there exists a unique MLE $\hat{\theta}_T$ with the form*

$$\hat{\theta}_T = \left(\frac{\int_0^T X_s^2 ds \int_0^T \frac{1}{X_s} (dX_s - r dJ_s) - \int_0^T X_s ds \int_0^T (dX_s - r dJ_s)}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2}, \frac{\int_0^T X_s ds \int_0^T \frac{1}{X_s} (dX_s - r dJ_s) - T \int_0^T (dX_s - r dJ_s)}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \right) \tag{3.3}$$

almost surely.

Proof By Hölder’s inequality, we have

$$\left(\int_0^T X_s ds \right)^2 \leq \left(\left(\int_0^T ds \right)^{1/2} \left(\int_0^T X_s^2 ds \right)^{1/2} \right)^2 = T \int_0^T X_s^2 ds$$

and

$$\mathbb{P}\left(T \int_0^T X_s^2 ds - \left(\int_0^T X_s ds\right)^2 = 0\right) = \mathbb{P}(X_s \equiv k, s \in [0, T], \text{ for some number } k).$$

From equation (1.1), we see that the constant solution is impossible. Hence,

$$T \int_0^T X_s^2 ds - \left(\int_0^T X_s ds\right)^2 > 0 \quad \text{a.s.}$$

It follows that (3.3) is well defined almost surely. Note that

$$J_t = \sum_{0 \leq s \leq t} \Delta J_s = \sum_{0 \leq s \leq t} \Delta X_s / r.$$

Hence, for $t \in [0, T]$, J_t is a measurable function of X^T , which implies that (3.3) is a true statistic. Next, we have

$$\begin{aligned} \frac{\partial}{\partial a} l_T(\theta; X^T) &= \int_0^T \frac{1}{X_s} (dX_s - r dJ_s) - aT + b \int_0^T X_s ds, \\ \frac{\partial}{\partial b} l_T(\theta; X^T) &= - \int_0^T (dX_s - r dJ_s) + a \int_0^T X_s ds - b \int_0^T X_s^2 ds. \end{aligned}$$

By direct calculation, we can get our desired result. □

4 Asymptotic properties

In order to get the asymptotic properties of our estimator, we need the following result.

Proposition 4.1 *Let assumptions (A1)–(A2) hold. Then, for any $x_0 \in \mathbb{R}_{++}$, there exists a positive constant C such that*

$$\limsup_{t \in \mathbb{R}_+} \frac{1}{t} \int_0^t X_s^2 ds \leq C \quad a.s.$$

Proof We follow the approach used in Lemma 4.1 of [4]. By the exponential martingale inequality, we get

$$\mathbb{P} \left(\sup_{0 \leq t \leq k} \left(\int_0^t \sigma X_s dW_s - \frac{\alpha}{2} \int_0^t \sigma^2 X_s^2 ds \right) > \frac{2}{\alpha} \log k \right) \leq \frac{1}{k^2},$$

where we choose $\alpha = b/(2\sigma^2)$. The well-known Borel–Cantelli lemma implies that for almost all $\omega \in \Omega$, there is a random integer $k_0 = k_0(\omega)$ such that

$$\int_0^t \sigma X_s dW_s \leq \frac{2}{\alpha} \log k + \frac{\alpha}{2} \int_0^t \sigma^2 X_s^2 ds$$

for all $t \in [0, k]$, $k \geq k_0$, almost surely. Substituting this to our equation (2.3), we have

$$\begin{aligned} X_t &\leq x_0 + \frac{2}{\alpha} \log k + \int_0^t \left(aX_s - \frac{3}{4} bX_s^2 + r \int_0^\infty zv(dz) \right) ds \\ &\quad + \int_0^t \int_0^\infty rz \tilde{N}(ds, dz) \end{aligned}$$

for all $t \in [0, k]$, $k \geq k_0$, almost surely. Hence

$$\begin{aligned} \frac{b}{2} \int_0^t X_s^2 ds &\leq x_0 + \frac{2}{\alpha} \log k + \int_0^t \left(aX_s - \frac{b}{4} X_s^2 + r \int_0^\infty zv(dz) \right) ds \\ &\quad + \int_0^t \int_0^\infty rz \tilde{N}(ds, dz) \\ &\leq x_0 + \frac{2}{\alpha} \log k + C_1 t + \int_0^t \int_0^\infty rz \tilde{N}(ds, dz) \end{aligned}$$

for all $t \in [0, k], k \geq k_0$, almost surely. Now, for almost all $\omega \in \Omega$, let $k \geq k_0$ and $k - 1 \leq t \leq k$, then

$$\frac{1}{t} \int_0^t X_s^2 ds \leq \frac{2}{(k-1)b} \left(x_0 + \frac{2}{\alpha} \log k + C_1 k + \int_0^k \int_0^\infty rz \tilde{N}(ds, dz) \right).$$

Letting $t \rightarrow \infty$ and hence $k \rightarrow \infty$, we obtain

$$\limsup_{t \rightarrow \infty} \frac{1}{t} \int_0^t X_s^2 ds \leq \frac{2}{b} \left(x_0 + C_2 + \limsup_{k \rightarrow \infty} \frac{1}{k} \int_0^k \int_0^\infty rz \tilde{N}(ds, dz) \right). \tag{4.1}$$

Under assumption (A2), note that $(\int_0^t \int_0^\infty rz \tilde{N}(ds, dz))_{t \in \mathbb{R}_+}$ is a local martingale with Meyer’s angle bracket process $(\int_0^t \int_0^\infty rz^2 \nu(dz) ds)_{t \in \mathbb{R}_+}$ and

$$\lim_{t \rightarrow \infty} \int_0^t \frac{\int_0^\infty rz^2 \nu(dz)}{(1+s)^2} ds < \infty.$$

By using the strong law of large numbers for local martingales (Lemma A.1), we get

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t \int_0^\infty rz \tilde{N}(ds, dz) = 0$$

almost surely. Hence, there exists a constant C_2 such that

$$\limsup_{k \rightarrow \infty} \frac{1}{k} \int_0^k \int_0^\infty rz \tilde{N}(ds, dz) \leq C_2$$

almost surely. Combining this with (4.1), we complete the proof. □

Corollary 4.2 *Suppose that assumptions (A1)–(A2) hold. The invariant measure π has a finite second moment, moreover*

$$\lim_{t \in \mathbb{R}_+} \frac{1}{t} \int_0^t X_s ds = \int_0^\infty y \pi(dy) \quad a.s.$$

and

$$\lim_{t \in \mathbb{R}_+} \frac{1}{t} \int_0^t X_s^2 ds = \int_0^\infty y^2 \pi(dy) \quad a.s.$$

Proof The proof of the first result is essentially the same as the proof of Theorem 4.2 in [4], and the second is the same as the proof in [20]. So, we omit them. □

In the following, we present the weak and strong consistency of our estimator.

Theorem 4.3 *Under assumption (A1), the estimator $\hat{\theta}_T = (\hat{a}_T, \hat{b}_T)'$ of $\theta = (a, b)'$ admits the weak consistency, i.e.,*

$$\hat{\theta}_T \xrightarrow{\mathbb{P}} \theta \quad \text{as } T \rightarrow \infty,$$

where $\xrightarrow{\mathbb{P}}$ denotes the convergence in probability. Under assumptions (A1)–(A2), the estimator $\hat{\theta}_T = (\hat{a}_T, \hat{b}_T)'$ of $\theta = (a, b)'$ admits the strong consistency, i.e.,

$$\hat{\theta}_T \rightarrow \theta \quad \text{a.s. as } T \rightarrow \infty.$$

Proof We have

$$\begin{aligned} \hat{a}_T &= a + \frac{\int_0^T X_s^2 ds \int_0^T dW_s - \int_0^T X_s ds \int_0^T X_s dW_s}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2}, \\ \hat{b}_T &= b + \frac{\int_0^T X_s ds \int_0^T dW_s - T \int_0^T X_s dW_s}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2}. \end{aligned}$$

Note that

$$\begin{aligned} \hat{a}_T - a &= \frac{\int_0^T X_s^2 ds \int_0^T dW_s - \int_0^T X_s ds \int_0^T X_s dW_s}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \\ &= \frac{\int_0^T X_s^2 ds \int_0^T dW_s}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} - \frac{\int_0^T X_s ds \int_0^T X_s dW_s}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \\ &:= I_1 - I_2. \end{aligned}$$

Case 1: Under assumption (A1), for I_1 , we have

$$|I_1| \leq \left| \int_0^T dW_s / T \right|.$$

According to the strong law of large numbers for continuous local martingales (Lemma A.2), we have

$$\lim_{T \rightarrow \infty} \int_0^T dW_s / T = 0 \quad \text{a.s.}$$

Then we obtain $\lim_{T \rightarrow \infty} I_1 = 0$, a.s. For I_2 , we have

$$|I_2| \leq \left| \frac{\int_0^T X_s ds}{T} \frac{\int_0^T X_s dW_s}{\int_0^T X_s^2 ds} \right|.$$

Note $(\frac{\int_0^T X_s ds}{T})_{T>0}$ is tight. Indeed, by Lemma 2.3, for $M > 0$, we have

$$\mathbb{P}\left(\frac{\int_0^T X_s ds}{T} > M\right) \leq \frac{\int_0^T \mathbb{E}X_s ds}{MT} \leq \frac{C}{M}. \tag{4.2}$$

On the other hand, by Proposition 2.5 and Proposition 2.6, we have

$$\lim_{T \rightarrow \infty} \frac{\int_0^T X_s^2 ds}{T} = \int_0^\infty y^2 \pi(dy) > 0,$$

where π is the unique invariant measure. It follows that

$$\lim_{T \rightarrow \infty} \int_0^T X_s^2 ds = \infty \quad \text{a.s.}$$

By Proposition 2.8, we also have

$$\int_0^T X_s^2 ds < \infty \quad \text{a.s.}$$

for each $T > 0$. Then, again by Lemma A.2, we get

$$\lim_{T \rightarrow \infty} \frac{\int_0^T X_s dW_s}{\int_0^T X_s^2 ds} = 0 \quad \text{a.s.} \tag{4.3}$$

From (4.2) and (4.3), we get $\lim_{T \rightarrow \infty} I_2 = 0$ in probability. Therefore, we obtain $\lim_{T \rightarrow \infty} \hat{a}_T = a$ in probability. Similarly, we can prove $\lim_{T \rightarrow \infty} \hat{b}_T = b$ in probability.

Case 2: Under assumptions (A1)–(A2). For I_1 , we have

$$I_1 = \frac{\int_0^T X_s^2 ds/T}{\int_0^T X_s^2 ds/T - (\int_0^T X_s ds/T)^2} \int_0^T dW_s/T.$$

According to Corollary 4.2 and Lemma A.2, we have $\lim_{T \rightarrow \infty} I_1 = 0$, a.s. For I_2 , we have

$$I_2 = \frac{\int_0^T X_s ds \int_0^T X_s^2 ds/T^2}{\int_0^T X_s^2 ds/T - (\int_0^T X_s ds/T)^2} \frac{\int_0^T X_s dW_s}{\int_0^T X_s^2 ds}.$$

Again by Corollary 4.2 and Lemma A.2, we immediately get $\lim_{T \rightarrow \infty} I_2 = 0$, a.s. Therefore, we obtain $\lim_{T \rightarrow \infty} \hat{a}_T = a$ a.s. Similarly, we can prove $\lim_{T \rightarrow \infty} \hat{b}_T = b$ a.s. We complete the proof. □

For simplicity of our notations, we denote $\mu_1 := \int_0^\infty y\pi(dy)$ and $\mu_2 := \int_0^\infty y^2\pi(dy)$. Now we present the following asymptotic normality.

Theorem 4.4 *Under assumptions (A1)–(A2). The estimator $\hat{\theta}_T$ of θ is asymptotically normal, i.e.,*

$$\sqrt{T}(\hat{\theta}_T - \theta) \xrightarrow{\mathcal{D}} N(0, \Sigma)$$

as $T \rightarrow \infty$, where $\xrightarrow{\mathcal{D}}$ denotes the convergence in distribution, $\Sigma = AA'$ and

$$A = \frac{1}{\mu_2 - \mu_1^2} \begin{pmatrix} \mu_2 + \mu_1^2 & \mu_1\sqrt{\mu_2} \\ 2\mu_1 & \sqrt{\mu_2} \end{pmatrix}.$$

By a random scaling, we also have

$$\sqrt{T} \begin{pmatrix} 1 & -\frac{\int_0^T X_s ds}{T} \\ -\frac{2 \int_0^T X_s ds/T}{\sqrt{\int_0^T X_s^2 ds/T}} & \frac{\int_0^T X_s^2 ds/T + (\int_0^T X_s ds/T)^2}{\sqrt{\int_0^T X_s^2 ds/T}} \end{pmatrix} (\hat{\theta}_T - \theta) \xrightarrow{\mathcal{D}} N(0, I)$$

as $T \rightarrow \infty$, where I is the identity matrix.

Proof We write our estimator in the matrix form

$$\hat{\theta} - \theta = \begin{pmatrix} \frac{\int_0^T X_s^2 ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} & \frac{\int_0^T X_s ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \\ \frac{\int_0^T X_s ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} & \frac{T}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \end{pmatrix} \begin{pmatrix} \int_0^T dW_s \\ -\int_0^T X_s dW_s \end{pmatrix}.$$

Let

$$M_t := \begin{pmatrix} \int_0^t dW_s \\ -\int_0^t X_s dW_s \end{pmatrix},$$

then $(M_t)_{t \in \mathbb{R}_+}$ is a 2-dimensional continuous local martingale with $M_0 = 0$ a.s. and with quadratic variation process

$$[M]_t = \begin{pmatrix} t & -\int_0^t X_s ds \\ -\int_0^t X_s ds & \int_0^t X_s^2 ds \end{pmatrix}.$$

Let

$$Q(t) := \begin{pmatrix} 1/\sqrt{t} & 0 \\ 0 & 1/\sqrt{t} \end{pmatrix}.$$

Then, by Corollary 4.2, we have

$$Q(t)[M]_t Q(t)^T \rightarrow \begin{pmatrix} 1 & \mu_1 \\ \mu_1 & \mu_2 \end{pmatrix} = \zeta \zeta' \quad \text{a.s. as } T \rightarrow \infty,$$

where

$$\zeta := \begin{pmatrix} 1 & 0 \\ \mu_1 & \sqrt{\mu_2} \end{pmatrix}.$$

By applying Lemma (A.3), we get

$$1/\sqrt{T} M_T \xrightarrow{\mathcal{D}} \zeta Z \quad \text{as } T \rightarrow \infty, \tag{4.4}$$

where Z is a 2-dimensional standard normal random vector. Note that, again by Corollary 4.2,

$$\begin{aligned} & T \begin{pmatrix} \frac{\int_0^T X_s^2 ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} & \frac{\int_0^T X_s ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \\ \frac{\int_0^T X_s ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} & \frac{T}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \end{pmatrix} \\ & \rightarrow \frac{1}{\mu_2 - \mu_1^2} \begin{pmatrix} \mu_2 & \mu_1 \\ \mu_1 & 1 \end{pmatrix} \quad \text{a.s. } T \rightarrow \infty. \end{aligned} \tag{4.5}$$

Combining (4.4) with (4.5), by using Slutsky’s lemma, we have

$$\begin{aligned} & \sqrt{T}(\hat{\theta}_T - \theta) \\ &= T \left(\begin{array}{cc} \frac{\int_0^T X_s^2 ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} & \frac{\int_0^T X_s ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \\ \frac{\int_0^T X_s ds}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} & \frac{T}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \end{array} \right) \cdot \frac{1}{\sqrt{T}} M_T \\ &\xrightarrow{\mathcal{D}} \frac{1}{\mu_2 - \mu_1^2} \begin{pmatrix} \mu_2 & \mu_1 \\ \mu_1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \mu_1 & \sqrt{\mu_2} \end{pmatrix} Z \\ &= \frac{1}{\mu_2 - \mu_1^2} \begin{pmatrix} \mu_2 + \mu_1^2 & \mu_1 \sqrt{\mu_2} \\ 2\mu_1 & \sqrt{\mu_2} \end{pmatrix} Z \\ &= AZ \end{aligned}$$

as $T \rightarrow \infty$. We have proved the first result. Next, it is easy to see that

$$\left(\begin{array}{cc} 1 & -\frac{\int_0^T X_s ds}{T} \\ -\frac{2 \int_0^T X_s ds / T}{\sqrt{\int_0^T X_s^2 ds / T}} & \frac{\int_0^T X_s^2 ds / T + (\int_0^T X_s ds / T)^2}{\sqrt{\int_0^T X_s^2 ds / T}} \end{array} \right) \rightarrow \left(\begin{array}{cc} 1 & -\mu_1 \\ -\frac{2\mu_1}{\sqrt{\mu_2}} & \frac{\mu_2 + \mu_1^2}{\sqrt{\mu_2}} \end{array} \right) = A^{-1}$$

a.s. $T \rightarrow \infty$. Again by Slutsky’s lemma, we have

$$\begin{aligned} & \sqrt{T} \left(\begin{array}{cc} 1 & -\frac{\int_0^T X_s ds}{T} \\ -\frac{2 \int_0^T X_s ds / T}{\sqrt{\int_0^T X_s^2 ds / T}} & \frac{\int_0^T X_s^2 ds / T + (\int_0^T X_s ds / T)^2}{\sqrt{\int_0^T X_s^2 ds / T}} \end{array} \right) (\hat{\theta}_T - \theta) \\ &\xrightarrow{\mathcal{D}} A^{-1}AZ = Z, \quad \text{as } T \rightarrow \infty. \end{aligned}$$

We finish the proof. □



5 Simulation results

In this section, we present some computer simulations. First, we apply Euler–Maruyama method to illustrate the stationary solution of equation (1.1) under assumption (A1). We consider the following two examples.

Examples 5.1 Let $a = 5, b = 1, \sigma = 1, r = 1$, and $x_0 = 10$ for equation (1.1). Let $(J_t)_{t \geq 0}$ be a Poisson process with intensity 1. Note that the Poisson process with intensity 1 is a subordinator with Lévy measure $\nu(dz) = \delta_1(dz)$. It follows from Proposition 2.5 there is a unique stationary distribution. We apply the Euler–Maruyama method to perform a computer simulation of 30,000 iterations of the single path of X_t with initial value $x_0 = 10, T = 30$, and step size $\Delta = 0.001$, which is shown in Fig. 1.

Examples 5.2 Let $a = 5, b = 1, \sigma = 1, r = 1$, and $x_0 = 10$ for equation (1.1). Let $(J_t)_{t \geq 0}$ be a compound Poisson process with exponentially distributed jump size, namely

$$\nu(dz) = c\lambda e^{-\lambda z} I_{(0,\infty)}(z) dz.$$



We set $c = 1$ and $\lambda = 10$. It is easy to see that ν satisfies assumption (A1). Again by Proposition 2.5 there is a unique stationary distribution. We apply the Euler–Maruyama method to perform a computer simulation of 2000 iterations of the single path of X_t with initial value $x_0 = 10$, $T = 20$, and step size $\Delta = 0.01$, which is shown in Fig. 2.


From the simulation paths of Fig. 1 and Fig. 2, we can see their stationary trends. The distributions implied by their histograms can be seen as the approximations of the stationary distributions.

Next, we exhibit the consistency of the MLE. It follows from Theorem 3.3 that our MLE is

$$\hat{\theta}_T = \begin{pmatrix} \frac{\int_0^T X_s^2 ds \int_0^T \frac{1}{X_s} (dX_s - r dJ_s) - \int_0^T X_s ds \int_0^T (dX_s - r dJ_s)}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \\ \frac{\int_0^T X_s ds \int_0^T \frac{1}{X_s} (dX_s - r dJ_s) - T \int_0^T (dX_s - r dJ_s)}{T \int_0^T X_s^2 ds - (\int_0^T X_s ds)^2} \end{pmatrix}. \tag{5.1}$$

Table 1 Mean and standard deviation of the estimators

Parameters	Ex.1					Ex.2			
	$\hat{a} - a$		$\hat{b} - b$			$\hat{a} - a$		$\hat{b} - b$	
	Mean	Stddev	Mean	Stddev	Mean	Stddev	Mean	Stddev	
$\Delta = 0.01$	$T = 10$	0.34571	1.07444	0.07110	0.21637	0.37412	1.01528	0.08287	0.21657
	$T = 10^2$	0.03285	0.30957	0.00639	0.06176	0.04797	0.30105	0.01051	0.06318
	$T = 10^3$	0.00469	0.09906	0.00082	0.01975	0.00384	0.09616	0.00091	0.01993



We perform 1000 Monte Carlo simulations of the sample paths generated by Example 5.1 and Example 5.2. The results are presented in Table 1. We see that the estimate errors become small as the observation time increases. This is consistent with our theoretical result.


Finally, we investigate the asymptotic distribution of the MLE in (5.1). That is, we will focus on the distribution of the following statistic:

$$\beta_T := \sqrt{T} \begin{pmatrix} 1 & -\frac{\int_0^T X_s ds}{T} \\ -\frac{2 \int_0^T X_s ds/T}{\sqrt{\int_0^T X_s^2 ds/T}} & \frac{\int_0^T X_s^2 ds/T + (\int_0^T X_s ds/T)^2}{\sqrt{\int_0^T X_s^2 ds/T}} \end{pmatrix} (\hat{\theta}_T - \theta).$$

We perform 1000 Monte Carlo simulations for Example 5.1 with $a = 1, b = 7, \sigma = 1, r = 1, T = 10,$ and $\Delta = 0.01$ and $x_0 = 10$. The 3D histogram of the 1000 simulations is presented in Fig. 3. By comparing the 3D histogram of the 1000 simulations to the 3D histogram of standard normal distribution (Fig. 3), we can see the tendency of the joint normality. The trend of normality of each element of the estimator β_T can be seen from Fig. 4, where the histogram of each element is given.

6 Conclusions

In this paper, we consider a stochastic Lotka–Volterra model with both multiplicative Brownian noises and additive jump noises. Some desired properties of the solution, such as existence and uniqueness of positive strong solution, unique stationary distribution, and exponential ergodicity, are proved. We also investigate the maximum likelihood estimation for the drift coefficients based on continuous time observations. The likelihood function and explicit estimator are derived by using semimartingale theory, and then con-



sistency and asymptotic normality of the estimator are proved. Finally, we give some computer simulations, which are consistent with our theoretical results. The case with multiplicative jump noises will be the subject of future investigation.

Appendix: Limit theorems for local martingales

In this section, we recall some limit theorems for local martingales. The first one is a strong law of large numbers for local martingales, e.g., [25].

Lemma A.1 *Let $(M_t)_{t \in \mathbb{R}_+}$ be a one-dimensional local martingale vanishing at time $t = 0$. For $t \in \mathbb{R}_+$, we define*

$$\rho_M(t) := \int_0^t \frac{1}{(1+s)^2} d\langle M \rangle_s,$$

where $(\langle M \rangle_t)_{t \in \mathbb{R}_+}$ is Meyer’s angle bracket process. Then

$$\lim_{t \rightarrow \infty} \rho_M(t) < \infty \quad a.s.$$

implies

$$\lim_{t \rightarrow \infty} \frac{M_t}{t} = 0 \quad a.s.$$

The next result is a strong law of large numbers for continuous local martingales, see, e.g., Lemma 17.4 of [26].

Lemma A.2 *Let $(M_t)_{t \in \mathbb{R}_+}$ be a one-dimensional square-integrable continuous local martingale vanishing at time $t = 0$. Let $([M]_t)_{t \in \mathbb{R}_+}$ be the quadratic variation process of M such that, for $t \in \mathbb{R}_+$,*

$$[M]_t < \infty \quad a.s.$$

and

$$[M]_t \rightarrow \infty \quad a.s. \text{ as } t \rightarrow \infty.$$

Then

$$\lim_{t \rightarrow \infty} \frac{M_t}{[M]_t} = 0 \quad a.s.$$

The last one is about the asymptotic behavior of continuous multivariate local martingales, see Theorem 4.1 of [27].

Lemma A.3 *Let $(M_t)_{t \in \mathbb{R}_+}$ be a d -dimensional square-integrable continuous local martingale vanishing at time $t = 0$. Suppose that there exists a function $Q : \mathbb{R}_+ \rightarrow \mathbb{R}^{d \times d}$ such that $Q(t)$ is an invertible (non-random) matrix for all $t \in \mathbb{R}_+$, $\lim_{t \rightarrow \infty} \|Q(t)\| = 0$ and*

$$Q(t)[M]_t Q(t)^T \xrightarrow{\mathbb{P}} \zeta \zeta^T \quad \text{as } t \rightarrow \infty,$$

where $\|Q(t)\| := \sup\{|Q(t)x| : x \in \mathbb{R}^d, |x| = 1\}$, $[M]_t$ is the quadratic variation process of M and ζ is a $d \times d$ random matrix. Then

$$Q(t)M_t \xrightarrow{\mathcal{D}} \zeta Z \quad \text{as } t \rightarrow \infty,$$

where Z is a d -dimensional standard normally distributed random vector independent of ζ .

Acknowledgements

This work was supported in part by the National Natural Science Foundation of China (11401029), Teacher Research Capacity Promotion Program of Beijing Normal University Zhuhai, the National Natural Science Foundation of China (11671104), the National Natural Science Foundation of China (71761019), and Jiangxi Provincial Natural Science Foundation (20171ACB21022). The authors appreciate the anonymous referees for their valuable suggestions and questions.

Competing interests

The authors declare that they have no competing interest.

Authors' contributions

The first author and the corresponding author contributed to Sects. 1, 2, 3, and 4. The third author contributed to Sect. 5. All authors read and approved the final manuscript.

Author details

¹School of Economics and Statistics, Guangzhou University, Guangzhou, P.R. China. ²Department of Statistics, Jiangxi Normal University, Jiangxi, P.R. China.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 27 October 2017 Accepted: 18 April 2018 Published online: 25 April 2018

References

1. Bahar, A., Mao, X.: Stochastic delay Lotka–Volterra model. *J. Math. Anal. Appl.* **292**(2), 364–380 (2004)
2. Bahar, A., Mao, X.: Stochastic delay population dynamics. *Int. J. Pure Appl. Math.* **11**, 377–400 (2004)
3. Mao, X., Marion, G., Renshaw, E.: Environmental Brownian noise suppresses explosions in population dynamics. *Stoch. Process. Appl.* **97**(1), 95–110 (2002)
4. Mao, X.: Stationary distribution of stochastic population systems. *Syst. Control Lett.* **60**(6), 398–405 (2011)
5. Bao, J., Mao, X., Yin, G., Yuan, C.: Competitive Lotka–Volterra population dynamics with jumps. *Nonlinear Anal., Theory Methods Appl.* **74**(17), 6601–6616 (2011)
6. Bao, J., Yuan, C.: Stochastic population dynamics driven by Lévy noise. *J. Math. Anal. Appl.* **391**(2), 363–375 (2012)
7. Tong, J., Zhang, Z., Bao, J.: The stationary distribution of the facultative population model with a degenerate noise. *Stat. Probab. Lett.* **83**(2), 655–664 (2013)
8. Zhang, Z., Zhang, X., Tong, J.: Exponential ergodicity for population dynamics driven by α -stable processes. *Stat. Probab. Lett.* **125**, 149–159 (2017)
9. Spagnolo, B., Valenti, D., Fiasconaro, A.: Noise in ecosystems: a short review. *Math. Biosci. Eng.* **1**(1), 185–211 (2004)
10. Valenti, D., Fiasconaro, A., Spagnolo, B.: Stochastic resonance and noise delayed extinction in a model of two competing species. *Phys. A, Stat. Mech. Appl.* **331**(3–4), 477–486 (2004)

11. La Cognata, A., Valenti, D., Dubkov, A.A., Spagnolo, B.: Dynamics of two competing species in the presence of Lévy noise sources. *Phys. Rev. E* **82**(1), 011121 (2010)
12. Ait-Sahalia, Y.: Transition densities for interest rate and other nonlinear diffusions. *J. Finance* **54**(4), 1361–1395 (1999)
13. Li, C.: Maximum-likelihood estimation for diffusion processes via closed-form density expansions. *Ann. Stat.* **41**(3), 1350–1380 (2013)
14. Li, C., Chen, D.: Estimating jump-diffusions using closed-form likelihood expansions. *J. Econom.* **195**(1), 51–70 (2016)
15. Barczy, M., Alaya, M.B., Kebaier, A., Pap, G.: Asymptotic properties of maximum likelihood estimator for the growth rate for a jump-type CIR process based on continuous time observations. *arXiv preprint. arXiv:1609.05865* (2016)
16. Li, Z., Ma, C.: Asymptotic properties of estimators in a stable Cox–Ingersoll–Ross model. *Stoch. Process. Appl.* **125**(8), 3196–3233 (2015)
17. Kutoyants, Y.A.: *Statistical Inference for Ergodic Diffusion Processes*. Springer, Berlin (2010)
18. Applebaum, D.: *Lévy Processes and Stochastic Calculus*, 2nd edn. Cambridge University Press, Cambridge (2009)
19. Valenti, D., Denaro, G., Spagnolo, B., Mazzola, S., Basilone, G., Conversano, F., Bonanno, A.: Stochastic models for phytoplankton dynamics in Mediterranean Sea. *Ecol. Complex.* **27**, 84–103 (2016)
20. Khasminskii, R.: *Stochastic Stability of Differential Equations*, vol. 66. Springer, Berlin (2011)
21. Meyn, S.P., Tweedie, R.L.: Stability of Markovian processes III: Foster–Lyapunov criteria for continuous-time processes. *Adv. Appl. Probab.* **25**(3), 518–548 (1993)
22. Jacod, J., Protter, P.: *Discretization of Processes. Stochastic Modelling and Applied Probability*, vol. 67. Springer, Berlin (2011)
23. Jacod, J., Shiryaev, A.N.: *Limit Theorems for Stochastic Processes*, vol. 288. Springer, Berlin (2013)
24. Sorensen, M.: Likelihood methods for diffusions with jumps. In: *Statistical Inference in Stochastic Processes*, pp. 67–105 (1991)
25. Liptser, R.S.: A strong law of large numbers for local martingales. *Stochastics* **3**(1–4), 217–228 (1980)
26. Liptser, R.S., Shiryaev, A.N.: *Statistics of Random Processes II. Applications*, 2nd edn. Springer, Berlin (2001)
27. van Zanten, H.: A multivariate central limit theorem for continuous local martingales. *Stat. Probab. Lett.* **50**(3), 229–235 (2000)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
